

DISTINGUISHED CATEGORIES AND THE ZILBER-PINK CONJECTURE

FABRIZIO BARROERO AND GABRIEL A. DILL

ABSTRACT. We propose an axiomatic approach towards studying unlikely intersections by introducing the framework of distinguished categories. This includes commutative algebraic groups and mixed Shimura varieties. It allows us to define all basic concepts of the field and prove some fundamental facts about them, e.g., the defect condition.

In some categories that we call very distinguished, we are able to show some implications between Zilber-Pink statements with respect to base change. This yields unconditional results, i.e., the Zilber-Pink conjecture for a complex curve in \mathcal{A}_2 that cannot be defined over $\bar{\mathbb{Q}}$, a complex curve in the g -th fibered power of the Legendre family, and a complex curve in the base change of a semiabelian variety over $\bar{\mathbb{Q}}$.

CONTENTS

1. Introduction	1
2. Distinguished categories	6
3. Connected mixed Shimura varieties	8
4. Intersections of special subvarieties of connected mixed Shimura varieties	13
5. Proof of Proposition 3.3	18
6. Special and weakly special subvarieties	25
7. Defect and defect condition	28
8. Optimality	32
9. Weak finiteness and very distinguished categories	33
10. Zilber-Pink and extensions of algebraically closed fields	37
11. Reduction to optimal singletons	41
12. Reduction of Zilber's formulation to Pink's formulation	42
13. Zilber-Pink for a complex curve in a fibered power of the Legendre family	44
14. Zilber-Pink for a complex curve in a semiabelian variety	45
Acknowledgements	45
References	45

1. INTRODUCTION

This article is inspired by recent advances in the field of unlikely intersections, most importantly by [HP16]. Its main purpose is to introduce the general framework of distinguished categories. A distinguished category is a category with a functor to the category of algebraic varieties over an algebraically closed field of characteristic 0 that satisfies four axioms (A1) to (A4); see Section 2 for the precise definition and the list of axioms. The functor is necessary in certain examples, but can often be taken to equal the inclusion functor. Algebraic varieties are always irreducible in this article; while (mixed) Shimura

Date: December 17, 2021.

2010 *Mathematics Subject Classification.* 11G18, 14G35, 14K10, 14L10.

Key words and phrases. Unlikely intersections, Zilber-Pink, semiabelian varieties, commutative algebraic groups, mixed Shimura varieties, Legendre family, special subvarieties.

varieties as usually defined might not be irreducible, we consider here only connected (mixed) Shimura varieties, which are irreducible. We sometimes omit “connected” when speaking informally. Fields are always of characteristic 0 and subvarieties are always closed and defined over the field of definition of the ambient variety.

Equivalents or close analogues of some results in this paper were obtained independently by Cassani [Cas21a, Cas21b] in the case of connected mixed Shimura varieties (of Kuga type).

The fundamental insight that underlies this article is that many statements about unlikely intersections are highly formal in nature and that one can prove them without using any more properties of the studied objects than the ones codified in the definition of a distinguished category.

To give an idea of the power of this approach, we note that it encompasses algebraic tori, (semi-)abelian varieties, pure and mixed Shimura varieties, and even powers of the additive group as well as general connected commutative algebraic groups. Many (but not all) of these examples of distinguished categories also satisfy a fifth axiom (A5), which is needed for some of the results in this article.

Such an axiomatic approach has advantages as well as disadvantages: In the case of Shimura varieties (pure or mixed), the method allows to dispense with their very technical definition once the axioms have been verified. Another obvious advantage is that certain statements can then be proven for all examples at once. As for disadvantages, the approach is far too general to grasp the arithmetic subtleties that have so far been crucial in proving deep results on unlikely intersections (lower bounds for the size of Galois orbits, height upper bounds, etc.). Nevertheless, it provides a streamlined approach to performing certain reduction steps. Other attempts to axiomatize basic facts about unlikely intersections have been made by Zilber [Zil16], Ullmo [Ull17], and Pila [Pil21].

We have already mentioned the article [HP16] by Habegger and Pila. This did not come out of the blue but, while of great importance, was just one of the last steps after a bit less than two decades of progress on the topic. Indeed, one of the first attempts to go beyond problems of Manin-Mumford-André-Oort type arguably dates back to the end of the 90s, when Bombieri, Masser and Zannier [BMZ99] showed that a curve in \mathbb{G}_m^n has finite intersection with the union of algebraic subgroups of codimension at least 2, provided that it is defined over the algebraic numbers and not contained in a proper coset of \mathbb{G}_m^n . Precursors of this result can be found in [Sch89] and in the appendix to [Sch00] by Zannier.

In [BMZ99], the authors mentioned possible analogues of their result in the setting of abelian varieties and families thereof, and so, while Bombieri, Masser and Zannier continued their study of intersections of curves in \mathbb{G}_m^n with algebraic subgroups, other people started considering the abelian analogues. The earliest work in this direction is from 2003, due to Viada [Via03] as well as Rémond and Viada [RV03].

In the meanwhile, Zilber [Zil02] was independently led to problems of this kind from a completely different viewpoint and with different motivations coming from model theory. He formulated a general conjecture for subvarieties of arbitrary dimension of semiabelian varieties.

Finally, in [Pin05a] and [Pin05b], Pink proposed very general conjectures in the context of mixed Shimura varieties. His most general conjecture contains all of the abovementioned results about \mathbb{G}_m^n and abelian varieties. At the same time, it provides an analogue of Zilber’s conjecture in the more general context of mixed Shimura varieties. Pink’s formulation of the conjecture seems to be slightly weaker than Zilber’s, but we will show in this article that they are often equivalent. Pink’s conjecture also implies the “modular analogue” of Zilber’s conjecture, which generalizes the André-Oort conjecture in the same way as Zilber’s conjecture generalizes the Manin-Mumford conjecture.

Related conjectures for powers of the multiplicative group were formulated by Bombieri, Masser and Zannier in [BMZ07]; see the Appendix to [BMZ08] for a proof of the equivalence of the various conjectures in the case of powers of the multiplicative group.

After the groundbreaking article [PZ08] of Pila and Zannier, in which they developed and applied a new strategy relying on point counting to reprove the Manin-Mumford conjecture, it gradually became clear that this new approach would lead to relevant progress towards the aforementioned conjectures. Indeed, in the last decade the method has been applied quite successfully in basically all different contexts included in Pink's formulation.

For the present work, the article [HP16] of Habegger and Pila is of particular importance. Although their main aim was to show that \mathfrak{o} -minimality, together with functional transcendence results and Galois bounds that are still largely open as of today, is sufficient to obtain new cases of the Zilber-Pink conjecture for abelian varieties and products of modular curves, they also introduced new ideas (e.g., the defect condition) that we have exploited in [BD19] to extend their result and prove the Zilber-Pink conjecture for curves in complex abelian varieties.

Going through the proofs in [HP16] and [BD19], one realizes that many steps are purely formal and here we propose a categorical setting in which basic objects (e.g., special and weakly special subvarieties) can be defined and facts about them (e.g., the defect condition) can be proved formally.

We show that basically all the abovementioned examples, prominently (connected) mixed Shimura varieties, satisfy our axioms. On the other hand, unlikely intersections have also been studied in various other settings and for some of these, it is unclear whether they could be integrated into the framework of distinguished categories: We mention here (families of) Drinfeld modules (see [GH13]), the results for affine space over function fields in [CGMM13], Klingler's mixed Hodge varieties in [Kli17], and Gao's enlarged mixed Shimura varieties in [Gao19] as well as the growing field of unlikely intersections in arithmetic dynamics (see the survey [BLJ⁺19]). We further refer to Section 3 of [GT21] for various examples that show the difficulty of defining special subvarieties in the context of arithmetic dynamics. While a common method has been applied to studying unlikely intersections in our examples of distinguished categories, namely using the Pila-Zannier strategy to combine \mathfrak{o} -minimal point counting, Galois bounds, and functional transcendence results of Ax-Schanuel type (see [Zan12]), the methods used in these other settings are sometimes very different.

In Section 2, we define our main object of study, distinguished categories, and we see that many of the abovementioned examples form distinguished categories.

We devote Sections 3, 4, and 5 to a detailed proof that the category of connected (mixed) Shimura varieties (of Kuga type) is distinguished. Unfortunately, slightly different definitions of connected mixed Shimura varieties appear in the literature. For this reason and since we have not been able to locate them in the literature, we decided to include proofs of some widely used and well-known facts about connected mixed Shimura varieties, for instance the fact that the irreducible components of an intersection of special subvarieties are special. We hope that this will make our text more accessible to readers who are not very familiar with this topic.

Many concepts from the field of unlikely intersections can be defined naturally in an arbitrary distinguished category: In Section 6, we introduce the concept of (weakly) special subvarieties. Formally, our definitions are identical with Pink's in [Pin05a] for the special case of the distinguished category of connected mixed Shimura varieties and they are equivalent to the usual definitions of (weakly) special subvarieties in the case of semiabelian varieties (see Remark 6.4). In the same section, we prove basic facts about (weakly) special subvarieties (e.g., special subvarieties are weakly special, irreducible components of intersections of weakly special subvarieties are weakly special) that are well-known for commutative algebraic groups and for mixed Shimura varieties, although in the latter case

we have not always been able to locate proofs in the literature. We also compare our approach with Ullmo's in [Ull17].

The so-called defect condition, introduced in [HP16] and conjectured there to hold in any mixed Shimura variety, is often a useful technical tool when studying unlikely intersections: In Section 7, we introduce the concept of (weak) defect and prove the defect condition in an arbitrary distinguished category (special cases were proven in [HP16], [DR18], [PT19], and [Cas21a]). The definition of defect is formally identical with the definition in [Pin05b].

In Section 8, we then introduce the concept of (weak) optimality in an arbitrary distinguished category. The definition of optimality goes back to [HP16] while weak optimality for subvarieties of \mathbb{G}_m^n already appears in [Poi01] under the name of cd-maximality.

In Section 9, we formulate the axiom (A5), which informally speaking says that weakly optimal subvarieties for a given subvariety come in finitely many "families". Given an extension $K \subset L$ of algebraically closed fields and a distinguished category over K , we can perform a base change to get a distinguished category over L . If (A5) is satisfied in any base change of the original category with respect to an extension of algebraically closed fields of finite transcendence degree, we call the category very distinguished. In our main examples of very distinguished categories (abelian varieties, tori, semiabelian varieties, connected Shimura varieties, and connected mixed Shimura varieties of Kuga type over algebraically closed fields of finite transcendence degree over \mathbb{Q}), the fact that they are very distinguished is usually deduced from a functional transcendence result of weak Ax-Schanuel type (called "Weak Complex Ax" in [HP16]) together with the facts that the corresponding uniformization map is definable in an o-minimal structure and that a countable definable set is finite.

In Section 10, we formulate a statement for arbitrary distinguished categories that corresponds to the Zilber-Pink conjecture. We then show in Theorem 10.7 that the Zilber-Pink statement for the base change of a very distinguished category with respect to an extension of algebraically closed fields follows from the same statement for the original category.

This reduction of the transcendence degree of the ground field in the Zilber-Pink statement is similar to the main result of [BD19], where we reduced the Zilber-Pink conjecture for abelian varieties from \mathbb{C} to $\bar{\mathbb{Q}}$. Note however that we do not recover the main result of [BD19]: If the ambient abelian variety A cannot be defined over $\bar{\mathbb{Q}}$, then the method we present here does not reduce the Zilber-Pink conjecture for A to the Zilber-Pink conjecture for an abelian variety over $\bar{\mathbb{Q}}$. Our method however allows to reduce to the case where the subvariety is defined over the algebraic closure of the field of definition of the abelian variety. The analogous statement for powers of the multiplicative group has been proven by Bombieri, Masser and Zannier in [BMZ08]. Our proof of Theorem 10.7 takes its inspiration from the proof of Proposition 4.1 in [BD19].

Recently, in [PS21], Pila and Scanlon proved an effective Zilber-Pink result over a transcendental field extension of \mathbb{C} as well as a reduction of the transcendental to the algebraic case in this context, considering only varieties associated to certain quotient spaces, transcendental points, and strongly special subvarieties. In their reduction step, they have to assume a formulation of the conjecture over the ground field that is a priori slightly stronger than what they obtain over the field extension (but we will prove in this paper that the two formulations are often equivalent). It seems likely that one could remove this restriction, even without using our work, and that their method could yield this reduction step in a very general context. However, the dimension of the variety that they need to apply the conjecture to over the ground field can be bigger than the dimension of the variety they started with. This is a fundamental difference to our method, which yields more refined results.

Theorem 10.7 also contains another statement that implies for example that the Zilber-Pink conjecture holds for any complex curve that cannot be defined over $\bar{\mathbb{Q}}$ inside the

moduli space \mathcal{A}_2 of principally polarized abelian surfaces over $\bar{\mathbb{Q}}$. We denote the base change of a K -variety V with respect to a field extension $K \subset L$ by V_L (see Definition 2.3).

The following theorem is a special case of Corollary 10.9.

Theorem 1.1. *Let K be an algebraically closed field. Let C be a curve in $(\mathcal{A}_2)_K$ that is not the base change of a curve in \mathcal{A}_2 (so cannot be defined over $\bar{\mathbb{Q}}$) and is not contained in a proper special subvariety of $(\mathcal{A}_2)_K$. Then C contains at most finitely many points that lie on a special subvariety of $(\mathcal{A}_2)_K$ of codimension at least 2.*

So far, only partial results for curves that are defined over $\bar{\mathbb{Q}}$ inside \mathcal{A}_2 are known (see [DO19a] and [DO19b]). Our proof does not use any information about \mathcal{A}_2 apart from the fact that it is a connected Shimura variety of dimension 3 over $\bar{\mathbb{Q}}$ for which the André-Oort conjecture is known to hold. Thus, we also recover Pila's Theorem 1.4 in [Pil17], which says that the Zilber-Pink conjecture holds for any complex curve that cannot be defined over $\bar{\mathbb{Q}}$ inside the cube of the moduli space of elliptic curves.

Let us now see an example. We consider the hyperelliptic curve of genus two defined by the equation

$$y^2 = x(x-1)(x-\pi t)(x-t^2)(x-t^4)$$

over the field $K(t)$, where $K = \overline{\mathbb{Q}(\pi)} \subset \mathbb{C}$ and t is an indeterminate. Let J_t be its Jacobian. This gives a curve C in $(\mathcal{A}_2)_K$ that cannot be defined over $\bar{\mathbb{Q}}$. There exists a finite set F such that, for each $t_0 \in \mathbb{C} \setminus F$, there is a specialized hyperelliptic curve and a specialized Jacobian J_{t_0} over \mathbb{C} .

By the Theorem in [Mas99], specializing π to 1, we know that J_t only has trivial endomorphisms, even over an algebraic closure of $K(t)$. This implies (see Table 1 on p. 11 of [DO19a]) that C is not contained in any proper special subvariety of $(\mathcal{A}_2)_K$. Theorem 1.1 then implies that the intersection of C with the union of all special subvarieties of $(\mathcal{A}_2)_K$ that have codimension at least 2 is finite. This in turn implies that there are at most finitely many complex numbers t_0 such that at least one of the following holds:

- (1) J_{t_0} is simple and its endomorphism ring is a \mathbb{Z} -module of rank four (i.e., J_{t_0} has quaternionic or complex multiplication),
- (2) J_{t_0} is isogenous to the square of an elliptic curve, or
- (3) J_{t_0} is isogenous to the product of an elliptic curve and a CM elliptic curve.

In Sections 11 and 12, we prove (again assuming (A5) most of the time) that the Zilber-Pink statement is equivalent to several seemingly weaker statements. In particular, while the statement is concerned with optimal subvarieties of arbitrary dimension, we show in Theorem 11.1 that it suffices to prove it for optimal singletons. This generalizes Theorem 8.3 in [DR18] (for Shimura varieties) and Theorem 6.1 in [BD19] (for abelian varieties); the latter theorem is an easy consequence of the results in [HP16].

Furthermore, the Zilber-Pink statement we commonly use in this article generalizes Zilber's formulation of the conjecture. However, we show in Theorem 12.4 that it is equivalent to the corresponding generalization of Pink's formulation of the conjecture. This generalizes Theorem 1.9 in [BD19], where we have shown (the non-trivial direction of) this equivalence in the case of abelian varieties. In the toric case, it had already earlier been pointed out by Ullmo and Zannier that the equivalence of the two formulations can be proven by imposing additional multiplicative relations on the positive-dimensional atypical intersections.

In both Theorem 11.1 and Theorem 12.4, we have to assume that (A5) is satisfied. Similar statements for connected mixed Shimura varieties of Kuga type have been proven by Cassani in [Cas21b].

In Section 13, we apply our results to prove that the Zilber-Pink conjecture holds for a complex curve in a fibered power of the Legendre family of elliptic curves, which is a

connected mixed Shimura variety of Kuga type. This was previously known only if the curve can be defined over $\bar{\mathbb{Q}}$, in which case it follows from combining the results of [RV03], [Via08], [Gal10], [BC16], and [Bar19]. We use these results together with our Theorem 10.7 and the fact that connected mixed Shimura varieties of Kuga type over $\bar{\mathbb{Q}}$ form a very distinguished category to deduce the \mathbb{C} -case.

The theorem below is a consequence of Theorem 13.1.

Theorem 1.2. *Let K be an algebraically closed field and let \mathcal{E}^g be the g -th fibered power of the Legendre family of elliptic curves over $\bar{\mathbb{Q}}$. Let C be a curve in \mathcal{E}_K^g that is not contained in a proper special subvariety of \mathcal{E}_K^g . Then C contains at most finitely many points that lie on a special subvariety of \mathcal{E}_K^g of codimension at least 2.*

In Section 14, we show that the Zilber-Pink conjecture holds for a complex curve in a semiabelian variety, provided that the semiabelian variety can be defined over $\bar{\mathbb{Q}}$.

Theorem 1.3. *Let K be an algebraically closed field and let G be a semiabelian variety over $\bar{\mathbb{Q}}$. Let C be a curve in G_K that is not contained in a proper algebraic subgroup of G_K . Then C contains at most finitely many points that lie on an algebraic subgroup of G_K of codimension at least 2.*

This follows from combining our Theorem 10.7 and the fact that semiabelian varieties over $\bar{\mathbb{Q}}$ form a very distinguished category with the recent work [BKS21] of the first-named author, Kühne, and Schmidt. See Theorem 14.1.

2. DISTINGUISHED CATEGORIES

Let K be an algebraically closed field of characteristic 0. Consider a category \mathfrak{C} with objects \mathfrak{V} and morphisms \mathfrak{M} together with a covariant functor \mathcal{F} from \mathfrak{C} to the category of varieties over K . Typically, this functor will correspond to forgetting some additional information.

We formulate the following axioms that this category may or may not satisfy:

- (A1) *Direct Products* - If $X, Y \in \mathfrak{V}$, then there exists $Z \in \mathfrak{V}$ and there exist morphisms $\pi_X : Z \rightarrow X$, $\pi_Y : Z \rightarrow Y$ in \mathfrak{M} such that the morphism $\mathcal{F}(Z) \rightarrow \mathcal{F}(X) \times_K \mathcal{F}(Y)$ induced by the morphisms $\mathcal{F}(\pi_X)$ and $\mathcal{F}(\pi_Y)$ is an isomorphism. We identify $\mathcal{F}(Z)$ with $\mathcal{F}(X) \times_K \mathcal{F}(Y)$. Furthermore, if $\phi : W \rightarrow X$ and $\psi : W \rightarrow Y$ are morphisms in \mathfrak{M} , then there exists a morphism $\chi : W \rightarrow Z$ in \mathfrak{M} such that $\pi_X \circ \chi = \phi$ and $\pi_Y \circ \chi = \psi$. We write $X \times Y$ or $X \times_K Y$ for Z and (ϕ, ψ) for χ .
- (A2) *Fibered Products* - If $\phi : X \rightarrow Z$ and $\psi : Y \rightarrow Z$ are morphisms in \mathfrak{M} , then there exists $n \in \mathbb{Z}$, $n \geq 0$, and there exist $X_1, \dots, X_n \in \mathfrak{V}$ and morphisms $\phi_i : X_i \rightarrow X \times Y$ in \mathfrak{M} ($i = 1, \dots, n$) such that $\bigcup_{i=1}^n \mathcal{F}(\phi_i)(\mathcal{F}(X_i)) = (\mathcal{F}(X) \times_{\mathcal{F}(Z)} \mathcal{F}(Y))_{\text{red}} \subset \mathcal{F}(X) \times_K \mathcal{F}(Y)$.
- (A3) *Final Object* - The category has a final object that is mapped to $\text{Spec } K$ by \mathcal{F} .
- (A4) *Fiber Dimension* - If $\phi : X \rightarrow Y$ is a morphism in \mathfrak{M} , then $\mathcal{F}(\phi)(\mathcal{F}(X))$ is closed in $\mathcal{F}(Y)$ and there exist morphisms $\phi_1 : W \rightarrow X$, $\phi_2 : W \rightarrow Z$, and $\phi_3 : Z \rightarrow Y$ in \mathfrak{M} such that $\phi \circ \phi_1 = \phi_3 \circ \phi_2$, $\mathcal{F}(\phi_1)$ is finite and surjective, $\mathcal{F}(\phi_3)$ has finite fibers, $\mathcal{F}(\phi_2)^{-1}(z)$ is irreducible for all $z \in \mathcal{F}(\phi_2)(\mathcal{F}(W))$, and $\dim_w \mathcal{F}(\phi_2)^{-1}(\mathcal{F}(\phi_2)(w))$ is a constant function for $w \in \mathcal{F}(W)$.

In other words, (A4) says that the image of $\mathcal{F}(\phi)$ is closed and ϕ factorizes as in the following diagram:

$$\begin{array}{ccc}
 & \text{non-empty fibers are irreducible} & \\
 & \text{of constant dimension} & \\
 W & \xrightarrow{\phi_2} & Z \\
 \phi_1 \downarrow \begin{array}{l} \text{finite,} \\ \text{surjective} \end{array} & & \phi_3 \downarrow \text{finite fibers} \\
 X & \xrightarrow{\phi} & Y,
 \end{array} \tag{2.1}$$

where the properties in the diagram are properties of the images of the morphisms under \mathcal{F} . Note that (A4) implies that $\dim_x \mathcal{F}(\phi)^{-1}(\mathcal{F}(\phi)(x))$ is a constant function for $x \in (\mathcal{F}(X))(K)$ and therefore for $x \in \mathcal{F}(X)$ (see Appendix E of [GW10]). By the same appendix, it suffices to show that all non-empty fibers of $\mathcal{F}(\phi_2)$ over closed points of $\mathcal{F}(Z)$ are irreducible of constant dimension in order to verify (A4).

Definition 2.1. *If \mathfrak{C} satisfies (A1) to (A4), we call it a distinguished category. The elements of \mathfrak{V} (which we will often identify with their images under \mathcal{F}) will then be called distinguished varieties while the elements of \mathfrak{M} (which we will often similarly identify) will be called distinguished morphisms.*

The domain and codomain of a distinguished morphism are automatically distinguished varieties.

Remark 2.2. We are going to see later that these axioms are all necessary in order to define fundamental concepts and prove basic facts. For the reader who is already familiar with the topic, we summarize here how they will be used:

- (A1) is needed to introduce (A2).
- (A2) is equivalent to the fact that irreducible components of intersections of special subvarieties are special.
- (A3) implies that special subvarieties are weakly special.
- (A4) together with (A2) implies that irreducible components of (pre-)images of (weakly) special subvarieties under distinguished morphisms are (weakly) special and irreducible components of intersections of weakly special subvarieties are weakly special. That the image is closed is necessary in order to be able to define special subvarieties at all. The constant fiber dimension is fundamental in the proof of the defect condition.

We are going to use the following notation throughout the article.

Definition 2.3. *If $K \subset L$ is an arbitrary field extension and V is a finite union of varieties over K , then $V_L = V \times_K L$ is called the base change of V to L . We use analogous notation for the base change of morphisms between varieties.*

We now give some examples of distinguished categories. If \mathcal{F} is not mentioned, then it is to be understood that \mathfrak{C} is a subcategory of the category of varieties over K and \mathcal{F} is just the inclusion functor.

A trivial but instructive example is the following: Let X be an arbitrary fixed variety over K . The category $\mathfrak{C}_{\text{triv}}(X)$ with objects $\mathfrak{V} = \{X^n; n \in \mathbb{Z}, n \geq 0\}$ and morphisms $\mathfrak{M} = \{\text{maps } X^m \rightarrow X^n, (x_1, \dots, x_m) \mapsto (x_{i_1}, \dots, x_{i_n})\}$ is distinguished.

The category of connected commutative algebraic groups gives an interesting example of a distinguished category $\mathfrak{C}_{\text{comm}}$. To be more precise, if we take \mathfrak{V} to be the connected commutative algebraic groups over K , \mathfrak{M} to consist of homomorphisms of algebraic groups composed with translations by torsion points, and \mathcal{F} to be the forgetful functor, we can easily see that axioms (A1) to (A4) follow from well-known properties of algebraic groups and homomorphisms between them.

In particular, also in view of [BD19], one can consider the full subcategories \mathfrak{C}_{ab} and $\mathfrak{C}_{\text{semiab}}$ of $\mathfrak{C}_{\text{comm}}$, consisting of (semi-)abelian varieties over K , or even just $\mathfrak{C}_{\text{tor}}$ given by $\mathfrak{V} = \{\mathbb{G}_{m,K}^n; n \in \mathbb{Z}, n \geq 0\}$ or $\mathfrak{C}_{\text{add}}$ given by $\mathfrak{V} = \{\mathbb{G}_{a,K}^n; n \in \mathbb{Z}, n \geq 0\}$.

In the next section we are going to see further examples of distinguished categories. We end this section by noting that given two distinguished categories \mathfrak{C} and \mathfrak{C}' over an algebraically closed field K , we can form a product category $\mathfrak{C} \times \mathfrak{C}'$ that is again distinguished: Its objects are pairs of objects of \mathfrak{C} and \mathfrak{C}' , its morphisms pairs of morphisms of \mathfrak{C} and \mathfrak{C}' , and its functor is defined by applying to each element of the pair of objects or morphisms the functor of the corresponding distinguished category and taking the direct

product of the two resulting varieties or the morphism between direct products induced by the two resulting morphisms of varieties respectively.

3. CONNECTED MIXED SHIMURA VARIETIES

In this section together with the next two ones, we are going to see that connected mixed Shimura varieties and the subcategories consisting of the pure and the Kuga type ones fit in our framework and satisfy axioms (A1) to (A4).

Different definitions of a (connected) mixed Shimura datum appear in the literature. We use the following. The Deligne torus $\mathrm{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m,\mathbb{C}}$ is denoted by \mathbb{S} , we identify $\mathbb{S}(\mathbb{R})$ with \mathbb{C}^* . We fix once and for all a complex square root of -1 that we denote by $\sqrt{-1}$. For the definition of a rational mixed Hodge structure, its type, and its weight filtration, we refer to Chapter 1 of [Pin90]. The definition of a Cartan involution can be found on pp. 274–275 of [Mil05].

Definition 3.1. *A connected mixed Shimura datum is a pair (P, X^+) , where P is a connected linear algebraic group over \mathbb{Q} with unipotent radical W and another algebraic subgroup $U \subset W$ that is normal in P and uniquely determined by X^+ and condition (MSD.c) below, and $X^+ \subset \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$ is a connected component of an orbit under conjugation (from the left) by the subgroup $P(\mathbb{R})U(\mathbb{C}) \subset P(\mathbb{C})$ such that for some (or equivalently for all) $x \in X^+$ we have that*

(MSD.a) *the composite homomorphism $\mathbb{S}_{\mathbb{C}} \xrightarrow{x} P_{\mathbb{C}} \rightarrow (P/U)_{\mathbb{C}}$ is defined over \mathbb{R} ,*

(MSD.b) *the adjoint representation induces on $\mathrm{Lie} P$ a rational mixed Hodge structure of type*

$$\{(-1, 1), (0, 0), (1, -1), (-1, 0), (0, -1), (-1, -1)\},$$

(MSD.c) *the weight filtration on $\mathrm{Lie} P$ is given by*

$$W_n(\mathrm{Lie} P) = \begin{cases} \{0\} & \text{if } n < -2, \\ \mathrm{Lie} U & \text{if } n = -2, \\ \mathrm{Lie} W & \text{if } n = -1, \\ \mathrm{Lie} P & \text{if } n \geq 0, \end{cases}$$

(MSD.d) *the conjugation by $x(\sqrt{-1})$ induces a Cartan involution on $(P/W)_{\mathbb{R}}^{\mathrm{ad}}$ and $(P/W)^{\mathrm{ad}}$ possesses no non-trivial \mathbb{Q} -factor H such that $H(\mathbb{R})$ is compact, and*

(MSD.e) *P/P^{der} is an almost-direct product of a \mathbb{Q} -split torus with a torus of compact type defined over \mathbb{Q} .*

If $U = \{1\}$, then (P, X^+) is called connected mixed Shimura datum of Kuga type, while if $W = \{1\}$ it is called connected pure Shimura datum or connected Shimura datum.

In Definition 2.4(1) in [Gao17b], X^+ is taken to be a $P(\mathbb{R})^+U(\mathbb{C})$ -orbit instead of a connected component of a $P(\mathbb{R})U(\mathbb{C})$ -orbit, where $P(\mathbb{R})^+$ denotes the identity component of $P(\mathbb{R})$. The two definitions are equivalent. Definition 2.1 in [Pin05a] is more restrictive than our definition here as Pink additionally demands that P possesses no proper normal subgroup P' , defined over \mathbb{Q} , such that x factors through $P'_{\mathbb{C}} \subset P_{\mathbb{C}}$. Note that the second part of (MSD.d) seems to be missing in Definition 2.1 in [Pin05a], but follows from Pink's additional condition (vi) together with the rest. Definition 2.1 in [Gao20] is equal to Definition 2.1 in [Pin05a] in the case of connected mixed Shimura data of Kuga type. Definition 2.1 in [Pin90] is a slightly more general analogue of our definition for mixed Shimura data instead of connected mixed Shimura data.

We now let $P(\mathbb{R})_+$ be the stabilizer of X^+ in $P(\mathbb{R})$ (it contains $P(\mathbb{R})^+$), and we let Γ be a congruence subgroup of $P(\mathbb{Q})$ that is contained in $P(\mathbb{Q})_+ := P(\mathbb{Q}) \cap P(\mathbb{R})_+$. By abuse of notation, we will say in this situation that Γ is a congruence subgroup of $P(\mathbb{Q})_+$. By Proposition 2.2 in [RS93], there always exists some such congruence subgroup. It follows

that for every congruence subgroup of $P(\mathbb{Q})$, its intersection with $P(\mathbb{Q})_+$ is a congruence subgroup of $P(\mathbb{Q})$ that is contained in $P(\mathbb{Q})_+$.

By [Pin90], 1.18(a), there is a canonical complex structure on X^+ . There exists an algebraic variety V over $\bar{\mathbb{Q}}$ such that $\Gamma \backslash X^+$ is canonically isomorphic to the analytification of $V_{\mathbb{C}}$ as a complex analytic space (see [Pin90], Proposition 3.3, Proposition 9.24, Definition 9.25, and Theorem 11.18).

The variety V is called a *connected mixed Shimura variety*. If (P, X^+) is of Kuga type or pure, then V is called *connected mixed Shimura variety of Kuga type* or *connected (pure) Shimura variety* respectively.

Condition (MSD.e) is usually dropped in the definition of a connected (pure) Shimura datum/variety. Nevertheless, in this article, connected (pure) Shimura data will satisfy (MSD.e). This does not cause any problems thanks to Remarks 3.5 and 4.4. If (P, X^+) satisfies (MSD.a) to (MSD.d) with $W = \{1\}$, then we can consider its quotient by the center of P (see Remark 3.5) and this quotient will satisfy (MSD.e). So (MSD.e) is not a big restriction in applications to unlikely intersections.

Definition 3.2. *Let (P, X^+) and (P', X'^+) be connected mixed Shimura data. If we have a homomorphism $\phi : P \rightarrow P'$ of algebraic groups over \mathbb{Q} which induces a map $X^+ \rightarrow X'^+$ through $x \mapsto \phi_{\mathbb{C}} \circ x$, we say that ϕ induces a Shimura morphism $(P, X^+) \rightarrow (P', X'^+)$ of connected mixed Shimura data.*

Suppose we are given a Shimura morphism $(P, X^+) \rightarrow (P', X'^+)$ induced by $\phi : P \rightarrow P'$ and congruence subgroups $\Gamma \subset P(\mathbb{Q})_+$ and $\Gamma' \subset P'(\mathbb{Q})_+$ with $\phi(\Gamma) \subset \Gamma'$. Then, this induces a holomorphic map $\Gamma \backslash X^+ \rightarrow \Gamma' \backslash X'^+$ between complex analytic spaces that in turn gives a morphism between the corresponding connected mixed Shimura varieties (see [Pin90], 3.4, Proposition 9.24, Proposition 11.10, and Theorem 11.18).

We are now in place to define a distinguished category $\mathfrak{C}_{\text{mSv}}$ over $\bar{\mathbb{Q}}$ where

- $\mathfrak{D}_{\text{mSv}}$ consists of triples (P, X^+, Γ) where (P, X^+) is a connected mixed Shimura datum and Γ is a congruence subgroup of $P(\mathbb{Q})_+$,
- $\mathfrak{M}_{\text{mSv}}$ consists of arrows $(P, X^+, \Gamma) \rightarrow (P', X'^+, \Gamma')$ given by a Shimura morphism $(P, X^+) \rightarrow (P', X'^+)$ induced by a homomorphism of algebraic groups $\phi : P \rightarrow P'$ that satisfies $\phi(\Gamma) \subset \Gamma'$ and composition of arrows is given by composition of Shimura morphisms, and
- the functor \mathcal{F}_{mSv} sends an element (P, X^+, Γ) of $\mathfrak{D}_{\text{mSv}}$ to the corresponding connected mixed Shimura variety V over $\bar{\mathbb{Q}}$. Moreover, it sends an element $(P, X^+, \Gamma) \rightarrow (P', X'^+, \Gamma')$ of $\mathfrak{M}_{\text{mSv}}$ to the corresponding morphism of connected mixed Shimura varieties.

By restricting the above definition to connected mixed Shimura data of Kuga type or connected pure Shimura data we can also define the full subcategories $\mathfrak{C}_{\text{mSvK}}$ and $\mathfrak{C}_{\text{pSv}}$ of $\mathfrak{C}_{\text{mSv}}$. By abuse of notation, we also call the elements of $\mathfrak{M}_{\text{mSv}}$, $\mathfrak{M}_{\text{mSvK}}$, and $\mathfrak{M}_{\text{pSv}}$ as well as their images under the respective functor Shimura morphisms.

We would like to remark that we cannot just take $\mathfrak{C}_{\text{mSv}}$ to be a subcategory of the category of varieties over $\bar{\mathbb{Q}}$ and take for \mathcal{F}_{mSv} the inclusion functor because of the following: There is a trivial connected Shimura datum that is associated to the trivial reductive group and admits Shimura morphisms from any other connected Shimura datum. Its associated connected Shimura variety consists only of a point. There are however also non-trivial connected Shimura data that are associated to certain (positive-dimensional) tori and do not admit Shimura morphisms from all other connected Shimura data, and in particular not from the trivial connected Shimura datum, but whose associated connected Shimura varieties nevertheless consist only of a point.

On the other hand, these non-trivial connected Shimura data can sometimes also admit Shimura morphisms to connected Shimura data with positive-dimensional associated

connected Shimura varieties (leading to special points on these), which the trivial connected Shimura datum cannot. This means that the existence or non-existence of Shimura morphisms between connected Shimura varieties depends not only on the variety structure, but also on the associated connected Shimura datum and congruence subgroup (and non-isomorphic pairs of connected Shimura data and congruence subgroups might yield connected Shimura varieties that are isomorphic as varieties). All of this holds also for connected mixed Shimura varieties or connected mixed Shimura varieties of Kuga type instead of connected Shimura varieties.

In the next two sections we are going to prove the following proposition:

Proposition 3.3. *The categories $\mathfrak{C}_{\text{mSv}}$, $\mathfrak{C}_{\text{mSvK}}$, and $\mathfrak{C}_{\text{pSv}}$ satisfy the axioms (A1) to (A4).*

Proposition 3.3 will allow us to forget the very technical definition of a connected (mixed) Shimura variety (of Kuga type) for a large part of this article and work only with the axioms (A1) to (A4).

We conclude this section with some definitions and a proposition concerning Shimura morphisms.

Definition 3.4. *Let ϕ be a Shimura morphism $(P, X^+, \Gamma) \rightarrow (Q, Y^+, \Delta)$. By abuse of notation we also denote the corresponding homomorphism of algebraic groups $P \rightarrow Q$ by ϕ . We say that $\mathcal{F}_{\text{mSv}}(\phi) : \mathcal{F}_{\text{mSv}}(P, X^+, \Gamma) \rightarrow \mathcal{F}_{\text{mSv}}(Q, Y^+, \Delta)$ is a*

- (1) Shimura submersion if the image of ϕ contains Q^{der} ,
- (2) quotient Shimura morphism if ϕ is surjective and in this case we call (Q, Y^+) a quotient Shimura datum of (P, X^+) ,
- (3) Shimura immersion if the identity component of $\ker \phi$ is a torus,
- (4) Shimura embedding if ϕ is injective and in this case we call (P, X^+) a Shimura subdatum of (Q, Y^+) , and
- (5) Shimura covering if it is a Shimura submersion and immersion.

These definitions concern morphisms between connected mixed Shimura varieties but actually depend on the underlying homomorphisms of algebraic groups (and only on these). We then also call Shimura submersions, quotient Shimura morphisms, Shimura immersions, embeddings, and coverings the corresponding Shimura morphisms between connected mixed Shimura data as well as the corresponding elements of $\mathfrak{M}_{\text{mSv}}$.

Remark 3.5. A remark about quotient Shimura data and morphisms is in order: In Proposition 2.9 in [Pin90], the quotient of a mixed Shimura datum (P, X) by a normal algebraic subgroup K of P is constructed; it satisfies a certain natural universal property. Thanks to Corollary 2.12 in [Pin90], this construction also yields quotients $(P, X^+)/K$ of connected mixed Shimura data (P, X^+) by such subgroups K that are again connected mixed Shimura data. One can check that the abovementioned slight difference between our definition and Pink's definition in [Pin90] is preserved under taking such quotients. Furthermore, one can check that a Shimura morphism $(P, X^+) \rightarrow (Q, Y^+)$ is a quotient Shimura morphism if and only if it induces an isomorphism between $(P, X^+)/K$ and (Q, Y^+) , where $K = \ker(P \rightarrow Q)$.

We recall some results from [Pin05a] that we will use when verifying that (A4) holds for our three categories $\mathfrak{C}_{\text{mSv}}$, $\mathfrak{C}_{\text{mSvK}}$, and $\mathfrak{C}_{\text{pSv}}$ and in Section 9.

Proposition 3.6 ([Pin05a], Facts 2.6). *The following hold:*

- (1) Shimura submersions are surjective.
- (2) Shimura immersions are finite.

As no proof of these facts is provided in [Pin05a] and the definition of a connected mixed Shimura datum in [Pin05a] is slightly different from our definition here, we give a proof. It relies on the proof of the proposition in the pure case that appears in [Orr13].

Proof. Let $\phi : P \rightarrow Q$ be a homomorphism of connected linear algebraic groups over \mathbb{Q} that induces a Shimura morphism of connected mixed Shimura data $(P, X^+) \rightarrow (Q, Y^+)$ and thereby induces a Shimura morphism $\mathcal{F}_{\text{mSv}}(P, X^+, \Gamma) \rightarrow \mathcal{F}_{\text{mSv}}(Q, Y^+, \Delta)$. We first note that surjectivity and finiteness may be checked after base change to \mathbb{C} . This is guaranteed by Propositions 14.48 and 14.51 in [GW10] (the morphism $\text{Spec } \mathbb{C} \rightarrow \text{Spec } \mathbb{Q}$ is faithfully flat and quasi-compact, see [GW10], Definition 10.1 and right after Definition 14.7). Surjectivity over \mathbb{C} may then be checked on \mathbb{C} -points thanks to Theorem 10.70 in [GW10].

We start with (1). Suppose the Shimura morphism $\mathcal{F}_{\text{mSv}}(P, X^+, \Gamma) \rightarrow \mathcal{F}_{\text{mSv}}(Q, Y^+, \Delta)$ induced by ϕ is a Shimura submersion. By Proposition 2.9 in [Gao17b], ϕ induces a Shimura morphism of connected pure Shimura data

$$(G_P, X_G^+) := (P, X^+)/W_P \rightarrow (Q, Y^+)/W_Q =: (G_Q, Y_G^+),$$

where W_P and W_Q are the unipotent radicals of P and Q respectively and $(P, X^+)/W_P$ and $(Q, Y^+)/W_Q$ are the induced quotient Shimura data. This Shimura morphism is also a Shimura submersion. Let $U_Q \subset W_Q$ and $U_P \subset W_P$ be the subgroups from the definition of a connected mixed Shimura datum. By Proposition 2.9 in [Gao17b], we have $\phi(U_P) \subset U_Q$.

By (MSD.e) we have that the quotient Q/Q^{der} is of multiplicative type. Moreover, the image of W_Q in such a quotient is unipotent and therefore trivial by [Mil17], Proposition 14.16. Therefore, we have $W_Q \subset Q^{\text{der}} \subset \phi(P)$.

Now we claim that $\phi|_{U_P} : U_P \rightarrow U_Q$ is surjective. If we take this for granted, then it follows from Proposition 5.1 in [Mil05] that $W_Q(\mathbb{R})U_Q(\mathbb{C}) \subset \phi(P(\mathbb{R})_+U_P(\mathbb{C}))$. Moreover, by the proof of Theorem 2.4(3) in [Orr13], the map $X_G^+ \rightarrow Y_G^+$ is surjective and $X^+ \rightarrow X_G^+$ is surjective by Proposition 5.1 in [Mil05]. Then [Pin90], 2.18, implies that the map $X^+ \rightarrow Y^+$ is surjective and therefore the induced map $\Gamma \backslash X^+ \rightarrow \Delta \backslash Y^+$ is surjective as well, from which (1) follows.

We are left with proving that $\phi|_{U_P} : U_P \rightarrow U_Q$ is surjective. We argue at the level of Lie algebras. The torus $\mathbb{S}_{\mathbb{C}}$ acts on $\text{Lie } P$ through the adjoint representation. We have $\mathbb{S}_{\mathbb{C}}$ -invariant subspaces $\text{Lie } \ker \phi \subset \text{Lie } \phi^{-1}(U_Q)$ and this gives an $\mathbb{S}_{\mathbb{C}}$ -invariant subspace $\text{Lie } \phi^{-1}(U_Q)/\text{Lie } \ker \phi$ of $\text{Lie } P/\text{Lie } \ker \phi$. Now $\text{Lie } \phi^{-1}(U_Q)/\text{Lie } \ker \phi$ is isomorphic to $\text{Lie}(U_Q \cap \phi(P))$ by Proposition 1.63 in [Mil17], so the action of $\mathbb{S}_{\mathbb{C}}$ on it is pure of type $\{(-1, -1)\}$ (as defined in [Pin90], 1.3, p. 9). Hence, $\text{Lie } \phi^{-1}(U_Q)/\text{Lie } \ker \phi$ must be contained in the $(-1, -1)$ -subspace of $\text{Lie } P/\text{Lie } \ker \phi$, which is equal to $\text{Lie}(\ker \phi)U_P/\text{Lie } \ker \phi$. It follows that $\text{Lie } \phi^{-1}(U_Q)$ is contained in $\text{Lie}(\ker \phi)U_P$. For the identity component $\phi^{-1}(U_Q)^0$, we have

$$\text{Lie}(\phi^{-1}(U_Q)^0 \cap (\ker \phi)U_P) = \text{Lie } \phi^{-1}(U_Q)^0 \cap \text{Lie}(\ker \phi)U_P = \text{Lie } \phi^{-1}(U_Q)^0,$$

see [Mil17], Chapter 10, p. 190. By [Mil17], Proposition 10.15, $\phi^{-1}(U_Q)^0$ is contained in $(\ker \phi)U_P$. But $\phi(\phi^{-1}(U_Q))$ is connected (being unipotent), so $\phi(\phi^{-1}(U_Q)) = \phi(\phi^{-1}(U_Q)^0)$ and hence $\phi^{-1}(U_Q) = (\ker \phi)\phi^{-1}(U_Q)^0 \subset (\ker \phi)U_P$. Since $U_Q \subset W_Q \subset \phi(P)$, this implies that $\phi|_{U_P} : U_P \rightarrow U_Q$ is surjective and we are done with (1).

We turn to (2), using the same notation as in (1), but now ϕ induces a Shimura immersion, i.e., the identity component of $\ker \phi$ is a torus. We want to show that the map $\Gamma \backslash X^+ \rightarrow \Delta \backslash Y^+$ of complex analytic spaces is finite, i.e., proper and separated with finite fibers. Note that separatedness is automatic. By GAGA (see [SGA03], Exposé XII, Proposition 3.2(vi)), this implies that the corresponding morphism of algebraic varieties over \mathbb{C} is also finite.

By Proposition 2.9 in [Gao17b], we have $\phi(W_P) \subset W_Q$. The kernel of $\phi|_{W_P}$ is a unipotent algebraic group whose identity component is a torus. Hence it is trivial, so $\phi|_{W_P}$ is injective. Again, there is an induced Shimura morphism $(G_P, X_G^+) \rightarrow (G_Q, Y_G^+)$. The kernel of the induced homomorphism $G_P \rightarrow G_Q$ is equal to $\phi^{-1}(W_Q)/W_P$. Since $\phi|_{W_P}$ is injective, the restriction of $P \rightarrow G_P$ to $\ker \phi$ is an isomorphism onto its image

$H \subset \phi^{-1}(W_Q)/W_P$. We have that $(\phi^{-1}(W_Q)/W_P)/H \simeq (W_Q \cap \phi(P))/\phi(W_P)$ is unipotent and normal in G_P/H . It follows from Theorem 22.42 in [Mil17] that G_P/H is reductive. Therefore, we must have $\phi^{-1}(W_Q)/W_P = H \simeq \ker \phi$. So the induced Shimura morphism $(G_P, X_G^+) \rightarrow (G_Q, Y_Q^+)$ is a Shimura immersion as well. Together with Theorem 2.4(2) in [Orr13], the injectivity of $\phi|_{W_P}$, and [Pin90], 2.18, this implies that the map $\Gamma \backslash X^+ \rightarrow \Delta \backslash Y^+$ has countable and therefore finite fibers. Note that Lemma 2.2 in [Orr13] (which is formulated for reductive groups, but holds with the same proof for arbitrary linear algebraic groups) allows us to find congruence subgroups of $G_P(\mathbb{Q})_+$ and $G_Q(\mathbb{Q})_+$ such that we can apply Theorem 2.4(2) in [Orr13].

It therefore remains to show that the continuous map $\Gamma \backslash X^+ \rightarrow \Delta \backslash Y^+$ is closed. A continuous map with compact – e.g., finite – fibers is proper if and only if it is closed.

We choose a Levi decomposition $P = G_P \times W_P$ (see Theorem 2.3 in [PR94]). Let $i_P : G_P \rightarrow P$ denote the induced homomorphism of algebraic groups and let $\pi_P : P \rightarrow G_P$ denote the canonical quotient homomorphism. Fix $x_G \in X_G^+$. It follows from the surjectivity of $X^+ \rightarrow X_G^+$ that there exists $\tilde{x} \in X^+$ such that $(\pi_P)_\mathbb{C} \circ \tilde{x} = x_G$. By Proposition 2.17 in [Pin90], i_P induces a morphism between the associated mixed Shimura data. Proposition 2.18 in [Pin90] then implies that $x = (i_P)_\mathbb{C} \circ x_G \in W_P(\mathbb{R})U_P(\mathbb{C})\tilde{x} \subset X^+$. If $\omega : \mathbb{G}_{m,\mathbb{R}} \rightarrow \mathbb{S}$ is defined by $t \in \mathbb{R}^* \mapsto t \in \mathbb{C}^* = \mathbb{S}(\mathbb{R})$, then the composition $x \circ \omega_\mathbb{C} = (i_P)_\mathbb{C} \circ x_G \circ \omega_\mathbb{C}$ is the base change of a homomorphism $x_\omega : \mathbb{G}_{m,\mathbb{Q}} \rightarrow P$ thanks to Remark 2.2(i) in [Gao17b] (note that $W_P \subset P^{\text{der}}$ and so the center of G_P is isogenous to $P/P^{\text{der}} \simeq G_P/(G_P)^{\text{der}}$).

Let $Z_P(x_\omega)$ denote the centralizer of $x_\omega(\mathbb{G}_{m,\mathbb{Q}})$ in P . The proof of Lemma 1.8 in [Pin90] shows that there is a Levi decomposition $P = G_P \times W_P$ such that $G_P \times \{0\} = Z_P(x_\omega)$ and we will assume that we have chosen this as the Levi decomposition of P . If $Z(x) \subset (Z_P(x_\omega))(\mathbb{R})^+ U_P(\mathbb{C})$ denotes the intersection with $P(\mathbb{R})^+ U_P(\mathbb{C})$ of the centralizer of $x(\mathbb{S}_\mathbb{C})$ in $P_\mathbb{C}$ and $Z(x_G)$ denotes the intersection with $G_P(\mathbb{R})^+$ of the centralizer of $x_G(\mathbb{S}_\mathbb{C})$ in $(G_P)_\mathbb{C}$, then this Levi decomposition induces a homeomorphism $X^+ \simeq P(\mathbb{R})^+ U_P(\mathbb{C})/Z(x) \rightarrow (G_P(\mathbb{R})^+/Z(x_G)) \times W_P(\mathbb{R})U_P(\mathbb{C}) \simeq X_G^+ \times W_P(\mathbb{R})U_P(\mathbb{C})$. The homeomorphism is $P(\mathbb{R})^+ U_P(\mathbb{C})$ -equivariant, where $P(\mathbb{R})^+ U_P(\mathbb{C})$ operates on $W_P(\mathbb{R})U_P(\mathbb{C})$ by multiplication from the left and its operation on X_G^+ is the canonical one.

Using $y = \phi_\mathbb{C} \circ x$, we get in the same way a Levi decomposition of Q and a homeomorphism $Y^+ \rightarrow Y_Q^+ \times W_Q(\mathbb{R})U_Q(\mathbb{C})$. Furthermore, we have $\phi(G_P \times \{0\}) \subset G_Q \times \{0\}$ for these Levi decompositions and the map $X_G^+ \times W_P(\mathbb{R})U_P(\mathbb{C}) \rightarrow Y_Q^+ \times W_Q(\mathbb{R})U_Q(\mathbb{C})$ induced by these homeomorphisms is the canonical one.

After maybe replacing Γ and Δ by smaller congruence subgroups, using again (a more general version of) Lemma 2.2 in [Orr13] in the process, we can assume without loss of generality that $\Gamma = \Gamma_G \times \Gamma_W$, where Γ_G is a neat congruence subgroup of $G_P(\mathbb{Q})_+$ (as defined in [Pin90], 0.5) and $\Gamma_W \subset W_P(\mathbb{Q})$ is a Γ_G -invariant congruence subgroup, and analogously $\Delta = \Delta_G \times \Delta_W$.

By Theorem 2.4(2) in [Orr13], the map $\Gamma_G \backslash X_G^+ \rightarrow \Delta_G \backslash Y_Q^+$ is proper. It follows that the same holds for the map

$$(\Gamma_G \backslash X_G^+) \times_{\Delta_G \backslash Y_Q^+} (\Delta \backslash Y^+) \rightarrow \Delta \backslash Y^+.$$

Consider now the map

$$\Gamma \backslash X^+ \rightarrow (\Gamma_G \backslash X_G^+) \times_{\Delta_G \backslash Y_Q^+} (\Delta \backslash Y^+).$$

Thanks to the homeomorphisms constructed above as well as Proposition 3.3(b) in [Pin90] with its proof together with the neatness of Γ_G and Δ_G , we can cover $\Gamma_G \backslash X_G^+$ with open sets \mathcal{U} over which $\Gamma \backslash X^+$ is homeomorphic to $\mathcal{U} \times (\Gamma_W \backslash W_P(\mathbb{R})U_P(\mathbb{C}))$ while $(\Gamma_G \backslash X_G^+) \times_{\Delta_G \backslash Y_Q^+} (\Delta \backslash Y^+)$ is homeomorphic to $\mathcal{U} \times (\Delta_W \backslash W_Q(\mathbb{R})U_Q(\mathbb{C}))$ (and the map between them is the

canonical one). It then suffices to show that the map

$$\Gamma_W \backslash W_P(\mathbb{R})U_P(\mathbb{C}) \rightarrow \Delta_W \backslash W_Q(\mathbb{R})U_Q(\mathbb{C})$$

is proper.

Set $V_P = W_P/U_P$ and $V_Q = W_Q/U_Q$. By [Pin90], 2.15, U_P and V_P are abelian and W_P is isomorphic to $U_P \times_{\mathbb{Q}} V_P$ as a \mathbb{Q} -variety in such a way that the induced morphisms $U_P \rightarrow U_P \times_{\mathbb{Q}} V_P \rightarrow V_P$ are the canonical ones; analogous statements hold for W_Q . Arguing as above for $\phi^{-1}(U_Q) \subset (\ker \phi)U_P$ in case (1) and using Proposition 2.9 in [Gao17b], we find that $\phi^{-1}(U_Q) \cap W_P = U_P$. The exact sequence $U_P \rightarrow W_P \rightarrow V_P$ induces an exact sequence of congruence subgroups $\Gamma_U \rightarrow \Gamma_W \rightarrow \Gamma_V$. Analogously, we get an exact sequence of congruence subgroups $\Delta_U \rightarrow \Delta_W \rightarrow \Delta_V$.

As $\phi^{-1}(U_Q) \cap W_P = U_P$, the induced homomorphism $\phi_V : V_P \rightarrow V_Q$ is injective. Since any basis of $\Delta_V \cap \phi_V(V_P(\mathbb{Q}))$ can be completed to a basis of Δ_V , the map $\phi_V^{-1}(\Delta_V) \backslash V_P(\mathbb{R}) \rightarrow \Delta_V \backslash V_Q(\mathbb{R})$ is a closed embedding. Furthermore, the map $\Gamma_V \backslash V_P(\mathbb{R}) \rightarrow \phi_V^{-1}(\Delta_V) \backslash V_P(\mathbb{R})$ is proper since Γ_V has finite index in $\phi_V^{-1}(\Delta_V)$. Hence, the map $\Gamma_V \backslash V_P(\mathbb{R}) \rightarrow \Delta_V \backslash V_Q(\mathbb{R})$ is proper and therefore the map

$$(\Gamma_V \backslash V_P(\mathbb{R})) \times_{\Delta_V \backslash V_Q(\mathbb{R})} (\Delta_W \backslash W_Q(\mathbb{R})U_Q(\mathbb{C})) \rightarrow \Delta_W \backslash W_Q(\mathbb{R})U_Q(\mathbb{C})$$

is proper as well.

Using that $W_P(\mathbb{R})U_P(\mathbb{C})$ and $W_Q(\mathbb{R})U_Q(\mathbb{C})$ are homeomorphic to $V_P(\mathbb{R}) \times U_P(\mathbb{C})$ and $V_Q(\mathbb{R}) \times U_Q(\mathbb{C})$ respectively and that the various quotient homomorphisms are covering maps in order to argue as above, we are reduced to showing that the map $\Gamma_U \backslash U_P(\mathbb{C}) \rightarrow \Delta_U \backslash U_Q(\mathbb{C})$ is proper. But this map is the composition of $\Gamma_U \backslash U_P(\mathbb{C}) \rightarrow \phi^{-1}(\Delta_U) \backslash U_P(\mathbb{C})$, which is proper since Γ_U has finite index in $\phi^{-1}(\Delta_U)$, and the closed embedding $\phi^{-1}(\Delta_U) \backslash U_P(\mathbb{C}) \rightarrow \Delta_U \backslash U_Q(\mathbb{C})$, hence proper. \square

Remark 3.7. The argument with Lie algebras in the proof of Proposition 3.6(1) shows that $\phi(P) \cap U_Q = \phi(\phi^{-1}(U_Q)) = \phi(U_P)$ for any Shimura morphism $(P, X^+) \rightarrow (Q, Y^+)$ induced by a homomorphism of algebraic groups $\phi : P \rightarrow Q$, where $U_P \subset P$ and $U_Q \subset Q$ denote the respective subgroups from the definition of a connected mixed Shimura datum. Combined with Proposition 2.9 in [Pin90] and its proof, this shows that being pure or of Kuga type is preserved under taking quotients. We will use these facts throughout the article.

4. INTERSECTIONS OF SPECIAL SUBVARIETIES OF CONNECTED MIXED SHIMURA VARIETIES

In this section and in the next one, we indicate by \mathfrak{O} , \mathfrak{M} , and \mathcal{F} respectively the class of objects, of morphisms, and the functor corresponding to any of the three categories we are considering in Proposition 3.3.

In order to prove (A2) we need to show that, in our context, irreducible components of intersections of special subvarieties are special. This is a well-known, widely used fact but we were unable to find a detailed proof of it in the literature and our definition of a connected mixed Shimura datum slightly differs from the ones in [Pin90] and [Pin05a]. For these reasons we include a proof here.

Before stating the theorem, we note that, for any $\phi \in \mathfrak{M}$, the image of $\mathcal{F}(\phi)$ is closed because of Remark 5.5 in [Gao17b] and Proposition 4.2 in [Pin05a] (which also holds with our definition). We can then call the image of a Shimura morphism a *special subvariety* (see Definition 6.1 below and Definition 5.9(1) in [Gao17b]).

Theorem 4.1. *Irreducible components of intersections of special subvarieties of connected (mixed) Shimura varieties (of Kuga type) are special.*

In order to prove the above theorem we need several auxiliary facts.

Lemma 4.2. *Let (P, X^+) be a connected mixed Shimura datum and let $H \subset P$ be a connected algebraic subgroup. Let U be the normal unipotent subgroup of P from the definition of a connected mixed Shimura datum. Set $X_H = \{x \in X^+; x(\mathbb{S}_{\mathbb{C}}) \subset H_{\mathbb{C}}\}$. Then X_H is a finite (possibly empty) union of $H(\mathbb{R})^+(U \cap H)(\mathbb{C})$ -orbits.*

Proof. We can assume without loss of generality that $X_H \neq \emptyset$.

Let $u : H \rightarrow H/(U \cap H)$ denote the quotient map, where we regard $H/(U \cap H)$ as an algebraic subgroup of P/U . Then, $u_{\mathbb{C}} \circ x$ can be defined over \mathbb{R} for each $x \in X_H$. Let W denote the unipotent radical of P and let $\pi : P \rightarrow P/W$ denote the quotient map. Set $(G, X_G^+) = (P, X^+)/W$ and $G_H = H/(W \cap H)$. The elements of X_G^+ are defined over \mathbb{R} , so we view X_G^+ as a subset of $\text{Hom}(\mathbb{S}, G_{\mathbb{R}})$.

Let X_G be the full $G(\mathbb{R})$ -conjugacy class of any $x \in X_G^+$. We first show that $X_{G_H} = \{x \in X_G; x(\mathbb{S}) \subset (G_H)_{\mathbb{R}}\}$ is a finite union of $G_H(\mathbb{R})$ -conjugacy classes.

For this, we reproduce an argument by Moonen in [Moo]: We first show that $X_H \neq \emptyset$ implies that G_H is reductive. It suffices to show that the image H' of G_H in G^{ad} is reductive since the center $Z(G)$ of G is of multiplicative type by Corollary 17.62(a) in [Mil17]. Let $x \in X_H$. By (MSD.d), conjugation by $x(\sqrt{-1})$ induces a Cartan involution θ on $G_{\mathbb{R}}^{\text{ad}}$, which yields an inner form $G_{\mathbb{R}}^{\text{ad}, \theta}$ of $G_{\mathbb{R}}^{\text{ad}}$ such that $G_{\mathbb{R}}^{\text{ad}, \theta}(\mathbb{R})$ is compact. Since $x \in X_H$, this involution restricts to an involution of $H'_{\mathbb{R}}$, which we also denote by θ . It yields an inner form $H'_{\mathbb{R}}{}^{\theta} \subset G_{\mathbb{R}}^{\text{ad}, \theta}$ of $H'_{\mathbb{R}}$. We deduce that $H'_{\mathbb{R}}{}^{\theta}(\mathbb{R})$ is compact as well, so $H'_{\mathbb{R}}$ has a compact inner form. It follows from Proposition 14.32 in [Mil17] that H' must be reductive and therefore G_H is reductive.

We introduce the following notation: Given an algebraic group Q over \mathbb{R} and a maximal torus $T \subset Q$, we write $RW_Q(T) = N_Q(T)(\mathbb{R})/Z_Q(T)(\mathbb{R})$ for the associated ‘‘real Weyl group’’, where $N_Q(T)$ and $Z_Q(T)$ denote the normalizer and the centralizer of T in Q respectively.

By Corollary 3 on p. 320 of [PR94], there exists a finite set $\{S_1, \dots, S_k\}$ of representatives for the conjugacy classes of maximal tori in $(G_H)_{\mathbb{R}}$ under conjugation by $G_H(\mathbb{R})$. Given a $G_H(\mathbb{R})$ -conjugacy class $X_{\alpha} \subset X_{G_H}$, we can find $i \in \{1, \dots, k\}$ such that some element of X_{α} factors through S_i . For each $G_H(\mathbb{R})$ -conjugacy class $X_{\alpha} \subset X_{G_H}$, we choose an index $i = i(\alpha)$ for which this holds. We also choose an element $x \in X_{\alpha}$ such that $x(\mathbb{S}) \subset S_i$.

Set $S'_i = \text{ad}_{\mathbb{R}}(S_i)$, where $\text{ad} : G \rightarrow G^{\text{ad}}$, and write $x' = \text{ad}_{\mathbb{R}} \circ x$. We claim that the class of x' in $RW_{H'_{\mathbb{R}}}(S'_i) \setminus \text{Hom}(\mathbb{S}, S'_i)$ depends only on X_{α} and the choice of i , but not on x . To see this, suppose we have another element $y \in X_{\alpha}$ such that y factors through S_i too. Choose $h \in G_H(\mathbb{R})$ such that $y = hx$. Let $M \subset (G_H)_{\mathbb{R}}$ be the centralizer of $x(\mathbb{S})$, which is a reductive subgroup of $(G_H)_{\mathbb{R}}$ by Corollary 17.59 in [Mil17]. Let M' be the image of M in $H'_{\mathbb{R}} \subset G_{\mathbb{R}}^{\text{ad}}$. By Proposition 17.20 in [Mil17], S'_i is a maximal torus of $H'_{\mathbb{R}}$ and therefore of M' . Since $h^{-1}S_i h$ is contained in the centralizer of $h^{-1}y = x$, $T := \text{ad}_{\mathbb{R}}(h)^{-1}S'_i \text{ad}_{\mathbb{R}}(h)$ is also a maximal torus of M' .

But $M'(\mathbb{R})$ is compact because the Cartan involution of $G_{\mathbb{R}}^{\text{ad}}$ that is induced by $x(\sqrt{-1})$ restricts to the identity on M' . Therefore, $S'_i(\mathbb{R})$ and $T(\mathbb{R})$ are conjugate in $M'(\mathbb{R})$ by Theorem 15 on p. 250 of [OV90]. By the classification of real tori up to isogeny, $S'_i(\mathbb{R})$ is Zariski dense in S'_i . Thus there exists $m \in M'(\mathbb{R})$ such that $\text{ad}_{\mathbb{R}}(h)^{-1}S'_i \text{ad}_{\mathbb{R}}(h) = m^{-1}S'_i m$. Then $z = \text{ad}_{\mathbb{R}}(h)m^{-1}$ is an element of $N_{H'_{\mathbb{R}}}(S'_i)(\mathbb{R})$, and because $\text{ad}_{\mathbb{R}} \circ y = \text{ad}_{\mathbb{R}} \circ hx = zx'$, we see that $\text{ad}_{\mathbb{R}} \circ y$ gives the same class in $RW_{H'_{\mathbb{R}}}(S'_i) \setminus \text{Hom}(\mathbb{S}, S'_i)$ as x' . So we get a well-defined class $\text{cl}(X_{\alpha}) \in RW_{H'_{\mathbb{R}}}(S'_i) \setminus \text{Hom}(\mathbb{S}, S'_i)$.

Choose now a maximal torus $T_i \subset G_{\mathbb{R}}$ containing S_i , and write $T_i^{\text{ad}} \subset G_{\mathbb{R}}^{\text{ad}}$ for its image under $\text{ad}_{\mathbb{R}}$. Applying the arguments of the previous paragraphs with G in place of G_H , X_G in place of X_{G_H} , T_i in place of S_i , and T_i^{ad} in place of S'_i , i.e., for $H = P$, we find that the $G(\mathbb{R})$ -conjugacy class X_G gives rise to a well-defined element of $RW_{G_{\mathbb{R}}^{\text{ad}}}(T_i^{\text{ad}}) \setminus \text{Hom}(\mathbb{S}, T_i^{\text{ad}})$.

Since $RW_{G_{\mathbb{R}}}^{\text{ad}}(T_i^{\text{ad}})$ is finite by Corollary 17.39(a) in [Mil17], this implies that, given an index i , there are only finitely many elements in $RW_{H_{\mathbb{R}}}^{\text{ad}}(S'_i) \setminus \text{Hom}(\mathbb{S}, S'_i)$ that can occur as $\text{cl}(X_{\alpha})$.

Suppose we have $G_H(\mathbb{R})$ -conjugacy classes X_{α} and X_{β} in X_{G_H} such that $i(\alpha) = i(\beta)$ and $\text{cl}(X_{\alpha}) = \text{cl}(X_{\beta})$. We want to show that X_{G_H} is a finite union of $G_H(\mathbb{R})$ -conjugacy classes. Choose $x \in X_{\alpha}$ and $y \in X_{\beta}$ such that x and y both factor through S_i . Since the group $RW_{H_{\mathbb{R}}}^{\text{ad}}(S'_i)$ is finite, it suffices to show that $\text{ad}_{\mathbb{R}} \circ x = \text{ad}_{\mathbb{R}} \circ y$ implies $X_{\alpha} = X_{\beta}$. If $\text{ad}_{\mathbb{R}} \circ x = \text{ad}_{\mathbb{R}} \circ y$, then x and y differ by a homomorphism $\chi : \mathbb{S} \rightarrow Z(G)_{\mathbb{R}} \cap S_i$, i.e., $y_{\mathbb{C}}(s) = x_{\mathbb{C}}(s) \cdot \chi_{\mathbb{C}}(s)$ for all $s \in \mathbb{S}(\mathbb{C})$. Now we look at the abelianization map $\text{ab} : G \rightarrow G/G^{\text{der}}$. Then $\text{ab}_{\mathbb{R}} \circ \xi$ is independent of the choice of $\xi \in X_G$. Hence we find that the image of χ is contained in $(Z(G) \cap G^{\text{der}})_{\mathbb{R}}$. But this is a finite group by Proposition 19.21(b) in [Mil17] and \mathbb{S} is connected, so it follows that χ is trivial and therefore $X_{\alpha} = X_{\beta}$.

It follows that X_{G_H} is a finite union of $G_H(\mathbb{R})$ -conjugacy classes. We deduce that there exists a finite subset $C_H \subset X_H$ such that for any $x \in X_H$, $\pi_{\mathbb{C}} \circ x$ is conjugate to $\pi_{\mathbb{C}} \circ x_0$ through $G_H(\mathbb{R})$ for some $x_0 \in C_H$. After maybe enlarging C_H , we can even assume that $\pi_{\mathbb{C}} \circ x$ and $\pi_{\mathbb{C}} \circ x_0$ are conjugate through the identity component $G_H(\mathbb{R})^+$.

We want to show that every $x \in X_H$ lies in the $H(\mathbb{R})^+(U \cap H)(\mathbb{C})$ -orbit of some $x_0 \in C_H$. By the above together with Proposition 5.1 in [Mil05], we can assume that $\pi_{\mathbb{C}} \circ x = \pi_{\mathbb{C}} \circ x_0$ for some $x_0 \in C_H$ after replacing x by an $H(\mathbb{R})^+$ -conjugate.

Let $\tilde{\pi} : P/U \rightarrow G$ be the canonical homomorphism and set $T = (\pi_{\mathbb{C}} \circ x_0)(\mathbb{S}_{\mathbb{C}})$. Both $(\tilde{\pi}_{\mathbb{C}})|_{(u_{\mathbb{C}} \circ x)(\mathbb{S}_{\mathbb{C}})}$ and $(\tilde{\pi}_{\mathbb{C}})|_{(u_{\mathbb{C}} \circ x_0)(\mathbb{S}_{\mathbb{C}})}$ are isomorphisms onto T . They each give a Levi decomposition $(\tilde{\pi}_{\mathbb{C}})|_{(H/(U \cap H))_{\mathbb{C}}}^{-1}(T) \simeq T \rtimes ((W \cap H)/(U \cap H))_{\mathbb{C}}$. Since everything is defined over \mathbb{R} here, both Levi decompositions are defined over \mathbb{R} . By Theorem 2.3 in [PR94], they are conjugate through an element of $((W \cap H)/(U \cap H))(\mathbb{R})$. By Proposition 5.1 in [Mil05] and Proposition 14.32 in [Mil17], we can assume that $u_{\mathbb{C}} \circ x = u_{\mathbb{C}} \circ x_0$ after replacing x by an $H(\mathbb{R})^+$ -conjugate. Applying a similar reasoning to $u : H \rightarrow H/(U \cap H)$ over \mathbb{C} finally shows that $x \in X_H$ lies in the $H(\mathbb{R})^+(U \cap H)(\mathbb{C})$ -orbit of $x_0 \in C_H$ as desired. \square

Lemma 4.3. *Let (P, X^+) be a connected mixed Shimura datum and let $H \subset P$ be a connected algebraic subgroup. Let U be the normal unipotent subgroup of P from the definition of a connected mixed Shimura datum. Let $X_H^+ \subset X^+$ be an $H(\mathbb{R})^+(U \cap H)(\mathbb{C})$ -orbit and suppose that H is the Mumford-Tate group of an element $x \in X_H^+$, i.e., the smallest algebraic subgroup of P such that $x(\mathbb{S}_{\mathbb{C}}) \subset H_{\mathbb{C}}$. Then (H, X_H^+) is a Shimura subdatum of (P, X^+) .*

Proof. The normal unipotent subgroup of H from the definition of a connected mixed Shimura datum will of course be $U \cap H$. (MSD.a) is then immediate.

The composition of x with the adjoint representation of $H_{\mathbb{C}}$ induces a representation of $\mathbb{S}_{\mathbb{C}}$ on $\text{Lie } H_{\mathbb{C}}$ and thus a decomposition of $\text{Lie } H_{\mathbb{C}}$ into character eigenspaces. This representation is a subrepresentation of the representation of $\mathbb{S}_{\mathbb{C}}$ on $\text{Lie } P_{\mathbb{C}}$ induced by the composition of x with the adjoint representation of $P_{\mathbb{C}}$. Let W denote the unipotent radical of P . Since the vector subspaces $\text{Lie}(U \cap H)$ and $\text{Lie}(W \cap H)$ of $\text{Lie } H$ are defined over \mathbb{Q} and the composition of x with $H_{\mathbb{C}} \rightarrow (H/(U \cap H))_{\mathbb{C}}$ is defined over \mathbb{R} , it follows from (MSD.b) and (MSD.c) for (P, X^+) that the decomposition of $\text{Lie } H_{\mathbb{C}}$ yields a rational mixed Hodge structure on $\text{Lie } H$ with $W_{-2}(\text{Lie } H) = \text{Lie}(U \cap H)$ and $W_{-1}(\text{Lie } H) = \text{Lie}(W \cap H)$. So (MSD.b) follows.

By the proof of Lemma 4.2, $G_H = H/(W \cap H)$ is reductive, so $W \cap H$ is the unipotent radical of H and (MSD.c) follows as well.

Set $G = P/W$. Let $w : H \rightarrow G_H \subset G$ and $\text{ad} : G \rightarrow G^{\text{ad}}$ denote the respective quotient homomorphisms, and let $Z(G)$ denote the center of G . Proposition 1.20 in [Mil05] shows that (MSD.d) for (P, X^+) implies that the adjoint representation $(\text{Lie } G^{\text{ad}})_{\mathbb{R}}$ of $G_{\mathbb{R}}^{\text{ad}}$ carries an $(\text{ad}_{\mathbb{C}} \circ w_{\mathbb{C}} \circ x)(\sqrt{-1})$ -polarization as defined in [Mil05]. Let ad_H denote

the subrepresentation $\mathrm{ad}(G_H) \rightarrow \mathrm{GL}(\mathrm{Lie}(\mathrm{ad}(G_H)))$ of the adjoint representation $G^{\mathrm{ad}} \rightarrow \mathrm{GL}(\mathrm{Lie} G^{\mathrm{ad}})$. It is equal to the adjoint representation of $\mathrm{ad}(G_H)$ and factors through $(G_H)^{\mathrm{ad}}$.

Since G_H is reductive, it follows from Theorem 22.42 in [Mil17] that also $\mathrm{ad}(G_H) = G_H/(Z(G) \cap G_H)$ is reductive. Let $Z(\mathrm{ad}(G_H))$ denote the center of $\mathrm{ad}(G_H)$. By Proposition 14.23 in [Mil17], $(\ker \mathrm{ad}_H)/Z(\mathrm{ad}(G_H))$ is unipotent (so in particular connected). But $\mathrm{ad}(G_H)/Z(\mathrm{ad}(G_H))$ is semisimple, so $(\ker \mathrm{ad}_H)/Z(\mathrm{ad}(G_H))$ is reductive by Corollary 21.53 in [Mil17]. It follows that $\ker \mathrm{ad}_H = Z(\mathrm{ad}(G_H))$, so we get a faithful representation of $\mathrm{ad}(G_H)/Z(\mathrm{ad}(G_H))$. There are canonical homomorphisms $\mathrm{ad}(G_H) \rightarrow (G_H)^{\mathrm{ad}} \rightarrow \mathrm{ad}(G_H)/Z(\mathrm{ad}(G_H))$. By Corollary 17.62(e) in [Mil17], $(G_H)^{\mathrm{ad}}$ has trivial center, so we get an isomorphism $(G_H)^{\mathrm{ad}} \simeq \mathrm{ad}(G_H)/Z(\mathrm{ad}(G_H))$ and hence a faithful representation of $((G_H)^{\mathrm{ad}})_{\mathbb{R}}$. The same Proposition 1.20 in [Mil05], applied the other way around, now shows that $x(\sqrt{-1})$ induces a Cartan involution on $((G_H)^{\mathrm{ad}})_{\mathbb{R}}$.

Suppose now that the group $(G_H)^{\mathrm{ad}}$ possesses a non-trivial \mathbb{Q} -factor H' such that $H'(\mathbb{R})$ is compact. Let ρ denote the adjoint representation $(G_H)^{\mathrm{ad}} \rightarrow \mathrm{GL}(\mathrm{Lie}(G_H)^{\mathrm{ad}})$. Since H' is normal in $(G_H)^{\mathrm{ad}}$, the subspace $\mathrm{Lie} H' \subset \mathrm{Lie}(G_H)^{\mathrm{ad}}$ is $(G_H)^{\mathrm{ad}}$ -invariant. Therefore, ρ induces a representation $\rho' : (G_H)^{\mathrm{ad}} \rightarrow \mathrm{GL}(\mathrm{Lie} H')$. Composing x with first the quotient map $H_{\mathbb{C}} \rightarrow ((G_H)^{\mathrm{ad}})_{\mathbb{C}}$ and then $\rho'_{\mathbb{C}}$ and noting that the resulting homomorphism is defined over \mathbb{R} by (MSD.a), we get a representation $\rho_{\mathbb{S}} : \mathbb{S} \rightarrow \mathrm{GL}(\mathrm{Lie} H')_{\mathbb{R}}$. Now conjugation by $x(\sqrt{-1})$ induces a Cartan involution on $((G_H)^{\mathrm{ad}})_{\mathbb{R}}$ and therefore on $H'_{\mathbb{R}}$. Since the identity is a Cartan involution on $H'_{\mathbb{R}}$, Corollary 4.3 in Chapter I of [Sat80] implies that the image of $x(\sqrt{-1})$ in $(G_H)^{\mathrm{ad}}(\mathbb{R})$ centralizes $H'_{\mathbb{R}}$. We deduce that $\sqrt{-1} \in (\ker \rho_{\mathbb{S}})(\mathbb{R})$. As the pure Hodge structure on $\mathrm{Lie}(G_H)^{\mathrm{ad}}$ induced by x is of type $\{(0, 0), (1, -1), (-1, 1)\}$, it follows from $\sqrt{-1} \in (\ker \rho_{\mathbb{S}})(\mathbb{R})$ that $\ker \rho_{\mathbb{S}} = \mathbb{S}$. Since H is the Mumford-Tate group of x , this implies that the representation $\rho' : (G_H)^{\mathrm{ad}} \rightarrow \mathrm{GL}(\mathrm{Lie} H')$ must be trivial. But the adjoint representation $H' \rightarrow \mathrm{GL}(\mathrm{Lie} H')$ of the semisimple algebraic group $H' \subset (G_H)^{\mathrm{ad}}$ factors through ρ' and arguing as above shows that its kernel is the center of H' , which is trivial by Corollary 17.62(e) and Theorem 21.51 in [Mil17] since H' is a \mathbb{Q} -factor of an adjoint reductive group. So we contradict the non-triviality of H' and therefore there is no such factor H' and (MSD.d) follows.

Set $W_H = W \cap H$, the unipotent radical of H . The algebraic group H acts trivially on $W_H/(W_H \cap H^{\mathrm{der}})$, but $\mathrm{Lie} W_H/(W_H \cap H^{\mathrm{der}}) \simeq \mathrm{Lie} W_H/\mathrm{Lie}(W_H \cap H^{\mathrm{der}})$ and so it follows from (MSD.c), already proved, that $W_H/(W_H \cap H^{\mathrm{der}}) = \{0\}$, i.e., $W_H \subset H^{\mathrm{der}}$. Therefore, $H/H^{\mathrm{der}} \simeq G_H/(G_H)^{\mathrm{der}}$, which is isogenous to the identity component $Z(G_H)^0$ of the center of G_H .

It therefore suffices to show that $Z(G_H)^0$ is a torus as in (MSD.e). Recall that $w : H \rightarrow G_H$ denotes the quotient map. By Lemma 13.3, Proposition 5.1, and Theorem 5.4 in [Mil05], there exists $y \in X_H^+$ and a maximal torus T_H in G_H such that $z = w_{\mathbb{C}} \circ y$ factors through $(T_H)_{\mathbb{C}}$. We have $Z(G_H)^0 \subset T_H$ by the maximality of T_H . We can find a maximal torus T in G that contains T_H and $Z(G)^0$. Since the category of \mathbb{Q} -tori up to isogeny is semi-simple (see Proposition 4.1 in [Bri17]), it suffices to show that T is a torus as in (MSD.e).

Recall that $G = G^{\mathrm{der}} Z(G)^0$. Therefore, $T = T^{\mathrm{der}} Z(G)^0$, where T^{der} by abuse of notation denotes the identity component $(T \cap G^{\mathrm{der}})^0$ and is a maximal torus in G^{der} . Since $Z(G)^0$ is isogenous to G/G^{der} , (MSD.e) for (P, X^+) shows that it suffices to prove that T^{der} is a torus as in (MSD.e). Now $T^{\mathrm{der}}(\mathbb{R})$ fixes z and so $T^{\mathrm{der}}(\mathbb{R})$ is contained in the stabilizer K_z of z in $G^{\mathrm{der}}(\mathbb{R})$.

The stabilizer K_z is a closed subset of $K_1(\mathbb{R})$, where K_1 denotes the algebraic subgroup of $G_{\mathbb{R}}^{\mathrm{der}}$ of elements commuting with $z(\sqrt{-1})$. Let K_2 denote the algebraic subgroup of $G_{\mathbb{R}}^{\mathrm{ad}}$ of elements commuting with the image of $z(\sqrt{-1})$ under the homomorphism $G_{\mathbb{R}} \rightarrow G_{\mathbb{R}}^{\mathrm{ad}}$. By (MSD.d), conjugation by $z(\sqrt{-1})$ induces a Cartan involution on $G_{\mathbb{R}}^{\mathrm{ad}}$, which implies

that $K_2(\mathbb{R})$ is compact. Clearly K_1 is mapped into K_2 by the homomorphism $G_{\mathbb{R}}^{\text{der}} \rightarrow G_{\mathbb{R}}^{\text{ad}}$. As the topological covering $G^{\text{der}}(\mathbb{C}) \rightarrow G^{\text{ad}}(\mathbb{C})$ has finite fibers, this implies that $K_1(\mathbb{R})$ is compact. Hence, K_z is a compact closed subgroup of $G^{\text{der}}(\mathbb{R})$ and so $T^{\text{der}}(\mathbb{R})$ is compact. This completes the proof. \square

Remark 4.4. By adapting the end of the proof of Lemma 4.3, we can also prove the following: Let (P, X^+) be a connected mixed Shimura datum and let $H \subset P$ be a connected algebraic subgroup. Let U be the normal unipotent subgroup of P from the definition of a connected mixed Shimura datum and let $X_H^+ \subset X^+$ be an $H(\mathbb{R})^+(U \cap H)(\mathbb{C})$ -orbit. If $W = \{1\}$ and (H, X_H^+) satisfies (MSD.a) to (MSD.d), then it also satisfies (MSD.e). This can also be proved if $W \neq \{1\}$, see Example 2.10 in [Pin90].

Lemma 4.5. *Let (P, X^+) be a connected mixed Shimura datum and let $H \subset P$ be a connected algebraic subgroup. Let U be the normal unipotent subgroup of P from the definition of a connected mixed Shimura datum. Let $X_H^+ \subset X^+$ be an $H(\mathbb{R})^+(U \cap H)(\mathbb{C})$ -orbit such that one (equivalently: any) $x \in X_H^+$ factors through $H_{\mathbb{C}}$. Then there exist finitely many $x_i \in X_H^+$ ($i = 1, \dots, n$) such that X_H^+ is the disjoint union of the $M_{x_i}(\mathbb{R})^+(U \cap M_{x_i})(\mathbb{C})x_i$, where $M_{x_i} \subset H$ denotes the Mumford-Tate group of x_i as defined in Lemma 4.3.*

Proof. We first show that $H(\mathbb{R})^+(U \cap H)(\mathbb{C})$ is canonically a connected real analytic manifold: Let W denote the unipotent radical of P . It follows from the proof of Lemma 4.2 that $W_H := W \cap H$ is the unipotent radical of H . Set $G_H = H/W_H$ and $U_H = U \cap H$, and fix a Levi decomposition $H = G_H \ltimes W_H$. We get a homeomorphism $H(\mathbb{R})^+U_H(\mathbb{C}) \rightarrow G_H(\mathbb{R})^+ \times W_H(\mathbb{R})U_H(\mathbb{C})$. Set $V_H = W_H/U_H$ and decompose $\text{Lie } W_H = V \oplus \text{Lie } U_H$ (as a vector space) such that the restriction of the differential $V \rightarrow \text{Lie } V_H$ is an isomorphism. Since the exponential map of a unipotent group is an isomorphism of schemes by Proposition 14.32 in [Mil17], we get an isomorphism of schemes $W_H \rightarrow \text{Lie } W_H \rightarrow \text{Lie } V_H \times_{\mathbb{Q}} \text{Lie } U_H \rightarrow V_H \times_{\mathbb{Q}} U_H$. By the commutative diagram before Proposition 14.32 on p. 289 in [Mil17], this yields a homeomorphism $W_H(\mathbb{R})U_H(\mathbb{C}) \rightarrow V_H(\mathbb{R}) \times U_H(\mathbb{C})$. This gives a canonical structure of a connected real analytic manifold on $H(\mathbb{R})^+U_H(\mathbb{C})$.

Fix now some $x_0 \in X_H^+$. For each $x \in X_H^+$, the set of $h \in H(\mathbb{R})^+U_H(\mathbb{C})$ such that hx_0 factors through the Mumford-Tate group of x is a closed real analytic subset of $H(\mathbb{R})^+U_H(\mathbb{C})$. Note that it suffices to check whether $(hx_0)(2)$ and $(hx_0)(\alpha)$ lie in the Mumford-Tate group of x for some fixed $\alpha \in S^1 \subset \mathbb{C}^* = \mathbb{S}(\mathbb{R})$ of infinite order since 2 and α then generate a Zariski dense subgroup of \mathbb{S} .

Now a connected real analytic manifold cannot be covered by countably many proper closed real analytic subsets by the Baire category theorem and the fact that any proper closed real analytic subset has empty interior by the identity theorem for real analytic functions. Since the set of possible Mumford-Tate groups of elements of X_H^+ is countable, there must be some $x_1 \in X_H^+$ such that all $x \in X_H^+$ factor through the Mumford-Tate group M_{x_1} of x_1 . It then follows from Lemma 4.2 that X_H^+ is a finite union of $M_{x_1}(\mathbb{R})^+(U \cap M_{x_1})(\mathbb{C})$ -orbits, among them $M_{x_1}(\mathbb{R})^+(U \cap M_{x_1})(\mathbb{C})x_1$.

If any such orbit, call it Ω , does not contain any element whose Mumford-Tate group is M_{x_1} , then we iterate the previous step to find that it contains an element x_2 such that all elements of Ω factor through the Mumford-Tate group M_{x_2} of x_2 . Furthermore, we have $M_{x_2} \subsetneq M_{x_1}$ and so $\dim M_{x_1} > \dim M_{x_2}$ since Mumford-Tate groups are connected. The orbit Ω is a finite union of $M_{x_2}(\mathbb{R})^+(U \cap M_{x_2})(\mathbb{C})$ -orbits by Lemma 4.2. Continuing in this way, we eventually prove the lemma as the dimension of the Mumford-Tate group cannot drop below 0. \square

We are now able to prove Theorem 4.1.

Proof of Theorem 4.1. Let S_1 and S_2 be two special subvarieties of a connected (mixed) Shimura variety (of Kuga type) $S = \mathcal{F}(P, X^+, \Gamma)$.

Let $i \in \{1, 2\}$. By definition, we have $S_i = \mathcal{F}(\phi_i)(\mathcal{F}(H_i, Y_i^+, \Gamma_i))$ for a Shimura morphism $\phi_i : (H_i, Y_i^+, \Gamma_i) \rightarrow (P, X^+, \Gamma)$. Let K_i denote the kernel of the homomorphism $H_i \rightarrow P$. By Proposition 2.9 in [Pin90], there is a quotient Shimura datum $(H_i, Y_i^+)/K_i$ and the Shimura morphism $(H_i, Y_i^+) \rightarrow (P, X^+)$ factors as $(H_i, Y_i^+) \rightarrow (H_i, Y_i^+)/K_i \rightarrow (P, X^+)$. Thanks to Lemma 2.2 in [Orr13] (which is formulated for reductive groups, but holds with the same proof for arbitrary linear algebraic groups), we can find a congruence subgroup Γ'_i of $(H_i/K_i)(\mathbb{Q})_+$ that gives a factorization of ϕ_i as

$$(H_i, Y_i^+, \Gamma_i) \rightarrow ((H_i, Y_i^+)/K_i, \Gamma'_i) \rightarrow (P, X^+, \Gamma).$$

Furthermore, being pure or of Kuga type is preserved under taking quotients by Remark 3.7. This shows that we can assume without loss of generality that (H_i, Y_i^+) is a Shimura subdatum of (P, X^+) .

There is a canonical identification $S(\mathbb{C}) = \Gamma \backslash X^+$. We denote the map $X^+ \rightarrow \Gamma \backslash X^+$ by unif . Since (H_i, Y_i^+) is a Shimura subdatum of (P, X^+) , we can identify Y_i^+ with its image in X^+ and it follows that $S_i(\mathbb{C}) = \text{unif}(Y_i^+)$ after this identification ($i = 1, 2$). We have

$$(S_1 \cap S_2)(\mathbb{C}) = \text{unif}(\text{unif}^{-1}(S_1(\mathbb{C})) \cap \text{unif}^{-1}(S_2(\mathbb{C}))) = \bigcup_{\gamma \in \Gamma} \text{unif}(Y_1^+ \cap \gamma Y_2^+).$$

Note that $(\gamma H_2 \gamma^{-1}, \gamma Y_2^+)$ is a Shimura subdatum of (P, X^+) for any $\gamma \in \Gamma$.

Let $U \subset P$ be the normal unipotent subgroup from the definition of a connected mixed Shimura datum. For $\gamma \in \Gamma$, let H_γ denote the identity component of $H_1 \cap \gamma H_2 \gamma^{-1}$. We have $Y_1^+ \cap \gamma Y_2^+ \subset \{x \in X^+; x(\mathbb{S}_{\mathbb{C}}) \subset (H_\gamma)_{\mathbb{C}}\}$. It follows from Remark 3.7 applied to $(\gamma H_i \gamma^{-1}, \gamma Y_i^+) \rightarrow (P, X^+)$ that $U \cap \gamma H_i \gamma^{-1}$ is the normal unipotent subgroup from the definition of a connected mixed Shimura datum for $(\gamma H_i \gamma^{-1}, \gamma Y_i^+)$ ($i = 1, 2, \gamma \in \Gamma$). Therefore, $Y_1^+ \cap \gamma Y_2^+$ is a union of $H_\gamma(\mathbb{R})^+(U \cap H_\gamma)(\mathbb{C})$ -orbits. By Lemma 4.2, it is a finite such union.

Combining Lemmas 4.3 and 4.5 shows that each such orbit is a finite union of sets Y'^+ such that (H', Y'^+) is a Shimura subdatum of (P, X^+) for a connected algebraic subgroup H' of P . If (P, X^+) is pure or of Kuga type, then (H', Y'^+) is also pure or of Kuga type respectively by Proposition 2.9 in [Gao17b].

This implies that $(S_1 \cap S_2)(\mathbb{C})$ is a countable union of sets $S'(\mathbb{C})$ for special subvarieties $S' \subset S$. Since the \mathbb{C} -points of an algebraic variety over \mathbb{C} cannot be covered by the \mathbb{C} -points of countably many proper subvarieties, each component of $S_1 \cap S_2$ must be special. \square

5. PROOF OF PROPOSITION 3.3

As one might expect, axioms (A1) and (A3) are the easiest to verify.

(A1) holds with $(P, X^+, \Gamma) \times (P', X'^+, \Gamma') = (P \times_{\mathbb{Q}} P', X^+ \times X'^+, \Gamma \times \Gamma')$, where $X^+ \times X'^+$ consists of the homomorphisms $(\sigma, \sigma') : \mathbb{S}_{\mathbb{C}} \rightarrow P_{\mathbb{C}} \times_{\mathbb{C}} P'_{\mathbb{C}}$ for $\sigma \in X^+$ and $\sigma' \in X'^+$ (cf. [Pin90], Definition 2.5).

(A3) is satisfied with P the trivial group, $X^+ = \text{Hom}(\mathbb{S}_{\mathbb{C}}, P_{\mathbb{C}})$, and $\Gamma = P(\mathbb{Q})$.

Let us now turn to (A2). We recall that a special subvariety is the image of a Shimura morphism.

Let $\phi : (P, X^+, \Gamma) \rightarrow (Q, Y^+, \Delta)$ and $\psi : (P', X'^+, \Gamma') \rightarrow (Q, Y^+, \Delta)$ be Shimura morphisms and set $V = \mathcal{F}(P, X^+, \Gamma)$, $V' = \mathcal{F}(P', X'^+, \Gamma')$, and $W = \mathcal{F}(Q, Y^+, \Delta)$. We want to prove that $(V \times_W V')_{\text{red}} \subset V \times_{\overline{\mathbb{Q}}} V'$ is a finite union of special subvarieties.

We let $U = V \times_{\overline{\mathbb{Q}}} V' \times_{\overline{\mathbb{Q}}} W \times_{\overline{\mathbb{Q}}} W$ and let $G \subset U$ be the graph of the Shimura morphism $\mathcal{F}(\phi) \times \mathcal{F}(\psi) : V \times_{\overline{\mathbb{Q}}} V' \rightarrow W \times_{\overline{\mathbb{Q}}} W$. We note that U is a connected (mixed) Shimura

variety (of Kuga type) and G is a special subvariety of U because it is the image of the Shimura morphism $(\mathrm{id}_{V \times_{\mathbb{Q}} V'}, \mathcal{F}(\phi) \times \mathcal{F}(\psi)) : V \times_{\mathbb{Q}} V' \rightarrow U$.

Let D be the diagonal in $W \times_{\mathbb{Q}} W$. It is special since it is the image of the Shimura morphism $(\mathrm{id}_W, \mathrm{id}_W) : W \rightarrow W \times_{\mathbb{Q}} W$. Hence, $D' := V \times_{\mathbb{Q}} V' \times_{\mathbb{Q}} D$ is a special subvariety of U .

We can apply Theorem 4.1 to G and D' and obtain that the irreducible components of $G \cap D'$ are special subvarieties of U .

Finally, if $\pi : U \rightarrow V \times_{\mathbb{Q}} V'$ is the projection, which is a Shimura morphism, we have that $(V \times_W V')_{\mathrm{red}}$ is the image of $G \cap D'$ via π and therefore a finite union of special subvarieties. We are done with (A2).

For proving (A4) we need several further auxiliary lemmas. We start by showing that, under certain conditions, a surjective homomorphism of algebraic groups gives a Shimura covering.

Lemma 5.1. *Suppose that (Q, Y^+) is a connected mixed Shimura datum and \tilde{Q} is an algebraic group over \mathbb{Q} . Suppose that there is a surjective homomorphism $\phi : \tilde{Q} \rightarrow Q$ whose kernel is a central torus in \tilde{Q} that is an almost-direct product of a \mathbb{Q} -split torus and a torus of compact type that is defined over \mathbb{Q} .*

If some $y \in Y^+$ factors through $\phi_{\mathbb{C}}$, then there exist a connected mixed Shimura datum (\tilde{Q}, \tilde{Y}^+) and a Shimura covering $(\tilde{Q}, \tilde{Y}^+) \rightarrow (Q, Y^+)$ associated to ϕ . If (Q, Y^+) is a connected pure Shimura datum or a connected mixed Shimura datum of Kuga type, then the same holds for (\tilde{Q}, \tilde{Y}^+) .

Proof. First of all, \tilde{Q} is a connected linear algebraic group by Proposition 8.1 in [Mil17].

Let $y \in Y^+ \subset \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, Q_{\mathbb{C}})$ such that there exists $\tilde{y} \in \mathrm{Hom}(\mathbb{S}_{\mathbb{C}}, \tilde{Q}_{\mathbb{C}})$ with $y = \phi_{\mathbb{C}} \circ \tilde{y}$.

Let W and \tilde{W} denote the unipotent radicals of Q and \tilde{Q} respectively. Let $U \subset W$ be the algebraic subgroup, normal in Q , from the definition of a connected mixed Shimura datum and set

$$\tilde{U} = (\phi|_{\tilde{W}})^{-1}(U) \subset \tilde{W}.$$

This will be the algebraic subgroup of \tilde{W} from the definition of a connected mixed Shimura datum. We start by verifying that $\phi|_{\tilde{W}}$ is an isomorphism onto W and that \tilde{U} is normal in \tilde{Q} .

Since ϕ is surjective, we have that $\phi(\tilde{W})$ is normal in Q . By Corollary 14.7 in [Mil17], $\phi(\tilde{W})$ is also unipotent and hence $\phi(\tilde{W}) \subset W$. By Corollary 14.17 in [Mil17] we have that $\tilde{W} \cap \ker \phi$ is trivial and so $\phi|_{\tilde{W}} : \tilde{W} \rightarrow Q$ is injective.

We have exact sequences

$$1 \rightarrow \ker \phi \rightarrow \phi^{-1}(W) \rightarrow W \rightarrow 1$$

and

$$1 \rightarrow \ker \phi \rightarrow \phi^{-1}(U) \rightarrow U \rightarrow 1.$$

The algebraic group $\phi^{-1}(W)$ is connected by Proposition 8.1 in [Mil17] and it admits a central subnormal series $\phi^{-1}(W) \supset \phi^{-1}(U) \supset \ker \phi \supset \{1\}$, because U is central in W and W/U is commutative by [Pin90], 2.15. Hence $\phi^{-1}(W)$ is nilpotent as defined in Definition 6.34 in [Mil17]. By Corollary 16.48 in [Mil17], $\phi^{-1}(W)$ is isomorphic to the product of a unipotent algebraic group and a torus. The first exact sequence above shows that $\phi^{-1}(W) \simeq (\ker \phi) \times_{\mathbb{Q}} W$. Therefore, \tilde{Q} contains a normal unipotent subgroup that is isomorphic to W . Hence $\dim \tilde{W} \geq \dim W$ and the injective homomorphism $\phi|_{\tilde{W}} : \tilde{W} \rightarrow W$ must be an isomorphism and the same for $\phi|_{\tilde{U}} : \tilde{U} \rightarrow U$.

Any conjugate of \tilde{U} in \tilde{Q} is contained in $\phi^{-1}(U)$ and unipotent. It follows as above from the second exact sequence that $\phi^{-1}(U) \simeq (\ker \phi) \times_{\mathbb{Q}} U \simeq (\ker \phi) \times_{\mathbb{Q}} \tilde{U}$ and hence \tilde{U} must be normal in \tilde{Q} .

We might be tempted to take \tilde{Y}^+ to be the connected component of the orbit of \tilde{y} under conjugation by $\tilde{Q}(\mathbb{R})\tilde{U}(\mathbb{C})$ that contains \tilde{y} , but this might not satisfy (MSD.a). To remedy this, we will multiply $\tilde{y} \in \text{Hom}(\mathbb{S}_{\mathbb{C}}, \tilde{Q}_{\mathbb{C}})$ by a suitable homomorphism $\mathbb{S}_{\mathbb{C}} \rightarrow (\ker \phi)_{\mathbb{C}}$ that we construct in the following. Let χ denote the composition of $y \in \text{Hom}(\mathbb{S}_{\mathbb{C}}, Q_{\mathbb{C}})$ with $Q_{\mathbb{C}} \rightarrow (Q/U)_{\mathbb{C}}$. Let T denote the real torus in $(Q/U)_{\mathbb{R}}$ such that $T_{\mathbb{C}}$ is equal to $\chi(\mathbb{S}_{\mathbb{C}})$. From now on, we view χ as a homomorphism $\mathbb{S} \rightarrow T$. Let $\psi : \tilde{Q}/\tilde{U} \rightarrow Q/U$ be the homomorphism induced by ϕ and set $\tilde{T} = \psi_{\mathbb{R}}^{-1}(T)$. We have an exact sequence

$$1 \rightarrow \ker \phi_{\mathbb{R}} \simeq \ker \psi_{\mathbb{R}} \rightarrow \tilde{T} \rightarrow T \rightarrow 1,$$

where the isomorphism between the two kernels is induced by the restriction of $\tilde{Q} \rightarrow \tilde{Q}/\tilde{U}$ to $\ker \phi$. By Proposition 8.1 and Theorem 15.39 in [Mil17], \tilde{T} is a torus as well.

In the next paragraphs, we will construct $\tilde{\chi} : \mathbb{S} \rightarrow \tilde{T}$ such that χ is equal to $\tilde{\chi}$ composed with $\tilde{T} \rightarrow T$. Let χ' denote the composition of \tilde{y} with $\tilde{Q}_{\mathbb{C}} \rightarrow (\tilde{Q}/\tilde{U})_{\mathbb{C}}$; its image is contained in $\tilde{T}_{\mathbb{C}}$. Then $\tilde{\chi}_{\mathbb{C}}\chi'^{-1}$ canonically yields a homomorphism $\mathbb{S}_{\mathbb{C}} \rightarrow \ker \phi_{\mathbb{C}}$. After replacing \tilde{y} by its product with that homomorphism, we can define \tilde{Y}^+ to be the connected component of the orbit of \tilde{y} under conjugation by $\tilde{Q}(\mathbb{R})\tilde{U}(\mathbb{C})$ that contains \tilde{y} . Condition (MSD.a) is then automatically satisfied.

It remains to find a lift $\tilde{\chi}$ of χ to \tilde{T} . Passing to the character modules and using Theorem 12.23 in [Mil17], we get an exact sequence

$$0 \rightarrow X^*(T) \rightarrow X^*(\tilde{T}) \rightarrow X^*(\ker \psi_{\mathbb{R}}) \rightarrow 0$$

of $\text{Gal}(\mathbb{C}/\mathbb{R})$ -modules and a $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant homomorphism $\chi^* = (\chi_1^*, \chi_2^*) : X^*(T) \rightarrow X^*(\mathbb{S}) \simeq \mathbb{Z}^2$ that we want to extend to a $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant homomorphism $X^*(\tilde{T}) \rightarrow \mathbb{Z}^2$ (where $\text{Gal}(\mathbb{C}/\mathbb{R})$ acts on \mathbb{Z}^2 by swapping the factors). We can find a free subgroup $F \subset X^*(\tilde{T})$ (isomorphic to $X^*(\ker \psi_{\mathbb{R}})$ as an abelian group) such that $X^*(\tilde{T}) \simeq X^*(T) \oplus F$ as abelian groups. Let $p_1 : X^*(\tilde{T}) \rightarrow X^*(T)$ and $p_2 : X^*(\tilde{T}) \rightarrow F$ denote the two projections; by abuse of notation, we will also view them as homomorphisms from $X^*(\tilde{T})$ to itself. If τ denotes the non-trivial element of $\text{Gal}(\mathbb{C}/\mathbb{R})$, which we also view as an automorphism of $X^*(\tilde{T})$, then we will show that an extension of χ^* is given by

$$v \in X^*(\tilde{T}) \mapsto ((\chi_1^* \circ p_1)(v), (\chi_2^* \circ p_1)(v) + (\chi_1^* \circ p_1 \circ \tau \circ p_2)(v)) \in \mathbb{Z}^2. \quad (5.1)$$

Clearly, this is a homomorphism of abelian groups that extends χ^* . We still have to verify that it is $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant.

Let $v \in X^*(\tilde{T})$. Since $\tau(X^*(T)) = X^*(T)$, χ^* is $\text{Gal}(\mathbb{C}/\mathbb{R})$ -equivariant, and $v = p_1(v) + p_2(v)$, we have

$$(\chi_1^* \circ p_1)(\tau(v)) = \chi_1^*(\tau(p_1(v) + p_2(v))) = (\chi_2^* \circ p_1)(v) + (\chi_1^* \circ p_1 \circ \tau \circ p_2)(v).$$

Furthermore, we have that

$$\begin{aligned} & (\chi_2^* \circ p_1)(\tau(v)) + (\chi_1^* \circ p_1 \circ \tau \circ p_2)(\tau(v)) = \\ & (\chi_2^* \circ p_1)(\tau(p_1(v))) + (\chi_2^* \circ p_1)(\tau(p_2(v))) + (\chi_1^* \circ p_1 \circ \tau \circ p_2)(\tau(v)) = \\ & (\chi_1^* \circ p_1)(v) + (\chi_2^* \circ p_1)(\tau(p_2(v))) + (\chi_1^* \circ p_1 \circ \tau \circ p_2)(\tau(v)) = \\ & (\chi_1^* \circ p_1)(v) + \chi_1^*((\tau \circ p_1 \circ \tau \circ p_2)(v) + (p_1 \circ \tau \circ p_2 \circ \tau \circ p_2)(v)). \end{aligned}$$

We can now use that

$$\begin{aligned} & \tau \circ p_1 \circ \tau \circ p_2 + p_1 \circ \tau \circ p_2 \circ \tau \circ p_2 = \\ & p_1 \circ \tau \circ p_1 \circ \tau \circ p_2 + p_1 \circ \tau \circ p_2 \circ \tau \circ p_2 = \\ & p_1 \circ \tau \circ \tau \circ p_2 = p_1 \circ p_2 = 0. \end{aligned}$$

So (5.1) is really the desired extension and we have proven that condition (MSD.a) is satisfied.

The composition of \tilde{y} with the adjoint representation yields a decomposition $\mathrm{Lie} \tilde{Q}_{\mathbb{C}} = \bigoplus_{(m,n) \in \mathbb{Z}^2} L^{m,n}$ into character eigenspaces for the induced representation of $\mathbb{S}_{\mathbb{C}}$. As \tilde{Q} is a central extension of Q and we have an $\mathbb{S}_{\mathbb{C}}$ -equivariant isomorphism $\mathrm{Lie} Q \simeq (\mathrm{Lie} \tilde{Q})/(\mathrm{Lie} \ker \phi)$, the numbers $\dim L^{m,n}$ are equal to the Hodge numbers $h^{m,n}$ (as defined in [Pin90], 1.1) of the rational mixed Hodge structure induced by y on $\mathrm{Lie} Q$ apart from $\dim L^{0,0}$, which equals $h^{0,0} + \dim \ker \phi$.

Recall that \tilde{U} is normal in \tilde{Q} and $\phi|_{\tilde{U}}$ is an isomorphism onto U . This yields an $\mathbb{S}_{\mathbb{C}}$ -equivariant isomorphism $\mathrm{Lie} \tilde{U}_{\mathbb{C}} \simeq \mathrm{Lie} U_{\mathbb{C}}$. Since $\dim L^{-1,-1} = h^{-1,-1}$ and $\mathrm{Lie} U = W_{-2}(\mathrm{Lie} Q)$, we must have $\mathrm{Lie} \tilde{U}_{\mathbb{C}} = L^{-1,-1}$. An analogous argument shows that $\mathrm{Lie} \tilde{W}_{\mathbb{C}} = L^{-1,-1} \oplus L^{0,-1} \oplus L^{-1,0}$. Finally, since the composition of \tilde{y} with $\tilde{Q}_{\mathbb{C}} \rightarrow (\tilde{Q}/\tilde{U})_{\mathbb{C}}$ is defined over \mathbb{R} , we have that complex conjugation acts on $\mathrm{Lie}(\tilde{Q}/\tilde{U})_{\mathbb{C}} \simeq L^{0,-1} \oplus L^{-1,0} \oplus L^{1,-1} \oplus L^{0,0} \oplus L^{-1,1}$ by sending $L^{m,n}$ to $L^{n,m}$.

It follows that conditions (MSD.b) and (MSD.c) are satisfied.

As $\ker \phi$ is central in \tilde{Q} and $\tilde{Q}/\tilde{W} \rightarrow Q/W$ is surjective, there are surjective homomorphisms

$$Q/W \simeq \tilde{Q}/((\ker \phi)\tilde{W}) \rightarrow (\tilde{Q}/\tilde{W})^{\mathrm{ad}} \rightarrow (Q/W)^{\mathrm{ad}}.$$

As the adjoint group of a reductive group is centerless by Corollary 17.62(e) in [Mil17], this shows that $(\tilde{Q}/\tilde{W})^{\mathrm{ad}}$ and $(Q/W)^{\mathrm{ad}}$ are canonically isomorphic. Condition (MSD.d) follows.

We set $\tilde{G} = \tilde{Q}/\tilde{W}$ and $G = Q/W$ and fix Levi decompositions $\tilde{Q} = \tilde{G} \times \tilde{W}$ and $Q = G \times W$. By condition (MSD.e) for (Q, Y^+) , we have $Q^{\mathrm{der}} = G^{\mathrm{der}} \times W$. Furthermore, we have $\tilde{Q}^{\mathrm{der}} = \tilde{G}^{\mathrm{der}} \times \tilde{V}$ for some algebraic subgroup $\tilde{V} \subset \tilde{W}$. As ϕ is surjective, so is $\phi|_{\tilde{Q}^{\mathrm{der}}} : \tilde{Q}^{\mathrm{der}} \rightarrow Q^{\mathrm{der}}$.

Arguing as in the proof that $\phi|_{\tilde{W}} : \tilde{W} \rightarrow W$ is an isomorphism with \tilde{Q}^{der} , Q^{der} , and the identity component $(\phi|_{\tilde{Q}^{\mathrm{der}}})^{-1}(W)^0$ of $(\phi|_{\tilde{Q}^{\mathrm{der}}})^{-1}(W)$ in place of \tilde{Q} , Q , and $\phi^{-1}(W)$ shows that $\phi|_{\tilde{V}} : \tilde{V} \rightarrow W$ is an isomorphism. It follows that $\tilde{V} = \tilde{W}$ and $\tilde{Q}^{\mathrm{der}} = \tilde{G}^{\mathrm{der}} \times \tilde{W}$. We deduce that $\tilde{Q}/\tilde{Q}^{\mathrm{der}} \simeq \tilde{G}/\tilde{G}^{\mathrm{der}}$ and $Q/Q^{\mathrm{der}} \simeq G/G^{\mathrm{der}}$.

Now $\tilde{G}/\tilde{G}^{\mathrm{der}}$ is isogenous to the identity component $Z(\tilde{G})^0$ of the center $Z(\tilde{G})$ of \tilde{G} and G/G^{der} is isogenous to $Z(G)^0$, see Example 19.25 in [Mil17]. Furthermore, ϕ induces a homomorphism $\bar{\phi} : \tilde{G} \rightarrow G$, whose kernel is a central torus isomorphic to $\ker \phi$, and we have an exact sequence

$$1 \rightarrow \ker \bar{\phi} \rightarrow Z(\tilde{G})^0 \rightarrow \bar{\phi}(Z(\tilde{G})^0) \rightarrow 1.$$

Since $\ker \bar{\phi}$ as well as $\bar{\phi}(Z(\tilde{G})^0) \subset Z(G)^0$ are almost-direct products of \mathbb{Q} -split tori with tori of compact type that are defined over \mathbb{Q} and the category of \mathbb{Q} -tori up to isogeny is semi-simple (see Proposition 4.1 in [Bri17]), this establishes condition (MSD.e) for (\tilde{Q}, \tilde{Y}^+) .

Hence, (\tilde{Q}, \tilde{Y}^+) is a connected mixed Shimura datum and ϕ induces a Shimura morphism $(\tilde{Q}, \tilde{Y}^+) \rightarrow (Q, Y^+)$. Since ϕ is surjective and its kernel is a torus, this Shimura morphism is a Shimura covering. Finally, if W is trivial, so is \tilde{W} , and the same holds for U and \tilde{U} . The lemma follows. \square

We are going to use the following lemma to construct the right vertical arrow in the diagram (2.1).

Lemma 5.2. *Suppose that (Q, Y^+) is a connected mixed Shimura datum. Then there is a connected mixed Shimura datum (\tilde{Q}, \tilde{Y}^+) and a Shimura covering $(\tilde{Q}, \tilde{Y}^+) \rightarrow (Q, Y^+)$ such that the associated homomorphism $\tilde{Q} \rightarrow Q$ is surjective and its kernel is a central torus of compact type that is defined over \mathbb{Q} . Furthermore, the derived subgroup \tilde{Q}^{der} of \tilde{Q} is essentially simply connected (as defined in [LV19]).*

If (Q, Y^+) is a connected pure Shimura datum or a connected mixed Shimura datum of Kuga type, then the same holds for (\tilde{Q}, \tilde{Y}^+) .

Proof. Let W be the unipotent radical of Q and set $G = Q/W$. There is an induced connected pure Shimura datum $(G, X^+) = (Q, Y^+)/W$ (see [Pin90], Proposition 2.9). We want to use the construction in Proposition 3.1 and Application 3.4 in Chapter V of [DMOS82] to get a connected pure Shimura datum (\tilde{G}, \tilde{X}^+) and a Shimura morphism $(\tilde{G}, \tilde{X}^+) \rightarrow (G, X^+)$ such that \tilde{G}^{der} is simply connected (as defined in Definition 18.5 in [Mil17]), the induced homomorphism $\tilde{G} \rightarrow G$ is surjective, and its kernel is a central torus in \tilde{G} .

However, the construction in [DMOS82] does not necessarily preserve condition (MSD.e) in the definition of a connected mixed Shimura datum as the kernel of $\tilde{G} \rightarrow G$ might have a non- \mathbb{Q} -split factor of non-compact type (up to isogeny). The construction in [DMOS82] therefore has to be modified.

The following argument was kindly provided to the authors by Chris Daw: By Lemma 13.3 and Theorem 5.4 in [Mil05], there exists $x \in X^+$ and a maximal \mathbb{Q} -torus T in G such that x factors through $T_{\mathbb{C}}$. Let Z denote the center of G and Z^0 its identity component. Recall that $G = G^{\text{der}}Z^0$ and $Z^0 \subset T$ by the maximality of T . Therefore, $T = T^{\text{der}}Z^0$, where T^{der} by abuse of notation denotes the identity component $(T \cap G^{\text{der}})^0$ and is a maximal \mathbb{Q} -torus in G^{der} . By Corollary 17.84 in [Mil17], we must have $T^{\text{der}} = T \cap G^{\text{der}}$. Since $Z^0 \subset T$, it follows that

$$T^{\text{der}} \cap Z^0 = G^{\text{der}} \cap Z^0. \quad (5.2)$$

We now show that $T^{\text{der}}(\mathbb{R})$ is compact in the same way as done at the end of the proof of Lemma 4.3: To start with, $T^{\text{der}}(\mathbb{R})$ fixes x and so $T^{\text{der}}(\mathbb{R})$ is contained in the stabilizer K_x of x in $G^{\text{der}}(\mathbb{R})$. The stabilizer K_x is a closed subset of $K_1(\mathbb{R})$, where K_1 denotes the algebraic subgroup of $G_{\mathbb{R}}^{\text{der}}$ of elements commuting with $x(\sqrt{-1})$. Let K_2 denote the algebraic subgroup of $G_{\mathbb{R}}^{\text{ad}}$ of elements commuting with the image of $x(\sqrt{-1})$ under the homomorphism $G_{\mathbb{R}} \rightarrow G_{\mathbb{R}}^{\text{ad}}$. By (MSD.d), conjugation by $x(\sqrt{-1})$ induces a Cartan involution on $G_{\mathbb{R}}^{\text{ad}}$, which implies that $K_2(\mathbb{R})$ is compact (see the definition in [Mil05], pp. 274–275). Clearly K_1 is mapped into K_2 by the homomorphism $G_{\mathbb{R}}^{\text{der}} \rightarrow G_{\mathbb{R}}^{\text{ad}}$. As the topological covering $G^{\text{der}}(\mathbb{C}) \rightarrow G^{\text{ad}}(\mathbb{C})$ has finite fibers, this implies that $K_1(\mathbb{R})$ is compact. Hence, K_x is a compact closed subgroup of $G^{\text{der}}(\mathbb{R})$ and so $T^{\text{der}}(\mathbb{R})$ is compact.

Now let $\pi : G' \rightarrow G^{\text{der}}$ denote the universal covering of G^{der} as defined in Definition 18.7 in [Mil17] (see, e.g., Theorem 18.25 and Remark 18.27 in [Mil17] for the existence) so that G' is a simply connected semisimple algebraic group. The kernel of π is finite and of multiplicative type. By Theorem 15.39 in [Mil17], $\pi^{-1}(T^{\text{der}})$ is of multiplicative type as well. Let T' denote the identity component $\pi^{-1}(T^{\text{der}})^0$. Then T' is a maximal \mathbb{Q} -torus in G' of compact type. Again by Corollary 17.84 in [Mil17], we have

$$T' = \pi^{-1}(T^{\text{der}}). \quad (5.3)$$

Since the homomorphism π has finite kernel, the induced map between cocharacter modules

$$X_*(T') \rightarrow X_*(T), \quad \chi \mapsto \pi_{\bar{\mathbb{Q}}} \circ \chi$$

is injective, and so we consider $X_*(T')$ as a submodule of $X_*(T)$.

In this proof we several times make use of the fact that cocharacter modules of tori are torsion-free, as \mathbb{G}_m is connected.

Now we consider the canonical maps $\text{ad} : G \rightarrow G^{\text{ad}}$ and $\nu : G \rightarrow C$, where $C = G/G^{\text{der}}$, and we let T^{ad} by abuse of notation denote the maximal \mathbb{Q} -torus $\text{ad}(T)$ of G^{ad} . We obtain an isogeny $T \rightarrow T^{\text{ad}} \times_{\mathbb{Q}} C$ (with kernel contained in $Z \cap G^{\text{der}}$) and hence another inclusion

$$X_*(T) \subset X_*(T^{\text{ad}}) \oplus X_*(C), \quad \chi \mapsto (\text{ad}_{\bar{\mathbb{Q}}} \circ \chi, \nu_{\bar{\mathbb{Q}}} \circ \chi).$$

By construction, $X_*(T')$ is contained in $X_*(T^{\text{ad}}) = X_*(T^{\text{ad}}) \oplus \{0\}$. Hence, we can form the following Cartesian square of Galois modules:

$$\begin{array}{ccc}
P_0 & \longrightarrow & F = X_*(T^{\text{ad}}) \oplus X_*(C) \\
\downarrow & & \downarrow \\
M = X_*(T)/X_*(T') & \hookrightarrow & (X_*(T^{\text{ad}})/X_*(T')) \oplus X_*(C).
\end{array}$$

Furthermore, since F is torsion-free, and the lower horizontal arrow is injective, P_0 is also torsion-free. Since the right vertical arrow is surjective, so too is the left vertical arrow.

Now we proceed exactly as in [DMOS82]. Let P_1 be the kernel of the left vertical arrow, and let P_2 be the fiber product of P_0 and $X_*(T)$ over M (the map $X_*(T) \rightarrow M$ being the projection to the quotient). We obtain the following diagram, having exact rows and columns:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \downarrow & & \downarrow & & \\
& & X_*(T') & \xlongequal{\quad} & X_*(T') & & \\
& & \downarrow & & \downarrow & & \\
0 & \longrightarrow & P_1 & \longrightarrow & P_2 & \longrightarrow & X_*(T) \longrightarrow 0 \\
& & \parallel & & \downarrow & & \downarrow \\
0 & \longrightarrow & P_1 & \longrightarrow & P_0 & \longrightarrow & M \longrightarrow 0 \\
& & & & \downarrow & & \downarrow \\
& & & & 0 & & 0
\end{array} \tag{5.4}$$

Note that P_2 is torsion-free, as it is a submodule of $P_0 \oplus X_*(T)$. Second, observe that P_1 is isomorphic to $X_*(T')$. To see this, suppose that $(0, f) \in P_1 \subset M \oplus F$. Then $f = (t, 0)$ such that $t \in X_*(T')$. Therefore, P_1 is contained in $\{0\} \oplus X_*(T') \oplus \{0\}$, and the reverse inclusion is immediate. Thus, P_1 is also torsion-free. Hence, the \mathbb{Z} -dual of the upper horizontal sequence is also exact. The category equivalence X^* (see [Mil17], Theorem 12.23) then yields a \mathbb{Q} -torus \tilde{T} and an exact sequence of \mathbb{Q} -tori

$$1 \rightarrow T' \rightarrow \tilde{T} \rightarrow T \rightarrow 1 \tag{5.5}$$

such that the \mathbb{Z} -dual of the upper horizontal sequence corresponds to the induced exact sequence of character modules of these \mathbb{Q} -tori.

Pulling back (5.5) along $Z^0 \subset T$ yields another exact sequence of tori

$$1 \rightarrow T' \rightarrow \tilde{Z} \xrightarrow{\eta} Z^0 \rightarrow 1.$$

The group G is isomorphic to $(G' \times_{\mathbb{Q}} Z^0)/E$, where the finite group $E = \pi^{-1}(G^{\text{der}} \cap Z^0) \subset G'$ embeds centrally into the product via $g \in E(\mathbb{Q}) \mapsto (g, \pi(g)^{-1}) \in G'(\mathbb{Q}) \times Z^0(\mathbb{Q})$. The embedding is central since E is mapped into the center of G^{der} by the homomorphism π , which has finite kernel.

The left column of (5.4) yields a homomorphism $T' \hookrightarrow \tilde{T}$ such that the composition $T' \rightarrow \tilde{T} \rightarrow T$ coincides with the morphism $\pi|_{T'}$, that we recall has finite kernel. Furthermore, we have $\tilde{Z} \subset \tilde{T}$ equal to the pre-image of $Z^0 \subset T$. Since $\pi(T') = T^{\text{der}}$, $T = T^{\text{der}} Z^0$, and $T^{\text{der}} \cap Z^0$ is finite, we get an isogeny $T' \times_{\mathbb{Q}} \tilde{Z} \rightarrow \tilde{T}$. Let \tilde{E} be its kernel. It projects bijectively onto its image in either factor.

If $(g, z) \in \tilde{E}(\mathbb{Q}) \subset G'(\mathbb{Q}) \times \tilde{Z}(\mathbb{Q})$, then we have $g \in E(\mathbb{Q})$, hence \tilde{E} is central in $G' \times_{\mathbb{Q}} \tilde{Z}$. We define $\tilde{G} = (G' \times_{\mathbb{Q}} \tilde{Z})/\tilde{E}$. The homomorphism $\text{id}_{G'} \times \eta : G' \times_{\mathbb{Q}} \tilde{Z} \rightarrow G' \times_{\mathbb{Q}} Z^0$ induces a surjective homomorphism $\rho : \tilde{G} \rightarrow G$. By construction, there is an isomorphism $\tilde{G}^{\text{der}} \simeq G'$.

Let $H \subset G' \times_{\mathbb{Q}} \tilde{Z}$ be the algebraic subgroup defined by

$$H(\bar{\mathbb{Q}}) = \{(g, z) \in G'(\bar{\mathbb{Q}}) \times \tilde{Z}(\bar{\mathbb{Q}}); g \in E(\bar{\mathbb{Q}}), \eta(z) = \pi(g)^{-1}\},$$

then $\tilde{E} \subset H$ and $\ker \rho = H/\tilde{E}$. Consider the image E' of \tilde{E} under projection onto $T' \subset G'$. Thanks to (5.2) and (5.3), we have

$$E' = (\pi|_{T'})^{-1}(T^{\text{der}} \cap Z^0) = \pi^{-1}(T^{\text{der}} \cap Z^0) = \pi^{-1}(G^{\text{der}} \cap Z^0) = E.$$

Since the map $\tilde{E} \rightarrow E'$ is bijective, this shows that $\ker \rho$ is isomorphic to $\ker \eta$, i.e., to T' , and central in \tilde{G} .

By construction, $\tilde{T} \simeq (T' \times_{\mathbb{Q}} \tilde{Z})/\tilde{E}$ embeds into \tilde{G} as a maximal torus with $\rho(\tilde{T}) = T'$ and $\ker \rho|_{\tilde{T}} \simeq \ker \rho \simeq T'$. Now $x \in \text{Hom}(\mathbb{S}_{\mathbb{C}}, G_{\mathbb{C}})$ is induced by $x \in \text{Hom}(\mathbb{S}_{\mathbb{C}}, T_{\mathbb{C}})$. Thanks to the exact sequence $1 \rightarrow \ker \rho \rightarrow \tilde{T} \rightarrow T' \rightarrow 1$ of tori, the latter lifts to $\tilde{x} \in \text{Hom}(S_{\mathbb{C}}, \tilde{T}_{\mathbb{C}})$ and hence to $\tilde{x} \in \text{Hom}(\mathbb{S}_{\mathbb{C}}, \tilde{G}_{\mathbb{C}})$. We then define $\tilde{Q} = \tilde{G} \times_G Q \simeq \tilde{G} \times W$. The kernel of the surjective homomorphism $\tilde{Q} \rightarrow Q$ is isomorphic to $\ker \rho$, which is isomorphic to the \mathbb{Q} -torus T' of compact type, and is central in \tilde{Q} . If $y \in Y^+$ maps to $x \in X^+$, then y lifts to $\tilde{y} \in \text{Hom}(\mathbb{S}_{\mathbb{C}}, \tilde{Q}_{\mathbb{C}})$. Most of Lemma 5.2 now follows from Lemma 5.1. That \tilde{Q}^{der} is essentially simply connected follows from the isomorphism $\tilde{Q}^{\text{der}} \simeq \tilde{G}^{\text{der}} \times W$ and the simply connectedness of $\tilde{G}^{\text{der}} \simeq G'$. \square

The following lemma gives almost all we need to deduce (A4).

Lemma 5.3. *Let $\phi : (P, X^+, \Gamma) \rightarrow (Q, Y^+, \Delta) \in \mathfrak{M}$. There are $\phi_1 : (\tilde{P}, \tilde{X}^+, \tilde{\Gamma}) \rightarrow (P, X^+, \Gamma)$, $\phi_2 : (\tilde{P}, \tilde{X}^+, \tilde{\Gamma}) \rightarrow (\tilde{Q}, \tilde{Y}^+, \tilde{\Delta})$, $\phi_3 : (\tilde{Q}, \tilde{Y}^+, \tilde{\Delta}) \rightarrow (Q, Y^+, \Delta) \in \mathfrak{M}$ such that $\phi \circ \phi_1 = \phi_3 \circ \phi_2$ and $\mathcal{F}(\phi_1)$ is a Shimura covering, $\mathcal{F}(\phi_2)$ is a Shimura submersion, and $\mathcal{F}(\phi_3)$ is a Shimura immersion. Moreover, $\phi_2 : \tilde{P} \rightarrow \tilde{Q}$ is surjective, its kernel is connected, and $\phi_2(\tilde{\Gamma}) = \tilde{\Delta}$.*

Proof. Let K be the identity component of $\ker \phi$. By Proposition 2.9 in [Pin90], there is a quotient Shimura datum $(P, X^+)/K$ and the Shimura morphism $(P, X^+) \rightarrow (Q, Y^+)$ factors as $(P, X^+) \rightarrow (P, X^+)/K \rightarrow (Q, Y^+)$, where the second Shimura morphism is a Shimura immersion as the homomorphism $P/K \rightarrow Q$ has finite kernel. Thanks to Lemma 2.2 in [Orr13] (which is formulated for reductive groups, but holds with the same proof for arbitrary linear algebraic groups), we can find a congruence subgroup Γ_K of $(P/K)(\mathbb{Q})_+$ that gives a factorization of ϕ as

$$(P, X^+, \Gamma) \rightarrow ((P, X^+)/K, \Gamma_K) \rightarrow (Q, Y^+, \Delta).$$

Here, we use that the intersection of a congruence subgroup of $(P/K)(\mathbb{Q})$ with $(P/K)(\mathbb{Q})_+$ is again a congruence subgroup of $(P/K)(\mathbb{Q})$ (see Section 3 of this article and Proposition 2.2 in [RS93]).

It suffices to prove the lemma for $(Q, Y^+) = (P, X^+)/K$. If (P, X^+) is pure or of Kuga type, then it follows from Remark 3.7 that the same holds for $(P, X^+)/K$. We then have by construction that the Shimura morphism $(P, X^+) \rightarrow (Q, Y^+)$ is a Shimura submersion, $\phi : P \rightarrow Q$ is surjective, and the kernel of ϕ is connected.

From Lemma 5.2, we get a connected mixed Shimura datum (\tilde{Q}, \tilde{Y}^+) and a Shimura covering $(\tilde{Q}, \tilde{Y}^+) \rightarrow (Q, Y^+)$ with surjective associated homomorphism $\phi_3 : \tilde{Q} \rightarrow Q$ and $\ker \phi_3$ a central torus of compact type that is defined over \mathbb{Q} . Furthermore, \tilde{Q}^{der} is essentially simply connected (as defined in [LV19]). We define $\tilde{P} = \tilde{Q} \times_Q P$. We automatically get homomorphisms $\phi_1 : \tilde{P} \rightarrow P$ and $\phi_2 : \tilde{P} \rightarrow \tilde{Q}$.

Note that the maps $\tilde{Y}^+ \rightarrow Y^+$ and $X^+ \rightarrow Y^+$ are surjective because of Proposition 5.1 in [Mil05]. This implies that given $y \in Y^+$ we can choose pre-images of y in \tilde{Y}^+ and X^+ and thus find $\tilde{x} \in \text{Hom}(\mathbb{S}_{\mathbb{C}}, \tilde{P}_{\mathbb{C}})$ such that $y = \phi_{\mathbb{C}} \circ (\phi_1)_{\mathbb{C}} \circ \tilde{x} = (\phi_3)_{\mathbb{C}} \circ (\phi_2)_{\mathbb{C}} \circ \tilde{x}$.

Now, $\ker \phi_1$ and $\ker \phi_3$ are isomorphic. Furthermore, ϕ_1 is surjective and $\ker \phi_1$ is central in \tilde{P} . Hence, ϕ_1 satisfies the hypotheses of Lemma 5.1. We then get a connected mixed Shimura datum (\tilde{P}, \tilde{X}^+) and a Shimura covering $(\tilde{P}, \tilde{X}^+) \rightarrow (P, X^+)$.

For $H \in \{\tilde{P}, P, \tilde{Q}, Q\}$, let W_H denote the unipotent radical of H , let U_H denote the unipotent subgroup from the definition of a connected mixed Shimura datum, and let $u_H : H \rightarrow H/U_H$ denote the corresponding quotient map. By construction, both $(u_P \circ \phi_1)_{\mathbb{C}} \circ \tilde{x}$ and $(u_{\tilde{Q}} \circ \phi_2)_{\mathbb{C}} \circ \tilde{x}$ are defined over \mathbb{R} . It follows from Remark 3.7, from the fact that $W_{\tilde{P}} \simeq W_{\tilde{Q}} \times_{W_Q} W_P$, and from the proof of Lemma 5.1 that we can assume that $U_{\tilde{P}} \simeq W_{\tilde{Q}} \times_{W_Q} U_P \simeq U_{\tilde{Q}} \times_{U_Q} U_P$ and that $U_{\tilde{Q}} \rightarrow U_Q$ is an isomorphism. We deduce that $\tilde{P}/U_{\tilde{P}} \simeq (\tilde{Q}/U_{\tilde{Q}}) \times_{Q/U_Q} (P/U_P)$. This implies that $(u_{\tilde{P}})_{\mathbb{C}} \circ \tilde{x}$ is also defined over \mathbb{R} . This means that in the proof of Lemma 5.1 we do not have to replace \tilde{x} for (MSD.a) to be satisfied and so we can assume that $\tilde{x} \in \tilde{X}^+$. We therefore also obtain a Shimura morphism $(\tilde{P}, \tilde{X}^+) \rightarrow (\tilde{Q}, \tilde{Y}^+)$ such that the obvious diagram commutes.

Both (\tilde{P}, \tilde{X}^+) and (\tilde{Q}, \tilde{Y}^+) are pure or of Kuga type if (P, X^+) and (Q, Y^+) both are.

Furthermore, ϕ_2 is surjective and its kernel is isomorphic to the kernel of ϕ , hence connected. In particular, $(\tilde{P}, \tilde{X}^+) \rightarrow (\tilde{Q}, \tilde{Y}^+)$ is a Shimura submersion.

We still have to find the congruence subgroups $\tilde{\Gamma}$ and $\tilde{\Delta}$.

We fix a Levi decomposition $\tilde{Q} = \tilde{G} \times W_{\tilde{Q}}$. Then we have $\tilde{Q}^{\text{der}} = \tilde{G}^{\text{der}} \times W_{\tilde{Q}}$ because of (MSD.e). Now $(\tilde{P}/\tilde{P}^{\text{der}})(\mathbb{Q}) \subset (\tilde{P}/\tilde{P}^{\text{der}})(\mathbb{A}_f)$ is discrete by (MSD.e) and Theorem 5.26 in [Mil05]. By choosing a sufficiently small congruence subgroup $\tilde{\Gamma}$ of $\tilde{P}(\mathbb{Q})_+$ inside the pre-image of Γ under ϕ_1 , we can hence suppose that $\tilde{\Gamma} \subset \tilde{P}^{\text{der}}(\mathbb{Q})$ and therefore $\tilde{\Delta} := \phi_2(\tilde{\Gamma}) \subset \tilde{Q}^{\text{der}}(\mathbb{Q})$. We now apply Proposition 0.1(ii) in [LV19] and Theorem 4.1 in [PR94] to the induced (surjective) homomorphism $\tilde{P}^{\text{der}} \rightarrow \tilde{Q}^{\text{der}}$ to deduce that $\tilde{\Delta}$ is a congruence subgroup of $\tilde{Q}^{\text{der}}(\mathbb{Q})$, using that \tilde{Q}^{der} is essentially simply connected (as defined in [LV19]) by construction. Furthermore, $\tilde{\Delta}$ is an arithmetic subgroup of $\tilde{Q}(\mathbb{Q})$ by Theorem 4.1 in [PR94]. We have $\tilde{\Delta} \subset \tilde{Q}(\mathbb{Q})_+$ since $\tilde{\Gamma} \subset \tilde{P}(\mathbb{Q})_+$.

Since $\tilde{Q}^{\text{der}} \subset \tilde{Q}$ is a closed algebraic subgroup, it follows that $\tilde{Q}^{\text{der}}(\mathbb{Q}) \subset \tilde{Q}(\mathbb{Q})$ is closed in the congruence topology. Since the congruence topology on $\tilde{Q}^{\text{der}}(\mathbb{Q})$ is induced by the congruence topology on $\tilde{Q}(\mathbb{Q})$, this implies that $\tilde{\Delta}$ is also closed with respect to the congruence topology in $\tilde{Q}(\mathbb{Q})$, so a congruence subgroup of $\tilde{Q}(\mathbb{Q})$. \square

We are now ready to verify (A4). Let (P, X^+, Γ) and (Q, Y^+, Δ) be two elements of \mathfrak{M} and let $\phi \in \mathfrak{M}$ be a morphism between them, induced by $\phi : P \rightarrow Q$. As already mentioned, the image of $\mathcal{F}(\phi)$ is closed by Remark 5.5 in [Gao17b].

We now apply Lemma 5.3 to our situation. This gives distinguished morphisms ϕ_1, ϕ_2, ϕ_3 with $\phi \circ \phi_1 = \phi_3 \circ \phi_2$ and such that $\mathcal{F}(\phi_1)$ is a Shimura covering and $\mathcal{F}(\phi_3)$ is a Shimura immersion. By Proposition 3.6, the morphism $\mathcal{F}(\phi_1)$ is finite and surjective and $\mathcal{F}(\phi_3)$ has finite fibers.

It remains to check that all fibers of the morphism $\mathcal{F}(\phi_2)$ are irreducible of the same dimension. This is equivalent to showing that the same holds for the corresponding morphism of complex analytic spaces $\tilde{\Gamma} \backslash \tilde{X}^+ \rightarrow \tilde{\Delta} \backslash \tilde{Y}^+$. By Lemma 5.3, the kernel of $\tilde{P} \rightarrow \tilde{Q}$ is connected. This implies that the fibers of the induced map $\tilde{X}^+ \rightarrow \tilde{Y}^+$ are irreducible as complex analytic spaces (cf. Remark 5.3(2) in [Gao17b]). Since all fibers of $\tilde{X}^+ \rightarrow \tilde{Y}^+$ are isomorphic as complex analytic spaces and $\phi_2(\tilde{\Gamma}) = \tilde{\Delta}$, (A4) now follows.

6. SPECIAL AND WEAKLY SPECIAL SUBVARIETIES

In this section as well as in the two following ones, we fix an algebraically closed field K of characteristic 0 and a distinguished category \mathfrak{C} with objects \mathfrak{M} , morphisms \mathfrak{M} , and functor \mathcal{F} to the category of algebraic varieties over K . From now on, we will drop the distinction between objects and morphisms in \mathfrak{C} and their images under \mathcal{F} in order to

avoid excessive notational baggage. The reader is advised to check themselves that all our constructions exist and make sense in \mathfrak{C} when necessary, and not only in the category of varieties over K .

We start by noticing that, by (A4), images of distinguished morphisms are closed subvarieties. Therefore, it makes sense to give the following definition.

Definition 6.1. *A subvariety of a distinguished variety X is called a special subvariety of X if it is the image of a distinguished morphism.*

We will define weakly special subvarieties to be irreducible components of images of pre-images of K -rational points via distinguished morphisms. With the following lemma we see that such images are actually finite unions of subvarieties.

Lemma 6.2. *If $\psi : Z \rightarrow Y$ and $\phi : Z \rightarrow X$ are distinguished morphisms and $y \in Y(K)$, then $\phi(\psi^{-1}(y))$ is closed in X .*

Proof. We consider the distinguished morphism $(\psi, \phi) : Z \rightarrow Y \times_K X$ and the special subvariety $S = (\psi, \phi)(Z) \subset Y \times_K X$. We have $\phi(\psi^{-1}(y)) = q^{-1}(S)$, where q is the (not necessarily distinguished) morphism $X \rightarrow Y \times_K X$ induced by the constant map $X \rightarrow \{y\}$ and the identity on X . Therefore $\phi(\psi^{-1}(y))$ is closed. \square

Definition 6.3. *A subvariety W of a distinguished variety X is called weakly special if there exist distinguished morphisms $\psi : Z \rightarrow Y$, $\phi : Z \rightarrow X$ and a point $y \in Y(K)$ such that W is an irreducible component of $\phi(\psi^{-1}(y))$.*

Remark 6.4. Note that in the category $\mathfrak{C}_{\text{mSv}}$ our definitions of special and weakly special subvarieties coincide with those in [Pin05a] (but recall that the definition of a connected mixed Shimura datum there is different from ours) and [Gao17b].

In the category $\mathfrak{C}_{\text{pSv}}$, they also coincide with the more usual definition as in [Orr13], where one does not assume (MSD.e). One can show this using Remarks 3.5 and 4.4 as well as the diagram in the proof of Proposition 2.6 at the bottom of p. 48 of [Orr13].

Moreover, in the category of connected commutative algebraic groups $\mathfrak{C}_{\text{comm}}$, one can easily check that weakly special subvarieties of a connected commutative algebraic group X are cosets xY where Y is a connected algebraic subgroup of X and $x \in X(K)$. The weakly special subvariety xY is special if and only if it contains a torsion point, i.e., if x can be chosen to have finite order.

The reader might not be very surprised by the following statement.

Lemma 6.5. *Special subvarieties are weakly special.*

Proof. This follows from (A3). \square

We are going to see below two fundamental lemmas saying that images via distinguished morphisms, components of pre-images via distinguished morphisms, and components of intersections of (weakly) special subvarieties are (weakly) special. First we need the following lemma that is also going to be useful later.

Lemma 6.6. *Let W be a weakly special subvariety of a distinguished variety X . Then there exist distinguished morphisms $\psi : Z \rightarrow Y$, $\phi : Z \rightarrow X$ and a point $y \in Y(K)$ such that $W = \phi(\psi^{-1}(y))$.*

Proof. By definition, there exist distinguished morphisms $\psi' : Z' \rightarrow Y'$, $\phi' : Z' \rightarrow X$ and a point $y' \in Y'(K)$ such that W is an irreducible component of $\phi'(\psi'^{-1}(y'))$.

We apply (A4) to get distinguished morphisms $\psi'_1 : U' \rightarrow Z'$, $\psi'_2 : U' \rightarrow V'$, and $\psi'_3 : V' \rightarrow Y'$ such that $\psi' \circ \psi'_1 = \psi'_3 \circ \psi'_2$, ψ'_1 is finite and surjective, ψ'_3 has finite fibers, and $(\psi'_2)^{-1}(\psi'_2(u'))$ is irreducible for all $u' \in U'(K)$.

It follows that

$$\psi'^{-1}(y') = \psi'_1((\psi'_3 \circ \psi'_2)^{-1}(y')) = \bigcup_{v' \in (\psi'_3)^{-1}(y')} \psi'_1((\psi'_2)^{-1}(v')),$$

and hence

$$\phi'(\psi'^{-1}(y')) = \bigcup_{v' \in (\psi'_3)^{-1}(y')} (\phi' \circ \psi'_1)((\psi'_2)^{-1}(v')).$$

As ψ'_3 has finite fibers and $(\psi'_2)^{-1}(\psi'_2(u'))$ is irreducible for all $u' \in U'(K)$, there are at most finitely many $v' \in (\psi'_3)^{-1}(y')$ and each $(\phi' \circ \psi'_1)((\psi'_2)^{-1}(v'))$ is either empty or a subvariety of X by Lemma 6.2. We deduce that

$$W = (\phi' \circ \psi'_1)((\psi'_2)^{-1}(v'))$$

for some $v' \in (\psi'_3)^{-1}(y')$ and the lemma follows. \square

Lemma 6.7. *Images and irreducible components of pre-images of (weakly) special subvarieties under distinguished morphisms are (weakly) special.*

Proof. For images, this follows directly from Lemma 6.6 and the fact that the composition of distinguished morphisms is distinguished.

We now turn to pre-images: Let $\psi : Y \rightarrow Z$, $\phi : U \rightarrow X$, and $\chi : Y \rightarrow X$ be distinguished morphisms. Thanks to Lemma 6.6, it suffices to prove that every irreducible component of $\phi^{-1}(\chi(Y))$ is special and that every irreducible component of $\phi^{-1}(\chi(\psi^{-1}(z)))$ is weakly special for every choice of $z \in Z(K)$.

Now $\phi^{-1}(\chi(Y))$ equals the projection of $(U \times_X Y)_{\text{red}} \subset U \times_K Y$ onto U . Therefore, all its irreducible components are special by (A2).

For the weakly special case, we note that $\phi^{-1}(\chi(\psi^{-1}(z)))$ equals the projection of $\Psi^{-1}(z) \cap (U \times_X Y)_{\text{red}} \subset U \times_K Y$ onto U , where $\Psi : U \times_K Y \rightarrow Z$ is the composition of the canonical projection with ψ (and as such a distinguished morphism).

By (A2), there exist distinguished varieties X_1, \dots, X_n and distinguished morphisms $\phi_i : X_i \rightarrow U \times_K Y$ ($i = 1, \dots, n$) such that $\bigcup_{i=1}^n \phi_i(X_i) = (U \times_X Y)_{\text{red}} \subset U \times_K Y$. It follows that every irreducible component of

$$\Psi^{-1}(z) \cap (U \times_X Y)_{\text{red}} = \bigcup_{i=1}^n \phi_i((\Psi \circ \phi_i)^{-1}(z))$$

is weakly special. The same follows for every irreducible component of $\phi^{-1}(\chi(\psi^{-1}(z)))$. \square

Lemma 6.8. *Every irreducible component of an intersection of two (weakly) special subvarieties is (weakly) special.*

While (A2) might seem slightly unnatural, it is unavoidable (given (A1)) if one wants Lemmas 6.7 and 6.8 to hold (cf. the proof of (A2) in Section 5 above).

Proof. Let X be a distinguished variety and S, S' two special subvarieties. There exists a distinguished morphism $\phi : Y \rightarrow X$ such that $S = \phi(Y)$. By Lemma 6.7, every irreducible component of $\phi^{-1}(S')$ is special. Applying Lemma 6.7 again shows that the same holds for every irreducible component of $\phi(\phi^{-1}(S')) = S \cap S'$.

Let now W, W' be two weakly special subvarieties of X . By Lemma 6.6, there exist distinguished morphisms $\phi : Y \rightarrow X$ and $\psi : Y \rightarrow Z$ as well as a point $z \in Z(K)$ such that $W = \phi(\psi^{-1}(z))$. We have $W \cap W' = \phi(\phi^{-1}(W') \cap \psi^{-1}(z))$. By Lemma 6.7, it suffices to show that every irreducible component of $\phi^{-1}(W') \cap \psi^{-1}(z) = (\phi, \psi)^{-1}(W' \times_K \{z\})$ is weakly special. Applying Lemma 6.7 again, we see that it suffices to show that $W' \times_K \{z\} \subset X \times_K Z$ is weakly special.

By Lemma 6.6, there exist distinguished morphisms $\phi' : Y' \rightarrow X$ and $\psi' : Y' \rightarrow Z'$ and a point $z' \in Z'(K)$ such that $W' = \phi'(\psi'^{-1}(z'))$. We then have that $W' \times_K \{z\} = (\phi' \times \text{id}_Z)((\psi' \times \text{id}_Z)^{-1}(z', z))$ is weakly special. \square

We conclude this section by comparing our approach with the concept of a special structure on a positive-dimensional quasi-projective algebraic variety X over \mathbb{C} as defined by Ullmo in Définition 2.1 in [Ull17]. Such a special structure is a collection of sets $\Sigma_i(X^n)$ ($n \in \mathbb{N} = \{1, 2, \dots\}$, $i = 0, \dots, n \dim X$) such that

- (a) the elements of $\Sigma_i(X^n)$ are subvarieties of X^n of dimension i (called special subvarieties of X^n),
- (b) $X^n \in \Sigma_{n \dim X}(X^n)$,
- (c) if $A \in \Sigma_i(X^n)$ and $B \in \Sigma_j(X^n)$, then every irreducible component of $A \cap B$ belongs to some $\Sigma_k(X^n)$,
- (d) $\Sigma(X^n) := \bigcup_{i=0}^{n \dim X} \Sigma_i(X^n)$ is countable,
- (e) every $Z \in \Sigma(X^n)$ contains a Zariski dense set of points that lie in $\Sigma_0(X^n)$,
- (f) $\Sigma_0(X^n) = \Sigma_0(X)^n$,
- (g) for every n -tuple of integers $(a_1, \dots, a_n) \in \{0, \dots, \dim X\}^n$, we have

$$\prod_{i=1}^n \Sigma_{a_i}(X) \subset \Sigma_{a_1 + \dots + a_n}(X^n),$$

where the inclusion map is the natural one,

- (h) the set $\Sigma_{\dim X}(X^2)$ contains infinitely many subvarieties of X^2 on which the projection to either factor restricts to a surjective and finite morphism,
- (i) for every $Y \in \Sigma(X) \setminus \Sigma_0(X)$, the sets

$$\Sigma_i(Y^n) := \{S \in \Sigma_i(X^n); S \subset Y^n\} \quad (n \in \mathbb{N}, i = 0, \dots, n \dim Y)$$

satisfy Properties (f) to (h) as well (with Y instead of X).

If we take $K = \mathbb{C}$ and take X to be an arbitrary distinguished variety in our setting, then all X^n ($n \in \mathbb{N}$) are distinguished varieties by (A1) and we can define $\Sigma_i(X^n)$ to be the set of special subvarieties of X^n (in our sense) of dimension i for $n \in \mathbb{N}$, $i = 0, \dots, n \dim X$. Properties (a) and (b) are then immediate. Property (c) follows from Lemma 6.8. Properties (f) and (g) follow directly from (A1). Properties (e) and (h) are not satisfied in general; $\mathfrak{C}_{\text{triv}}(Y)$ provides a counterexample (with $X = Y$). Property (d) is also not satisfied in general; $\mathfrak{C}_{\text{add}}$ provides a counterexample (with $X = \mathbb{G}_{a,\mathbb{C}}$) since $\mathbb{G}_{a,\mathbb{C}}^2$ contains uncountably many distinct lines through the origin.

Conversely, given a special structure on some X as defined by Ullmo, it is unclear how to obtain a distinguished category in our sense. So the two approaches have some overlap, which is to be expected as they attempt to formalize the same phenomenon, but they are not equivalent.

7. DEFECT AND DEFECT CONDITION

Lemma 6.8 shows that the following definition makes sense:

Definition 7.1. *For a subvariety V of a distinguished variety X , let $\langle V \rangle$ denote the smallest special subvariety of X that contains V and let $\langle V \rangle_{\text{ws}}$ denote the smallest weakly special subvariety of X that contains V . We call $\delta(V) = \dim \langle V \rangle - \dim V$ the defect of V and $\delta_{\text{ws}}(V) = \dim \langle V \rangle_{\text{ws}} - \dim V$ the weak defect of V .*

As already mentioned in the introduction, the concept of defect is not new as it appears already in [Pin05b]. In [HP16], the authors introduce as geodesic defect what we call here weak defect.

Theorem 7.2 (Defect Condition). *Let X be a distinguished variety and $V' \subset V \subset X$ a sequence of nested subvarieties. Then*

$$\dim \langle V \rangle - \dim \langle V \rangle_{\text{ws}} \leq \dim \langle V' \rangle - \dim \langle V' \rangle_{\text{ws}},$$

or, equivalently,

$$\delta(V) - \delta_{\text{ws}}(V) \leq \delta(V') - \delta_{\text{ws}}(V').$$

Thanks to Proposition 3.3, this proves Conjecture 4.4 in [HP16], at least for an ambient connected mixed Shimura variety or semiabelian variety. Note that, in the case of connected pure Shimura varieties, this is essentially Proposition 4.4 in [DR18]. Other special cases of Theorem 7.2 were established in [HP16] itself as well as in [PT19]. For connected mixed Shimura varieties, Cassani [Cas21a] has recently provided a proof of the defect condition.

Before proving Theorem 7.2, we collect some preliminary results that are going to be useful here and in the next section.

Theorem 7.3 (Fiber Dimension Theorem). *Let K be an algebraically closed field and let $f : V \rightarrow V'$ be a dominant morphism of varieties over K . The following hold:*

- (1) *For every $v' \in V'$, every irreducible component of $f^{-1}(v')$ has dimension at least $\dim V - \dim V'$.*
- (2) *There exists $U \subset V'$ open and dense in V' such that for every $v' \in U$, we have $\dim f^{-1}(v') = \dim V - \dim V'$.*

Proof. This follows from Corollary 14.116 and Remark 14.117 in [GW10] and [GW21]. \square

The following two lemmas both assert that a certain function is constant on all (possibly non-closed) points of a given variety. By constructibility (see Appendix E of [GW10]), it suffices to show that the respective function is constant on the closed points of the variety and we will prove the lemmas in this modified form.

Lemma 7.4. *Let $\phi : X \rightarrow Y$ be a distinguished morphism and $S \subset X$ a special subvariety. Then the function $\dim_s \phi|_S^{-1}(\phi(s))$ is constant on S .*

Proof. We have $S = \psi(Z)$ for a distinguished morphism $\psi : Z \rightarrow X$. Let F be an irreducible component of $\phi|_S^{-1}(y)$ for some $y \in (\phi(S))(K)$. Note that $\phi(S) = (\phi \circ \psi)(Z)$ is a subvariety of Y by (A4). By the Fiber Dimension Theorem, we have $\dim F \geq \dim S - \dim \phi(S)$. We want to show that $\dim F = \dim S - \dim \phi(S)$.

All irreducible components of $(\phi \circ \psi)^{-1}(y) = (\phi|_S \circ \psi)^{-1}(y)$ are weakly special of dimension

$$\dim Z - \dim(\phi \circ \psi)(Z) = \dim Z - \dim \phi(S)$$

by (A4) and the Fiber Dimension Theorem. By Lemma 6.7, their images under ψ are closed in X .

We consider the set \mathcal{I} of all such irreducible components that are mapped onto F by ψ . As $S = \psi(Z)$, we have $\mathcal{I} \neq \emptyset$.

We now assume that $\dim F > \dim S - \dim \phi(S)$ and aim for a contradiction. By the Fiber Dimension Theorem, applied to $\psi|_V : V \rightarrow F$ for each $V \in \mathcal{I}$, there exists a point $x \in F(K)$ that is not contained in the image of any irreducible component of $(\phi \circ \psi)^{-1}(y)$ not belonging to \mathcal{I} such that

$$\dim \psi|_V^{-1}(x) = \dim V - \dim F < \dim V - \dim S + \dim \phi(S) = \dim Z - \dim S$$

for all $V \in \mathcal{I}$. As x is not contained in the image of any irreducible component of $(\phi \circ \psi)^{-1}(y)$ not belonging to \mathcal{I} , we have

$$\psi^{-1}(x) = \bigcup_{V \in \mathcal{I}} \psi|_V^{-1}(x),$$

and therefore $\dim \psi^{-1}(x) < \dim Z - \dim S = \dim Z - \dim \psi(Z)$. But this contradicts the Fiber Dimension Theorem. The lemma follows. \square

Lemma 7.5. *Let $\phi : X \rightarrow Y$ be a distinguished morphism and $W \subset X$ a weakly special subvariety. Then the function $\dim_w \phi|_W^{-1}(\phi(w))$ is constant on W .*

Proof. By Lemma 6.6, we can write $W = \psi(\chi^{-1}(z))$ for distinguished morphisms $\psi : U \rightarrow X$, $\chi : U \rightarrow Z$ and $z \in Z(K)$. Let Δ be the diagonal in $U \times_K U$.

Let F be an irreducible component of $\phi|_W^{-1}(y)$ for some $y \in (\phi(W))(K)$. Then $F \times_K \{z\}$ is an irreducible component of

$$F' = (\phi^{-1}(y) \cap \psi(\chi^{-1}(z))) \times_K \{z\} = (\phi \times \text{id}_Z)|_{(\psi \times \chi)(\Delta)}^{-1}(y, z),$$

where $\phi \times \text{id}_Z : X \times_K Z \rightarrow Y \times_K Z$ is a distinguished morphism.

By Lemma 6.7, $(\psi \times \chi)(\Delta)$ is special since Δ is special as the image of the distinguished morphism $(\text{id}_U, \text{id}_U)$. It follows from Lemma 7.4 that every irreducible component of F' has the same dimension, independently of the choice of y . The lemma follows. \square

Lemma 7.6. *Let $f : V \rightarrow V'$ be a dominant morphism of varieties over K such that $\dim_v f^{-1}(f(v))$ is constant on V . Let $V'_1 \subset V'$ be a subvariety of V' and suppose that V_1 is an irreducible component of $f^{-1}(V'_1)$ that dominates V'_1 . Then $\dim V - \dim V' = \dim V_1 - \dim V'_1$.*

Proof. The set of irreducible components of $f^{-1}(V'_1)$ that dominate V'_1 is in bijection with the set of irreducible components of the generic fiber of $f|_{f^{-1}(V'_1)} : f^{-1}(V'_1) \rightarrow V'_1$, see (2.1.8) in [Gro60]. By hypothesis and the Fiber Dimension Theorem, these are all of the same dimension $\dim V - \dim V'$. It follows from the Fiber Dimension Theorem that all irreducible components of $f^{-1}(V'_1)$ that dominate V'_1 , so in particular V_1 , are of dimension $\dim V'_1 + \dim V - \dim V'$. \square

Proof of Theorem 7.2. Note that $\langle V \rangle = \langle \langle V \rangle_{\text{ws}} \rangle$. Therefore it is enough to show that, if $W' \subset W \subset X$ is a sequence of nested subvarieties, with W, W' weakly special, then

$$\dim \langle W \rangle - \dim W \leq \dim \langle W' \rangle - \dim W'.$$

We can assume without loss of generality that W' is an irreducible component of $\langle W' \rangle \cap W$.

We will choose the notation such that \tilde{S}' denotes a special subvariety while all letters derived from W denote weakly special subvarieties. They will be defined by taking irreducible components of intersections or pre-images under distinguished morphisms of (weakly) special subvarieties and we tacitly use Lemmas 6.7 and 6.8 to see that they are (weakly) special.

By Lemma 6.6, there exist distinguished morphisms $\phi : X' \rightarrow X$, $\psi : X' \rightarrow Y'$ and a point $y' \in Y'(K)$ such that $W = \phi(\psi^{-1}(y'))$.

For reference, the following diagram shows most of the weakly special subvarieties that will appear:

$$\begin{array}{ccccc} \{y'\} & \subset & \psi(\tilde{W}) & \supset & \psi(\tilde{W}') \\ \uparrow & & \uparrow & & \uparrow \\ \hat{W} & \subset & \tilde{W} & \supset & \tilde{W}' \\ \downarrow & \swarrow & & \swarrow & \\ W & \supset & & \supset & W'. \end{array}$$

Here, the varieties in the top row are contained in Y' , those in the middle row in X' , and those in the bottom row in X . Furthermore, all maps are surjective and induced by either ϕ (middle row to bottom row) or ψ (middle row to top row).

Let \hat{W} be an irreducible component of $\psi^{-1}(y')$ such that $\phi(\hat{W}) = W$. As $W \subset \phi(\langle \hat{W} \rangle)$ and $\phi(\langle \hat{W} \rangle)$ is special by Lemma 6.7, we must have $\langle W \rangle \subset \phi(\langle \hat{W} \rangle)$. On the other hand, $\langle \hat{W} \rangle$ is contained in $\phi^{-1}(\langle W \rangle)$ by Lemma 6.7, so it follows that $\phi(\langle \hat{W} \rangle) = \langle W \rangle$.

Let \tilde{W} be an irreducible component of $\phi|_{\langle \hat{W} \rangle}^{-1}(W) = \phi^{-1}(W) \cap \langle \hat{W} \rangle$ that contains \hat{W} . We deduce that $\phi(\tilde{W}) = W$. By Lemmas 7.4 and 7.6, applied to $\phi|_{\langle \hat{W} \rangle}$, W , and \tilde{W} , we have

$$\dim \tilde{W} - \dim W = \dim \langle \hat{W} \rangle - \dim \langle W \rangle \quad (7.1)$$

or equivalently

$$\dim \langle W \rangle - \dim W = \dim \langle \hat{W} \rangle - \dim \tilde{W}. \quad (7.2)$$

Applying Lemmas 7.4 and 7.5 together with the Fiber Dimension Theorem to $\psi|_{\langle \hat{W} \rangle}$ and $\psi|_{\tilde{W}}$ respectively and noting that $\langle \hat{W} \rangle$ as well as \tilde{W} contain an irreducible component of $\psi^{-1}(y')$, we find that

$$\dim \langle \hat{W} \rangle - \dim \psi(\langle \hat{W} \rangle) = \dim \psi^{-1}(y') = \dim \tilde{W} - \dim \psi(\tilde{W}),$$

which implies

$$\dim \langle \hat{W} \rangle - \dim \tilde{W} = \dim \psi(\langle \hat{W} \rangle) - \dim \psi(\tilde{W}). \quad (7.3)$$

Let \tilde{S}' be an irreducible component of $\phi|_{\langle \hat{W} \rangle}^{-1}(\langle W' \rangle) = \phi^{-1}(\langle W' \rangle) \cap \langle \hat{W} \rangle$ containing an irreducible component of $\phi|_{\tilde{W}'}^{-1}(W')$ that contains an irreducible component of $\phi|_{\tilde{W}'}^{-1}(W')$ that surjects onto W' . Since $W' \subset \phi(\tilde{S}') \subset \langle W' \rangle$ and $\phi(\tilde{S}')$ is special by Lemma 6.7, we must have $\phi(\tilde{S}') = \langle W' \rangle$.

It follows from Lemmas 7.4 and 7.6, applied to $\phi|_{\langle \hat{W} \rangle}$, $\langle W' \rangle$, and \tilde{S}' , that

$$\dim \tilde{S}' - \dim \langle W' \rangle = \dim \langle \hat{W} \rangle - \dim \langle W \rangle. \quad (7.4)$$

Let \tilde{W}' be an irreducible component of $\tilde{S}' \cap \tilde{W}$ that contains an irreducible component of $\phi|_{\tilde{W}'}^{-1}(W')$ that surjects onto W' . Since W' is an irreducible component of $\langle W' \rangle \cap W$ and $\phi(\tilde{W}') \subset \phi(\tilde{S}') \cap \phi(\tilde{W}) = \langle W' \rangle \cap W$, we must then have that $\phi(\tilde{W}') = W'$. It follows that \tilde{W}' in fact is an irreducible component of $\phi|_{\tilde{W}'}^{-1}(W')$ that surjects onto W' . It follows from Lemmas 7.5 and 7.6, applied to $\phi|_{\tilde{W}'}$, W' , and \tilde{W}' , that

$$\dim \tilde{W}' - \dim W' = \dim \tilde{W} - \dim W$$

and therefore, by (7.1),

$$\dim \tilde{W}' - \dim W' = \dim \langle \hat{W} \rangle - \dim \langle W \rangle. \quad (7.5)$$

Combining (7.4) and (7.5), we find that

$$\dim \tilde{S}' - \dim \tilde{W}' = \dim \langle W' \rangle - \dim W'. \quad (7.6)$$

We have $\hat{W} \cap \tilde{S}' \neq \emptyset$ and therefore $y' \in \psi(\tilde{S}')$. We deduce that $\hat{W} \subset \psi^{-1}(\psi(\tilde{S}'))$ and therefore $\langle \hat{W} \rangle \subset \psi^{-1}(\psi(\tilde{S}'))$ by Lemma 6.7. Since $\tilde{S}' \subset \langle \hat{W} \rangle$, it follows that we must have $\psi(\tilde{S}') = \psi(\langle \hat{W} \rangle)$. As $\tilde{W}' \subset \tilde{W}$, we deduce that

$$\dim \psi(\langle \hat{W} \rangle) - \dim \psi(\tilde{W}) \leq \dim \psi(\tilde{S}') - \dim \psi(\tilde{W}'),$$

which by (7.2) and (7.3) implies that

$$\dim \langle W \rangle - \dim W \leq \dim \psi(\tilde{S}') - \dim \psi(\tilde{W}'). \quad (7.7)$$

Thanks to Lemmas 7.4 and 7.5, applied to $\psi|_{\tilde{S}'}$ and $\psi|_{\tilde{W}'}$ respectively, together with the Fiber Dimension Theorem and $\tilde{W}' \subset \tilde{S}'$, it follows that

$$\dim \tilde{W}' - \dim \psi(\tilde{W}') \leq \dim \tilde{S}' - \dim \psi(\tilde{S}')$$

and hence

$$\dim \psi(\tilde{S}') - \dim \psi(\tilde{W}') \leq \dim \tilde{S}' - \dim \tilde{W}'.$$

In combination with (7.6) and (7.7), this yields the theorem. \square

8. OPTIMALITY

In this section, we introduce (weak) optimality. As for the defect, this is not a new concept. In [HP16], weak optimality is called geodesic-optimality while in [Poi01] it is called cd-maximality.

Definition 8.1. *Let V be a subvariety of a distinguished variety X . A subvariety $W \subset V$ is called optimal for V in X if $\delta(U) > \delta(W)$ for every subvariety U such that $W \subsetneq U \subset V$. It is called weakly optimal for V in X if it satisfies the same property with δ_{ws} in place of δ .*

Proposition 8.2. *Let V be a subvariety of a distinguished variety X . If $W \subset V$ is optimal for V in X , then it is weakly optimal for V in X .*

Proof. This can be deduced purely formally from Theorem 7.2 as in the proof of Proposition 4.5 in [HP16]. For the reader's convenience, we reproduce the proof here: Let U be a subvariety of X such that $W \subset U \subset V$, $\delta_{\text{ws}}(U) \leq \delta_{\text{ws}}(W)$, and U is weakly optimal for V in X . Then it follows from Theorem 7.2, applied to $W \subset U$, that $\delta(U) - \delta_{\text{ws}}(U) \leq \delta(W) - \delta_{\text{ws}}(W)$ and so

$$\delta(U) = \delta_{\text{ws}}(U) + \delta(U) - \delta_{\text{ws}}(U) \leq \delta_{\text{ws}}(W) + \delta(W) - \delta_{\text{ws}}(W) = \delta(W).$$

Since W is optimal for V in X , we deduce that $U = W$ and so W is weakly optimal for V in X . \square

The following lemma is a weaker analogue of Lemma 2.6 in [BD19] (see Lemma 10.5 for the missing part).

Lemma 8.3. *Let $\psi : X \rightarrow X'$ be a distinguished morphism. Let $W \subset V \subset X$ be a sequence of nested subvarieties. Let V' be the closure of $\psi(V)$ and let W' be the closure of $\psi(W)$. Then the following hold:*

- (1) *Suppose that $V_0 \subset V$ is open and dense in V such that $\psi(V_0)$ is open and dense in V' and $\psi|_{V_0} : V_0 \rightarrow \psi(V_0)$ is smooth. If W is optimal for V in X , W intersects V_0 , and $\langle W \rangle$ contains an irreducible component of a fiber of ψ , then W' has defect at most $\delta(W)$ and is optimal for V' in X' .*
- (2) *If ψ has finite fibers, then $\delta(W') = \delta(W)$.*

Proof. For (1), let W be optimal for V in X such that $W \cap V_0 \neq \emptyset$, where $\psi|_{V_0} : V_0 \rightarrow \psi(V_0)$ is smooth of relative dimension n . Let U' be a subvariety of X' such that $W' \subset U' \subset V'$ and $\delta(U') \leq \delta(W')$. Since $\langle W \rangle$ contains an irreducible component of a fiber of ψ , it follows from Lemma 7.4 that

$$\dim \langle W \rangle - \dim \psi(\langle W \rangle) = \dim X - \dim \psi(X) =: m.$$

Since $\psi(\langle W \rangle)$ furthermore contains $\langle W' \rangle$ thanks to Lemma 6.7, we obtain that

$$\delta(U') \leq \delta(W') \leq \dim \psi(\langle W \rangle) - \dim W' = \dim \langle W \rangle - \dim W' - m.$$

Let now U be an irreducible component of $\psi^{-1}(U') \cap V = \psi|_V^{-1}(U')$ that contains W . We have $U \cap V_0 \neq \emptyset$ since it contains $W \cap V_0$. Furthermore, $U \cap V_0$ is an irreducible component of $\psi|_{V_0}^{-1}(U' \cap \psi(V_0))$. Since $\psi|_{V_0} : V_0 \rightarrow \psi(V_0)$ is smooth of relative dimension n , it follows that $\dim U = \dim(U \cap V_0) = \dim(U' \cap \psi(V_0)) + n = \dim U' + n$. Taking $U' = W'$ and using that $W \subset U$ shows that $\dim W \leq \dim W' + n$. If U' is again an arbitrary subvariety as above, we deduce that

$$\delta(U') \leq \delta(W') \leq \dim \langle W \rangle - m - \dim W + n. \tag{8.1}$$

After noting that $n \leq \dim X - \dim \psi(X)$, we deduce that the defect of W' is at most the defect of W .

Assume now that W' is not optimal for V' in X' . It follows that we can choose U' as above with $W' \subsetneq U'$. Since $\dim U' > \dim W'$, we have

$$\dim U = \dim U' + n > \dim W' + n \geq \dim W.$$

It follows that $W \subsetneq U$ and so $\delta(U) > \delta(W)$ by the optimality of W . As any irreducible component of $\psi^{-1}(\langle U' \rangle)$ is special by Lemma 6.7, we have $\dim \langle U \rangle \leq \dim \langle U' \rangle + (\dim X - \dim \psi(X))$. It follows that $\delta(U) \leq \delta(U') + (\dim X - \dim \psi(X)) - n$.

Using (8.1), we deduce that

$$\delta(U') \leq \delta(W) - m + n < \delta(U) - m + n \leq \delta(U'),$$

a contradiction.

For (2), we note that $\dim W' = \dim W$. Furthermore, we have $\langle W' \rangle \subset \psi(\langle W \rangle) \subset \psi(X)$ and $\langle W \rangle \subset \psi^{-1}(\langle W' \rangle)$. It follows that $\dim \langle W' \rangle \leq \dim \psi(\langle W \rangle)$ and

$$\dim \psi(\langle W \rangle) = \dim \langle W \rangle \leq \dim \psi^{-1}(\langle W' \rangle) = \dim \langle W' \rangle.$$

Hence, we have that $\langle W' \rangle = \psi(\langle W \rangle)$ and $\delta(W') = \delta(W)$. \square

9. WEAK FINITENESS AND VERY DISTINGUISHED CATEGORIES

We can now introduce a fifth axiom that a distinguished category can satisfy:

- (A5) *Weak Finiteness* - If X is a distinguished variety and $V \subset X$ is a subvariety, then there exists a finite set of pairs (ϕ, ψ) of distinguished morphisms $\phi : Y_\phi \rightarrow X$ and $\psi : Y_\psi \rightarrow Z_\psi$ such that for every subvariety $W \subset V$ that is weakly optimal for V in X there exists a pair (ϕ, ψ) in this set and $z \in Z_\psi(K)$ such that ϕ has finite fibers and $\langle W \rangle_{\text{ws}}$ is an irreducible component of $\phi(\psi^{-1}(z))$.

Axiom (A5) is prominently not satisfied by $\mathfrak{C}_{\text{add}}$ (or $\mathfrak{C}_{\text{comm}}$): Consider $X = \mathbb{G}_{a,K}^3$ with affine coordinates x, y, z and $V \subset X$ defined by $x^2 + y^2 = z^2$. Then V is covered by lines, but is not a plane, so every line it contains is weakly optimal for V in X . If (A5) were satisfied, then the set of the slopes of all these lines would be finite, but it is not.

Unlike (A1) to (A4), Axiom (A5) is not preserved when taking products of distinguished categories as defined at the end of Section 2: We will see later that \mathfrak{C}_{ab} over $\bar{\mathbb{Q}}$ satisfies (A5). Let E be an elliptic curve over $\bar{\mathbb{Q}}$ and consider the object (E^2, E^2) of $\mathfrak{C}_{\text{ab}} \times \mathfrak{C}_{\text{ab}}$. It is sent to $E^2 \times_{\bar{\mathbb{Q}}} E^2$ by the functor associated to $\mathfrak{C}_{\text{ab}} \times \mathfrak{C}_{\text{ab}}$. Let $\Delta \subset E^2 \times_{\bar{\mathbb{Q}}} E^2$ denote the diagonal. For every one-dimensional abelian subvariety $A \subset E^2$, we have that $W = (A \times_{\bar{\mathbb{Q}}} A) \cap \Delta$ is weakly optimal for Δ in $E^2 \times_{\bar{\mathbb{Q}}} E^2$ with $\langle W \rangle_{\text{ws}} = A \times_{\bar{\mathbb{Q}}} A$ and there are infinitely many such A .

Let $K \subset L$ be an extension of algebraically closed fields. Given a distinguished category \mathfrak{C} over K with associated functor \mathcal{F} , one can check that (A1) to (A4) are preserved under base change (see Table 1 on pp. 306–307 of [Poo17]) and we naturally obtain a distinguished category \mathfrak{C}_L over L by composing \mathcal{F} with the base change functor. Distinguished varieties and morphisms in \mathfrak{C}_L are then base changes of distinguished varieties and morphisms in \mathfrak{C} . Note that special subvarieties in \mathfrak{C}_L are base changes of special subvarieties in \mathfrak{C} while the same fact clearly does not hold for weakly special subvarieties.

We can now introduce a subclass of distinguished categories that will be used in the following.

Definition 9.1. *Let K be an algebraically closed field. A very distinguished category over K is a distinguished category \mathfrak{C} over K such that for every extension of algebraically closed fields $K \subset L$ of finite transcendence degree, the base change \mathfrak{C}_L satisfies (A5).*

As $\mathfrak{C}_{\text{add}}$ and $\mathfrak{C}_{\text{comm}}$ do not satisfy (A5) as noted above, they certainly are not very distinguished. On the other hand, it is trivial that $\mathfrak{C}_{\text{triv}}(X)$ is very distinguished. We do not know if there are distinguished categories that satisfy (A5), but are not very

distinguished (the question was raised by Orr). Similarly, we do not know whether weak optimality is preserved under base change in an arbitrary distinguished category.

The categories \mathfrak{C}_{ab} , $\mathfrak{C}_{\text{tor}}$, and $\mathfrak{C}_{\text{semiab}}$ are very distinguished as well: For \mathfrak{C}_{ab} , see [Rém09] and [HP16]. For $\mathfrak{C}_{\text{tor}}$, see Corollaire 3.7 in [Poi01] (a similar finiteness statement is proven in [BMZ07]; see also Corollary 3 in [Zil02]). These results are over \mathbb{C} , but they imply the same statement over an arbitrary algebraically closed field (see the proof of Theorem 9.3 below). This also justifies a step in the proof of Proposition 4.1 in [BD19].

In Theorem 9.3, we will show that $\mathfrak{C}_{\text{semiab}}$ over an algebraically closed field is very distinguished, using a result due to Kirby. This directly implies the same for \mathfrak{C}_{ab} and $\mathfrak{C}_{\text{tor}}$. We first prove the following theorem.

Theorem 9.2. *Let X be a semiabelian variety over \mathbb{C} and $V \subset X$ a subvariety. There is a finite set Σ of semiabelian subvarieties of X , depending only on X and V , such that for every $W \subset V$ that is weakly optimal for V in X there exist $Y \in \Sigma$ and $x \in W(\mathbb{C})$ with $\langle W \rangle_{\text{ws}} = xY$.*

Proof. We induct on the dimension of V , the case $\dim V = 0$ being trivial. Let $W \subset V$ be weakly optimal for V in X . We can assume without loss of generality that $W \neq V$. This implies that W is an atypical component of the intersection $\langle W \rangle_{\text{ws}} \cap V$ as defined in Définition 2.1 in [CL12].

By Théorème 2.3 in [CL12], due to Kirby in [Kir09], there is a finite set Φ of semiabelian subvarieties of X , depending only on X and V , such that $W \subset xY$ for some $Y \in \Phi$ and $x \in X(\mathbb{C})$ and furthermore

$$\dim Y + \dim W = \dim \langle W \rangle_{\text{ws}} + \dim(V \cap xY).$$

This is equivalent to

$$\delta_{\text{ws}}(W) = \dim Y - \dim(V \cap xY).$$

We can assume without loss of generality that $x \in W(\mathbb{C})$. We want to prove that $\langle W \rangle_{\text{ws}} = xY$.

Consider the projection $\pi : V \rightarrow VY/Y \subset X/Y$. We have $\pi^{-1}(\pi(v)) = V \cap vY$ for $v \in V(\mathbb{C})$. It follows from the Fiber Dimension Theorem that there exists a closed subset $V_1 \subsetneq V$, the preimage of the complement of the open subset of VY/Y from the theorem, such that $\pi^{-1}(\pi(V_1)) = V_1$ and $\pi^{-1}(\pi(v))$ is equidimensional for $v \in (V \setminus V_1)(\mathbb{C})$. Note that W is contained in a fiber of π .

If $W \subset V_1$ (or equivalently: if $x \in V_1(\mathbb{C})$), then W is weakly optimal for a component of V_1 in X and we are done by induction. Otherwise, let \tilde{W} be an irreducible component of $V \cap xY$ that contains W . As $V \cap xY = \pi^{-1}(\pi(x))$, it follows from the above that

$$\delta_{\text{ws}}(W) = \dim Y - \dim(V \cap xY) = \dim Y - \dim \tilde{W} \geq \delta_{\text{ws}}(\tilde{W}).$$

It follows from the weak optimality of W that $W = \tilde{W}$, so $\dim Y = \dim \langle W \rangle_{\text{ws}}$. As $\langle W \rangle_{\text{ws}} \subset xY$, we have $\langle W \rangle_{\text{ws}} = xY$. \square

Theorem 9.3. *The distinguished category of semiabelian varieties over an algebraically closed field K is very distinguished.*

Proof. Let $K \subset L$ be an extension of algebraically closed fields and let X be a semiabelian variety over K . There exists an algebraically closed subfield $K_0 \subset K$ and a semiabelian variety X_0 over K_0 such that $X = (X_0)_K$ and K_0 embeds into \mathbb{C} . We fix one such embedding. Let $V \subset X_L$ be a subvariety. Again, there exists an algebraically closed subfield $L_0 \subset L$, containing K_0 , and a subvariety V_0 of $(X_0)_{L_0}$ such that $V = (V_0)_L$ and there is an embedding of L_0 into \mathbb{C} that extends the one of K_0 . We also fix one such embedding. The theorem now follows from Theorem 9.2, applied to $(V_0)_{\mathbb{C}} \subset (X_0)_{\mathbb{C}}$, together with the two following facts:

- (1) All semiabelian subvarieties of $(X_0)_{\mathbb{C}}$ are base changes of semiabelian subvarieties of X_0 since the torsion points of any semiabelian variety are dense in it and are defined over the algebraic closure of its field of definition, and
- (2) if $W \subset V$ is a subvariety, then there exists an algebraically closed subfield $L_W \subset L$, containing L_0 , and a subvariety W_0 of $(V_0)_{L_W}$ such that $W = (W_0)_L$ and there is an embedding of L_W into \mathbb{C} that extends the one of L_0 . Furthermore, if $(W_0)_{\mathbb{C}} \subset x(Y_0)_{\mathbb{C}}$ for some $x \in X_0(\mathbb{C})$ and some semiabelian subvariety Y_0 of X_0 , then x can be chosen as the base change of an L_W -point of W_0 . In particular, weak optimality is preserved under base change. \square

We now want to apply a theorem of Gao in [Gao20] to see that the distinguished categories of connected pure Shimura varieties and connected mixed Shimura varieties of Kuga type over \mathbb{Q} are very distinguished. Unfortunately, the definition of a connected mixed Shimura datum of Kuga type and the definition of a connected pure Shimura datum in [Gao20] do not coincide with our definitions here (which are the same as in [Gao17b]). In fact, the definitions in [Gao20] give rise to full subcategories $\mathfrak{C}_{\text{mSvK}}^0$ and $\mathfrak{C}_{\text{pSv}}^0$ of $\mathfrak{C}_{\text{mSvK}}$ and $\mathfrak{C}_{\text{pSv}}$ respectively, where we consider only those mixed Shimura data that have generic Mumford-Tate group as defined in Definition 2.5 in [Gao17b]. By analogy, we also get a full subcategory $\mathfrak{C}_{\text{mSv}}^0$ of $\mathfrak{C}_{\text{mSv}}$. We do not know if these subcategories of distinguished categories are distinguished themselves, but we can still apply Definitions 6.1 and 6.3 to define (weakly) special subvarieties in them.

The following two lemmas will solve all arising technical issues. We continue identifying objects of $\mathfrak{C}_{\text{mSv}}$ (or of one of its subcategories) with their images under \mathcal{F}_{mSv} .

Lemma 9.4. *Let K be an algebraically closed field and let $\mathfrak{C}, \mathfrak{C}'$ be two categories among $(\mathfrak{C}_{\text{mSv}})_K, (\mathfrak{C}_{\text{mSvK}})_K, (\mathfrak{C}_{\text{pSv}})_K, (\mathfrak{C}_{\text{mSv}}^0)_K, (\mathfrak{C}_{\text{mSvK}}^0)_K, (\mathfrak{C}_{\text{pSv}}^0)_K$. Let $\mathfrak{Y}, \mathfrak{Y}'$ denote their respective classes of objects and suppose that $\mathfrak{Y}' \subset \mathfrak{Y}$. Let $X \in \mathfrak{Y}'$. Then the (weakly) special subvarieties of X as an object of \mathfrak{C}' are the same as the (weakly) special subvarieties of X as an object of \mathfrak{C} .*

Proof. It is sufficient to prove the lemma if either \mathfrak{C} is among the categories $(\mathfrak{C}_{\text{mSv}})_K, (\mathfrak{C}_{\text{mSvK}})_K, (\mathfrak{C}_{\text{pSv}})_K$ and $\mathfrak{C}' = \mathfrak{C}^0$ is the corresponding subcategory as defined above (case 1) or if $\mathfrak{C}, \mathfrak{C}'$ are both among the categories $(\mathfrak{C}_{\text{mSv}})_K, (\mathfrak{C}_{\text{mSvK}})_K, (\mathfrak{C}_{\text{pSv}})_K$ (case 2).

As every morphism of \mathfrak{C}' is also a morphism of \mathfrak{C} , it is clear that all (weakly) special subvarieties of X as an object of \mathfrak{C}' are also (weakly) special subvarieties of X as an object of \mathfrak{C} . What remains to be proved is the converse direction. Let $X \in \mathfrak{Y}'$, let $\phi : Y \rightarrow X, \psi : Y \rightarrow Z$ be morphisms of \mathfrak{C} , and let $z \in Z(K)$. We want to show that $\phi(Y)$ is a special subvariety of X as an object of \mathfrak{C}' and that every irreducible component W of $\phi(\psi^{-1}(z))$ is a weakly special subvariety of X as an object of \mathfrak{C}' .

In case 1, Proposition 2.6(1) in [Gao17b], Proposition 2.2 in [RS93], Lemma 2.2 in [Orr13] (which is formulated for reductive groups, but holds with the same proof for arbitrary linear algebraic groups), and the definition of having generic Mumford-Tate group yield (base changes of) surjective Shimura embeddings $\phi_Y : Y^0 \rightarrow Y, \psi_Z : Z^0 \rightarrow Z$ in \mathfrak{C} and a morphism $\psi^0 : Y^0 \rightarrow Z^0$ of \mathfrak{C}' such that $Y^0, Z^0 \in \mathfrak{Y}'$ and $\psi \circ \phi_Y = \psi_Z \circ \psi^0$. Hence, $\phi \circ \phi_Y$ is a morphism of \mathfrak{C}' and $\phi(Y) = (\phi \circ \phi_Y)(Y^0)$ is a special subvariety of X as an object of \mathfrak{C}' . Furthermore, we have

$$\phi(\psi^{-1}(z)) = (\phi \circ \phi_Y)((\psi \circ \phi_Y)^{-1}(z)) = \bigcup_{z^0 \in \psi_Z^{-1}(z)} (\phi \circ \phi_Y)((\psi^0)^{-1}(z^0)).$$

Since ψ_Z is a (base change of a) Shimura embedding, $\psi_Z^{-1}(z)$ is finite by Proposition 3.6 and the desired claim follows.

In case 2, note that, by Proposition 2.9 in [Gao17b], being pure and being of Kuga type are inherited by Shimura subdata. These properties are also inherited by quotient

Shimura data by Remark 3.7. The desired claim for special subvarieties then follows from Lemma 5.11 in [Gao17b].

After replacing K by an algebraically closed subfield of finite transcendence degree over $\bar{\mathbb{Q}}$ such that z is the base change of a point defined over that subfield, we can embed K into \mathbb{C} . By Proposition 5.4 in [Gao17b], we have that $W_{\mathbb{C}}$ is an irreducible component of $\phi'_{\mathbb{C}}((\psi'_{\mathbb{C}})^{-1}(z'))$, where $\phi' : Y' \rightarrow X$ and $\psi' : Y' \rightarrow Z'$ are (base changes of) a Shimura embedding and a quotient Shimura morphism respectively (in particular, ϕ', ψ' are morphisms of \mathfrak{C}' and ϕ' is finite by Proposition 3.6) and $z' \in Z'(\mathbb{C})$. It follows from the finiteness of ϕ' that $\{z'\} = \psi'_{\mathbb{C}}(\tilde{W})$ for some irreducible component \tilde{W} of $(\phi'_{\mathbb{C}})^{-1}(W_{\mathbb{C}})$. We therefore deduce that z' is the base change of a K -point of Z' and the desired claim follows. \square

Lemma 9.5. *Let K be an algebraically closed field, let \mathfrak{C} be one of the categories $(\mathfrak{C}_{mSv})_K$, $(\mathfrak{C}_{mSvK})_K$, $(\mathfrak{C}_{pSv})_K$, and let \mathfrak{C}^0 be the corresponding subcategory as defined above. Let $\mathfrak{X}, \mathfrak{X}^0$ denote the respective classes of objects of $\mathfrak{C}, \mathfrak{C}^0$. Let $X \in \mathfrak{X}$. Then there exists a (base change of a) surjective Shimura embedding $\phi : X^0 \rightarrow X$ such that $X^0 \in \mathfrak{X}^0$. If V is a subvariety of X and W is weakly optimal for V in X , then there exist irreducible components V^0, W^0 of $\phi^{-1}(V), \phi^{-1}(W)$ respectively such that $W^0 \subset V^0$, W^0 is weakly optimal for V^0 in X^0 , and $\langle W \rangle_{\text{ws}} = \phi(\langle W^0 \rangle_{\text{ws}})$.*

Thanks to Lemma 9.4, the operator $\langle \cdot \rangle_{\text{ws}}$ associated to X^0 as well as the concept of weak optimality for subvarieties of X^0 here do not depend on whether we regard X^0 as an object of \mathfrak{C}^0 or as an object of \mathfrak{C} .

Proof. The existence of ϕ follows from Proposition 2.6(1) in [Gao17b]. If $U^0 \subset X^0$ is any subvariety, then Lemma 6.7 implies that $\langle \phi(U^0) \rangle_{\text{ws}} \subset \phi(\langle U^0 \rangle_{\text{ws}})$ as well as $\langle U^0 \rangle_{\text{ws}} \subset \phi^{-1}(\langle \phi(U^0) \rangle_{\text{ws}})$. Since ϕ is surjective, we deduce that $\langle \phi(U^0) \rangle_{\text{ws}} = \phi(\langle U^0 \rangle_{\text{ws}})$. Since ϕ is also finite by Proposition 3.6, this implies that $\delta_{\text{ws}}(\phi(U^0)) = \delta_{\text{ws}}(U^0)$.

Since ϕ is surjective, we can now choose an irreducible component V^0 of $\phi^{-1}(V)$ such that $\phi(V^0) = V$. We can then choose an irreducible component W^0 of $\phi|_{V^0}^{-1}(W)$ such that $\phi(W^0) = W$. Since ϕ is finite, W^0 must be an irreducible component of $\phi^{-1}(W)$ as well. Together with the finiteness of ϕ , the previous paragraph implies that W^0 is weakly optimal for V^0 in X^0 and $\langle W \rangle_{\text{ws}} = \langle \phi(W^0) \rangle_{\text{ws}} = \phi(\langle W^0 \rangle_{\text{ws}})$. \square

We can now prove the following theorem:

Theorem 9.6 (Gao). *The distinguished categories of connected pure Shimura varieties and connected mixed Shimura varieties of Kuga type over $\bar{\mathbb{Q}}$ are very distinguished.*

For pure Shimura varieties, this was essentially already established in [DR18], later refined in [BD21]. With the new results obtained independently by Chiu [Chi21] and Gao-Klingler [GK21], we expect that the proof of Theorem 9.6 can be extended to show the same for the distinguished category of connected mixed Shimura varieties.

Proof. Note that, by Proposition 2.9 in [Gao17b], being pure and being of Kuga type are inherited by Shimura subdata. These properties are also inherited by quotient Shimura data by Remark 3.7. The required finiteness statement over \mathbb{C} then follows from Theorem 8.2 in [Gao20] together with Remark 5.3 and Lemma 5.6 in [Gao17b], as well as Lemma 9.5.

We can go from \mathbb{C} to an arbitrary algebraically closed field of finite transcendence degree over $\bar{\mathbb{Q}}$ thanks to the following: Let X be a connected mixed Shimura variety. Let $U \subset W \subset X_{\mathbb{C}}$ be a chain of subvarieties, where W is weakly special. By Proposition 5.4 in [Gao17b], we have that W is an irreducible component of $\phi_{\mathbb{C}}(\psi_{\mathbb{C}}^{-1}(z))$, where $\phi : Y \rightarrow X$ and $\psi : Y \rightarrow Z$ are a Shimura embedding and a quotient Shimura morphism respectively (in particular, ϕ is finite by Proposition 3.6) and $z \in Z(\mathbb{C})$. It follows from the finiteness

of ϕ that $\{z\} = \psi_{\mathbb{C}}(\tilde{U})$ for some irreducible component \tilde{U} of $\phi_{\mathbb{C}}^{-1}(U)$. If U can be defined over some algebraically closed subfield of \mathbb{C} , we can therefore deduce the same for z . This implies that weak optimality is preserved under base change. The theorem then also follows since Theorem 8.2 in [Gao20] yields precisely such pairs of Shimura embeddings and quotient Shimura morphisms. \square

10. ZILBER-PINK AND EXTENSIONS OF ALGEBRAICALLY CLOSED FIELDS

We now introduce a notation that allows us to easily express the fact that the analogue of the Zilber-Pink conjecture holds in a certain setting.

Definition 10.1. *Let K be an algebraically closed field and let \mathfrak{C} be a distinguished category over K . Let m, d be non-negative integers and let X be a distinguished variety. We say that $\text{ZP}(X, m, d)$ holds if every subvariety of X of dimension at most m contains at most finitely many optimal subvarieties of defect at most d .*

The distinguished category $\mathfrak{C}_{\text{triv}}(Y)$ (for a fixed variety Y over K) trivially satisfies $\text{ZP}(X, m, d)$ for all non-negative integers m and d and all distinguished varieties X since every distinguished variety in $\mathfrak{C}_{\text{triv}}(Y)$ has only finitely many special subvarieties and an optimal subvariety for a subvariety V in X must be an irreducible component of the intersection of V and some special subvariety of X .

We collect some auxiliary results that are going to be useful later: The following lemma is Lemma 2.2 in [BD19].

Lemma 10.2. *Let $K \subset L$ be an extension of algebraically closed fields such that L has transcendence degree 1 over K . Let V be a variety over K and let W be a subvariety of V_L . Then there exists a subvariety W' of V with $\dim W' \leq \dim W + 1$ such that $W \subset W'_L$.*

The following lemma is Lemma 2.5 in [BD19].

Lemma 10.3. *Let K be an algebraically closed field and let $f : V \rightarrow W$ be a dominant morphism of algebraic varieties, defined over K . Then there exists $V_0 \subset V$ open and dense such that $f(V_0)$ is open and dense in W and $f|_{V_0} : V_0 \rightarrow f(V_0)$ is smooth.*

The following lemma is the analogue of Lemma 2.11 in [BD19] in our setting.

Lemma 10.4. *Let $K \subset L$ be an extension of algebraically closed fields. Let \mathfrak{C} be a distinguished category over K , let X be a distinguished variety, and let V be a subvariety of X . If W is an optimal subvariety for V_L in X_L , then there exists an optimal subvariety W' for V in X such that $W = (W')_L$ and $\delta(W) = \delta(W')$.*

Proof. Since a special subvariety is the image of a distinguished morphism, we have that any special subvariety of X_L is the base change of a special subvariety of X . Therefore, if V is a subvariety of X , any optimal subvariety for V_L in X_L is an irreducible component of an intersection $V_L \cap S_L$ for some special subvariety S of X and is then the base change of a subvariety $W' \subset V$ that must be optimal for V in X and of the same defect. \square

The following lemma is an analogue of Lemma 2.6(2) in [BD19].

Lemma 10.5. *Let m and d be non-negative integers. Let K be an algebraically closed field and let \mathfrak{C} be a distinguished category over K . Let $\phi : X \rightarrow Y$ be a distinguished morphism with finite fibers. Then $\text{ZP}(Y, m, d) \Rightarrow \text{ZP}(X, m, d)$.*

Proof. We induct on m , the case $m = 0$ being trivial. Let $V \subset X$ be a subvariety of dimension m . Let V' be the closure of $\phi(V)$. By Lemma 10.3 and the Fiber Dimension Theorem, we can find $V_0 \subset V$ open and dense such that $\phi(V_0)$ is open and dense in V' and $\phi|_{V_0} : V_0 \rightarrow \phi(V_0)$ is smooth of relative dimension $\dim V_0 - \dim \phi(V_0)$, which must be 0 as ϕ has finite fibers.

Let now $W \subset V$ be an optimal subvariety for V in X whose defect is at most d . If $W \cap V_0 = \emptyset$, then W is optimal for a component of $V \setminus V_0$ and we are done by induction on m . Otherwise we can apply Lemma 8.3(1) to deduce that the closure W' of $\phi(W)$ is optimal for V' in Y and the defect of W' is at most d . It follows from $\text{ZP}(Y, m, d)$ that there are at most finitely many possibilities for W' and hence for W , which is an irreducible component of $\phi^{-1}(W')$. \square

Lemma 10.6. *Let m and d be non-negative integers. Let K be an algebraically closed field and let \mathfrak{C} be a distinguished category over K . Let $\phi : X \rightarrow Y$ be a surjective distinguished morphism with finite fibers. Then $\text{ZP}(X, m, d) \Leftrightarrow \text{ZP}(Y, m, d)$.*

Proof. Thanks to Lemma 10.5, it suffices to prove the implication “ \Rightarrow ”.

We induct on m , the case $m = 0$ being trivial. Let $V \subset Y$ be a subvariety of dimension m . Let \tilde{V} be an irreducible component of $\phi^{-1}(V)$ that dominates V . We have $\dim \tilde{V} = \dim V \leq m$. By Lemma 10.3 and the Fiber Dimension Theorem, we can find $\tilde{V}_0 \subset \tilde{V}$ open and dense such that $\phi(\tilde{V}_0)$ is open and dense in V and $\phi|_{\tilde{V}_0} : \tilde{V}_0 \rightarrow \phi(\tilde{V}_0)$ is smooth of relative dimension $\dim \tilde{V}_0 - \dim \phi(\tilde{V}_0) = 0$.

Let now $W \subset V$ be an optimal subvariety for V in Y whose defect is at most d . If $W \cap \phi(\tilde{V}_0) = \emptyset$, then W is optimal for a component of $V \setminus \phi(\tilde{V}_0)$ and we are done by induction on m . Otherwise $W \cap \phi(\tilde{V}_0)$ is dense in W . Let \tilde{W} be the closure in \tilde{V} of a component of $\phi|_{\tilde{V}_0}^{-1}(W)$ that dominates W . By Lemma 8.3(2), we have $\delta(\tilde{W}) = \delta(W) \leq d$.

Let \tilde{U} be a subvariety of \tilde{V} such that $\tilde{W} \subset \tilde{U} \subset \tilde{V}$, \tilde{U} is optimal for \tilde{V} in X , and $\delta(\tilde{U}) \leq \delta(\tilde{W})$. We have $\tilde{U} \cap \tilde{V}_0 \neq \emptyset$ since $\tilde{W} \cap \tilde{V}_0 \neq \emptyset$. By Lemma 8.3, the closure U of $\phi(\tilde{U})$ is optimal for V in Y of defect

$$\delta(U) = \delta(\tilde{U}) \leq \delta(\tilde{W}) = \delta(W).$$

Furthermore, we have $W \subset U \subset V$. Since W is optimal for V in Y , it follows that $U = W$. Hence $\tilde{W} = \tilde{U}$ is optimal for \tilde{V} in X and its defect is at most d . By $\text{ZP}(X, m, d)$, there are at most finitely many possibilities for \tilde{W} and hence also for W . \square

The following theorem, which is one of our main results, is an analogue of Proposition 4.1 in [BD19]. In its statement, we use the notation from Definition 10.1.

Theorem 10.7. *Let m and d be non-negative integers. Let $K \subset L$ be an extension of algebraically closed fields and let \mathfrak{C} be a very distinguished category over K . Let X be a distinguished variety. Let \mathcal{S} be a class of distinguished varieties such that $X \in \mathcal{S}$ and for every $X' \in \mathcal{S}$, every field extension $K \subset K'$ of finite transcendence degree, and every subvariety V of $X'_{K'}$, the set in (A5) can be chosen with all Y_ϕ, Z_ψ equal to base changes of distinguished varieties belonging to \mathcal{S} . Then the following hold:*

- (1) *Suppose that $\text{ZP}(X', m, d)$ holds for all $X' \in \mathcal{S}$. Then $\text{ZP}(X_L, m, d)$ holds as well.*
- (2) *Suppose that $\text{ZP}(X', m - 1, d)$ and $\text{ZP}(X', m, d - 1)$ hold for all $X' \in \mathcal{S}$. Let V be a subvariety of X_L of dimension at most m which is not the base change of a subvariety of X (in particular $K \neq L$). Then V contains at most finitely many optimal subvarieties of defect at most d .*

It is of course possible to take the whole class of objects of \mathfrak{C} as \mathcal{S} , but for applications, it can be useful to take \mathcal{S} smaller as we will see later.

Proof. We prove (1) and (2) in parallel: Let V be a subvariety of X_L . We can find an algebraically closed subfield L_1 of L that has finite transcendence degree over K and a subvariety V_1 of X_{L_1} such that $V = (V_1)_L$. If W is any optimal subvariety for V in X_L , then by Lemma 10.4 it is equal to $(W_1)_L$ for an optimal subvariety W_1 for V_1 in X_{L_1} such that $\delta(W_1) = \delta(W)$. Hence, it suffices to prove the theorem under the assumption that L has finite transcendence degree over K .

We will actually prove the stronger statement that the conclusions of (1) and (2) hold with any $X' \in \mathcal{S}$ in place of X . Arguing by induction on the transcendence degree of L over K , one can see that it is enough to prove (1) and (2) when L has transcendence degree 1 over K .

We proceed by induction on m . Clearly $\text{ZP}(X_L, 0, d)$ holds for all d . Let V be a subvariety of X_L of dimension $m > 0$. We will deduce that V contains at most finitely many optimal subvarieties of defect at most d from either of the following hypotheses:

- (1) $\text{ZP}(X'_L, m - 1, d)$ and $\text{ZP}(X', m, d)$ hold for all $X' \in \mathcal{S}$, or
- (2) $\text{ZP}(X'_L, m - 1, d)$ and $\text{ZP}(X', m, d - 1)$ hold for all $X' \in \mathcal{S}$ and V is not the base change of a subvariety of X .

As all $X' \in \mathcal{S}$ satisfy the same hypotheses as X , it will follow that the conclusions of (1) and (2) hold for all $X' \in \mathcal{S}$ as desired.

If $V = V'_L$ for some $V' \subset X$, then we must be in case (1) and are done by Lemma 10.4 and $\text{ZP}(X, m, d)$. We will then assume that this is not the case.

Let V'_L be the smallest subvariety of X_L that is the base change of some $V' \subset X$ and contains V . It exists and has dimension m or $m + 1$ by Lemma 10.2 but the first case is not possible because it would imply that $V = V'_L$.

Let $W \subset V$ be an optimal subvariety for V in X_L that has defect at most d . We can assume without loss of generality that $W \neq V$.

We let W'_L be the smallest subvariety of X_L that is the base change of some $W' \subset X$ and contains W . By Lemma 10.2, we have either $W = W'_L$ or $\dim W'_L = \dim W + 1$.

If $W = W'_L$, then W is contained in $Z'_L \subset V$ for $Z' \subset X$ maximal among all finite unions of subvarieties $Z'' \subset X$ with $Z''_L \subset V$ (and equal to the closure of the union of all such Z''). Since $V \neq V'_L$, the dimension of Z'_L is at most $\dim V - 1$. Of course, W is also optimal for the component of Z'_L that contains it. It follows that W lies in a finite set because $\text{ZP}(X_L, m - 1, d)$ holds. We can therefore assume that $W \subsetneq W'_L$ and so $\dim W'_L = \dim W + 1$.

Recall that, by Lemma 10.4, an optimal subvariety for V'_L in X_L is the base change of an optimal subvariety for V' in X . Let U'_L be such an optimal subvariety for V'_L in X_L that contains W'_L and satisfies $\delta(U'_L) \leq \delta(W'_L)$. Note that $\langle V \rangle = \langle V'_L \rangle$ and $\langle W \rangle = \langle W'_L \rangle$ because, for instance, $V'_L \subset \langle V \rangle \cap V'_L$ by definition. It follows that $\delta(W'_L) = \delta(W) - 1$, so $\delta(U'_L) \leq d - 1$.

We claim that $U'_L \neq V'_L$. If not, we could deduce that $\delta(V'_L) = \delta(U'_L) \leq \delta(W'_L)$. It would then follow that $\delta(V) = \delta(V'_L) + 1 \leq \delta(W'_L) + 1 = \delta(W)$, which contradicts the optimality of $W \subsetneq V$ for V .

We deduce that $U'_L \subsetneq V'_L$ and hence $U'_L \cap V \subsetneq V$, otherwise $U'_L \supset V$ would contradict the minimality of V'_L . Since $W \subset U'_L \cap V$ and W has defect at most d and is optimal for a component of $U'_L \cap V$ in X_L , it suffices to show that U' and therefore U'_L belongs to a finite set. Then we are done as any component of $U'_L \cap V$ that contains W has dimension at most $m - 1$ and $\text{ZP}(X_L, m - 1, d)$ holds.

By the optimality of U' for V' in X and Proposition 8.2, we have that U' is also weakly optimal for V' in X . It follows from (A5) that there exists a finite set of pairs (ϕ, ψ) of distinguished morphisms $\phi : Y_\phi \rightarrow X$ and $\psi : Y_\phi \rightarrow Z_\psi$, depending only on V' , such that there exists a pair (ϕ, ψ) in this set and $z \in Z_\psi(K)$ such that ϕ has finite fibers and $\langle U' \rangle_{\text{ws}}$ is an irreducible component of $\phi(\psi^{-1}(z))$. By our hypothesis, we can assume that $Y_\phi, Z_\psi \in \mathcal{S}$. Since ϕ and ψ vary in a finite set, we can assume them fixed.

Let \hat{U} be an irreducible component of $\psi^{-1}(z) \cap \phi^{-1}(U')$ such that $\phi(\hat{U})$ is dense in U' . Since ϕ has finite fibers, we deduce that $\dim \hat{U} = \dim U'$ and therefore \hat{U} must be an irreducible component of $\phi^{-1}(U')$. Let \hat{V} be an irreducible component of $\phi^{-1}(V')$ that contains \hat{U} . As ϕ has finite fibers, it follows that $\dim \hat{V} \leq \dim V' \leq m + 1$.

Furthermore, \hat{U} is optimal for \hat{V} : Otherwise, we could find \tilde{U} such that $\hat{U} \subsetneq \tilde{U} \subset \hat{V}$ and $\delta(\tilde{U}) \leq \delta(\hat{U})$. Applying ϕ and taking closures yields a contradiction to the optimality of U' for V' thanks to Lemma 8.3(2). The same lemma shows that $\delta(\hat{U}) = \delta(U')$.

We have $\hat{U} \subset \psi^{-1}(z) \cap \hat{V}$ by construction. If \hat{U} were not equal to an irreducible component of this intersection, then we could find a subvariety \tilde{U} such that $\hat{U} \subsetneq \tilde{U} \subset \psi^{-1}(z) \cap \hat{V}$. Since ϕ has finite fibers, it would follow that the closure of $\phi(\tilde{U})$ strictly contains U' , while still being contained in $\phi(\psi^{-1}(z))$ as well as in V' .

Since every irreducible component of $\phi(\psi^{-1}(z))$ has the same dimension by (A4) and the fact that ϕ has finite fibers, this contradicts the fact that U' is weakly optimal for V' . It follows that \hat{U} is an irreducible component of $\psi^{-1}(z) \cap \hat{V}$.

We get a subvariety V'' of Z_ψ , equal to the closure of $\psi(\hat{V})$. Furthermore, we know that $\{z\} = \psi(\hat{U})$.

By Lemma 10.3 and the Fiber Dimension Theorem, we can find $\hat{V}_0 \subset \hat{V}$ open and dense such that $\psi(\hat{V}_0)$ is open and dense in V'' and $\psi|_{\hat{V}_0} : \hat{V}_0 \rightarrow \psi(\hat{V}_0)$ is smooth of relative dimension $n = \dim \hat{V}_0 - \dim \psi(\hat{V}_0) = \dim \hat{V} - \dim V''$. If \hat{U} is contained in $\hat{V} \setminus \hat{V}_0$, then \hat{U} is contained in one of finitely many subvarieties of \hat{V} of dimension at most m . As \hat{U} is optimal for \hat{V} , it is also optimal for that subvariety. We have $\delta(\hat{U}) = \delta(U') \leq \delta(W') = \delta(W) - 1 \leq d - 1$.

Since $Y_\phi \in \mathcal{S}$, $\text{ZP}(Y_\phi, m, d - 1)$ holds. It follows that there are then only finitely many possibilities for \hat{U} and hence for U' .

Hence we can assume that $\hat{U} \cap \hat{V}_0 \neq \emptyset$. Since \hat{U} is an irreducible component of $\psi^{-1}(z) \cap \hat{V} = \psi|_{\hat{V}}^{-1}(z)$, $\hat{U} \cap \hat{V}_0$ is then an irreducible component of $\psi|_{\hat{V}_0}^{-1}(z)$. It follows that $n = \dim(\hat{U} \cap \hat{V}_0) = \dim \hat{U}$.

Since we know that $W \neq W'_L$, we have $\dim W' > 0$ and hence $n = \dim \hat{U} = \dim U' > 0$.

The special subvariety $\langle \hat{U} \rangle$ contains $\langle \hat{U} \rangle_{\text{ws}}$. By construction, $\langle \hat{U} \rangle_{\text{ws}}$ is contained in an irreducible component of $\psi^{-1}(z)$. As $\langle U' \rangle_{\text{ws}}$ is equal to an irreducible component of $\phi(\psi^{-1}(z))$, $\langle \hat{U} \rangle_{\text{ws}}$ must actually be equal to a component of $\psi^{-1}(z)$: Otherwise, it would follow for dimension reasons that $U' \subset \phi(\langle \hat{U} \rangle_{\text{ws}}) \cap \langle U' \rangle_{\text{ws}} \subsetneq \langle U' \rangle_{\text{ws}}$ since ϕ has finite fibers, a contradiction with Lemmas 6.7 and 6.8.

Since \hat{U} is optimal for \hat{V} and has defect at most $d - 1$, it then follows from Lemma 8.3(1) that $\{z\} = \psi(\hat{U})$ is optimal for V'' in Z_ψ and has defect at most $d - 1$. Furthermore, V'' is a subvariety of Z_ψ of dimension $\dim \hat{V} - n \leq \dim \hat{V} - 1 \leq \dim V' - 1 = m$.

As Z_ψ lies in \mathcal{S} , $\text{ZP}(Z_\psi, m, d - 1)$ holds. Thus, z lies in a finite set. As \hat{U} is a component of $\psi^{-1}(z) \cap \hat{V}$, it lies in a finite set as well and therefore the same holds for U' . \square

We will now apply Theorem 10.7(2) to certain connected mixed Shimura varieties of Kuga type. For this, we first prove a lemma:

Lemma 10.8. *Let K be an algebraically closed field and let \mathfrak{C} be a distinguished category over K . Let X be a distinguished variety and let $\phi : X \rightarrow Y$ be a surjective distinguished morphism. If $\text{ZP}(X, m, 0)$ holds for all non-negative integers m , then the same holds for Y .*

Proof. Let $V \subset Y$ be a subvariety. If $S \subset V$ is optimal for V in Y and $\delta(S) = 0$, then S is a maximal special subvariety of V . The lemma now follows from the fact that any irreducible component of $\phi^{-1}(S)$ is a maximal special subvariety of some irreducible component of $\phi^{-1}(V)$. \square

In the following corollary, \mathbb{H}_g denotes the Siegel upper half space of dimension $\frac{g(g+1)}{2}$ and $(\text{GSp}_{2g}, \mathbb{H}_g)$ denotes the connected pure Shimura datum associated to the moduli space of principally polarized abelian varieties of dimension g .

Corollary 10.9. *Let $X = \mathcal{F}_{\text{mSvK}}(P, X^+, \Gamma)$ be a connected mixed Shimura variety of Kuga type of dimension 3. Let W be the unipotent radical of P and let $(P, X^+)/W$ denote the quotient Shimura datum (see [Pin90], Proposition 2.9). Suppose that $(P, X^+)/W$ is a Shimura subdatum of $(\text{GSp}_{2g}, \mathbb{H}_g)$ for some $g \in \mathbb{N}$. Let $\bar{\mathbb{Q}} \subset K$ be an extension of algebraically closed fields and let $C \subset X_K$ be a curve that is not the base change of a curve in X . Then C contains at most finitely many optimal subvarieties.*

Corollary 10.9 applies for example to the moduli space of principally polarized abelian surfaces or to the (direct) product of the Legendre family of elliptic curves (see Section 13) with the moduli space of elliptic curves. It also applies to the cube of the moduli space of elliptic curves, for which the corresponding statement has been proven by Pila in Theorem 1.4 in [Pil17].

Proof. As $\delta(C) \leq 2$, any proper optimal subvariety of C has defect at most 1. By Theorem 9.6, the category of connected mixed Shimura varieties of Kuga type is very distinguished.

We want to apply Theorem 10.7(2) with $m = d = 1$, choosing the class \mathcal{S} there as small as possible. Note that $\text{ZP}(X', 0, 1)$ is trivially satisfied for every $X' \in \mathcal{S}$ while $\text{ZP}(X', 1, 0)$ is the André-Oort conjecture for curves in X' . We have $\text{ZP}(X, m, 0)$ for all non-negative integers m thanks to Theorem 5.2 in [Tsi18] and Theorem 13.6 in [Gao17b].

Theorem 8.2 in [Gao20] and the proof of Theorem 9.6 show that we first have to include in \mathcal{S} all triples $((Q, Y^+)/N, *)$, where (Q, Y^+) is a Shimura subdatum of (P, X^+) , N is a normal algebraic subgroup of Q whose reductive part is semisimple, and $(Q, Y^+)/N$ denotes the quotient Shimura datum (see [Pin90], Proposition 2.9). This step must then be iterated for every connected mixed Shimura datum thus obtained.

By Lemma 10.5, $\text{ZP}(*, m, 0)$ is preserved when passing to a Shimura subdatum (by Lemma 10.6, the validity of $\text{ZP}(*, m, d)$ is independent from the choice of congruence subgroup). If we have $\text{ZP}(*, m, 0)$ for all non-negative integers m , this is also preserved when passing to a quotient Shimura datum by Lemma 10.8. Hence, we have $\text{ZP}(X', 1, 0)$ for all $X' \in \mathcal{S}$ as desired. \square

11. REDUCTION TO OPTIMAL SINGLETONS

The following theorem is an analogue of Theorem 8.3 in [DR18] and Theorem 6.1 in [BD19] (see also [HP16]) and is similar to what was recently done by Cassani in [Cas21b] for connected mixed Shimura varieties of Kuga type.

Theorem 11.1. *Let m and d be non-negative integers. Let K be an algebraically closed field and let \mathfrak{C} be a distinguished category over K that satisfies (A5). Let X be a distinguished variety and let \mathcal{S} be a class of distinguished varieties such that $X \in \mathcal{S}$ and for every $X' \in \mathcal{S}$ and every subvariety of X' the set in (A5) can be chosen with all $Y_\phi, Z_\psi \in \mathcal{S}$. Suppose that for all $X' \in \mathcal{S}$ every subvariety of X' of dimension at most m contains at most finitely many optimal singletons of defect at most d . Then $\text{ZP}(X, m, d)$ holds.*

Proof. We proceed by induction on m . Clearly $\text{ZP}(X, 0, d)$ holds. We are going to deduce $\text{ZP}(X, m, d)$ from $\text{ZP}(X, m-1, d)$ and the fact that all subvarieties of dimension at most m of any $X' \in \mathcal{S}$ contain at most finitely many optimal singletons of defect at most d .

We fix a subvariety V of X with $\dim V \leq m$ and a subvariety $W \subset V$ optimal for V in X with $\delta(W) \leq d$.

By Proposition 8.2, W is weakly optimal for V in X and (A5) implies that there is a finite set of pairs (ϕ, ψ) of distinguished morphisms $\phi : Y_\phi \rightarrow X$ and $\psi : Y_\phi \rightarrow Z_\psi$, depending only on V , such that there exists a pair (ϕ, ψ) in this set and $z \in Z_\psi(K)$ such that ϕ has finite fibers and $\langle W \rangle_{\text{ws}}$ is an irreducible component of $\phi(\psi^{-1}(z))$. By our hypothesis, we can assume that $Y_\phi, Z_\psi \in \mathcal{S}$. As the pair (ϕ, ψ) varies in a finite set, we can assume it is fixed.

We now set \hat{W} to be an irreducible component of $\psi^{-1}(z) \cap \phi^{-1}(W)$ such that $\phi(\hat{W})$ is dense in W and \hat{V} to be an irreducible component of $\phi^{-1}(V)$ that contains \hat{W} . We recall that ϕ has finite fibers and thus $\dim \hat{W} = \dim W$ and $\dim \hat{V} \leq \dim V$. Moreover, Lemma 8.3(2) implies that $\delta(\hat{W}) = \delta(W)$.

We claim that \hat{W} is optimal for \hat{V} in Y_ϕ . If this were not the case, there would exist a $\hat{U} \supsetneq \hat{W}$ optimal for \hat{V} and with $\delta(\hat{U}) \leq \delta(\hat{W})$. Using Lemma 8.3(2), we see that this would contradict the optimality of W for V .

By definition we have that \hat{W} is contained in an irreducible component of $\psi^{-1}(z) \cap \hat{V}$. If this containment were strict, we could find a subvariety \hat{U} with $\hat{W} \subsetneq \hat{U} \subset \psi^{-1}(z) \cap \hat{V}$ but then, as ϕ has finite fibers, W would be strictly contained in the Zariski closure of $\phi(\hat{U}) \subset \phi(\psi^{-1}(z)) \cap V$. This contradicts the weak optimality of W because every irreducible component of $\phi(\psi^{-1}(z))$ has the same dimension. Thus \hat{W} has to be an irreducible component of $\psi^{-1}(z) \cap \hat{V}$.

We now set $\tilde{V} \subset Z_\psi$ to be the Zariski closure of $\psi(\hat{V})$. Moreover, recall that $\psi(\hat{W}) = \{z\}$. By Lemma 10.3 and the Fiber Dimension Theorem, we can find $\hat{V}_0 \subset \hat{V}$ open and dense such that $\psi(\hat{V}_0)$ is open and dense in \tilde{V} and $\psi|_{\hat{V}_0} : \hat{V}_0 \rightarrow \psi(\hat{V}_0)$ is smooth of relative dimension $n = \dim \hat{V}_0 - \dim \psi(\hat{V}_0) = \dim \hat{V} - \dim \tilde{V}$. If $\hat{W} \subset \hat{V} \setminus \hat{V}_0$, then \hat{W} is contained in one of finitely many proper subvarieties of \hat{V} and is optimal for that subvariety. By Lemma 10.5 and $\text{ZP}(X, m-1, d)$, after recalling that $\delta(\hat{W}) \leq d$, we have at most finitely many possibilities for \hat{W} and therefore at most finitely many possibilities for W .

We are then allowed to assume that $\hat{W} \cap \hat{V}_0 \neq \emptyset$. Since \hat{W} is an irreducible component of $\psi|_{\hat{V}_0}^{-1}(z)$, we must have that $\hat{W} \cap \hat{V}_0$ is an irreducible component of $\psi|_{\hat{V}_0}^{-1}(z)$ and thus $n = \dim(\hat{W} \cap \hat{V}_0) = \dim \hat{W} = \dim W$. Note that $\dim \tilde{V} = \dim \hat{V} - n \leq \dim V - n \leq m$.

We now want to apply Lemma 8.3(1) to ψ and $\hat{W} \subset \hat{V} \subset Y_\phi$. For this we need to verify that $\langle \hat{W} \rangle$ contains an irreducible component of a fiber of ψ . This fiber can be taken to be nothing but $\psi^{-1}(z)$: we have that $\langle \hat{W} \rangle_{\text{ws}} \subset \langle \hat{W} \rangle$ is contained in an irreducible component of $\psi^{-1}(z)$ but if this containment were strict, since $\langle W \rangle_{\text{ws}}$ is an irreducible component of $\phi(\psi^{-1}(z))$ and ϕ has finite fibers, we would have that $W \subset \phi(\langle \hat{W} \rangle_{\text{ws}}) \cap \langle W \rangle_{\text{ws}} \subsetneq \langle W \rangle_{\text{ws}}$. By Lemmas 6.7 and 6.8, every irreducible component of $\phi(\langle \hat{W} \rangle_{\text{ws}}) \cap \langle W \rangle_{\text{ws}}$ is weakly special, a contradiction.

We can then apply Lemma 8.3(1) and obtain that $\{z\} = \psi(\hat{W})$ is optimal for \tilde{V} in Z_ψ with $\delta(\{z\}) \leq \delta(\hat{W}) \leq d$. By our hypothesis, z lies in a finite set.

Finally, recall that \hat{W} is a component of $\psi^{-1}(z) \cap \hat{V}$, thus lies in a finite set and the same holds for W . \square

12. REDUCTION OF ZILBER'S FORMULATION TO PINK'S FORMULATION

In this section, we introduce two different formulations of the Zilber-Pink conjecture for a distinguished category and show that they are equivalent in the presence of (A5).

Conjecture 12.1. *Let K be an algebraically closed field and let \mathfrak{C} be a distinguished category over K . For every distinguished variety X and all non-negative integers m and d , $\text{ZP}(X, m, d)$ holds.*

This is Zilber's version of the conjecture (see [Zil02]), in the formulation by Habegger and Pila in [HP16] (see also there for a form of the conjecture that is closer to Zilber's original conjecture; their proof of the equivalence of the two versions works in any distinguished category). The following is a version that corresponds to Pink's Conjecture 1.2 in [Pin05b]. To formulate it, we introduce the following notation: If X is a distinguished variety and $k \in \{0, \dots, \dim X\}$, then $X^{[k]}$ denotes the union of all special subvarieties of X of codimension at least k .

Conjecture 12.2. *Let K be an algebraically closed field, let \mathfrak{C} be a distinguished category over K , and let m and d be non-negative integers. Let X be a distinguished variety and $V \subset X$ a subvariety of dimension at most m . Then $V \cap X^{[\dim X - \min\{\delta(V) - 1, d\}]}$ is not Zariski dense in V .*

We point out that Conjectures 12.1 and 12.2 certainly do not hold in an arbitrary distinguished category. Indeed, in $\mathfrak{C}_{\text{add}}$, we can consider $X = \mathbb{G}_{a,K}^4$ with affine coordinates x, y, z, w and $V \subset X$ defined by $x^2 + y^2 = z^2$, $x^3 + y^3 = w^3$. Then V is covered by lines passing through the origin that are all optimal for V in X , which contradicts Conjecture 12.1. Furthermore, any such line is of dimension $1 = \delta(V) - 1$, so we contradict Conjecture 12.2 as well.

Lemma 12.3. *Let K be an algebraically closed field, let \mathfrak{C} be a distinguished category over K , let X be a distinguished variety, and let m and d be non-negative integers. Then $\text{ZP}(X, m, d)$ implies Conjecture 12.2 for the given \mathfrak{C} , X , m , and d .*

Proof. Let X be a distinguished variety and $V \subset X$ a subvariety of dimension at most m . Let W be an irreducible component of the intersection of V with a special subvariety of codimension at least $\dim X - \min\{\delta(V) - 1, d\}$. Then

$$\delta(W) \leq \min\{\delta(V) - 1, d\} - \dim W \leq \min\{\delta(V) - 1, d\}.$$

There is a subvariety $U \subset V$, optimal for V in X , such that $W \subset U$ and $\delta(U) \leq \delta(W) \leq d$. As $\delta(W) < \delta(V)$, we have $U \neq V$. So W is contained in the union of all proper optimal subvarieties for V in X of defect at most d , which by $\text{ZP}(X, m, d)$ is a finite union.

We deduce that $V \cap X^{[\dim X - \min\{\delta(V) - 1, d\}]}$ is not dense in V . \square

The following theorem is an analogue of Theorem 1.9 in [BD19]. A similar statement for connected mixed Shimura varieties of Kuga type has recently been established by Cassani in [Cas21b].

Theorem 12.4. *Let K be an algebraically closed field, let \mathfrak{C} be a distinguished category over K that satisfies (A5), and let m and d be non-negative integers. Then Conjectures 12.1 and 12.2 for \mathfrak{C} , m , and d are equivalent.*

Proof. By Lemma 12.3, it suffices to show that Conjecture 12.2 for \mathfrak{C} , m , and d implies $\text{ZP}(X, m, d)$ for every distinguished variety X . By Theorem 11.1, it is then enough to show that every subvariety of dimension at most m of a distinguished variety contains at most finitely many optimal singletons of defect at most d .

For this, we follow the proof of Theorem 1.9 in [BD19]: Let X be a distinguished variety and let V be a subvariety of X of dimension at most m . We show the following claim by induction on $j \in \{0, \dots, \dim V\}$:

Claim. The optimal singletons for V in X of defect at most d are contained in a finite union of subvarieties of V of dimension at most $\dim V - j$.

This is obvious for $j = 0$.

Suppose that the claim has been proven for some $j < \dim V$. Let W be one of the finitely many subvarieties of V of dimension at most $\dim V - j$ that contain the optimal singletons for V in X of defect at most d . We can assume without loss of generality that $\dim W = \dim V - j$.

If $\{p\} \subset W$ is an optimal singleton for W in X of defect at most d , then $\langle\{p\}\rangle \subset \langle W \rangle$. Since $\dim W = \dim V - j > 0$, we have that $\{p\} \subsetneq W$ and therefore $\delta(\{p\}) = \dim \langle\{p\}\rangle < \delta(W)$. It follows that the codimension of $\langle\{p\}\rangle$ is greater than or equal to $k := \dim X - \min\{\delta(W) - 1, d\}$. So the optimal singletons for W in X of defect at most d are contained in $W \cap X^{[k]}$.

It then follows from Conjecture 12.2 for X , d , and W that $W \cap X^{[k]}$ is not dense in W . Together with the above, this implies that the optimal singletons for W in X of defect at most d are contained in a proper closed subset of W as desired. This establishes the claim by induction.

Now taking $j = \dim V$ in the claim shows that the number of optimal singletons for V in X of defect at most d is finite. \square

13. ZILBER-PINK FOR A COMPLEX CURVE IN A FIBERED POWER OF THE LEGENDRE FAMILY

Let $Y(2) = \mathbb{A}_{\mathbb{Q}}^1 \setminus \{0, 1\}$. Let \mathcal{E} denote the Legendre family of elliptic curves, defined in $Y(2) \times_{\mathbb{Q}} \mathbb{P}_{\mathbb{Q}}^2$ by $Y^2Z = X(X-Z)(X-\lambda Z)$, where $[X : Y : Z]$ are the projective coordinates on $\mathbb{P}_{\mathbb{Q}}^2$ and λ is the affine coordinate on $Y(2) \subset \mathbb{A}_{\mathbb{Q}}^1$. Let \mathcal{E}^g denote the g -th fibered power of \mathcal{E} over $Y(2)$ ($g \in \mathbb{N}$).

Set $I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $J_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Let $P = \mathrm{GL}_{2,\mathbb{Q}} \times \mathbb{Q}^{2g}$, where $\mathrm{GL}_{2,\mathbb{Q}}$ acts on $\mathbb{Q}^{2g} = (\mathbb{Q}^2)^{\oplus g}$ by acting tautologically on each of the g summands. Then \mathcal{E}^g is canonically a connected mixed Shimura variety of Kuga type, associated to the triple $(P, X^+, \Gamma \times \mathbb{Z}^{2g})$, where X^+ is a connected component of the $P(\mathbb{R})$ -conjugacy class of the homomorphism $h : \mathbb{S}_{\mathbb{C}} \rightarrow \mathrm{GL}_{2,\mathbb{C}} \times \mathbb{C}^{2g}$ defined by

$$\alpha \in \mathbb{S}(\mathbb{R}) = \mathbb{C}^* \mapsto h(\alpha) = ((\mathrm{Re} \alpha)I_2 + (\mathrm{Im} \alpha)J_2, 0)$$

and

$$\Gamma = \left\{ A \in \mathrm{SL}_2(\mathbb{Z}); A \equiv \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{2} \right\}.$$

As a complex manifold, X^+ is isomorphic to $\mathbb{H} \times \mathbb{C}^g$.

Theorem 13.1. *Let K be an algebraically closed field. Then $\mathrm{ZP}(\mathcal{E}_K^g, 1, d)$ holds for all $g \in \mathbb{N}$ and all non-negative integers d .*

Proof. By Theorem 9.6, the category of connected mixed Shimura varieties of Kuga type over $\bar{\mathbb{Q}}$ is very distinguished. We want to apply Theorem 10.7(1), choosing the class \mathcal{S} there as small as possible. Looking at the proof of Theorem 9.6 and at Theorem 8.2 in [Gao20] shows that we first have to include all triples $((Q, Y^+)/N, *)$, where (Q, Y^+) is a Shimura subdatum of (P, X^+) and N is a connected algebraic subgroup of Q^{der} , normal in Q (equivalently: N is a normal connected algebraic subgroup of Q whose reductive part is semisimple). Note that we can assume that N is connected in the conclusion of Theorem 8.2 in [Gao20] since replacing N by its identity component does not change $N(\mathbb{R})^+$.

By Proposition 1.2.16 and its proof in [Gao14] (cf. Proposition 3.4 in [Gao17a]), we have $Q = (I_2, v)(G_Q \times V)(I_2, v)^{-1}$, where $G_Q \subset \mathrm{GL}_{2,\mathbb{Q}}$ is a reductive subgroup, V is a G_Q -invariant vector subspace of \mathbb{Q}^{2g} , and $v \in \mathbb{Q}^{2g}$. Up to isomorphism, we can assume that $v = 0$. Furthermore, $G_{Q,\mathbb{R}}$ contains a 2-dimensional torus (the image of \mathbb{S}) that contains the center of $\mathrm{GL}_{2,\mathbb{R}}$. It follows from Theorem 1 in [NvdPT08] that G_Q is either a 2-dimensional torus T or $G_Q = \mathrm{GL}_{2,\mathbb{Q}}$. Thanks to Proposition 4.17 in [Mil17], we can then assume (up to isomorphism) that V is the direct sum of g_Q summands of the direct sum $(\mathbb{Q}^2)^{\oplus g}$ for some non-negative integer g_Q as these summands are all simple as representations of either T or $\mathrm{GL}_{2,\mathbb{Q}}$.

Since (Q, Y^+) is a connected mixed Shimura datum, we have $Q^{\mathrm{der}} = (G_Q)^{\mathrm{der}} \times V$, so $Q^{\mathrm{der}} = V$ in case $G_Q = T$ and $Q^{\mathrm{der}} = \mathrm{SL}_{2,\mathbb{Q}} \times V$ if $G_Q = \mathrm{GL}_{2,\mathbb{Q}}$.

We set $V_N = N \cap V$ and $G_N = N/V_N$. Thanks to Proposition 4.17 in [Mil17], we can find a G_Q -invariant vector subspace $V_N^{\perp} \subset V$ such that $V = V_N \oplus V_N^{\perp}$ and $Q/N \simeq (G_Q/G_N) \times V_N^{\perp}$, where the action of G_Q/G_N on $V_N^{\perp} \simeq V/V_N$ is induced by the action of

G_Q on V . Furthermore, V_N^\perp can be chosen as the direct sum of some of the summands in the direct sum $(\mathbb{Q}^2)^{\oplus g_Q}$.

In case $G_Q = T$, we have that $N = V_N$ is a T -invariant vector subspace of V and therefore $Q/N \simeq T \ltimes V_N^\perp$.

If $G_Q = \mathrm{GL}_{2,\mathbb{Q}}$, then $G_N \subset \mathrm{SL}_{2,\mathbb{Q}}$ is normal and connected and $V_N \subset V$ is a $\mathrm{GL}_{2,\mathbb{Q}}$ -invariant vector subspace. Since $\mathrm{SL}_{2,\mathbb{Q}}$ is almost-simple (as defined in Definition 19.7 in [Mil17]), it follows that $G_N = \{1\}$ or $G_N = \mathrm{SL}_{2,\mathbb{Q}}$.

In the first case, $N = V_N$ is a $\mathrm{GL}_{2,\mathbb{Q}}$ -invariant vector subspace of V . Then, $Q/N \simeq \mathrm{GL}_{2,\mathbb{Q}} \ltimes V_N^\perp$.

We are then left with the case $G_N = \mathrm{SL}_{2,\mathbb{Q}}$. We have that $(-I_2, w) \in N(\mathbb{C})$ for some $w \in V(\mathbb{C})$ and hence

$$(-I_2, w)(I_2, v)(-I_2, w)(I_2, -v) = (I_2, -2v) \in N(\mathbb{C})$$

for every $v \in V(\mathbb{C})$. This implies that $N = \mathrm{SL}_{2,\mathbb{Q}} \ltimes V = Q^{\mathrm{der}}$ and therefore $Q/N \simeq \mathbb{G}_{m,\mathbb{Q}}$.

Hence there are the following possibilities for Q/N (up to isomorphism): $T \ltimes V_N^\perp$, $\mathrm{GL}_{2,\mathbb{Q}} \ltimes V_N^\perp$, and $\mathbb{G}_{m,\mathbb{Q}}$. In principle, we now have to iterate this step and again look at all quotients of Shimura subdata as above of these connected mixed Shimura data. But this yields nothing new, so we have found our \mathcal{S} . The above considerations also show that the special subvarieties of \mathcal{E}^g are precisely the irreducible components of flat subgroup schemes (see Section 2 of [BC16]) and algebraic subgroups of CM fibers.

Instead of checking $\mathrm{ZP}(X, 1, d)$ for all $X \in \mathcal{S}$, we can use Lemma 10.6 to fix one congruence subgroup for each connected mixed Shimura datum that occurs in \mathcal{S} . To be able to apply Theorem 10.7(1), we therefore only have to check that $\mathrm{ZP}(X, 1, d)$ holds for X equal to a point, a j -th power of a CM elliptic curve over $\bar{\mathbb{Q}}$ ($1 \leq j \leq g$), or \mathcal{E}^j ($1 \leq j \leq g$). This follows from the results of [RV03], [Via08], [Gal10], [BC16], and [Bar19]. \square

14. ZILBER-PINK FOR A COMPLEX CURVE IN A SEMIABELIAN VARIETY

Theorem 14.1. *Let K be an algebraically closed field and let G be a semiabelian variety over $\bar{\mathbb{Q}}$. Then $\mathrm{ZP}(G_K, 1, d)$ holds for all non-negative integers d .*

Proof. By Theorem 9.3, the category of semiabelian varieties over $\bar{\mathbb{Q}}$ is very distinguished. Theorem 14.1 then follows from Theorem 10.7(1) combined with the main result of the recent work of the first-named author, Kühne, and Schmidt [BKS21]. \square

ACKNOWLEDGEMENTS

We thank Chris Daw for providing us with the crucial part of the proof of Lemma 5.2 and David Masser for pointing out his paper [Mas99] to us. We thank Cassani, Ziyang Gao, Philipp Habegger, Lars Kühne, Ben Moonen, Martin Orr, Jonathan Pila, Harry Schmidt, Christian Urech, and Immanuel van Santen for relevant and helpful discussions and correspondence. The second-named author was supported during his work on this article by the Swiss National Science Foundation as part of the project ‘‘Diophantine Problems, o-Minimality, and Heights’’, no. 200021_165525, as well as through the Early Postdoc.Mobility grant no. P2BSP2_195703. He thanks the Mathematical Institute of the University of Oxford and his host there, Jonathan Pila, for hosting him as a visitor for the duration of this grant.

REFERENCES

- [Bar19] F. Barroero, *CM relations in fibered powers of elliptic families*, J. Inst. Math. Jussieu **18** (2019), no. 5, 941–956.
- [BC16] F. Barroero and L. Capuano, *Linear relations in families of powers of elliptic curves*, Algebra Number Theory **10** (2016), no. 1, 195–214.

- [BD19] F. Barroero and G. A. Dill, *On the Zilber-Pink conjecture for complex abelian varieties*, to appear in Ann. Sci. Éc. Norm. Supér. (4), <https://arxiv.org/abs/1909.01271>, 2019.
- [BD21] G. Binyamini and C. Daw, *Effective computations for weakly optimal subvarieties*, <https://arxiv.org/abs/2105.12760>, 2021.
- [BIJ⁺19] R. Benedetto, P. Ingram, R. Jones, M. Manes, J. H. Silverman, and T. J. Tucker, *Current trends and open problems in arithmetic dynamics*, Bull. Amer. Math. Soc. (N.S.) **56** (2019), no. 4, 611–685.
- [BKS21] F. Barroero, L. Kühne, and H. Schmidt, *Unlikely intersections of curves with algebraic subgroups in semiabelian varieties*, <https://arxiv.org/abs/2108.12405>, 2021.
- [BMZ99] E. Bombieri, D. Masser, and U. Zannier, *Intersecting a curve with algebraic subgroups of multiplicative groups*, Int. Math. Res. Not. IMRN (1999), no. 20, 1119–1140.
- [BMZ07] ———, *Anomalous subvarieties—structure theorems and applications*, Int. Math. Res. Not. IMRN (2007), no. 19, Art. ID rnm057.
- [BMZ08] ———, *On unlikely intersections of complex varieties with tori*, Acta Arith. **133** (2008), no. 4, 309–323.
- [Bri17] M. Brion, *Commutative algebraic groups up to isogeny*, Doc. Math. **22** (2017), 679–725.
- [Cas21a] Cassani, *The Defect Condition*, MODNET eprint, 2021.
- [Cas21b] ———, *The Finiteness Result à la Bogomolov*, MODNET eprint, 2021.
- [CGMM13] Z. Chatzidakis, D. Ghioca, D. Masser, and G. Maurin, *Unlikely, likely and impossible intersections without algebraic groups*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **24** (2013), no. 4, 485–501.
- [Chi21] K. C. T. Chiu, *Ax-Schanuel for variations of mixed Hodge structures*, <https://arxiv.org/abs/2101.10968>, 2021.
- [CL12] A. Chambert-Loir, *Relations de dépendance et intersections exceptionnelles*, no. 348, 2012, Séminaire Bourbaki: Vol. 2010/2011. Exposés 1027–1042, pp. Exp. No. 1032, viii, 149–188.
- [DMOS82] P. Deligne, J. S. Milne, A. Ogus, and K. Shih, *Hodge cycles, motives, and Shimura varieties*, Lecture Notes in Mathematics, vol. 900, Springer-Verlag, Berlin-New York, 1982.
- [DO19a] C. Daw and M. Orr, *Unlikely intersections with $E \times CM$ curves in \mathcal{A}_2* , to appear in Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), <https://arxiv.org/abs/1902.10483>, 2019.
- [DO19b] ———, *Quantitative reduction theory and unlikely intersections*, to appear in Int. Math. Res. Not. IMRN, <https://arxiv.org/abs/1911.05618>, 2019.
- [DR18] C. Daw and J. Ren, *Applications of the hyperbolic Ax-Schanuel conjecture*, Compos. Math. **154** (2018), no. 9, 1843–1888.
- [Gal10] A. Galateau, *Une minoration du minimum essentiel sur les variétés abéliennes*, Comment. Math. Helv. **85** (2010), no. 4, 775–812.
- [Gao14] Z. Gao, *The mixed Ax-Lindemann theorem and its applications to the Zilber-Pink conjecture*, Ph.D. thesis, Universiteit Leiden/Université Paris-Sud, 2014.
- [Gao17a] ———, *A special point problem of André-Pink-Zannier in the universal family of Abelian varieties*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **17** (2017), no. 1, 231–266.
- [Gao17b] ———, *Towards the Andre-Oort conjecture for mixed Shimura varieties: the Ax-Lindemann theorem and lower bounds for Galois orbits of special points*, J. Reine Angew. Math. **732** (2017), 85–146.
- [Gao19] ———, *Enlarged mixed Shimura varieties, bi-algebraic system and some Ax type transcendental results*, Forum Math. Sigma **7** (2019), Paper No. e16, 65.

- [Gao20] ———, *Mixed Ax-Schanuel for the universal abelian varieties and some applications*, *Compos. Math.* **156** (2020), no. 11, 2263–2297.
- [GH13] D. Ghioca and L.-C. Hsia, *Torsion points in families of Drinfeld modules*, *Acta Arith.* **161** (2013), no. 3, 219–240.
- [GK21] Z. Gao and B. Klingler, *The Ax-Schanuel conjecture for variations of mixed Hodge structures*, <https://arxiv.org/abs/2101.10938>, 2021.
- [Gro60] A. Grothendieck, *Éléments de géométrie algébrique. I. Le langage des schémas*, *Inst. Hautes Études Sci. Publ. Math.* (1960), no. 4, 5–214.
- [GT21] D. Ghioca and T. J. Tucker, *A reformulation of the dynamical Manin-Mumford conjecture*, *Bull. Aust. Math. Soc.* **103** (2021), no. 1, 154–161.
- [GW10] U. Görtz and T. Wedhorn, *Algebraic geometry I*, *Advanced Lectures in Mathematics*, Vieweg + Teubner, Wiesbaden, 2010, Schemes with examples and exercises.
- [GW21] U. Görtz and T. Wedhorn, *Errata for “Algebraic geometry I (Edition 1)”*, <https://algebraic-geometry.de/errata/1/>, 2021.
- [HP16] P. Habegger and J. Pila, *O-minimality and certain atypical intersections*, *Ann. Sci. Éc. Norm. Supér. (4)* **49** (2016), no. 4, 813–858.
- [Kir09] J. Kirby, *The theory of the exponential differential equations of semiabelian varieties*, *Selecta Math. (N.S.)* **15** (2009), no. 3, 445–486.
- [Kli17] B. Klingler, *Hodge loci and atypical intersections: conjectures*, to appear in the book “Motives and complex multiplication” (Birkhäuser), <https://arxiv.org/abs/1711.09387>, 2017.
- [LV19] A. Lubotzky and T. N. Venkataramana, *The congruence topology, Grothendieck duality and thin groups*, *Algebra Number Theory* **13** (2019), no. 6, 1281–1298.
- [Mas99] D. W. Masser, *Specializations of some hyperelliptic Jacobians*, *Number theory in progress, Vol. 1 (Zakopane-Kościelisko, 1997)*, de Gruyter, Berlin, 1999, pp. 293–307.
- [Mil05] J. S. Milne, *Introduction to Shimura varieties*, *Harmonic analysis, the trace formula, and Shimura varieties*, *Clay Math. Proc.*, vol. 4, Amer. Math. Soc., Providence, RI, 2005, pp. 265–378.
- [Mil17] ———, *Algebraic groups*, *Cambridge Studies in Advanced Mathematics*, vol. 170, Cambridge University Press, Cambridge, 2017, The theory of group schemes of finite type over a field.
- [Moo] B. Moonen, *Letter to Christophe Cornut*, private communication.
- [NvdPT08] K. A. Nguyen, M. van der Put, and J. Top, *Algebraic subgroups of $GL_2(\mathbb{C})$* , *Indag. Math. (N.S.)* **19** (2008), no. 2, 287–297.
- [Orr13] M. Orr, *The André–Pink conjecture : Hecke orbits and weakly special subvarieties*, Ph.D. thesis, Université Paris Sud – Paris XI, 2013.
- [OV90] A. L. Onishchik and È. B. Vinberg, *Lie groups and algebraic groups*, *Springer Series in Soviet Mathematics*, Springer-Verlag, Berlin, 1990, Translated from the Russian and with a preface by D. A. Leites.
- [Pil17] J. Pila, *On a modular Fermat equation*, *Comment. Math. Helv.* **92** (2017), no. 1, 85–103.
- [Pil21] ———, *Unpublished manuscript*, private communication, 2021.
- [Pin90] R. Pink, *Arithmetical compactification of mixed Shimura varieties*, *Bonner Mathematische Schriften [Bonn Mathematical Publications]*, vol. 209, Universität Bonn, Mathematisches Institut, Bonn, 1990, Dissertation, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, 1989.
- [Pin05a] ———, *A combination of the conjectures of Mordell–Lang and André–Oort*, *Geometric methods in algebra and number theory*, *Progr. Math.*, vol. 235,

- Birkhäuser Boston, Boston, MA, 2005, pp. 251–282.
- [Pin05b] ———, *A common generalization of the conjectures of André-Oort, Manin-Mumford, and Mordell-Lang*, <https://people.math.ethz.ch/~pink/ftp/AOMMML.pdf>, 2005.
- [Poi01] B. Poizat, *L'égalité au cube*, J. Symbolic Logic **66** (2001), no. 4, 1647–1676.
- [Poo17] B. Poonen, *Rational points on varieties*, Graduate Studies in Mathematics, vol. 186, American Mathematical Society, Providence, RI, 2017.
- [PR94] V. Platonov and A. Rapinchuk, *Algebraic groups and number theory*, Pure and Applied Mathematics, vol. 139, Academic Press, Inc., Boston, MA, 1994, Translated from the 1991 Russian original by Rachel Rowen.
- [PS21] J. Pila and T. Scanlon, *Effective transcendental Zilber-Pink for variations of Hodge structures*, <https://arxiv.org/abs/2105.05845>, 2021.
- [PT19] J. Pila and J. Tsimerman, *Independence of CM points in Elliptic Curves*, to appear in J. Eur. Math. Soc. (JEMS), <https://arxiv.org/abs/1907.02737>, 2019.
- [PZ08] J. Pila and U. Zannier, *Rational points in periodic analytic sets and the Manin-Mumford conjecture*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **19** (2008), no. 2, 149–162.
- [Rém09] G. Rémond, *Intersection de sous-groupes et de sous-variétés. III*, Comment. Math. Helv. **84** (2009), no. 4, 835–863.
- [RS93] J. Rohlf's and J. Schwermer, *Intersection numbers of special cycles*, J. Amer. Math. Soc. **6** (1993), no. 3, 755–778.
- [RV03] G. Rémond and E. Viada, *Problème de Mordell-Lang modulo certaines sous-variétés abéliennes*, Int. Math. Res. Not. IMRN (2003), no. 35, 1915–1931.
- [Sat80] I. Satake, *Algebraic structures of symmetric domains*, Kanô Memorial Lectures, vol. 4, Iwanami Shoten, Tokyo; Princeton University Press, Princeton, N.J., 1980.
- [Sch89] A. Schinzel, *Reducibility of lacunary polynomials. X*, Acta Arith. **53** (1989), no. 1, 47–97.
- [Sch00] ———, *Polynomials with special regard to reducibility*, Encyclopedia of Mathematics and its Applications, vol. 77, Cambridge University Press, Cambridge, 2000, With an appendix by Umberto Zannier.
- [SGA03] *Revêtements étales et groupe fondamental (SGA 1)*, Documents Mathématiques (Paris), vol. 3, Société Mathématique de France, Paris, 2003, Séminaire de géométrie algébrique du Bois Marie 1960–61, directed by A. Grothendieck, with two papers by M. Raynaud, updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin].
- [Tsi18] J. Tsimerman, *The André-Oort conjecture for \mathcal{A}_g* , Ann. of Math. (2) **187** (2018), no. 2, 379–390.
- [Ull17] E. Ullmo, *Structures spéciales et problème de Zilber-Pink*, Around the Zilber-Pink conjecture/Autour de la conjecture de Zilber-Pink, Panor. Synthèses, vol. 52, Soc. Math. France, Paris, 2017, pp. 1–30.
- [Via03] E. Viada, *The intersection of a curve with algebraic subgroups in a product of elliptic curves*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **2** (2003), no. 1, 47–75.
- [Via08] ———, *The intersection of a curve with a union of translated codimension-two subgroups in a power of an elliptic curve*, Algebra Number Theory **2** (2008), no. 3, 249–298.
- [Zan12] U. Zannier, *Some Problems of Unlikely Intersections in Arithmetic and Geometry*, Annals of Mathematics Studies, vol. 181, Princeton University Press, 2012, With appendixes by David Masser.
- [Zil02] B. Zilber, *Exponential sums equations and the Schanuel conjecture*, J. London Math. Soc. (2) **65** (2002), no. 1, 27–44.

[Zil16] ———, *Model theory of special subvarieties and Schanuel-type conjectures*, Ann. Pure Appl. Logic **167** (2016), no. 10, 1000–1028.

UNIVERSITÀ DEGLI STUDI ROMA TRE, DIPARTIMENTO DI MATEMATICA E FISICA, LARGO SAN MURIALDO 1, 00146 ROMA, ITALY

Email address: `fbarroero@gmail.com`

UNIVERSITY OF OXFORD, MATHEMATICAL INSTITUTE, ANDREW WILES BUILDING, RADCLIFFE OBSERVATORY QUARTER, WOODSTOCK ROAD, OXFORD OX2 6GG

Email address: `gabriel.dill@maths.ox.ac.uk`