

STABILITY MANIFOLDS OF VARIETIES WITH FINITE ALBANESE MORPHISMS

LIE FU, CHUNYI LI, AND XIAOLEI ZHAO

ABSTRACT. For a smooth projective complex variety whose Albanese morphism is finite, we show that every Bridgeland stability condition on its bounded derived category of coherent sheaves is geometric, in the sense that all skyscraper sheaves are stable with the same phase. Furthermore, we describe the stability manifolds of irregular surfaces and abelian threefolds with Picard rank one, and show that they are connected and contractible.

CONTENTS

1. Introduction	1
2. Geometric stability conditions	3
3. Surface case	9
4. Abelian threefold case	12
References	21

1. INTRODUCTION

Let X be a smooth projective variety over the field of complex numbers \mathbb{C} . Denote by $D^b(X)$ the bounded derived category of coherent sheaves on X . The notion of stability conditions on triangulated categories was introduced by Bridgeland in [Bri07] (see Section 2.1 for a recap). Let $\text{Stab}(X)$ be the set of (full locally-finite numerical) stability conditions on $D^b(X)$. By the seminal result in [Bri07], $\text{Stab}(X)$ is naturally endowed with a structure of complex manifold with local coordinates given by the central charge. We call $\text{Stab}(X)$ the *stability manifold* of X .

A complete description of the stability manifold has been worked out only for curves and abelian surfaces. More precisely,

- $\text{Stab}(\mathbb{P}^1) \cong \mathbb{C}^2$, see [Oka06].
- $\text{Stab}(C) \cong \mathbb{C} \times \mathbb{H}$ for any smooth projective curve C of genus ≥ 1 , see [Bri07, Mac07]; here \mathbb{H} denotes the upper half-plane.
- Abelian surfaces, see [Bri08, HMS08].

In any of the following cases, a principal connected component of the stability manifold is determined, which is expected to be the whole stability manifold:

- K3 surfaces with Picard rank one, see [Bri08, BB17].

Date: October 12, 2021.

Key words and phrases. Bridgeland stability conditions, irregular surfaces, Albanese morphism, abelian threefolds.

- The projective plane \mathbf{P}^2 , see [Li17].
- Abelian threefolds with Picard rank one, see [MP15, MP13, BMS16].

Moreover, in each case above, the stability manifold (or the principal component) is simply connected and contains an open subset consisting of the so-called *geometric* stability conditions, meaning that all skyscraper sheaves are stable with the same phase (see Definition 2.8). Note that the condition of equal phase turns out to be automatic, see Proposition 2.9.

In this paper, we provide more instances of algebraic varieties whose stability manifolds have the same feature as the aforementioned examples. Our first main result reads as follows:

Theorem 1.1 (Corollary 2.15). *Let X be a connected smooth projective variety over \mathbb{C} . If the Albanese morphism of X is finite, then every numerical stability condition on $D^b(X)$ is geometric.*

Here we briefly recall some examples of varieties with finite Albanese morphisms. The most basic ones are abelian varieties and curves of genus ≥ 1 . To produce new examples out of old ones, observe that this property is stable under products and is inherited by subvarieties and finite ramified covers. See [Kaw81] for more on the structure of a variety with finite Albanese morphism.

Also, note that a smooth projective surface S of Picard rank one has finite Albanese morphism if and only if it is *irregular*, that is, $q(S) := \dim H^1(S, \mathcal{O}_S) \neq 0$. One expects that a generic minimal surface of irregularity $q \geq 2$ is of this type. Here are some geometrically interesting examples: the Fano surface of lines on a generic cubic threefold ([Rou11]), the Fano surface of planes on a generic cubic fivefold.

Theorem 1.2 (Corollary 3.8). *The stability manifold of an irregular surface of Picard rank one is connected and contractible.*

One should compare Theorem 1.2 with [HMS08, Theorem 1] on abelian surfaces. Finally, we establish the following result on abelian threefolds.

Theorem 1.3 (Corollary 4.9). *Let (A, H) be a polarized abelian threefold, then $\text{Stab}_H(A) = \tilde{\mathfrak{P}}$ as that constructed in [BMS16, Theorem 9.1]. In particular, when A is with Picard rank one, $\tilde{\mathfrak{P}}$ is the whole stability manifold and it is connected and contractible.*

This completes the result of [MP15, MP13, BMS16] on the stability manifold of abelian threefolds of Picard rank one.

Sketch of the proof. Theorem 1.1 mainly relies on the result in [Pol07] and the uniqueness of Harder–Narasimhan filtration. By [Pol07, Corollary 3.5.2], the natural action of the group $\text{Pic}^0(X)$ on the stability manifold is trivial. Hence the Harder–Narasimhan factors of every $\text{Pic}^0(X)$ -invariant object, e.g. a skyscraper sheaf, must also be $\text{Pic}^0(X)$ -invariant. It follows by the assumption on the Albanese morphism that such factors of a skyscraper sheaf may only be skyscraper sheaves. In particular, all skyscraper sheaves are stable. On the other hand, we prove in general that all skyscraper sheaves are of the same phase when they are all stable, see Proposition 2.9. This seemingly-simple statement is probably known to some experts, but we found no proof in the literature.

Theorem 1.2 on irregular surfaces then follows from the same observation as that in [Bri08, Section 10], which implies that every geometric stability condition can be constructed via tilting heart up to the $\widetilde{\text{GL}}_2^+(\mathbb{R})$ -action.

The proof of Theorem 1.3 on abelian threefolds is more involved. First, for any abelian variety A , using a variant of the Fourier–Mukai transform between $D^b(A)$ and $D^b(A^\vee)$, we show that all simple semi-homogeneous vector bundles are stable with respect to all stability conditions, see Corollary 2.17. Second, specializing to abelian threefolds, we show that every stability condition is with the same central charge as that constructed in [BMS16]. The actual difficulty of the whole argument is to prove that their heart structures are also the same. Proposition 4.3, Lemma 4.6 and Lemma 4.7 are to deal with this issue. Roughly speaking, we show that there are enough morphisms from simple semi-homogeneous vector bundles to other stable objects so that a stability condition can be uniquely determined by the phases of all semi-homogeneous vector bundles and the central charge.

Acknowledgements. We are very grateful to Arend Bayer, Emanuele Macrì, and Paolo Stellari for pointing out references and enlightening comments. In particular, the shortcut proof of Lemma 2.14 is due to Arend Bayer. L. F. is supported by the Radboud Excellence Initiative program. C. L. is a University Research Fellow supported by the Royal Society URF\R1\201129 “Stability condition and application in algebraic geometry”. X. Z. is partially supported by the Simons Collaborative Grant 636187.

2. GEOMETRIC STABILITY CONDITIONS

2.1. Generalities on stability conditions. Let X be a smooth projective variety over \mathbb{C} . Denote by $D^b(X)$ the bounded derived category of coherent sheaves on X . We recall some basic notions of stability conditions.

Definition 2.1. A *slicing* \mathcal{P} on $D^b(X)$ is a collection of full additive subcategories $\mathcal{P}(\phi) \subset D^b(X)$ indexed by all $\phi \in \mathbb{R}$ such that

- (a) $\mathcal{P}(\phi)[1] = \mathcal{P}(\phi + 1)$;
- (b) if $\phi_1 > \phi_2$ and $F_i \in \text{Obj}(\mathcal{P}(\phi_i))$, then $\text{Hom}(F_1, F_2) = 0$;
- (c) for any object E in $D^b(X)$, there are real numbers $\phi_1 > \cdots > \phi_m$, objects E_i in $D^b(X)$, and a collection of distinguished triangles

$$\begin{array}{ccccccc}
 0 = E_0 & \longrightarrow & E_1 & \longrightarrow & E_2 & \rightarrow \cdots \rightarrow & E_{m-1} & \longrightarrow & E_m = E \\
 & & \swarrow & & \swarrow & & \swarrow & & \swarrow \\
 & & A_1 & & A_2 & & A_m & & \\
 & \nwarrow & & \nwarrow & & \nwarrow & & \nwarrow & \\
 & & +1 & & +1 & & +1 & &
 \end{array}$$

such that $A_i = \text{Cone}(E_{i-1} \rightarrow E_i)$ is an object in $\mathcal{P}(\phi_i)$ for every $1 \leq i \leq m$.

Remark 2.2. Non-zero objects of $\mathcal{P}(\phi)$ are called *semistable* with phase ϕ ; simple objects of $\mathcal{P}(\phi)$ are called *stable*. The sequence of maps in (c) is called the *Harder–Narasimhan (HN) filtration* of E . It is not hard to see that the HN filtration is unique up to isomorphism. We will call each A_i the *HN factor* of E . We denote $\phi_{\mathcal{P}}^+(E) := \phi_1$ and $\phi_{\mathcal{P}}^-(E) := \phi_m$; when E is semistable, we denote $\phi_{\mathcal{P}}(E)$ the phase of E (or simply $\phi(E)$ when there is no ambiguity).

Notation 2.3. Let \mathcal{P} be a slicing. For every interval $I \subset \mathbb{R}$, we define a *truncated HN factor* $\text{HN}_{\mathcal{P}}^I(E)$ of E as follows. In the HN filtration 2.1(c) of E , let $a, b \in \{1, 2, \dots, m\}$ be such that

$\phi_i \in I$ when and only when $a \leq i \leq b$, then $\text{HN}_{\mathcal{P}}^I(E)$ is defined to be $\text{Cone}(E_{a-1} \rightarrow E_b)$ (and to be 0 if no ϕ_i is contained in I). We denote by $\mathcal{P}(I)$ the full subcategory of all objects $E \in D^b(X)$ such that $\phi_{\mathcal{P}}^{\pm}(E) \in I$.

Denote by $K(X)$ the Grothendieck group of $D^b(X)$.

Definition 2.4. A *Bridgeland pre-stability condition* on $D^b(X)$ is a pair $\sigma = (\mathcal{P}, Z)$, where

- \mathcal{P} is a slicing of $D^b(X)$;
- $Z : K(X) \rightarrow \mathbb{C}$ is a group homomorphism, called the *central charge*;

such that for any non-zero object E in $\mathcal{P}(\phi)$, we have $Z([E]) = m(E)e^{i\pi\phi}$ for some $m(E) \in \mathbb{R}_{>0}$.

The set of all pre-stability conditions has a natural topology induced by the following generalised metric function: for two pre-stability conditions $\sigma_1 = (\mathcal{P}_1, Z_1)$ and $\sigma_2 = (\mathcal{P}_2, Z_2)$, their distance is defined as

$$\text{dist}(\sigma_1, \sigma_2) := \sup_{\substack{E \in D^b(X) \\ E \neq 0}} \left\{ \left| \phi_{\mathcal{P}_1}^-(E) - \phi_{\mathcal{P}_2}^-(E) \right|, \left| \phi_{\mathcal{P}_1}^+(E) - \phi_{\mathcal{P}_2}^+(E) \right|, \|Z_1 - Z_2\| \right\} \in [0, +\infty].$$

We have the following two natural group actions on the set of pre-stability conditions; see [MS17, Remark 5.14] for more details.

- The universal covering $\widetilde{\text{GL}}_2^+(\mathbb{R})$ of the group $\text{GL}_2^+(\mathbb{R})$ acts on the right of this set. This $\widetilde{\text{GL}}_2^+(\mathbb{R})$ -action does not affect the (semi)stability of any object.
- The group $\text{Aut}(D^b(X))$ of autoequivalences of $D^b(X)$ acts on the left of this set.

Let Λ be a free abelian group of finite rank and let $\lambda : K(X) \twoheadrightarrow \Lambda$ be a surjective homomorphism. A typical choice (by default) for Λ is the numerical Grothendieck group $K_{\text{num}}(X)$ and λ is the natural projection $K(X) \twoheadrightarrow K_{\text{num}}(X)$.

Definition 2.5 ([Bri07, KS08]). We say that a pre-stability condition (\mathcal{P}, Z) satisfies the *support property* (with respect to Λ , or rather λ) if the central charge Z factors through $\lambda : K(X) \twoheadrightarrow \Lambda$ and there is a quadratic form Q_{Λ} on $\Lambda \otimes \mathbb{R}$ such that

- (a) the kernel $\text{Ker } Z \subset \Lambda \otimes \mathbb{R}$ of the central charge is negative definite with respect to Q_{Λ} ;
- (b) for any semistable object E , we have $Q_{\Lambda}(E) \geq 0$.

A pre-stability condition satisfying the support property is called a *stability condition*, and we denote the set of all stability conditions (with respect to Λ , or rather λ) as $\text{Stab}_{\Lambda}(X)$.

When λ factors through $K(X) \twoheadrightarrow K_{\text{num}}(X)$, we will call the stability condition *numerical*.

Theorem 2.6 ([Bri07, Theorem 7.1], [Bay19, Theorem 1.2]). *The space $\text{Stab}_{\Lambda}(X)$ of stability conditions is naturally a complex manifold of dimension $\text{rank}(\Lambda)$, such that the map forgetting the slicing*

$$\pi_Z : \text{Stab}_{\Lambda}(X) \rightarrow \text{Hom}_{\mathbb{Z}}(\Lambda, \mathbb{C}) \quad \sigma = (Z, \mathcal{P}) \mapsto Z$$

is a local biholomorphic isomorphism at every point of $\text{Stab}_{\Lambda}(X)$.

Remark 2.7. Note that the notion of pre-stability condition in this paper (as well as many others) is called stability condition in Bridgeland’s original paper [Bri07].

When the lattice Λ is the numerical Grothendieck group, our definition of stability conditions is the same as that of the full locally-finite numerical stability conditions in [Bri08]. We denote the space of such stability conditions as $\text{Stab}(X)$ and call it the *stability manifold* of X .

2.2. Geometric stability conditions.

Definition 2.8. A stability condition σ on $D^b(X)$ is called *geometric* (with respect to X) if for each point $p \in X$, the skyscraper sheaf \mathcal{O}_p is σ -stable, and all skyscraper sheaves are of the same phase.

The definition is similar to that of [Bri08, Definition 10.2]. The only difference is that in [Bri08], the author further assumes a ‘goodness’ condition. In the next section, we will see that the ‘goodness’ condition is redundant at least in the surface case. In the next proposition, we show that the condition of having the same phase is redundant for numerical stability conditions when X is a smooth connected projective variety. The result might be known to some experts, but there is no proof written down as far as we know.

Proposition 2.9. *Let X be a connected smooth projective variety. Let σ be a numerical stability condition on $D^b(X)$ such that every skyscraper sheaf is σ -stable. Then all skyscraper sheaves are of the same phase, in other words, σ is geometric.*

We first establish the following technical lemma.

Lemma 2.10. *Let σ be a stability condition on $D^b(X)$. Let $g: F \rightarrow E$ be a morphism and $\tilde{F} := \text{Cone}(F \xrightarrow{g} E)[-1]$ in $D^b(X)$.*

1. *Then $\text{HN}_\sigma^a(F) \cong \text{HN}_\sigma^a(\tilde{F})$ for every $a \notin [\phi^-(E) - 1, \phi^+(E)]$.*
2. *If E is non-zero and the second smallest phase of the HN factors of E is greater than $\phi^-(E) + 1$, then $\text{HN}_\sigma^{\leq \phi^-(E)}(F) \not\cong \text{HN}_\sigma^{\leq \phi^-(E)}(\tilde{F})$.*

Proof. As for the first statement, we consider the following diagram of distinguished triangles:

$$\begin{array}{ccccccc}
0 & \longrightarrow & \text{HN}_\sigma^{>\phi^+(E)}(\tilde{F}) & \xrightarrow{\text{id}} & \text{HN}_\sigma^{>\phi^+(E)}(\tilde{F}) & \longrightarrow & 0 \\
\downarrow & & \downarrow \text{can} & & \downarrow f & & \downarrow \\
E[-1] & \longrightarrow & \tilde{F} & \xrightarrow{g'} & F & \xrightarrow{g} & E \\
\downarrow \text{id} & & \downarrow & & \downarrow & & \downarrow \text{id} \\
E[-1] & \longrightarrow & \text{HN}_\sigma^{\leq \phi^+(E)}(\tilde{F}) & \longrightarrow & K & \longrightarrow & E
\end{array}$$

The morphism f is the composition of g' and can . The square on the top-right commutes since $g \circ f \in \text{Hom}(\text{HN}_\sigma^{>\phi^+(E)}(\tilde{F}), E) = 0$. The sequence on the bottom is the cone of distinguished triangles on the top and middle, hence it is a distinguished triangle by the octahedral axiom.

For any σ -semistable object A with phase $> \phi^+(E)$, by applying $\text{Hom}(A, -)$ to the distinguished triangle on the bottom, we have $\text{Hom}(A, K) = 0$. Therefore, the object K is in $\mathcal{P}_\sigma(\leq \phi^+(E))$.

Note that $K = \text{Cone}(\text{HN}_\sigma^{>\phi^+(E)}(\tilde{F}) \xrightarrow{f} F)$. It follows $K \cong \text{HN}_\sigma^{\leq\phi^+(E)}(F)$ and

$$\text{HN}_\sigma^{>\phi^+(E)}(\tilde{F}) \cong \text{HN}_\sigma^{>\phi^+(E)}(F).$$

By a similar argument for the diagram

$$\begin{array}{ccccccc} E[-1] & \longrightarrow & L & \longrightarrow & \text{HN}_\sigma^{>\phi^-(E)-1}(F) & \longrightarrow & E \\ \downarrow \text{id} & & \downarrow & & \downarrow & & \downarrow \text{id} \\ E[-1] & \longrightarrow & \tilde{F} & \longrightarrow & F & \longrightarrow & E \\ \downarrow & & \downarrow & & \downarrow \text{can} & & \downarrow \\ 0 & \longrightarrow & \text{HN}_\sigma^{\leq\phi^-(E)-1}(F) & \xrightarrow{\text{id}} & \text{HN}_\sigma^{\leq\phi^-(E)-1}(F) & \longrightarrow & 0, \end{array}$$

we get $\text{HN}_\sigma^{\leq\phi^-(E)-1}(\tilde{F}) \cong \text{HN}_\sigma^{\leq\phi^-(E)-1}(F)$. The first statement holds.

As for the second statement, note that $\mathcal{A} := \mathcal{P}_\sigma((\phi^-(E) - 1, \phi^-(E)])$ is a heart on $\text{D}^b(X)$. Apply $\mathcal{H}_\mathcal{A}^\bullet(-)$ to the distinguished triangle $\tilde{F} \rightarrow F \xrightarrow{g} E$, we get the long exact sequence

$$\mathcal{H}_\mathcal{A}^{-1}(E) \rightarrow \mathcal{H}_\mathcal{A}^0(\tilde{F}) \rightarrow \mathcal{H}_\mathcal{A}^0(F) \rightarrow \mathcal{H}_\mathcal{A}^0(E) \rightarrow \mathcal{H}_\mathcal{A}^1(\tilde{F}) \xrightarrow{h} \mathcal{H}_\mathcal{A}^1(F) \rightarrow \mathcal{H}_\mathcal{A}^1(E).$$

Note that for every object B , $\mathcal{H}_\mathcal{A}^i(B) = \text{HN}_\sigma^{(\phi^-(E)-i-1, \phi^-(E)-i]}(B)[i]$. By the assumption on E , we have $\mathcal{H}_\mathcal{A}^{\pm 1}(E) = 0$ and $\mathcal{H}_\mathcal{A}^0(E) = \text{HN}^-(E) := \text{HN}_\sigma^{\phi^-(E)}(E)$.

If the map h is not an isomorphism, then the kernel of h in \mathcal{A} is a quotient object of $\text{HN}^-(E)$, which must be in $\mathcal{P}_\sigma(\phi^-(E))$. We get a short exact sequence

$$0 \rightarrow \ker h \rightarrow \text{HN}_\sigma^{\phi^-(E)-1}(\tilde{F})[1] \xrightarrow{h} \text{HN}_\sigma^{\phi^-(E)-1}(F)[1] \rightarrow 0,$$

which implies $\text{HN}_\sigma^{\phi^-(E)-1}(\tilde{F}) \not\cong \text{HN}_\sigma^{\phi^-(E)-1}(F)$ as $[\ker h]$ is not 0 in $\text{K}_{\text{num}}(X)$.

If the map h is an isomorphism, then we have the short exact sequence

$$0 \rightarrow \mathcal{H}_\mathcal{A}^0(\tilde{F}) \rightarrow \mathcal{H}_\mathcal{A}^0(F) \rightarrow \mathcal{H}_\mathcal{A}^0(E) \rightarrow 0.$$

Since $[\mathcal{H}_\mathcal{A}^0(E)] \neq 0$ in $\text{K}_{\text{num}}(X)$, we have

$$\text{HN}_\sigma^{(\phi^-(E)-1, \phi^-(E))}(\tilde{F}) = \mathcal{H}_\mathcal{A}^0(\tilde{F}) \not\cong \mathcal{H}_\mathcal{A}^0(F) = \text{HN}_\sigma^{(\phi^-(E)-1, \phi^-(E))}(F).$$

In any case, we always have $\text{HN}_\sigma^{[\phi^-(E)-1, \phi^-(E)]}(F) \not\cong \text{HN}_\sigma^{[\phi^-(E)-1, \phi^-(E)]}(\tilde{F})$. \square

Proof of Proposition 2.9. For a smooth connected curve C on X , suppose not all skyscraper sheaves of the points on the curve C are with the same phase. Then there exist a large enough integer m and points $p_1, \dots, p_m, q_1, \dots, q_m$ on C such that

- $\mathcal{O}_C(-p_1 - \dots - p_m) \cong \mathcal{O}_C(-q_1 - \dots - q_m) =: \mathcal{L}$;

- $\phi(\mathcal{O}_{p_1}) < \phi(\mathcal{O}_{q_i})$ for every $1 \leq i \leq m$.

In particular, as the central charge factors via $K_{\text{num}}(X)$, $\phi(\mathcal{O}_{p_1}) \leq \phi(\mathcal{O}_{q_i}) - 2$ for every i . We may also assume $\phi(\mathcal{O}_{p_1}) = \min\{\phi(\mathcal{O}_{p_i})\}$.

Note that \mathcal{L} has two expressions $\text{Cone}(\mathcal{O}_C \rightarrow \mathcal{O}_{p_1} \oplus \cdots \oplus \mathcal{O}_{p_m})[-1]$ and $\text{Cone}(\mathcal{O}_C \rightarrow \mathcal{O}_{q_1} \oplus \cdots \oplus \mathcal{O}_{q_m})[-1]$. By Lemma 2.10.1, the HN factor HN^a of \mathcal{L} is isomorphic to that of \mathcal{O}_C for every $a < \min\{\phi(\mathcal{O}_{q_i})\} - 1$. By Lemma 2.10.2, $\text{HN}^{\leq \phi(\mathcal{O}_{p_1})}(\mathcal{L}) \not\cong \text{HN}^{\leq \phi(\mathcal{O}_{p_1})}(\mathcal{O}_C)$, which leads to a contradiction.

Hence, the phase of all skyscraper sheaves of points on C are the same. As X is connected and projective, this implies that all skyscraper sheaves are of the same phase. \square

The following lemma is an immediate generalization of [Bri08, Lemma 10.1] in the higher dimensional case.

Lemma 2.11 ([Bri08, Lemma 10.1]). *Let X be an n -dimensional smooth projective variety and $\sigma = (\mathcal{P}, \mathcal{Z})$ be a geometric stability condition on X such that $\mathcal{O}_p \in \mathcal{P}(1)$. Let F be an object of $\text{D}^b(X)$. Then*

- (a) *if $F \in \mathcal{P}((0, 1])$, then $\mathcal{H}^i(F)$ vanishes unless $-n + 1 \leq i \leq 0$, and moreover, $\mathcal{H}^{-n+1}(F)$ is torsion-free;*
- (b) *if $F \in \text{Coh}(X)$, then $F \in \mathcal{P}((-n + 1, 1])$; if F is a torsion sheaf, then $F \in \mathcal{P}((-n + 2, 1])$.*

Proof. (a) One may assume that F is stable and is not a skyscraper sheaf. For any point $p \in X$, since \mathcal{O}_p is stable with phase 1, $\text{Hom}(\mathcal{O}_p, F[i]) = \text{Hom}(F, \mathcal{O}_p[i - 1]) = 0$ for all $i \leq 0$. By Serre duality, $\text{Hom}(F, \mathcal{O}_p[i]) = 0$ unless $0 \leq i \leq n - 1$.

It follows from [BM02, Proposition 5.4] that F is quasi-isomorphic to a length n complex of locally free sheaves. So F satisfies the condition in part (a).

(b) Let F be a coherent sheaf. For every object $E \in \mathcal{P}(> 1)$, by part (a), $\mathcal{H}^i(E) = 0$ when $i \geq 0$. So $\text{Hom}(E, F) = 0$. For every object $G \in \mathcal{P}(\leq -n + 1)$, by part (a), $\mathcal{H}^i(E) = 0$ when $i \leq 0$. So $\text{Hom}(F, G) = 0$. It follows that $F \in \mathcal{P}((-n + 1, 1])$.

Now let F be a torsion sheaf. We have a distinguished triangle

$$E \rightarrow F \rightarrow G \xrightarrow{+1}$$

for some $E \in \mathcal{P}((-n + 2, 1])$ and $G \in \mathcal{P}((-n + 1, -n + 2])$. By part (a), $\mathcal{H}^i(G) = 0$ unless $0 \leq i \leq n - 1$, and $\mathcal{H}^0(G)$ is torsion-free. Since F is a torsion sheaf, $\text{Hom}(F, G) = 0$. This can only happen when $G = 0$, so $F \in \mathcal{P}((-n + 2, 1])$. \square

2.3. Stability of skyscraper sheaves. The following basic result is used in the study of homogeneous vector bundles on abelian varieties.

Lemma 2.12. *Let A be an abelian variety and $E \in \text{D}^b(A)$. Further assume that for any $\xi \in A^\vee$, we have that $E \otimes P_\xi \simeq E$, where P_ξ is the line bundle on A parameterized by ξ . Then the support of E is finite.*

Proof. See for example the proof of [Pol03, Proposition 11.8]. \square

Theorem 2.13. *Let X be a smooth projective variety such that its Albanese morphism is finite, then skyscraper sheaves are all stable with respect to every numerical stability condition on $D^b(X)$.*

Proof. Fix an arbitrary stability condition on X . Let $p \in X$ be a closed point. Let E_1, \dots, E_n be the Jordan–Hölder factors in the Harder–Narasimhan filtration associated with \mathcal{O}_p . By Polishchuk [Pol07, Corollary 3.5.2], the group $\text{Pic}^0(X) \subset \text{Aut}(D^b(X))$ acts trivially on the stability manifold. In particular, the Harder–Narasimhan filtration, as well as the Jordan–Hölder factors, are preserved by the action of $\text{Pic}^0(X)$. As a consequence, each E_i satisfies the condition that $E_i \otimes L \simeq E_i$ for any $L \in \text{Pic}^0(X)$.

Let $a : X \rightarrow \text{Alb}(X)$ be the Albanese morphism. For any $\xi \in \text{Pic}^0(\text{Alb}(X))$, denote by P_ξ the corresponding line bundle on $\text{Alb}(X)$. Then for any i , we have $E_i \otimes a^*(P_\xi) \simeq E_i$. By the projection formula,

$$Ra_*(E_i) \otimes P_\xi \simeq Ra_*(E_i),$$

which implies that the support of $Ra_*(E_i)$ is finite by Lemma 2.12. Combined with the assumption that a is a finite morphism, we see that the support of E_i is also finite, for any i .

The theorem then follows by Lemma 2.14. \square

Lemma 2.14. *Let X be a smooth variety and σ a stability condition on $D^b(X)$. Assume that the Jordan–Hölder factors of \mathcal{O}_p have finite support. Then \mathcal{O}_p is stable.*

Proof. Suppose \mathcal{O}_p is not stable, then there exists a stable object $E \neq \mathcal{O}_p$ supported at p . Let k, l be the maximal and minimal non-vanishing cohomology degrees of E . Then there exists composition of morphisms:

$$E \xrightarrow{\text{can}} \mathcal{H}^k(E)[-k] \xrightarrow{\iota_1} \mathcal{O}_p[-k] \xrightarrow{\iota_2} \mathcal{H}^l(E)[-k] \xrightarrow{\text{can}} E[l-k].$$

As both $\mathcal{H}^k(E)$ and $\mathcal{H}^l(E)$ are supported at p , we may let ι_i 's be non-zero. Their composition $\iota_2 \circ \iota_1$ is then non-zero as well.

The whole composition is non-zero because it induces a non-zero map from the term of E with maximal cohomology degree to the term of $E[l-k]$ with minimal cohomology degree. Since $k \geq l$, by the stability of E , we may only have $k = l$, and $E = \mathcal{H}^k(E)[-k] = \mathcal{O}_p[-k]$ which is stable. This leads to a contradiction. \square

Corollary 2.15. *Let X be a connected smooth projective variety such that its Albanese morphism is finite. Then all numerical stability conditions on $D^b(X)$ are geometric.*

Proof. It is a combination of Proposition 2.9 and Theorem 2.13. \square

2.4. Some applications. We give here several interesting applications of Corollary 2.15. The first one is a slightly weaker version of [KO18] and [Pir21, Theorem 1.4] on the indecomposibility of derived categories.

Proposition 2.16. *Let X be a connected smooth projective variety such that its Albanese morphism is finite. Then $D^b(X)$ does not admit any semiorthogonal decomposition for which each component admits a numerical stability condition.*

Proof. Suppose that there is a non-trivial semiorthogonal decomposition $D^b(X) = \langle \mathcal{D}_1, \mathcal{D}_2 \rangle$, and that \mathcal{D}_1 and \mathcal{D}_2 admit stability conditions σ_1 and σ_2 respectively. Since X is irreducible, $\langle \mathcal{D}_1, \mathcal{D}_2 \rangle$ is not a complete orthogonal decomposition. There exists σ_i -stable objects F_i satisfying $\text{Hom}(F_1, F_2) \neq 0$.

Replace σ_1 by its sufficiently negative integral shift, by [CP10, Theorem 3.6], one can have a glued stability condition σ of $\sigma_1[-m]$ and σ_2 satisfying $\phi_\sigma(F_2) - \phi_\sigma(F_1) > 2 \dim X + 2$.

By Corollary 2.15, the stability condition σ is geometric. By Lemma 2.11, for any two σ -stable objects F and G with $\phi_\sigma(G) - \phi_\sigma(F) \geq 2n + 1$, we have $\text{Hom}(F, G) = 0$. This leads to a contradiction. \square

Our second application concerns the stability of vector bundles on abelian varieties. Let A be an abelian variety. For every element $\frac{[L]}{t} \in \text{NS}(A) \otimes_{\mathbb{Z}} \mathbb{Q}$, there exist simple semi-homogeneous vector bundles with $\frac{\det}{\text{rk}} = \frac{[L]}{t}$ in the sense of Mukai [Muk78], see [Orl02, Section 4].

Corollary 2.17. *Let A be an abelian variety. Then all simple semi-homogeneous vector bundles are stable with respect to any numerical stability condition on $D^b(A)$.*

Proof. Denote by M the moduli space parameterizing all simple semi-homogeneous vector bundles with $\frac{\det}{\text{rk}} = \frac{[L]}{t}$. In particular, $M \cong A^\vee$ by [Muk78, Theorem 7.11]. Denote by $\mathcal{E}_{\frac{[L]}{t}}$ the universal family on $A \times M$. By [Orl02, Lemma 4.8], and [Bri99, Theorem 5.1 and 5.4], the Fourier–Mukai transformation $\Phi_{\mathcal{E}_{\frac{[L]}{t}}}$ is an equivalence between $D^b(A)$ and $D^b(M)$.

Note that the property of being stable with respect to any stability condition is preserved under any auto-equivalence. The skyscraper sheaves on M are mapped to all simple semi-homogeneous vector bundles with $\frac{\det}{\text{rk}} = \frac{[L]}{t}$. The statement now follows from Theorem 2.13. \square

3. SURFACE CASE

Let (X, H) be a polarized smooth projective variety. We fix the following surjection from $K(X)$, whose image, denoted by Λ_H , is clearly a lattice in \mathbb{R}^n :

$$\lambda_H : K(X) \twoheadrightarrow \Lambda_H; [E] \mapsto (H^n \text{rk}(E), H^{n-1} \text{ch}_1(E), H^{n-2} \text{ch}_2(E), \dots, \text{ch}_n(E)),$$

where n is the dimension of X . We denote the set of all (geometric) stability conditions with respect to Λ_H as $\text{Stab}_H(X)$ (resp. $\text{Stab}_H^{\text{Geo}}(X)$). In particular, if S is a surface of Picard rank one, then $\text{Stab}_H(S)$ is $\text{Stab}(S)$, the whole stability manifold of S .

3.1. Le Potier function. Recall that the H -slope of a coherent sheaf F on X is defined as

$$\mu_H(F) := \begin{cases} \frac{H^{n-1} \text{ch}_1(F)}{H^n \text{rk}(F)}, & \text{if } \text{rk}(F) > 0; \\ +\infty, & \text{if } \text{rk}(F) = 0. \end{cases}$$

A coherent sheaf F is called H -(semi)stable if for every proper non-zero subsheaf E , one has

$$\mu_H(E) < (\leq) \mu_H(F/E).$$

Definition 3.1. Let (S, H) be a polarized surface. We define the *Le Potier function* $\Phi_{S,H} : \mathbb{R} \rightarrow \mathbb{R}$ as:

$$\Phi_{S,H}(x) := \overline{\lim}_{\mu \rightarrow x} \left\{ \frac{\text{ch}_2(F)}{H^2 \text{rk}(F)} \mid F \text{ is } H\text{-semistable with } \mu_H(F) = \mu \right\}.$$

Proposition 3.2. *The Le Potier function is well-defined satisfying $\Phi_{S,H}(x) \leq \frac{x^2}{2}$. It is the smallest upper semi-continuous function satisfying*

$$\frac{\text{ch}_2(F)}{H^2 \text{rk}(F)} \leq \Phi_{S,H} \left(\frac{H \text{ch}_1(F)}{H^2 \text{rk}(F)} \right)$$

for every torsion-free H -stable (or semistable) sheaf F .

Proof. By [HL10, Theorem 5.2.5], for every rational number μ , there exists H -stable sheaf F with $\mu_H(F) = \mu$. The value of the function at every point is therefore in $\mathbb{R} \cup \{+\infty\}$.

The Bogomolov inequality states that

$$2 \text{rk}(F) \text{ch}_2(F) \leq \text{ch}_1(F)^2.$$

for every H -semistable sheaf F . By Hodge Index Theorem, we have

$$2H^2 \text{rk}(F) \text{ch}_2(F) \leq (H \text{ch}_1(F))^2$$

for every H -semistable sheaf F . When F is torsion-free, we can divided both sides by $(H^2 \text{rk}(E))^2$, which implies $\Phi_{S,H}(x) \leq \frac{x^2}{2}$. In particular, the function is well-defined.

The last statement is directly by the definition of $\Phi_{S,H}$. \square

Remark 3.3. The Le Potier function, which can be viewed as the sharp Bogomolov type inequality, is only known for very few polarized surfaces. When S is an abelian surface, it is known that $\Phi_{S,H}(x) = \frac{x^2}{2}$. When S is the projective plane, the function is known thanks to the work [DLP85]. The explicit formula for $\Phi_{\mathbb{P}^2}$ is very complicated, see [Li17] for more details. The function is also known or partially known for polarized K3, del Pezzo surfaces, and few sporadic surfaces, for example, the intersection of quadric and quintic $S_{2,5}$ in \mathbb{P}^4 , see [Li19]. We will discuss more properties of this function for hypersurfaces in a future project.

3.2. Tilting construction. In this section, we recall the tilting construction of stability conditions on polarized surfaces. More details are referred to the lecture notes [MS17, Section 6], or [AB13] for a more original treatment. The same construction gives weak stability conditions on threefolds or higher dimensional varieties. We will only use the threefold case in the next section. More details are referred to [BMT14, BBMT14].

Let (X, H) be a polarized variety. For every $\beta \in \mathbb{R}$, we define a pair of subcategories:

$$\mathcal{T}_\beta := \{F \in \text{Coh}(X) \mid \text{any } H\text{-semistable factor of } F \text{ satisfies } \mu_H(F) > \beta\};$$

$$\mathcal{F}_\beta := \{F \in \text{Coh}(X) \mid \text{any } H\text{-semistable factor of } F \text{ satisfies } \mu_H(F) \leq \beta\}.$$

This is a torsion pair on $\text{Coh}(X)$. We denote the tilted heart as

$$\text{Coh}^\beta(X) := \langle \mathcal{T}_\beta, \mathcal{F}_\beta[1] \rangle = \left\{ F \in \text{D}^b(X) \mid \begin{array}{l} \mathcal{H}^i(F) = 0 \text{ for } i \neq 0, -1; \\ \mathcal{H}^0(F) \in \mathcal{T}_\beta; \\ \mathcal{H}^{-1}(F) \in \mathcal{F}_\beta. \end{array} \right\}.$$

Recall from [MS17, Lemma 5.11] that giving a stability condition (\mathcal{P}, Z) on a triangulated category is equivalent to giving (\mathcal{A}, Z) , where $\mathcal{A} = \mathcal{P}((0, 1])$ is the heart structure.

Now let (S, H) be a polarized surface. For every $\alpha \in \mathbb{R}$, we define the central charge on $D^b(S)$ as

$$Z_{\alpha, \beta}(F) := (-\text{ch}_2(F) + \alpha H^2 \text{rk}(F)) + i(H \text{ch}_1(F) - \beta H^2 \text{rk}(F)).$$

Theorem 3.4 ([MS17, Theorem 6.10]). *For every $\alpha > \Phi_{S,H}(\beta)$, the pair $\sigma_{\alpha, \beta} = (\text{Coh}^\beta(S), Z_{\alpha, \beta})$ is a geometric stability condition (with respect to Λ_H) on S . Moreover, the map*

$$\Sigma : \{(\alpha, \beta) \in \mathbb{R}^2 \mid \alpha > \Phi_{S,H}(\beta)\} \rightarrow \text{Stab}_H(S) : (\alpha, \beta) \mapsto \sigma_{\alpha, \beta}$$

is a continuous embedding.

Remark 3.5. The theorem is slightly different as that in [MS17, Theorem 6.10] in the sense that we include stability conditions $\sigma_{\alpha, \beta}$ for some $\alpha \leq \frac{\beta^2}{2}$ when $\Phi_{S,H}(x) \neq \frac{x^2}{2}$. We briefly explain why the construction still works.

For every pair (α_0, β_0) satisfying $\alpha_0 > \Phi_{S,H}(\beta_0)$, since $\Phi_{S,H}$ is upper semi-continuous, there exists sufficiently small $\delta > 0$ satisfying

$$\delta(x - \beta_0)^2 + \alpha_0 - \delta \geq \Phi_{S,H}(x)$$

for every $x \in \mathbb{R}$. In other words, every H -semistable coherent sheaf F satisfies the Bogomolov type inequality:

$$\delta (H \text{ch}_1(F) - \beta H^2 \text{rk}(F))^2 - \text{rk}(F) (\text{ch}_2(F) - (\alpha_0 - \delta) \text{rk}(F)) \geq 0.$$

By exactly the same argument as that for [MS17, Theorem 6.10] (or see [PT19, Section 2] for a more general set-up), the pair $\sigma_{\alpha_0, \beta_0}$ is a stability condition and the embedding Σ is continuous at (α_0, β_0) .

Note that skyscraper sheaves are all simple in $\text{Coh}^{\beta_0}(S)$ by definition. They are all $\sigma_{\alpha_0, \beta_0}$ -stable with phase 1. Hence the stability condition $\sigma_{\alpha_0, \beta_0}$ is geometric.

3.3. Geometric stability conditions on surfaces.

Proposition 3.6 ([Bri08, Section 10]). *Let $\sigma = (\mathcal{P}, Z)$ be a geometric stability condition in $\text{Stab}_H^{\text{Geo}}(S)$. Then $\sigma = \sigma_{\alpha, \beta} g$ for some $\alpha > \Phi_{S,H}(\beta)$ and $g \in \widetilde{\text{GL}}_2^+(\mathbb{R})$.*

Proof. Applying an element of $\widetilde{\text{GL}}_2^+(\mathbb{R})$ one can assume that $Z(\mathcal{O}_p) = -1$ and $\mathcal{O}_p \in \mathcal{P}(1)$ for all $p \in X$. In particular, the central charge is of the form

$$-\text{ch}_2 + aH \text{ch}_1 + bH^2 \text{rk} + i(cH \text{ch}_1 + dH^2 \text{rk})$$

for some $a, b, c, d \in \mathbb{R}$.

By [Bri08, Lemma 10.1.c], the torsion sheaf $\mathcal{O}_C(mH) \in \mathcal{P}((0, 1])$ for every curve C on S and $m \in \mathbb{Z}$. Therefore, the coefficient $c > 0$. Applying an element of $\widetilde{\text{GL}}_2^+(\mathbb{R})$ again, we may assume the central charge is of the form $Z_{\alpha, \beta}$ for some $\alpha, \beta \in \mathbb{R}$. By [MS17, Lemma 6.20] (note that the argument holds for all $\alpha \in \mathbb{R}$), the heart $\mathcal{P}((0, 1]) = \text{Coh}^\beta(S)$.

Now we only need to show that $\alpha > \Phi_{S,H}(\beta)$. By [MS17, Proposition 5.27], there is an open neighbourhood U of σ in $\text{Stab}_H(S)$ where all skyscraper sheaves are stable. By Theorem 2.6,

there is an open neighborhood W of (α, β) in \mathbb{R}^2 such that for every $(\alpha', \beta') \in W$, there exists a stability condition $\sigma' = (\mathcal{P}', Z') \in U$ with

$$\ker Z' = (1, \beta', \alpha') \cdot \mathbb{R} \subset \Lambda_H \otimes \mathbb{R}.$$

Suppose $\alpha \leq \Phi_{S,H}(\beta)$, by Definition 3.1, there exists an H -semistable sheaf F and $(\alpha_0, \beta_0) \in W$ such that

$$H \operatorname{ch}_1(F) = \beta_0 H^2 \operatorname{rk}(F) \text{ and } \operatorname{ch}_2(F) > \alpha_0 H^2 \operatorname{rk}(F).$$

Let $\sigma_0 = (\mathcal{P}_0, Z_0)$ be a stability condition in U with $\ker Z_0 = (1, \beta_0, \alpha_0) \cdot \mathbb{R}$. Applying an element of $\widetilde{\operatorname{GL}}_2^+(\mathbb{R})$ on σ_0 , one can assume that $Z_0 = Z_{\alpha_0, \beta_0}$. By [MS17, Lemma 6.20], $\mathcal{P}_0((0, 1]) = \operatorname{Coh}^{\beta_0}(S) \ni F[1]$. This leads to a contradiction that $Z_{\alpha_0, \beta_0}(F[1]) \in \mathbb{R}_{>0}$. Therefore, we must have $\alpha > \Phi_{S,H}(\beta)$. \square

Corollary 3.7. *Let (S, H) be a smooth polarized surface such that its Albanese morphism is finite, then $\operatorname{Stab}_H(S)$ is connected and contractible.*

Proof. By Corollary 2.15 and Proposition 3.6, $\operatorname{Stab}_H(S)$ is homeomorphic to a $\widetilde{\operatorname{GL}}_2^+(\mathbb{R})$ -principal bundle over $\{(\alpha, \beta) \in \mathbb{R}^2 \mid \alpha > \Phi_{S,H}(\beta)\}$, which is contractible. \square

Corollary 3.8. *The space of stability conditions of an irregular surface with Picard rank one is connected and contractible.*

Proof. The Albanese morphism of the irregular surface S is non-trivial. It does not contract any curve, since otherwise its Picard rank is at least 2. Let H be an ample divisor, then $\operatorname{Stab}(S) = \operatorname{Stab}_H(S)$, which is contractible by Corollary 3.7. \square

4. ABELIAN THREEFOLD CASE

Let (A, H) be a polarized abelian threefold. In this section, we show that the principal component of the stability manifold of A as that constructed in [BMS16, Theorem 9.1] is the whole space $\operatorname{Stab}_H(A)$.

4.1. Review: stability conditions on abelian threefolds. We briefly recall the construction of stability conditions on abelian threefolds as that in [MP13, MP15] and [BMS16].

For every $\beta \in \mathbb{R}$, recall from the previous section that we have the heart

$$\operatorname{Coh}^\beta(A) := \langle \mathcal{T}_\beta, \mathcal{F}_\beta[1] \rangle.$$

We will always consider the twisted Chern character $\operatorname{ch}^\beta(F) := e^{-\beta H} \operatorname{ch}(F)$. More explicitly, we have

$$\begin{aligned} \operatorname{ch}_1^\beta &= \operatorname{ch}_1 - \beta H \operatorname{rk}; & \operatorname{ch}_2^\beta &= \operatorname{ch}_2 - \beta H \operatorname{ch}_1 + \frac{\beta^2}{2} H^2 \operatorname{rk}; \\ \operatorname{ch}_3^\beta &= \operatorname{ch}_3 - \beta H \operatorname{ch}_2 + \frac{\beta^2}{2} H^2 \operatorname{ch}_1 - \frac{\beta^3}{6} H^3 \operatorname{rk}. \end{aligned}$$

Note that for every object F in $\operatorname{Coh}^\beta(A)$, we have $H^2 \operatorname{ch}_1^\beta(F) \geq 0$.

For every $\alpha > 0$, we consider the slope function $\mu_{\alpha,\beta}$ on $\text{Coh}^\beta(X)$ defined as

$$\mu_{\alpha,\beta}(F) := \begin{cases} \frac{H \text{ch}_2^\beta(F) - \frac{\alpha^2}{2} \text{rk}(F)}{H^2 \text{ch}_1^\beta(F)}, & \text{if } H^2 \text{ch}_1^\beta(F) > 0; \\ +\infty, & \text{if } H^2 \text{ch}_1^\beta(F) = 0. \end{cases}$$

A non-zero object $F \in \text{Coh}^\beta(A)$ is called $\mu_{\alpha,\beta}$ -(semi)stable if for every proper non-zero subobject $E \hookrightarrow F$ in $\text{Coh}^\beta(A)$, one has

$$\mu_{\alpha,\beta}(E) < (\leq) \mu_{\alpha,\beta}(F/E).$$

Consider the pair of subcategories:

$$\mathcal{T}_{\alpha,\beta} := \{F \in \text{Coh}^\beta(A) \mid \text{any quotient object } F \twoheadrightarrow G \text{ in } \text{Coh}^\beta(A) \text{ satisfies } \mu_{\alpha,\beta}(G) > 0\};$$

$$\mathcal{F}_{\alpha,\beta} := \{F \in \text{Coh}^\beta(A) \mid \text{any non-zero subobject } E \hookrightarrow F \text{ in } \text{Coh}^\beta(A) \text{ satisfies } \mu_{\alpha,\beta}(E) \leq 0\}.$$

This is a torsion pair on $\text{Coh}^\beta(A)$. We denote the tilted heart as

$$\text{Coh}^{\alpha,\beta}(A) := \langle \mathcal{T}_{\alpha,\beta}, \mathcal{F}_{\alpha,\beta}[1] \rangle.$$

For every $b \in \mathbb{R}$ and $a > \frac{\alpha^2}{6} + \frac{1}{2}|b|\alpha$, we define the central charge as

$$Z_{\alpha,\beta}^{a,b} := -\text{ch}_3^\beta + bH \text{ch}_2^\beta + aH^2 \text{ch}_1^\beta + i \left(H \text{ch}_2^\beta - \frac{1}{2}\alpha^2 H^3 \text{rk} \right).$$

We will write $\mu_{\alpha,\beta}^{a,b} := -\frac{\text{Re}(Z_{\alpha,\beta}^{a,b})}{\text{Im}(Z_{\alpha,\beta}^{a,b})}$ for the slope function. Now we may summarize the construction of stability conditions on $D^b(A)$ as follows.

Theorem 4.1 ([BMS16, Section 8 and Theorem 9.1]). *Let (A, H) be a polarized abelian threefold. Then for every $\alpha, \beta, a, b \in \mathbb{R}$ satisfying $\alpha > 0$ and $a > \frac{1}{6}\alpha^2 + \frac{1}{2}|b|\alpha$, the pair $\sigma_{\alpha,\beta}^{a,b} = (\text{Coh}^\beta(A), Z_{\alpha,\beta}^{a,b})$ is a stability condition on $D^b(A)$ in $\text{Stab}_H(A)$. Moreover, the map*

$$\begin{aligned} \Sigma : \{(a, b, \alpha, \beta) \in \mathbb{R}^4 \mid \alpha > 0, a > \frac{\alpha^2}{6} + \frac{1}{2}|b|\alpha\} &\rightarrow \text{Stab}_H(A) \\ (a, b, \alpha, \beta) &\mapsto \sigma_{\alpha,\beta}^{a,b} \end{aligned}$$

is a continuous embedding.

Denote $\tilde{\mathfrak{P}} := \widetilde{\text{GL}}_2^+(\mathbb{R}) \cdot (\text{im}(\Sigma))$, then $\text{im}(\Sigma)$ is a slice of the $\widetilde{\text{GL}}_2^+(\mathbb{R})$ -action on $\tilde{\mathfrak{P}}$. The space $\tilde{\mathfrak{P}}$ is a connected component in $\text{Stab}_H(A)$.

Our goal is to show that $\tilde{\mathfrak{P}}$ is actually the unique connected component of $\text{Stab}_H(A)$.

4.2. Semi-homogeneous vector bundles. Let (A, H) be a polarized abelian threefold. We will make use of simple semi-homogeneous vector bundles with $\frac{\det}{\text{rk}} = sH$ for $s \in \mathbb{Q}$. To simplify the notations, we will denote such a simple semi-homogeneous vector bundle as E_s .

Write $s = \frac{p}{q}$ for some $q \in \mathbb{Z}_{>0}$ and q, p coprime. The Chern characters of E_s are

$$\text{ch}(E_s) = \text{rk}(E_s)e^{sH} = \text{rk}(E_s)(1, sH, \frac{s^2}{2}H^2, \frac{s^3}{6}H^3).$$

For every object $F \in D^b(A)$, it follows from Hirzebruch–Riemann–Roch that

$$\chi(E_s, F) = \text{rk}(E_s) \text{ch}_3^s(F).$$

Every E_s can be constructed as the push-forward of line bundles via an isogeny $Y \rightarrow X$, see [Muk78, Theorem 5.8].

Lemma 4.2. *Let (A, H) be a polarized abelian threefold. If $s < t$, then $\text{Hom}(E_s, E_t[i]) \neq 0$ if and only if $i = 0$.*

Proof. By Corollary 2.17, both E_s and E_t are $\sigma_{\alpha, \beta}^{a, b}$ -stable for $\beta = \frac{s+t}{2}$, $\alpha < \frac{t-s}{2}$, $b = 0$, and $a = \frac{\alpha^2}{6} + \epsilon$. In particular, $E_t, E_s[2] \in \text{Coh}^{\beta, \alpha}(A)$, and

$$\mu_{\alpha, \beta}^{\frac{\alpha^2}{6} + \epsilon, 0}(E_s[2]) = \frac{s}{3} < \frac{t}{3} = \mu_{\alpha, \beta}^{\frac{\alpha^2}{6} + \epsilon, 0}(E_t)$$

when ϵ tends to 0. Therefore, $\text{Hom}(E_t, E_s[i]) = 0$ when $i \leq 2$. By Serre duality, $\text{Hom}(E_s, E_t[i]) = 0$ unless $i = 0$.

By the HRR formula

$$\chi(E_s, E_t) = \text{rk}(E_s) \text{rk}(E_t) \frac{(t-s)^3}{6} > 0.$$

So $\text{Hom}(E_t, E_s) \neq 0$. □

Denote by \mathcal{C} the set of all (shifted) skyscraper sheaves $\mathcal{O}_p[t]$ and all simple semi-homogeneous vector bundle $E_s[t]$'s. By Corollary 2.15 and 2.17, all elements in \mathcal{C} are stable with respect to every stability condition. The following proposition is the key input to our main result.

Proposition 4.3. *Let $\sigma = \sigma_{\alpha, \beta}^{a, b}$ be a stability condition on $D^b(A)$ and $F \in \mathcal{P}_\sigma(\theta)$. Then*

$$\begin{aligned} \inf_{E \in \mathcal{C}} \{\theta - \phi_\sigma(E) \mid \text{Hom}(E, F) \neq 0\} &\leq 1; \\ \inf_{E \in \mathcal{C}} \{\phi_\sigma(E) - \theta \mid \text{Hom}(F, E) \neq 0\} &\leq 1. \end{aligned}$$

Proof. We may assume that F is σ -stable and is not in \mathcal{C} . Since the set \mathcal{C} is closed under homological shift, we may also assume $\theta \in (0, 1]$.

We first deal with the case when $\theta \in (0, 1)$ and $H^{3-\bullet} \text{ch}_\bullet(F) \notin \mathbb{R} \cdot (1, t, \frac{t^2}{2}, \frac{t^3}{6})$. Denote by $C := \mu(F) = -\cot(\pi\theta) \in \mathbb{R}$ the slope of F . It follows that

$$(1) \quad \text{ch}_3^\beta(F) - bH \text{ch}_2^\beta(F) - aH^2 \text{ch}_1^\beta(F) + C \left(H \text{ch}_2^\beta(F) - \frac{1}{2} \alpha^2 H^3 \text{rk}(F) \right) = 0.$$

Consider the equation

$$(2) \quad f(x) := \frac{x^3}{6} + (C - b) \frac{x^2}{2} - ax - \frac{1}{2} \alpha^2 C = 0.$$

By the assumption that $a > \frac{1}{6} \alpha^2 + \frac{1}{2} |b| \alpha$ and $\alpha > 0$, we have

$$f(\alpha) = \alpha \left(\frac{1}{6} \alpha^2 - \frac{1}{2} b \alpha - a \right) < 0 < \alpha \left(a - \frac{1}{6} \alpha^2 - \frac{1}{2} b \alpha \right) = f(-\alpha).$$

Note that $\lim_{x \rightarrow \pm\infty} f(x) = \pm\infty$, we have

$$f(s_j) = 0 \text{ for some } s_1 < -\alpha < s_2 < \alpha < s_3.$$

It worth to notice that $f(x) = 0$ if and only if $\mu(e^{(\beta+x)H}) = C$.

Now assume that $\text{ch}_3^{\beta+s_1}(F) > 0$, we will show that under this assumption, we have

$$\inf_{E \in \mathcal{C}} \{\phi_\sigma(E) - \theta | \text{Hom}(F, E) \neq 0\} \leq 1.$$

Remark 4.4. *When $\beta + s_1$ is rational, we may simply notice that*

$$\chi(F, E_{\beta+s_1}) = -\text{rk}(E_{\beta+s_1}) \text{ch}_3^{\beta+s_1}(F) < 0.$$

It then follows that $\text{Hom}(F, E_{\beta+s_1}[3]) \neq 0$, which implies the conclusion.

As $\beta + s_1$ can be an irrational number, we need to deform the stability condition so that there exists $E_{\beta+s'_1}$ in \mathcal{C} . Yet this deformation is not completely straightforward, as at certain point we need the assumption that $H^{3-\bullet} \text{ch}_\bullet(F) \notin \mathbb{R} \cdot (1, t, \frac{t^2}{2}, \frac{t^3}{6})$. The only purpose of next few paragraphs is to deal with this issue. For the convenience of the readers, it is harmless to skip these paragraphs and simply assume that $E_{\beta+s_i} \in \mathcal{C}$ exists.

By [MS17, Proposition 5.27], there is an open neighbourhood W of (α, β, a, b) such that F is $\sigma_{\alpha', \beta'}^{a', b'}$ -stable for every $(\alpha', \beta', a', b') \in W$. For every $\epsilon > 0$, there exists $\delta_0 > 0$ such that for every $|\delta|, |\delta'|, |\delta''| < \delta_0$ we have

- $(\alpha + \delta'', \beta, a + \delta, b + \delta') \in W$;
- $\text{ch}_3^{\beta+s_1+\delta}(F) > 0$;
- $\text{dist}(\sigma_{\alpha, \beta}^{a, b}, \sigma_{\alpha, \beta}^{a+\delta, b+\delta'}) < \frac{\epsilon}{2}$.

It follows from $f(s_1) = 0$ that $\mu(e^{(\beta+s_1)H}) = C$. Note that the function $B(x)$ satisfying

$$\mu_{\alpha, \beta}^{a, B(x)}(e^{(\beta+x)H}) = \mu_{\alpha, \beta}^{a, B(x)}(F)$$

is well-defined and continuous when $x \neq \pm \sqrt{\frac{2H \text{ch}_2^\beta(F)}{H^3 \text{rk}(F)}}$. When $s_1 \neq \pm \sqrt{\frac{2H \text{ch}_2^\beta(F)}{H^3 \text{rk}(F)}}$, $B(s_1) = b$. There exists $|\delta| < \delta_0$ such that $E_{\beta+s_1+\delta} \in \mathcal{C}$ and $|B(s_1 + \delta) - b| < \delta_0$.

If $s_1 = \pm \sqrt{\frac{2H \text{ch}_2^\beta(F)}{H^3 \text{rk}(F)}}$ but $s_1 \neq \frac{H^2 \text{ch}_1^\beta(F)}{H^3 \text{rk}(F)}$, then the function $A(x)$ satisfying

$$\mu_{\alpha, \beta}^{A(x), b}(e^{(\beta+x)H}) = \mu_{\alpha, \beta}^{A(x), b}(F)$$

is well-defined and continuous at a small neighbourhood of $x = s_1$. There exists $|\delta| < \delta_0$ such that $E_{\beta+s_1+\delta} \in \mathcal{C}$ and $|A(s_1 + \delta) - a| < \delta_0$.

If $s_1 = \pm \sqrt{\frac{2H \text{ch}_2^\beta(F)}{H^3 \text{rk}(F)}} = \frac{H^2 \text{ch}_1^\beta(F)}{H^3 \text{rk}(F)}$, since $H^{3-\bullet} \text{ch}_\bullet(F) \notin \mathbb{R} \cdot (1, t, \frac{t^2}{2}, \frac{t^3}{6})$, we may deform α as that for a or b .

In any case, we may assume there exists $E_{\beta+s'_1} \in \mathcal{C}$ and (α', β, a', b') such that

- $s'_1 < -\alpha'$, $E_{\beta+s'_1}[2] \in \mathcal{P}_\sigma((0, 1])$ and $\text{ch}^{\beta+s'_1}(F) > 0$;

- F is $\sigma_{\alpha',\beta}^{a',b}$ -stable and $\mu_{\alpha',\beta}^{a',b'}(F) = \mu_{\alpha',\beta}^{a',b'}(E_{\beta+s'_1})$;
- $\left| \phi_\sigma(E_{\beta+s'_1}[2]) - \phi_\sigma(F) \right| < \epsilon$.

Note that $\phi_{\alpha',\beta}^{a',b'}(F) = \phi_{\alpha',\beta}^{a',b'}(E_{\beta+s'_1}[2])$ and $F \neq E_{\beta+s'_1}[2]$, we have

$$\mathrm{Hom}(E_{\beta+s'_1}, F[t]) = (\mathrm{Hom}(F, E_{\beta+s'_1}[3-t]))^* = 0$$

for all $t \neq 0, -1$. Therefore,

$$\begin{aligned} & \mathrm{Hom}(F, E_{\beta+s'_1}[4]) - \mathrm{Hom}(F, E_{\beta+s'_1}[3]) \\ &= \chi(F, E_{\beta+s'_1}) = -\mathrm{rk}(E_{\beta+s'_1}) \mathrm{ch}^{\beta+s'_1}(F) < 0. \end{aligned}$$

It follows that $\mathrm{Hom}(F, E_{\beta+s'_1}[3]) \neq 0$. In particular,

$$\inf_{E \in \mathcal{C}} \{ \phi_\sigma(E) - \theta \mid \mathrm{Hom}(F, E) \neq 0 \} \leq \phi_\sigma(E_{\beta+s'_1}[3]) - \theta < 1 + \epsilon.$$

As ϵ tends to 0, we have $\inf_{E \in \mathcal{C}} \{ \phi_\sigma(E) - \theta \mid \mathrm{Hom}(F, E) \neq 0 \} \leq 1$.

By the same argument, if any value of $\{(-1)^{i+1} \mathrm{ch}_3^{\beta+s_i}(F)\}_{i=1,2,3}$ is positive, then

$$\inf_{E \in \mathcal{C}} \{ \phi_\sigma(E) - \theta \mid \mathrm{Hom}(F, E) \neq 0 \} \leq 1.$$

If any value of $\{(-1)^{i+1} \mathrm{ch}_3^{\beta+s_i}(F)\}_{i=1,2,3}$ is negative, then

$$\inf_{E \in \mathcal{C}} \{ \theta - \phi_\sigma(E) \mid \mathrm{Hom}(E, F) \neq 0 \} \leq 1.$$

To conclude the argument for the case that $H^{3-\bullet} \mathrm{ch}_\bullet(F) \notin \mathbb{R} \cdot (1, t, \frac{t^2}{2}, \frac{t^3}{6})$ and $\theta \neq 1$, we need the following property for the values of $\mathrm{ch}_3^{\beta+s_i}(F)$.

Lemma 4.5. *The values $\{(-1)^j \mathrm{ch}_3^{\beta+s_i}(F)\}_{i=1,2,3}$ cannot be all positive (or negative) at the same. Moreover, if all of them are non-negative (or non-positive), then they must be all 0 and*

$$(H^3 \mathrm{rk}(F), H^2 \mathrm{ch}_1^\beta(F), H \mathrm{ch}_2^\beta(F), \mathrm{ch}_3^\beta(F)) = \lambda \cdot (1, b - C, -a, -\frac{C}{2}\alpha^2)$$

for some $\lambda \in \mathbb{R}^\times$.

The proof for the lemma is elementary, we postpone it after the proof of this proposition.

By Lemma 4.5, the only outstanding case is when $\mathrm{ch}_3^{\beta+s_i}(F)$ are all zero. In this case, for every $\epsilon > 0$, we may deform a to an $a' \neq a$ such that

- $\mathrm{dist}(\sigma, \sigma_{\alpha,\beta}^{a',b}) < \epsilon$;
- F is $\sigma_{\alpha,\beta}^{a',b}$ -stable and is not in $\mathcal{P}_{\sigma_{\alpha,\beta}^{a',b}}(1)$.

The characters of F cannot satisfy the condition

$$(H^3 \mathrm{rk}(F), H \mathrm{ch}_2^\beta(F)) = \lambda \cdot (1, -a')$$

as that in Lemma 4.5 for any $\lambda \in \mathbb{R}^\times$. Therefore,

$$\begin{aligned} & \inf_{E \in \mathcal{C}} \{ \phi_\sigma(E) - \phi_\sigma(F) \mid \text{Hom}(F, E) \neq 0 \} \\ & < \inf_{E \in \mathcal{C}} \left\{ \phi_{\sigma_{\alpha, \beta}^{a', b}}(E) - \phi_{\sigma_{\alpha, \beta}^{a', b}}(F) + 2\epsilon \mid \text{Hom}(F, E) \neq 0 \right\} \leq 1 + 2\epsilon. \end{aligned}$$

As ϵ tends to 0, we get the inequality for σ . The statement holds for all $F \in \mathcal{P}_\sigma(\theta)$ with $\theta \notin \mathbb{Z}$ and $H^{3-\bullet} \text{ch}_\bullet(F) \neq \mathbb{R} \cdot (1, t, \frac{t^2}{2}, \frac{t^3}{6})$.

In the case that $\theta = 1$, if F is a skyscraper sheaf, then the statement hold automatically as $F \in \mathcal{C}$. Otherwise, $\text{ch}(F) \neq (0, 0, 0, a)$, we can deform σ in any small open neighbourhood so that F is still stable but not with phase 1. The inequalities in the statement hold.

In the case that $H^{3-\bullet} \text{ch}_\bullet(F) = \mathbb{R} \cdot (1, t, \frac{t^2}{2}, \frac{t^3}{6})$ for some $t \in \mathbb{R}$, we have $Q_K^\beta(H^{3-\bullet} \text{ch}_\bullet(F)) = 0$ for all quadratic form $Q_K^\beta = K \Delta_H + \nabla_H^\beta$. By [BMS16, Proposition A.8], they are stable with respect to all stability conditions as that in Theorem 4.1. In particular, $F[-i] = F' \in \text{Coh}(A)$ where $i = 0$ when $t > \beta + \alpha$; $i = 1$ when $\beta - \alpha < t \leq \beta + \alpha$; and $i = 2$ when $t \leq \beta - \alpha$. By the same argument as that in Lemma 4.2, $\text{Hom}(E_s, F') \neq 0$ when $s < t$; and $\text{Hom}(F', E_s) \neq 0$ when $s > t$. It is clear that both infimums for F in the statement are 0. \square

Proof for Lemma 4.5. As s_i 's are the solutions to $f(x) = 0$, we have

$$\begin{aligned} (3) \quad & s_1 + s_2 + s_3 = 3b - 3C; \\ (4) \quad & s_1 s_2 + s_2 s_3 + s_3 s_1 = -6a; \\ (5) \quad & s_1 s_2 s_3 = 3\alpha^2 C. \end{aligned}$$

Substitute (1) into $\text{ch}_3^{\beta+s_i}(F)$'s, we may replace the term $\text{ch}_3^\beta(F)$ by other terms:

$$(6) \quad \text{ch}_3^{\beta+s_i}(F) = (b - C - s_i)H \text{ch}_2^\beta(F) + (a + \frac{s_i^2}{2})H^2 \text{ch}_1^\beta(F) + (\frac{1}{2}C\alpha^2 - \frac{s_i^3}{6})H^3 \text{rk}(F).$$

To simplify the notations, we denote

$$(7) \quad L_i(x, y, z) := (b - C - s_i)x + (a + \frac{s_i^2}{2})y + (\frac{1}{2}C\alpha^2 - \frac{s_i^3}{6})z.$$

Substitute $(x, y, z) = (\frac{s_i^2}{2}, s_i, 1)$ into L_j 's, we have

$$\begin{aligned} L_j(\frac{s_i^2}{2}, s_i, 1) &= (b - C)\frac{s_i^2}{2} - \frac{s_j s_i^2}{2} + a s_i + \frac{s_j^2 s_i}{2} + \frac{\alpha^2 C}{2} - \frac{s_j^3}{6} \\ &= \frac{s_i^3}{6} - \frac{s_j s_i^2}{2} + \frac{s_j^2 s_i}{2} - \frac{s_j^3}{6} \\ &= \frac{1}{6}(s_i - s_j)^3. \end{aligned}$$

Substitute $(x, y, z) = (-a, b - C, 1)$ into L_i 's, we have

$$\begin{aligned} L_i(-a, b - C, 1) &= as_i + \frac{s_i^2}{2}(b - C) + \frac{1}{2}C\alpha^2 - \frac{s_i^3}{6} \\ &= -\frac{1}{6}((s_1s_2 + s_2s_3 + s_3s_1)s_i - s_i^2(s_1 + s_2 + s_3) - s_1s_2s_3 + s_i^3) \\ &= 0. \end{aligned}$$

Therefore, L_i 's are linear dependant, and we have

$$(s_3 - s_1)^3 L_2 = (s_3 - s_2)^3 L_1 + (s_2 - s_1)^3 L_3.$$

Since $s_1 < s_2 < s_3$, the values $\{(-1)^i L_i(x, y, z)\}$ are all non-negative (or non-positive) if and only if $(x, y, z) \in \mathbb{R} \cdot (-a, b - C, 1)$. \square

4.3. Stability manifold of abelian threefolds. Given stability conditions σ_1 and σ_2 on $D^b(X)$, we may consider the following generalised metric function on them as that defined in [Bri07, Section 6]:

$$d(\sigma_1, \sigma_2) := \sup_{0 \neq E \in D^b(X)} \{|\phi_{\sigma_1}^-(E) - \phi_{\sigma_2}^-(E)|, |\phi_{\sigma_1}^+(E) - \phi_{\sigma_2}^+(E)|\} \in [0, +\infty].$$

Lemma 4.6. *Let σ and τ be two stability conditions and $\delta \in [0, 1]$ such that for every σ -stable object E we have*

$$\phi_{\tau}^+(E) \geq \phi_{\sigma}(E) - \delta \text{ and } \phi_{\tau}^-(E) \leq \phi_{\sigma}(E) + \delta.$$

Then $d(\sigma, \tau) \leq \delta$.

Proof. For every τ -stable object F , we have the distinguished triangle

$$(8) \quad E \rightarrow F \rightarrow G \xrightarrow{+},$$

where $E = \text{HN}_{\sigma}^{\phi_{\sigma}^+(F)}(F)$ and $G = \text{HN}_{\sigma}^{<\phi_{\sigma}^+(F)}(F)$.

Suppose $\phi_{\sigma}^+(F) > \phi_{\tau}(F) + \delta$. Then by the assumption of the statement,

$$\phi_{\tau}^+(E) \geq \phi_{\sigma}(E) - \sigma > \max\{\phi_{\tau}(F), \phi_{\sigma}^+(G) - 1\}.$$

Apply $\text{Hom}(\text{HN}_{\tau}^{\phi_{\tau}^+(E)}(E), -)$ to (8), we have an exact sequence

$$\rightarrow 0 = \text{Hom}(\text{HN}_{\tau}^{\phi_{\tau}^+(E)}(E), G[-1]) \rightarrow \text{Hom}(\text{HN}_{\tau}^{\phi_{\tau}^+(E)}(E), E) \rightarrow \text{Hom}(\text{HN}_{\tau}^{\phi_{\tau}^+(E)}(E), F) = 0 \rightarrow,$$

which is a contradiction. Therefore, we must have $\phi_{\sigma}^+(F) \leq \phi_{\tau}(F) + \delta$ for every τ -stable object F . For the same reason, $\phi_{\sigma}^-(F) \leq \phi_{\tau}(F) - \delta$ for every τ -stable object F . By [Bri07, Lemma 6.1], we have $d(\sigma, \tau) \leq \delta$. \square

Lemma 4.7. *Let σ and τ be two stability conditions with the same central charge. If $d(\sigma, \tau) \leq 1$, then $\sigma = \tau$.*

Proof. For every σ -stable object F , we first show that $\phi_{\tau}^+(F) \geq \phi_{\sigma}(F)$.

Suppose $\phi_{\tau}^+(F) < \phi_{\sigma}(F)$, then since $d(\sigma, \tau) \leq 1$, the object $F \in \mathcal{P}_{\tau}([\phi_{\sigma}(F) - 1, \phi_{\sigma}(F)])$. Therefore,

$$Z_{\tau}(F) = me^{\pi i \theta} \text{ for some } m > 0 \text{ and } \theta \in [\phi_{\sigma}(F) - 1, \phi_{\sigma}(F)],$$

which cannot equal $Z_\sigma(F) = m'e^{\pi i\phi_\sigma(F)}$ for any m' . It follows that $\phi_\tau^+(F) \geq \phi_\sigma(F)$. Due to the same argument, for every τ -stable object E , $\phi_\sigma^+(E) \geq \phi_\tau(E)$.

For every σ -stable object F , we then show that $\phi_\tau^+(F) = \phi_\sigma(F)$. Note that we have the distinguished triangle

$$(9) \quad E \rightarrow F \rightarrow G \xrightarrow{+},$$

where $E = \mathrm{HN}_\tau^{\phi_\tau^+(F)}(F)$ and $G = \mathrm{HN}_\tau^{<\phi_\tau^+(F)}(F)$. Suppose $\phi_\tau^+(F) > \phi_\sigma(F)$, then

$$\phi_\sigma^+(E) \geq \phi_\tau(E) = \phi_\tau^+(F) > \phi_\sigma(F).$$

In particular, we have $\mathrm{Hom}(\mathrm{HN}_\sigma^{\phi_\sigma^+(E)}(E), F) = 0$.

Since $d(\sigma, \tau) \leq 1$, we have

$$\phi_\sigma^+(G[-1]) < \phi_\tau^+(F) \leq \phi_\sigma^+(E).$$

Therefore, we have $\mathrm{Hom}(\mathrm{HN}_\sigma^{\phi_\sigma^+(E)}(E), G[-1]) = 0$. Apply $\mathrm{Hom}(\mathrm{HN}_\sigma^{\phi_\sigma^+(E)}(E), -)$ to (9), we get the contradiction. Hence, we must have $\phi_\tau^+(F) = \phi_\sigma(F)$.

By the same argument, $\phi_\tau^-(F) = \phi_\sigma(F)$ for every σ -stable object F . By [Bri07, Lemma 6.1], $d(\sigma, \tau) = 0$. As σ and τ also have the same central charge, they are the same stability condition. \square

Theorem 4.8. *Let (A, H) be a polarized abelian threefold and $\sigma = (\mathcal{P}, Z)$ be a stability condition in $\mathrm{Stab}_H(A)$. Then $\sigma = \sigma_{\alpha, \beta}^{a, b} g$ for some $\alpha > 0$, $a > \frac{1}{6}\alpha^2 + \frac{1}{2}|b|\alpha$, and $g \in \widetilde{\mathrm{GL}}_2^+(\mathbb{R})$.*

Proof. Applying an element of $\widetilde{\mathrm{GL}}_2^+(\mathbb{R})$ one can assume that $Z(\mathcal{O}_p) = -1$ and $\mathcal{O}_p \in \mathcal{P}(1)$ for all $p \in A$. In particular, the central charge is of the form

$$Z = -\mathrm{ch}_3 + l_1 H \mathrm{ch}_2 + l_2 H^2 \mathrm{ch}_1 + l_3 H^3 \mathrm{rk} + i(m_1 H \mathrm{ch}_2 + m_2 H^2 \mathrm{ch}_1 + m_3 H^3 \mathrm{rk})$$

for some $l_i, m_i \in \mathbb{R}$.

We first show that $m_1 > 0$. Suppose $m_1 \leq 0$, then we may deform the stability condition so that $\mathcal{O}_p \in \mathcal{P}(1)$ and $m_1 < 0$. Note that

$$\lim_{t \rightarrow \pm\infty} \mathrm{Im}(Z(e^{tH})) = -\infty,$$

it follows from Corollary 2.17, Lemma 4.2 and 2.11 that all $E_t \in \mathcal{C}$ are in the heart $\mathcal{P}((-1, 0])$. Note that

$$\lim_{t \rightarrow +\infty} -\frac{\mathrm{Re}(Z(e^{tH}))}{\mathrm{Im}(Z(e^{tH}))} = -\infty \text{ and } \lim_{t \rightarrow -\infty} -\frac{\mathrm{Re}(Z(e^{tH}))}{\mathrm{Im}(Z(e^{tH}))} = +\infty,$$

we have $\lim_{t \rightarrow +\infty} (\phi_\sigma(E_t)) = -1$ and $\lim_{t \rightarrow -\infty} (\phi_\sigma(E_t)) = 0$. This contradicts with Corollary 2.17 and Lemma 4.2. Therefore, we must have $m_1 > 0$.

Applying an element of $\widetilde{\mathrm{GL}}_2^+(\mathbb{R})$ we may further assume that the central charge is of the form

$$Z = -\mathrm{ch}_3^\beta + bH \mathrm{ch}_2^\beta + aH^2 \mathrm{ch}_1^\beta + cH^3 \mathrm{rk} + i(H \mathrm{ch}_2^\beta + dH^3 \mathrm{rk})$$

for some $a, b, c, d, \beta \in \mathbb{R}$.

We then show that d must be positive. Suppose $d \geq 0$, we may deform the stability condition so that $d < 0$. Note that $\text{Im}(Z(e^{tH})) > 0$ for all $t \in \mathbb{Q}$, it follows from Corollary 2.17 and Lemma 4.2 that there exists $\beta' \in \mathbb{R} \cup \{\pm\infty\}$ such that $E_t \in \mathcal{P}((-2, -1])$ for all $t < \beta'$ and $E_t \in \mathcal{P}((0, 1])$ for all $t > \beta'$.

We show that β' must be β in this case. Suppose $\beta' < \beta$, then there exists $\frac{p}{q}$ such that $\beta' < \frac{p}{q} < \frac{p+1}{q} < \beta$, and both p and $p+1$ are coprime with q . Let $E_{\frac{p}{q}}, E_{\frac{p+1}{q}} \in \mathcal{C}$. Since both of them are the push-forward of line bundle from the same isogeny mop, there exist an injective map $f : E_{\frac{p}{q}} \rightarrow E_{\frac{p+1}{q}}$. The cokernel sheaf F is in $\mathcal{P}((0, 2])$. By Lemma 2.11 part (b), the coherent sheaf F is in $\mathcal{P}((0, 1])$. We get the contradiction by computing

$$\text{Im}(Z(F)) = \text{Im}(Z(E_{\frac{p+1}{q}})) - \text{Im}(Z(E_{\frac{p}{q}})) = \frac{1}{2q} \left(\frac{2p+1}{q} - \beta \right) H^3 < 0.$$

Suppose $\beta' > \beta$, then there exists $\frac{p}{q}$ such that $\beta < \frac{p}{q} < \frac{p+1}{q} < \beta'$, and both p and $p+1$ are coprime with q . The cokernel sheaf F must be in $\mathcal{P}((-2, 0])$. As F is a torsion sheaf, it is in $\mathcal{P}((-1, 0])$ by Lemma 2.11 part (b). We get the contradiction by computing

$$\text{Im}(Z(F)) = \text{Im}(Z(E_{\frac{p+1}{q}})) - \text{Im}(Z(E_{\frac{p}{q}})) = \frac{1}{2q} \left(\frac{2p+1}{q} - \beta \right) H^3 > 0.$$

Therefore, $\beta' = \beta$. In particular, for any $s < \beta < t$, we have $\phi_\sigma(E_s) \leq -1 < 0 < \phi_\sigma(E_t)$. We may deform the stability condition in a small neighbourhood (say, radius less than $\frac{1}{4}$) to another stability condition σ' such that $\mathcal{O}_p \in \mathcal{P}(1)$ and $\text{Im} Z' = H \text{ch}_2^{\beta_0} + d' H^3 \text{rk}$ for some $\beta_0 > \beta$ and $d' > 0$. By the same argument, we have $\phi_{\sigma'}(E_t) \leq -1 < 0 < \phi(E_t)$ for any rational number $\beta < t < \beta_0$. Hence $\text{dist}(\sigma, \sigma') \geq 1$ which leads to the contradiction.

Applying an element of $\widetilde{\text{GL}}_2^+(\mathbb{R})$ we may further assume that the central charge is of the form

$$Z = -\text{ch}_3^\beta + bH \text{ch}_2^\beta + aH^2 \text{ch}_1^\beta + i(H \text{ch}_2^\beta - \frac{\alpha^2}{2} H^3 \text{rk})$$

for some $a, b, \beta \in \mathbb{R}$ and $\alpha > 0$.

We now show that $a > \frac{\alpha^2}{6} + \frac{1}{2}|b|\alpha$. Suppose $a \leq \frac{\alpha^2}{6} + \frac{1}{2}|b|\alpha$, by deforming the stability condition, we may assume $\alpha, \beta \in \mathbb{Q}$, and $a < \frac{\alpha^2}{6} + \frac{1}{2}|b|\alpha$. Note that $\text{Im}(Z(E_t)) \geq 0$ when and only when $|t - \beta| \geq \alpha$. By Corollary 2.17, Lemma 4.2 and 2.11, we have $E_t \in \mathcal{P}((0, 1])$ when $t > \beta + \alpha$; $E_t \in \mathcal{P}((-1, 0])$ when $\beta - \alpha < t \leq \beta + \alpha$; and $E_t \in \mathcal{P}((-2, -1])$ when $t \leq \beta - \alpha$. Note that $\text{Im}(Z(E_{\beta \pm \alpha})) = 0$ and

$$\begin{aligned} \text{Re}(Z(E_{\beta+\alpha}[1])) &= -\text{rk}(E_{\beta+\alpha})H^3\alpha\left(a + \frac{1}{2}b\alpha - \frac{\alpha^2}{6}\right); \\ \text{Re}(Z(E_{\beta-\alpha}[2])) &= -\text{rk}(E_{\beta-\alpha})H^3\alpha\left(a - \frac{1}{2}b\alpha - \frac{\alpha^2}{6}\right). \end{aligned}$$

At least one of them is positive, which leads to the contradiction.

Finally, applying an element of $\widetilde{\mathrm{GL}}_2^+(\mathbb{R})$, we may assume that $\mathcal{O}_p \in \mathcal{P}(1)$, and σ has the same central charge as that of $\sigma_{\alpha,\beta}^{a,b}$ for some $\alpha > 0$ and $a > \frac{\alpha^2}{6} + \frac{1}{2}|b|\alpha$. Moreover, by Corollary 2.17, Lemma 4.2 and 2.11, for every object $E \in \mathcal{C}$, we have $\phi(E) = \phi_{\sigma_{\alpha,\beta}^{a,b}}(E)$.

By Proposition 4.3, for every $\sigma_{\alpha,\beta}^{a,b}$ -stable object F , we have

$$\phi_{\sigma}^+(F) \geq \phi_{\sigma_{\alpha,\beta}^{a,b}}(F) - 1 \text{ and } \phi_{\sigma}^-(F) \leq \phi_{\sigma_{\alpha,\beta}^{a,b}}(F) + 1.$$

By Lemma 4.6, we have $d(\sigma, \sigma_{\alpha,\beta}^{a,b}) \leq 1$. By Lemma 4.7, we have $\sigma = \sigma_{\alpha,\beta}^{a,b}$. \square

Corollary 4.9. *Let (A, H) be a polarized abelian threefold, then $\mathrm{Stab}_H(A) = \tilde{\mathfrak{B}}$.*

4.4. Further questions.

Question 4.10. *Let (X, H) be a smooth projective variety whose Albanese morphism is finite, then is $\mathrm{Stab}_H(X)$ also contractible?*

This is already non-trivial for the threefold case. Unlike Theorem 4.8 of the abelian threefold case, there are examples of different geometric stability conditions on the projective space with the same central charge. However, we don't expect such examples exist when the Albanese morphism of the variety is finite. In particular, we expect [BMS16, Conjecture 1.7] holds in an even stronger sense that $\tilde{\mathfrak{B}}_n$ is the whole space $\mathrm{Stab}_H(X)$ when the Albanese morphism of X is finite.

Question 4.11. *Let X be a smooth projective variety whose Albanese morphism is not finite. Then does there always exist non-geometric stability conditions?*

In other words, we expect the finite Albanese morphism condition is necessary and sufficient for all stability conditions being geometric. When X is of dimension one, the answer is affirmative by [Mac07, Theorem 2.7]. When X is of dimension greater than or equal to three, the question is far beyond reach, as even the existence of stability conditions are only known in few cases, see [BMS16, Li19, Liu20] for more details.

This makes surfaces the most interesting case, which is also already highly non-trivial. By gluing stability conditions with respect to the Orlov semiorthogonal decomposition for blow-ups, there are always non-geometric stability conditions on non-minimal surfaces. One may therefore always assume the surface is minimal. Among all such surfaces, the most interesting case is when the Albanese morphism of X is trivial, in other words, $H^1(X, \mathcal{O}_X) = 0$. We have the following conjecture for the Le Potier function of them.

Conjecture. Let (S, H) be a smooth polarized surface with zero irregularity, then the Le Potier function $\Phi_{S,H}$ is not continuous at 0.

Admitting this conjecture, we expect that there always exist stability conditions as that discussed in [Bri08, Theorem 12.1] when S has zero irregularity. We leave details to a project in the future.

REFERENCES

- [AB13] Daniele Arcara and Aaron Bertram. Bridgeland-stable moduli spaces for K -trivial surfaces. *J. Eur. Math. Soc. (JEMS)*, 15(1):1–38, 2013, arXiv:0708.2247. With an appendix by Max Lieblich.
- [Bay19] Arend Bayer. A short proof of the deformation property of Bridgeland stability conditions. *Math. Ann.*, 375(3-4):1597–1613, 2019.

- [BB17] Arend Bayer and Tom Bridgeland. Derived automorphism groups of K3 surfaces of Picard rank 1. *Duke Math. J.*, 166(1):75–124, 2017.
- [BBMT14] Arend Bayer, Aaron Bertram, Emanuele Macrì, and Yukinobu Toda. Bridgeland stability conditions on threefolds II: An application to Fujita’s conjecture. *J. Algebraic Geom.*, 23(4):693–710, 2014, arXiv:1106.3430.
- [BM02] Tom Bridgeland and Antony Maciocia. Fourier-Mukai transforms for K3 and elliptic fibrations. *J. Algebraic Geom.*, 11(4):629–657, 2002, arXiv:math/9908022.
- [BMS16] Arend Bayer, Emanuele Macrì, and Paolo Stellari. The space of stability conditions on abelian threefolds, and on some Calabi-Yau threefolds. *Inventiones Mathematicae*, 206:1–65, 2016.
- [BMT14] Arend Bayer, Emanuele Macrì, and Yukinobu Toda. Bridgeland stability conditions on threefolds I: Bogomolov-Gieseker type inequalities. *J. Algebraic Geom.*, 23(1):117–163, 2014, arXiv:1103.5010.
- [Bri99] Tom Bridgeland. Equivalences of triangulated categories and Fourier-Mukai transforms. *Bull. London Math. Soc.*, 31(1):25–34, 1999, arXiv:math/9809114.
- [Bri07] Tom Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007, arXiv:math/0212237.
- [Bri08] Tom Bridgeland. Stability conditions on K3 surfaces. *Duke Math. J.*, 141(2):241–291, 2008, arXiv:math/0307164.
- [CP10] John Collins and Alexander Polishchuk. Gluing stability conditions. *Adv. Theor. Math. Phys.*, 14(2):563–607, 2010, arXiv:0902.0323.
- [DLP85] J.-M. Drezet and J. Le Potier. Fibrés stables et fibrés exceptionnels sur \mathbf{P}_2 . *Ann. Sci. École Norm. Sup. (4)*, 18(2):193–243, 1985.
- [HL10] Daniel Huybrechts and Manfred Lehn. *The geometry of moduli spaces of sheaves*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010.
- [HMS08] Daniel Huybrechts, Emanuele Macrì, and Paolo Stellari. Stability conditions for generic K3 categories. *Compos. Math.*, 144(1):134–162, 2008, arXiv:math/0608430.
- [Kaw81] Yujiro Kawamata. Characterization of abelian varieties. *Compositio Math.*, 43(2):253–276, 1981.
- [KO18] Kotaro Kawatani and Shinnosuke Okawa. Nonexistence of semiorthogonal decompositions and sections of the canonical bundle, 2018, arXiv:1508.00682.
- [KS08] Maxim Kontsevich and Yan Soibelman. Stability structures, motivic Donaldson-Thomas invariants and cluster transformations, 2008, arXiv:0811.2435.
- [Li17] Chunyi Li. The space of stability conditions on the projective plane. *Selecta Math. (N.S.)*, 23(4):2927–2945, 2017.
- [Li19] Chunyi Li. On stability conditions for the quintic threefold. *Invent. Math.*, 218(1):301–340, 2019.
- [Liu20] Yucheng Liu. Stability conditions on product varieties, 2020, arXiv:1907.09326.
- [Mac07] Emanuele Macrì. Stability conditions on curves. *Math. Res. Lett.*, 14(4):657–672, 2007, arXiv:0705.3794.
- [MP13] Antony Maciocia and Dulip Piyaratne. Fourier-Mukai transforms and Bridgeland stability conditions on abelian threefolds II. *Internat. J. Math.*, 27(1):1650007, 27, 2013, arXiv:1310.0299.
- [MP15] Antony Maciocia and Dulip Piyaratne. Fourier-Mukai transforms and Bridgeland stability conditions on abelian threefolds. *Algebr. Geom.*, 2(3):270–297, 2015.
- [MS17] Emanuele Macrì and Benjamin Schmidt. Lectures on Bridgeland stability. In *Moduli of curves*, volume 21 of *Lect. Notes Unione Mat. Ital.*, pages 139–211. Springer, Cham, 2017.
- [Muk78] Shigeru Mukai. Semi-homogeneous vector bundles on an Abelian variety. *J. Math. Kyoto Univ.*, 18(2):239–272, 1978.
- [Oka06] So Okada. Stability manifold of \mathbb{P}^1 . *J. Algebraic Geom.*, 15(3):487–505, 2006, arXiv:math/0411220.
- [Orl02] D. O. Orlov. Derived categories of coherent sheaves on abelian varieties and equivalences between them. *Izv. Ross. Akad. Nauk Ser. Mat.*, 66(3):131–158, 2002.
- [Pir21] Dmitrii Pirozhkov. Stably semiorthogonally indecomposable varieties, 2021, arXiv:2011.12743.
- [Pol03] Alexander Polishchuk. *Abelian varieties, theta functions and the Fourier transform*, volume 153 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2003.

- [Pol07] A. Polishchuk. Constant families of t -structures on derived categories of coherent sheaves. *Mosc. Math. J.*, 7(1):109–134, 167, 2007, arXiv:math/0606013.
- [PT19] Dulip Piyaratne and Yukinobu Toda. Moduli of Bridgeland semistable objects on 3-folds and Donaldson-Thomas invariants. *J. Reine Angew. Math.*, 747:175–219, 2019.
- [Rou11] Xavier Roulleau. Fano surfaces with 12 or 30 elliptic curves. *Michigan Math. J.*, 60(2):313–329, 2011.

L. F.: IMAPP, RADBOD UNIVERSITY, NIJMEGEN, NETHERLANDS AND INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD LYON 1, FRANCE

Email address: fu@math.univ-lyon1.fr

URL: <http://math.univ-lyon1.fr/~fu/index.en.html/>

C. L.: MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL, UNITED KINGDOM

Email address: C.Li.25@warwick.ac.uk

URL: <https://sites.google.com/site/chunyili0401/>

X. Z.: DEPARTMENT OF MATHEMATICS, SOUTH HALL 6607, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, CA 93106, USA

Email address: xlzhao@math.ucsb.edu

URL: <https://sites.google.com/site/xiaoleizhaoswebsite/>