

KINK-LIKE SOLUTION FOR THE LORENTZ-VIOLATING ϕ^4 -THEORY EQUATION OF MOTION WITH DISSIPATION

M.A. Knyazev

*Belarusian National Technical University,
65, Independence Ave., Minsk, 220013, Belarus
e-mail: maknyazev@bntu.by*

June 11, 2021

Abstract

A (1+1)-dimension equation of motion for ϕ^4 -theory is considered for the case of simultaneous taking into account the processes of dissipation and violation the Lorentz-invariance. A topological non-trivial solution of one-kink type for this equation is constructed in an analytical form. The modified Hirota method for a solving the nonlinear partial derivatives equations is applied. A modification of the method led to the special conditions on parameters of the model and solution.

Key words: kink, Lorentz-violating model, ϕ^4 -theory, dissipation, modified Hirota method.

PACS number(s): 02.30.Jr, 47.27.Ak, 52.25.-b

The Lorentz invariance of the field theory equations of motion is a common place in modern physics. This takes a place for the ϕ^4 -theory also. This theory is widely used in the different fields of physics: classical and quantum field theories, physics of the elementary particles, theory of the phase transitions in condensed matter, physics of magnetism, inflation theory, theory of formation the topological defects in cosmology and so on [1].

Last years the physical processes with a violation of the Lorentz-invariance attract an attention of researches. One can mention the Maxwell's electrodynamics which takes into account the Chern-Simons interaction, an accounting the members violating Lorentz invariance in a low-energetic limit of the standard model and high-energetic limit in the string models, non-commutative field theories, supersymmetric theories, a possible limits on the Lorentz invariance from the radioastronomy and so on. In the scalar field theory models, including multi-components models, an equation of motion usually has a kink or kink-like solution. A violation of the Lorentz invariance for such equation takes a place for the Kondo effect, a capture of fermions, in the problems of entropy of various informative processes and others. A review of the modern state of art in this field are presented in paper [2] (see also, [3]).

In this paper (1+1)-dimensional equations of motion in ϕ^4 -theory are considered. Below for convenience where it is possible the coefficients in all equations of motion will be equal to unity.

The static equation of motion in ϕ^4 -theory has a form

$$\phi_{xx} + \phi - \phi^3 = 0, \quad (1)$$

where $\phi_{xx} = \frac{\partial^2 \phi}{\partial x^2}$ and so on. The solution of this equation is well-known [4]

$$\phi(x)_{st} = \pm \tanh\left(\frac{x - x_0}{\sqrt{2}}\right). \quad (2)$$

Here sign plus corresponds to kink and sign minus - to antikink. x_0 is an initial phase and determines a coordinate of the centrum of kink.

For dynamical case Eq.(1) takes a form

$$\phi_{tt} - \phi_{xx} - \phi + \phi^3 = 0. \quad (3)$$

Eq.(3) is not an integrable one. There is only one-kink (one-antikink) solution for this equation. There are no solutions correspond to the coupled states of any kinks, antikinks or kinks and antikinks [5], [6]. Eq.(3) possesses the Lorentz invariance. Hence, its solution may be written in a form

$$\phi(x, t) = \pm \tanh\left(\frac{x - x_0 - ut}{\sqrt{2(1 - u^2)}}\right), \quad (4)$$

where $-1 < u < 1$ is a kink velocity.

If one takes into account a dissipation the equation of motion may be written in a form

$$\phi_{tt} - \phi_{xx} + \alpha\phi_t - \phi + \phi^3 = 0, \quad (5)$$

where α is a damping coefficient. This equation is the Lorentz invariant one. It is possible to solve Eq.(5) by using the elliptic functions [7] or the Hirota direct method for nonlinear partial differential equations [1], [6]. The solution has a form

$$\phi(x, t) = \frac{1}{2}\{1 - \tanh[\alpha(x - ut) + x_0]\}. \quad (6)$$

This solution is not a kink, but kink-like object. For Eq.(5) it is also impossible to construct the solutions which correspond to coupled states of any kinks, antikinks or kinks and antikinks. Hence, Eq. (5) is also not integrable one.

Consider the equation of motion for ϕ^4 -theory in which the Lorentz invariance is violated. This equation has a form

$$\phi_{tt} - \phi_{xx} + 2\beta\phi_{xt} - \phi + \phi^3 = 0. \quad (7)$$

A constant coefficient $\beta > 0$ describes a violation of the Lorentz invariance. A case $\beta = 0$ is trivial one. A coefficient 2 is introduced for a convenience of calculation. It is clear that term $2\beta\phi_{xt}$ does not play any role in a static case. In paper [3] it is shown that Eq.(3) is invariant under a transformation

$$x' = \gamma(x_0 - ut), \gamma = \frac{1}{\sqrt{1 - u^2 + 2\beta u}}. \quad (8)$$

For such a case the topological nontrivial solution for Eq.(8) has a form

$$\phi(x, t) = \tanh[\gamma(x - ut - x_0)]. \quad (9)$$

A positive value of velocity correspond a motion of kink to the right, a negative - to the left. The problem of integrability for Eq.(9) is still open now.

Eqs.(3), (5) and (7) possesses a some similarity. At the same time the differences between them consist in different linear terms. Hence, these terms do not influence essentially on nonlinear properties of equation and a form of their solutions. The solutions of these equations are determined by a hyperbolic tangent and represent a kink or kink-like object.

A further development of studies of the equation of movement for ϕ^4 -theory consists in its generalization. Eqs. (1),(3), (5) and (7) have the same potential. A modification of the potential by taking into account a higher degrees of ϕ is one of the ways of generalization. There are many papers on this topic (see, for example, [8], [9]). One another way of generalization is a deformation of known potential [10], [11].

A generalization of the equation of motion in which a dissipation and violation of the Lorentz invariance take into account simultaneously have a significant interest also. Such equation may be written in a form [12]

$$\phi_{tt} - \phi_{xx} + \alpha\phi_t + 2\beta\phi_{xt} - \phi + \phi^3 = 0. \quad (10)$$

In this paper a one-kink-like solution for Eq.(10) is constructed. The direct Hirota method is applied for this purpose but some modification of the method was realized.

Introduce a new dependent variable according to the Hirota method

$$\phi_x = \sigma \frac{F_x}{F}, \quad (11)$$

where $F = F(x, t)$ is a new unknown function, σ is a constant will be determined below. Now, Eq.(10) may be written in a form

$$\begin{aligned} \frac{F_{xt}}{F} - 2\frac{F_{xt}F_t}{F^2} - \frac{F_x F_{tt}}{F^2} + 2\frac{F_x F_t^2}{F^3} - \frac{F_{xxx}}{F} + 3\frac{F_x F_{xx}}{F^2} - 2\frac{F_x^3}{F^3} + 2\beta\frac{F_{xxt}}{F} - 2\beta\frac{F_{xx}F_t}{F^2} - \\ 4\beta\frac{F_x F_{xt}}{F^2} + 4\beta\frac{F_x^2 F_t}{F^3} + \alpha\frac{F_{xt}}{F} - \alpha\frac{F_x F_t}{F^2} - \sigma\frac{F_x}{F} + \sigma^2\frac{F_x^3}{F^3} = 0. \end{aligned} \quad (12)$$

One looks for a partial solution, hence the next condition may be accepted

$$(\sigma^2 - 2)\frac{F_x^3}{F^3} = 0. \quad (13)$$

From this equation one obtains $\sigma = \sqrt{2}$. Usually, the Hirota method may be used successfully if after condition Eq.(13) the equation under study contains the second order term only. If there are the third order terms, the equation may be solved by the Hirota method for very limited and special cases [13], [14]. In our problem after an accounting Eq.(13), two third order terms are

stayed in Eq.(12). It is clear, that an additional condition is needed. Let write the such condition in a form

$$F_t + 2\beta F_x = 0. \quad (14)$$

After an applying these two conditions to Eq.(12) it contains the second order (bilinear) terms only:

$$\begin{aligned} F_{xtt}F - 2F_{xt}F_t - F_xF_{tt} - F_{xxx}F + 3F_xF_{xx} + 2\beta F_{xxt}F - 2\beta F_{xx}F_t - \\ 4\alpha F_xF_{xt} + \alpha F_{xt}F - \alpha F_xF_t - F_xF = 0. \end{aligned} \quad (15)$$

From now all the steps in constructing of the one-kink-like solution are standard ones for the Hirota method. As a result one obtains:

$$\phi(x, t) = \frac{1}{k} \left[1 + \tanh\left(\frac{kx - \omega t + \eta_0}{\sqrt{2}}\right) \right], \quad (16)$$

where $k = -\frac{3}{4\alpha\beta}$, $\omega = -\frac{3}{2\alpha}$. η_0 describes an initial position of solution. A dispersion relation has a form

$$\omega^2 - k^2 - 2\beta k\omega - \alpha\omega - 1 = 0. \quad (17)$$

As to condition (14) it takes a form

$$\omega = 2\beta k. \quad (18)$$

The last condition leads to relation $\alpha^2\beta^2 > 1$.

Substitute Eq.(16) into Eq.(10). Then for $\omega = -\frac{3}{2\alpha}$ Eq.(16) will be a solution of Eq.(10) for $k^2 = \frac{1}{2}$ only. This condition is in agreement with relation $\alpha^2\beta^2 > 1$. Namely, $\alpha^2\beta^2 = \frac{9}{8}$.

In this work a one-kink-like solution is constructed for the Lorentz-violating ϕ^4 -theory equation of motion with dissipation. It was done by some modification of the Hirota direct method for nonlinear equation. This modification led to the special conditions on coefficients of equation and parameters of solution.

References

- [1] M. A. Knyazev, *Kinks in Scalar Model with Damping*, Technologiya, Minsk, 2013 [in Russian]
- [2] Haobo Yan, Yuan Zhong, Yu-Xiao Lui, Kei-ichi Maeda, *Kink-antikink collision in a Lorentz-violating ϕ^4 model*, Phys. Lett. B **807**(2020),135542 (7 pp.), doi:10.1016/j.physletb.2020.135542
- [3] M.N. Barreto, D. Bazeia, R. Menezes, *Defect structures in Lorentz and CPT violating scenarios*, arXiv:hep-th/0506262, doi:10.1103/PhysRevD.73.065015
- [4] R. Rajaraman, *Solitons and Instantons. An Introduction to Solitons and Instantons in Quantum Field Theory*, North-Holland Publ. Company, Amsterdam et al., 1982
- [5] A.C. Newell, *Solitons in Mathematics and Physics*, SIAM, 1985, doi:10.1137/1.981611970227
- [6] M.J. Ablowitz, H. Segur, *Solitons and Inverse Scattering Transform*, SIAM, 1982
- [7] J.M. Cervero, P.G. Estevez, *General elliptic solution for the cubic equation with damping*, Phys. Lett. A **114** (1986), 435-436.
- [8] V.A. Gani, Aliakbar Moradi Marjaneh, P.A. Blinov, *Explicit kinks in higher-order field theories*, arXiv:hep-th/2002.09981, doi:10.1103/PhysRevD.101.12507
- [9] I.C. Christov, R.J. Decker, A. Demirkaya, V.A. Gani, P.G. Kevrekidis, *Kink-antikink collisions and multi-bounce resonance windows in higher-order field theories*, arXiv:hep-th/2005.00154
- [10] D. Bazeia, E. Belendryasova, V. Gani, *Scattering of kinks of the sinh-deformed ϕ^4 model*, arXiv:hep-th/1710.04993, doi:10.1140/epic/s10052-018-5815-z
- [11] D. Bazeia, E. Belendryasova, V. Gani, *Scattering of kinks in a non-polynomial model*, arXiv:hep-th/1710.07788,doi:10.1088/1742-6596/1/012032

- [12] M.A. Knyazev, *A problem on dissipative equation of movement for ϕ^4 theory with a violating of the Lorentz invariance*, Proc. of the 13 Int. Sci. and Tech. Conf. "Instrumentation Engineering-2020" (BNTU, Minsk, 2020), 256-257
- [13] J. Hietarinta, B. Grammaticos, A. Ramani, *Integrable trilinear PDE's*, arXiv:solv-int/9411003
- [14] P.P. Goldstein, *Hints on Hirota Bilinear Method*, Acta Physica Polonica A **112** (2007),1171 (14 pp.), doi:10.12693/APhysPolA.1121171