

AN EXPLICIT ESTIMATE OF THE BERGMAN KERNEL FOR POSITIVE LINE BUNDLES

XU WANG

Dedicated to Bo Berndtsson on occasion of his 70th birthday

ABSTRACT. We shall give an explicit estimate of the lower bound of the Bergman kernel associated to a positive line bundle. In the compact Riemann surface case, our result can be seen as an explicit version of Tian's partial C^0 -estimate.

CONTENTS

1. Introduction	1
2. Hessian comparison theorem	3
3. An Ohsawa–Takegoshi type theorem	4
3.1. Proof of Theorem A, B, An, Bn	5
4. Another proof of a weaker version of Theorem A, B	5
4.1. The Blocki–Zwonek estimate	6
4.2. Isoperimetric inequality	7
References	8

1. INTRODUCTION

Let $(L, e^{-\phi})$ be a positive line bundle over an n -dimensional complex manifold X . Let m be a positive integer. Let K_X be the canonical line bundle over X . We call

$$(1.1) \quad K_{m\phi}(x) := \sup_{u \in H^0(X, K_X + mL)} \frac{u(x) \wedge \overline{u(x)} e^{-m\phi(x)}}{\int_X u \wedge \bar{u} e^{-m\phi}},$$

the *Bergman kernel forms* and

$$(1.2) \quad B_{m\phi}(x) := \sup_{u \in H^0(X, mL)} \frac{|u(x)|^2 e^{-m\phi(x)}}{\int_X |u|^2 e^{-m\phi}}, \quad MA_{m\phi} := \frac{(i\partial\bar{\partial}(m\phi))^n}{n!},$$

the *Bergman kernel functions*. In [19] Tian proved that if X is compact then

$$(1.3) \quad \lim_{m \rightarrow \infty} \frac{K_{m\phi}}{MA_{m\phi}} = \lim_{m \rightarrow \infty} B_{m\phi} = \frac{1}{(2\pi)^n}.$$

Date: January 6, 2022.

Effective lower bound estimate (with Ricci curvature, diameter and volume assumptions) for $B_{m\phi}$ is known as Tian's *partial C^0 -estimate* [20]. The first general result is obtained by Donaldson–Sun [9] using proof by contradiction. Our main results are the followings:

Theorem A. *Let $(L, e^{-\phi})$ be a positive line bundle over a compact Riemann surface X . Put $\omega := \text{MA}_\phi = i\partial\bar{\partial}\phi$. Denote by $\text{Ric}\omega := i\partial\bar{\partial}\log\omega$ the Ricci form of ω . Assume that*

$$\text{Ric}\omega \leq \omega, \quad L_0 \geq 2\pi,$$

where L_0 denotes the infimum of the length of closed geodesics in X , then $K_\phi/\text{MA}_\phi \geq \frac{1}{8\pi}$.

Theorem B. *Let $(L, e^{-\phi})$ be a positive line bundle over a compact Riemann surface X . If*

$$-\omega/2 \leq \text{Ric}\omega \leq \omega/2, \quad L_0 \geq 2\pi\sqrt{2},$$

then $B_\phi \geq \frac{1}{16\pi}$.

Remark. *In case $X = \mathbb{P}^1$ and $\omega = 2 \cdot i\partial\bar{\partial}\log(1 + |z|^2)$ we have*

$$\text{Ric}\omega = \omega, \quad L_0 = 2\pi,$$

a direct computation gives $L = -K_X$ and $K_\phi/\text{MA}_\phi = \frac{1}{4\pi}$. We do not know whether

$$K_\phi/\text{MA}_\phi \geq \frac{1}{4\pi}$$

is always true with the assumptions in Theorem A. On the other hand, Theorem A implies $K_{m\phi}/\text{MA}_{m\phi} \geq 1/(8\pi)$ for every positive integer m . This is also near optimal since by (1.3)

$$\lim_{m \rightarrow \infty} K_{m\phi}/\text{MA}_{m\phi} = 1/(2\pi).$$

In case $\text{Ric}\omega \leq 0$, $L_0/2$ is equal to the injectivity radius. For example if $X = \mathbb{C}/\Gamma$ is a torus and $\omega = i\partial\bar{\partial}(|z|^2/2)$ then $\text{Ric}\omega = 0$ and $L_0 = \inf_{0 \neq \gamma \in \Gamma} |\gamma|$.

In the first version of this paper, a weaker version of the above theorems is proved using an Ohsawa–Takegoshi type theorem, a variant of the Blocki–Zwonek estimate [5] and the isoperimetric inequality. Later we find that one may use the Hessian comparison theorem to simplify the proof and generalize the above theorems to the followings higher dimensional cases.

Theorem An. *Let $(L, e^{-\phi})$ be a positive line bundle over an n -dimensional compact complex manifold X . Assume that the sectional curvature of $\omega := i\partial\bar{\partial}\phi$ is bounded above by $1/(4n)$ and $L_0 \geq 2\pi\sqrt{n}$ then $K_\phi/\text{MA}_\phi \geq \frac{1}{2} \frac{1}{(4\pi n)^n}$.*

Theorem Bn. *Let $(L, e^{-\phi})$ be a positive line bundle over an n -dimensional compact manifold X . Assume that the sectional curvature of $\omega := i\partial\bar{\partial}\phi$ is bounded above by $1/(8n)$, $L_0 \geq 2\pi\sqrt{2n}$ and $\text{Ric}\omega \geq -\omega/2$. Then $B_\phi \geq \frac{1}{2} \frac{1}{(8\pi n)^n}$.*

Remark. *Since the Ricci curvature is certain sums of sectional curvatures, the curvature assumptions in Theorem Bn also imply a lower bound of the sectional curvature. Hence one may use [12, Corollary 2.3.2] to find a lower bound of L_0 in terms of the lower bound of the volume and the upper bound of the diameter. Thus, except the upper bound of the sectional curvature, the assumptions in Theorem Bn follow from the standard assumptions in Tian's partial C^0 -estimate (for results on Tian's partial C^0 -estimate, see [1, 6, 7, 13, 14, 15, 18, 21, 22, 24], etc). Our*

main contribution is the explicit constant in the estimate. Moreover, our estimate implies that $B_{m\phi} \geq \frac{1}{2} \frac{1}{(8\pi n)^n}$ for all positive integers m . If we understand [9] correctly, for general positive line bundles, this estimate can not be true with just Ricci curvature, diameter and volume assumptions (see the explanation at the end of [9]).

Acknowledgment: It is a pleasure to thank Bo Berndtsson for stimulating discussions related to the topics of this article.

2. HESSIAN COMPARISON THEOREM

Definition 2.1. Let X be a Riemann manifold. Denote by $K(V, W)$ the sectional curvature of the tangent plane spanned by V, W . Fix $x \in X$, the injectivity radius at $x \in X$ is defined as

$$\text{inj}(x) := \sup\{r > 0 : \exp_x|_{B(0,r)} \text{ is diffeomorphism}\}$$

where $\exp_x : T_x X \rightarrow X$ denotes the exponential map at x and $B(0, r)$ denotes the ball of radius r around $0 \in T_x X$. We call $\text{inj}_X := \inf_{x \in X} \text{inj}(x)$ the injectivity radius of X .

The following $\partial\bar{\partial}$ -comparison theorem is a direct consequence of the Hessian comparison theorem (see [10, Lemma 1.13 in page 14 and Theorem A in page 19]).

Theorem 2.1. Let X_1, X_2 be Kähler manifolds. Let $\gamma_1 : [0, b] \rightarrow X_1$ and $\gamma_2 : [0, b] \rightarrow X_2$ be unit speed geodesics. With the definition above, suppose that

$$(2.1) \quad b \leq \min\{\text{inj}(\gamma_1(0)), \text{inj}(\gamma_2(0))\}$$

and for all $t \in [0, b]$, $v_1 \perp \dot{\gamma}_1(t)$ and $v_2 \perp \dot{\gamma}_2(t)$,

$$(2.2) \quad K(\dot{\gamma}_1(t), v_1) \leq K(\dot{\gamma}_2(t), v_2).$$

Let $d_j := d(\cdot, \gamma_j(0))$ be distance functions. If $f : (0, b) \rightarrow \mathbb{R}$ is smooth and increasing then

$$(2.3) \quad i\partial\bar{\partial}(f \circ d_1)(V_1, V_1) \geq i\partial\bar{\partial}(f \circ d_2)(V_2, V_2)$$

for all $t \in (0, b)$, $V_j \in T_{\gamma_j(t)} X_j$, $j = 1, 2$, such that $|V_1| = |V_2|$ and

$$(\dot{\gamma}_1(t), V_1) = (\dot{\gamma}_2(t), V_2), \quad (\dot{\gamma}_1(t), JV_1) = (\dot{\gamma}_2(t), JV_2).$$

We shall apply the above theorem to $X_2 = \mathbb{P}^n$ with the Fubini study metric form $\omega_2 = 2i\partial\bar{\partial} \log(1 + |z|^2)$. A direct computation gives

$$(2.4) \quad K(V, W) = \frac{1}{4} (1 + 3(V, JW)^2), \quad \forall V \perp W, \quad \text{inj}(x) = \pi.$$

In particular, $K(V, W) = 1$ in case $n = 1$ and $K(V, W) \geq 1/4$ in case $n \geq 2$. We also need the following distance function formula on $\mathbb{C}^n \subset \mathbb{P}^n$

$$d_2 := d(0, z) = 2 \int_0^{|z|} \frac{dx}{1+x^2} = 2 \arctan |z|.$$

Put

$$\psi = \log \sin^2(d_2/2)$$

we have

$$\psi = \log \frac{\tan^2(d_2/2)}{1 + \tan^2(d_2/2)} = \log |z|^2 - \log(1 + |z|^2) \leq 0$$

and

$$i\partial\bar{\partial}\psi \geq -i\partial\bar{\partial}\log(1 + |z|^2) = -\omega_2/2.$$

Apply the above theorem to $f(x) = \log \sin^2(x/2)$ and $b = \pi$ we get:

Corollary 2.2. *Let (X_1, ω_1) be an n -dimension Kähler manifold. Fix $x \in X_1$. Assume that the injectivity radius at x is no less than π . Put*

$$\psi(z) := \begin{cases} \log \sin^2(d(z, x)/2) & d(z, x) \leq \pi \\ 0 & d(z, x) > \pi. \end{cases}$$

If

$$\begin{cases} \text{the sectional curvature of } \omega_1 \text{ is no bigger than } 1, & n = 1 \\ \text{the sectional curvature of } \omega_1 \text{ is no bigger than } 1/4, & n \geq 2 \end{cases}$$

then $i\partial\bar{\partial}\psi \geq -\omega_1/2$ on X .

Proof. Apply the theorem above, we get that $i\partial\bar{\partial}\psi \geq -\omega_1/2$ when $d(z, x) < \pi$. Hence the corollary follows since the gradient of ψ vanishes at $d(z, x) = \pi$. \square

3. AN OHSAWA–TAKEGOSHI TYPE THEOREM

We shall use the following Ohsawa–Takegoshi type theorem [17], which is a special case of the main theorem in [11].

Theorem 3.1. *Let $(L, e^{-\phi})$ be a positive line bundle on an n -dimensional compact complex manifold. Fix $x \in X$. Assume that there is a non-positive function G smooth outside x such that $G(z) - \log |z - x|^{2n}$ is smooth near x and*

$$i\partial\bar{\partial}\phi + \lambda i\partial\bar{\partial}G \geq 0$$

on X for some constant $\lambda > 1$. Then K_ϕ in (1.1) satisfies

$$(3.1) \quad K_\phi(x) \geq \frac{\lambda - 1}{\lambda} \lim_{t \rightarrow -\infty} \frac{\text{MA}_\phi(x)}{e^{-t} \int_{G < t} \text{MA}_\phi},$$

where MA_ϕ is defined in (1.2).

Proof. Let us rephrase the proof in [17]. Our curvature assumption implies that

$$\phi^{t+is} := \phi + \lambda \max\{G - t, 0\}$$

defines a singular metric on $\mathbb{C}_{t+is} \times L$ with non-negative curvature current. Hence Berndtsson's theorem [2] implies that $\log K_{\phi^t}(x)$ is a convex function of t . By a direct computation (see the appendix in [17] or Theorem 3.8 in [3]) we find that

$$\lim_{t \rightarrow -\infty} e^t K_{\phi^t}(x) = \frac{\lambda - 1}{\lambda} \lim_{t \rightarrow -\infty} \frac{\text{MA}_\phi(x)}{e^{-t} \int_{G < t} \text{MA}_\phi}$$

is finite since $G(z) - \log|z - x|^{2n}$ is smooth near x . Hence $e^t K_{\phi^t}(x) = e^{t + \log K_{\phi^t}(x)}$, as a convex function of t bounded near $-\infty$, must be increasing. Thus

$$K_{\phi}(x) = e^0 K_{\phi^0}(x) \geq \lim_{t \rightarrow -\infty} e^t K_{\phi^t}(x) = \frac{\lambda - 1}{\lambda} \lim_{t \rightarrow -\infty} \frac{\text{MA}_{\phi}(x)}{e^{-t} \int_{G < t} \text{MA}_{\phi}}$$

gives our estimate. \square

3.1. Proof of Theorem A, B, An, Bn.

Proof of Theorem A, An. By Klingenberg's estimate (see [23, Corollary 1.2]), if the sectional curvature is no bigger than $1/c$ then we have

$$(3.2) \quad \min\{L_0/2, \sqrt{c}\pi\} \leq \text{inj}_X \leq L_0/2.$$

Hence our assumptions implies that the injectivity radius of $(X, \omega/n)$, $\omega := i\partial\bar{\partial}\phi$, is no less than π . Thus one may apply Corollary 2.2 to $(X_1, \omega_1) = (X, \omega/n)$. Put $G = n\psi$. Corollary 2.2 implies that

$$i\partial\bar{\partial}\phi + 2i\partial\bar{\partial}G \geq 0.$$

Hence Theorem 3.1 ($\lambda = 2$) gives Theorem A, An. \square

Proof of Theorem B, Bn. By the Ricci curvature assumption, ϕ and $i\partial\bar{\partial}\phi$ defines a metric on $L - K_X$ with curvature

$$i\Theta = \omega + \text{Ric}\omega \geq \omega/2.$$

Thus one may apply Corollary 2.2 to $(X_1, \omega_1) = (X, \omega/(2n))$. Put $G = n\psi$ then

$$i\Theta + 2i\partial\bar{\partial}G \geq 0.$$

Apply Theorem 3.1 to $L - K_X$, we get Theorem B, Bn. \square

4. ANOTHER PROOF OF A WEAKER VERSION OF THEOREM A, B

In this section we shall give another proof of Theorem A, B with an extra volume assumption.

Theorem C. *Let $(L, e^{-\phi})$ be a positive line bundle over a compact Riemann surface X . If*

$$\int_X \omega \geq 8\pi, \quad \text{Ric}\omega \leq \omega, \quad L_0 \geq 2\pi,$$

then $K_{\phi}/\text{MA}_{\phi} \geq \frac{1}{8\pi}$.

Theorem D. *Let $(L, e^{-\phi})$ be a positive line bundle over a compact Riemann surface X . If*

$$\int_X \omega \geq 16\pi, \quad -\omega/2 \leq \text{Ric}\omega \leq \omega/2, \quad L_0 \geq 2\pi\sqrt{2},$$

then $B_{\phi} \geq \frac{1}{16\pi}$.

4.1. The Blocki–Zwonek estimate. We shall study the right hand side of (3.1) using a variant of Blocki–Zwonek’s estimate [5, Proof of Theorem 3] (see also [4, section 10] for related results).

Lemma 4.1. *With the notation in Theorem 3.1. Let ω be an arbitrary Kähler form on X . Then*

$$\frac{d}{dt} \int_{G < t} \omega_n \geq \frac{\sigma(G = t)^2}{2 \int_{G < t} i\partial\bar{\partial}G \wedge \omega_{n-1}}, \quad \omega_q := \omega^q/q!,$$

where

$$\sigma(G = t) := \int_{G=t} d\sigma, \quad d\sigma := \sqrt{2} \sum \frac{G_{\bar{\alpha}} \omega^{\bar{\alpha}\beta}}{|\bar{\partial}G|_{\omega}} \frac{\partial}{\partial z^{\beta}} \rfloor \omega_n,$$

is the measure of the hypersurface $\{G = t\}$ with respect to ω

Proof. Notice that

$$V := \frac{\partial}{\partial t} + \sum \frac{G_{\bar{\alpha}} \omega^{\bar{\alpha}\beta}}{|\bar{\partial}G|_{\omega}^2} \frac{\partial}{\partial z^{\beta}}$$

satisfies $V(G - t) = 0$, hence it can be used to compute $\frac{d}{dt} \int_{G < t}$, in particular, we have

$$\frac{d}{dt} \int_{G < t} \omega_n = \int_{G < t} L_V \omega_n = \int_{G=t} V \rfloor \omega_n = \frac{1}{\sqrt{2}} \int_{G=t} \frac{d\sigma}{|\bar{\partial}G|_{\omega}}.$$

Hence the Cauchy–Schwarz inequality gives

$$\frac{d}{dt} \int_{G < t} \omega_n \geq \frac{\sigma(G = t)^2}{\sqrt{2} \int_{G=t} |\bar{\partial}G|_{\omega} d\sigma} = \frac{\sigma(G = t)^2}{2 \int_{G=t} i\bar{\partial}G \wedge \omega_{n-1}} = \frac{\sigma(G = t)^2}{2 \int_{G < t} i\partial\bar{\partial}G \wedge \omega_{n-1}},$$

where we use the Stokes theorem in the last equality. \square

Since

$$G_1 \leq G_2 \Rightarrow \int_{G_2 < t} \omega_n \leq \int_{G_1 < t} \omega_n,$$

in order to get the best estimate from (3.1), one should choose G to be the following *envelope*, say $g_{\phi, x, \lambda}$, defined by

$$(4.1) \quad \sup\{G \leq 0 : G(z) - \log|z - x|^{2n} \text{ is smooth near } x \text{ and } i\partial\bar{\partial}\phi + \lambda i\partial\bar{\partial}G \geq 0\}.$$

It is known that (see [8])

$$(4.2) \quad \epsilon_x(L) := \sup\{\lambda \geq 0 : \text{there exists } G \leq 0 \text{ on } X \text{ such that the blue part in (4.1) holds}\}$$

is equal to the *Seshadri constant* up to a constant factor n . If $0 < \lambda < \epsilon_x(L)$ then $g_{\phi, x, \lambda}$, as an envelope, must satisfy

$$(4.3) \quad (i\partial\bar{\partial}\phi + \lambda i\partial\bar{\partial}g_{\phi, x, \lambda})^n = (2\pi n\lambda)^n \delta_x$$

on $\{g_{\phi,x,\lambda} < 0\}$, where δ_x is the Dirac measure defined by $\int_X f \delta_x = f(x)$. Thus if we choose $G = g_{\phi,x,\lambda}$ and $\omega = i\partial\bar{\partial}\phi$ then

$$(4.4) \quad \int_{G<t} i\partial\bar{\partial}G \wedge \omega_{n-1} = \int_{G<t} (i\partial\bar{\partial}G + \omega/\lambda - \omega/\lambda) \wedge \omega_{n-1}$$

$$(4.5) \quad = \int_{G<t} (i\partial\bar{\partial}G + \omega/\lambda) \wedge \omega_{n-1} - \frac{n}{\lambda} \int_{G<t} \omega_n$$

$$(4.6) \quad \leq \int_{G<0} (i\partial\bar{\partial}G + \omega/\lambda) \wedge \omega_{n-1} - \frac{n}{\lambda} \int_{G<t} \omega_n$$

$$(4.7) \quad = \frac{n}{\lambda} \left(\int_{G<0} \omega_n - \int_{G<t} \omega_n \right),$$

where we use $\int_{G<0} i\partial\bar{\partial}G \wedge \omega_{n-1} = 0$ (since $G = 0$ outside $\{G < 0\}$) in (4.7). In case $n = 1$, (4.3) directly gives

$$(4.8) \quad \int_{G<t} i\partial\bar{\partial}G = \int_{G<t} i\partial\bar{\partial}G + \omega/\lambda - \omega/\lambda = 2\pi - \int_{G<t} \omega/\lambda.$$

Hence Lemma 4.1 implies

$$(4.9) \quad \frac{d}{dt} \int_{G<t} \omega \geq \frac{\sigma(G=t)^2}{4\pi - \frac{2}{\lambda} \int_{G<t} \omega}.$$

4.2. Isoperimetric inequality. We shall use the following result (see inequality (5.4) in [16, Proposition 5.2]).

Lemma 4.2 (Isoperimetric inequality). *Let U be an open subset of a compact Riemann surface (X, ω) . Assume that*

$$A := \int_U \omega \leq \frac{1}{2} \int_X \omega, \quad \text{Ric } \omega \leq k \omega.$$

Then

$$(4.10) \quad \sigma(\partial U)^2 \geq \min\{L_0^2, A(4\pi - kA)\},$$

where L_0 denotes the infimum of the length of simple closed geodesics in X .

Proof. By the definition of the Seshadri constant in (4.2), we know that in case $n = 1$, the Hodge decomposition gives

$$(4.11) \quad \epsilon_x = \text{deg}(L) := \int_X c_1(L) = \frac{1}{2\pi} \int_X \omega.$$

Hence if $\int_X \omega \geq 8\pi$ then $\epsilon_x \geq 4$. Hence we can take $\lambda = 2$ in (4.9), which gives

$$(4.12) \quad \frac{d}{dt} \int_{G<t} \omega \geq \frac{\sigma(G=t)^2}{4\pi - \int_{G<t} \omega}.$$

Now, since $\text{Ric } \omega \leq \omega$, $L_0^2 \geq 4\pi^2$ and

$$\int_{G<t} \omega \leq \int_{G<0} \omega = 2\pi\lambda = 4\pi \leq \frac{1}{2} \int_X \omega,$$

one may apply the above lemma to $U = \{G < t\}$, then by (4.10) we have

$$\sigma(G = t)^2 \geq \min\{4\pi^2, A(4\pi - A)\} = A(4\pi - A), \quad A := \int_{G < t} \omega.$$

Thus (4.12) gives

$$\frac{d}{dt} \int_{G < t} \omega \geq \int_{G < t} \omega,$$

which implies that $e^{-t} \int_{G < t} \omega$ is increasing with respect to $t < 0$. Hence

$$\lim_{t \rightarrow -\infty} e^{-t} \int_{G < t} \omega \leq e^{-0} \int_{G < 0} \omega = 2\pi\lambda = 4\pi.$$

Then by (3.1) we have $K_\phi \geq \frac{\omega}{8\pi}$. □

Theorem D follows by a similar argument (see the difference between the proof of Theorem A, B above).

REFERENCES

- [1] R. Bamler, *Convergence of Ricci flows with bounded scalar curvature*, Ann. of Math. **188** (2018), 753–831.
- [2] B. Berndtsson, *Curvature of vector bundles associated to holomorphic fibrations*, Ann. of Math. **169** (2009), 531–560.
- [3] B. Berndtsson and L. Lempert, *A proof of the Ohsawa–Takegoshi theorem with sharp estimates*, J. Math. Soc. Japan **68** (2016), 1461–1472.
- [4] Z. Blocki, *Cauchy-Riemann meet Monge-Ampere*, Bull. Math. Sci. **4** (2014), 433–480.
- [5] Z. Blocki and W. Zwonek, *Estimates for the Bergman Kernel and the Multidimensional Suita Conjecture*, arXiv:1404.7692.
- [6] X. X. Chen and B. Wang, *Space of Ricci flows I*, Comm. Pure Appl. Math. **65** (2012), 1399–1457.
- [7] X. X. Chen and B. Wang, *Space of Ricci flows II*, arXiv:1405.6797.
- [8] J. P. Demailly, *Singular hermitian metrics on positive line bundles*, Lecture Notes in Math. 1992, 87–104.
- [9] S. Donaldson and S. Sun, *Gromov–Hausdorff limits of Kähler manifolds and algebraic geometry*, Acta. Math. **213** (2014), 63–106.
- [10] R. E. Greene and H. Wu, *Function theory on manifolds which possess a pole*, Lecture Notes in Mathematics, vol. 699, Springer, Berlin, 1979.
- [11] Q. Guan and X. Zhou, *A solution of an L^2 extension problem with an optimal estimate and applications*, Ann. of Math. (2), **181** (2015), 1139–1208.
- [12] E. Heintze and H. Karcher, *A general comparison theorem with applications to volume estimates for submanifolds*, Ann. Sci. Ecole Norm. Sup. **11** (1978), 451–470.
- [13] W. Jiang, *Bergman kernel along the Kähler-Ricci flow and Tian’s conjecture*, J. Reine Angew. Math. **717** (2016), 195–226.
- [14] W. Jiang, F. Wang and X. Zhu, *Bergman kernels for a sequence of almost Kähler-Ricci solitons*, Ann. Inst. Fourier. **67** (2017), 1279–1320.
- [15] G. Liu and G. Székelyhidi, *Gromov-Hausdorff limits of Kähler manifolds with Ricci curvature bounded below*, arXiv:1804.08567.
- [16] F. Morgan and D. L. Johnson, *Some sharp isoperimetric theorems for Riemannian manifolds*, Indiana Univ. Math. J. **49** (2000), 1017–1041.
- [17] T. Nguyen and X. Wang, *A remark to the Berndtsson–Lempert proof of the Ohsawa–Takegoshi extension theorem*, to appear.
- [18] G. Székelyhidi, *The partial C^0 -estimate along the continuity method*, J. Amer. Math. Soc. **29** (2016), 537–560.
- [19] G. Tian, *On a set of polarized Kähler metrics on algebraic manifolds*, J. Differential Geom. **32** (1990), 99–130.

- [20] G. Tian, *On Calabi's conjecture for complex surfaces with positive first Chern class*, Invent. Math. **101** (1990), 101–172.
- [21] G. Tian, *Partial C^0 -estimates for Kähler-Einstein metrics*, Commun. Math. Stat. **1** (2013), 105–113.
- [22] F. Wang and X. Zhu, *Tian's partial C^0 -estimate implies Hamilton-Tian's conjecture*, arXiv:2002.05501.
- [23] Z. Wang, *The sphere theorem*, <http://staff.ustc.edu.cn/~wangzuoq/Courses/16S-RiemGeom/Notes/Lec22.pdf>
- [24] K. Zhang, *Some refinements of the partial C^0 -estimate*, arXiv:1911.11328.

DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, NO-7491 TRONDHEIM, NORWAY

Email address: `xu.wang@ntnu.no`