

BIRATIONAL KLEINIAN GROUPS

SHENGYUAN ZHAO

ABSTRACT. In this paper we initiate the study of birational Kleinian groups, i.e. groups of birational transformations of complex projective varieties acting in a free, properly discontinuous and cocompact way on an open subset of the variety with respect to the usual topology. We obtain a classification in dimension two.

CONTENTS

1. Introduction	1
2. Varieties of non-negative Kodaira dimension	6
3. Complex projective Kleinian groups	8
4. Preliminaries on groups of birational transformations of surfaces	10
5. Birational Kleinian groups are elementary	13
6. Foliated surfaces	19
7. Invariant rational fibration I	22
8. Invariant rational fibration II: elliptic base	29
9. Invariant rational fibration III: rational base	31
10. Classification and proof	43
References	47

1. INTRODUCTION

1.1. Birational Kleinian groups.

1.1.1. *Definitions.* According to Klein's 1872 Erlangen program a geometry is a space with a nice group of transformations. One of the implementations of this idea is Ehresmann's notion of geometric structures ([Ehr36]) which models a space locally on a geometry in the sense of Klein. In his Erlangen program, Klein conceived a geometry modeled upon birational transformations:

Of such a geometry of rational transformations as must result on the basis of the transformations of the first kind, only a beginning has so far been made... ([Kle72], [Kle93]8.1)

In this paper we study the group of birational transformations of a variety from Klein and Ehresmann's point of view.

Let Y be a smooth complex projective variety and $U \subset Y$ a connected open set in the usual topology. Let $\Gamma \subset \text{Bir}(Y)$ be an *infinite* group of birational transformations. We impose the following conditions on Γ :

2010 *Mathematics Subject Classification.* 57S30, 14E07, 32Q30, 32M99, 57M50, 37F75.

- (1) the points of indeterminacy of Γ are disjoint from U and Γ preserves U , in other words each element of Γ induces a holomorphic diffeomorphism of U ;
- (2) the action of Γ on U is free, properly discontinuous and cocompact.

Therefore U is identified with an infinite Galois covering of the quotient compact complex manifold $X = U/\Gamma$ and Γ is identified with the group of deck transformations. We call the data (Y, Γ, U, X) a *birational kleinian group*. When the context is clear, we will just call Γ a birational kleinian group.

We say that two birational kleinian groups (Y, Γ, U, X) and (Y', Γ', U', X') are *geometrically conjugate* if there exists a birational map $\phi : Y \dashrightarrow Y'$ such that

- (1) ϕ is regular on U and sends U biholomorphically to U' ;
- (2) $\Gamma' = \phi\Gamma\phi^{-1}$.

The corresponding quotient manifolds X and X' are isomorphic. If (Y, Γ, U, X) is a birational kleinian group and if Γ' is a finite index subgroup of Γ , then (Y, Γ', U, X') is also a birational kleinian group where X' is a finite unramified covering of X .

Two typical families of examples are given by complex tori and quotients of bounded symmetric domains:

- (1) $Y = \mathbb{P}^n$, $U = \mathbb{C}^n$, Γ is a lattice in \mathbb{C}^n and X is a complex torus.
- (2) U is a hermitian symmetric space of noncompact type, Y is the associated hermitian symmetric space of compact type and Γ is a cocompact lattice in the corresponding Lie group.

The problem of classification of birational kleinian groups was transmitted to the author by Serge Cantat. Serge Cantat learned it from Fedor Bogomolov who learned it from Shing-Tung Yau.

1.1.2. Comparison with classical Kleinian groups. Nowadays a *Kleinian group* means an arbitrary discrete subgroup of $\mathrm{PGL}(2, \mathbb{C})$. In this text we will call them *classical Kleinian groups* to avoid confusions. If $\Gamma \subset \mathrm{PGL}(2, \mathbb{C})$ is a classical Kleinian group, then there is a unique maximal Γ -invariant open (not necessarily connected) subset $\Omega_\Gamma \subset \mathbb{P}^1$ on which Γ acts discontinuously; it is called the *domain of discontinuity* of Γ . The complement $\mathbb{P}^1 \setminus \Omega_\Gamma$ is called the *limit set* of Γ .

If Γ is a torsion free finitely generated classical Kleinian group then Ahlfors finiteness theorem asserts that Ω_Γ/Γ is a finite union of Riemann surfaces of finite type, i.e. smooth quasiprojective curves (see [Ahl64], [Sul85]). If Γ is a torsion free finitely generated classical Kleinian group such that Ω_Γ has an invariant connected component Ω_Γ^0 , then Ω_Γ^0/Γ is a connected smooth quasiprojective curve. Our definition of birational Kleinian group is a direct generalization of such classical Kleinian groups.

For more general spaces the notion of domain of discontinuity and that of limit set are not well defined (see [Kul78]). In dimension ≥ 2 a group of automorphisms of a projective variety may act in a free, properly discontinuous and cocompact way on different invariant open subsets. For example let a Fuchsian surface group act diagonally on $\mathbb{P}^1 \times \mathbb{P}^1$, then both $\mathbb{D} \times \mathbb{P}^1$ and $\mathbb{P}^1 \times \mathbb{D}$ can serve as domain of discontinuity (see [CNS13], [KLP18] for more examples). This is why we define a birational Kleinian group as a quadruple (Y, Γ, U, X) : Y, Γ, U determine X but U is not always uniquely determined by Γ .

1.1.3. Complex projective Kleinian groups and Anosov subgroups. If a birational Kleinian group (Y, Γ, U, X) satisfies that $Y = \mathbb{P}^n$ and $\Gamma \subset \mathrm{PGL}_{n+1}(\mathbb{C})$, then we call it a *complex projective Kleinian group*. Complex projective Kleinian groups have been classified in dimension two by Kobayashi-Ochiai, Mok-Yeung, Klingler and Cano-Seade (see [KO80],

[MY93], [Kli98], [CS14]). If the quotient manifold is projective then they have been almost classified in any dimension ≥ 2 (see [JR15]). A reference on the subject is the book [CNS13] where the authors use the terminology “complex Kleinian group”.

Another class of birational Kleinian groups that has been studied is the so called *Anosov subgroups*. The theory of Anosov representation developed in recent years leads to the construction by Guichard-Wienhard [GW12] and Kapovich-Leeb-Porti [KLP18] of many birational Kleinian groups (Y, Γ, U, X) such that Y is a flag variety G/P , the quotient of a complex linear group G by a parabolic subgroup P , and Γ is some discrete subgroup of G called Anosov subgroup. Note that in most known cases of complex projective Kleinian groups (see [CNS13]) or Anosov subgroups (see [DS20]), the quotient compact complex manifolds are not Kähler.

For complex projective Kleinian groups or Anosov subgroups, transformations in Γ are in the neutral component of the automorphism group of the variety which is a Lie group. Such an automorphism acts trivially on the cohomology of the variety. An automorphism acting non trivially on cohomology has more complicated dynamical properties (see [Can18a]). In general the group of birational transformations does not even act directly on the cohomology. When $n \geq 2$ the group of birational transformations of \mathbb{P}^n is much larger than $\mathrm{PGL}_n(\mathbf{C})$ and has complicated topological structure (see [BF13]). In particular one needs an infinite number of parameters to describe it. We emphasize also the fact that the action of a group of birational transformations is not really a set-theoretic group action on the whole variety: a birational transformation could fail to be well defined at some points, or fail to be locally a diffeomorphism at some points where it is defined. These are some of the reasons why birational Kleinian groups seem to be much more complicated. Beyond complex projective Kleinian groups and Anosov subgroups, even if we restrict to automorphisms, there is essentially nothing known.

1.1.4. *Uniformization and geometric structures.* Koebe-Poincaré’s uniformization theorem asserts that every simply connected Riemann surface is biholomorphic to \mathbb{P}^1 , \mathbf{C} or the unit disk \mathbb{D} ; it plays an omnipresent role in the study of classical Kleinian groups. Universal covers of projective varieties of dimension ≥ 2 can be very complicated, without speaking about general compact complex manifolds. A reasonable way to study them is by adding additional hypothesis. The three elements of the uniformization are: the fundamental group, the universal cover and the action of the fundamental group on it. There are various way to specify these data. For example saying that a hyperbolic Riemann surface is covered by the unit disk with deck transformations contained in $\mathrm{PSL}_2(\mathbf{R})$ does not tell the arithmetic properties of the coefficients of the matrices.

From this point of view, our paper is similar in the theme to Claudon-Höring-Kollár’s work [CHK13]. In [CHK13] it is proved under the abundance conjecture that the only projective varieties with quasi-projective universal covers are bundles over abelian varieties or their finite quotients. Kollár-Pardon investigated in [KP12] the situation where we replace quasi-projective universal covers with semi-algebraic universal covers, and abelian varieties with quotients of noncompact hermitian symmetric spaces; a similar statement was proved with more additional assumptions. Recall that abelian varieties and locally symmetric varieties are two basic examples of birational Kleinian groups.

The results of [CHK13] and [KP12] suggest that requiring universal covers to be algebraic in some sense is very restrictive. Our study amounts to put, in the problem of uniformization, an algebraic hypothesis on the action of deck transformations instead of on the universal cover itself. We should think of it as uniformization with geometric models. More precisely it is a uniformization with Ehresmann’s geometric structures, i.e. the

so called (G, X) -structures. A (G, X) -structure on a manifold is an atlas of charts with values in X and locally constant changes of coordinates in G . Classically G is a Lie group acting transitively on some model manifold X . Here we consider the more general notion of $(\text{Bir}(Y), Y)$ -structures. For example a $(\text{Bir}(\mathbb{P}^n), \mathbb{P}^n)$ -structure is an atlas of charts such that the changes of coordinates can be written as rational fractions. Quotient manifolds of birational Kleinian groups can be considered as the simplest $(\text{Bir}(Y), Y)$ -manifolds; they have injective developing maps. We refer to [Dlo16] and [Zhac] for $(\text{Bir}(Y), Y)$ -structures. A similar notion called bi-algebraic structure has been introduced in [KUY16].

There are very few known examples of domains in algebraic varieties of dimension > 1 that are infinite covers of projective varieties. We expect such domains to be holomorphically convex (see [EKPR12]). Let $m \geq 2$. Wong [Won77] and Rosay [Ros79] proved that if a bounded domain in \mathbf{C}^m covers a projective variety and has smooth boundary then it is biholomorphic to the unit ball. Frankel [Fra89] proved that the only convex bounded domains in \mathbf{C}^m that cover projective varieties are bounded symmetric domains. These results suggest that domains covering projective varieties are hard to describe if they are not bounded symmetric domains. The only other examples of domains that the author knows are universal covers of Kodaira fibrations and their variants. A Kodaira fibration is a projective surface with a submersive non-isotrivial fibration onto a hyperbolic Riemann surface which is a differentiable fiber bundle (see [Kod67]). It follows from Bers' work on Teichmüller theory that the universal cover of a Kodaira fibration is biholomorphic to a bounded domain in \mathbf{C}^2 (see [Gri71]). We will see that there is no birational Kleinian group (Y, Γ, U, X) such that X is a Kodaira fibration.

1.2. Main results. First we observe that birational Kleinian groups only exist on those varieties whose groups of birational transformations are large enough.

Theorem A. *There are no birational Kleinian groups acting on projective varieties of Kodaira dimension ≥ 0 .*

The core of our paper concerns only dimension two. We obtain an almost complete classification of birational Kleinian groups in dimension two. See Section 10.1 for the full result. Our result is new even for groups of automorphisms. Here we state a short version of our classification for birational Kleinian groups with Kähler quotients and simply connected domains. There are two hermitian symmetric spaces of noncompact type in dimension two: the Euclidean ball \mathbb{B}^2 in \mathbb{P}^2 and the bidisk $\mathbb{D} \times \mathbb{D}$ in $\mathbb{P}^1 \times \mathbb{P}^1$. Roughly speaking our classification says that apart from these two cases of hermitian symmetric spaces and the case of complex tori, all other birational Kleinian groups having Kähler quotients come from one-dimensional classical Kleinian groups.

Theorem B. *Let (Y, U, Γ, X) be a birational Kleinian group in dimension two such that U is simply connected and X is Kähler. Then up to geometric conjugation and up to taking a finite index subgroup, we are in one of the following six situations:*

- (1) $Y = \mathbb{P}^2$, $U = \mathbf{C}^2$, Γ is a group of translations isomorphic to \mathbf{Z}^4 and X is a complex torus.
- (2) $Y = \mathbb{P}^1 \times \mathbb{P}^1$, U is the bidisk $\mathbb{D} \times \mathbb{D}$ and Γ is a torsion free irreducible cocompact lattice in $\text{PSL}(2, \mathbf{R}) \times \text{PSL}(2, \mathbf{R}) \subset \text{Aut}(\mathbb{P}^1) \times \text{Aut}(\mathbb{P}^1)$.
- (3) $Y = \mathbb{P}^2$, U is the Euclidean ball \mathbb{B}^2 , Γ is a torsion free cocompact lattice in $\text{PU}(1, 2) \subset \text{PGL}(3, \mathbf{C}) = \text{Aut}(\mathbb{P}^2)$.
- (4) $\Gamma = \Gamma_1 \times \Gamma_2$ is a product of two classical Kleinian groups (one possibly trivial), U is a product $D_1 \times D_2$ embedded in $Y = \mathbb{P}^1 \times \mathbb{P}^1$, where $D_i \subset \mathbb{P}^1$ is a simply connected invariant component of the domain of discontinuity of Γ_i .

- (5) $Y = \mathbb{P}^1 \times \mathbb{P}^1$, $U = D_1 \times \mathbb{C}$ where D_1 is a simply connected invariant component of the domain of discontinuity of a classical Kleinian group Γ_1 . The projection of $\mathbb{P}^1 \times \mathbb{P}^1$ onto the first factor is equivariant under a surjective homomorphism $\Gamma \rightarrow \Gamma_1$. The quotient surface X is a principal elliptic bundle over D_1/Γ_1 . Any element of Γ has the form $(x, y) \mapsto (\gamma_1(x), y + R(x))$ where $\gamma_1 \in \Gamma_1$ and $R \in \mathbb{C}(x)$ has no poles in D_1 .
- (6) $Y = \mathbb{P}^1 \times \mathbb{P}^1$, $U = D_1 \times \mathbb{P}^1$ where D_1 is a simply connected invariant component of the domain of discontinuity of a classical Kleinian group Γ_1 . The projection of $\mathbb{P}^1 \times \mathbb{P}^1$ onto the first factor is equivariant under an isomorphism $\Gamma \rightarrow \Gamma_1$. The quotient surface X is a geometrically ruled surface over D_1/Γ_1 .

Classical Kleinian groups that appear in Theorem B have simply connected invariant components in their domains of discontinuity. Examples are fuchsian groups, quasi-fuchsian groups and their limits ([Ber70], [Mas88]). Any such group is a limit of quasi-fuchsian groups (see [BCM12]).

1.3. Strategy and plan.

1.3.1. *Strategy.* Let (Y, U, Γ, X) be a birational Kleinian group. For simplicity and to explain the strategy in the most important case, we assume that Y is a rational surface, X is Kähler and U is simply connected. Thus Γ can be identified with the fundamental group $\pi_1(X)$. We give here a sketch of the main steps of the proof of Theorem B. The proof is a mixture of three topics: dynamics of birational transformations, fundamental groups of compact Kähler manifolds (a.k.a Kähler groups) and holomorphic foliations. We explain here briefly how they interact in our study.

Manin [Man86] introduced a faithful action of the group of birational transformations $\text{Bir}(Y)$ by isometries on an infinite dimensional hyperbolic space \mathbb{H}_Y (see [Can11]). Thus we have an action of $\Gamma = \pi_1(X)$ on \mathbb{H}_Y . Very strong restrictions on isometric actions of Kähler groups on finite dimensional hyperbolic spaces are found by Carleson-Toledo [CT89] and they are generalized by Delzant-Py [DP12] to infinite dimensional hyperbolic spaces; their works are based on the very developed theory of harmonic mappings on Kähler manifolds. We combine Delzant-Py's results with the Kleinian property of Γ to prove that the action of Γ on \mathbb{H}_Y has fixed points in \mathbb{H}_Y or on the boundary of \mathbb{H}_Y . Cantat proved in [Can11] that the actions on Y of finitely generated groups of birational transformations with fixed points in $\mathbb{H}_Y \cup \partial\mathbb{H}_Y$ have highly constrained geometric features. In our case of birational Kleinian groups, we deduce from Cantat's result that either Γ is conjugate to a group of automorphisms, or to a group preserving a rational fibration.

If Γ is conjugate to a group of automorphisms, then we can reduce to the case of complex projective Kleinian groups. We can thus apply the classification of complex projective Kleinian groups done by Kobayashi-Ochiai [KO80], Mok-Yeung [MY93], Klingler [Kli98] and Cano-Seade [CS14].

Consider now the case where Γ preserves a rational fibration. Then we observe that the rational fibration induces a regular holomorphic foliation on X . Brunella did an almost complete classification of regular holomorphic foliations on compact complex surfaces in [Bru97]; however there is one missing case. The foliations appearing in our study have additional structures: they support foliated $(\text{PGL}(2, \mathbb{C}), \mathbb{C})$ -structures along the leaves. Recently in [DG21] Deroin-Guillot proved a property of foliated $(\text{PGL}(2, \mathbb{C}), \mathbb{C})$ -structures that the missing case in Brunella's classification cannot satisfy. Combining

Derooin-Guillot's theorem with Brunella's classification we have a detailed list of possible foliations appearing in our study. Then we examine whether and how each foliation on X is realized. This analysis will occupy half of the paper.

1.3.2. *Plan.* In Section 2 we prove Theorem A. In the rest of the paper we only work in dimension two. In Section 3 we present the known classification of complex projective Kleinian groups. Preliminaries on groups of birational transformations of projective surfaces are introduced in Section 4. In Section 5 we prove that birational Kleinian groups have fixed points on $\mathbb{H}_Y \cup \partial\mathbb{H}_Y$. Section 6 is a glossary of holomorphic foliations on surfaces. In Sections 7, 8 and 9 we study birational Kleinian groups preserving a rational fibration. We consider in Section 7 the case where each fiber of the the rational fibration is invariant, in Section 8 the case where the base is an elliptic curve and in Section 9 the case where the base is rational. At last Section 10 consists of our classification of birational Kleinian groups in dimension two. Results from previous chapters are assembled. It contains the long version of Theorem B and serves as an outline of the materials in previous sections.

Acknowledgements. Most of the work presented in this paper was done at Institut de Recherches Mathématiques de Rennes. I would like to thank warmly Serge Cantat for introducing me to this subject and his constant support. I would like to thank Benoît Claudon, Laura DeMarco, Bertrand Derooin, Adolfo Guillot, Bruno Klingler, Frank Loray, Frédéric Touzet, Junyi Xie for helpful discussions and comments on previous versions of the paper.

2. VARIETIES OF NON-NEGATIVE KODAIRA DIMENSION

2.1. **Invariant measure.** A measure which is absolutely continuous with respect to the Lebesgue measure associated with some Riemannian metric does not charge algebraic sets of codimension ≥ 1 . Therefore the push-forward of such a measure by a birational transformation is well-defined and it makes sense to talk about Γ -invariant measure. More precisely for ν such a measure, $\gamma \in \Gamma$ a birational transformation and $E \subset Y$ a subset, $\gamma_*\nu(E) = \nu(\gamma^{-1}(E \setminus \text{Ind}(\gamma^{-1})))$ where Ind means the set of indeterminacy points. The objective of this section is to prove:

Lemma 2.1.1. *Let Y be a smooth projective variety of non-negative Kodaira dimension. Let Γ be a subgroup of $\text{Bir}(Y)$. Then there is a Γ -invariant probability measure of total support on Y which is absolutely continuous with respect to the Lebesgue measure.*

We now deduce Theorem A from Lemma 2.1.1.

Proof. Let (Y, Γ, U, X) be a birational Kleinian group such that Y has non-negative Kodaira dimension. There exists an open subset W of Y such that $\gamma(W) \cap W = \emptyset$ for any $\gamma \neq e$. Let μ be a Γ -invariant probability measure of total support which is absolutely continuous with respect to the Lebesgue measure. Then $\mu(U) \geq \sum_{\gamma \in \Gamma} \mu(\gamma(W)) = +\infty$, contradiction. \square

Another consequence of the Lemma 2.1.1 is:

Corollary 2.1.2. *Let Y be a smooth projective variety of non-negative Kodaira dimension. Let $f \in \text{Bir}(Y)$ be a birational transformation of infinite order. Then for any non-empty f -invariant Borel subset V of Y , the action of f on V is recurrent. In particular there is no wandering open subset.*

Proof. Let V be a non-empty f -invariant Borel subset. Let $V' = V - \cup_n f^{-n}(\text{Ind}(f))$ be the complement of indeterminacy points of all iterates of f in V . Let μ be a Γ -invariant probability measure of total support which is absolutely continuous with respect to the Lebesgue measure. Then V' is f -invariant and $\mu(V) = \mu(V')$. Poincaré recurrence theorem says that a measure preserving dynamical system on a space of finite measure is always recurrent. \square

Here f -invariant means that f and f^{-1} are regular on V and $f(V) = V$; recurrent means that for any open subset $W \subset V$, for almost all $x \in W$ (with respect to the Lebesgue measure associated with a Riemannian metric), there are infinitely many $n \in \mathbf{N}$ such that $f^n(x) \in W$. An open subset is wandering if its iterates by f are disjoint from each other.

To prove Lemma 2.1.1, let us explain first the construction of the measure. When Y is an abelian variety or a Calabi-Yau variety, the canonical bundle is trivial and we can use the measure associated with the natural non-vanishing volume form defined up to a multiplicative scalar. In the general case we take a non-zero pluricanonical section $\alpha \in H^0(Y, mK_Y)$ which in local coordinates can be written as $\alpha(z_1, \dots, z_n)(dz_1 \wedge \dots \wedge dz_n)^m$ where n is the dimension of Y . Then the differential $2n$ -form

$$(1) \quad (\alpha \wedge \bar{\alpha})^{\frac{1}{m}} = i^{n^2} |\alpha(z)|^{\frac{2}{m}} dz_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_1 \wedge \dots \wedge d\bar{z}_n$$

is a singular volume form which is > 0 on the complement of the divisor $\{\alpha = 0\}$. The finite measure defined by this singular volume form has full support and is absolutely continuous with respect to the Lebesgue measure associated with any Riemannian metric on Y . Hence the proof of Lemma 2.1.1 follows from the existence of a $\text{Bir}(Y)$ -invariant pluricanonical form. This is an immediate consequence of a theorem of Nakamura-Ueno ([Uen75] Chapter VI) saying that the linear representation of $\text{Bir}(Y)$ on the space of pluricanonical forms has finite image. We give here an elementary argument without using the full strength of Nakamura-Ueno's theorem.

For a pluricanonical form $\alpha \in H^0(Y, mK_Y)$ and a birational transformation $f \in \text{Bir}(Y)$, we can pull back α by f outside the indeterminacy locus of f and then extend it in a unique way by Hartogs' Principle. In this way we obtain a linear representation

$$\rho : \text{Bir}(Y) \rightarrow \text{GL}(H^0(Y, mK_Y)).$$

If we pull back the measure associated with a pluricanonical form, then we get the measure associated with the pulled-back pluricanonical form. Lemma 2.1.1 is a consequence of the following statement:

Lemma 2.1.3. *Let Y be a smooth projective variety of non-negative Kodaira dimension. Let Γ be a subgroup of $\text{Bir}(Y)$. Then there is a singular volume form Ξ such that for any $\gamma \in \Gamma$ we have $\gamma^*\Xi = \Xi$.*

Proof. By the assumption on Kodaira dimension, there exists $m \in \mathbf{N}^*$ such that $H^0(Y, K_Y^m)$ is not trivial. There is a norm on the vector space $H^0(Y, K_Y^m)$:

$$\|\alpha\|^2 = \int_Y (\alpha \wedge \bar{\alpha})^{\frac{1}{m}}.$$

For $f \in \text{Bir}(Y)$, a change of variables gives

$$\|\alpha\|^2 = \int_Y (\alpha \wedge \bar{\alpha})^{\frac{1}{m}} = \int_Y (f^*\alpha \wedge \overline{f^*\alpha})^{\frac{1}{m}} = \|f^*\alpha\|^2.$$

This means that the image of the representation $\rho : \text{Bir}(Y) \rightarrow \text{GL}(H^0(Y, K_Y^m))$ is contained in the compact group preserving the norm $\|\cdot\|$. Denote by G the Zariski closure

of $\rho(\text{Bir}(Y))$ in $\text{GL}(\text{H}^0(Y, K_Y^m))$. The group G is compact and admits a left and right invariant Haar measure ν . Let α be a non-zero element of $\text{H}^0(Y, K_Y^m)$. Then

$$\mathfrak{E} = \int_G g^* \left((\alpha \wedge \bar{\alpha})^{\frac{1}{m}} \right) d\nu(g)$$

is a desired singular volume form. \square

2.2. Genus one fibrations. Every surface of Kodaira dimension 1 has a genus one fibration. In this concrete case we present an elementary topological argument to illustrate Theorem A. We will also need this argument later while dealing with Halphen twists on rational surfaces.

Lemma 2.2.1. *Let Λ be an infinite group of automorphisms of an elliptic curve E . Then the closure of the orbit of a point $x \in E$ is a finite union of real subtori.*

Proof. The automorphism group of E is the extension of a finite group by E . Some finite index subgroup $\Lambda' \subset \Lambda$ is thus a subgroup of the Lie group E . The closure $\overline{\Lambda'}$ in E is a Lie group. It is thus the product of a connected real subtorus F with a finite group G isomorphic to $\overline{\Lambda'}/F$. For any $x \in E$ the closure of the orbit $\Lambda' \cdot x$ is $\overline{\Lambda'}x$ which is the union of $|G|$ translates of F . \square

Lemma 2.2.2. *Let $\varphi : Y \rightarrow C$ be a genus one fibration on a surface Y . Let U be an open set of Y which is preserved by an infinite subgroup Γ of $\text{Bir}(Y)$. Suppose that Γ preserves φ and that the induced action on C is finite. Then the action of Γ on U is not discrete.*

Proof. By hypothesis, a finite index subgroup Γ_0 of Γ preserves each fiber of φ . Since this group is infinite and abelian, it contains a countable infinite subgroup Γ_1 . There is a fiber $E = \varphi^{-1}(c)$ of genus 1 which intersects U and on which Γ_1 induces an infinite group of automorphisms; indeed, U is open, so $\varphi(U)$ is uncountable, and if $\gamma \in \Gamma_1$ is not the identity there are at most countably many points $z \in C$ for which $\gamma|_{\varphi^{-1}(z)}$ is the identity. Then, the result follows from Lemma 2.2.1, applied to E and $\Gamma_1|_E$. \square

If Y has Kodaira dimension one then the genus one fibration $\pi : Y \rightarrow C$ is induced by a pluricanonical linear system (see [Iit70]). Every birational transformation of Y preserves π . The induced action of $\text{Bir}(Y)$ on C is finite by Nakamura-Ueno Theorem. Therefore Lemma 2.2.2 implies Theorem A for these surfaces.

3. COMPLEX PROJECTIVE KLEINIAN GROUPS

3.1. Classification. This section is a presentation of the known classification of complex projective Kleinian groups in dimension two. We first state the classification then describe its cases.

Theorem 3.1.1 (Kobayashi-Ochiai [KO80], Mok-Yeung [MY93], Klingler [Kli98], Cano-Seade [CS14]). *Let $(\mathbb{P}^2, U, \Gamma, X)$ be a birational Kleinian group such that $\Gamma \subset \text{PGL}(3, \mathbb{C})$. Then up to taking finite index subgroup and up to geometric conjugation inside $\text{PGL}(3, \mathbb{C})$, we are in one of the following situations:*

- (1) $U = \mathbb{C} \times \mathbb{C}, \mathbb{C} \times \mathbb{C}^*$ or $\mathbb{C}^* \times \mathbb{C}^*$ and X is a complex torus; here $\mathbb{C} \times \mathbb{C}, \mathbb{C} \times \mathbb{C}^*, \mathbb{C}^* \times \mathbb{C}^*$ are the standard Zariski open subsets of \mathbb{P}^2 and Γ is a lattice in U viewed as a Lie group.
- (2) U is the Euclidean ball \mathbb{B}^n embedded in the standard way in \mathbb{P}^2 and X is a ball quotient.

- (3) $U = \mathbf{C}^2 \setminus \{0\}$ where \mathbf{C}^2 is embedded in the standard way in \mathbb{P}^2 as a Zariski open set and X is a Hopf surface.
- (4) U is the standard Zariski open set $\mathbf{C}^2 \subset \mathbb{P}^2$ and X is a primary Kodaira surface.
- (5) $U = \mathbb{H} \times \mathbf{C}$ embedded in the standard way in the Zariski open subset \mathbf{C}^2 , X is an Inoue surface and Γ is the group defining the Inoue surface, see Paragraph 3.2.5.
- (6) U is biholomorphic to $\mathbb{H} \times \mathbf{C}^*$ and X is an affine-elliptic bundle; here $\mathbb{H} \times \mathbf{C}^*$ is not embedded in a standard way in \mathbf{C}^2 , see Paragraph 3.2.6.

Remark that the cases of complex tori and ball quotients are the only cases where X is Kähler. In higher dimension, if we want X to be projective, then there is only one more possibility: X can be the total space of a family of abelian varieties over a Shimura curve (Jahnke-Radloff [JR15]).

The quotient surface of a complex projective Kleinian group is equipped with a $(\mathrm{PGL}(3, \mathbf{C}), \mathbb{P}^3)$ -structure. The proof of Theorem 3.1.1 has two main steps. The first one is to determine which surfaces admit $(\mathrm{PGL}(3, \mathbf{C}), \mathbb{P}^3)$ -structures and the second one is to find all $(\mathrm{PGL}(3, \mathbf{C}), \mathbb{P}^3)$ -structures on those candidates. The first problem was studied by Gunning [Gun78] and Kobayashi-Ochiai [KO80]; it is completed by Klingler [Kli98]. In [Kli98], the second problem is also investigated and is almost completed, except the case of ball quotients. The fact that the natural complex projective structure on a ball quotient is the only one is proved by Mok-Yeung in [MY93]; later Klingler gave a different proof in [Kli01]. The version of the classification that we state here can be found in the paper of Cano-Seade [CS14] which contains more precise information, in particular when the group has torsion.

Though not every $(\mathrm{PGL}(3, \mathbf{C}), \mathbb{P}^3)$ -structure arises from a complex projective Kleinian group, it was easier to study first the geometry of surfaces with complex projective structures than to look directly at the dynamics of subgroups of $\mathrm{PGL}(3, \mathbf{C})$. However this is not the strategy that we will use for general birational Kleinian groups: we are based on a good understanding of groups of birational transformations while $(\mathrm{Bir}(Y), Y)$ -structures seem to be more difficult to study.

3.2. Examples. Here we describe all cases in Theorem 3.1.1. Some of them exist in any dimension.

3.2.1. Complex tori. A complex torus is a quotient of the commutative Lie group \mathbf{C}^2 by a lattice, thus is naturally realized as the quotient of a complex affine Kleinian group. The affine varieties $\mathbf{C} \times \mathbf{C}$, $\mathbf{C} \times \mathbf{C}^*$, $\mathbf{C}^* \times \mathbf{C}^*$ are also infinite covers of complex tori and the corresponding groups of deck transformations are also birational.

3.2.2. Ball quotients. Let \mathbb{B}^2 be the unit ball $\{[z_0; \dots; z_2]; |z_0|^2 - |z_1|^2 - |z_2|^2 > 0\} \subset \mathbb{P}^2$. The group of biholomorphisms of \mathbb{B}^2 is $\mathrm{PU}(1, 2)$. If Γ is a torsion-free cocompact lattice of $\mathrm{PU}(1, 2)$, then \mathbb{B}^2/Γ is a projective manifold of general type equipped with a $(\mathrm{PGL}(3, \mathbf{C}), \mathbb{P}^2)$ -structure.

3.2.3. Hopf surfaces. Let $\gamma \in \mathrm{GL}_2(\mathbf{C})$ be a contracting linear automorphism of \mathbf{C}^2 such that $\gamma^n(x) \rightarrow 0$ for any $x \in \mathbf{C}^2 \setminus \{0\}$. Then the quotient of $\mathbf{C}^2 \setminus \{0\}$ by the action of γ is a compact surface with a complex affine structure. Such surfaces are among the so called *Hopf surfaces* and are not Kähler.

3.2.4. *Primary Kodaira surfaces.* Consider the subgroup G of the group of affine transformations $\text{Aff}(2, \mathbf{C})$ generated by the following four elements

$$\begin{aligned} z &\mapsto z + \begin{pmatrix} 0 \\ 1 \end{pmatrix}, & z &\mapsto \begin{pmatrix} 1 & 0 \\ c & 1 \end{pmatrix} z + \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ z &\mapsto z + \begin{pmatrix} 0 \\ a \end{pmatrix}, & z &\mapsto \begin{pmatrix} 1 & 0 \\ d & 1 \end{pmatrix} z + \begin{pmatrix} b \\ 0 \end{pmatrix}. \end{aligned}$$

The quotient \mathbf{C}^2/G is a primary Kodaira surface, i.e. a non Kähler principal bundle of elliptic curves over an elliptic curve; its Kodaira dimension is zero (see [BHPVdV04]). Note that the group G is not unique up to conjugation for a fixed primary Kodaira surface, it has one parameter of freedom (see [Kli98]).

3.2.5. *Inoue surfaces.* Inoue surfaces are discovered in [Ino74] where they were constructed as quotients of complex affine Kleinian groups. There are three types of Inoue surfaces. Their universal covers are all $\mathbb{H} \times \mathbf{C}$ where \mathbb{H} is the Poincaré upper half plane.

An *Inoue surface of type S^0* is a quotient $\mathbb{H} \times \mathbf{C}/G$ where the group G is a subgroup of $\text{Aff}(2, \mathbf{C})$ generated by

$$\begin{aligned} g_0 &: (x, y) \mapsto (\alpha x, \beta y) \\ g_i &: (x, y) \mapsto (x + a_i, y + b_i) \quad \text{for } i = 1, 2, 3. \end{aligned}$$

An *Inoue surface of type S^+* is a quotient $\mathbb{H} \times \mathbf{C}/G$ where G is generated by

$$\begin{aligned} g_0 &: (x, y) \mapsto (\alpha x, y + t) \\ g_i &: (x, y) \mapsto (x + a_i, y + b_i x + c_i) \quad i = 1, 2 \\ g_3 &: (x, y) \mapsto (x, y + \frac{b_1 a_2 - b_2 a_1}{n}) \end{aligned}$$

An Inoue surface of type S^- has a double cover which is an Inoue surface of type S^+ ; in the above formulas it suffices to replace $g_0 : (x, y) \mapsto (\alpha x, y + t)$ by $(x, y) \mapsto (\alpha x, -y)$. We refer to [Ino74] for the precise conditions that the parameters in the formulas have to satisfy.

3.2.6. *Affine-elliptic bundles.* This example is constructed in [Kli98]. Let $S = \mathbb{H}/\pi_1(S)$ be a compact hyperbolic Riemann surface. Let $\bar{\rho} : \pi_1(S) \rightarrow \text{PGL}(2, \mathbf{C})$ and $\phi : \mathbb{H} \rightarrow \mathbb{P}^1$ be the holonomy and developing map of some $(\text{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure on S . By [Gun67], $\bar{\rho}$ always lifts to a representation $\rho : \pi_1(S) \rightarrow \text{GL}_2(\mathbf{C})$. Consider $\mathbf{C}^2 \setminus \{0\}$ as a \mathbf{C}^* -bundle over \mathbb{P}^1 and denote it by W . Consider the \mathbf{C}^* -bundle ϕ^*W over \mathbb{H} obtained by pulling back W via ϕ . The natural complex affine structure on W as an open set of \mathbf{C}^2 induces a complex affine structure on ϕ^*W . The group $\pi_1(S)$ acts on ϕ^*W via its natural action on \mathbb{H} and the representation ρ . This action preserves the affine structure on ϕ^*W . Multiplication in the fibers also preserves this affine structure. Hence the quotient of ϕ^*W by the group generated by $\pi_1(S)$ and a multiplication in the fibers is a compact elliptic surface with a complex affine structure. We call them *affine-elliptic bundles* as in [Kli98]. If the $(\text{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure on S came from a classical Kleinian group then this construction gives rise to a complex affine Kleinian group.

4. PRELIMINARIES ON GROUPS OF BIRATIONAL TRANSFORMATIONS OF SURFACES

This section is a glossary of notions and theorems on groups of birational transformations of surfaces. Most materials presented in this section can be found in the survey [Can18b]. We refer to [Bea96] and [BHPVdV04] for background on birational geometry of surfaces.

4.1. Two subgroups of the Cremona group. The group $\text{Bir}(\mathbb{P}^2)$ is called the *Cremona group*. In homogeneous coordinates a *Cremona transformation* can be written as

$$[x_0; x_1; x_2] \mapsto [P_0; P_1; P_2]$$

where P_1, P_2, P_3 are homogeneous polynomials in x_0, x_1, x_2 and it has an inverse map of the same form. Here are two subgroups of $\text{Bir}(\mathbb{P}^2)$ that we will encounter several times in this text.

4.1.1. The toric subgroup. Fix a system of homogeneous coordinates on the projective plane \mathbb{P}^2 . The complement of three coordinates axes is isomorphic to $\mathbf{C}^* \times \mathbf{C}^*$. A matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}(2, \mathbf{Z})$ acts by automorphism on $\mathbf{C}^* \times \mathbf{C}^*$ as follows: $(x, y) \mapsto (x^a y^b, x^c y^d)$. The group $\mathbf{C}^* \times \mathbf{C}^*$ acts on itself by multiplication. We obtain in this way an embedding of the semi-direct product $\mathbf{C}^* \times \mathbf{C}^* \rtimes \text{GL}(2, \mathbf{Z})$ into the Cremona group $\text{Bir}(\mathbb{P}^2)$. We call its image *the toric subgroup* of the Cremona group. The toric subgroup depends on a choice of homogeneous coordinates in \mathbb{P}^2 . Different choices of coordinates yield conjugate subgroups of the Cremona group and the conjugation is by an element of $\text{PGL}(3, \mathbf{C})$.

4.1.2. The Jonquière's group. Fix an affine chart of \mathbb{P}^2 with coordinates (x, y) . *The Jonquière's group* Jonq is the subgroup of the Cremona group of all transformations of the form

$$(x, y) \mapsto \left(\frac{ax+b}{cx+d}, \frac{A(x)y+B(x)}{C(x)y+D(x)} \right), \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PGL}(2, \mathbf{C}), \quad \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \text{PGL}(2, \mathbf{C}(x)).$$

In other words, Jonq is the group of all birational transformations of $\mathbb{P}^1 \times \mathbb{P}^1$ permuting the fibers of the projection onto the first factor; it is isomorphic to the semi-direct product $\text{PGL}(2, \mathbf{C}) \rtimes \text{PGL}(2, \mathbf{C}(x))$. The Jonquière's group is defined only if a system of coordinates is chosen; a different choice of the affine chart yields a conjugation by an element of $\text{PGL}(3, \mathbf{C})$.

4.2. Hyperbolic geometry and birational transformations. Let Y be a smooth complex projective surface. Manin [Man86] constructed an infinite dimensional hyperbolic space \mathbb{H}_Y and an injective homomorphism from $\text{Bir}(Y)$ to the isometry group $\text{Isom}(\mathbb{H}_Y)$. For $f \in \text{Bir}(Y)$ we denote by f_* the corresponding element of $\text{Isom}(\mathbb{H}_Y)$. There is a dictionary between the dynamics of f_* on \mathbb{H}_Y and the dynamics of f on Y .

Let $f \in \text{Bir}(Y)$. Let $\kappa \in H^{1,1}(Y, \mathbf{R})$ be an ample class of self-intersection 1. The *degree* of f with respect to κ , denoted by $\deg_\kappa(f)$, is the intersection number $f_* \kappa \cdot \kappa$. The *translation length* of f_* on \mathbb{H}_Y is $L(f_*) = \inf_{x \in \mathbb{H}_Y} d(x, f_*(x))$. The sequence $(\deg_\kappa(f^n))^{\frac{1}{n}}$ converges to a real number $\lambda(f) \geq 1$, called the dynamical degree of f (see [DF01]); its logarithm $\log(\lambda(f))$ is the translation length $L(f_*)$.

Elements of $\text{Bir}(Y)$ are classified into four types (see [DF01] and previous works [Giz80], [Can01a], [Can01b]):

- (1) The sequence $\deg_\kappa(f^n)$ is bounded, f is birationally conjugate to an automorphism of a smooth projective surface birational to Y and a positive iterate of f lies in the connected component of identity of the automorphism group of that surface. The isometry f_* is an elliptic isometry of \mathbb{H}_Y . We call f an *elliptic element*.
- (2) The sequence $\deg_\kappa(f^n)$ grows linearly, f preserves a unique pencil of rational curves and f is not conjugate to an automorphism of any projective surface. The isometry f_* is a parabolic isometry of \mathbb{H}_Y . We call f a *Jonquière's twist*.

- (3) The sequence $\deg_k(f^n)$ grows quadratically, f is conjugate to an automorphism of a projective surface preserving a unique genus one fibration. The isometry f_* is a parabolic isometry of \mathbb{H}_Y . We call f a *Halphen twist*.
- (4) The sequence $\deg_k(f^n)$ grows exponentially, i.e. $\lambda(f) > 1$. The isometry f_* is a loxodromic isometry of \mathbb{H}_Y . We call f a *loxodromic element*.

4.3. Tits alternative. A group G is said to satisfy *the Tits alternative* if any subgroup of G contains either a solvable subgroup of finite index, or a non-abelian free subgroup. Tits [Tit72] proved that any linear algebraic group over a field of characteristic zero satisfies the Tits alternative. Lamy [Lam01] proved that the group $\text{Aut}(\mathbf{C}^2)$ of polynomial automorphisms of the affine plane \mathbf{C}^2 satisfies the Tits alternative. For $\text{Bir}(\mathbb{P}^2)$, the Tits alternative has been proved by Cantat [Can11] for finitely generated subgroups and in general by Urech [Ure21]. Furthermore for $\text{Bir}(\mathbb{P}^2)$ the Tits alternative has been improved to the following classification of infinite subgroups; one proves the Tits alternative by proving it for each case in the classification. The bulk of the classification is done by Cantat [Can11] and it is finished by Urech [Ure21]. Besides [Can11] and [Ure21], the works of Weil [Wei55], Gizatullin [Giz80], Blanc-Cantat [BC16] and Déserti [Dés15] are respectively the main input for case 1), 4), 5), 6) in the classification.

Theorem 4.3.1 (Strong Tits alternative for $\text{Bir}(Y)$). *Let Y be a smooth complex projective surface birational to \mathbb{P}^2 and let Γ be an infinite subgroup of $\text{Bir}(Y)$. Then up to taking a finite index subgroup Γ fits in one of the following mutually exclusive situations:*

- (1) *There is a projective surface Y' and a birational map $\phi : Y \dashrightarrow Y'$ such that $\Gamma' = \phi\Gamma\phi^{-1}$ is contained in $\text{Aut}^0(Y')$.*
- (2) *There is a projective surface Y' and a birational map $\phi : Y \dashrightarrow Y'$ such that $\Gamma' = \phi\Gamma\phi^{-1}$ preserves a rational fibration on Y' . All elements of Γ are elliptic but their degrees are not uniformly bounded and the group Γ is not conjugate to a group of automorphisms of any projective surface.*
- (3) *There is a projective surface Y' and a birational map $\phi : Y \dashrightarrow Y'$ such that $\Gamma' = \phi\Gamma\phi^{-1}$ preserves a rational fibration on Y' . At least one element of Γ is a Jonquieres twist.*
- (4) *There is a projective surface Y' and a birational map $\phi : Y \dashrightarrow Y'$ such that $\Gamma' = \phi\Gamma\phi^{-1}$ acts by automorphisms and preserves a genus one fibration. The group Γ is virtually a free abelian group of rank ≤ 8 formed by Halphen twists.*
- (5) *Γ is virtually a cyclic group generated by a loxodromic element.*
- (6) *Γ is solvable but not virtually abelian. There is a birational map $\phi : Y \dashrightarrow \mathbf{C}^* \times \mathbf{C}^*$ such that $\Gamma' = \phi\Gamma\phi^{-1}$ is virtually contained in the toric subgroup $\text{Aut}(\mathbf{C}^* \times \mathbf{C}^*) = (\mathbf{C}^* \times \mathbf{C}^*) \rtimes \text{GL}_2(\mathbf{Z})$. The loxodromic elements of Γ form an infinite cyclic group.*
- (7) *Γ contains a non-abelian free group all of whose non-trivial elements are loxodromic.*
- (8) *Γ is a torsion group, is not finitely generated, and is not conjugate to a group of automorphisms of any projective surface.*

A subgroup is called *elementary* if it fixes a point or preserves a geodesic in \mathbb{H}_Y ; otherwise it is called *non-elementary*. In the above theorem Γ is non-elementary only in the seventh case.

In the first case Γ is called an *elliptic subgroup* of $\text{Bir}(Y)$ because it fixes a point in \mathbb{H}_Y ; in the third or the fourth case, Γ is called a *parabolic subgroup* because it fixes a point in $\partial\mathbb{H}_Y$. It is still an open question whether the second case is possible for finitely generated

subgroups. There are finitely generated examples for all other cases. We will only need Theorem 4.3.1 for finitely generated subgroups.

4.4. Centralizers of birational transformations preserving a rational fibration. Here we collect a series of technical results that we will use in various concrete situations. They can all be found with proofs in [Zhab]. Let Y be a smooth rational surface and $r : Y \rightarrow \mathbb{P}^1$ be a rational fibration. The subgroup of $\text{Bir}(Y)$ that preserves the fibration r will be identified with the Jonquières group. For an element $f \in \text{Jonq}$ (resp. a subgroup $\Gamma \subset \text{Jonq}$) we denote by f_B (resp. Γ_B) the element (resp. subgroup) of $\text{Aut}(\mathbb{P}^1)$ induced by the action on the base of the fibration. The centralizer in $\text{Bir}(Y)$ of an element $f \in \text{Jonq}$ is denoted by $\text{Cent}(f)$. If f is a Jonquières twist then $\text{Cent}(f)$ is necessarily a subgroup of Jonq and we denote by $\text{Cent}_0(f)$ the normal subgroup of $\text{Cent}(f)$ that preserves fiberwise the fibration r . If f is a Jonquières twist then we have an exact sequence

$$\{1\} \rightarrow \text{Cent}_0(f) \rightarrow \text{Cent}(f) \rightarrow \text{Cent}_B(f) \rightarrow \{1\}.$$

Theorem 4.4.1 (Blanc-Déserti [BD15]). *Let $f \in \text{Bir}(\mathbb{P}^2)$ be an elliptic element of infinite order. There exists an affine chart with affine coordinates (x, y) on which f acts by automorphism of the following form:*

- (1) $(x, y) \mapsto (\alpha x, \beta y)$ where $\alpha, \beta \in \mathbf{C}^*$ are such that the kernel of the group homomorphism $\mathbf{Z}^2 \rightarrow \mathbf{C}^*$, $(i, j) \mapsto \alpha^i \beta^j$ is generated by $(k, 0)$ for some $k \in \mathbf{Z}$;
- (2) $(x, y) \mapsto (\alpha x, y + 1)$ where $\alpha \in \mathbf{C}^*$.

The centralizer $\text{Cent}(f)$ of f in $\text{Bir}(Y)$ is respectively described as follows:

- (1) $\text{Cent}(f) = \{(x, y) \mapsto (\eta(x), yR(x^k)) \mid R \in \mathbf{C}(x), \eta \in \text{PGL}(2, \mathbf{C}), \eta(\alpha x) = \alpha \eta(x)\}$.
If α is not a root of unity, i.e. if $k = 0$, then R must be constant.
- (2) $\text{Cent}(f) = \{(x, y) \mapsto (\eta(x), y + R(x)) \mid \eta \in \text{PGL}(2, \mathbf{C}), \eta(\alpha x) = \alpha \eta(x), R \in \mathbf{C}(x), R(\alpha x) = R(x)\}$. If α is not a root of unity then R must be constant and $\eta(x) = \beta x$ for some $\beta \in \mathbf{C}^*$.

Theorem 4.4.2 ([Zhab]). *Let $f \in \text{Jonq}$ be a Jonquières twist such that f_B has infinite order. Then $\text{Cent}_B(f)$ is isomorphic to the product of a finite cyclic group with \mathbf{Z} . The infinite cyclic subgroup generated by f_B has finite index in $\text{Cent}_B(f)$.*

We will also use the above two theorems in the following form:

Theorem 4.4.3 ([Zhab]). *Let $\Gamma \subset \text{Jonq}$. Suppose that Γ is isomorphic to \mathbf{Z}^n and that Γ_B is infinite. Then there is a subgroup Γ' of finite index of Γ with generators $\gamma_1, \dots, \gamma_n$ such that one of the following assertions holds up to conjugation in Jonq :*

- (1) For any i , γ_i has the form $(x, y) \mapsto (a_i x, b_i y)$ with $a_i, b_i \in \mathbf{C}^*$.
- (2) For any i , γ_i has the form $(x, y) \mapsto (x + a_i, b_i y)$ with $a_i \in \mathbf{C}$ and $b_i \in \mathbf{C}^*$.
- (3) For any i , γ_i has the form $(x, y) \mapsto (a_i x, y + b_i)$ with $a_i \in \mathbf{C}^*$ and $b_i \in \mathbf{C}$.
- (4) For any i , γ_i has the form $(x, y) \mapsto (x + a_i, y + b_i)$ with $a_i, b_i \in \mathbf{C}$.
- (5) γ_1 is a Jonquières twist of the form $(x, y) \mapsto (\eta(x), yR(x))$ where $\eta \in \text{PGL}(2, \mathbf{C})$ and $R \in \mathbf{C}(x)^*$. For any $i > 1$, γ_i has the form $(x, y) \mapsto (x, b_i y)$ with $b_i \in \mathbf{C}^*$.
- (6) γ_1 is a Jonquières twist of the form $(x, y) \mapsto (\eta(x), y + R(x))$ where $\eta \in \text{PGL}(2, \mathbf{C})$ and $R \in \mathbf{C}(x)$. For any $i > 1$, γ_i has the form $(x, y) \mapsto (x, y + b_i)$ with $b_i \in \mathbf{C}$.

5. BIRATIONAL KLEINIAN GROUPS ARE ELEMENTARY

5.1. Isometric actions of Kähler groups on hyperbolic spaces. The goal of this section is to show

Theorem 5.1.1. *Let (Y, U, Γ, X) be a birational Kleinian group in dimension two. If X is not a surface of class VII then Γ is an elementary subgroup of $\text{Bir}(Y)$.*

Recall that a non-elementary subgroup of $\text{Bir}(Y)$ is a subgroup that fixes a point or preserves a geodesic in \mathbb{H}_Y (cf. Section 4.3). A surface of class VII is a compact complex surface with Kodaira dimension $-\infty$ and first Betti number 1; they are all non-Kähler. In the next two subsections we will deduce Theorem 5.1.1 from the following theorem of Delzant-Py (the terminology *factors through a fibration* is explained below):

Theorem 5.1.2 (Delzant-Py [DP12]). *Let X be a compact Kähler manifold and Y a rational surface. Let $\rho : \pi_1(X) \rightarrow \text{Bir}(Y)$ be a non-elementary representation, then one of the following two cases occurs:*

- (1) *There is a birational map $\varphi : Y \dashrightarrow \mathbf{C}^* \times \mathbf{C}^*$ such that $\varphi\rho\varphi^{-1}$ is a representation into the toric group $\mathbf{C}^* \times \mathbf{C}^* \rtimes \text{GL}_2(\mathbf{Z})$.*
- (2) *There is a finite index subgroup Γ' of Γ corresponding to a finite étale cover X' of X such that the induced representation $\pi_1(X') \rightarrow \Gamma' \rightarrow \text{Bir}(\mathbb{P}^2)$ factors through a fibration $X' \rightarrow \Sigma$ onto a hyperbolic orbicurve Σ .*

In this article a *hyperbolic orbicurve* Σ will mean a Riemann surface with finitely many marked points with multiplicities, obtained as a quotient of the Poincaré half-plane by the action of a cocompact lattice in $\text{PSL}(2, \mathbf{R})$; marked points are images of the fixed points of the action and multiplicities are orders of the stabilizers. The cocompact lattice is isomorphic to the orbifold fundamental group $\pi_1^{orb}(\Sigma)$ of Σ . A continuous map from a complex manifold X to Σ is holomorphic if it lifts to a holomorphic map from the universal cover \tilde{X} of X to the half-plane, i.e. if there exists a holomorphic map from \tilde{X} to \mathbb{H} such that the following diagram is commutative:

$$\begin{array}{ccc} \tilde{X} & \longrightarrow & \mathbb{H} \\ \downarrow & & \downarrow \\ X & \longrightarrow & \Sigma \end{array}$$

A *fibration* of X onto Σ is a holomorphic surjective map $f : X \rightarrow \Sigma$ with connected fibers. A fibration $f : X \rightarrow \Sigma$ induces a homomorphism $f_* : \pi_1(X) \rightarrow \pi_1^{orb}(\Sigma)$. We say that a group representation $\rho : \pi_1(X) \rightarrow G$ *factors through the fibration* f if there exists a homomorphism $\hat{\rho} : \pi_1^{orb}(\Sigma) \rightarrow G$ such that $\rho = \hat{\rho} \circ f_*$.

Non-elementary isometric actions of Kähler groups on hyperbolic spaces are known to factor through fibrations since the pioneer work of Carlson-Toledo [CT89]. Delzant-Py's Theorem is obtained by generalizing Carlson-Toledo's work to infinite dimension.

5.2. Conjugation. The goal of this subsection is to prove a lemma which will be used in the proof of Theorem 5.1.1 and also in the sequel of our paper. Roughly speaking the lemma says that a conjugation of birational Kleinian groups is always a geometric conjugation. Let (Y, Γ, U, X) be a birational Kleinian group.

Remark 5.2.1. Since Γ acts regularly on U , the indeterminacy points and contracted curves of elements of Γ are disjoint from U .

Lemma 5.2.2. *Suppose that there exists a birational map $\phi : Y \dashrightarrow Y'$ to a second surface Y' and a non-empty Zariski open set $Z' \subset Y'$ such that $\Gamma' = \phi \circ \Gamma \circ \phi^{-1}$ is in $\text{Aut}(Z')$. Then there exist a third surface Y'' , a non-empty Zariski open set $Z'' \subset Y''$ and a birational Kleinian group $(Y'', \Gamma'', U'', X'')$ geometrically conjugate to (Y, Γ, U, X) such that $\Gamma'' \in \text{Aut}(Z'')$.*

The Zariski open set Z'' can be obtained by blowing up Z' . If $Y' = Z'$, then $Y'' = Z''$; in this case if moreover $\Gamma' \subset \text{Aut}^0(Y')$, then a finite index subgroup of Γ'' is in $\text{Aut}^0(Y'')$.

Proof. We first claim that ϕ^{-1} does not contract any curve intersecting Z' onto a point in U . Suppose by contradiction that $u \in U$ is an indeterminacy point of ϕ and a curve C' intersecting Z' is contracted onto u by ϕ^{-1} . Pick a non trivial element $\gamma \in \Gamma$. Denote $\phi \circ \gamma \circ \phi^{-1}$ by γ' . As γ acts freely by diffeomorphism on U , $\gamma(u)$ is a point different from u . And as γ' is an automorphism of Z' , the strict transform $\gamma'(C')$ is still a curve which intersects Z' . Thus $\gamma'(C')$ is contracted by ϕ^{-1} onto $\gamma(u)$. This implies that $\{\gamma(u) | \gamma \in \Gamma\}$ is an infinite set of indeterminacy points of ϕ , which is impossible.

Replacing Y with the blow up at the indeterminacy points of ϕ outside U , we can assume from the beginning that ϕ^{-1} does not contract any curve intersecting Z' . We can also assume that no (-1) -curves contracted by ϕ onto a point of Z' lies outside U . This means that the connected chains of (rational) curves contracted by ϕ onto a point of Z' intersect U in their components which are (-1) -curves. In particular if an indeterminacy point of ϕ^{-1} is in Z' then it is the image of a point in U at which ϕ is regular.

Pick $u' \in Z'$ an indeterminacy point of ϕ^{-1} . Let C be the connected chain of curves on Y contracted onto u' . Then $C \cap U \neq \emptyset$ and for $\gamma \in \Gamma$ the strict transform $\gamma(C) = \overline{\gamma(C \setminus \{\text{ind points of } \gamma\})}$ is still a chain of curves intersecting U . As γ' acts by automorphism on Z' , $\gamma'(u')$ is still a point in Z' and $\gamma(C)$ is contracted onto it. Therefore Γ' permutes the indeterminacy points of ϕ^{-1} in Z' .

Let $Y_1 \xrightarrow{\phi_1} Y'$ be the blow-up at the indeterminacy points of ϕ^{-1} that are in Z' . Let Z_1 be the preimage $\phi_1^{-1}(Z') \subset Y_1$. Since Γ' permutes the indeterminacy points of ϕ^{-1} in Z' , the group $\Gamma_1 = \phi_1^{-1} \circ \Gamma' \circ \phi_1$ acts by automorphisms on Z_1 . Some finite index subgroup of Γ_1 is in $\text{Aut}^0(Y_1)$ if we had $Y' = Z'$ and $\Gamma' \subset \text{Aut}(Y')$. Then we continue the process for Y_1, Z_1 and Γ_1 . After finitely many times, say m times, we get Y_m, Z_m, Γ_m such that the birational map $\phi : Y_m \dashrightarrow Y$ induced by ϕ^{-1} has no indeterminacy points in Z_m . Then our hypothesis at the beginning of the second paragraph implies that ϕ restricted to Z_m is an isomorphism onto image. Hence Γ acts by automorphisms on the Zariski open set $\phi(Z_m) \subset Y$. The proof is finished. \square

5.3. From Delzant-Py's theorem to birational Kleinian groups. This subsection is a proof of Theorem 5.1.1. Throughout the section (Y, Γ, U, X) will be a birational Kleinian group in dimension two. We want to show that Γ is elementary. By Theorem A the Kodaira dimension of Y is necessarily $-\infty$. If Y is not rational, then it is a ruled surface and $\text{Bir}(Y)$ preserves the ruling. This means that $\text{Bir}(Y)$ does not have any non-elementary subgroup. Thus we can and will assume that Y is a *rational surface*.

The open set U is an infinite intermediate Galois covering of X and Γ , being the deck transformation group of the covering, is a quotient of $\pi_1(X)$. Thus we have a representation $\rho : \pi_1(X) \rightarrow \text{Bir}(Y)$, composition of the quotient homomorphism $\pi_1(X) \twoheadrightarrow \Gamma$ and the inclusion $\Gamma \hookrightarrow \text{Bir}(Y)$.

The proof of Theorem 5.1.1 is divided into three parts. In the first part we will show that Γ is not conjugate to a non-elementary subgroup of the toric group. In the second part we show that the representation ρ does not factor through a curve. In the third part we investigate the situation where the quotient X is neither Kähler nor class VII.

If X is Kähler then we can apply Theorem 5.1.2 to the representation ρ . If ρ is non-elementary then there are two possible cases : either it factors through an orbicurve or it is conjugate to a subgroup of the toric subgroup. Thus for Kähler X the first and the second part of this subsection will be sufficient for the proof of Theorem 5.1.1.

5.3.1. *Toric subgroup.* A valuation on the function field $\mathbf{C}(Y) = \mathbf{C}(x, y)$ is a \mathbf{Z} -valued function v on $\mathbf{C}(Y)^*$ such that

- (1) $v(a) = 0$ for any $a \in \mathbf{C}^*$;
- (2) $v(PQ) = v(P) + v(Q)$ and $v(P + Q) \geq \min(v(P), v(Q))$ for any $P, Q \in \mathbf{C}(Y)^*$;
- (3) $v(\mathbf{C}(Y)^*) = \mathbf{Z}$.

The elements of $\mathbf{C}(Y)^*$ with valuations ≥ 0 together with 0 form the valuation ring A_v ; those with valuations > 0 form a maximal ideal M_v . If the residue field A_v/M_v has transcendence degree 1 over \mathbf{C} , then v is called a *divisorial valuation*. The set of divisorial valuations is in bijection with the set of irreducible hypersurfaces in all birational models of Y (cf. [ZS75]). If v_E is the divisorial valuation associated with an irreducible hypersurface E in some birational model of Y , then for any $P \in \mathbf{C}(Y)^*$ the value $v_E(P)$ is the vanishing order of P along E . We can identify $\text{Bir}(Y)$ with the group of automorphisms of $\mathbf{C}(Y)$. Thus $\text{Bir}(Y)$ acts on the set of valuations by precomposition. It preserves the subset of divisorial valuations.

Lemma 5.3.1. *Suppose that Y is a smooth compactification of $\mathbf{C}^* \times \mathbf{C}^*$. Let γ be a loxodromic element in the toric subgroup. Then any irreducible component of $Y \setminus (\mathbf{C}^* \times \mathbf{C}^*)$ is contracted by a power of γ .*

Proof. The complement $Y \setminus (\mathbf{C}^* \times \mathbf{C}^*)$ has only finitely many irreducible components. Let v_E be a divisorial valuation with E outside of $\mathbf{C}^* \times \mathbf{C}^*$. It is sufficient to prove that v_E has an infinite orbit under γ . Let (x, y) be the standard coordinates on $\mathbf{C}^* \times \mathbf{C}^*$. As E is outside $\mathbf{C}^* \times \mathbf{C}^*$, at least one of x, y has non-zero valuation.

The transformation γ can be written as $(x, y) \mapsto (\alpha x^a y^b, \beta x^c y^d)$ with $\alpha, \beta \in \mathbf{C}^*$ and $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbf{Z})$. The matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is hyperbolic because γ is loxodromic. We have for any $n \in \mathbf{Z}$

$$\begin{pmatrix} (\gamma^n \cdot v_E)(x) \\ (\gamma^n \cdot v_E)(y) \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}^n \begin{pmatrix} v_E(x) \\ v_E(y) \end{pmatrix}.$$

Since $\begin{pmatrix} v_E(x) \\ v_E(y) \end{pmatrix}$ is a non-zero integer vector and $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is a hyperbolic matrix, the orbit of $\begin{pmatrix} v_E(x) \\ v_E(y) \end{pmatrix}$ under $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is infinite. This implies that the orbit of v_E under γ is infinite. \square

Lemma 5.3.2. *Suppose that there is a birational map $\phi : Y \dashrightarrow \mathbf{C}^* \times \mathbf{C}^*$ such that $\phi\Gamma\phi^{-1}$ is contained in the toric subgroup. Suppose that Γ is not virtually a cyclic group. Then (Y, Γ, U, X) is geometrically conjugate to a birational Kleinian group (Y', Γ', U', X') such that Y' is a compactification of $\mathbf{C}^* \times \mathbf{C}^*$ and $\Gamma' \subset \text{Aut}(\mathbf{C}^* \times \mathbf{C}^*)$.*

Proof. By Lemma 5.2.2 we can and will assume that there exists a blow up Z of $\mathbf{C}^* \times \mathbf{C}^*$ such that Y a compactification of Z and $\Gamma \subset \text{Aut}(Z)$. Consider the blow down map $g : Z \rightarrow \mathbf{C}^* \times \mathbf{C}^*$ which is equivariant under the action of Γ .

Suppose that C is a connected curve on Z such that $g(C)$ is a point. Up to replacing Γ by a finite index subgroup, we can and will assume that each irreducible component of a curve contracted by g is invariant under Γ . Thus $g(C)$ is a fixed point of Γ in $\mathbf{C}^* \times \mathbf{C}^*$. Without loss of generality we can assume that $g(C)$ is the point $(1, 1)$. The stabilizer of $(1, 1)$ in the toric group is $\text{GL}(2, \mathbf{Z})$. Thus $\Gamma \subset \text{GL}(2, \mathbf{Z})$. Let Z_1 be the blow up of $\mathbf{C}^* \times \mathbf{C}^*$ at $(1, 1)$ and let $E_1 \subset Z_1$ be the exceptional curve. The induced action of $\text{GL}(2, \mathbf{Z})$ on E_1 is exactly the usual action of $\text{GL}(2, \mathbf{Z})$ on \mathbb{P}^1 . Note that g factors through Z_1 . If $C \neq E_1$ then

Γ must have a fixed point in E_1 to be blown up further. This would imply that Γ is cyclic because $\Gamma \subset \mathrm{GL}(2, \mathbf{Z})$. Therefore we deduce that $C = E_1$.

If $C = E_1$ does not intersect U , then the contraction of C would be a geometric conjugation of our birational Kleinian group and the proof is finished. Suppose by contradiction that $U_1 = U \cap C$ is not empty. Then U_1/Γ is a compact curve on X . In other words U_1 is a union of invariant components of a classical Kleinian group isomorphic to Γ . However this classical Kleinian group is contained in $\mathrm{GL}(2, \mathbf{Z})$. It must be cyclic, contradiction. \square

Proposition 5.3.3. *Suppose that there is a birational map $\phi : Y \dashrightarrow \mathbf{C}^* \times \mathbf{C}^*$ such that $\phi\Gamma\phi^{-1}$ is contained in the toric subgroup. Suppose that Γ is not virtually a cyclic group generated by a loxodromic element. Then Γ contains no loxodromic element.*

Proof. By Lemma 5.3.2 we can suppose that Y is a compactification of $\mathbf{C}^* \times \mathbf{C}^*$ and $\Gamma \subset \mathbf{C}^* \times \mathbf{C}^* \rtimes \mathrm{GL}_2(\mathbf{Z})$. Suppose by contradiction that Γ contains a loxodromic element η . Then by Lemma 5.3.1 any irreducible component of $Y \setminus (\mathbf{C}^* \times \mathbf{C}^*)$ is contracted by a power of η . This implies that the open set U on which η acts by biholomorphism is a subset of $\mathbf{C}^* \times \mathbf{C}^*$.

Consider the exponential map π from \mathbf{C}^2 to $\mathbf{C}^* \times \mathbf{C}^*$; it is a covering map. The exponential also gives rise to a homomorphism $\rho : \mathrm{Aff}(2, \mathbf{C}) = \mathbf{C}^2 \rtimes \mathrm{GL}(2, \mathbf{Z}) \rightarrow \mathbf{C}^* \times \mathbf{C}^* \rtimes \mathrm{GL}(2, \mathbf{Z})$. Then π is ρ -equivariant. Consider a connected component \bar{U} of $\pi^{-1}(U)$ and the subgroup $\bar{\Gamma}$ of $\rho^{-1}(\Gamma)$ that preserves \bar{U} . Then $\bar{\Gamma}$ is a complex affine Kleinian group. We can apply Theorem 3.1.1 to $\bar{\Gamma}$. There are five possibilities for X : it is a torus, a Hopf surface, an Inoue surface, a primary Kodaira surface or an affine-elliptic bundle. From the descriptions that we give in Section 3.2, we see that, for the linear part of $\bar{\Gamma}$ to have a hyperbolic matrix, the only possibilities are Inoue surfaces and affine-elliptic bundles. But in these two cases the group is not contained in $\mathrm{GL}(2, \mathbf{Z})$. For Inoue surfaces we can see this from the arithmetic properties of the coefficients in the formulas (see [Ino74]). For affine-elliptic bundles it is because $\mathrm{GL}(2, \mathbf{Z})$ has no Kleinian subgroup with compact hyperbolic Riemann surface as quotient. Hence the assumption that Γ contains a loxodromic element is absurd. \square

5.3.2. Factorization through curves.

Lemma 5.3.4. *If Γ contains a loxodromic element of $\mathrm{Bir}(Y)$, then for any compact curve C on X the image of the composition $\pi_1(C) \rightarrow \pi_1(X) \twoheadrightarrow \Gamma$ is infinite.*

Proof. By Theorem A we can assume that Y is a birationally ruled surface. All birational transformations of a non rational ruled surface preserve the ruling, thus are not loxodromic. We can and will further assume that Y is rational.

Suppose by contradiction that C is a compact curve on X such that $\pi_1(C) \rightarrow \Gamma$ has finite image. Denote by D the normalization of C . Then there is a finite unramified cover $\bar{D} \rightarrow D$ such that the composition $\pi_1(\bar{D}) \rightarrow \pi_1(D) \rightarrow \pi_1(X) \twoheadrightarrow \Gamma$ is trivial. This implies the existence of a map $\iota : \bar{D} \rightarrow U$ lifting the composition map $\bar{D} \rightarrow D \rightarrow C \rightarrow X$. Only a finite subgroup of Γ preserves $\iota(\bar{D})$ by Lemma 5.3.5 below. Therefore $\{\gamma(\iota(\bar{D}))\}_{\gamma \in \Gamma}$ form an infinite family of disjoint smooth compact curves in $U \subset Y$. Every element of Γ permutes these curves. Denote by $\alpha_j, j \in \mathbf{N}^*$ the classes of these curves in the Néron-Severi group of Y . Note that the Picard group is isomorphic to the Néron-Severi group because Y is rational. The intersection number $\alpha_i \cdot \alpha_j$ is zero for $i \neq j$ since the corresponding curves are disjoint. As the Néron-Severi group has finite rank, we can suppose that for some $r \in \mathbf{N}^*$, the classes $\alpha_1, \dots, \alpha_r$ are linearly independent and for any $n > r$ the class

α_n is equal to a linear combination of the $\alpha_j, j \leq r$. Among the $\alpha_j, j \leq r$, at most one has zero self-intersection because otherwise there would exist a two-dimensional totally isotropic space which contradicts the Hodge index theorem. When we write α_n as a linear combination of the $\alpha_j, j \leq r$, if the coefficient before α_j is non-zero then α_j has zero self-intersection because $\alpha_n \cdot \alpha_j = 0$. This implies that all but finitely many of the α_j are equal to a class β of zero self-intersection. Thus the linear system associated with β has dimension ≥ 1 . Since the group Γ permutes the curves $\gamma(\iota(\bar{D}))$, the class β is Γ -invariant. Hence we obtain a pencil of curves invariant under Γ . However a loxodromic birational transformation preserves no pencils of curves. \square

Lemma 5.3.5. *An infinite order automorphism of an irreducible compact curve cannot be free and properly discontinuous.*

Proof. An automorphism of a singular irreducible curve permutes the singular points and is not free. An automorphism of \mathbb{P}^1 has a fixed point. An automorphism of infinite order of a genus one curve without fixed points is an irrational translation; its action is not properly discontinuous by Lemma 2.2.1. An automorphism of a general type curve has finite order. \square

Proposition 5.3.6. *Suppose that Γ is a non-elementary subgroup. Then the representation $\rho : \pi_1(X) \rightarrow \Gamma \hookrightarrow \text{Bir}(Y)$ does not factor through a hyperbolic orbicurve.*

Proof. Suppose by contradiction that ρ is non-elementary and factors through a hyperbolic orbicurve Σ . Let F be a general fibre of the fibration $X \rightarrow \Sigma$. The image of $\phi : \pi_1(F) \rightarrow \pi_1(X)$ is in the kernel of $\pi_1(X) \rightarrow \pi_1^{\text{orb}}(\Sigma)$, thus in the kernel of ρ . This contradicts Lemma 5.3.4. \square

5.3.3. *Elliptic surfaces.* According to Kodaira's classification of surfaces (cf. [BHPVdV04]), a non-Kähler surface which is not of class VII fits in one of the two following possibilities: either X is a primary or secondary Kodaira surface, or X is an elliptic surface with Kodaira dimension 1. The fundamental group of a Kodaira surface is solvable and has no non-elementary representations into $\text{Bir}(Y)$. To finish the proof of Theorem 5.1.1, it remains to consider the case where X is an elliptic surface of Kodaira dimension 1.

Assume that $X \rightarrow \Sigma$ is a genus one fibration and that X has Kodaira dimension one. We have an exact sequence

$$1 \rightarrow H \rightarrow \pi_1(X) \xrightarrow{\varphi} \pi_1^{\text{orb}}(\Sigma) \rightarrow 1$$

where H is the image of the fundamental group of a regular fiber in $\pi_1(X)$ (see [GS85] Theorem 1).

Suppose by contradiction that the representation $\rho : \pi_1(X) \rightarrow \Gamma \hookrightarrow \text{Bir}(Y)$ is non-elementary. By Proposition 5.3.4 $\rho(H)$ is not finite. As a regular fiber is an elliptic curve, $\rho(H)$ must be an infinite abelian group. Since Γ is non-elementary, there exists at least one element $a \in \pi_1(X)$ such that $\varphi(a)$ has infinite order and $\rho(a)$ is a loxodromic element. Consider $\langle \varphi(a) \rangle$ the infinite cyclic subgroup of $\pi_1(\Sigma)$ generated by $\varphi(a)$. Denote by G the subgroup $\varphi^{-1}(\langle \varphi(a) \rangle)$ of $\pi_1(X)$. Then G is an extension of $\langle \varphi(a) \rangle$ by H ; in particular it is solvable. The group $\rho(G)$ is solvable but not virtually cyclic; it contains a loxodromic element $\rho(a)$. By Theorem 4.3.1 we infer that a conjugate of $\rho(G)$ is contained in the toric subgroup and that $\rho(H)$ is an infinite elliptic subgroup. The whole group Γ normalizes $\rho(H)$. By a theorem of Cantat (see Appendix of [DP12]) we infer that Γ is up to conjugation contained in the toric subgroup. This contradicts Proposition 5.3.3.

5.4. Surfaces of class VII. It would be nice to drop the hypothesis that X is not a class VII surface in Theorem 5.1.1. Note that no known surface of class VII has non-solvable fundamental group. Thus an analogue of Theorem 5.1.1 for class VII surfaces could be a vacuous statement. On the other hand in [CT97] Carleson-Toledo applied the same method of [CT89] to study isometric actions of fundamental groups of class VII surfaces on finite dimensional hyperbolic spaces, by replacing harmonic mappings with hermitian harmonic mappings. So one could expect an analogue of Delzant-Py's theorem for class VII surfaces.

6. FOLIATED SURFACES

This section is a glossary of holomorphic foliations on surfaces. We first present Brunella's classification in 6.1. In 6.2 we state Deroin-Guillot's theorem on foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structures. In this paper we only consider holomorphic foliations without singularities, i.e. regular holomorphic foliations.

6.1. Brunella's Theorem. The vocabulary in the following theorem will be explained below.

Theorem 6.1.1 (Brunella [Bru97]). *Let X be a compact complex surface and \mathcal{F} a regular holomorphic foliation on X . Then one of the following situations holds:*

- (1) \mathcal{F} comes from a fibration of X onto a curve;
- (2) X is a complex torus and \mathcal{F} is an irrational linear foliation;
- (3) \mathcal{F} is an obvious foliation on a non-elliptic Hopf surface;
- (4) \mathcal{F} is an obvious foliation on an Inoue surface;
- (5) \mathcal{F} is an infinite suspension of \mathbb{P}^1 or of an elliptic curve over a compact Riemann surface;
- (6) \mathcal{F} is a turbulent foliation with at least one invariant fiber;
- (7) \mathcal{F} is a transversely hyperbolic foliation with dense leaves whose universal cover is a fibration of disks over a disk.

The possible overlaps in the above list are: case 2 with case 5, case 3 with case 6, case 5 with case 6. When a complex torus admits an elliptic fibration, it is straightforward to see that an irrational linear foliation is also a suspension of an elliptic curve. For overlaps between case 3 and case 6, or between case 5 and case 6, see Section 9.1.1.

6.1.1. Fibrations. Let X be compact complex surface and let $f : X \rightarrow B$ be a fibration whose singular fibers are all multiples of smooth curves. The fibration equips X with a regular foliation whose leaves are the underlying manifolds of the fibers. Let mF_b be a fiber with multiplicity m lying over a point $b \in B$. Let w be a local coordinate on B which vanishes at b , let h be a local equation of F_b in X , then $f^*(dw)/h^{m-1}$ is a local differential form defining the foliation.

6.1.2. Linear foliations on tori. Let Λ be a lattice in \mathbf{C}^2 . The quotient $X = \mathbf{C}^2/\Lambda$ is a two dimensional complex torus. Let (z, w) be the natural coordinates on \mathbf{C}^2 . A constant holomorphic differential $adz + bdw$ on \mathbf{C}^2 with $a, b \in \mathbf{C}$ and $ab \neq 0$ descends onto X and defines a regular foliation \mathcal{F} on X ; this is called a *linear foliation* on the torus X . Choosing such a linear foliation amounts to choose a one dimensional \mathbf{C} -linear subspace $W = \ker(adz + bdw)$ of \mathbf{C}^2 . If $W \cap \Lambda$ is a lattice in W , then the leaves of \mathcal{F} are elliptic curves and the foliation is a fibration. If $W \cap \Lambda$ is non empty but not a lattice, then the leaves of \mathcal{F} are biholomorphic to \mathbf{C}^* . If $W \cap \Lambda$ is empty, then the leaves of \mathcal{F} are biholomorphic to \mathbf{C} and are dense in X . When the leaves of \mathcal{F} are not compact, we say the linear foliation \mathcal{F} is *irrational*.

6.1.3. *Obvious foliations on Hopf surfaces.* A *Hopf surface* is a compact complex surface covered by $\mathbf{C}^2 \setminus \{0\}$. A *primary Hopf surface* is a surface biholomorphic to the quotient of $\mathbf{C}^2 \setminus \{0\}$ by a transformation of the form

$$H_{(\alpha, \beta, \gamma, m)} : (z, w) \mapsto (\alpha z + \gamma w^m, \beta w), \quad \alpha, \beta, \gamma \in \mathbf{C}, m \in \mathbf{N}^*, 0 < |\alpha| \leq |\beta| < 1$$

with $\alpha = \beta^m$ if $\gamma \neq 0$. We call the model $(\mathbf{C}^2 \setminus \{0\}) / \langle H_{(\alpha, \beta, \gamma, m)} \rangle$ a standard primary Hopf surface. Any Hopf surface has a finite unramified cover which is a primary Hopf surface. See [Kod66] for the above assertion and the following:

- (1) The standard primary Hopf surface has an elliptic fibration if and only if $\gamma = 0$ and $\alpha^k = \beta^l$ for some $k, l \in \mathbf{N}^*$.
- (2) If $\gamma = 0$ but $\alpha^k \neq \beta^l$ for any $k, l \in \mathbf{N}^*$, then the two smooth elliptic curves given by $z = 0$ and $w = 0$ are the only curves on the primary Hopf surface.
- (3) If $\gamma \neq 0$ then the smooth elliptic curve given by $w = 0$ is the only curve on the primary Hopf surface.

Suppose that $\gamma = 0$. The complex lines in \mathbf{C}^2 parallel to $z = 0$ and to $w = 0$ form two regular foliations on $\mathbf{C}^2 \setminus \{0\}$ that descend to regular foliations on the Hopf surface. A vector field of the form $az \frac{\partial}{\partial z} + bw \frac{\partial}{\partial w}$ with $a, b \in \mathbf{C}$ and $ab \neq 0$ is invariant under $(z, w) \mapsto (\alpha z, \beta w)$, thus descends to the Hopf surface. It gives rise to a regular foliation whose leaves in $\mathbf{C}^2 \setminus \{0\}$ which are different from the two axes are given by (e^{az}, ce^{bz}) . When $a = b$ this is just the foliation by complex vector lines.

Suppose that $\gamma \neq 0$. The complex lines parallel to $w = 0$ give a regular foliation on the Hopf surface. There exist also regular foliations associated with certain vectors fields on $\mathbf{C}^2 \setminus \{0\}$ of Poincaré-Dulac's form $(mz + aw^m) \frac{\partial}{\partial z} + w \frac{\partial}{\partial w}$. When $m = 1$ the normal form $(z, w) \mapsto (\alpha z + \gamma w, \beta w)$ is a linear transformation and the foliation by complex vector lines in $\mathbf{C}^2 \setminus \{0\}$ is one of these foliations.

All the foliations described above will be called *obvious foliations on Hopf surfaces*.

6.1.4. *Obvious foliations on Inoue surfaces.* See Paragraph 3.2.5 for the formulas defining Inoue surfaces. The vertical and horizontal foliations on $\mathbb{H} \times \mathbf{C}$ descend to two regular foliations on an Inoue surface of type S^0 . Only the vertical foliation on $\mathbb{H} \times \mathbf{C}$ descends to a regular foliation on an Inoue surface of type S^+ or S^- . The foliations described above will be called *obvious foliations on Inoue surfaces*.

6.1.5. *Suspensions.* Let M be a Riemann surface. We denote by \hat{M} a Galois covering of M with deck transformation group G . Let N be another Riemann surface. Let $\alpha : G \rightarrow \text{Aut}(N)$ be a homomorphism of groups. The group G acts on $\hat{M} \times N$ in the following way: $g \cdot (m, n) = (g \cdot m, \alpha(g) \cdot n)$. Then the quotient $X = (\hat{M} \times N) / G$ fibers onto $M = \hat{M} / G$ via the first projection; the fibers are all isomorphic to N . The foliation by $\{\hat{M} \times \{n\}\}_{n \in N}$ descends to a foliation on X which is everywhere transverse to the fibration. We call such a foliation a *suspension of N over M* with monodromy $\alpha : G \rightarrow \text{Aut}(N)$. If M, N are compact, then X is compact; this is the only case we will consider. If moreover $\alpha(G)$ is a finite subgroup of $\text{Aut}(N)$, then the leaves of the foliation are compact and the foliation is in fact a fibration. When $\alpha(G)$ is infinite (since N is compact this is only possible if N is \mathbb{P}^1 or an elliptic curve) the leaves are non-compact and we call such a suspension *infinite*.

Remark that if X is a compact complex surface with an infinite suspension foliation, then X is Kähler. This is clear if it is a suspension of \mathbb{P}^1 because then X is ruled. Assume that it is a suspension of an elliptic curve. We use the notations in the previous paragraph. Up to replacing G by a subgroup of finite index we can assume that $\alpha(G) \subset \text{Aut}(N)$ is an abelian group of translations on the elliptic curve N . Thus there is an action of N on X by

translations in the fibers of the elliptic bundle $X \rightarrow M$. In other words $X \rightarrow M$ is a principal elliptic bundle (see [BHPVdV04] V.5.1). Furthermore the suspension process says exactly that the bundle $X \rightarrow M$ is defined by a locally constant cocycle. By [BHPVdV04] V.5.1 and V.5.3 we infer that the second Betti number of X is even. This implies that X is Kähler (see [BHPVdV04] IV.3.1).

6.1.6. Turbulent foliations. Let X be a compact complex surface and $X \rightarrow B$ an elliptic fibration with constant functional invariant (i.e. all regular fibers are isomorphic) whose singular fibers are all multiples of smooth elliptic curves. Let \mathcal{F} be a regular foliation on X . If a finite number of fibers of the elliptic fibration are \mathcal{F} -invariant and all other fibers are transverse to \mathcal{F} , then \mathcal{F} is called a *turbulent foliation*. The underlying elliptic fibration is locally trivial outside the invariant fibers of \mathcal{F} ; the trivialization is given by the foliation. The invariant fibers are regular or multiples of elliptic curves. Locally around an invariant fiber, let (z, w) be a system of local coordinates such that the fibration is given by $(z, w) \mapsto z^m$ where m is the multiplicity of the fiber. Then the foliation \mathcal{F} is locally defined by a local differential form $dz - A(z)dw$ where A is a holomorphic function vanishing at 0.

6.1.7. Transversely hyperbolic foliations with dense leaves. This is the most difficult type of foliations to study in Brunella's theorem. We refer to [Bru97] and [Tou16] for precise definition and descriptions of transversely hyperbolic foliations. The only known example is as follows. Let Γ be a torsion free cocompact irreducible lattice in $\mathrm{PSL}(2, \mathbf{R}) \times \mathrm{PSL}(2, \mathbf{R})$, then the bidisk quotient $\mathbb{D} \times \mathbb{D}/\Gamma$ is a general type surface with two foliations induced by the product structure of $\mathbb{D} \times \mathbb{D}$. Each leaf is dense in $\mathbb{D} \times \mathbb{D}/\Gamma$ because the projection of an irreducible lattice onto a factor $\mathrm{PSL}(2, \mathbf{R})$ is dense. It is not known whether there exists other examples of regular foliations with dense leaves on general type surfaces. It is in this sense that Brunella's classification is still incomplete. Thanks to Deroin-Guillot's work that we will see shortly, these mysterious foliations will not appear in our study of birational Kleinian groups.

6.2. Foliated complex projective structures.

6.2.1. Deroin-Guillot's Theorem. Let \mathcal{F} be a regular holomorphic foliation on a complex manifold X . A *foliated* $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure on \mathcal{F} is an open covering $\{U_i\}$ of X and submersions $\phi_i : U_i \rightarrow \mathbb{P}^1$, transverse to \mathcal{F} , injective on local leaves, such that the restrictions on a leaf L satisfy $\phi_i|_L \circ \phi_j|_L^{-1} \in \mathrm{PGL}(2, \mathbf{C})$. Obvious foliations on Hopf and Inoue surfaces, turbulent foliations and suspension foliations carry natural foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structures (see [DG21]). Though every Riemann surface carries a $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure, not every foliation carries a foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure. The existence of a *foliated* $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure imposes strong restrictions not only on the foliation but also on the topology of the manifold. We state one of the main result of [DG21] in the generality that we need:

Theorem 6.2.1 (Deroin-Guillot). *If X is a compact complex surface having a regular foliation with a foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure, then the Chern classes of X satisfy $c_1(X)^2 = 2c_2(X)$.*

The strict inequality $c_1^2 > 2c_2$ is satisfied by any Kodaira fibration (see [BHPVdV04] V.14) and by any general type surface equipped with a regular foliation with dense leaves which is not a bidisk quotient (see [Bru97] the last paragraph). Therefore the following statement is a consequence of Deroin-Guillot's theorem (see [Zhaa] for a different proof for Kodaira fibrations):

Corollary 6.2.2. *If X is a general type surface having a regular foliation with a foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure, then the universal cover of X is the bidisk.*

It's well known that all $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structures on a given Riemann surface form an affine space directed by the vector space of quadratic differentials. For a foliation \mathcal{F} , let $T_{\mathcal{F}}$ be the tangent bundle of \mathcal{F} . Consider holomorphic sections of $T_{\mathcal{F}}^{*2}$. They are quadratic differentials along the leaves of \mathcal{F} . If \mathcal{F} admits at least one foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure, then all foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structures on \mathcal{F} form an affine space directed by the vector space of holomorphic sections of $T_{\mathcal{F}}^{*2}$ (see [DG21]).

Proposition 6.2.3. *Let \mathcal{F} be a regular foliation on a surface X defined by a fibration $f : X \rightarrow C$. If \mathcal{F} has a foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure then regular fibers of f are isomorphic. If moreover the fibers of f have genus ≥ 2 , then they have the same $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure.*

Proof. Since \mathcal{F} is a foliation without singularities, the singular fibers of f can only be multiples of smooth curves. Such a fibration is not isotrivial if and only if it is a Kodaira fibration. However we just said that there is no foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure on a Kodaira fibration.

If fibers of f have genus ≥ 2 , then there exists a finite unramified cover \hat{C} of C such that the fibration $\hat{f} : \hat{X} \rightarrow \hat{C}$ obtained by base change has no multiple fibers and is in fact a product. Consider the induced foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure on \hat{X} . The conclusion follows because $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structures on a given Riemann surface form an affine space and there is no non constant holomorphic map from \hat{C} to an affine space. \square

Let Γ be a torsion free cocompact irreducible lattice in $\mathrm{PSL}(2, \mathbf{R}) \times \mathrm{PSL}(2, \mathbf{R})$ and consider the bidisk quotient $X = \mathbb{D} \times \mathbb{D} / \Gamma$. Both foliations on X are equipped naturally with a foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure. All leaves are biholomorphic to the disk and have the same $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure: the standard one of a round disk in \mathbb{P}^1 .

Proposition 6.2.4. *If \mathcal{F} is one of the natural foliation on the quotient of the bidisk by a torsion free cocompact irreducible lattice, then the natural foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure is the unique foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure on \mathcal{F} .*

Proof. It suffices to show that $H^0(X, T_{\mathcal{F}}^{*2}) = \{0\}$. In fact for any positive integer n the line bundle $T_{\mathcal{F}}^{*n}$ has no non-zero sections because $T_{\mathcal{F}}^*$ has Kodaira dimension $-\infty$ by [Bru15] Chapter 9.5 Example 9.3. \square

7. INVARIANT RATIONAL FIBRATION I

In this and the next two sections we suppose that (Y, Γ, U, X) is a birational Kleinian group and there is a Γ -invariant rational fibration $r : Y \rightarrow B$ over a smooth projective curve B . Note that when B is not a rational curve, any birational transformation group of Y preserves automatically r . We have a group homomorphism $\Gamma \rightarrow \mathrm{Aut}(B)$ whose image will be denoted by Γ_B . The ruling r is equivariant with respect to the action of Γ on Y and that of Γ_B on B . If Γ_B is infinite, then B is \mathbb{P}^1 or an elliptic curve. We will study in this section the case where Γ_B is finite. The case where Γ_B will be studied in Sections 8 and 9.

By contracting (-1) -curves which are contained in the fibers of r but are disjoint from U , we can and will assume, without loss of generality, that if a fiber of r intersects U , then its irreducible components of self-intersection -1 all intersect U .

7.1. From invariant fibration to foliation. We recall a basic fact about non relatively minimal ruled surface:

Fact 7.1.1. Let F be a singular fiber of a rational fibration. Then F is a tree of rational curves whose components are of self-intersection ≤ -1 .

Proof. The singular fiber F is obtained by successive blow-ups from a regular fiber of a relatively minimal rational fibration, thus a tree of rational curves. The second assertion holds because original regular fiber is of self-intersection 0 and that a blow-up decreases the self-intersection of the components. \square

Lemma 7.1.2. Let F be a fiber which intersects U . Then no element of Γ has an indeterminacy point on F and no component of F is contracted by an element of Γ .

Proof. Let $\gamma \in \Gamma$ be a non trivial birational transformation. Let us first show that γ does not contract any component of F . If F is a regular fiber, then it has only one irreducible component which is itself and this component intersects U by hypothesis. Since the action of Γ on U is regular, F cannot be contracted.

Now assume that F is a singular fiber. Let $Y \xleftarrow{\varepsilon} Z \xrightarrow{\delta} Y$ be the minimal resolution of indeterminacy of γ . Here Z is a smooth projective surface, ε, δ are birational morphisms and $\gamma = \delta \circ \varepsilon^{-1}$. Suppose by contradiction that some component of F is contracted by γ . Then there exists a component C of F such that the strict transform of C in Z has self-intersection (-1) and is contracted by δ . By Fact 7.1.1 C has self-intersection ≤ -1 . As the self-intersection number decreases after a blow up, the strict transform of C in Z has self-intersection (-1) if and only if C has self-intersection (-1) and no point in C is blown up by ε . But by the hypothesis of our initial setting the (-1) -components of F all intersect U and can not be contracted by γ . This means δ does not contract C , contradiction.

We remark that the total transform of F by γ is also a fiber of r which intersects U . The above reasoning, applied to γ^{-1} , says that γ^{-1} does not contract anything onto F , i.e. γ does not have any indeterminacy point on F . \square

We deduce immediately from Lemma 7.1.2 the following corollary:

Corollary 7.1.3. The group Γ acts by holomorphic diffeomorphisms on $r^{-1}(r(U))$. In particular if $r(U) = B$ then $\Gamma \subset \text{Aut}(Y)$.

The following proposition shows how holomorphic foliations come in our study.

Proposition 7.1.4. Let (Y, Γ, U, X) be a birational kleinian group. Suppose $\Gamma \subset \text{Bir}(Y)$ preserves a fibration $r : Y \rightarrow B$. If a fiber F of the invariant fibration intersects U , then $F \cap U$ contains no singular point of F . Therefore the fibration descends to a regular holomorphic foliation on X .

Proof. Suppose by contradiction that F is a singular fiber and $p \in F \cap U$ is a singular point of F . If $\gamma \in \Gamma$, then $\gamma(p)$ is a singular point of the singular fiber $\gamma(F)$ since Γ preserves the fibration and acts by holomorphic diffeomorphisms on U . However a fibration has only finitely many singular fibers and each singular fiber has only finitely many singular points. Thus the infinite group Γ does not act freely on U , contradiction. \square

From now on we will denote by \mathcal{F} the foliation on X induced by the ruling r .

Proposition 7.1.5. In the setting of Proposition 7.1.4, if moreover r has no singular fibers then the foliation \mathcal{F} is equipped with a foliated $(\text{PGL}(2, \mathbb{C}), \mathbb{P}^1)$ -structure along the leaves.

Proof. The open set $r^{-1}(r(U))$ is biholomorphic to a \mathbb{P}^1 -bundle over $r(U)$. According to Corollary 7.1.3 an element of Γ sends a fiber over a point of $r(U)$ to another fiber by a Möbius transformation. Then the proposition follows directly from the definition of foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure. \square

7.2. Finite action on the base.

Theorem 7.2.1. *Let (Y, Γ, U, X) be a birational Kleinian group on a surface Y which is ruled over a curve B . Assume that Γ preserves the ruling, and Γ induces a finite action on the base curve B . Then up to geometric conjugation and up to taking finite index subgroup, we are in one of the following situations:*

- (1) $Y = B \times \mathbb{P}^1$, $\Gamma \subset \{\mathrm{Id}\} \times \mathrm{PGL}(2, \mathbf{C})$ and $U = B \times D$ where $D \subset \mathbb{P}^1$ is an invariant component of the domain of discontinuity of Γ viewed as a classical Kleinian group.
- (2) Y is $\mathbb{P}(\mathcal{E})$ where \mathcal{E} is an extension of \mathcal{O}_B by \mathcal{O}_B . The extension determines a section s of the ruling. We have $U = Y - s$. The subgroup of $\mathrm{Aut}_B(Y)$ fixing s is isomorphic to \mathbf{C} in which Γ is a lattice. The group Γ is isomorphic to \mathbf{Z}^2 and the surface X is a principal elliptic fiber bundle.
- (3) Y is obtained by blowing up a decomposable ruled surface, U is a Zariski open set of Y whose intersection with each fiber is biholomorphic to \mathbf{C}^* . The group Γ is cyclic and is generated by an automorphism which acts by multiplication in each fiber. The quotient surface X is an elliptic fibration over B with isomorphic regular fibers and whose only singular fibers are multiples of smooth elliptic curves.

The rest of Section 7 is the proof of Theorem 7.2.1; it is divided into three parts. In the first part we will study how Γ acts on each fiber of the ruling. In the second part we present some preliminaries on automorphisms of ruled surfaces. In the third part we finish the proof of Theorem 7.2.1.

7.3. Fiberwise dynamics. We start the proof of Theorem 7.2.1 by making the following reduction: replacing Γ with a finite index subgroup, we can and will assume that $\Gamma_B = \{\mathrm{Id}\}$.

Lemma 7.3.1. *We have $r(U) = B$, $\Gamma \subset \mathrm{Aut}(Y)$. The foliation \mathcal{F} on X is a fibration.*

Proof. The set $V = r(U)$ is an open subset of B because a holomorphic map is open. Since $\Gamma_B = \{\mathrm{Id}\}$ the fibration r induces a commutative diagram:

$$\begin{array}{ccc} U & \xrightarrow{\pi} & X \\ \downarrow & & \downarrow \\ V & \longrightarrow & V \end{array}$$

As X is compact, V should be compact too. This implies that V is the whole curve B . By Corollary 7.1.3, we have $\Gamma \in \mathrm{Aut}(Y)$. The leaves of the foliation on X are the fibers of the map $X \rightarrow V$; they are compact. Thus the foliation is a fibration. \square

Since $\Gamma \subset \mathrm{Aut}(Y)$, we can and will assume, up to replacing Γ by a finite index subgroup, that any irreducible curve contained in a fiber is Γ -invariant. In particular every singular point of a singular fiber is fixed by Γ .

Proposition 7.3.2. *If $F \cong \mathbb{P}^1$ is a regular fiber of $r : Y \rightarrow B$, then $F \cap U$, as a subset of \mathbb{P}^1 , is a union of components of the domain of discontinuity of a classical Kleinian group. If moreover $r : Y \rightarrow B$ has at least one singular fiber, then for any fiber F (singular or non-singular), the intersection $F \cap U$ is biholomorphic to \mathbf{C}^* and Γ is cyclic.*

Proof. Let F_b be the (possibly singular) fiber over a point $b \in B$ such that $F_b \cap U \neq \emptyset$. There is a homomorphism $\Gamma \rightarrow \text{Aut}(F_b)$ with image Γ_b and Γ_b preserves $U \cap F_b$. The group Γ_b is isomorphic to Γ because the action of Γ on U is free. We have a commutative diagram

$$(2) \quad \begin{array}{ccc} U & \xrightarrow{\pi} & X \\ \downarrow r & & \downarrow f \\ B & \longrightarrow & B \end{array}$$

where f is a surjective proper holomorphic map. By [BHPVdV04] III.11 the sheaf $f_*\mathcal{O}_X$ is locally free and the non-singular fibers of f have the same number of connected components. If F_b is a non-singular fiber then it is isomorphic to the projective line \mathbb{P}^1 . The quotient of $U \cap F_b$ by the action of Γ_b is a general fiber of f , that is, a disjoint union of smooth compact curves. This means that Γ_b is a classical Kleinian group and $U \cap F_b$ is a union of connected components of the discontinuity set of Γ_b .

Suppose that the fiber F_b is singular. Every irreducible component of F_b contains at least one singular point. If an irreducible component of F_b intersects U then it contains at most two singular points because otherwise it would be pointwise fixed by Γ_b . In particular Γ_b restricted to each component of F_b is isomorphic to a solvable subgroup of $\text{PGL}(2, \mathbb{C})$. So if there exists at least one singular fiber, then the group Γ is solvable. A solvable torsion free classical Kleinian group is isomorphic to \mathbf{Z} or \mathbf{Z}^2 and its set of discontinuity is biholomorphic to \mathbb{C} or \mathbb{C}^* . This implies that the intersection of U with any fiber is biholomorphic to \mathbb{C} or \mathbb{C}^* . In particular $f : X \rightarrow B$ has connected fibers. Thus for the singular fiber F_b the intersection $F_b \cap U$ is connected and is contained in one component C_b of F_b . The component C_b is necessarily a (-1) -curve because by our hypothesis made at the beginning of Section 7 every (-1) -curve in the fiber F_b intersects U .

Under the hypothesis that there is a singular fiber F_b , we need to rule out the situation where Γ is isomorphic to \mathbf{Z}^2 and the intersection of U with any fiber is biholomorphic to \mathbb{C} . Suppose to the contrary that Γ is isomorphic to \mathbf{Z}^2 . Then the fibration $f : X \rightarrow B$ is a genus one fibration whose singular fibers are multiples of smooth elliptic curves. Non-singular fibers of such a genus one fibration are all isomorphic to each other. Let Δ be a small disk in B centred at b and let (x, y) be the local coordinates around a point of the singular fiber F_b where r is given by x and F_b is defined by $x = 0$. The isotriviality of f implies that, locally over $\Delta \setminus \{0\}$, up to a holomorphic change of coordinates the action of Γ is generated by two transformations $\gamma_j : (x, y) \mapsto (x, y + a_j)$, $j = 1, 2$ where $\mathbf{Z}a_1 + \mathbf{Z}a_2$ is a lattice in \mathbb{C} . In other words there is a sequence of blow-downs $r^{-1}(\Delta) \rightarrow \Delta \times \mathbb{P}^1$ which sends F_b to \mathbb{P}^1 such that the action of Γ on $r^{-1}(\Delta)$ is the lift of the action of \mathbf{Z}^2 on $\Delta \times \mathbb{P}^1$ generated by $\gamma_j : (x, y) \mapsto (x, y + a_j)$, $j = 1, 2$. To obtain $r^{-1}(\Delta)$ from $\Delta \times \mathbb{P}^1$ we blow up the common fixed (infinitely near) points of γ_j , $j = 1, 2$. However the action of γ_j , $j = 1, 2$ on the exceptional divisors are always trivial. Thus there was no blowup so that $r^{-1}(\Delta)$ is biholomorphic to $\Delta \times \mathbb{P}^1$. This contradicts that F_b is a singular fiber. \square

7.4. Automorphism groups of geometrically ruled surfaces. Before continuing our study of birational Kleinian groups preserving a rational fibration, we collect some preliminaries on automorphism groups of geometrically ruled surfaces, following Maruyama's paper [Mar71]. Our notations are those of [Har77] but are different from those of [Mar71]. Let Y be a geometrically ruled surface. The geometrically ruled surface Y over B is isomorphic to the projective bundle $\mathbb{P}(\mathcal{E})$ where \mathcal{E} is a rank two vector bundle (not unique) over B such that $\text{H}^0(B, \mathcal{E}) \neq \{0\}$ and $\text{H}^0(B, \mathcal{E} \otimes \mathcal{L}) = \{0\}$ for every line bundle \mathcal{L} with $\text{deg}(\mathcal{L}) < 0$. The opposite of the degree of $\wedge^2 \mathcal{E}$, denoted by e , is an invariant of Y ; the number $-e$ is the

minimal self-intersection number of a section of $Y \rightarrow B$. A section with self-intersection number $-e$ is called a minimal section; the line sub-bundle of \mathcal{E} which corresponds to the minimal section is called a maximal line bundle. The geometrically ruled surface Y is called *decomposable* if there exists a decomposition $\mathcal{E} = \mathcal{L}_1 \oplus \mathcal{L}_2$ where $\mathcal{L}_1, \mathcal{L}_2$ are line bundles, otherwise Y is called indecomposable. There exist two sections which do not intersect each other if and only if Y is decomposable.

The group of automorphisms of Y which preserve fiberwise the ruling is denoted by $\text{Aut}_B(Y)$.

Theorem 7.4.1 (Maruyama [Mar71]). *(1) If $e < 0$, then $\text{Aut}_B(Y)$ is a finite group.*

(2) If $e \geq 0$ and if Y is indecomposable, then $\text{Aut}_B(Y) \cong \mathbf{C}^k$ where $k = \dim H^0(B, (\det \mathcal{E})^{-1} \otimes \mathcal{L}^2)$ and $\mathcal{L} \subset \mathcal{E}$ is the unique maximal line bundle.

(3) If $Y \cong \mathbb{P}^1 \times B$, then $\text{Aut}_B(Y) \cong \text{PGL}(2, \mathbf{C})$.

(4) If $Y \cong \mathbb{P}(\mathcal{E})$ with $\mathcal{E} = \mathcal{L} \oplus (\mathcal{L} \otimes \mathcal{N})$, $\mathcal{N}^2 = \mathcal{O}_B$ and $\mathcal{N} \neq \mathcal{O}_B$, then $\text{Aut}_B(Y)$ is an extension of \mathbf{C}^ by $\mathbf{Z}/2\mathbf{Z}$.*

(5) If Y is decomposable and does not fit in the previous two cases, then $\text{Aut}_B(Y)$ is isomorphic to the following subgroup of $\text{GL}_{k+1}(\mathbf{C})$:

$$H_k = \left\{ \begin{pmatrix} \alpha & t_1 & \cdots & \cdots & t_k \\ 0 & 1 & 0 & \cdots & 0 \\ \vdots & & \ddots & & \vdots \\ \vdots & & & \ddots & \vdots \\ 0 & \cdots & 0 & & 1 \end{pmatrix} \in \text{GL}(k+1, \mathbf{C}) \right\}$$

where $k = \dim H^0(B, (\det \mathcal{E})^{-1} \otimes \mathcal{L}^2)$ and \mathcal{L} is a maximal line bundle.

In the second and last case, the action of an element of H_k on the fiber F_b over $b \in B$ is basically $x \mapsto \alpha x + t_1 l_1(b) + \cdots + t_k l_k(b)$ where l_1, \dots, l_k form a base of $H^0(B, (\det \mathcal{E})^{-1} \otimes \mathcal{L}^2)$ (see [Mar71] for the exact meaning of the formula).

We deduce immediately from Maruyama's theorem the following:

Corollary 7.4.2. *If $\text{Aut}_B(Y)$ is infinite and not solvable, then $Y = B \times \mathbb{P}^1$.*

7.5. Proof of Theorem 7.2.1. We turn back to the proof of Theorem 7.2.1 without assuming that Y is geometrically ruled. We still assume that $\Gamma_B = \{\text{Id}\}$ and that any component of any singular fiber of r is fixed by Γ . By Proposition 7.3.2 we know that if Y is not geometrically ruled then the intersection of U with a regular fiber is biholomorphic to \mathbf{C}^* .

7.5.1. Non-abelian case.

Proposition 7.5.1. *Suppose that Γ is not solvable. Then $Y = \mathbb{P}^1 \times B$ and $U = W \times B$ where $W \subset \mathbb{P}^1$ is an invariant component of Γ , viewed as a non-elementary classical Kleinian group. The quotient surface X is the product of B with a hyperbolic curve.*

Proof. In particular Γ is not cyclic and Proposition 7.3.2 implies that Y is geometrically ruled. By Corollary 7.4.2 $Y = \mathbb{P}^1 \times B$. In this case $\text{Aut}_B(Y) = \text{PGL}_2 \mathbf{C}$ and the intersection of U with each fiber is a union of components of some classical Kleinian group. This classical Kleinian group does not depend on the fiber. Since U is connected, the intersection of U with each fiber is the same component of the discontinuity set. \square

7.5.2. *Translation in the fibers.* As Γ is isomorphic to a classical Kleinian group by Proposition 7.3.2, if it is solvable then it is isomorphic to \mathbf{Z}^2 or \mathbf{Z} , depending on whether the intersection of U with a fiber is \mathbf{C} or \mathbf{C}^* .

We first investigate the case where $\Gamma = \mathbf{Z}^2$; in this case Γ acts on each fiber by translations (translation means parabolic automorphism of \mathbb{P}^1). In case 2 (resp. case 5) of Maruyama's theorem, the action is given, in local coordinate z of the fiber, by

$$(3) \quad z \mapsto z + t_1 e_1 + \cdots + t_k e_k \quad (\text{resp. } z \mapsto \alpha z + t_1 e_1 + \cdots + t_k e_k)$$

where e_1, \dots, e_k extend to a global base of $H^0(B, (\det \mathcal{E})^{-1} \otimes \mathcal{L}^2)$. Since a line bundle has a nowhere vanishing section if and only if it is trivial, an automorphism can act by non-trivial translation in every fiber only if the line bundle $(\det \mathcal{E})^{-1} \otimes \mathcal{L}^2$ is trivial. Writing $0 \rightarrow \mathcal{L} \rightarrow \mathcal{E} \rightarrow \mathcal{L}' \rightarrow 0$, we have $(\det \mathcal{E})^{-1} \otimes \mathcal{L}^2 = \mathcal{L} \otimes (\mathcal{L}')^{-1}$. Therefore the triviality of $(\det \mathcal{E})^{-1} \otimes \mathcal{L}^2$ implies that $\mathcal{L}^{-1} \otimes \mathcal{E}$ is an extension of two trivial line bundles. Thus we obtain:

Proposition 7.5.2. *Suppose that the intersection of U with a fiber is \mathbf{C} . There is a rank two vector bundle \mathcal{E} on B such that*

- (1) *Y is isomorphic to $\mathbb{P}(\mathcal{E})$;*
- (2) *We can write \mathcal{E} as an extension $0 \rightarrow \mathcal{O}_B \rightarrow \mathcal{E} \rightarrow \mathcal{O}_B \rightarrow 0$ such that $U = Y - s(B)$ where $s : B \rightarrow Y$ is the section determined by this extension, i.e. $s \mapsto s(b)$ with $s(b) = \mathbb{P}(\mathcal{O}_b) \in \mathbb{P}(\mathcal{E}_b)$.*

As $(\det \mathcal{E})^{-1} \otimes \mathcal{L}^2 = \mathcal{O}_B$, the subgroup of $\text{Aut}_B(Y)$ fixing $s(B)$ is isomorphic to \mathbf{C} by Theorem 7.4.1. We can identify Γ as a lattice in this subgroup. The surface X is an elliptic fiber bundle over B with fibers isomorphic to \mathbf{C}/Γ .

The action of \mathbf{C} on U descends to an action of \mathbf{C}/Γ on X ; this implies that X is a principal elliptic bundle by Lemma V.5.1 of [BHPVdV04]. A principal elliptic bundle is topologically a product of the circle S^1 with an S^1 -bundle (Proposition V.5.2 of [BHPVdV04]). We claim that in our case the corresponding S^1 -bundle is a product. The reason is that the rank two vector bundle \mathcal{E} is topologically trivial because an extension of a trivial bundle by a trivial bundle always splits in the topological category. If $B = \mathbb{P}^1$ and X is Kähler, then the only possibility is $X = (\mathbf{C}/\Gamma) \times \mathbb{P}^1$; if B is an elliptic curve and X is Kähler, then X is a complex torus or a hyperelliptic surface (see [BHPVdV04] Chapter V).


7.5.3. *Multiplication in the fibers.* We now investigate the case where $\Gamma = \mathbf{Z}$. Let us see first that it is indeed possible that Y is not geometrically ruled (cf. Proposition 7.3.2).

Given a decomposable geometrically ruled surface $Y_0 = \mathbb{P}(\mathcal{E})$ over B , up to tensoring \mathcal{E} with a line bundle, we can suppose that it decomposes as $\mathcal{E} = \mathcal{O}_B \oplus \mathcal{L}$. The decomposition determines two sections S_1, S_2 on Y_0 without intersection. Pick a finite family of fibers F_1, \dots, F_n over $b_1, \dots, b_n \in B$. Choose a positive integer $m \leq n$. Denote by p_1, \dots, p_m the intersection points of F_1, \dots, F_m with S_1 and p_{m+1}, \dots, p_n the intersection points of F_{m+1}, \dots, F_n with S_2 . Construct a birationally ruled surface Y over B as follows: Y is obtained from Y_0 by successive blow-ups at p_1, \dots, p_n and some infinitely near points. For $i \leq m$ (resp. $i > m$) the successive blow-ups on the fiber over p_i are executed either at the intersection point of two irreducible components of the fiber or at the intersection point of the fiber with (the strict transform of) S_1 (resp. S_2). We denote by $\pi : Y \rightarrow Y_0$ the contraction map. The singular fibers of $Y \xrightarrow{\phi} B$ are chains of smooth rational curves, i.e.

the singular fiber corresponding to F_i is

$$\tilde{F}_i = \sum_{k=0}^{l(i)} d_{i,k} F_{i,k};$$

here $F_{i,0}$ is the strict transform of F_i and the $F_{i,k}$ are all smooth rational curves among which the only non vanishing intersection pairings are $F_{i,k} \cdot F_{i,k+1} = 1$. We have $d_{i,0} = 1$ and $d_{i,1}, \dots, d_{i,l(i)} \in \mathbf{N}^+$. Each $F_{i,k}$ has two distinguished points $\{0_{i,k}, \infty_{i,k}\}$ which are the two intersection points of $F_{i,k}$ with other components of the fiber or with (strict transforms of) the two sections S_1, S_2 . Let's say that the one which points to S_1 is $0_{i,k}$.



By Theorem 7.4.1, the subgroup of $\text{Aut}_B(Y_0)$ fixing S_1, S_2 is isomorphic to \mathbf{C}^* ; it acts by multiplication in the fibers. The \mathbf{C}^* -action lifts to an action on Y by automorphisms because the blow-ups which we did to obtain Y are all at the fixed points of the \mathbf{C}^* -action. An element $\lambda \in \mathbf{C}^*$ acts on a component $F_{i,k}$ of the singular fiber \tilde{F}_i by multiplication by $\lambda^{d_{i,k}}$ with fixed points $\{0_{i,k}, \infty_{i,k}\}$.

Now for each \tilde{F}_i , choose a component $F_{i,k_0(i)}$ and consider the open set

$$U = Y \setminus \left(S_1 \cup S_2 \cup \left(\bigcup_i \bigcup_{k \neq k_0(i)} F_{i,k} \right) \right).$$

We remark that U is Zariski open in Y . The ruling ϕ restricted to U is a surjective holomorphic map onto B with fibers biholomorphic to \mathbf{C}^* . The quotient of U by $\lambda \in \mathbf{C}^*, |\lambda| \neq 1$ is a compact complex surface X which admits an elliptic fibration over B . If a fiber of $X \rightarrow B$ comes from a non-singular fiber of $\phi : Y \rightarrow B$, i.e. the fiber over a point that is not one of the p_1, \dots, p_n , then it is isomorphic to the elliptic curve $\mathbf{C}^* / \langle \lambda \rangle$. If a fiber of $X \rightarrow B$ comes from F_i , then it is the quotient of $\mathbf{C}^* = F_{i,k_0(i)} \setminus \{0_{i,k_0(i)}, \infty_{i,k_0(i)}\}$ by $\langle \lambda^{d_{i,k_0(i)}} \rangle$. Hence the only singular fibers of the elliptic fibration $X \rightarrow B$ are multiples of smooth elliptic curves, the multiplicities being the $d_{i,k_0(i)}$.

Actually the above construction exhausts all possibilities:

Proposition 7.5.3. *If the intersection of U with a fiber is \mathbf{C}^* , i.e. if $\Gamma = \mathbf{Z}$ then X is obtained from the above process. In particular the surface X is an elliptic fibration over B whose singular fibers are all multiples of smooth elliptic curves.*

Proof. Recall that by Lemma 7.3.1 $\Gamma \subset \text{Aut}(Y)$. We have also assumed that Γ preserves any component of any fiber. Therefore the action of Γ descends to an action by automorphisms, after contracting some exceptional curves in a fiber. Continuing the contraction until a geometrically ruled surface, we infer that the action of Γ on Y comes from a \mathbf{C}^* -action on a geometrically ruled surface Y_0 ; and Y is obtained from Y_0 by blowing-up fixed points (or infinitely near fixed points) of the \mathbf{C}^* -action. This is exactly how the above process works. \square

We denote by X_0 the quotient of $Y_0 \setminus (S_1 \cup S_2)$ by $\langle \lambda \rangle$. It is a principal elliptic fiber bundle over B because the \mathbf{C}^* -action descends to X_0 (Proposition V.5.2 of [BHPVdV04]). The surface X is obtained from X_0 by logarithmic transformations which replace the fibers over $b_1, \dots, b_n \in B$ with $d_{1,k_0(1)}E_1, \dots, d_{n,k_0(n)}E_n$ where $E_i = \mathbf{C}^* / \langle \lambda^{d_{i,k_0(i)}} \rangle$ (see [Kod64] for logarithmic transformations).

8. INVARIANT RATIONAL FIBRATION II: ELLIPTIC BASE

The base curve B is assumed to be an elliptic curve in this section. We prove the following statement which encapsulates both the case where Γ_B is finite and the case where it is infinite:

Theorem 8.0.1. *Let (Y, Γ, U, X) be a birational Kleinian group on a surface Y which is ruled over an elliptic curve B . Then up to geometric conjugation and up to taking finite index subgroup, we are in one of the following situations:*

- (1) $Y = \mathbb{P}^1 \times B$, $U = D \times B$ where $D \subset \mathbb{P}^1$ is an invariant component of the domain of discontinuity of a classical Kleinian group. X is a fiber bundle over B or a suspension of B .
- (2) $Y = R_0$, $U = Y - s$ and U has a structure of algebraic group which is isomorphic to $\text{Aut}^0(Y)$. As a complex analytic Lie group U is isomorphic to $\mathbf{C}^* \times \mathbf{C}^*$ and the group Γ is isomorphic to a lattice in U . The quotient X is a complex torus.
- (3) Y is decomposable with $e = 0$, $U = Y - (s_1 \cup s_2)$ and U has a structure of algebraic group which is isomorphic to $\text{Aut}^0(Y)$. The quotient X is a complex torus.
- (4) As in Case 3) of Theorem 7.2.1.

In any case of the above theorem, the open set U is not simply connected. Except in the first case, U is a Zariski open set.

In the rest of this section (Y, Γ, U, X) will be a birational Kleinian group satisfying the hypothesis of the theorem.

Lemma 8.0.2. *Suppose that Γ_B is an infinite group. Up to geometric conjugation of birational Kleinian groups, Y is geometrically ruled, $r(U) = B$ and $\Gamma \subset \text{Aut}(Y)$.*

Proof. The open set $r(U)$ is invariant under the infinite group Γ_B . Suppose by contradiction that $r(U) \neq B$. Then according to Lemma 2.2.1, it is a connected component of the complement of a finite union of real subtori, thus a band. The action of Γ_B on the band $r(U)$ is by translations along the direction of the band. Such an action on a band is never cocompact, i.e. the union of the images under Γ of a compact subset of the band never covers the band. This contradicts the cocompactness of the action of Γ on U .

By Corollary 7.1.3, the fact $r(U) = B$ implies that $\Gamma \subset \text{Aut}(Y)$. Since Γ_B is infinite and B is an elliptic curve, every point of B has an infinite Γ_B -orbit. If there existed a singular fiber of r , then its translations by elements of Γ would give rise to an infinite number of singular fibers. Therefore Y is geometrically ruled. \square

The ruling $r : Y \rightarrow B$ is identified with the Albanese morphism and is preserved by each element of Γ . We have an exact sequence

$$0 \rightarrow \Gamma_r \rightarrow \Gamma \rightarrow \Gamma_B \rightarrow 0$$

where Γ_r is the subgroup preserving fiberwise the fibration.

Proposition 8.0.3. *Suppose that Γ_B is infinite. Then X is a complex torus. The foliation \mathcal{F} is a linear irrational foliation or a suspension of B . In the two cases we have $r(U) = B$.*

Proof. We apply Theorem 6.1.1 to \mathcal{F} . The orbit of a point on B under Γ_B is not discrete and its closure contains a real subtorus (cf. Lemma 2.2.1). This implies that no leaf of \mathcal{F} is a submanifold of X . Thus the foliation is neither a fibration, nor a turbulent foliation, nor an obvious foliation on a Hopf surface. Here the base curve B is elliptic so that \mathcal{F} is transversely Euclidean. This rules out obvious foliations on Inoue surfaces (cf. [Zhae]). The foliation \mathcal{F} cannot be a suspension of \mathbb{P}^1 because the only \mathbb{P}^1 in Y are the fibers of the rational fibration. There are only two possibilities left: either X is a complex torus with an irrational linear foliation or X is an infinite suspension of a hyperbolic curve over an elliptic curve. \square

We continue to use the notations of section 7.4. It is well known that, up to isomorphisms, there are only two indecomposable geometrically ruled surfaces over an elliptic curve B : R_0 with $e = 0$ and R_1 with $e = 1$ (see [Har77] Chapter 5). Now we describe the whole automorphism group of the geometrically ruled surface Y according to Maruyama [Mar71]:

- (1) If $Y = \mathbb{P}^1 \times B$, then $\text{Aut}(Y) = \text{PGL}(2, \mathbf{C}) \times \text{Aut}(B)$.
- (2) If Y is decomposable and $e \neq 0$, then we have an exact sequence of algebraic groups

$$0 \rightarrow \text{Aut}_B(Y) \rightarrow \text{Aut}(Y) \rightarrow G \rightarrow 0$$

where G is a finite subgroup of $\text{Aut}(B)$.

- (3) If Y is decomposable with $e = 0$ and if $Y \neq \mathbb{P}^1 \times B$, then we have an exact sequence of algebraic groups

$$0 \rightarrow \text{Aut}_B(Y) \rightarrow \text{Aut}(Y) \rightarrow G \rightarrow 0$$

where $\text{Aut}_B(Y)$ is isomorphic to \mathbf{C}^* or an extension of \mathbf{C}^* by $\mathbf{Z}/2\mathbf{Z}$, and G is a subgroup of $\text{Aut}(B)$ containing $\text{Aut}^0(B)$. If s_1, s_2 are the minimal sections, then $Y - (s_1 \cup s_2)$ is a principal \mathbf{C}^* -bundle over B which has a structure of commutative algebraic group. The algebraic group $Y - (s_1 \cup s_2)$ is isomorphic to the connected component of identity $\text{Aut}^0(Y)$.

- (4) If $Y = R_0$, then we have an exact sequence of algebraic groups

$$0 \rightarrow \mathbf{C} \rightarrow \text{Aut}(Y) \rightarrow \text{Aut}(B) \rightarrow 0.$$

If s is the minimal section, then $Y - s$ is a principal \mathbf{C} -bundle over B which has a structure of commutative algebraic group. The algebraic group $Y - s$ is isomorphic to the connected component of identity $\text{Aut}^0(Y)$.

- (5) If $Y = R_1$, then we have

$$0 \rightarrow \text{Aut}_B(Y) \rightarrow \text{Aut}(Y) \rightarrow \text{Aut}(B) \rightarrow 0$$

where $\text{Aut}_B(Y)$ is isomorphic to $\mathbf{Z}/2\mathbf{Z} \times \mathbf{Z}/2\mathbf{Z}$.

Proof of Theorem 8.0.1. We already treated the case where Γ_B is finite in theorem 7.2.1. This is covered by case 1), 2), 4) of Theorem 8.0.1. From now on we assume that Γ_B is infinite. We analyse the situation case by case according to the above description of automorphism group. From the infiniteness of $\text{Aut}_B(Y)$ we infer that Y is not decomposable with $e \neq 0$. The surface Y is not R_1 because $\text{Aut}(R_1)$ is compact and does not contain infinite discrete subgroup.

Assume that Y is decomposable with $e = 0$. The two minimal sections s_1 and s_2 are disjoint from U because they are invariant under $\text{Aut}(Y)$. The universal cover of the complex analytic Lie group $Y - (s_1 \cup s_2)$ is \mathbf{C}^2 because it is commutative. Hence we obtain immediately that $U = Y - (s_1 \cup s_2)$ and X is a complex torus. If $Y = R_0$, then the same reasoning tells us that $U = Y - s$ and X is a complex torus.

Assume that $Y = \mathbb{P}^1 \times B$. By Proposition 8.0.3 the foliation on X induced by $Y \rightarrow B$ is a linear foliation on a torus or a suspension of B over a hyperbolic curve. If X is a torus then the intersection of U with any fiber of $Y \rightarrow B$ is biholomorphic to \mathbf{C} or \mathbf{C}^* ; the conclusion follows immediately in this case. Assume that X is a suspension over a hyperbolic curve. The projection $Y \rightarrow \mathbb{P}^1$ induces a second regular foliation on X which is transverse to the first one; this second foliation must be the elliptic fibration subjacent to the suspension. The conclusion follows. \square

9. INVARIANT RATIONAL FIBRATION III: RATIONAL BASE

In this section we work under the assumption that $B = \mathbb{P}^1$ and Γ_B is infinite. We will investigate the foliation \mathcal{F} case by case according to Theorem 6.1.1 and Corollary 6.2.2.

The situation where \mathcal{F} is an obvious foliation on an Inoue surface is treated in our previous paper [Zhac] where results stronger than what we need here are proved. We see from the description of Inoue surfaces in Section 3.2.5 that their standard realizations by complex affine Kleinian group fit in our setting of this section. The main result of [Zhac] implies:

Theorem 9.0.1 ([Zhac]). *Let S be an Inoue surface. Then up to geometric conjugation there is only one birational Kleinian group (Y, Γ, U, X) such that $X = S$. It corresponds to the standard construction of the Inoue surface by taking the quotient of $\mathbb{H} \times \mathbf{C}$ by a group of affine transformations.*

9.1. Turbulent foliations, Hopf surfaces, suspensions of \mathbb{P}^1 .

9.1.1. *Examples.* The Hirzebruch surface \mathcal{H}_n is the projectivization of the rank two vector bundle $\mathcal{O} \oplus \mathcal{O}(n)$ on \mathbb{P}^1 . The subbundles \mathcal{O} and $\mathcal{O}(n)$ determine two sections. The complement of these two sections, denoted by W_n , is isomorphic to the principal \mathbf{C}^* -bundle over \mathbb{P}^1 associated with the line bundle $\mathcal{O}(n)$. The quasi-projective variety W_0 is a product $\mathbf{C}^* \times \mathbb{P}^1$. The quasi-projective variety W_1 is isomorphic to $\mathbf{C}^2 \setminus \{0\}$. For $n \geq 1$, W_1 is an unramified cover of degree n of W_n ; the cover is given in each fiber by $w \mapsto w^n$.

The automorphism group of \mathcal{H}_n is connected and fits in the following exact sequence (cf. [Mar71]):

$$1 \rightarrow H_{n+1} \rightarrow \text{Aut}(\mathcal{H}_n) \rightarrow \text{Aut}(\mathbb{P}^1) \rightarrow 1$$

where H_{n+1} is the group defined in Theorem 7.4.1. Using coordinates we have

$$\text{Aut}(\mathcal{H}_n) = \left\{ (x, y) \mapsto \left(\frac{ax+b}{cx+d}, \frac{y+t_0+t_1x+\dots+t_nx^n}{(cx+d)^n} \right) \mid \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}(2, \mathbf{C}), t_0, \dots, t_n \in \mathbf{C} \right\}.$$

The action around the fiber at infinity is obtained by the change of coordinates $(x, y) \mapsto (\frac{1}{x}, \frac{y}{x^n})$. In particular for the automorphism $(x, y) \mapsto (ax, by)$ with $a, b \in \mathbf{C}^*$, the action on

the fiber at infinity is multiplication by $\frac{b}{a^n}$; for the automorphism $(x, y) \mapsto (x + c, by)$ with $a, b \in \mathbf{C}^*$, the action on the fiber at infinity is multiplication by b .

Example 9.1.1. Take $a, b \in \mathbf{C}^*$ such that $0 < |b| < 1$ and $|b| < |a|^n$. The automorphism $f_{n,a,b} : (x, y) \mapsto (ax, by)$ of \mathcal{H}_n preserves W_n and acts freely properly discontinuously and cocompactly on W_n . The action of $f_{1,a,b}$ on W_1 is nothing else but the linear contraction $(z, w) \mapsto (bz, \frac{b}{a}w)$ on $\mathbf{C}^2 \setminus \{0\}$. Let X be the quotient surface $= W_n / \langle f_{n,a,b} \rangle$ and \mathcal{F} be the foliation on X induced by the \mathbf{C}^* -bundle structure of W_n .

Assume that a is not a root of unity. When $n = 0$, the quotient is a ruled surface over the elliptic curve $\mathbf{C}^* / \langle b \rangle$; the foliation \mathcal{F} is a suspension of \mathbb{P}^1 . When $n > 0$ the surface X is a Hopf surface with two elliptic curves isomorphic to $\mathbf{C}^* / \langle b \rangle$ and $\mathbf{C}^* / \langle \frac{b}{a^n} \rangle$; the leaves of \mathcal{F} come from vector lines on the universal cover $\mathbf{C}^2 \setminus \{0\}$. No matter what n is, a and b are multiplicatively dependent if and only if X is equipped with an elliptic fibration; in this case \mathcal{F} is also a turbulent foliation with respect to that elliptic fibration.

We can modify the above examples by adapting the construction in Section 7.5.3. We blow up \mathcal{H}_n at points in the fibers over $x = 0$ or $x = \infty$ to obtain a new surface Y ; the points we blow up are intersection points of irreducible components with other irreducible components or with the strict transforms of the two sections of \mathcal{H}_n . The points we blowed up are fixed points of the $f_{n,a,b}$. Thus $f_{n,a,b}$ acts also by automorphism on Y ; its action on an irreducible component of a fiber is by multiplication. Let $U \subset Y$ be a \mathbf{C}^* -fibration over \mathbb{P}^1 which is the complement in Y of the union of strict transforms of the two sections and some components of the singular fibers, as in Section 7.5.3. Then for suitable choice of a, b , the action of $f_{n,a,b}$ on U is free, properly discontinuous and cocompact. The quotient is either a Hopf surface or a ruled surface over an elliptic curve. If the quotient is a ruled surface then U is foliated by rational curves of self-intersection 0; this forces U to be isomorphic to the product $\mathbb{P}^1 \times \mathbf{C}^*$. Hence the quotient is a ruled surface if and only if the birational Kleinian group is geometrically conjugate to $\langle f_{0,a,b} \rangle$ on $\mathcal{H}_0 = \mathbb{P}^1 \times \mathbb{P}^1$.

Example 9.1.2. Take $0 < |b| < 1$ and $c \in \mathbf{C}^*$. The automorphism $g_{n,b,c} : (x, y) \mapsto (x + c, by)$ of \mathcal{H}_n preserves W_n ; its action on the fiber at infinity is multiplication by b . If $n = 0$ then the quotient of W_0 by $g_{0,b,c}$ is a ruled surface over an elliptic curve equipped with a suspension foliation. The action of $g_{1,b,c}$ on W_1 is nothing else but the contraction $(z, w) \mapsto (bz + bcw, bw)$ on $\mathbf{C}^2 \setminus \{0\}$. For $n \geq 1$ the quotient of W_n by $g_{n,b,c}$ is a Hopf surface with only one elliptic curve. It has an unramified cover which is a primary Hopf surface corresponding to the normal form $(z, w) \mapsto (bz + \gamma w, bw)$ where γ is some number depending on n, b, c . Remark that this construction does not give all primary Hopf surfaces with only one elliptic curve, but only those with linear normal form. The foliation induced by the rational fibration corresponds to the foliation by lines in $\mathbf{C}^2 \setminus \{0\}$. As in Example 9.1.1 we can modify the above construction by blowing up the invariant fiber.

Example 9.1.3. Consider two automorphisms $f_i : (x, y) \mapsto (x + a_i, y + b_i)$, $i = 1, 2$ of $\mathbb{P}^1 \times \mathbb{P}^1$. Suppose that $\mathbf{Z}b_1 + \mathbf{Z}b_2$ is a lattice in \mathbf{C} . Then f_1 and f_2 preserve $U = \mathbb{P}^1 \times \mathbf{C} \subset \mathbb{P}^1 \times \mathbb{P}^1$; the data $(\mathbb{P}^1 \times \mathbb{P}^1, \langle f_1, f_2 \rangle, U)$ give a birational Kleinian group. The quotient $X = U / \langle f_1, f_2 \rangle$ is a ruled surface over the elliptic curve $\mathbf{C} / (\mathbf{Z}b_1 + \mathbf{Z}b_2)$; the foliation \mathcal{F} is a suspension. If moreover $\mathbf{Z}a_1 + \mathbf{Z}a_2$ is a lattice in \mathbf{C} such that the elliptic curve $\mathbf{C} / (\mathbf{Z}a_1 + \mathbf{Z}a_2)$ is isogeneous to $\mathbf{C} / (\mathbf{Z}b_1 + \mathbf{Z}b_2)$, then X is equipped with an elliptic fibration so that the foliation is turbulent with one compact leaf.

Consider two automorphisms g_1, g_2 of $\mathbb{P}^1 \times \mathbb{P}^1$ defined by $g_1 : (x, y) \mapsto (x, y + b_1)$ and $g_2 : (x, y) \mapsto (ax, y + b_2)$. Suppose that $\mathbf{Z}b_1 + \mathbf{Z}b_2$ is a lattice in \mathbf{C} . Then g_1 and g_2 preserve $U = \mathbb{P}^1 \times \mathbf{C} \subset \mathbb{P}^1 \times \mathbb{P}^1$; the data $(\mathbb{P}^1 \times \mathbb{P}^1, \langle g_1, g_2 \rangle, U)$ give a birational Kleinian group.

The quotient $X = U / \langle g_1, g_2 \rangle$ is a ruled surface over the elliptic curve $\mathbf{C}/(\mathbf{Z}b_1 + \mathbf{Z}b_2)$; the foliation \mathcal{F} is a suspension. If moreover $\langle a \rangle$ is a lattice in \mathbf{C}^* such that the elliptic curve $\mathbf{C}^*/\langle a \rangle$ is isogeneous to $\mathbf{C}/(\mathbf{Z}b_1 + \mathbf{Z}b_2)$, then X is equipped with an elliptic fibration so that the foliation is turbulent with two compact leaves.

Remark 9.1.4. There are no analogues of Example 9.1.3 for \mathcal{H}_n with $n \geq 1$. For $n \geq 1$ the action of $(x, y) \mapsto (x + a, y + b)$ on the fiber at infinity is trivial and $(x, y) \mapsto (ax, y + b)$ acts as multiplication by $\frac{1}{a^n}$ on the fiber at infinity.

We can see in another way that there are no such analogues when $n \geq 1$. Let U_n be the complement of the minimal section in \mathcal{H}_n ; it is a fibration over \mathbb{P}^1 with fibers biholomorphic to \mathbf{C} . If the quotient of U_n by a \mathbf{Z}^2 -action was a compact surface then this compact surface would be Kähler because it has even second Betti number (cf. Theorem IV.3.1 [BHPVdV04]). However U_n is not the universal cover of any compact Kähler surface: if the universal cover of a compact Kähler surface is non-compact and quasi-projective then it is \mathbf{C}^2 or $\mathbf{C} \times \mathbb{P}^1$ (cf. [Cla], [CHK13]).

9.1.2. Characterizations.

Proposition 9.1.5. *Suppose that $r(U) = B = \mathbb{P}^1$. Then \mathcal{F} is a turbulent foliation, an obvious foliation on a Hopf surface or a suspension of \mathbb{P}^1 .*

Proof. Assume that \mathcal{F} is a linear foliation on a torus. The leaves of a linear foliation on \mathbf{C}^2 are fibers of a linear map $h : \mathbf{C}^2 \rightarrow \mathbf{C}$. The group $\pi_1(X)$ acts on the target \mathbf{C} by translations so that h is equivariant. The image of $\pi_1(X)$ in $\text{Aut}(\mathbf{C})$ is a free abelian group of rank 2, 3 or 4, depending on whether the leaves of \mathcal{F} are elliptic curves, \mathbf{C}^* or \mathbf{C} . The open set $r(U)$ is the set of leaves of U . We obtain thus a surjective holomorphic map $\mathbf{C} \rightarrow r(U)$ that is equivariant with respect to the homomorphism $\pi_1(X) \rightarrow \Gamma_B$. This forces $r(U)$ to be \mathbf{C} or \mathbf{C}^* . Contradiction.

Assume that X is a bidisk quotient and that \mathcal{F} is one of the natural foliations. Consider the composition $\pi_1(X) \rightarrow \Gamma \rightarrow \Gamma_B \subset \text{PGL}(2, \mathbf{C})$. Since $\pi_1(X)$ is an irreducible lattice in $\text{PSL}(2, \mathbf{R}) \times \text{PSL}(2, \mathbf{R})$, Margulis superrigidity ([Mar91]) says that Γ_B is up to conjugation in $\text{PGL}(2, \mathbf{C})$ a dense subgroup of $\text{PSL}(2, \mathbf{R})$ and that the quotient homomorphism $\pi_1(X) \rightarrow \Gamma_B$ is injective. This means that U is biholomorphic to the bidisk, the universal cover of X . Therefore $r(U) \neq B$.

\mathcal{F} is not a fibration by Lemma 9.3.1 that we prove later. Inoue surfaces are excluded by Theorem 9.0.1.

By Corollary 7.1.3 and Proposition 7.1.5 \mathcal{F} has a foliated projective structure. Combining Theorem 6.1.1 with Corollary 6.2.2 we see that the remaining possibilities are those listed in the proposition. \square

Proposition 9.1.6. *Suppose that \mathcal{F} is a turbulent foliation. Then up to a geometric conjugation preserving the rational fibration and up to replacing Γ with a finite index subgroup, we are in one of the cases described in Examples 9.1.1 and 9.1.3.*

Proof. Let $f : X \rightarrow C$ be the genus one fibration subjacent to the turbulent foliation. At least one fiber of f is a leaf of \mathcal{F} ; let E be such a fiber. Denote by π the covering map from U to X . We have an exact sequence $1 \rightarrow H \rightarrow \pi_1(X) \rightarrow \pi_1^{orb}(C) \rightarrow 1$ where H is the image of the fundamental group of a regular fiber (see [GS85] Theorem 1). If E_m is a multiple fiber of f then the image of $\pi_1(E_m)$ in $\pi_1(X)$ contains H as a subgroup of finite index. Let $G \subset \Gamma$ be the image of the composition $H \rightarrow \pi_1(X) \rightarrow \Gamma$; it is cyclic or isomorphic to \mathbf{Z}^2 . Then G is a normal subgroup of Γ ; it is the image of $\pi_1(E)$ in Γ if E is non-multiple and is a finite index subgroup of it if E is multiple. For any fiber E' of f , for any connected

component Ω of $\pi^{-1}(E')$, G preserves Ω and Ω/G is an elliptic curve which is E' if E' is non multiple. Thus the intermediate cover U/G is a genus one fibration over a possibly non-compact base.

Denote by G_B the image of G in Γ_B . We claim that G_B is infinite. Take a regular fiber E' of f which is transverse to \mathcal{F} . Let $\Omega \subset U$ be a connected component of $\pi^{-1}(E')$. Ω is everywhere transverse to the rational fibration r and r restricted to Ω is a covering map. Therefore $r(\Omega)$ is an open subset biholomorphic to \mathbf{C}^* or \mathbf{C} in $B = \mathbb{P}^1$. The claim follows because $\Omega/G = E'$ and $r|_{\Omega}$ is equivariant under the homomorphism $G \rightarrow G_B$.

We claim that G is a subgroup of Γ of finite index. Let $\gamma \in \Gamma$ be an element of infinite order. Let F_E be a connected component of $\pi^{-1}(E)$; it is a copy of \mathbf{C} or \mathbf{C}^* contained in a G -invariant fiber F of r . The image $\gamma(F_E)$ is also a connected component of $\pi^{-1}(E)$. It is preserved by G because G is a normal subgroup of Γ . This implies in particular $\gamma(F)$ is a G -invariant fiber. As G_B is infinite, there are at most two G -invariant fibers of r . Up to replacing Γ with an index two subgroup we can assume $\gamma(F) = F$. The possibly singular fiber F has only finitely many irreducible components; one of them contains F_E . Up to replacing γ by an iterate, we can assume that γ preserves F_E . As F_E/G is already an elliptic curve, Γ/G must be finite.

Up to replacing Γ with a finite index subgroup of G , we can and will assume that $G = \Gamma$ and that Γ, Γ_B are free abelian groups of rank one or two. We choose coordinates in $B = \mathbb{P}^1$ so that the fiber F containing E is over $\infty \in B$. Denote by R the set of fixed points of G_B in B . It contains ∞ and has at most two elements; we let the other element be 0 if it exists. The complement $r(U) \setminus R$ is \mathbf{C} or \mathbf{C}^* ; Γ_B acts respectively by translations or multiplications. Any point in $r(U) \setminus R$ has an infinite Γ_B -orbit. Thus by Corollary 7.1.3 r has no singular fibers over $r(U) \setminus R$. The fibers over points of R correspond to fibers of $f : X \rightarrow C$ that are leaves of \mathcal{F} . Denote by S the set of points in C over which the fibers of f are leaves of \mathcal{F} ; it has the same cardinality as R . The foliation \mathcal{F} restricted to $f^{-1}(C \setminus S)$ is everywhere transverse to the elliptic fibration f , thus is a suspension. We have

$$\pi^{-1}(f^{-1}(C \setminus S)) = r^{-1}(r(U) \setminus R) \cap U$$

and we denote this open set by M .

Let E' be a fiber of f over $C \setminus S$. Let Ω be its preimage in M ; it is biholomorphic to \mathbf{C} if Γ has rank two and to \mathbf{C}^* if Γ has rank one. It is everywhere transverse to the rational fibration r . It is invariant under Γ ; the quotient is E' . The projection $\Omega \rightarrow r(U) \setminus R$ induced by r is equivariant with respect to the homomorphism $\Gamma \rightarrow \Gamma_B$ and is an unramified covering. This implies that $(r(U) \setminus R)/\Gamma_B$ is an elliptic curve isogeneous to E' . We are in one of the following situations:

- (1) Γ and Γ_B are cyclic, $r(U) \setminus R$ is \mathbf{C}^* ;
- (2) Γ and Γ_B have rank two, $r(U) \setminus R$ is \mathbf{C} ;
- (3) Γ has rank two, Γ_B is cyclic, $r(U) \setminus R$ is \mathbf{C}^* .

Let us consider first the first two cases. In these two cases the covering map $\Omega \rightarrow r(U) \setminus R$ is finite. For any $x \in r(U) \setminus R$, $r^{-1}(x) \cap \Omega$ is a finite set. Thus a leaf of \mathcal{F} transverse to f intersects a fixed fiber of f only a finite number of times. Thus such a leaf is a finite unramified cover of $C \setminus S$. This implies that for any $x \in r(U) \setminus R$, $r^{-1}(x) \cap U$ is a finite unramified cover of $C \setminus S$, thus a Riemann surface of finite type. This is only possible if $C = \mathbb{P}^1$ because $r^{-1}(x) \cap U$ is an open subset of \mathbb{P}^1 . In the first case we obtain that $R = \{0, \infty\}$, $r(U) = B$ and $r^{-1}(b) \cap U$ is biholomorphic to \mathbf{C}^* . In the second case we obtain that $R = \{\infty\}$, $r(U) = B$ and $r^{-1}(b) \cap U$ is biholomorphic to \mathbf{C} . The fact $r(U) = B$ allows us to suppose that $\Gamma \subset \text{Aut}(Y)$ by Corollary 7.1.3.

Recall that Y has no singular fibers over $r(U) \setminus R$. The first case corresponds exactly to Example 9.1.1. In the second case U is a fibration over \mathbb{P}^1 with fibers biholomorphic to \mathbf{C} . It is necessarily a Zariski open set. Then Remark 9.1.4 allows us to conclude that we are in Example 9.1.3.

Finally let us consider the third case. In this case $r^{-1}(\infty) \cap U$ is isomorphic to \mathbf{C} and Γ acts by translations on it. Let Γ_r be the kernel of the projection $\Gamma \rightarrow \Gamma_B$; it preserves fiberwise r . The projection $\Omega \rightarrow r(U) \setminus R$ is an infinite cyclic covering, i.e. the exponential map from \mathbf{C} to \mathbf{C}^* . As in the previous case, using the fact the foliation on $\pi(M) = f^{-1}(C \setminus S)$ is a suspension of E' over $C \setminus S$, we deduce that for any $b \in r(U) \setminus R$, $r^{-1}(b) \cap U$ is an infinite cyclic cover of $C \setminus S$. The quotient of $r^{-1}(b) \cap U$ by Γ_r is a finite cover of $C \setminus S$, thus a quasi-projective curve. Therefore Γ_r can be viewed as a cyclic classical Kleinian group (not necessarily with compact quotient) in the fiber $r^{-1}(b)$. As S has cardinality 1 or 2, we have two possibilities: 1) the cyclic Kleinian group Γ_r is generated by a parabolic automorphism, $r^{-1}(b) \cap U$ is \mathbf{C} , $C = \mathbb{P}^1$ and $C \setminus S$ is \mathbf{C}^* ; 2) Γ_r is generated by a loxodromic automorphism, C is an elliptic curve, $r^{-1}(b) \cap U$ is the complement in \mathbf{C}^* of one or two Γ_r -orbits.

Suppose by contradiction that C is an elliptic curve. Let γ_1 be a generator of Γ_r ; its action on a fiber of r contained in M is a loxodromic automorphism. Let γ_2 be an element of Γ whose image in Γ_B generates Γ_B ; its action on B is a multiplication. Then $\Gamma = \mathbf{Z}^2$ is generated by γ_1 and γ_2 . By Theorem 4.4.3 (the information we use here is due to [CD12]), γ_1, γ_2 can be written respectively as $(x, y) \mapsto (x, ay)$ and $(x, y) \mapsto (bx, R(x)y)$ with $a, b \in \mathbf{C}^*$ and $R \in \mathbf{C}(x)^*$. Recall that we have a Γ -invariant fiber F over $x = \infty$ and an open subset F_E of F biholomorphic to \mathbf{C} such that Γ acts by translations on F_E . This gives a contradiction because γ_1 acts on any irreducible component of F by multiplication.

Therefore we obtain that $C = \mathbb{P}^1$, $C \setminus S = \mathbf{C}^*$ and $r^{-1}(b) \cap U = \mathbf{C}$ for any $b \notin R$. We have $R = \{0, \infty\}$ and $r(U) = B$. Then U is a fibration over \mathbb{P}^1 with fibers isomorphic to \mathbf{C} . Again $r(U) = B$ implies that $\Gamma \subset \text{Aut}(Y)$ by Corollary 7.1.3. This implies again by Remark 9.1.4 that Y is $\mathbb{P}^1 \times \mathbb{P}^1$. Hence we are in Example 9.1.3. \square

Proposition 9.1.7. *Suppose that \mathcal{F} is an obvious foliation on a non-elliptic Hopf surface. Then up to a geometric conjugation preserving the rational fibration and up to replacing Γ with a finite index subgroup we are in one of the following situations:*

- (1) X has two elliptic curves and we are in Example 9.1.1.
- (2) X has one elliptic curve and we are in Example 9.1.2.
- (3) U is the standard open set $\mathbf{C}^2 \setminus \{0\}$ of $Y = \mathbb{P}^1 \times \mathbb{P}^1$ and Γ is generated by the normal form $(x, y) \mapsto (\alpha x + \gamma y^m, \beta y)$.

Proof. Since the fundamental group of Hopf surface is virtually cyclic, the open set U is a finite quotient of $\mathbf{C}^2 \setminus \{0\}$. If \mathcal{F} comes from the foliation by parallel lines on $\mathbf{C}^2 \setminus \{0\}$, then the intersection of U with a fiber of r is \mathbf{C} except for one which is \mathbf{C}^* . Such an open set is necessarily Zariski open and isomorphic to $\mathbf{C}^2 \setminus \{0\}$. As the normal form is already birational and preserves a rational fibration, our situation is necessarily conjugate to it.

Assume now that \mathcal{F} is not a foliation by parallel lines. The space of leaves of the corresponding foliation on any infinite cover of X is \mathbb{P}^1 . Thus $r(U) = \mathbb{P}^1$. By Corollary 7.1.3 this implies $\Gamma \subset \text{Aut}(Y)$. The open set is necessarily one of the Zariski open sets appeared in Examples 9.1.1 and 9.1.2 because every leaf is biholomorphic to \mathbf{C}^* . Hence we are in Example 9.1.1 if Γ_B is generated by a multiplication and in Example 9.1.2 if Γ_B is generated by a translation. \square

Proposition 9.1.8. *Suppose that $r(U) = B$ and that \mathcal{F} is a suspension foliation. Then it is a suspension of \mathbb{P}^1 . Up to a geometric conjugation preserving the rational fibration and up to replacing Γ with a finite index subgroup, we have $Y = \mathbb{P}^1 \times \mathbb{P}^1$ and $\Gamma \subset \mathrm{PGL}(2, \mathbf{C}) \times \mathrm{PGL}(2, \mathbf{C})$. The projection of Γ onto the second factor is a classical Kleinian group; we have $U = \mathbb{P}^1 \times D$ where $D \subset \mathbb{P}^1$ is an invariant component of that classical Kleinian group.*

Proof. Again $r(U) = B$ implies that $\Gamma \subset \mathrm{Aut}(Y)$ by Corollary 7.1.3. The preimages of the leaves of \mathcal{F} in U are everywhere transverse to r . So they are unramified covers of $B = \mathbb{P}^1$, thus are isomorphic to \mathbb{P}^1 . In other words U is covered by rational curves of self-intersection 0 which are transverse to r . This implies that up to geometric conjugation $Y = \mathbb{P}^1 \times \mathbb{P}^1$; the first projection is r and the the rational curves in U are fibers of the second projection. The conclusion follows. \square

9.2. Suspensions. We denote the covering map $U \rightarrow X$ by π , and by Γ_r the kernel of the homomorphism $\Gamma \rightarrow \Gamma_B$. We now consider the case where there exists compact Riemann surfaces N, M such that the foliation \mathcal{F} on X is an infinite suspension of N over M . Either N is an elliptic curve or it is \mathbb{P}^1 . We have a fibration $f : X \rightarrow C$ whose fibers are all isomorphic to N . The fibration f is everywhere transverse to \mathcal{F} . It lifts to a foliation \mathcal{G} on U , everywhere transverse to the rational fibration r . By the transversality, we infer that, restricted to every leaf of \mathcal{G} , r is an unramified covering map onto $r(U)$. The covering map π restricted to a leaf of \mathcal{G} is a covering map onto a fiber of f , which is isomorphic to N . From this we deduce immediately:

Lemma 9.2.1. *If N is \mathbb{P}^1 then $r(U) = B$ and the situation is described by Proposition 9.1.8. If N is an elliptic curve then $r(U)$ is isomorphic to \mathbf{C} or \mathbf{C}^* .*

From now on we assume that N is an elliptic curve. Recall the construction of suspension. There exists an infinite covering \tilde{M} of M with deck transformation group G and a representation $\alpha : G \rightarrow \mathrm{Aut}(N)$ such that X is the quotient of $\tilde{M} \times N$ by the action of G defined by $g \cdot (m, n) = (g \cdot m, \alpha(g) \cdot n)$. Up to replacing \tilde{M} with a quotient, we can assume that α is injective. Up to replacing G with a subgroup of finite index, we can assume that it is an abelian group, i.e. $\alpha(G)$ is a group of translations on N . The fundamental group of $\tilde{M} \times N$ is $\pi_1(\tilde{M}) \times \pi_1(N)$ and the fundamental group of X fits in the exact sequence

$$1 \rightarrow \pi_1(\tilde{M}) \times \pi_1(N) \rightarrow \pi_1(X) \rightarrow G \rightarrow 1.$$

Denote by \tilde{M} the universal cover of M . The universal cover of X is $\tilde{M} \times \mathbf{C}$. Any element of $\pi_1(X)$ acts on $\tilde{M} \times \mathbf{C}$ in a diagonal way; the action on the \mathbf{C} factor is a translation. Thus $\pi_1(N)$ is in the center of $\pi_1(X)$. Our birational Kleinian group Γ is a quotient of $\pi_1(X)$. Denote respectively by Γ_1 and Γ_2 the images of $\pi_1(N)$ and $\pi_1(\tilde{M})$ in Γ ; they are normal subgroups of Γ . Up to replacing Γ by a subgroup of finite index, we can and will assume that Γ_1 is a free abelian group of rank one or two. The center of Γ contains Γ_1 . Note that we do not have necessarily $\Gamma_1 \cap \Gamma_2 = \{1\}$ because we are taking quotients. Denote by Γ_3 the quotient $\Gamma/(\Gamma_1\Gamma_2)$; it is isomorphic to a quotient of G . Denote by q the epimorphism $\Gamma \rightarrow \Gamma_3$.

Lemma 9.2.2. (1) Γ_1 preserves each leaf of the foliation \mathcal{G} . The image Γ_{1B} of Γ_1 in Γ_B is an infinite group of translations if $r(U) = \mathbf{C}$ or an infinite group of multiplications if $r(U) = \mathbf{C}^*$.

- (2) r has no singular fibers over $r(U)$ and $r^{-1}(r(U))$ is biholomorphic to $r(U) \times \mathbb{P}^1$.
(3) Let Ω be a connected component of the intersection of U with a fiber of r and let Γ_Ω be the subgroup of Γ that preserves Ω . Then $\Gamma_\Omega = \Gamma_2$ and $\Gamma_2 \subset \Gamma_r$.

Proof. The foliation \mathcal{G} on U is the pull-back of the elliptic fibration on X . The first assertion is a consequence of the following two observations that we made earlier: 1) r restricted to a leaf of \mathcal{G} is a covering map onto $r(U)$ and $r(U)$ is \mathbf{C}^* or \mathbf{C} ; 2) π restricted to a leaf of \mathcal{G} is a covering map onto a fiber of the elliptic fibration f .

By Corollary 7.1.3 Γ acts by holomorphic diffeomorphisms on $r^{-1}(r(U))$. If there was a singular fiber then there would be an infinite number of them by the first assertion. Hence the second assertion.

The image $\pi(\Omega) = \Omega/\Gamma_\Omega$ is a leaf of \mathcal{F} . It is biholomorphic to \bar{M} . Thinking of a fundamental group as a deck transformation group, it follows from the definition of Γ_2 that $\Gamma_\Omega = \Gamma_2$. As a consequence Γ_Ω is a normal subgroup of Γ and does not depend on Ω . This implies that Γ_Ω preserves r fiberwise, i.e. $\Gamma_\Omega \subset \Gamma_r$. \square

Lemma 9.2.3. Γ_B is not virtually cyclic.

Proof. Suppose by contradiction that Γ_B is virtually cyclic. We know already that Γ_{1B} is infinite. Thus Γ_{1B} is a subgroup of finite index of Γ_B . By the first assertion in Lemma 9.2.2 the action of Γ_B on $r(U)$ is discrete. Therefore $r(U)/\Gamma_B$ is an orbifold Riemann surface and we have a surjective holomorphic map from X to $r(U)/\Gamma_B$ whose fibers are leaves of \mathcal{F} . This contradicts that \mathcal{F} is an infinite suspension foliation. \square

Lemma 9.2.4. *Up to geometric conjugation realized by elementary transformations, we have $Y = \mathbb{P}^1 \times \mathbb{P}^1$ (where r is the projection onto the first factor) and Γ_1 is a subgroup of one of the following four groups: $\{(x, y) \mapsto (ax, by) | a, b \in \mathbf{C}^*\}$, $\{(x, y) \mapsto (ax, y + b) | a \in \mathbf{C}^*, b \in \mathbf{C}\}$, $\{(x, y) \mapsto (x + a, by) | a \in \mathbf{C}, b \in \mathbf{C}^*\}$ or $\{(x, y) \mapsto (x + a, y + b) | a, b \in \mathbf{C}\}$. In the first two cases $r(U) = \mathbf{C}^*$; in the last two cases $r(U) = \mathbf{C}$.*

Proof. Recall that Γ_1 is in the center of Γ . The image in $\text{Aut}(B)$ of the centralizer of a Jonquières twist is virtually cyclic by Theorem 4.4.2. Thus by Lemma 9.2.3 we conclude that Γ_1 contains no Jonquières twists and is an elliptic free abelian group. Then there exists a birational map $\phi : Y \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$, composition of elementary transformations, such that $\phi\Gamma_1\phi^{-1}$ is contained in one of the four groups in the statement (see Theorem 4.4.3). It remains to show that the conjugation ϕ can be chosen to be a geometric conjugation; the reason is as follows. Up to post-composition with an automorphism of $\mathbb{P}^1 \times \mathbb{P}^1$ we observe that the elementary transformations in ϕ can be made over any point in B . As $r(U)$ is a proper subset of B , we can do elementary transformations on a fiber over a point outside $r(U)$. Such an elementary transformation has no effects on U or on the dynamics of Γ on U ; in other words elementary transformations outside U realize a geometric conjugation of Γ . \square

Proposition 9.2.5. *Up to geometric conjugation, Γ is a subgroup of $\text{PGL}(2, \mathbf{C}) \times \text{PGL}(2, \mathbf{C})$ acting on $Y = \mathbb{P}^1 \times \mathbb{P}^1$. The open subset $U \subset Y$ is $\Omega_1 \times \Omega_2$ where Ω_2 is \mathbf{C} or \mathbf{C}^* and $\Omega_1 \subset \mathbb{P}^1$ is biholomorphic to \bar{M} .*

Proof. We see from Lemma 9.2.4 that Γ_1 acts separately on the x and y coordinates, the action on the x coordinate being Γ_{1B} . Assume first that its action on the y coordinate is finite. Since Γ_1 is a central abelian subgroup of Γ , we can replace Γ with a subgroup of finite index to assume that the action of Γ_1 on the y coordinate is trivial. Let L be a leaf of \mathcal{G} . Recall that L is biholomorphic to \mathbf{C} or \mathbf{C}^* , is transverse to the rational fibration $(x, y) \mapsto x$, is preserved by Γ_1 and L/Γ_1 is isomorphic to N . This forces L to be $r(U) \times \{y\}$ for some $y \in \mathbb{P}^1$. Hence the open set U is as described in the statement. We obtain also that the foliation \mathcal{G} is induced by the projection of $\mathbb{P}^1 \times \mathbb{P}^1 = Y$ onto the second factor. Therefore

Γ preserves this second rational fibration as well. This implies that Γ is a subgroup of $\mathrm{PGL}(2, \mathbf{C}) \times \mathrm{PGL}(2, \mathbf{C})$.

Now assume that the action of Γ_1 on the y coordinate has transformations of infinite order. Then Γ_1 contains elements with infinite actions on both x and y coordinates. Recall that Γ_1 is in the center of Γ . Thus by Theorem 4.4.1, Γ is a subgroup of $\mathrm{PGL}(2, \mathbf{C}) \times \mathrm{PGL}(2, \mathbf{C})$. Therefore the projection of $\mathbb{P}^1 \times \mathbb{P}^1 = Y$ onto the second factor induces a second foliation \mathcal{F}' on X . This foliation is transverse to \mathcal{F} and is not the elliptic fibration f because the sets $\{y = \text{constant}\}$ are not invariant under Γ_1 . Remark that X has Kodaira dimension ≥ 0 because it is a genus one bundle over M which is a compact Riemann surface with infinite fundamental group. Proposition 4.3 of [Bru97] tells us that a regular foliation on a genus one bundle of non-negative Kodaira dimension is either the genus one fibration or a (not necessarily infinite) suspension or a turbulent foliation. We said that \mathcal{F}' is not the original fibration. It is not a turbulent foliation by Proposition 9.1.6. Therefore \mathcal{F}' is another (possibly finite) suspension foliation. Consider the three foliations on X : \mathcal{F} , \mathcal{F}' and the fibration $f : X \rightarrow C$. They are transverse to each other. Thus these three foliations trivialize the projectivized tangent bundle of X . Consequently X has trivial first Chern class, and is a complex torus or a K3 surface. As there are no regular foliations on a K3 surface (cf. [Bru97] 5.Proposition 7), X is a complex torus. As a consequence, \mathcal{F} is a irrational linear foliation on X and its leaves are biholomorphic to \mathbf{C} or \mathbf{C}^* . This implies that the intersection of U with a fiber of r is \mathbf{C} or \mathbf{C}^* . Together with the fact $\Gamma \subset \mathrm{PGL}(2, \mathbf{C}) \times \mathrm{PGL}(2, \mathbf{C})$, this implies that U is either $\mathbf{C} \times \mathbf{C}$, or $\mathbf{C} \times \mathbf{C}^*$, or $\mathbf{C}^* \times \mathbf{C}$, or $\mathbf{C}^* \times \mathbf{C}^*$. Thus in the case where Γ_1 has infinite actions on the y coordinate, the conclusion also holds. \square

9.3. Fibrations. In this subsection we consider the case where the induced foliation \mathcal{F} on X is a fibration $f : X \rightarrow C$. We still denote the covering map $U \rightarrow X$ by π , and by Γ_r the kernel of the homomorphism $\Gamma \rightarrow \Gamma_B$. By Corollary 7.1.3 Γ acts by holomorphic diffeomorphisms on $r^{-1}(r(U))$. Up to replacing Γ with a finite index subgroup, we can assume that Γ fixes any singular point of any fiber in $r^{-1}(r(U))$. For a point $x \in r(U)$ we denote by Γ_x the subgroup of Γ that preserves the fiber $r^{-1}(x)$. The group Γ_x acts by automorphisms on $r^{-1}(x)$. The group Γ_r is a normal subgroup of Γ_x for all x .

- Lemma 9.3.1.** (1) For any $x \in r(U)$, the intersection of U with $r^{-1}(x)$ has finitely many connected components.
(2) If Ω is a connected component of $U \cap r^{-1}(x)$, then the subgroup Γ_Ω of Γ preserving Ω is a finite index subgroup of Γ_x .
(3) For any $x \in r(U)$, the fiber $r^{-1}(x)$ is non-singular, i.e. $r^{-1}(x) = \mathbb{P}^1$.
(4) $r(U)$ is a proper subset of $B = \mathbb{P}^1$.

Proof. Let Ω be a connected component of $U \cap r^{-1}(x)$ for some point $x \in r(U)$. It is a leaf of the foliation $\pi^* \mathcal{F}$ on U , i.e. $\pi(\Omega)$ is a fiber of $f : X \rightarrow C$. We assume that it is not a multiple fiber. Denote by Γ_Ω the subgroup of Γ that preserves Ω ; it is a subgroup of Γ_x . The quotient of Ω by Γ_Ω is the fiber $\pi(\Omega)$. The group Γ_Ω is the image of the composition $\pi_1(\pi(\Omega)) \rightarrow \pi_1(X) \rightarrow \Gamma$. Therefore Γ_Ω is a normal subgroup of Γ and it does not depend on the leaf Ω as long as $\pi(\Omega)$ is a regular fiber of $f : X \rightarrow C$. In particular Γ_Ω preserves fiberwise the rational fibration, i.e. $\Gamma_\Omega \subset \Gamma_r$. For a leaf Ω' such that $\pi(\Omega')$ is a multiple fiber, Γ_Ω is a subgroup of finite index of $\Gamma_{\Omega'}$.

Suppose that $r^{-1}(x)$ is a singular fiber of the rational fibration. By our hypothesis at the beginning of the section, Γ_Ω fixes all singular points of $r^{-1}(x)$. Thus it is solvable. Consequently Ω is biholomorphic to \mathbf{C} or \mathbf{C}^* and fibers of f are elliptic curves. This

implies that the intersection of U with an irreducible component of $r^{-1}(x)$ is either empty or biholomorphic to \mathbf{C} or \mathbf{C}^* . Thus the first assertion of our lemma is true if $r^{-1}(x)$ is a singular fiber. Let us assume now that $r^{-1}(x)$ is not singular, i.e. is isomorphic to \mathbb{P}^1 . Then we can think of Γ_Ω as a classical Kleinian group with an invariant component Ω . The intersection of U with $r^{-1}(x)$ is a union of connected components of Γ_Ω . By the above discussion all of them are either Γ_Ω -invariant, or invariant under a finite index subgroup of Γ_Ω if $\pi(\Omega)$ is a multiple fiber. Ahlfors's finiteness theorem (see [Ahl64], [Sul85]) says that $(U \cap r^{-1}(x))/\Gamma_\Omega$ is a finite union of compact Riemann surfaces. This implies that $r^{-1}(x) \cap U$ has only finitely many connected components.

In the first paragraph of the proof we inferred $\Gamma_\Omega \subset \Gamma_r$; we also have an inclusion $\Gamma_r \subset \Gamma_x$. The above finiteness of connected components implies that Γ_Ω has finite index in Γ_x . From this and the infiniteness of Γ_B we deduce that the orbit of any point $x \in r(U)$ has an infinite orbit under Γ_B . As Γ acts by holomorphic diffeomorphisms on $r^{-1}(r(U))$, we conclude that r has no singular fibers over $r(U)$.

Any element of Γ_B has a fixed point in $B = \mathbb{P}^1$. If $r(U) = B$ then a fixed point x of some element of infinite order of Γ_B would give rise to a Γ_x such that Γ_r has infinite index in Γ_x , contradicting the second point of the lemma. \square

Up to making elementary transformations on a fiber over a point outside $r(U)$ we can and will assume that $Y = \mathbb{P}^1 \times \mathbb{P}^1$ and that r is the projection onto the first factor. As a finitely generated subgroup of $\mathrm{PGL}(2, \mathbf{C})$, Γ_B has a torsion free subgroup of finite index by Selberg's lemma (see [Sel60]). Up to replacing Γ with the preimage of such a subgroup of Γ_B , we can and will assume that Γ_B is *torsion free*. This is equivalent to say that *for any* $x \in B$, *we have* $\Gamma_x = \Gamma_r$.

By Proposition 7.1.5, \mathcal{F} is equipped with a foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure. We claim that regular fibers of f are isomorphic to each other and have the same $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure. This is clear if the fibers are \mathbb{P}^1 . If a fiber of f is an elliptic curve, then its $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure is induced by one of its infinite covers \mathbf{C} or \mathbf{C}^* , depending on whether Γ_r has rank one or two. If fibers of f have genus ≥ 2 , then the claim follows from Proposition 6.2.3.

We will think of Γ_r as a family of classical Kleinian groups parametrized by $r(U)$. Let Ω be a connected component of the preimage of a regular fiber of f ; it is contained in a fiber of r . Denote by Γ_Ω the subgroup of Γ that preserves Ω . The pair (Ω, Γ_Ω) gives the developing map and holonomy representation of the $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure on the corresponding fiber of f . Therefore up to conjugation they do not depend on the fiber containing Ω . More precisely there exists a holomorphic map $H : r(U) \rightarrow \mathrm{PGL}(2, \mathbf{C})$ such that $H\Gamma_\Omega H^{-1}$ is a trivial family of classical Kleinian groups. Since Γ_Ω is a subgroup of $\mathrm{PGL}(2, \mathbf{C}(x))$, the conjugation H is actually algebraic, i.e. it is also an element of $\mathrm{PGL}(2, \mathbf{C}(x))$. In other words H realizes a geometric conjugation of our birational Kleinian group so that Γ_Ω becomes contained in the subgroup $\{\mathrm{Id}\} \times \mathrm{PGL}(2, \mathbf{C})$ of $\mathrm{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$. Moreover we obtain $U = r(U) \times \Omega$ by connectedness of U . Thus $\Gamma_r = \Gamma_\Omega$. Consequently $f : X \rightarrow C$ is a locally trivial fibration without multiple fibers and $r(U)/\Gamma_B = C$. In summary we have:

Proposition 9.3.2. *Up to geometric conjugation and up to taking finite index subgroup, $Y = \mathbb{P}^1 \times \mathbb{P}^1$, $U = r(U) \times \Omega$ and $\Gamma_r = \Gamma_\Omega$ is a subgroup of $\{\mathrm{Id}\} \times \mathrm{PGL}(2, \mathbf{C})$.*

Finally we distinguish three situations according to whether the fibers of f are \mathbb{P}^1 , elliptic curves or hyperbolic Riemann surfaces. If the fibers of f are \mathbb{P}^1 , then U is necessarily $r(U) \times \mathbb{P}^1$.

Example 9.3.3. Consider the cyclic group Γ generated by

$$(x, y) \mapsto (bx, cx^k y), \quad b, c \in \mathbf{C}^*, k \in \mathbf{Z}^* \mid b \neq 1.$$

The quotient $\mathbf{C}^* \times \mathbb{P}^1 / \Gamma$ is a geometrically ruled surface over an elliptic curve; it is a decomposable ruled surface because of the two disjoint sections $\{y = 0\}, \{y = \infty\}$. This is our only example of birational Kleinian group for which Γ is not conjugate to a group of automorphisms of a projective surface and X is Kähler.

Proposition 9.3.4. *Suppose that the fibers of f have genus 1. Then up to geometric conjugation and up to finite index subgroup we are in one of the two following situations*

- (1) $Y = \mathbb{P}^1 \times \mathbb{P}^1$, $U = D_1 \times \mathbf{C}^*$ and D_1 is an invariant component of the classical Kleinian group Γ_B . Γ is a subgroup of $J^+ := \{(x, y) \mapsto (\eta(x), R(x)y) \mid \eta \in \mathrm{PGL}(2, \mathbf{C}), R \in \mathbf{C}(x)^*\} = \mathbf{C}(x)^* \rtimes \mathrm{PGL}(2, \mathbf{C})$. $\Gamma_r \subset \{\mathrm{Id}\} \times \mathrm{PGL}(2, \mathbf{C})$ is a cyclic central subgroup of Γ .
- (2) $Y = \mathbb{P}^1 \times \mathbb{P}^1$, $U = D_1 \times \mathbf{C}$ and D_1 is an invariant component of the classical Kleinian group Γ_B . Γ is a subgroup of $J^* := \{(x, y) \mapsto (\eta(x), y + R(x)) \mid \eta \in \mathrm{PGL}(2, \mathbf{C}), R \in \mathbf{C}(x)\} = \mathbf{C}(x) \rtimes \mathrm{PGL}(2, \mathbf{C})$. $\Gamma_r \subset \{\mathrm{Id}\} \times \mathrm{PGL}(2, \mathbf{C})$ is a central subgroup of Γ isomorphic to \mathbf{Z}^2 .

Proof. By the exact sequence $1 \rightarrow \Gamma_r \rightarrow \Gamma \rightarrow \Gamma_B \rightarrow 1$ the group Γ_B acts by outer automorphisms on the abelian group Γ_r . Any such outer automorphism induces an automorphism of a fiber of f ; it is an automorphism of an elliptic curve fixing the origin. Therefore up to replacing Γ by a finite index subgroup, we can and will assume that Γ_r is central in Γ . This is equivalent to say that we replace X with a finite unramified cover which is a principal elliptic bundle (cf. [BHPVdV04] V.5).

It remains to show that Γ is included in J^+ or J^* . This is because the centralizer of Γ_r is respectively J^+ or J^* by Theorem 4.4.1. \square

Let us see two examples where the group Γ is not conjugate to a subgroup of $\mathrm{PGL}(2, \mathbf{C}) \times \mathrm{PGL}(2, \mathbf{C})$. The first is given by the standard construction of primary Kodaira surfaces (see Paragraph 3.2.4). In the second example Γ is not conjugate to a group of automorphisms of a projective surface:

Example 9.3.5. Consider the group Γ generated by the following two elements

$$(x, y) \mapsto (x, ay)$$

$$(x, y) \mapsto (bx, cx^k y), \quad a, b, c \in \mathbf{C}^*, k \in \mathbf{Z}^* \mid |a|, |b| \neq 1.$$

By lifting the deck transformations to \mathbf{C}^2 we see that the quotient $\mathbf{C}^* \times \mathbf{C}^* / \Gamma$ is a primary Kodaira surface. Note Γ is included in the toric subgroup. The birational transformation $(x, y) \mapsto (bx, cx^k y)$ has linear degree growth and is not birationally conjugate to an automorphism of a projective surface.

Proposition 9.3.6. *Suppose that the fibers of f have genus ≥ 2 . Then up to geometric conjugation and up to finite index subgroup we have: $Y = \mathbb{P}^1 \times \mathbb{P}^1$, $\Gamma = \Gamma_B \times \Gamma_r$ and $U = D_1 \times D_2$ where D_1, D_2 are respectively invariant components of the classical Kleinian groups Γ_B, Γ_r .*

Proof. For $x \in r(U)$, we denote by \mathbb{P}_x^1 the fiber of r over x and by Γ_x the restriction of Γ_r on \mathbb{P}_x^1 . The classical Kleinian groups Γ_x are the same if we identify the different \mathbb{P}_x^1 .

Let γ be an element of Γ_B . We write it as $\gamma: (x, y) \mapsto (\gamma_B(x), \rho(x)(y))$ where γ_B is the image of γ in Γ_B and ρ is a rational map from B to $\mathrm{PGL}(2, \mathbf{C})$. Since Γ_r is a normal

subgroup, we have $\gamma\Gamma_r\gamma^{-1} = \Gamma_r$. If we think of Γ_r as a classical Kleinian group, then we have $\rho(x)\Gamma_r\rho(x)^{-1} = \Gamma_r$. This means that for any $x \in r(U)$, $\rho(x)$ is in the normalizer Γ_r in $\mathrm{PGL}(2, \mathbf{C})$. As the fibers of f are hyperbolic, Γ_r is a non-elementary Kleinian group and its normalizer is discrete (see [Mas88] Chapter V E.10). Hence ρ is constant and $\gamma \in \mathrm{PGL}(2, \mathbf{C}) \times \mathrm{PGL}(2, \mathbf{C})$

Therefore Γ is a subgroup of $\mathrm{PGL}(2, \mathbf{C}) \times \mathrm{PGL}(2, \mathbf{C})$. Let Γ_2 be its projection onto the second factor; it is contained in the normalizer of Γ_r . Since the fibers of the locally trivial fibration $f : X \rightarrow C$ have genus ≥ 2 , there exists a finite unramified cover \bar{C} of C such that the pullback fibration over \bar{C} is simply a product (see [BHPVdV04] V.6). In particular a finite unramified cover of X is a product of two curves. This implies Γ_2/Γ_r is a finite group. Hence up to replacing Γ with a finite index subgroup, we have $\Gamma_2 = \Gamma_r$ and $\Gamma = \Gamma_B \times \Gamma_r$. \square

9.4. Complex tori.

Proposition 9.4.1. *Assume that $B = \mathbb{P}^1$, Γ_B is infinite and X is a complex torus. Then up to geometric conjugation the birational Kleinian group (Y, Γ, U, X) satisfies that*

- $Y = \mathbb{P}^1 \times \mathbb{P}^1$. U is one of the three Zariski open sets: \mathbf{C}^2 , $\mathbf{C} \times \mathbf{C}^*$ or $\mathbf{C}^* \times \mathbf{C}^*$.
- Γ can be identified with a lattice in U with its natural structure of algebraic group, i.e. the elements of Γ have respectively the form $(x, y) \mapsto (x + a, y + b)$, $(x, y) \mapsto (x + a, by)$ or $(x, y) \mapsto (ax, by)$.

Proof. The fundamental group of a complex torus is isomorphic to \mathbf{Z}^4 . After replacing Γ with a subgroup of finite index we can and will assume that Γ and Γ_B are free abelian groups. Remark that the only regular foliations on complex tori are turbulent foliations and linear foliations. We know by Proposition 9.1.6 that \mathcal{F} is linear.

A linear foliation on \mathbf{C}^2 is induced by a linear map from \mathbf{C}^2 to \mathbf{C} equivariant under $\pi_1(X)$, the action on \mathbf{C} being by translations. The image of $\pi_1(X)$ in $\mathrm{Aut}(\mathbf{C})$ is a free abelian group of rank 2, 3 or 4, depending on whether the leaves of \mathcal{F} are elliptic curves, \mathbf{C}^* or \mathbf{C} . We obtain a surjective holomorphic map $\mathbf{C} \rightarrow r(U)$ that is equivariant with respect to the homomorphism $\pi_1(X) \rightarrow \Gamma_B$. This forces $r(U)$ to be \mathbf{C} or \mathbf{C}^* . Then Γ_B is respectively a discrete group of translations or multiplications. In particular every point of $r(U)$ has an infinite Γ_B -orbit. Thus r does not have singular fibers over $r(U)$ by Corollary 7.1.3. Consequently U is a \mathbf{C} or \mathbf{C}^* bundle over \mathbf{C} or \mathbf{C}^* . Being an infinite cover of a torus, U is biholomorphic to \mathbf{C}^2 , $\mathbf{C} \times \mathbf{C}^*$ or $\mathbf{C}^* \times \mathbf{C}^*$. In other words the complement $r^{-1}(r(U)) \setminus U$ is one or two holomorphic sections of r over $r(U)$. We need to prove that these sections are algebraic. We know that Γ is a free abelian group of rank ≥ 2 .

We assume first that Γ_B is free of rank ≥ 2 . In this case the descriptions of Γ and U given in the statement follow from Theorem 4.4.3. We only need to observe that the conjugation that we need in Theorem 4.4.3 is a geometric conjugation of birational Kleinian groups. This is because the conjugation is a sequence of elementary transformations. If we let the elementary transformations to be done outside U , then the conjugation is geometric. Remark that the hypothesis that the rank of Γ_B is ≥ 2 is always satisfied when \mathcal{F} is irrational. This is because if \mathcal{F} is irrational then the action of Γ_B on $r(U)$ is not discrete.

Consider now the case where Γ_B is cyclic. In this case $r(U)$ is necessarily \mathbf{C}^* , \mathcal{F} is an elliptic fibration and Γ_r , the subgroup of Γ preserving fiberwise the rational fibration, is a free abelian group of rank 1 or 2. If Γ_r has rank 1 then $r^{-1}(x) \cap U = \mathbf{C}^*$ for any $x \in r(U)$ and Γ_r acts by multiplications in the fibers; if Γ_r has rank 2 then $r^{-1}(x) \cap U = \mathbf{C}$ for any $x \in r(U)$ and Γ_r acts by translations in the fibers.

First consider the case where Γ_r has rank 1. By Theorem 4.4.3 up to conjugation Γ_r is generated by an element $\gamma_1 : (x, y) \mapsto (x, ay)$ and Γ is generated by γ_1 and $\gamma_2 : (x, y) \mapsto (bx, R(x)y)$ with $R \in \mathbf{C}(x)^*$; being a sequence of elementary transformations the conjugation can be done outside U so that it is a geometric conjugation of birational Kleinian groups. Therefore U is the standard Zariski open set $\mathbf{C}^* \times \mathbf{C}^*$. As γ_2 acts regularly over $r(U) = \mathbf{C}^*$, the rational function R has neither zeros nor poles over \mathbf{C}^* . Thus $R(x) = x^k$ for some $k \in \mathbf{Z}$. If $k \neq 0$ then the quotient $\mathbf{C}^* \times \mathbf{C}^*$ would be a primary Kodaira surface (cf. Example 9.3.5). Hence the conclusion.

Finally consider case where Γ_r has rank 2. By Theorem 4.4.3 up to conjugation Γ_r is generated by two elements $\gamma_j : (x, y) \mapsto (x, y + a_j)$, $j = 1, 2$ and Γ is generated by γ_1, γ_2 and $\gamma_3 : (x, y) \mapsto (bx, y + R(x))$ with $R \in \mathbf{C}(x)$. Again the conjugation can be chosen to be geometric conjugation. Therefore U is the standard Zariski open set $\mathbf{C}^* \times \mathbf{C}$. As γ_3 acts in a regular way over $r(U) = \mathbf{C}^*$, the rational function R has no poles over \mathbf{C}^* . Thus $R(x) = \frac{Q(x)}{x^k}$ for some $k \in \mathbf{Z}$ and $Q \in \mathbf{C}[x]$. The iterates of γ_3 can be written as

$$\gamma_3^n : (x, y) \mapsto (b^n x, y + S(x)) \quad \text{where} \quad S(x) = \frac{1}{x^k} \sum_{j=0}^{n-1} \frac{Q(b^j x)}{b^k}.$$

Thus γ_3 has bounded degree growth and is an elliptic element. By conjugating Γ with elements of the form $(x, y) \mapsto (x, y + \frac{P(x)}{x^k})$ with $P \in \mathbf{C}[x]$, we do not change γ_1, γ_2 and can put γ_3 in the form $(x, y) \mapsto (bx, y)$. Hence the conclusion. \square

9.5. Bidisk quotients. Finally we consider the case where \mathcal{F} is a transversely hyperbolic foliation with dense leaves. By Proposition 9.1.5 $r(U)$ is a proper subset of \mathbb{P}^1 . Since every leaf of \mathcal{F} is dense in X , any Γ_B -orbit in $r(U)$ is infinite. This implies by Corollary 7.1.3 that $r^{-1}(r(U))$ is biholomorphic to $r(U) \times \mathbb{P}^1$. Thus by Proposition 7.1.5 \mathcal{F} is equipped with a foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure. By Corollary 6.2.2 X is a bidisk quotient.

Consider the composition $\pi_1(X) \rightarrow \Gamma \rightarrow \Gamma_B \subset \mathrm{PGL}(2, \mathbf{C})$. Since $\pi_1(X)$ is an irreducible lattice in $\mathrm{PSL}(2, \mathbf{R}) \times \mathrm{PSL}(2, \mathbf{R})$, Margulis superrigidity ([Mar91]) says that the quotient homomorphism $\pi_1(X) \rightarrow \Gamma_B$ is an isomorphism. Up to conjugation in $\mathrm{PGL}(2, \mathbf{C})$ and up to a Galois conjugate, Γ_B is exactly the projection of the lattice into one of the factors $\mathrm{PSL}(2, \mathbf{R})$. In particular Γ_B is a dense subgroup of $\mathrm{PSL}(2, \mathbf{R})$. This implies that $r(U)$ is a round disk \mathbb{D} in \mathbb{P}^1 and that U is biholomorphic to the bidisk, the universal cover of X .

By Proposition 6.2.4 the foliated $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -structure on \mathcal{F} is the natural one; every leaf of \mathcal{F} can be identified with the unit disk as $(\mathrm{PGL}(2, \mathbf{C}), \mathbb{P}^1)$ -manifold. Consequently $r^{-1}(x) \cap U$ is a round disk for any $x \in r(U)$. Furthermore there exists a holomorphic map $h : \mathbb{D} \rightarrow \mathrm{PGL}(2, \mathbf{C})$ and a $\pi_1(X)$ -equivariant biholomorphism $H : \mathbb{D} \times \mathbb{P}^1 \rightarrow \mathbb{D} \times \mathbb{P}^1$ defined by $(x, y) \mapsto (x, h(x)(y))$ which sends U to $\mathbb{D} \times \mathbb{D}$.

Proposition 9.5.1. *If \mathcal{F} is a transversely hyperbolic foliation with dense leaves, then up to geometric conjugation we are in the standard example of bidisk quotient.*

Proof. We would like to show that H is the restriction of a birational transformation preserving the rational fibration, i.e. an element of the Jonquières group. Then it realizes the desired geometric conjugation of birational Kleinian groups.

Our group Γ is isomorphic to a cocompact irreducible lattice in $\mathrm{PSL}(2, \mathbf{R}) \times \mathrm{PSL}(2, \mathbf{R})$. We will use two properties of irreducible lattices:

- (1) Γ has subgroups isomorphic to \mathbf{Z}^2 because $\mathrm{PSL}(2, \mathbf{R}) \times \mathrm{PSL}(2, \mathbf{R})$ has real rank two (see [PR72]).

(2) Elements Γ_B , are either elliptic or loxodromic elements of $\text{PSL}(2, \mathbf{R})$ because every element of a cocompact lattice is semisimple (see [BHC62], [MT62]).

Let $\gamma_1, \gamma_2 \in \Gamma$ such that $\langle \gamma_1, \gamma_2 \rangle$ a subgroup isomorphic to \mathbf{Z}^2 . Theorem 4.4.3 together with the second property above allow us to write γ_1, γ_2 , up to conjugation in Jonq , as $\gamma_i : (x, y) \mapsto (a_i x, b_i y), i = 1, 2$ or $\gamma_i : (x, y) \mapsto (a_i x, b_i + y), i = 1, 2$. Here the latter case is impossible because the holomorphic conjugation H is supposed to make $\gamma_i, i = 1, 2$ also loxodromic in the y coordinate. Hence we can write them as $\gamma_i : (x, y) \mapsto (a_i x, b_i y), i = 1, 2$.

Denote by ∂U the boundary of U in $\mathbb{P}^1 \times \mathbb{P}^1$ and by $\partial_h(U)$ the boundary of U in $\mathbb{D} \times \mathbb{P}^1$. The two horizontal disks $S_1 = \{y = 0, x \in \mathbb{D}\}, S_2 = \{y = \infty, x \in \mathbb{D}\}$ are invariant under $\gamma_i, i = 1, 2$. We want to show that they are contained in $\partial_h(U)$. Since Γ_B is isomorphic to Γ , the subgroup of \mathbf{C}^* generated by a_1, a_2 is isomorphic to \mathbf{Z}^2 , thus dense. Therefore there is a non-trivial sequence of birational transformations (δ_n) in $\langle \gamma_1, \gamma_2 \rangle$ such that δ_{nB} tends to the identity in $\text{PGL}(2, \mathbf{C})$. Up to extracting a subsequence, $\delta_n|_U$ converges to a holomorphic map from U to $U \cap \partial U$. By discontinuity of the action of Γ on U , the limit map has values in ∂U ; as δ_{nB} tends to the identity, the image of the limit map is in $\partial_h(U)$. Since each δ_n has the form $(x, y) \mapsto (c_n x, d_n y)$, the limit disk is either S_1 or S_2 . By considering the sequence (δ_n^{-1}) , we infer that S_1 and S_2 are both in $\partial_h(U)$.

Consider the conjugates $H \circ \delta_n \circ H^{-1}$. The sequence $H \circ \delta_n \circ H^{-1}|_{\mathbb{D} \times \mathbb{D}}$ converges to a horizontal disk as well. Therefore we can assume that $H(S_i) = S_i, i = 1, 2$. In other words the holomorphic conjugation map H has the form $(x, y) \mapsto (x, A(x)y)$ where A is a holomorphic function $\mathbb{D} \rightarrow \mathbf{C}^*$. To conclude we need to show that A is algebraic. It suffices to exhibit other disks in $\partial_h(U)$ defined by algebraic functions. We can conjugate the γ_1, γ_2 in Γ to obtain $\gamma_3, \gamma_4 \in \Gamma$ such that $\langle \gamma_3, \gamma_4 \rangle$ is isomorphic to \mathbf{Z}^2 but γ_3, γ_4 do not commute with γ_1, γ_2 . We apply the above discussion to γ_3, γ_4 and obtain other algebraic disks in $\partial_h(U)$. The proof is finished. \square

10. CLASSIFICATION AND PROOF

10.1. Classification.

Theorem C. *Let (Y, U, Γ, X) be a birational Kleinian group in dimension 2. Suppose that*

- (1) Γ is not a cyclic group generated by a loxodromic element, and
- (2) if X is a class VII surface then Γ contains no loxodromic elements.

Then up to geometric conjugation and up to taking a finite index subgroup, we are in one of the cases in the following table:

Birational Kleinian groups in dimension two				
	Y	U	Γ	X
1	$B \times \mathbb{P}^1$ where B is compact Riemann surface	$B \times D_1$ where D_1 is an invariant component of a classical Kleinian group Γ_1	$\{\text{Id}\} \times \Gamma_1 \subset \text{Aut}(B) \times \text{Aut}(\mathbb{P}^1)$	$B \times (D_1/\Gamma_1)$
2	$\mathbb{P}(\mathcal{E})$ where \mathcal{E} is an extension of \mathcal{O}_B by \mathcal{O}_B and B is compact Riemann surface	the complement of a section of the ruling	isomorphic to \mathbf{Z}^2	a principal elliptic bundle

	Y	U	Γ	X
3	a blow-up of a decomposable ruled surface	a Zariski open subset whose intersection with each fiber of the ruling is \mathbf{C}^*	isomorphic to \mathbf{Z}	an elliptic fibration with only type mI_0 singular fibers
4	$B \times \mathbb{P}^1$ where B is an elliptic curve	$B \times D_1$ where D_1 is an invariant component of a classical Kleinian group Γ_1	$\exists \rho : \Gamma_1 \rightarrow \text{Aut}(B), \forall \gamma \in \Gamma, \exists \gamma_1 \in \Gamma_1, \gamma = (\rho(\gamma_1), \gamma_1) \in \text{Aut}(B) \times \text{Aut}(\mathbb{P}^1)$	B -bundle over D_1/Γ_1
5	an indecomposable geometrically ruled surface over an elliptic curve with a section of zero self-intersection	the complement of the section of zero self-intersection	isomorphic to \mathbf{Z}^2	complex torus
6	a decomposable geometrically ruled surface over an elliptic curve	the complement of two disjoint sections	isomorphic to \mathbf{Z}	complex torus
7	\mathbb{P}^2	\mathbf{C}^2	isomorphic to \mathbf{Z}^4	complex torus
8	\mathbb{P}^2	$\mathbf{C} \times \mathbf{C}^*$	isomorphic to \mathbf{Z}^3	complex torus
9	\mathbb{P}^2	$\mathbf{C}^* \times \mathbf{C}^*$	isomorphic to \mathbf{Z}^2	complex torus
10	\mathbb{P}^2	\mathbf{C}^2	a group of affine transformations, an extension of \mathbf{Z}^2 by \mathbf{Z}^2	primary Kodaira surface
11	\mathbb{P}^2	$\mathbf{C}^2 \setminus \{0\}$	isomorphic to \mathbf{Z}	Hopf surface
12	\mathbb{P}^2	$\mathbb{H} \times \mathbf{C}$	a solvable group of affine transformations	Inoue surface
13	\mathbb{P}^2	\mathbb{B}^2	a cocompact lattice in $\text{PU}(1, 2)$	ball quotient
14	$\mathbb{P}^1 \times \mathbb{P}^1$	$\mathbb{D}^1 \times \mathbb{D}^1$	an irreducible cocompact lattice in $\text{PSL}(2, \mathbf{R}) \times \text{PSL}(2, \mathbf{R})$	bidisk quotient
15	a Hirzebruch surface blown-up at at most two fibers	a Zariski open subset whose intersection with each fiber of the ruling is \mathbf{C}^*	a cyclic group generated by $(x, y) \mapsto (ax, by)$	Hopf surface
16	a Hirzebruch surface blown-up at at most one fiber	a Zariski open subset whose intersection with each fiber of the ruling is \mathbf{C}^*	a cyclic group generated by $(x, y) \mapsto (x + a, by)$	Hopf surface

	Y	U	Γ	X
17	$\mathbb{P}^1 \times \mathbb{P}^1$	$D_1 \times D_2$ where D_i is an invariant component of a classical Kleinian group Γ_i	$\Gamma_1 \times \Gamma_2 \subset \text{PGL}(2, \mathbf{C}) \times \text{PGL}(2, \mathbf{C})$	$(D_1/\Gamma_1) \times (D_2/\Gamma_2)$
18	$\mathbb{P}^1 \times \mathbb{P}^1$	$D_1 \times \mathbf{C}^*$ where D_1 is an invariant component of a classical Kleinian group Γ_1	isomorphic to a central extension of Γ_1 by \mathbf{Z} , any element has the form $(x, y) \mapsto (\gamma_1(x), R(x)y)$ where $\gamma_1 \in \Gamma_1$ and $R \in \mathbf{C}(x)^*$	principal elliptic bundle over D_1/Γ_1
19	$\mathbb{P}^1 \times \mathbb{P}^1$	$D_1 \times \mathbf{C}$ where D_1 is an invariant component of a classical Kleinian group Γ_1	isomorphic to a central extension of Γ_1 by \mathbf{Z}^2 , any element has the form $(x, y) \mapsto (\gamma_1(x), y + R(x))$ where $\gamma_1 \in \Gamma_1$ and $R \in \mathbf{C}(x)$	principal elliptic bundle over D_1/Γ_1
20	$\mathbb{P}^1 \times \mathbb{P}^1$	$D_1 \times \mathbb{P}^1$ where D_1 is an invariant component of a classical Kleinian group Γ_1	isomorphic to Γ_1 , preserves the projection of $\mathbb{P}^1 \times \mathbb{P}^1$ onto the first factor	geometrically ruled surface over D_1/Γ_1

The surface Y is rational unless in the first six cases. The group Γ is a group of automorphisms except in the last three cases. In cases 18), 19), 20), Γ may and may not be conjugate to a group of automorphisms of any projective surface.

We explain how Theorem C implies Theorem B. Under the hypothesis that U is simply connected and X is Kähler, the group Γ is isomorphic to the Kähler group $\pi_1(X)$. By Hodge theorem the abelianization of a Kähler group has even rank, thus is not cyclic. Hence the hypothesis of Theorem B implies the hypothesis of Theorem C. It suffices thus to remark that in the list given by Theorem C the only cases where U is simply connected and X is Kähler are the seven cases listed in Theorem B.

When X is an Inoue surface, we proved in our previous paper [Zhaç] that there is no cyclic birational Kleinian group having quotient surface X ; in other words the two hypothesis in Theorem C can be dropped. The proof uses particular geometric features of Inoue surfaces as well as ergodic properties of loxodromic birational transformations. Known Class VII surfaces have rather small fundamental groups, typically cyclic. The author does not know any example of cyclic birational Kleinian group generated by a loxodromic element.

10.2. Proof. Let (Y, Γ, U, X) be a birational Kleinian group in dimension two. By Theorem A Y has Kodaira dimension $-\infty$. Thus it is either rational or birational to a ruled surface over a non-rational curve.

Suppose that Y is not rational. Then there is a rational fibration $r: Y \rightarrow B$ onto a curve B of genus ≥ 1 . Every birational transformation of Y preserves the fibration r . If Γ_B is a finite

group then all possibilities are classified by Theorem 7.2.1; this gives the first three cases in Theorem C. If Γ_B is infinite then B is an elliptic curve and all possibilities are classified by Theorem 8.0.1; this gives cases 1), 4), 5) and 6) in Theorem C.

From now on we assume that Y is a rational surface. We apply Theorem 4.3.1 to Γ . We will first deal with the cases where a rational fibration is preserved by Γ up to conjugation, then deal with other cases of Theorem 4.3.1.

10.2.1. *Rational surfaces I: invariant rational fibrations.* Assume that Γ preserves a rational fibration up to conjugation, i.e. Γ preserves a pencil of rational curves. This hypothesis always holds in Cases 2), 3) of Theorem 4.3.1 and can hold in case 1). Since the action of Γ on U is free, the set of base points of the invariant pencil does not intersect U . We can blow up the base points to get an invariant fibration. Thus we have:

Lemma 10.2.1. *If $\Gamma \subset \text{Bir}(Y)$ is conjugate in $\text{Bir}(Y)$ to a group preserving a fibration, then the conjugation can be chosen to be geometric.*

Therefore we can and will assume that Γ preserves a rational fibration $r : Y \rightarrow B$ where $B = \mathbb{P}^1$. As before we denote by Γ_B the image of Γ in $\text{Aut}(B)$. By Proposition 10.2.1 r induces on X a regular holomorphic foliation \mathcal{F} . If Γ_B is a finite group then all possibilities are classified by Theorem 7.2.1; this gives the first three cases in Theorem C. In this case \mathcal{F} is a fibration.

Assume now that Γ_B is infinite. We distinguish two cases according to whether $r(U) = B$ or $r(U)$ is a proper subset of B .

First case. Suppose that $r(U) = B$. Then by Proposition 9.1.5 the foliation \mathcal{F} is a turbulent foliation, an obvious foliation on a Hopf surface or a suspension of \mathbb{P}^1 .

Assume that \mathcal{F} is a turbulent foliation. Then by Proposition 9.1.6, up to taking a subgroup of finite index of Γ , we are in Examples 9.1.1 and 9.1.3. In the situation of these two examples the quotient X is a Hopf surface or a geometrically ruled surface over an elliptic curve. If X is a Hopf surface then this is Case 15) of Theorem C. If X is a geometrically ruled surface over an elliptic curve, then the situations described in Examples 9.1.1 and 9.1.3 are particular subcases of Case 20) in Theorem C.

Assume that \mathcal{F} is an obvious foliation on a non-elliptic Hopf surface. Then Proposition 9.1.7 classifies all possibilities. There are three cases in Proposition 9.1.7. The last case does not satisfy $r(U) = B$. The first two cases satisfy $r(U) = B$ and they correspond to Cases 15) and 16) of Theorem C.

Assume that \mathcal{F} is a suspension of \mathbb{P}^1 . Then Proposition 9.1.8 says that we are in Case 20) of Theorem C

Second case. Assume now that $r(U) \neq B$. By Propositions 9.1.7 and 9.1.8 \mathcal{F} is neither a turbulent foliation nor a suspension of \mathbb{P}^1 .

If \mathcal{F} is an obvious foliation on an Inoue surface then Theorem 9.0.1 says that we are in Case 12) of Theorem C. If \mathcal{F} is an obvious foliation on a non-elliptic Hopf surface then Proposition 9.1.7 classifies all possibilities; only the last case of Proposition 9.1.7 satisfies $r(U) \neq B$ and it corresponds to Case 11) of Theorem C. If \mathcal{F} is a suspension of an elliptic curve, then Proposition 9.2.5 says that the situation is a particular subcase of Case 18) or 19).

Assume that \mathcal{F} is a fibration. If the genus of a fiber is ≥ 2 then Proposition 9.3.6 says that we are in Case 17) of Theorem C. If the fibers are elliptic curves then Proposition 9.3.4 says that we are in Cases 18) and 19). If the fibers are \mathbb{P}^1 then Proposition 9.3.2 says that we are in Case 20).

If \mathcal{F} is an irrational linear foliation on a complex torus, then Proposition 9.4.1 says that we are in Cases 7), 8) or 9). If \mathcal{F} is a transversely hyperbolic foliation with dense leaves, then Proposition 9.5.1 says that we are in Case 14).

10.2.2. *Rational surfaces II: other cases in strong Tits alternative.* Now we consider Cases 1), 4), 6) and 7) of Theorem 4.3.1 for Γ . Case 5) of Theorem 4.3.1 is excluded by our hypothesis in Theorem C. Note that we do not need to consider Case 8) of Theorem 4.3.1 because being a quotient of the fundamental group of a compact manifold, Γ is finitely generated.

Genus one fibration. Let us consider the fourth case of Theorem 4.3.1. The elements of Γ are Halphen twists. There is a rational surface Y' and a birational map $\phi : Y \rightarrow Y'$ such that $\Gamma' = \phi\Gamma\phi^{-1}$ acts by automorphisms and preserves a genus one fibration $\varphi : Y' \rightarrow C$. By Lemma 5.2.2 or Lemma 10.2.1 we can choose Y' and ϕ so that the conjugation ϕ is a geometric conjugation of birational Kleinian groups. For any Halphen twist f preserving φ , the induced action of f on C has finite order (see [CF03] Proposition 3.6). This is impossible by Lemma 2.2.2.

Elliptic group. Let us consider the first case of Theorem 4.3.1. In this case there is a projective surface Y' and a birational map $\phi : Y \rightarrow Y'$ such that $\Gamma' = \phi\Gamma\phi^{-1}$ is a group of automorphisms of Y' and a finite index subgroup of Γ' is in $\text{Aut}^0(Y')$. By Lemma 5.2.2 we can choose Y' and ϕ so that the conjugation ϕ is a geometric conjugation of birational Kleinian groups. Thus we can and will assume that $\Gamma \subset \text{Aut}^0(Y)$.

Assume that Y is not \mathbb{P}^2 . Then there is a birational morphism $\varphi : Y \rightarrow Y'$ to a Hirzebruch surface Y' and we have $\varphi \circ \text{Aut}^0(Y) \circ \varphi^{-1} \subset \text{Aut}^0(Y')$. Since the Aut^0 of a Hirzebruch surface preserves a rational fibration, Γ preserves a rational fibration on Y . Then all possibilities are already classified in 10.2.1.

In the remaining case $Y = \mathbb{P}^2$ and $\Gamma \subset \text{PGL}_3(\mathbb{C}) = \text{Aut}(\mathbb{P}^2)$. This is exactly the case of complex projective Kleinian groups. The complete classification is given by Theorem 3.1.1; it corresponds to Cases 7)–13) in Theorem C.

Toric subgroup. Proposition 5.3.3 says that, under the hypothesis that Γ is not cyclic, if Γ is conjugate in $\text{Bir}(Y)$ to a subgroup of the toric subgroup then it contains no loxodromic elements. Thus Case 6) of Theorem 4.3.1 is impossible.

Non-elementary subgroup. Theorem 5.1.1 says that Γ is not a non-elementary subgroup of $\text{Bir}(Y)$ under the hypothesis that X is not of class VII. Thus Case 7) of Theorem 4.3.1 is ruled out.

REFERENCES

- [Ahl64] Lars V. Ahlfors. Finitely generated Kleinian groups. *Amer. J. Math.*, 86:413–429, 1964.
- [BC16] Jérémy Blanc and Serge Cantat. Dynamical degrees of birational transformations of projective surfaces. *J. Am. Math. Soc.*, 29(2):415–471, 2016.
- [BCM12] Jeffrey Brock, Richard Canary, and Yair Minsky. The classification of Kleinian surface groups, II: The ending lamination conjecture. *Ann. of Math. (2)*, 176(1):1–149, 2012.
- [BD15] Jérémy Blanc and Julie Déserti. Degree growth of birational maps of the plane. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 14(2):507–533, 2015.
- [Bea96] Arnaud Beauville. *Complex algebraic surfaces*, volume 34 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, second edition, 1996.
- [Ber70] Lipman Bers. On boundaries of Teichmüller spaces and on Kleinian groups. I. *Ann. of Math. (2)*, 91:570–600, 1970.
- [BF13] Jérémy Blanc and Jean-Philippe Furter. Topologies and structures of the Cremona groups. *Ann. of Math. (2)*, 178(3):1173–1198, 2013.

- [BHC62] Armand Borel and Harish-Chandra. Arithmetic subgroups of algebraic groups. *Ann. of Math. (2)*, 75:485–535, 1962.
- [BHPVdV04] Wolf P. Barth, Klaus Hulek, Chris A. M. Peters, and Antonius Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge*. Springer-Verlag, Berlin, second edition, 2004.
- [Bru97] Marco Brunella. Feuilletages holomorphes sur les surfaces complexes compactes. *Ann. Sci. École Norm. Sup. (4)*, 30(5):569–594, 1997.
- [Bru15] Marco Brunella. *Birational geometry of foliations*, volume 1 of *IMPA Monographs*. Springer, Cham, 2015.
- [Can01a] Serge Cantat. Dynamique des automorphismes des surfaces $K3$. *Acta Math.*, 187(1):1–57, 2001.
- [Can01b] Serge Cantat. Sur la dynamique du groupe d’automorphismes des surfaces $K3$. *Transform. Groups*, 6(3):201–214, 2001.
- [Can11] Serge Cantat. Sur les groupes de transformations birationnelles des surfaces. *Ann. of Math. (2)*, 174(1):299–340, 2011.
- [Can18a] Serge Cantat. Automorphisms and dynamics. In *Proceedings of International Congress of mathematicians -2018 Rio de Janeiro*, Vol. II, pages 619–634. 2018.
- [Can18b] Serge Cantat. The Cremona group. In *Algebraic geometry: Salt Lake City 2015*, volume 97 of *Proc. Sympos. Pure Math.*, pages 101–142. Amer. Math. Soc., Providence, RI, 2018.
- [CD12] Dominique Cerveau and Julie Déserti. Centralisateurs dans le groupe de Jonquières. *Michigan Math. J.*, 61(4):763–783, 2012.
- [CF03] Serge Cantat and Charles Favre. Symétries birationnelles des surfaces feuilletées. *J. Reine Angew. Math.*, 561:199–235, 2003.
- [CHK13] Benoît Claudon, Andreas Höring, and János Kollár. Algebraic varieties with quasi-projective universal cover. *J. Reine Angew. Math.*, 679:207–221, 2013.
- [Cla] Benoît Claudon. Non algebraicity of universal covers of kähler surfaces. <https://arxiv.org/abs/1001.2379>.
- [CNS13] Angel Cano, Juan Pablo Navarrete, and José Seade. *Complex Kleinian groups*, volume 303 of *Progress in Mathematics*. Birkhäuser/Springer Basel AG, Basel, 2013.
- [CS14] Angel Cano and José Seade. On discrete groups of automorphisms of $\mathbb{P}_{\mathbb{C}}^2$. *Geom. Dedicata*, 168:9–60, 2014.
- [CT89] James A. Carlson and Domingo Toledo. Harmonic mappings of Kähler manifolds to locally symmetric spaces. *Inst. Hautes Études Sci. Publ. Math.*, pages 173–201, 1989.
- [CT97] James A. Carlson and Domingo Toledo. On fundamental groups of class VII surfaces. *Bull. London Math. Soc.*, 29(1):98–102, 1997.
- [Dés15] Julie Déserti. On solvable subgroups of the Cremona group. *Illinois J. Math.*, 59(2):345–358, 2015.
- [DF01] Jeffrey Diller and Charles Favre. Dynamics of bimeromorphic maps of surfaces. *Amer. J. Math.*, 123(6):1135–1169, 2001.
- [DG21] Bertrand Deroin and Adolfo Guillot. Foliated affine and projective structures. *preprint*, (1), 2021.
- [Dlo16] Georges Dloussky. Special birational structures on non-Kählerian complex surfaces. *J. Math. Pures Appl. (9)*, 106(1):76–122, 2016.
- [DP12] Thomas Delzant and Pierre Py. Kähler groups, real hyperbolic spaces and the Cremona group. *Compos. Math.*, 148(1):153–184, 2012.
- [DS20] David Dumas and Andrew Sanders. Geometry of compact complex manifolds associated to generalized quasi-Fuchsian representations. *Geom. Topol.*, 24(4):1615–1693, 2020.
- [Ehr36] Charles Ehresmann. Sur les espaces localement homogènes. *Enseign. Math.*, 35:317–333, 1936.
- [EKPR12] Philippe Eyssidieux, Ludmil Katzarkov, Tony Pantev, and Mahan Ramachandran. Linear Shafarevich conjecture. *Ann. of Math. (2)*, 176(3):1545–1581, 2012.
- [Fra89] Sidney Frankel. Complex geometry of convex domains that cover varieties. *Acta Math.*, 163(1-2):109–149, 1989.
- [Giz80] Marat Gizatullin. Rational G -surfaces. *Izv. Akad. Nauk SSSR Ser. Mat.*, 44(1):110–144, 239, 1980.
- [Gri71] Phillip A. Griffiths. Complex-analytic properties of certain Zariski open sets on algebraic varieties. *Ann. of Math. (2)*, 94:21–51, 1971.
- [GS85] R. V. Gurjar and A. R. Shastri. Covering spaces of an elliptic surface. *Compositio Math.*, 54(1):95–104, 1985.
- [Gun67] Robert C. Gunning. Special coordinate coverings of Riemann surfaces. *Math. Ann.*, 170:67–86, 1967.

- [Gun78] Robert C. Gunning. On uniformization of complex manifolds: The role of connections. *Mathematical Notes*. Princeton, New Jersey: Princeton University Press. Tokyo: University of Tokyo Press. II, 141 p. \$ 9.50 (1978)., 1978.
- [GW12] Olivier Guichard and Anna Wienhard. Anosov representations: domains of discontinuity and applications. *Invent. Math.*, 190(2):357–438, 2012.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [Iit70] Shigeru Iitaka. Deformations of compact complex surfaces. II. *J. Math. Soc. Japan*, 22:247–261, 1970.
- [Ino74] Masahisa Inoue. On surfaces of Class VII₀. *Invent. Math.*, 24:269–310, 1974.
- [JR15] Priska Jahnke and Ivo Radloff. Projective uniformization, extremal Chern classes and quaternionic Shimura curves. *Math. Ann.*, 363(3-4):753–776, 2015.
- [Kle72] Felix Klein. Vergleichende Betrachtungen über neuere geometrische Forschungen. Programm zum Eintritt in die philosophische Facultät und den Senat der Universität zu Erlangen. Erlangen. A. Deichert (1872)., 1872.
- [Kle93] Felix Klein. A comparative review of recent researches in geometry. *Bull. Amer. Math. Soc.*, 2(10):215–249, 1893.
- [Kli98] Bruno Klingler. Structures affines et projectives sur les surfaces complexes. *Ann. Inst. Fourier (Grenoble)*, 48(2):441–477, 1998.
- [Kli01] Bruno Klingler. Un théorème de rigidité non-métrique pour les variétés localement symétriques hermitiennes. *Comment. Math. Helv.*, 76(2):200–217, 2001.
- [KLP18] Michael Kapovich, Bernhard Leeb, and Joan Porti. Dynamics on flag manifolds: domains of proper discontinuity and cocompactness. *Geom. Topol.*, 22(1):157–234, 2018.
- [KO80] Shoshichi Kobayashi and Takushiro Ochiai. Holomorphic projective structures on compact complex surfaces. *Math. Ann.*, 249(1):75–94, 1980.
- [Kod64] Kunihiko Kodaira. On the structure of compact complex analytic surfaces. I. *Amer. J. Math.*, 86:751–798, 1964.
- [Kod66] Kunihiko Kodaira. On the structure of compact complex analytic surfaces. II. *Amer. J. Math.*, 88:682–721, 1966.
- [Kod67] Kunihiko Kodaira. A certain type of irregular algebraic surfaces. *J. Analyse Math.*, 19:207–215, 1967.
- [KP12] János Kollár and John Pardon. Algebraic varieties with semialgebraic universal cover. *J. Topol.*, 5(1):199–212, 2012.
- [Kul78] Ravindra S. Kulkarni. Groups with domains of discontinuity. *Math. Ann.*, 237(3):253–272, 1978.
- [KUY16] Bruno Klingler, Emmanuel Ullmo, and Andrei Yafaev. The hyperbolic Ax-Lindemann-Weierstrass conjecture. *Publ. Math. Inst. Hautes Études Sci.*, 123:333–360, 2016.
- [Lam01] Stéphane Lamy. L’alternative de Tits pour $\text{Aut}[\mathbb{C}^2]$. *J. Algebra*, 239(2):413–437, 2001.
- [Man86] Yuri I. Manin. *Cubic forms*, volume 4 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, second edition, 1986.
- [Mar71] Masaki Maruyama. On automorphism groups of ruled surfaces. *J. Math. Kyoto Univ.*, 11:89–112, 1971.
- [Mar91] Gregory Margulis. *Discrete subgroups of semisimple Lie groups*, volume 17 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1991.
- [Mas88] Bernard Maskit. *Kleinian groups*, volume 287 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1988.
- [MT62] George Mostow and Tsuneo Tamagawa. On the compactness of arithmetically defined homogeneous spaces. *Ann. of Math. (2)*, 76:446–463, 1962.
- [MY93] Ngaiming Mok and Sai-Keek Yeung. Geometric realizations of uniformization of conjugates of Hermitian locally symmetric manifolds. In *Complex analysis and geometry*, Univ. Ser. Math., pages 253–270. Plenum, New York, 1993.
- [PR72] Gopal Prasad and M. S. Raghunathan. Cartan subgroups and lattices in semi-simple groups. *Ann. of Math. (2)*, 96:296–317, 1972.
- [Ros79] Jean-Pierre Rosay. Sur une caractérisation de la boule parmi les domaines de \mathbb{C}^n par son groupe d’automorphismes. *Ann. Inst. Fourier (Grenoble)*, 29(4):ix, 91–97, 1979.
- [Sel60] Atle Selberg. On discontinuous groups in higher-dimensional symmetric spaces. In *Contributions to function theory (internat. Colloq. Function Theory, Bombay, 1960)*, pages 147–164. Tata Institute of Fundamental Research, Bombay, 1960.

- [Sul85] Dennis Sullivan. Quasiconformal homeomorphisms and dynamics. I. Solution of the Fatou-Julia problem on wandering domains. *Ann. of Math. (2)*, 122(3):401–418, 1985.
- [Tit72] Jacques Tits. Free subgroups in linear groups. *J. Algebra*, 20:250–270, 1972.
- [Tou16] Frédéric Touzet. On the structure of codimension 1 foliations with pseudoeffective conormal bundle. In *Foliation theory in algebraic geometry*, Simons Symp., pages 157–216. Springer, Cham, 2016.
- [Uen75] Kenji Ueno. *Classification theory of algebraic varieties and compact complex spaces. Notes written in collaboration with P. Cherenack.*, volume 439. Springer, Cham, 1975.
- [Ure21] Christian Urech. Subgroups of elliptic elements of the Cremona group. *J. Reine Angew. Math.*, 770:27–57, 2021.
- [Wei55] André Weil. On algebraic groups of transformations. *Amer. J. Math.*, 77:355–391, 1955.
- [Won77] Bun Wong. Characterization of the unit ball in \mathbf{C}^n by its automorphism group. *Invent. Math.*, 41(3):253–257, 1977.
- [Zhaa] Shengyuan Zhao. a note on holomorphic families of complex projective structures. *preprint*.
- [Zhab] Shengyuan Zhao. Centralizers of elements of infinite order in plane Cremona groups. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, to appear.
- [Zhac] Shengyuan Zhao. Uniqueness of birational structures on Inoue surfaces. *preprint*, <https://arxiv.org/abs/1909.13266>.
- [ZS75] Oscar Zariski and Pierre Samuel. *Commutative algebra. Vol. II*. Springer-Verlag, New York-Heidelberg, 1975. Reprint of the 1960 edition, Graduate Texts in Mathematics, Vol. 29.

INSTITUTE FOR MATHEMATICAL SCIENCES, STONY BROOK UNIVERSITY, STONY BROOK NY 11794-3660, USA

Email address: shengyuan.zhao@stonybrook.edu