

Zoology of Graviton non-Gaussianities

Giovanni Cabass^a

^aSchool of Natural Sciences, Institute for Advanced Study, 1 Einstein Dr, Princeton, NJ 08540

E-mail: gcabass@ias.edu

Abstract. We characterize graviton non-Gaussianities in models of inflation where de Sitter boosts are spontaneously broken. We discuss which of the symmetry breaking patterns studied in Nicolis et al., 2015 [1] can sustain a period of quasi de Sitter expansion, and show that the symmetry breaking pattern of Solid Inflation allows the most freedom for graviton interactions. We comment on the phenomenological consequences of some of these interactions. As a byproduct of this analysis we construct the “EFT of Solid Inflation”, which has all the useful features of the EFT of Inflation but for the case of broken spatial diffeomorphisms.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 1 |
| 2 | “Zoology of condensed matter” and coupling with gravity | 2 |
| 2.1 | The eightfold (sixfold) way | 3 |
| 2.2 | Coupling with gravity | 4 |
| 3 | Inflationary solutions and unitary-gauge actions | 4 |
| 3.1 | Type-I superfluids | 4 |
| 3.2 | Solids and fluids | 5 |
| 3.3 | Type-II framid (\supset type-I framid) | 6 |
| 3.4 | A word on the Goldstone modes | 7 |
| 4 | Main results | 8 |
| 4.1 | Most general graviton interactions | 8 |
| 4.2 | A bit of phenomenology | 9 |
| 5 | Conclusions | 12 |
| A | An “EFT of Inflation” for solids | 12 |
| B | Cubic graviton interaction in Solid Inflation | 14 |

1 Introduction

Upcoming CMB experiments will target the tensor-to-scalar ratio r to a sensitivity of $\sigma_r \sim 10^{-3}$ [2, 3, 4, 5, 6, 7, 8, 9, 10]. If primordial B modes are detected, the way towards constraints on tensor non-Gaussianities, and then on the interactions of the graviton during inflation, will open.

Even in the less exciting case that these experiments will only improve the current upper bounds, it is still very interesting to look at what are the possible interactions in quasi de Sitter backgrounds. This is because of recent developments in the “Cosmological Bootstrap” program [11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21]. Indeed, Ref. [21] has discussed how to extend the bootstrap rules to the scenario where de Sitter boosts are broken, more precisely to the phenomenologically-interesting case of single-clock inflation. Working at the level of the Lagrangian can help in developing the bootstrap rules, and also identify regions of parameter space where non-Gaussianities can be enhanced, as is the case for the scalar non-Gaussianity in $P(X, \phi)$ theories.

The case of single-clock inflation is particularly simple to study since only one Goldstone mode is present. Once we move away from this symmetry breaking pattern the number of Goldstone modes increases, and a study of all the possible non-Gaussianities (scalar, vector and tensor) becomes complicated. Since restricting to graviton interactions still allows to highlight the new features of the other symmetry breaking patterns, we have decided to study these first.

Graviton non-Gaussianities in single-clock inflation have been investigated in detail in [11, 22]. What happens when couplings with the foliation are included has been discussed

in [23, 24, 25, 26]. How do we extend the analysis to different symmetry breaking patterns? Ref. [1] provides a guide for how to achieve this. On scales shorter than the horizon the de Sitter isometries reduce to the Poincaré group (special conformal transformations and “de Sitter dilations” – the combination of time translations with spatial dilations – reduce to Lorentz boosts and time translations, respectively). Then, we can gain intuition from the flat-space analysis of [1], in which the various ways to break spacetime symmetries while preserving spatial rotations and translations (both space and time) have been classified.

The structure of this paper is as follows. In Section 2 we review the results of [1] and discuss the coupling with gravity. In Section 3 we discuss which of the different symmetry breaking patterns admit a quasi de Sitter solution once coupled to gravity, and consequently build the unitary-gauge action. The main section of the paper is, then, Section 4: there we study what are the graviton interactions in the different scenarios and show that the symmetry breaking pattern of solids, of which Solid Inflation [27] is the lowest order in a derivative expansion, is the one that allows the most freedom. In this section we also discuss some phenomenological consequences of these interactions, and make a comparison to the symmetry breaking pattern of a superfluid. For simplicity, when discussing phenomenology we focus on quadratic and cubic graviton self-couplings up to cubic order in derivatives: for each of these we list all the corresponding operators in the unitary-gauge action. We conclude in Section 5.

Appendices A and B contain some details of the calculations carried out in Sections 3 and 4. More precisely, Appendix A shows how to construct the equivalent of the Effective Field Theory (EFT) of Inflation [28] for the case of broken spatial diffeomorphisms, with special emphasis on the property of “tadpole cancellation” (i.e. the fact that we want to write an EFT for fluctuations around a FLRW metric).

Summary of main results We summarize the main results of the paper in the compact table below. First, we list the different ways to break de Sitter boosts and what they correspond to once we make gravity dynamical. For each of them we show what are the available (\checkmark) building blocks in terms of the transverse and traceless graviton field γ_{ij} . These building blocks can be combined by contracting indices in an $SO(3)$ -invariant way. Given that for all the symmetry breaking patterns it is possible to take further time and spatial derivatives of the available building blocks, there is no loss of generality in stopping at the order in derivatives we show in the table.

| medium | broken gauge symmetry | γ_{ij} | $a^{-1}\partial_k\gamma_{ij}$ | $a^{-2}\partial_k\partial_l\gamma_{ij}$ | $\dot{\gamma}_{ij}$ |
|----------------|----------------------------|---------------|-------------------------------|---|---------------------|
| superfluid | time diff.s | \times | \times | \checkmark | \checkmark |
| type-I framid | local Lorentz boosts | \times | \times | \checkmark | \checkmark |
| type-II framid | local boosts and rotations | \times | \checkmark | \checkmark | \checkmark |
| solid | spatial diff.s | \checkmark | \checkmark | \checkmark | \checkmark |

2 “Zoology of condensed matter” and coupling with gravity

Let us review the analysis of [1]. We are interested in classifying all the symmetry breaking patterns that can be associated with a static, homogeneous, and isotropic medium in a

relativistic theory.

This implies that, together with the Poincaré generators P_μ, K_i, J_i of spacetime transformations, there is a set of translation and rotation generators \bar{P}_μ, \bar{J}_i that govern the excitations inside the medium and that leave the ground state invariant. These satisfy the same algebra as the Poincaré generators, i.e.

$$[\bar{J}_i, \bar{J}_j] = i\epsilon_{ijk}\bar{J}_k, \quad [\bar{J}_i, \bar{P}_j] = i\epsilon_{ijk}\bar{P}_k, \quad (2.1)$$

but they need not be the same as P_μ, J_i . Indeed, the different systems are classified by whether or not \bar{P}_μ, \bar{J}_i contain additional symmetries, generated by Q, Q_i, \tilde{Q}_i :

$$\bar{P}_0 = P_0 + Q, \quad \bar{P}_i = P_i + Q_i, \quad \bar{J}_i = J_i + \tilde{Q}_i. \quad (2.2)$$

2.1 The eightfold (sixfold) way

Ref. [1] identifies eight types of media, depending on which of the generators Q, Q_i, \tilde{Q}_i are non-vanishing. However, only six of them can be realized by having these generators to be internal symmetries: the remaining two require them to not commute with the Poincaré generators. These two media are dubbed “galileids”, since they are based on the galileon symmetry. Since, as the authors themselves argue, it is difficult to extend this symmetry to the case where gravity is dynamical, we will not consider them in the following.

Type-I framids: $\bar{P}_0 = P_0, \quad \bar{P}_i = P_i, \quad \bar{J}_i = J_i$

This is the simplest scenario, since it does not involve any internal symmetry. It can be realized by having a vector field that acquires a vacuum expectation value $\langle A^\mu(x) \rangle = \delta_0^\mu$. Ref. [1] shows that there are three Goldstone modes present: they can be thought as the rapidity of a boost acting on δ_0^μ .

Type-I superfluids: $\bar{P}_0 = P_0 + Q, \quad \bar{P}_i = P_i, \quad \bar{J}_i = J_i$

The minimal setup to realize the superfluid scenario is to have a “phase” field $\psi(x)$ with vacuum expectation value $\langle \psi(x) \rangle = t$. The Goldstone mode is the fluctuation of this phase field, $\psi(x) = t + \pi(x)$.

Type-II framids: $\bar{P}_0 = P_0, \quad \bar{P}_i = P_i, \quad \bar{J}_i = J_i + \tilde{Q}_i$

As we will see in the rest of the paper, this is an extension of the type-1 framid. It can be realized by a triplet of vector fields that rotate under an internal $SO(3)$ with vacuum expectation value $\langle A_i^\mu(x) \rangle = \delta_i^\mu$. There are six Goldstone modes, which can be identified with the rapidity and Euler angles of boosts and rotations acting on δ_i^μ .

Type-II superfluids: $\bar{P}_0 = P_0 + Q, \quad \bar{P}_i = P_i, \quad \bar{J}_i = J_i + \tilde{Q}_i$

This symmetry breaking pattern can be realized by a combination of a type-1 superfluid and a type-2 framid, with a total of seven Goldstone modes.

Solids and fluids: $\bar{P}_0 = P_0, \quad \bar{P}_i = P_i + Q_i, \quad \bar{J}_i = J_i$

This is the case of isotropic solids (that is, solids with no preferred axes, or “jellies”). The low-energy effective field theory can be characterized by a triplet of scalar fields $\phi^i(x)$ with vacuum expectation value $\langle \phi^i(x) \rangle = x^i$ (the Lagrangian and Eulerian coordinates of the volume elements of the solid, respectively). There are three Goldstone modes, $\phi^i(x) = x^i + \pi^i(x)$. Fluids are obtained by imposing a symmetry under volume-preserving internal diffeomorphisms.

Supersolids: $\bar{P}_0 = P_0 + Q, \quad \bar{P}_i = P_i + Q_i, \quad \bar{J}_i = J_i + \tilde{Q}_i$

Supersolids (with finite-temperature superfluids as a subset) are obtained by adding a type-I superfluid phase to the above case.

2.2 Coupling with gravity

Let us now discuss how to couple these systems with gravity. The precise derivation would involve the Callan-Coleman-Wess-Zumino (CCWZ) construction [29, 30, 31, 32], also known as coset construction, similarly to what has been done in [33] for Einstein gravity. Here we take a much faster route: we just make an educated guess on the final result based on [34, 35]. Indeed, these papers employ the CCWZ construction to couple to gravity two of the patterns of the previous section.

- Ref. [35] showed that the coset construction for the type-I superfluid gives the EFT of Inflation [28], in which time diffeomorphisms are broken.
- Ref. [34], instead, showed that the type-I framid results in a breaking of local Lorentz boosts, intended as transformations in the tangent space: for example, given the vierbein e_A^μ , objects like $\nabla_\mu e_0^\mu$ can now appear in the action (μ, ν, \dots and A, B, \dots are diffeomorphism indices and local Lorentz ones, respectively).

How to couple the remaining symmetry breaking patterns to gravity is then clear.

- The case of a generic fluid, and more generally of a solid, amounts to the breaking of spatial diffeomorphisms, i.e. to Solid Inflation [27] at lowest order in a derivative expansion.
- Type-II framids also involve the breaking of local Lorentz transformations. While in the case of the type-I framid only e_0^μ is allowed, now the dreibein e_I^μ ($I = 1, 2, 3$) is what can appear in the action.
- In both these cases, the fact that we want spatial and internal rotations to be broken to the diagonal subgroup amounts to contracting the indices I, J, K, \dots in an $SO(3)$ -invariant way.
- Finally, the finite-temperature superfluid results in the breaking of all diffeomorphisms, i.e. in Supersolid Inflation at lowest order in derivatives [36, 37, 38, 39], while type-II superfluids break time diffeomorphisms and local Lorentz transformations.

3 Inflationary solutions and unitary-gauge actions

We are now ready to construct the unitary-gauge action for the different symmetry breaking patterns. We focus on the cases of type-I superfluids, solids and framids only. As discussed at the end of the previous section, the remaining two patterns can be obtained straightforwardly from these four.

3.1 Type-I superfluids

The breaking of time diffeomorphisms results in the EFT of Inflation, for which we refer to the original paper [28]. The building blocks in this case are all diffeomorphism-invariant quantities constructed from the metric, together with all objects that can be constructed from $\partial_\mu t$, or equivalently the normal $n_\mu = -\partial_\mu t / \sqrt{-g^{00}}$ to the hypersurfaces of constant clock.

What is important to emphasize in this case is the following. The high degree of symmetry of the FLRW background ensures that is possible to obtain ‘‘tadpole cancellation’’ at all orders in derivatives. Let us consider the ‘‘slow-roll action’’, i.e. [28]

$$S = \int d^4x \sqrt{-g} \left\{ \frac{M_{\text{P}}^2}{2} R - c(t)g^{00} - \Lambda(t) \right\}. \quad (3.1)$$

At any order in derivatives one can show that additional operators can be split in a perturbation and a background part, without introducing new degrees of freedom, in such a way that the background part is always reabsorbed by the free functions c and Λ (possibly after integration by parts). The chief example is that of the extrinsic curvature of constant- t hypersurfaces, $K_{\mu\nu} = \delta K_{\mu\nu} + H h_{\mu\nu}$ (where $h_{\mu\nu} = g_{\mu\nu} + n_\mu n_\nu$).

3.2 Solids and fluids

When breaking spatial diffeomorphisms, the building blocks are again all diffeomorphism-invariant quantities constructed from the metric, together with all objects constructed from $\partial_\mu x^i$ (see also [40] for a discussion focused on the scalar and vector modes).

At leading order in derivatives, the action is

$$S = \int d^4x \sqrt{-g} \left\{ \frac{M_{\text{P}}^2}{2} R + F(X, Y, Z) \right\}, \quad (3.2)$$

where

$$X = g^{ii}, \quad Y = \frac{g^{ij} g^{ji}}{X^2}, \quad Z = \frac{g^{ij} g^{jk} g^{ki}}{X^3}. \quad (3.3)$$

Choosing F and $\partial F/\partial X$ appropriately, one can solve the equations of motion on the background for any value of the energy density and pressure. More precisely, quasi de Sitter expansion $\varepsilon = -\dot{H}/H^2$ can be achieved if

$$\varepsilon = \frac{\partial \ln F}{\partial \ln X} \ll 1, \quad (3.4)$$

and the role of the clock determining when inflation ends is played by $X = 3/a^2(t)$.

Compared to the EFT of Inflation, there are however some subtleties involved when we want to construct the action at higher order in derivatives.

- First, whenever we add new operators we want to make sure that we are not strongly breaking the de Sitter dilation symmetry. While in the EFT of Inflation this is automatically ensured if we use $\partial_\mu t$, which is invariant under $t \rightarrow t + c$, things are different if we use $\partial_\mu x^i$, which is not invariant under scale transformations $x^i \rightarrow \lambda x^i$. More prosaically, the diffeomorphism-breaking operators should be constructed from the unit-norm one-forms $\partial_\mu x^i/\sqrt{X}$, normal to the worldlines $x^i = \text{const.}$ of the volume elements of the solids.
- Second is the problem of tadpole subtraction. Once we include higher-derivative operators, it is not possible to reabsorb their background in the leading-order action of Eq. (3.2). Let us consider, for example, the equivalent of the extrinsic curvature. Given the worldlines $x^i = \text{const.}$, we can construct the unit vector parallel to them as [41]

$$O^\mu = \frac{e^{\mu\nu\rho\sigma} \epsilon_{ijk} \partial_\nu x^i \partial_\rho x^j \partial_\sigma x^k}{6\sqrt{\det(g^{mn})}}, \quad (3.5)$$

where $e_{\mu\nu\rho\sigma} = \sqrt{-g} \epsilon_{\mu\nu\rho\sigma}$ ($e^{\mu\nu\rho\sigma} = -\epsilon_{\mu\nu\rho\sigma}/\sqrt{-g}$) is the volume form. Given the projector $h_{\mu\nu} = g_{\mu\nu} + O_\mu O_\nu$, the tensor

$$\mathcal{K}_{\mu\nu} = h_{(\mu}{}^\rho \nabla_\rho O_{\nu)} \quad (3.6)$$

is then the equivalent of $K_{\mu\nu}$, to which it would reduce if $e^{\mu\nu\rho\sigma} O_\nu \nabla_\rho O_\sigma = 0$.¹ It is now impossible, however, to write its background without breaking time diffeomorphisms. One

¹We refer to [41] for a more detailed discussion about vorticity, which is outside of the scope of this work.

can construct an object starting at linear order in perturbations by taking the combination

$$\delta\mathcal{K}_{\mu\nu} = \mathcal{K}_{\mu\nu} - \frac{\nabla_\rho O^\rho}{3} h_{\mu\nu} . \quad (3.7)$$

However when we construct, for example, the operator $\mathcal{K}_{\mu\nu}\mathcal{K}^{\mu\nu}$, we run into the issue that the operator $(\nabla_\rho O^\rho)^2$ is not contained in the action of Eq. (3.2). This way of phrasing the problem suggests its solution: it should be enough to add a dependence of F on a single additional operator, $\nabla_\rho O^\rho$.

While the problem of tadpole subtraction is immaterial if one wants to study graviton interactions only, as we will see in detail in Section 4, it is still worth to keep it in mind as a difference between this symmetry breaking pattern and the EFT of Inflation.

3.3 Type-II framid (\supset type-I framid)

Let us first focus on the case of type-II framids. As it is easy to imagine, the fact that we do not have a clock tells us that we cannot write an effective theory for perturbations around a generic FLRW spacetime.

Thus, if we wanted to use this symmetry breaking pattern as a legit inflationary model, we should add new degrees of freedom to exit inflation (the simplest thing to do is to add a clock, which would result in the symmetry breaking pattern of type-II superfluids). Let us nevertheless press forward and see how one can obtain a pure de Sitter solution, since this allows us to introduce the geometric objects that will be used in Section 4.

The quickest way to show this is by looking at the different operators one can write down, order-by-order in derivatives. First, given a dreibein e_I^μ , we can immediately obtain the unit vector orthogonal to e_I^μ via the volume form as

$$e_0^\mu = -\frac{g^{\mu\nu}}{6}\epsilon_{IJK}e_{\nu\alpha\beta\gamma}e_I^\alpha e_J^\beta e_K^\gamma . \quad (3.8)$$

Importantly, this unit vector is not only orthogonal to e_I^μ , but one can show that if we define the tetrad as $e_A^\mu = \{e_0^\mu, e_I^\mu\}$, and transform e_I^μ as $e_I'^\mu = \Lambda_I^B e_B^\mu$ (where Λ_A^B is a local Lorentz transformation), we have that

$$e_0'^\mu = -\frac{g^{\mu\nu}}{6}\epsilon_{IJK}e_{\nu\alpha\beta\gamma}e_I'^\alpha e_J'^\beta e_K'^\gamma = \Lambda_0^A e_A^\mu . \quad (3.9)$$

What operators can we construct from the tetrad e_A^μ , then?

- At zeroth order in derivatives, the only non-gauge-invariant quantity we can use is e_A^μ itself. Since greek indices must be contracted, the only object we can use is $g_{\mu\nu}e_A^\mu e_B^\nu = \eta_{AB}$, where tetrad indices must be contracted in an $SO(3)$ -invariant way. These only modify the cosmological constant.
- At first order in derivatives we have four additional building blocks. These are the tensors

$$e_0^\nu \nabla_\nu e_0^\mu , \quad e_0^\nu \nabla_\nu e_J^\mu , \quad e_I^\nu \nabla_\nu e_0^\mu , \quad e_I^\nu \nabla_\nu e_J^\mu . \quad (3.10)$$

It is straightforward to see that around a de Sitter spacetime we can subtract the background of these tensors without introducing new degrees of freedom. More precisely, defining the dual one-forms

$$\omega_\mu^A = \eta_{AB} g_{\mu\nu} e_B^\nu , \quad (3.11)$$

we can isolate the fluctuations as

$$\delta(e_I^\nu \nabla_\nu e_0^\mu) = e_I^\nu \nabla_\nu e_0^\mu - H e_I^\mu, \quad (3.12a)$$

$$\delta(e_I^\nu \nabla_\nu e_J^\mu) = e_I^\nu \nabla_\nu e_J^\mu - H \delta_{IJ} e_0^\mu \quad (3.12b)$$

and

$$\delta(\nabla_\nu e_0^\mu) = \nabla_\nu e_0^\mu - H(\delta_\nu^\mu - \omega_\nu^0 e_0^\mu). \quad (3.13)$$

These imply that $e_0^\nu \nabla_\nu e_0^\mu$ vanishes for a de Sitter metric, as does $e_0^\nu \nabla_\nu e_J^\mu$.

- At higher order in derivatives, via the product rule we can always rewrite the background of non-gauge-invariant tensors in terms of Eqs. (3.12), (3.13).

One can also check how these objects transform under the de Sitter dilation symmetry. It is easy to see that, in de Sitter spacetime, the Lie derivative of e_I^μ along the vector field $\delta_0^\mu - H x^i \delta_i^\mu$ is equal to zero.

What about type-I framids? Given Eqs. (3.8), (3.9), we see that this symmetry breaking pattern is a subset of the type-II framid: for this reason we will not discuss them further. It is interesting to notice that if we require a symmetry under the global transformation

$$e_I^\mu \rightarrow M_I^J e_J^\mu, \quad (3.14)$$

with M_I^J a three-dimensional matrix having unit determinant, it seems we can single out e_0^μ as the only vector from which to construct non-gauge-invariant operators. However this works only at leading (zeroth) order in derivatives. Given that finding such symmetry is not necessary for the purposes of this paper, we leave this question to future work.

3.4 A word on the Goldstone modes

Before proceeding, we find it useful to quickly summarize how the Stueckelberg trick to reintroduce the Goldstone modes works for these different symmetry breaking patterns.

- In the superfluid case we are breaking time diffeomorphisms. The Stueckelberg trick, then, simply amounts to performing a time diffeomorphism $t \rightarrow t + \pi$, as explained in detail in [28]. After the Stueckelberg trick, the “flat-gauge metric” can be written in the usual ADM decomposition, with a spatial part that only contains the graviton (i.e. no vector or scalar modes).
- In the case of a solid we are breaking spatial diffeomorphisms. The Stueckelberg trick is a spatial diffeomorphism $x^i \rightarrow x^i + \pi^i$, where π^i can be decomposed in a scalar and a vector part. Again, the spatial metric only contains the graviton after performing the Stueckelberg trick.
- In the case of framids the Stueckelberg trick consists in doing a local Lorentz transformation. As we have already discussed in Section 2.1, the parameters of this transformation are the rapidity and the three Euler angles, i.e. we have six degrees of freedom (two scalars and two vectors). Also in this case the metric after the Stueckelberg trick can be decomposed in ADM variables, with only the graviton present in g_{ij} (see e.g. [34] for the case of type-I framids).

Importantly, in none of the symmetry breaking patterns we have additional spin-2 degrees of freedom besides the graviton.

4 Main results

In this section we collect the main results of this paper. First, we show that the solid is the symmetry breaking that allows the most freedom for graviton interactions (Section 4.1). Then, we study some phenomenological consequences of these interactions (Section 4.2).

4.1 Most general graviton interactions

The best way to see what is the symmetry breaking pattern that allows the most freedom is working at the level of linear fluctuations. That is, we work directly with the graviton field γ_{ij} , and check what are the “minimal” building blocks in the different symmetry breaking patterns.

Since we only focus on the graviton, we can take the line element to be

$$ds^2 = -dt^2 + a^2(e^\gamma)_{ij}dx^i dx^j, \quad (4.1)$$

that is we set the lapse and shift variables to zero, together with the scalar and vector modes that are introduced by making broken diffeomorphisms or broken local Lorentz transformations (see Section 3.4 for details).² From this, we can write the tetrad as

$$e_0^\mu = \delta_0^\mu, \quad e_I^0 = 0, \quad e_I^i = a^{-1}(e^{-\frac{\gamma}{2}})_{iI}. \quad (4.2)$$

Type-I superfluids Let us start from looking at the building blocks for the EFT of Inflation. At linear order, γ_{ij} can only appear with derivatives: time derivatives enter as $\delta K_i^j = \dot{\gamma}_{ij}/2$, while spatial derivatives $a^{-2}\partial_k\partial_l\gamma_{ij}$ enter via ${}^{(3)}R_{kl}^{ij}$, the Riemann curvature tensor on constant-time hypersurfaces. More precisely, $a^{-2}\partial_k\partial_l\gamma_{ij}$ can only appear in the combination

$${}^{(3)}R_{kl}^{ij} = a^{-2}(\partial_k\partial_{(i}\gamma_{j)l} - \partial_l\partial_{(i}\gamma_{j)k}). \quad (4.3)$$

It is then possible to take any number of spatial and time derivatives of these building blocks by using the covariant derivative D_μ on the $t = \text{const.}$ hypersurfaces and $n^\mu\nabla_\mu$, respectively.

Type-II framids We then move to the type-II framid. First, we see that using e_0^μ allows us to reproduce all the geometric objects of the EFT of Inflation. Then, given that at lowest order in derivatives we cannot write any non-gauge-invariant object – we always end up with a cosmological constant – it is still impossible to write γ_{ij} without derivatives. Unlike the case of broken time diffeomorphisms, however, it is now possible to use $a^{-1}\partial_k\gamma_{ij}$ as a building block. More precisely, we can use the combination

$$-\omega_\mu^K \delta(e_J^\nu \nabla_\nu e_I^\mu) = -\omega_\mu^K e_J^\nu \nabla_\nu e_I^\mu = a^{-1}\partial_{[K}\gamma_{I]J}. \quad (4.4)$$

Here we notice the overall factor of a^{-1} , that we expect from the discussion on invariance under de Sitter dilations of Section 3.3.

Solids When we break spatial diffeomorphisms we can still reproduce all the operators of the EFT of Inflation. This is achieved by using the vector O^μ defined in Eq. (3.5). Let us go back to the problem of the cancellation of tadpoles. We know that in the EFT of Inflation the background of all operators can be constructed from the normal vector to the constant-time hypersurfaces (whose role here is played by O^μ), and the Hubble rate with its time derivatives.

²When looking at interactions higher than cubic one also needs to include the constraints (which otherwise need only to be solved at linear order [22], and consequently do not contain γ_{ij}). These terms do not change the conclusions of this paper, so we leave their detailed study to future work.

We have seen, e.g. in Eq. (3.7), that the four-divergence of O^μ can play the role of the Hubble rate. We can then use $O^\mu \nabla_\mu$ to take time derivatives of this proxy for H . There is no loss of generality in doing this if we can show that these objects do not contain the graviton. Luckily, this is easy to prove. From the definition of Eq. (3.5), we see that O^μ does not contain γ_{ij} since neither $e^{\mu\nu\rho\sigma}$ nor $\sqrt{\det(g^{mn})}$ do. Using the formula

$$\nabla_\mu O^\mu = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} O^\mu) \quad (4.5)$$

we see that its four-divergence also does not contain the graviton. Then, given that to take its time derivatives we are only acting with $O^\mu \nabla_\mu$ on a scalar, we can replace ∇_μ with ∂_μ (which also does not contain γ_{ij}).

What about spatial derivatives? It is straightforward to see that we can use directly $a^{-1} \partial_k \gamma_{ij}$ (where, unlike Eq. (4.4), we do not need to take any antisymmetrization over k and j) by considering

$$\Gamma^{kij} \equiv -\frac{g^{\mu k} g^{\nu(i} \nabla_\mu \nabla_{\nu} x^j)}{(X/3)^{3/2}}, \quad (4.6)$$

a quantity that starts linear in perturbations. As discussed in Section 3.2, we divide by the appropriate power of X to ensure invariance under spatial dilations. This in turn ensures that we reproduce the expected power of the scale factor when we have one spatial derivative acting on γ_{ij} , as in Eq. (4.4). Indeed, we find

$$\frac{1}{2a} \partial_k \gamma_{ij} = \Gamma^{kij}. \quad (4.7)$$

Finally, there is one more additional freedom with respect to the case of type-II framids. While there we can only contract spatial indices via the Kronecker delta, here we can use γ_{ij} itself through

$$\Gamma^{ij} \equiv \delta_{ij} - \frac{3g^{ij}}{X}, \quad (4.8)$$

which is equal to γ_{ij} at linear order in perturbations.

4.2 A bit of phenomenology

Let us then discuss the phenomenology of the solid symmetry breaking pattern. Given the large amount of freedom we have, characterizing all the signatures is a complicated task. For this reason, we think it is more useful to compare with the EFT of Inflation.

While we have shown that the symmetry breaking pattern of a solid captures the most general graviton interactions at *all* orders in perturbations and derivatives, when discussing the phenomenology it is useful to work in a derivative expansion. We stop at second order in this expansion. Moreover, we focus on operators that start from cubic order in perturbations at most, i.e. we look at the tree-level graviton bispectrum.

At lowest (zeroth) order in derivatives, we have a small mass of order εH^2 for the graviton, and a cubic vertex $\gamma_{ij} \gamma_{jk} \gamma_{ki}$ that instead is not slow-roll-suppressed (see [27, 42], for example: we review this in Appendix B). The latter has been shown in [42] to give a graviton bispectrum that peaks in the squeezed limit, and that does not asymptote a constant at late times even at leading order in ε (at this order we can use de Sitter modes to compute the in-in integral: given that the interaction is purely local, it is easy to see that there are divergences of the form $\ln(-k\eta)$, where η is the conformal time $-1/aH$).

Let us go to second order in derivatives (i.e. consider the new operators discussed in Section 4.1). At quadratic order in perturbations, we can look at the different ways in which the speed of sound of the graviton can be modified. Besides the operators of the EFT of Inflation, like the Ricci scalar on the constant-time hypersurfaces ${}^{(3)}R$ and the square of the extrinsic curvature $\delta\mathcal{K}_{\mu\nu}\delta\mathcal{K}^{\mu\nu}$, there are many new operators that reduce to $\dot{\gamma}_{ij}^2$ and $a^{-2}(\partial_k\gamma_{ij})^2$ at quadratic order in γ_{ij} , but differ from the EFT of Inflation ones at cubic order (and higher). Two simple examples are

$$(O^\mu\nabla_\mu\Gamma^{ij})(O^\nu\nabla_\nu\Gamma^{ij}), \quad \Gamma^{kij}\Gamma^{kij}. \quad (4.9)$$

An operator that instead is essentially equivalent to the three-Ricci scalar is the one studied in [37], i.e. G^{ii} , where $G_{\mu\nu}$ is the four-dimensional Einstein tensor (indeed, Ref. [37] shows that the resulting graviton bispectrum has the same shape as that from Einstein gravity).

A phenomenologically interesting observation comes when we turn from the quadratic action to the interactions. One of the main points of [23] is that by appropriately redefining the lightcone and consequently putting the tensor sound speed to 1, one also removes completely all graviton self-interactions beyond the ones from Einstein gravity. This relies on the fact that there are only two operators to remove, ${}^{(3)}R$ and $\delta\mathcal{K}_{\mu\nu}\delta\mathcal{K}^{\mu\nu}$, and two free coefficients in the lightcone redefinition (those of a conformal and disformal transformation of the metric, see also [24] for details). Now the situation is different simply because we have more operators to play with, but the same number of free coefficients in the lightcone redefinition.³ This tells us that, while there is no loss of generality in putting the tensor speed of sound to 1, doing so does *not* remove also the interaction terms.

Another important phenomenological difference with the EFT of Inflation is the following. As we have seen, all interactions in that case come from operators starting quadratic in γ_{ij} : there are no operators starting at cubic order in γ_{ij} at second order in derivatives. If we break spatial diffeomorphisms the situation changes. For example, we can combine Eq. (4.8) with Eq. (4.9) to construct

$$\gamma_{jl}\dot{\gamma}_{ij}\dot{\gamma}_{il} \quad \text{and} \quad a^{-2}\gamma_{kl}\partial_k\gamma_{ij}\partial_l\gamma_{ij}, \quad (4.10)$$

where we notice that the second structure appears directly in the Einstein-Hilbert action (it is contained in ${}^{(3)}R$ together with $a^{-2}\gamma_{kl}\partial_k\gamma_{ij}\partial_j\gamma_{il}$), while the first one does not. This confirms that, if spatial diffeomorphisms are broken, the three-point function of the graviton is not controlled only by coefficients entering in the graviton power spectrum. This is already apparent at the zeroth order in derivatives (for example, we can see this by comparing Eq. (B.3) with Eq. (B.4) in Appendix B): here we arrive at the same conclusion at second order.

This is a fast way to see that the consistency relations are not respected if we break spatial diffeomorphisms. Showing that the bispectrum depends on additional free parameters that are absent from the power spectrum bypasses the need to check any soft limit (we refer to [42, 43, 44] for a discussion of soft limits involving scalar and vector modes in Solid Inflation).

It is straightforward to compute the graviton three-point function from the operators of Eq. (4.10). Using the standard Fourier decomposition for the graviton,⁴ together with the

³Indeed, at this order in derivatives and perturbations the only objects we can use are the vector O^μ and the normalized one-forms $\partial_\mu x^i/\sqrt{X}$. Because we need to contract latin indices in an $SO(3)$ -invariant way, we can only use the combination $\partial_\mu x^i\partial_\nu x^i/X$, which is essentially equivalent to $g_{\mu\nu} + O_\mu O_\nu$.

⁴This is

$$\gamma_{ij}(t, \mathbf{x}) = \int \frac{d^3k}{(2\pi)^3} \sum_s \epsilon_{ij}^s(\mathbf{k}) \gamma_{\mathbf{k}}^s(t) e^{i\mathbf{k}\cdot\mathbf{x}}, \quad (4.11)$$

where the traceless polarization tensors ϵ_{ij}^s satisfy $k^i\epsilon_{ij}^s(\mathbf{k}) = 0$.

three elementary symmetric polynomials

$$k_T = e_1 = k_1 + k_2 + k_3 , \quad (4.12a)$$

$$e_2 = k_1 k_2 + k_2 k_3 + k_3 k_1 , \quad (4.12b)$$

$$e_3 = k_1 k_2 k_3 , \quad (4.12c)$$

we find

$$\langle \gamma_{\mathbf{k}_1}^{s_1} \gamma_{\mathbf{k}_2}^{s_2} \gamma_{\mathbf{k}_3}^{s_3} \rangle'_{\gamma \dot{\gamma} \dot{\gamma}} = \frac{H^4 M^2}{2M_{\text{P}}^6} \frac{k_T e_2^2 + e_2 e_3 - 2k_T^2 e_3}{e_3^3 k_T^2} \epsilon_{ij}^{s_1}(-\mathbf{k}_1) \epsilon_{jk}^{s_2}(-\mathbf{k}_2) \epsilon_{ki}^{s_3}(-\mathbf{k}_3) , \quad (4.13a)$$

$$\langle \gamma_{\mathbf{k}_1}^{s_1} \gamma_{\mathbf{k}_2}^{s_2} \gamma_{\mathbf{k}_3}^{s_3} \rangle'_{\gamma \partial \gamma \partial \gamma} = \frac{H^4 M^2}{2M_{\text{P}}^6} \frac{k_T e_2 + e_3 - k_T^3}{e_3^3 k_T^2} \epsilon_{il}^{s_1}(-\mathbf{k}_1) \epsilon_{jm}^{s_2}(-\mathbf{k}_2) \epsilon_{kn}^{s_3}(-\mathbf{k}_3) T_{ijk}^{lmn}(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3) , \quad (4.13b)$$

with

$$T_{ijk}^{lmn}(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3) = k_2^i k_3^l \delta_{jk} \delta_{mn} + k_3^j k_1^m \delta_{ki} \delta_{nl} + k_1^k k_2^n \delta_{ij} \delta_{lm} . \quad (4.14)$$

Here M^2 is a mass scale that makes the operators have mass dimension 4. We have used de Sitter mode functions to compute these bispectra: given that the mass of the graviton is slow-roll-suppressed, this is correct at leading order in ε .

A final observation that is worth making is the following. If we break parity, in the EFT of Inflation it is only possible to modify the quadratic action of the graviton starting from third order in derivatives, via operators that reduce to

$$a^{-1} \epsilon_{ijk} \partial_i \dot{\gamma}_{jl} \dot{\gamma}_{lk} \quad \text{and} \quad a^{-3} \epsilon_{ijk} \partial_i \partial_l \gamma_{jm} \partial_l \gamma_{mk} \quad (4.15)$$

at quadratic order in γ_{ij} . Now it is possible to write operators both at first and second order in derivatives, via

$$\epsilon_{ijk} \Gamma^{il} \Gamma^{jlk} = a^{-1} \epsilon_{ijk} \gamma_{il} \partial_j \gamma_{lk} , \quad (4.16a)$$

$$\epsilon_{ijk} \Gamma^{jlk} O^\mu \nabla_\mu \Gamma^{il} = a^{-1} \epsilon_{ijk} \dot{\gamma}_{il} \partial_j \gamma_{lk} . \quad (4.16b)$$

These two operators split the graviton helicities. However, one can easily see that they lead to serious instabilities on subhorizon scales. Given the circularly-polarized polarization tensors $\epsilon_{ij}^\pm(\mathbf{k})$, and using the property $ik^l \epsilon_{jlm} \epsilon_{im}^s(\mathbf{k}) = \lambda_s k \epsilon_{ij}^s(\mathbf{k})$, $\lambda_\pm = \pm 1$, we see that the contribution of these two operators to the dispersion relation of the two helicities is

$$\omega^2 = k^2 + \lambda_s M k , \quad (4.17a)$$

$$\omega^2 = k^2 + ic \lambda_s \omega k . \quad (4.17b)$$

Here M and c (of mass dimension 1 and 0 respectively) are the coefficients controlling the operators of Eq. (4.16), we have neglected the mass of the graviton since it is of order εH^2 , and we assume that $M \gg H$. We see that the second operator essentially gives a complex speed of sound to the graviton (one can see this in the limit $c \ll 1$), while the first operator makes the frequency of one of the two helicities purely imaginary for $k \ll M$. In summary, these operators lead to instabilities on short scales, and it is therefore unclear how to make sense of the Bunch-Davies vacuum if we turn them on. This is also why we do not study parity-breaking operators at higher order in perturbations: it does not make much sense to do so while putting to zero the operators of Eqs. (4.16), since the cubic and higher-order operators could generate these via loop effects.⁵ We leave a more detailed investigation of this issue for a future work.

⁵Indeed, there is no symmetry besides parity that is recovered if we put M or c to zero.

5 Conclusions

In this paper we have studied graviton non-Gaussianities in models of the early universe in which de Sitter boosts are spontaneously broken, showing that the case where the epoch of de Sitter expansion is driven by a solid is the one with the more freedom for graviton self-interactions.

This freedom makes systematically studying all the possible correlation functions very complicated, and we expect that a full characterization of non-Gaussianities involving also the Goldstone modes (a scalar and a vector for a solid) to be even more difficult. For this reason it would be interesting to extend the approach of the “Boostless Bootstrap” recently put forward in [21] to this symmetry breaking pattern. The well-known fact that cosmological perturbations are not conserved in Solid Inflation, even at leading order in the slow-roll expansion (for the graviton this is exemplified by the unsuppressed cubic interaction $\propto \gamma_{ij}\gamma_{jk}\gamma_{ki}$, and see e.g. [45, 46, 47, 48, 49, 50, 51, 52] for studies of mixed three-point functions involving curvature perturbations), makes it difficult to set up a theory of correlators on the boundary of de Sitter: we leave investigations of such issues to future work.

Finally, we emphasize that in this work we have focused on *isotropic* solids, i.e. where we have invariance under the full $SO(3)$ group. Refs. [53, 54, 55] study what happens if this assumption is relaxed to invariance under a discrete subgroup of rotations. It would be interesting to study how our results generalize to their setup.

Acknowledgements

It is a pleasure to thank Misha Ivanov, Mehrdad Mirbabayi, Enrico Pajer, Gianmassimo Tasinato, Matias Zaldarriaga, and especially Sadra Jazayeri for useful discussions. I also thank Mehrdad Mirbabayi and Enrico Pajer for very useful comments on the draft. I acknowledge support from the Institute for Advanced Study.

A An “EFT of Inflation” for solids

As emphasized in the main text, and made clear in the original paper [28], a key ingredient of the EFT of Inflation is the tadpole cancellation. This is what allows to write an effective theory for metric fluctuations, which are the quantities we constrain with late-time cosmological observations, dependent on free EFT coefficients *and* the expansion history during inflation. Isolating all the non-gauge-invariant operators, as done e.g. in [40] (and as we have also done in Sections 3.2 and 4.1), is not enough.

Can we achieve this in the case of broken spatial diffeomorphisms? We have seen in Section 3.2 that all the geometric objects constructed from O^μ can have their background subtracted by using suitable powers of $\theta \equiv \nabla_\mu O^\mu$. While this is correct in spirit, we need to be more precise and characterize the operators at all orders in derivatives.

Zeroth order in derivatives At this order we have Solid Inflation. Using the fact that the derivative of Y and Z with respect to $g^{\mu\nu}$ vanishes for an FLRW metric, there is no loss of generality if we rewrite the non-Einstein-Hilbert part of Eq. (3.2) as

$$\int d^4x \sqrt{-g} \{ \mathcal{L}_0(X) + M^4(X, \delta Y, \delta Z) \} , \quad (\text{A.1})$$

where we have defined $\delta Y \equiv Y - 1/3$, $\delta Z \equiv Z - 1/9$. The form of \mathcal{L}_0 is fixed at any given order in slow-roll parameters by solving the background Einstein equations. They take the form

$$3M_{\text{p}}^2 H^2 = -\mathcal{L}_0, \quad (\text{A.2a})$$

$$-\frac{\dot{H}}{H^2} = \frac{d \ln \mathcal{L}_0}{d \ln X}. \quad (\text{A.2b})$$

Then, we have in mind that the function M^4 is expanded in powers of δY and δZ , with coefficients depending on X . This dependence should be mild if we want the de Sitter dilation symmetry to be softly broken. This expansion is similar to that of the EFT of Inflation in powers of $g^{00} + 1$, with EFT coefficients that depend only mildly on time. From now on we do not worry about M^4 or any other terms in the Lagrangian that start at second order in perturbations.

Up to second order in derivatives Let us see how at this order we can get away with just adding a dependence of \mathcal{L}_0 on θ . The only operators that are not accounted are those of the form $O^\mu \nabla_\mu \theta$ times a function of X . We can integrate these by parts, and then use the fact that we can rewrite $O^\mu \nabla_\mu X$ as a function of X times θ^2 plus terms starting quadratic in perturbations. What do the equations of motion look like? Using the relation (valid for any vector V^μ)

$$\nabla_\mu V^\mu = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} V^\mu), \quad (\text{A.3})$$

together with Eq. (3.5), we have

$$\delta\theta = \frac{1}{2} \theta g_{\mu\nu} \delta g^{\mu\nu} - \frac{1}{2} \nabla_\mu (O^\mu g_{ij} \delta g^{ij}). \quad (\text{A.4})$$

Correspondingly, Eqs. (A.2) become

$$3M_{\text{p}}^2 H^2 = \frac{\partial \mathcal{L}_0}{\partial \ln \theta} - \mathcal{L}_0, \quad (\text{A.5a})$$

$$-\frac{\dot{H}}{H^2} = \frac{\partial \ln \mathcal{L}_0}{\partial \ln X} \left(1 - \frac{\partial \ln \mathcal{L}_0}{\partial \ln \theta} \right)^{-1} \left(1 - \frac{3}{2} \frac{(\mathcal{L}_0)_{\theta\theta}}{M_{\text{p}}^2} \right)^{-1}, \quad (\text{A.5b})$$

where $(\mathcal{L}_0)_{\theta\theta}$ is the second derivative of \mathcal{L}_0 with respect to θ . These relations are used in the same way as in Solid Inflation: one expands \mathcal{L}_0 in perturbations around an FLRW metric, and the derivatives with respect to X and θ can be replaced by the expansion history using Eqs. (A.5).⁶

Up to fourth order in derivatives At this order we cannot get away anymore with just adding a dependence on θ . This is exemplified by the operator

$$(O^\rho \nabla_\rho \mathcal{K}_{\mu\nu})(O^\sigma \nabla_\sigma \mathcal{K}^{\mu\nu}), \quad (\text{A.7})$$

whose background contains $(O^\mu \nabla_\mu \theta)^2$. Still, it is enough to include a dependence on the operator $\dot{\theta} \equiv O^\mu \nabla_\mu \theta$ (and on this operator only) in \mathcal{L}_0 .

⁶Higher derivatives with respect to these variables can be computed recursively. For example, at zeroth order in derivatives (i.e. in Solid Inflation) one has

$$\frac{X^2 (\mathcal{L}_0)_{XX}}{X (\mathcal{L}_0)_X} = -1 + \varepsilon - \frac{\eta}{2}. \quad (\text{A.6})$$

Higher orders The same happens at higher orders. We always need to add a finite number of operators to \mathcal{L}_0 at any order in derivatives: these are essentially “time derivatives” of θ . The tadpole cancellation can be implemented in a similar way as the procedure leading to Eqs. (A.5).

In this way we can write an “EFT of Inflation” for solids, with all the good properties of the one for broken time diffeomorphisms. More precisely, we write

$$S = S_{\text{EH}} + \underbrace{\int d^4x \sqrt{-g} \mathcal{L}_0(X, \theta, \dot{\theta}, \dots)}_{\equiv S_0} + \text{EFT operators} , \quad (\text{A.8})$$

where the “EFT operators” start at second or higher order in perturbations around FLRW. One can turn on a time dependence of the coefficients of these operators by making them dependent on X , similarly to how the coefficients in the EFT of Inflation are allowed to be dependent on time. In S_0 we keep a finite number of derivatives of θ depending on the order in derivatives at which we stop in the overall action. The only complication is that the background equations, which determine the form of S_0 , must be solved every time. This is only a slight complication given the very simple form of the functional derivatives of $X, \theta, \dot{\theta}, \dots$ around a metric as symmetric as FLRW.

It would be interesting to study how the additional operators $\theta, \dot{\theta}, \dots$ present in S_0 affects the dynamics of the Goldstone modes. This is equivalent in studying the predictions of the “slow-roll action” of the EFT of Inflation. Unlike what happens here, in the EFT of Inflation S_0 is fixed at all orders in derivatives: it is just Eq. (3.1) with $c = -M_{\text{P}}^2 \dot{H}$ and $\Lambda = M_{\text{P}}^2(3H^2 + \dot{H})$. We leave this to future work.

B Cubic graviton interaction in Solid Inflation

Let us quickly show that the cubic graviton action at zeroth order in derivatives is not suppressed by ε . Expanding Eq. (3.2) in perturbations, one finds (using the notation $\partial F/\partial X = F_X$, $\partial F/\partial Y = F_Y$, $\partial F/\partial Z = F_Z$ for compactness and dropping integral signs)

$$\frac{S_{\gamma\gamma}}{a^3} \supset \left(\frac{1}{6} \frac{\partial F}{\partial \ln X} + \frac{1}{9} F_Y + \frac{1}{9} F_Z \right) \gamma_{ij} \gamma_{ij} , \quad (\text{B.1a})$$

$$\frac{S_{\gamma\gamma\gamma}}{a^3} \supset - \left(\frac{1}{18} \frac{\partial F}{\partial \ln X} + \frac{1}{9} F_Y + \frac{4}{27} F_Z \right) \gamma_{ij} \gamma_{jk} \gamma_{ki} . \quad (\text{B.1b})$$

Using Eq. (3.4) and the relation $3M_{\text{P}}^2 H^2 = -F$, together with the fact that the propagation speeds c_{L}^2 and c_{T}^2 of the longitudinal and transverse part of π^i are [27]

$$c_{\text{L}}^2 = \frac{1}{3} + \frac{8}{9} \frac{F_Y + F_Z}{X F_X} , \quad (\text{B.2a})$$

$$c_{\text{T}}^2 = 1 + \frac{2}{3} \frac{F_Y + F_Z}{X F_X} , \quad (\text{B.2b})$$

we see that Eq. (B.1a) is a small mass

$$m_{\gamma}^2 = -4\dot{H}c_{\text{T}}^2 , \quad (\text{B.3})$$

while Eq. (B.1b) gives

$$\frac{S_{\gamma\gamma\gamma}}{M_{\text{P}}^2 a^3} \supset -\frac{1}{9} \frac{F_Y}{F} H^2 \gamma_{ij} \gamma_{jk} \gamma_{ki} + \frac{1}{2} \varepsilon H^2 c_{\text{L}}^2 \gamma_{ij} \gamma_{jk} \gamma_{ki} . \quad (\text{B.4})$$

The first term here reproduces Eq. (A.6) of [42].

References

- [1] A. Nicolis, R. Penco, F. Piazza, and R. Rattazzi, “Zoology of condensed matter: Framids, ordinary stuff, extra-ordinary stuff,” [JHEP 06 \(2015\) 155](#), [arXiv:1501.03845 \[hep-th\]](#).
- [2] T. Matsumura et al., “Mission design of LiteBIRD,” [arXiv:1311.2847 \[astro-ph.IM\]](#). [*J. Low. Temp. Phys.*176,733(2014)].
- [3] **CMB-S4** Collaboration, K. N. Abazajian et al., “CMB-S4 Science Book, First Edition,” [arXiv:1610.02743 \[astro-ph.CO\]](#).
- [4] **CORE** Collaboration, F. Finelli et al., “Exploring cosmic origins with CORE: Inflation,” [JCAP 1804 \(2018\) 016](#), [arXiv:1612.08270 \[astro-ph.CO\]](#).
- [5] **CORE** Collaboration, J. Delabrouille et al., “Exploring cosmic origins with CORE: Survey requirements and mission design,” [JCAP 1804 no. 04, \(2018\) 014](#), [arXiv:1706.04516 \[astro-ph.IM\]](#).
- [6] **Simons Observatory** Collaboration, P. Ade et al., “The Simons Observatory: Science goals and forecasts,” [JCAP 1902 \(2019\) 056](#), [arXiv:1808.07445 \[astro-ph.CO\]](#).
- [7] **NASA PICO** Collaboration, S. Hanany et al., “PICO: Probe of Inflation and Cosmic Origins,” [arXiv:1902.10541 \[astro-ph.IM\]](#).
- [8] S. Shandera et al., “Probing the origin of our Universe through cosmic microwave background constraints on gravitational waves,” [Bull. Am. Astron. Soc. 51 \(2019\) 338](#), [arXiv:1903.04700 \[astro-ph.CO\]](#).
- [9] K. Abazajian et al., “CMB-S4 Science Case, Reference Design, and Project Plan,” [arXiv:1907.04473 \[astro-ph.IM\]](#).
- [10] **LiteBIRD** Collaboration, M. Hazumi et al., “LiteBIRD: JAXA’s new strategic L-class mission for all-sky surveys of cosmic microwave background polarization,” [Proc. SPIE Int. Soc. Opt. Eng. 11443 \(2020\) 114432F](#), [arXiv:2101.12449 \[astro-ph.IM\]](#).
- [11] J. M. Maldacena and G. L. Pimentel, “On graviton non-Gaussianities during inflation,” [JHEP 09 \(2011\) 045](#), [arXiv:1104.2846 \[hep-th\]](#).
- [12] I. Mata, S. Raju, and S. Trivedi, “CMB from CFT,” [JHEP 07 \(2013\) 015](#), [arXiv:1211.5482 \[hep-th\]](#).
- [13] A. Ghosh, N. Kundu, S. Raju, and S. P. Trivedi, “Conformal Invariance and the Four Point Scalar Correlator in Slow-Roll Inflation,” [JHEP 07 \(2014\) 011](#), [arXiv:1401.1426 \[hep-th\]](#).
- [14] E. Pajer, G. L. Pimentel, and J. V. S. Van Wijck, “The Conformal Limit of Inflation in the Era of CMB Polarimetry,” [JCAP 1706 no. 06, \(2017\) 009](#), [arXiv:1609.06993 \[hep-th\]](#).
- [15] N. Arkani-Hamed, D. Baumann, H. Lee, and G. L. Pimentel, “The Cosmological Bootstrap: Inflationary Correlators from Symmetries and Singularities,” [JHEP 04 \(2020\) 105](#), [arXiv:1811.00024 \[hep-th\]](#).
- [16] D. Green and E. Pajer, “On the Symmetries of Cosmological Perturbations,” [JCAP 09 \(2020\) 032](#), [arXiv:2004.09587 \[hep-th\]](#).
- [17] D. Baumann, C. Duaso Pueyo, A. Joyce, H. Lee, and G. L. Pimentel, “The Cosmological Bootstrap: Spinning Correlators from Symmetries and Factorization,” [arXiv:2005.04234 \[hep-th\]](#).

- [18] E. Pajer, D. Stefanyshyn, and J. Supel, “The Boostless Bootstrap: Amplitudes without Lorentz boosts,” [arXiv:2007.00027 \[hep-th\]](#).
- [19] H. Goodhew, S. Jazayeri, and E. Pajer, “The Cosmological Optical Theorem,” [arXiv:2009.02898 \[hep-th\]](#).
- [20] S. Céspedes, A.-C. Davis, and S. Melville, “On the time evolution of cosmological correlators,” [JHEP 02 \(2021\) 012](#), [arXiv:2009.07874 \[hep-th\]](#).
- [21] E. Pajer, “Building a Boostless Bootstrap for the Bispectrum,” [arXiv:2010.12818 \[hep-th\]](#).
- [22] J. M. Maldacena, “Non-Gaussian features of primordial fluctuations in single field inflationary models,” [JHEP 05 \(2003\) 013](#), [arXiv:astro-ph/0210603 \[astro-ph\]](#).
- [23] P. Creminelli, J. Gleyzes, J. Noreña, and F. Vernizzi, “Resilience of the standard predictions for primordial tensor modes,” [Phys. Rev. Lett. 113 no. 23, \(2014\) 231301](#), [arXiv:1407.8439 \[astro-ph.CO\]](#).
- [24] L. Bordin, G. Cabass, P. Creminelli, and F. Vernizzi, “Simplifying the EFT of Inflation: generalized disformal transformations and redundant couplings,” [JCAP 1709 no. 09, \(2017\) 043](#), [arXiv:1706.03758 \[astro-ph.CO\]](#).
- [25] L. Bordin and G. Cabass, “Graviton non-Gaussianities and Parity Violation in the EFT of Inflation,” [JCAP 07 \(2020\) 014](#), [arXiv:2004.00619 \[astro-ph.CO\]](#).
- [26] N. Bartolo, L. Caloni, G. Orlando, and A. Ricciardone, “Tensor non-Gaussianity in chiral scalar-tensor theories of gravity,” [arXiv:2008.01715 \[astro-ph.CO\]](#).
- [27] S. Endlich, A. Nicolis, and J. Wang, “Solid Inflation,” [JCAP 10 \(2013\) 011](#), [arXiv:1210.0569 \[hep-th\]](#).
- [28] C. Cheung, P. Creminelli, A. L. Fitzpatrick, J. Kaplan, and L. Senatore, “The Effective Field Theory of Inflation,” [JHEP 03 \(2008\) 014](#), [arXiv:0709.0293 \[hep-th\]](#).
- [29] S. R. Coleman, J. Wess, and B. Zumino, “Structure of phenomenological Lagrangians. 1.,” [Phys. Rev. 177 \(1969\) 2239–2247](#).
- [30] C. G. Callan, Jr., S. R. Coleman, J. Wess, and B. Zumino, “Structure of phenomenological Lagrangians. 2.,” [Phys. Rev. 177 \(1969\) 2247–2250](#).
- [31] D. V. Volkov, “Phenomenological Lagrangians,” [Fiz. Elem. Chast. Atom. Yadra 4 \(1973\) 3–41](#).
- [32] E. A. Ivanov and V. I. Ogievetsky, “The Inverse Higgs Phenomenon in Nonlinear Realizations,” [Teor. Mat. Fiz. 25 \(1975\) 164–177](#).
- [33] L. V. Delacrétaz, S. Endlich, A. Monin, R. Penco, and F. Riva, “(Re-)Inventing the Relativistic Wheel: Gravity, Cosets, and Spinning Objects,” [JHEP 11 \(2014\) 008](#), [arXiv:1405.7384 \[hep-th\]](#).
- [34] L. V. Delacrétaz, T. Noumi, and L. Senatore, “Boost Breaking in the EFT of Inflation,” [JCAP 02 \(2017\) 034](#), [arXiv:1512.04100 \[hep-th\]](#).
- [35] L. Bordin, P. Creminelli, A. Khmelnitsky, and L. Senatore, “Light Particles with Spin in Inflation,” [JCAP 1810 no. 10, \(2018\) 013](#), [arXiv:1806.10587 \[hep-th\]](#).
- [36] N. Bartolo, D. Cannone, A. Ricciardone, and G. Tasinato, “Distinctive signatures of space-time diffeomorphism breaking in EFT of inflation,” [JCAP 03 \(2016\) 044](#), [arXiv:1511.07414 \[astro-ph.CO\]](#).

- [37] A. Ricciardone and G. Tasinato, “Primordial gravitational waves in supersolid inflation,” [Phys. Rev. D](#) **96** no. 2, (2017) 023508, [arXiv:1611.04516 \[astro-ph.CO\]](#).
- [38] G. Domènech, T. Hiramatsu, C. Lin, M. Sasaki, M. Shiraishi, and Y. Wang, “CMB Scale Dependent Non-Gaussianity from Massive Gravity during Inflation,” [JCAP](#) **05** (2017) 034, [arXiv:1701.05554 \[astro-ph.CO\]](#).
- [39] M. Celoria, D. Comelli, L. Pilo, and R. Rollo, “Boosting GWs in Supersolid Inflation,” [JHEP](#) **01** (2021) 185, [arXiv:2010.02023 \[gr-qc\]](#).
- [40] C. Lin and L. Z. Labun, “Effective Field Theory of Broken Spatial Diffeomorphisms,” [JHEP](#) **03** (2016) 128, [arXiv:1501.07160 \[hep-th\]](#).
- [41] S. Dubovsky, T. Grégoire, A. Nicolis, and R. Rattazzi, “Null energy condition and superluminal propagation,” [JHEP](#) **03** (2006) 025, [arXiv:hep-th/0512260](#).
- [42] S. Endlich, B. Horn, A. Nicolis, and J. Wang, “Squeezed limit of the solid inflation three-point function,” [Phys. Rev. D](#) **90** no. 6, (2014) 063506, [arXiv:1307.8114 \[hep-th\]](#).
- [43] L. Bordin, P. Creminelli, M. Mirbabayi, and J. Noreña, “Solid Consistency,” [JCAP](#) **03** (2017) 004, [arXiv:1701.04382 \[astro-ph.CO\]](#).
- [44] S. Jazayeri, E. Pajer, and D. van der Woude, “Solid Soft Theorems,” [JCAP](#) **06** (2019) 011, [arXiv:1902.09020 \[hep-th\]](#).
- [45] N. Bartolo, S. Matarrese, M. Peloso, and A. Ricciardone, “Anisotropy in solid inflation,” [JCAP](#) **08** (2013) 022, [arXiv:1306.4160 \[astro-ph.CO\]](#).
- [46] M. Akhshik, R. Emami, H. Firouzjahi, and Y. Wang, “Statistical Anisotropies in Gravitational Waves in Solid Inflation,” [JCAP](#) **09** (2014) 012, [arXiv:1405.4179 \[astro-ph.CO\]](#).
- [47] N. Bartolo, M. Peloso, A. Ricciardone, and C. Unal, “The expected anisotropy in solid inflation,” [JCAP](#) **11** (2014) 009, [arXiv:1407.8053 \[astro-ph.CO\]](#).
- [48] E. Dimastrogiovanni, M. Fasiello, D. Jeong, and M. Kamionkowski, “Inflationary tensor fossils in large-scale structure,” [JCAP](#) **12** (2014) 050, [arXiv:1407.8204 \[astro-ph.CO\]](#).
- [49] M. Akhshik, “Clustering Fossils in Solid Inflation,” [JCAP](#) **05** (2015) 043, [arXiv:1409.3004 \[astro-ph.CO\]](#).
- [50] A. A. Abolhasani, M. Akhshik, R. Emami, and H. Firouzjahi, “Primordial Statistical Anisotropies: The Effective Field Theory Approach,” [JCAP](#) **03** (2016) 020, [arXiv:1511.03218 \[astro-ph.CO\]](#).
- [51] L. Bordin, P. Creminelli, M. Mirbabayi, and J. Noreña, “Tensor Squeezed Limits and the Higuchi Bound,” [JCAP](#) **1609** no. 09, (2016) 041, [arXiv:1605.08424 \[astro-ph.CO\]](#).
- [52] P. Mészáros, “Inflationary tensor fossils deformed by solid matter - scalar field interaction,” [JCAP](#) **09** (2019) 048, [arXiv:1905.03544 \[gr-qc\]](#).
- [53] J. Kang and A. Nicolis, “Platonic solids back in the sky: Icosahedral inflation,” [JCAP](#) **03** (2016) 050, [arXiv:1509.02942 \[hep-th\]](#).
- [54] J. Kang and A. Nicolis, “Anisotropic tensor modes from icosahedral inflation,” [arXiv:1807.03924 \[hep-th\]](#).
- [55] A. Nicolis and G. Sun, “Scalar-tensor mixing from icosahedral inflation,” [arXiv:2011.04687 \[hep-th\]](#).