

**$(G, \chi_\phi)$ -EQUIVARIANT  $\phi$ -COORDINATED QUASI MODULES FOR VERTEX ALGEBRAS**FULIN CHEN<sup>1</sup>, XIAOLING LIAO, SHAOBIN TAN<sup>2</sup>, AND QING WANG<sup>3</sup>

ABSTRACT. To give a unified treatment on the association of Lie algebras and vertex algebras, we study  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for vertex algebras, where  $G$  is a group with  $\chi_\phi$  a linear character of  $G$  and  $\phi$  is an associate of the one-dimensional additive formal group. The theory of  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for nonlocal vertex algebra is established in [JKLT]. In this paper, we concentrate on the context of vertex algebras. We establish several conceptual results, including a generalized commutator formula and a general construction of vertex algebras and their  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules. Furthermore, for any conformal algebra  $\mathcal{C}$ , we construct a class of Lie algebras  $\widehat{\mathcal{C}}_\phi[G]$  and prove that restricted  $\widehat{\mathcal{C}}_\phi[G]$ -modules are exactly  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for the universal enveloping vertex algebra of  $\mathcal{C}$ . As an application, we determine the  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for affine and Virasoro vertex algebras.

## 1. INTRODUCTION

Modules and twisted modules have been extensively studied in vertex algebra theory. We usually associate Lie algebras with vertex algebras through modules or twisted modules (see [FLM, FZ, Li1, Li6], etc). But these theories are not valid for some algebras which generating functions are not “local”, we need generalize the notion of modules. A notion of quasi module of vertex algebra was introduced in [Li5] to associate more infinite dimensional Lie algebras like quantum torus Lie algebras with vertex algebras. In order to associate quantum affine algebras with quantum vertex algebras, a theory of  $\phi$ -coordinated module for the quantum vertex algebra was developed in [Li2]. The parameter  $\phi$  is an associate of the one-dimensional additive formal group  $F(x, y) = x + y$ . When  $\phi = x + z$ , a  $\phi$ -coordinated module of vertex algebra is just a module. To establish the explicit connections of quantum vertex algebras to various algebraic structures like quantum affine algebras, a notion of  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi module

---

2010 *Mathematics Subject Classification.* 17B65 & 17B69.

<sup>1</sup>Partially supported by China NSF grant (No.11971397) and the Fundamental Research Funds for the Central Universities (No.20720190069).

<sup>2</sup>Partially supported by China NSF grants (No.11971397).

<sup>3</sup>Partially supported by China NSF grants (No.12071385) and the Fundamental Research Funds for the Central Universities (No.20720200067).

for nonlocal vertex algebras was introduced in [JKLT], where  $G$  is a group and  $\chi_\phi$  is a linear character of  $G$ . In [Li2], the author proved that for every  $p(x) \in \mathbb{C}((x))$ ,  $\phi = e^{zp(x)\frac{d}{dx}}(x)$  is an associate of  $F(x, y)$  and every associate is of this form. Moreover, an explicit association between quantum vertex algebra and certain quantum  $\beta\gamma$ -system was established in [Li2] through  $\phi$ -coordinated modules, where  $\phi = xe^z$ , i.e.,  $p(x) = x$ . Recently, the  $\phi_\epsilon$ -coordinated modules of vertex algebras have been studied precisely in [BLP] for  $p(x) = x^\epsilon$  with  $\epsilon$  an integer, and they use the theory of  $\phi_\epsilon$ -coordinated modules to associate vertex algebras with Novikov algebras. Specifically, for  $\phi = xe^z$ , a notion of  $G$ -equivariant  $\phi$ -coordinated quasi module for quantum vertex algebras was introduced in [Li4]. A commutator formula for  $G$ -equivariant  $\phi$ -coordinated quasi module was obtained and a natural connection between the deformed Virasoro algebra  $Vir_{p,-1}$  of a certain Clifford vertex superalgebra through  $G$ -equivariant  $\phi$ -coordinated quasi module was established therein. Later in [JKLT], a generalized commutator formula and a general construction of  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for nonlocal vertex algebras were established, and by using these results, they give a construction of (equivariant)  $\phi$ -coordinated quasi modules for lattice vertex algebras. We note that if  $\phi = xe^z$ , the notion of  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi module coincides with the notion of  $G$ -equivariant  $\phi$ -coordinated quasi module. Motivated by these works, in this paper, we focus on the frame of vertex algebra. We expect to unify the association of Lie algebras and vertex algebras though  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules.

Next, we briefly outline our ideas. Let  $\phi = e^{zp(x)\frac{d}{dx}}(x)$  for  $0 \neq p(x) \in \mathbb{C}((x))$ . A  $\phi$ -coordinated module for a vertex algebra  $V$  is a vector space  $W$  with a linear map  $Y_W(\cdot, z)$  from  $V$  to  $\text{Hom}(W, W((z)))$  satisfying the conditions that  $Y_W(\mathbf{1}, z) = 1_W$  and that for  $u, v \in V$ , there exists a nonnegative integer  $k$  such that

$$\begin{aligned} (z_1 - z_2)^k Y_W(u, z_1) Y_W(v, z_2) &\in \text{Hom}(W, W((z_1, z_2))), \\ (\phi(z_2, z_0) - z_2)^k Y_W(Y(u, z_0)v, z_2) &= ((z_1 - z_2)^k Y_W(u, z_1) Y_W(v, z_2)) \Big|_{z_1=\phi(z_2, z_0)}. \end{aligned}$$

By generalizing the corresponding result in [BLP], we prove that if  $(W, Y_W)$  is a  $\phi$ -coordinated  $V$ -module, then for  $u, v \in V$ , we have the commutator formula

$$[Y_W(u, z_1), Y_W(v, z_2)] = \sum_{i \geq 0} \frac{1}{i!} Y_W(u_i v, z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^i p(z_1) z_1^{-1} \delta \left( \frac{z_2}{z_1} \right).$$

For any vector space  $W$ , we prove that any (quasi) local subset of  $\mathcal{E}(W)$  (=  $\text{Hom}(W, W((z)))$ ) can generate a vertex algebra with  $W$  is naturally a  $\phi$ -coordinated (quasi) module of it. To give a unified treatment on the association of Lie algebras and vertex algebras, we need the notion of conformal algebra which was introduced in [K] (also known as a vertex Lie algebra in [DLM, P]) as a bridge. A conformal algebra  $\mathcal{C}$  is a  $\mathbb{C}[\partial]$ -module with a  $\mathbb{C}$ -bilinear product  $a_n b$  for each  $n \in \mathbb{N}$  satisfying

some axioms. Define a multiplication on the formal loop space  $\mathcal{C} \otimes \mathbb{C}((x))$  by

$$[a \otimes f(x), b \otimes g(x)] = \sum_{i \geq 0} \frac{1}{i!} (a_i b) \otimes \left( \left( p(x) \frac{d}{dx} \right)^i f(x) \right) g(x)$$

for  $a, b \in \mathcal{C}$  and  $f(x), g(x) \in \mathbb{C}((x))$ . Then the quotient space

$$\widehat{\mathcal{C}}_\phi := \mathcal{C} \otimes \mathbb{C}((x)) / \text{Im} \left( \partial \otimes 1 + 1 \otimes p(x) \frac{d}{dx} \right)$$

with this multiplication form a Lie algebra [K]. For  $a \in \mathcal{C}$ , we set

$$a_\phi(z) = \sum_{n \in \mathbb{Z}} \overline{a \otimes x^n p(x)} z^{-n-1} \in \widehat{\mathcal{C}}_\phi[[z, z^{-1}]],$$

where  $\overline{a \otimes x^n p(x)}$  is the image of  $a \otimes x^n p(x)$  in  $\widehat{\mathcal{C}}_\phi$ . Then for  $a, b \in \mathcal{C}$ ,

$$[a_\phi(z), b_\phi(w)] = \sum_{i \geq 0} \frac{1}{i!} (a_i b)_\phi(w) \left( p(w) \frac{\partial}{\partial w} \right)^i p(z) z^{-1} \delta \left( \frac{w}{z} \right).$$

We prove that restricted  $\widehat{\mathcal{C}}_\phi$ -modules are exactly  $\phi$ -coordinated modules for the universal enveloping vertex algebra  $V_{\mathcal{C}}$  of  $\mathcal{C}$ .

Let  $G$  be a group with a linear character  $\chi$ . Following [JKLT], a  $(G, \chi)$ -vertex algebra is a vertex algebra  $V$  with a  $G$ -action  $R$  on  $V$  such that  $R_g(\mathbf{1}) = \mathbf{1}$  and

$$R_g Y(v, z) R_g^{-1} = Y(R_g(v), \chi(g)z) \quad \text{for } g \in G, v \in V.$$

The notion of  $(G, \chi)$ -vertex algebra was introduced in [Li3] (cf. [Li5]). A  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi module is defined for a  $(G, \chi)$ -vertex algebra and from [JKLT], we may always assume that

$$\phi(x, \chi(g)z) = \chi_\phi(g) \phi(\chi_\phi(g)^{-1}x, z) \quad \text{for } g \in G.$$

In this paper, we obtain a general construction of vertex algebras and their  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules based on [JKLT]. Let  $(W, Y_W)$  be a  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V$ -module, we prove the following generalized commutator formula:

$$\begin{aligned} & [Y_W(u, z_1), Y_W(v, z_2)] \\ &= \text{Res}_{z_0} \sum_{g \in \psi(\chi_\phi(G))} Y_W(Y(R_{g^{-1}}u, z_0)v, z_2) e^{z_0 p(z_2) \frac{\partial}{\partial z_2}} \left( \chi(g) p(z_1) z_1^{-1} \delta \left( \frac{\chi_\phi(g) z_2}{z_1} \right) \right), \end{aligned}$$

where  $\psi : \chi_\phi(G) \rightarrow G$  is a section of  $\chi_\phi$ . The notion of  $(G, \chi)$ -conformal algebra was introduced in [Li3] which generalizes the notion of  $\Gamma$ -conformal algebra in [G-K-K]. A  $(G, \chi)$ -conformal algebra is a conformal algebra  $\mathcal{C}$  with a  $G$ -action on  $\mathcal{C}$  satisfying

$$R_g(a_m b) = \chi(g)^{-(m+1)} (R_g a)_m (R_g b) \quad \text{for any } a, b \in \mathcal{C}, m \in \mathbb{N},$$

and some other compatible conditions. Let  $(\mathcal{C}, R)$  be a  $(G, \chi)$ -conformal algebra, for any  $g \in G$ , we define a linear automorphism  $\hat{R}_g$  on  $\mathcal{C} \otimes \mathbb{C}((x))$  by

$$\hat{R}_g(a \otimes f(x)) = \chi(g)^{-1} R_g(a) \otimes f(\chi_\phi(g)^{-1}x), \quad a \in \mathcal{C}, f(x) \in \mathbb{C}((x)).$$

Then we define a multiplication on  $\hat{\mathcal{C}}_\phi$  by

$$(u, v) \mapsto \sum_{g \in G} [\hat{R}_g(u), v], \quad u, v \in \hat{\mathcal{C}}_\phi.$$

It follows that the quotient space

$$\hat{\mathcal{C}}_\phi[G] := \hat{\mathcal{C}}_\phi / \text{span}\{\hat{R}_g(u) - u \mid g \in G, u \in \hat{\mathcal{C}}_\phi\}$$

is a Lie algebra with this new multiplication. For  $a \in \mathcal{C}$ ,  $f(x) \in \mathbb{C}((x))$ , we still denote the image of  $\overline{a \otimes f(x)}$  in  $\hat{\mathcal{C}}_\phi[G]$  by  $\overline{a \otimes f(x)}$ . For  $a \in \mathcal{C}$ , we set

$$a_\phi^G(z) = \sum_{n \in \mathbb{Z}} \overline{a \otimes x^n p(x)} z^{-n-1} \in \hat{\mathcal{C}}_\phi[G][[z, z^{-1}]].$$

Then for  $a, b \in \mathcal{C}$ , we have

$$\begin{aligned} & [a_\phi^G(z), b_\phi^G(w)] \\ &= \sum_{g \in G} \sum_{i \geq 0} \frac{1}{i!} ((R_{g^{-1}}a)_i b)_\phi^G(w) \left( p(w) \frac{\partial}{\partial w} \right)^i \left( \chi(g) p(z) z^{-1} \delta \left( \frac{\chi_\phi(g)w}{z} \right) \right). \end{aligned}$$

Let  $H = \ker \chi_\phi$  and let  $\bar{\chi}_\phi$  be the linear character of  $G/H$  induced by  $\chi_\phi$ . From [Li3], the quotient space

$$\mathcal{C}/H = \mathcal{C} / \text{span}\{R_h a - a \mid h \in H, a \in \mathcal{C}\}$$

has a natural  $(G/H, \bar{\chi})$ -conformal algebra structure, where  $\bar{\chi}$  is a linear character of  $G/H$  induced by  $\chi$ . Then we prove that restricted  $\hat{\mathcal{C}}_\phi[G]$ -modules are exactly  $(G/H, \bar{\chi}_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V_{\mathcal{C}/H}$ -modules. Finally, we apply these theories on affine and Virasoro vertex algebras, and the vertex algebras associated to Novikov algebras.

This paper is organized as follows. In Section 2, we recall  $\phi$ -operations on  $\mathcal{E}(W)$  and establish some technical lemmas. In Section 3, we recall the notion of  $\phi$ -coordinated (quasi) module for a vertex algebra and establish some conceptual results. In Section 4, we prove that restricted  $\hat{\mathcal{C}}_\phi$ -modules are exactly  $\phi$ -coordinated  $V_{\mathcal{C}}$ -modules. In Section 5, we establish a generalized commutator formula and a general construction of  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules of vertex algebras. Furthermore, we construct a class of Lie algebras  $\hat{\mathcal{C}}_\phi[G]$  and prove that restricted  $\hat{\mathcal{C}}_\phi[G]$ -modules are exactly  $(G/H, \bar{\chi}_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V_{\mathcal{C}/H}$ -modules. In Section 6, we determine a class of  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for affine vertex algebras, and determine the  $\phi$ -coordinated modules for Virasoro vertex algebra and the vertex algebras associated to Novikov algebras.

In this paper, we let  $\mathbb{Z}, \mathbb{Z}^\times, \mathbb{Z}_+, \mathbb{N}$  and  $\mathbb{C}$  be the sets of integers, nonzero integers, positive integers, nonnegative integers and complex numbers, respectively. All the Lie algebras in this paper are over the field of complex numbers. For any linear endomorphism  $\varphi$  of a vector space, we set  $\varphi^{(k)} = \frac{\varphi^k}{k!}$ .

## 2. $\phi$ -OPERATIONS ON $\mathcal{E}(W)$

In this section, we recall the  $\phi$ -operations following [Li2] and prove some technical Lemmas. We begin by recalling the notion of an associate for the one-dimensional additive formal group (law)  $F(x, y) = x + y$ . Following [Li2], an associate  $\phi(x, z)$  (or simply  $\phi$ ) of  $F(x, y)$  by definition is a formal series in  $\mathbb{C}((x))[[z]]$  satisfying

$$\phi(x, 0) = x, \quad \phi(\phi(x, z_1), z_2) = \phi(x, z_1 + z_2).$$

It was proved therein that for every  $p(x) \in \mathbb{C}((x))$ ,  $\phi(x, z) = e^{zp(x)\frac{d}{dx}}(x)$  is an associate of  $F(x, y)$  and every associate is of this form with  $p(x)$  uniquely determined. Throughout this paper, we fix an associate

$$(2.1) \quad \phi = \phi(x, z) = e^{zp(x)\frac{d}{dx}}(x) \quad \text{with} \quad 0 \neq p(x) \in \mathbb{C}((x)).$$

Throughout this section let  $W$  be a fixed vector space and set

$$\mathcal{E}(W) = \text{Hom}(W, W((z))) \subset (\text{End}W)[[z, z^{-1}]].$$

A pair  $(a(z), b(z))$  in  $\mathcal{E}(W)$  is said to be *quasi local* if there exists a nonzero polynomial  $q(z_1, z_2) \in \mathbb{C}[z_1, z_2]$  such that

$$(2.2) \quad q(z_1, z_2) [a(z_1), b(z_2)] = 0,$$

and is said to be *local* if the polynomial  $q(z_1, z_2)$  is assumed to a polynomial of the form  $(z_1 - z_2)^n$  for some  $n \in \mathbb{N}$ . The main goal of this section is to establish a commutator formula for quasi local pairs in  $\mathcal{E}(W)$  in terms of the following  $\phi$ -operations introduced in [Li2]:

**Definition 2.1.** Let  $(a(z), b(z))$  be a quasi local pair in  $\mathcal{E}(W)$  such that (2.2) holds. We define

$$Y_{\mathcal{E}}^\phi(a(z), z_0)b(z) = \sum_{n \in \mathbb{Z}} a(z)_n^\phi b(z) z_0^{-n-1} \in \mathcal{E}(W)((z_0))$$

by the following rule:

$$Y_{\mathcal{E}}^\phi(a(z), z_0)b(z) = q(\phi(z, z_0), z)^{-1} (q(z_1, z) a(z_1) b(z)) |_{z_1 = \phi(z, z_0)}.$$

It was proved therein that the definition of  $Y_{\mathcal{E}}^\phi$  dose not depend on the choice of  $q(z_1, z_2)$ . In what follows, we establish some technical lemmas.

**Lemma 2.2.** *Let  $j, k$  be two nonnegative integers with  $j > k$ . Then for any nonzero complex number  $\lambda$ , we have*

$$(2.3) \quad (z_1 - \lambda z_2)^j \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(k)} \delta \left( \frac{\lambda z_2}{z_1} \right) = 0.$$

*Proof.* Note that we may write

$$\left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(k)} = f_k(z_2) \left( \frac{\partial}{\partial z_2} \right)^{(k)} + \cdots + f_1(z_2) \frac{\partial}{\partial z_2} + f_0(z_2)$$

for some uniquely determined  $f_0(z_2), \dots, f_k(z_2) \in \mathbb{C}((z_2))$ . Then the assertion follows from the fact that  $(z_1 - \lambda z_2)^j \left( \frac{\partial}{\partial z_2} \right)^{(i)} \delta \left( \frac{\lambda z_2}{z_1} \right) = 0$  for  $i = 0, \dots, k$ .  $\square$

For convenience, we set

$$\bar{p}(x) = x^{-1}p(x) \in \mathbb{C}((x)).$$

**Lemma 2.3.** *Let  $\lambda_1, \dots, \lambda_r$  be distinct nonzero complex numbers,  $k_1, \dots, k_r \in \mathbb{Z}_+$  and  $A_{ij}(z) \in \mathcal{E}(W)$ , where  $i = 1, \dots, r$ , and for a fixed  $i$ ,  $0 \leq j \leq k_i - 1$ . Then*

$$(2.4) \quad \sum_{i=1}^r \sum_{j=0}^{k_i-1} A_{ij}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_i^{-1} z_1) \delta \left( \frac{\lambda_i z_2}{z_1} \right) = 0$$

*if and only if  $A_{ij}(z) = 0$  for all  $i, j$ .*

*Proof.* For  $1 \leq i \leq r$ , set

$$f_i(z) = \frac{\prod_{j=1}^r (z - \lambda_j)^{k_j}}{(z - \lambda_i)^{k_i}} \in \mathbb{C}[z].$$

In view of Lemma 2.2, for  $1 \leq i \neq s \leq r$  and  $0 \leq j \leq k_s - 1$  we have

$$(2.5) \quad f_i(z_1/z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_s^{-1} z_1) \delta \left( \frac{\lambda_s z_2}{z_1} \right) = 0.$$

As  $f_1(z), \dots, f_r(z)$  are pairwise relatively prime, we may take suitable polynomials  $q_1(z), \dots, q_r(z) \in \mathbb{C}[z]$  such that

$$q_1(z)f_1(z) + \cdots + q_r(z)f_r(z) = 1.$$

This together with (2.5) gives that for  $1 \leq s \leq r$ ,  $j = 0, \dots, k_s - 1$ ,

$$(2.6) \quad \begin{aligned} & A_{sj}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_s^{-1} z_1) \delta \left( \frac{\lambda_s z_2}{z_1} \right) \\ &= \left( \sum_{i=1}^r f_i(z_1/z_2) q_i(z_1/z_2) \right) A_{sj}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_s^{-1} z_1) \delta \left( \frac{\lambda_s z_2}{z_1} \right) \\ &= A_{sj}(z_2) f_s(z_1/z_2) q_s(z_1/z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_s^{-1} z_1) \delta \left( \frac{\lambda_s z_2}{z_1} \right). \end{aligned}$$

Now we assume that (2.4) holds and fix  $s$  with  $1 \leq s \leq r$ . Then from (2.5) and (2.6) we have

$$\begin{aligned} 0 &= \sum_{i=1}^r \sum_{j=0}^{k_i-1} A_{ij}(z_2) f_s(z_1/z_2) q_s(z_1/z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_i^{-1} z_1) \delta \left( \frac{\lambda_i z_2}{z_1} \right) \\ &= \sum_{j=0}^{k_s-1} A_{sj}(z_2) f_s(z_1/z_2) q_s(z_1/z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_s^{-1} z_1) \delta \left( \frac{\lambda_s z_2}{z_1} \right) \\ &= \sum_{j=0}^{k_s-1} A_{sj}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_s^{-1} z_1) \delta \left( \frac{\lambda_s z_2}{z_1} \right), \end{aligned}$$

which amounts to

$$(2.7) \quad \sum_{j=0}^{k_s-1} A_{sj}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} p(z_1) z_1^{-1} \delta \left( \frac{z_2}{z_1} \right) = 0.$$

For  $j = 0, \dots, k_s - 1$ , write

$$A_{sj}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} = \sum_{n=0}^j g_{jn}(z_2) \left( \frac{\partial}{\partial z_2} \right)^{(n)}$$

for some uniquely determined  $g_{jn}(z_2) \in \mathbb{C}((z_2))$ . In view of (2.7) we have

$$\sum_{n=0}^{k_s-1} \left( \sum_{j=n}^{k_s-1} g_{jn}(z_2) \right) \left( \frac{\partial}{\partial z_2} \right)^{(n)} z_1^{-1} \delta \left( \frac{z_2}{z_1} \right) = 0.$$

By [Li1, Lemma 2.1.4], this implies  $\sum_{j=n}^{k_s-1} g_{jn}(z_2) = 0$  for  $0 \leq n \leq k_s - 1$ , and hence implies  $A_{sj}(z) = 0$  for  $0 \leq j \leq k_s - 1$ , as desired.  $\square$

The following result is a slight generalization of [Li4, Lemma 2.3].

**Lemma 2.4.** *Let  $a(z), b(z) \in \mathcal{E}(W)$  and let*

$$(2.8) \quad q(z) = (z - \lambda_1)^{k_1} \cdots (z - \lambda_r)^{k_r} \in \mathbb{C}[z],$$

where  $\lambda_1, \dots, \lambda_r \in \mathbb{C}^\times$  are distinct and  $k_1, \dots, k_r \in \mathbb{Z}_+$ . Then

$$(2.9) \quad q(z_1/z_2) [a(z_1), b(z_2)] = 0$$

if and only if

$$(2.10) \quad [a(z_1), b(z_2)] = \sum_{i=1}^r \sum_{j=0}^{k_i-1} A_{ij}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_i^{-1} z_1) \delta \left( \frac{\lambda_i z_2}{z_1} \right)$$

for some uniquely determined  $A_{ij}(z) \in \mathcal{E}(W)$ .

*Proof.* The proof is similar to that of [Li4, Lemma 2.3] and we give a sketch of it. The uniqueness of  $A_{ij}(z)$  follows from Lemma 2.3. For the existence, write

$$\frac{1}{q(x)} = \sum_{i=1}^r \sum_{j=1}^{k_i} \frac{a_{ij}}{(x - \lambda_i)^j}$$

for some  $a_{ij} \in \mathbb{C}$ . Then from (2.9) we obtain (cf. [Li4, Lemma 2.3])

$$(2.11) \quad [a(z_1), b(z_2)] = \sum_{i=1}^r \sum_{j=0}^{k_i-1} a_{i,j+1} \lambda_i^{-j} z_2^{j+1} A(z_1, z_2) \left( \frac{\partial}{\partial z_2} \right)^{(j)} z_1^{-1} \delta \left( \frac{\lambda_i z_2}{z_1} \right),$$

where  $A(z_1, z_2) = q(z_1/z_2)a(z_1)b(z_2) \in \text{Hom}(W, W((z_1, z_2)))$ .

Note that for any  $n \in \mathbb{N}$ , there exist  $f_0(z_2), \dots, f_n(z_2) \in \mathbb{C}((z_2))$  such that

$$\left( \frac{\partial}{\partial z_2} \right)^{(n)} = f_n(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(n)} + f_{n-1}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(n-1)} + \dots + f_0(z_2).$$

An induction argument shows that for any  $B(z_1, z_2) \in \text{Hom}(W, W((z_1, z_2)))$  and  $n \in \mathbb{N}$ , there exist  $B_0(z), \dots, B_n(z) \in \mathcal{E}(W)$  such that

$$B(z_1, z_2) \left( \frac{\partial}{\partial z_2} \right)^{(n)} z_1^{-1} \delta \left( \frac{\lambda_i z_2}{z_1} \right) = \sum_{i=0}^n B_i(z_2) \left( \frac{\partial}{\partial z_2} \right)^{(i)} \bar{p}(\lambda_i^{-1} z_1) \delta \left( \frac{\lambda_i z_2}{z_1} \right).$$

Thus (2.10) follows.  $\square$

We fix a  $\hat{p}(x) \in \mathbb{C}((x))$  such that  $\frac{d}{dx} \hat{p}(x) = p^{-1}(x)$ , and set

$$f_{p(x)}(x, z) := \hat{p}(x(1+z)) - \hat{p}(x).$$

Then we can deduce that

$$(2.12) \quad \phi(x, f_{p(x)}(x, z)) = x(1+z).$$

We have (cf. [BLP, Li2]):

**Lemma 2.5.** *Let*

$$A(z_1, z_2) \in \text{Hom}(W, W((z_1))((z_2))), \quad B(z_1, z_2) \in \text{Hom}(W, W((z_2))((z_1))), \\ C(z_0, z_2) \in (\text{Hom}(W, W((z_2))))((z_0)).$$

*If there exists a nonnegative integer  $k$  such that*

$$(z_1 - z_2)^k A(z_1, z_2) = (z_1 - z_2)^k B(z_1, z_2), \\ ((z_1 - z_2)^k A(z_1, z_2))|_{z_1=\phi(z_2, z_0)} = (\phi(z_2, z_0) - z_2)^k C(z_0, z_2),$$

*then*

$$(z_2 z)^{-1} \delta \left( \frac{z_1 - z_2}{z_2 z} \right) A(z_1, z_2) - (z_2 z)^{-1} \delta \left( \frac{z_2 - z_1}{-z_2 z} \right) B(z_1, z_2) \\ = z_1^{-1} \delta \left( \frac{z_2(1+z)}{z_1} \right) C(f_{p(z_2)}(z_2, z), z_2).$$



*Proof.* When  $p(x) = x^\epsilon$  for some  $\epsilon \in \mathbb{Z}$ , the assertion is proved in [BLP, Lemma 3.16], where  $f_{p(x)}(x, z)$  is denoted as  $f_\epsilon(x, z)$ . In view of (2.12), the proof therein also valid for a general  $p(x)$ .  $\square$

Now we are in a position to state the main result of this section, which says that those  $A_{ij}(z)$  in (2.10) are nothing but the  $\phi$ -products  $a(\lambda_i z)_j^\phi b(z)$ .

**Proposition 2.6.** *Let  $(a(z), b(z))$  be a pair in  $\mathcal{E}(W)$  satisfying*

$$q(z_1/z_2)[a(z_1), b(z_2)] = 0$$

for some  $q(z) \in \mathbb{C}[z]$ . Then

$$[a(z_1), b(z_2)] = \sum_{\lambda \in \mathbb{C}^\times} \sum_{j \geq 0} a(\lambda z_2)_j^\phi b(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda^{-1} z_1) \delta \left( \frac{\lambda z_2}{z_1} \right),$$

which is a finite sum. Furthermore, the coefficients  $a(\lambda z_2)_j^\phi b(z_2) \neq 0$  only if  $\lambda$  is a root of  $q(z)$  and  $j$  is less than the multiplicity of  $\lambda$ .

*Proof.* We assume that  $q(z)$  has the form as in (2.8) and fix an  $s$  with  $1 \leq s \leq r$ . From (2.10), we only need to prove that

$$(2.13) \quad A_{sj}(z_2) = a(\lambda_s z_2)_j^\phi b(z_2), \quad j = 0, \dots, k_s - 1.$$

Let us introduce the polynomials

$$q_s(z) = \prod_{1 \leq i \neq s \leq r} ((z - \lambda_s)^{k_s} - (\lambda_i - \lambda_s)^{k_s})^{k_i}.$$

Note that  $(z - \lambda_i)^{k_i}$  divides  $q_s(z)$  and so there is a  $g_s(z) \in \mathbb{C}[z]$  such that

$$(2.14) \quad q_s(z)(z - \lambda_s)^{k_s} = g_s(z)q(z).$$

Then from (2.14) and (2.3), we have

$$(2.15) \quad \begin{aligned} & q_s(z_1/z_2)[a(z_1), b(z_2)] \\ &= \sum_{j=0}^{k_s-1} q_s(z_1/z_2) A_{sj}(z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda_s^{-1} z_1) \delta \left( \frac{\lambda_s z_2}{z_1} \right). \end{aligned}$$

By replacing  $z_1$  with  $\lambda_s z_1$  in (2.15) and multiplying  $(z_1/z_2 - 1)^{k_s}$  on both sides of (2.15) we have

$$(z_1/z_2 - 1)^{k_s} q_s(\lambda_s z_1/z_2)[a(\lambda_s z_1), b(z_2)] = 0.$$

Thus we have (see Definition 2.1)

$$\begin{aligned} & (\phi(z, z_0)/z - 1)^{k_s} q_s(\lambda_s \phi(z, z_0)/z) Y_{\mathcal{E}}^\phi(a(\lambda_s z), z_0) b(z) \\ &= (z_1/z - 1)^{k_s} q_s(\lambda_s z_1/z) a(\lambda_s z_1) b(z) \Big|_{z_1=\phi(z, z_0)}. \end{aligned}$$

In view of Lemma 2.5, this implies that

$$\begin{aligned} & (z_2 z)^{-1} \delta \left( \frac{z_1 - z_2}{z_2 z} \right) q_s(\lambda_s z_1 / z_2) a(\lambda_s z_1) b(z_2) \\ & \quad - (z_2 z)^{-1} \delta \left( \frac{z_2 - z_1}{-z_2 z} \right) q_s(\lambda_s z_1 / z_2) b(z_2) a(\lambda_s z_1) \\ & = z_1^{-1} \delta \left( \frac{z_2(1+z)}{z_1} \right) q_s(\lambda_s z_1 / z_2) Y_{\mathcal{E}}^{\phi}(a(\lambda_s z_2), f_{p(z_2)}(z_2, z)) b(z_2). \end{aligned}$$

By taking  $\text{Res}_{z_2}$  in the above identity we find

$$\begin{aligned} & q_s(\lambda_s z_1 / z_2) [a(\lambda_s z_1), b(z_2)] \\ & = \text{Res}_{z_2} z_1^{-1} \delta \left( \frac{z_2(1+z)}{z_1} \right) z_2 q_s(\lambda_s z_1 / z_2) Y_{\mathcal{E}}^{\phi}(a(\lambda_s z_2), f_{p(z_2)}(z_2, z)) b(z_2) \\ & = \text{Res}_{z_0} \frac{\partial}{\partial z_0} \left( \frac{\phi(z_2, z_0)}{z_2} - 1 \right) z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) z_2 q_s(\lambda_s z_1 / z_2) Y_{\mathcal{E}}^{\phi}(a(\lambda_s z_2), z_0) b(z_2) \\ & = \text{Res}_{z_0} p(\phi(z_2, z_0)) z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) q_s(\lambda_s z_1 / z_2) Y_{\mathcal{E}}^{\phi}(a(\lambda_s z_2), z_0) b(z_2) \\ & = \text{Res}_{z_0} p(z_1) z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) q_s(\lambda_s z_1 / z_2) Y_{\mathcal{E}}^{\phi}(a(\lambda_s z_2), z_0) b(z_2) \\ & = \sum_{j \geq 0} \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \left( \bar{p}(z_1) \delta \left( \frac{z_2}{z_1} \right) \right) q_s(\lambda_s z_1 / z_2) a(\lambda_s z_2)_j^{\phi} b(z_2), \end{aligned}$$

where in the second equality we have used the fact (2.12). Combining this with (2.15) (replacing  $z_1$  with  $\lambda_s z_1$  therein) we obtain

$$(2.16) \quad \sum_{j=0}^{k_s-1} q_s \left( \lambda_s \frac{z_1}{z_2} \right) (A_{sj}(z_2) - a(\lambda_s z_2)_j^{\phi} b(z_2)) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(z_1) \delta \left( \frac{z_2}{z_1} \right) = 0.$$

Set

$$\alpha = \prod_{1 \leq i \neq s \leq r} (-1)^{k_i} (\lambda_i - \lambda_s)^{k_s k_i} \in \mathbb{C}^{\times}.$$

Note that there is an  $h(z) \in \mathbb{C}[z]$  such that

$$q_s(\lambda_s z) = \prod_{1 \leq i \neq s \leq r} ((\lambda_s z - \lambda_s)^{k_s} - (\lambda_i - \lambda_s)^{k_s})^{k_i} = (z-1)^{k_s} h(z) + \alpha.$$

In view of this and (2.3), (2.16) can be rewritten as

$$\sum_{j=0}^{k_s-1} \alpha (A_{sj}(z_2) - a(\lambda_s z_2)_j^{\phi} b(z_2)) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(z_1) \delta \left( \frac{z_2}{z_1} \right) = 0.$$

Note that  $\alpha \neq 0$ , this together with Lemma 2.3 proves (2.13).  $\square$

**Remark 2.7.** For  $p(x) = 1$  and  $p(x) = x$ , Proposition 2.6 was proved in [Li5] and [Li4] respectively.

### 3. $\phi$ -COORDINATED MODULES OF VERTEX ALGEBRAS

In this section we establish some conceptual results on  $\phi$ -coordinated (quasi) modules for vertex algebras. Let  $(V, Y, \mathbf{1})$  be a vertex algebra, the following notion was introduced in [Li2].

**Definition 3.1.** A  $\phi$ -coordinated quasi  $V$ -module is a vector space  $W$  equipped with a linear map

$$\begin{aligned} Y_W(\cdot, z) : V &\rightarrow \text{Hom}(W, W((z))) \subset (\text{End}W)[[z, z^{-1}]], \\ v &\mapsto Y_W(v, z) = \sum_{n \in \mathbb{Z}} v_n z^{-n-1}, \end{aligned}$$

satisfying the conditions that  $Y_W(\mathbf{1}, z) = 1_W$  and that for  $u, v \in V$ , there exists a polynomial  $f(z_1, z_2) \in \mathbb{C}[z_1, z_2]$  such that

$$\begin{aligned} f(z_1, z_2) Y_W(u, z_1) Y_W(v, z_2) &\in \text{Hom}(W, W((z_1, z_2))), \\ f(\phi(z_2, z_0), z_2) Y_W(Y(u, z_0)v, z_2) &= (f(z_1, z_2) Y_W(u, z_1) Y_W(v, z_2)) \Big|_{z_1=\phi(z_2, z_0)}. \end{aligned}$$

A  $\phi$ -coordinated  $V$ -module is defined as above except that  $f(z_1, z_2)$  is assumed to be a polynomial of the form  $(z_1 - z_2)^k$  with  $k \in \mathbb{N}$ .

**Remark 3.2.** When  $p(x) = 1$ , a  $\phi$ -coordinated  $V$ -module is just a  $V$ -module.

In what follows we collect some basic properties of  $\phi$ -coordinated quasi  $V$ -modules for later use.

**Lemma 3.3.** [Li2] *Let  $(W, Y_W)$  be a  $\phi$ -coordinated quasi  $V$ -module and let  $u, v \in V$ . Assume that there exists a nonzero polynomial  $q(z_1, z_2) \in \mathbb{C}[z_1, z_2]$  such that*

$$q(z_1, z_2) Y_W(u, z_1) Y_W(v, z_2) \in \text{Hom}(W, W((z_1, z_2))).$$

*Then*

$$(3.1) \quad q(z_1, z_2) [Y_W(u, z_1), Y_W(v, z_2)] = 0,$$

$$(3.2) \quad q(\phi(z_2, z_0), z_2) Y_W(Y(u, z_0)v, z_2) = (q(z_1, z_2) Y_W(u, z_1) Y_W(v, z_2)) \Big|_{z_1=\phi(z_2, z_0)}.$$

We define the linear operator  $\mathcal{D}$  on  $V$  by  $\mathcal{D}v = v_{-2}\mathbf{1}$  for  $v \in V$ .

**Lemma 3.4.** [Li2] *Let  $(W, Y_W)$  be a  $\phi$ -coordinated quasi  $V$ -module. Then*

$$(3.3) \quad Y_W(\mathcal{D}v, z) = \left( p(z) \frac{d}{dz} \right) Y_W(v, z), \quad v \in V,$$

$$(3.4) \quad Y_W(Y(u, z_0)v, z) = Y_{\mathcal{E}}^\phi(Y_W(u, z), z_0) Y_W(v, z), \quad u, v \in V.$$

The following is an analogue of Borcherd's commutator formula (cf. [BLP]).

**Proposition 3.5.** *Let  $(W, Y_W)$  be a  $\phi$ -coordinated  $V$ -module. Then for  $u, v \in V$ ,*

$$[Y_W(u, z_1), Y_W(v, z_2)] = \sum_{j \geq 0} Y_W(u_j v, z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(z_1) \delta \left( \frac{z_2}{z_1} \right).$$

*Conversely, if  $W$  is faithful and*

$$[Y_W(u, z_1), Y_W(v, z_2)] = \sum_{j \geq 0} Y_W(c_j, z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(z_1) \delta \left( \frac{z_2}{z_1} \right),$$

*for some  $c_j \in V$ , then  $c_j = u_j v$  for all  $j \geq 0$ .*

*Proof.* The second assertion follows from Lemma 2.3. For the first one, let  $u, v \in V$ . In view of (3.1), there is a nonnegative integer  $k$  such that

$$(z_1 - z_2)^k [Y_W(u, z_1), Y_W(v, z_2)] = 0.$$

It then follows from Proposition 2.6 that

$$[Y_W(u, z_1), Y_W(v, z_2)] = \sum_{j \geq 0} Y_W(u, z_2)_j^\phi Y_W(v, z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(z_1) \delta \left( \frac{z_2}{z_1} \right).$$

This together with (3.4) proves the first assertion.  $\square$

Next we are going to give a conceptual construction of  $\phi$ -coordinated quasi modules for vertex algebras. The following is a  $\phi$ -generalization of [Li5, Proposition 4.3].

**Proposition 3.6.** *Let  $a(z), b(z), c(z) \in \mathcal{E}(W)$ . Assume that*

$$f(z_1, z_2)[a(z_1), b(z_2)] = g(z_1, z_2)[a(z_1), c(z_2)] = h(z_1, z_2)[b(z_1), c(z_2)] = 0$$

*for some nonzero polynomials  $f(z_1, z_2), g(z_1, z_2), h(z_1, z_2)$ . Then for any  $n \in \mathbb{Z}$ , there exists a nonnegative integer  $k$  such that*

$$g(z_1, z_2)^k h(z_1, z_2) [a(z_1)_n^\phi b(z_1), c(z_2)] = 0.$$

*Proof.* Fix an  $n \in \mathbb{Z}$  and take a nonnegative integer  $k$  such that

$$z_0^{k+n} f(\phi(z, z_0), z)^{-1} \in \mathbb{C}((z))[[z_0]].$$

Using this and the fact

$$\phi(\phi(z, z_0), -z_0) = \phi(z, 0) = z = \phi(\phi(z, -z_0), z_0),$$

we obtain

$$\begin{aligned} & g(z_2, z_3)^k h(z_2, z_3) (a(z_2)_n^\phi b(z_2)) c(z_3) \\ &= \text{Res}_{z_0} z_0^n g(z_2, z_3)^k h(z_2, z_3) \left( Y_{\mathcal{E}}^\phi(a(z_2), z_0) b(z_2) \right) c(z_3) \\ &= \text{Res}_{z_1} \text{Res}_{z_0} z_0^n g(z_2, z_3)^k h(z_2, z_3) f(\phi(z_2, z_0), z_2)^{-1} \end{aligned}$$

$$\begin{aligned}
& z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) (f(z_1, z_2) a(z_1) b(z_2)) c(z_3) \\
&= \text{Res}_{z_1} \text{Res}_{z_0} z_0^n g(\phi(z_1, -z_0), z_3)^k h(z_2, z_3) f(\phi(z_2, z_0), z_2)^{-1} \\
& \quad z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) (f(z_1, z_2) a(z_1) b(z_2)) c(z_3) \\
&= \text{Res}_{z_1} \text{Res}_{z_0} z_0^n \left( e^{-z_0 p(z_1) \frac{\partial}{\partial z_1}} g(z_1, z_3)^k \right) h(z_2, z_3) f(\phi(z_2, z_0), z_2)^{-1} \\
& \quad z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) (f(z_1, z_2) a(z_1) b(z_2)) c(z_3) \\
&= \text{Res}_{z_1} \text{Res}_{z_0} \sum_{i=0}^{k-1} z_0^{n+i} \left( \left( -p(z_1) \frac{\partial}{\partial z_1} \right)^{(i)} g(z_1, z_3)^k \right) h(z_2, z_3) f(\phi(z_2, z_0), z_2)^{-1} \\
& \quad z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) (f(z_1, z_2) a(z_1) b(z_2)) c(z_3) \\
&= \text{Res}_{z_1} \text{Res}_{z_0} \sum_{i=0}^{k-1} z_0^{n+i} \left( \left( -p(z_1) \frac{\partial}{\partial z_1} \right)^{(i)} g(z_1, z_3)^k \right) h(z_2, z_3) c(z_3) \\
& \quad f(\phi(z_2, z_0), z_2)^{-1} z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) (f(z_1, z_2) a(z_1) b(z_2)) \\
&= \text{Res}_{z_1} \text{Res}_{z_0} z_0^n \left( e^{-z_0 p(z_1) \frac{\partial}{\partial z_1}} g(z_1, z_3)^k \right) h(z_2, z_3) c(z_3) f(\phi(z_2, z_0), z_2)^{-1} \\
& \quad z_1^{-1} \delta \left( \frac{\phi(z_2, z_0)}{z_1} \right) (f(z_1, z_2) a(z_1) b(z_2)) \\
&= g(z_2, z_3)^k h(z_2, z_3) c(z_3) (a(z_2) \phi_n b(z_2)).
\end{aligned}$$

□

A subset  $S$  of  $\mathcal{E}(W)$  is said to be (*resp. quasi*) *local* if any pair in  $S$  is (*resp. quasi*) local, and a (*resp. quasi*) local subspace  $U$  of  $\mathcal{E}(W)$  is said to be  $Y_{\mathcal{E}}^\phi$ -*closed* if

$$a(z) \phi_n b(z) \in U \quad \text{for all } a(z), b(z) \in U, n \in \mathbb{Z}.$$

In view of Proposition 3.6, we can formulate the following notion.

**Definition 3.7.** Let  $S$  be a quasi local subset of  $\mathcal{E}(W)$ . We define  $\langle S \rangle_\phi$  to be the smallest  $Y_{\mathcal{E}}^\phi$ -closed quasi local subspace of  $\mathcal{E}(W)$  which contains  $S$  and  $1_W$ .

We need the following  $\phi$ -generalization of [Li5, Proposition 4.8] for later use.

**Proposition 3.8.** For  $a(z), b(z) \in \mathcal{E}(W)$ , if

$$f(z_1, z_2) [a(z_1), b(z_2)] = 0$$

for some  $f(z_1, z_2) \in \mathbb{C}[z_1, z_2]$ , then

$$(z_1 - z_2)^k [Y_{\mathcal{E}}^\phi(a(z), z_1), Y_{\mathcal{E}}^\phi(b(z), z_2)] = 0,$$

where  $k$  is the order of the zero of  $f(z_1, z_2)$  at  $z_1 = z_2$ .

*Proof.* Let  $a(z), b(z), f(z_1, z_2)$  and  $k$  be as in the proposition. Then it follows from [JKLT, Proposition 2.18] that

$$(3.5) \quad f(\phi(z, z_1), \phi(z, z_2)) [Y_{\mathcal{E}}^{\phi}(a(z), z_1), Y_{\mathcal{E}}^{\phi}(b(z), z_2)] = 0.$$

Write

$$f(z_1, z_2) = (z_1 - z_2)^k f_0(z_1, z_2)$$

for some polynomial  $f_0(z_1, z_2)$  with  $f_0(z_1, z_1) \neq 0$ . Note that  $\phi(z, 0) = z$ . This implies that  $f_0(\phi(z, 0), \phi(z, 0))$  is nonzero and hence  $f_0(\phi(z, z_1), \phi(z, z_2))$  is invertible in  $\mathbb{C}((z))[[z_1, z_2]]$ . On the other hand, from the definition of  $\phi(z, z_1) = e^{z_1 p(z) \frac{d}{dz}}(z)$ , we have

$$\phi(z, z_1) - \phi(z, z_2) = (z_1 - z_2)g(z, z_1, z_2)$$

for some  $g(z, z_1, z_2) \in \mathbb{C}((z))[[z_1, z_2]]$ . Furthermore, we have  $g(z, 0, 0) = p(z) \neq 0$  and hence  $g(z, z_1, z_2)$  is also invertible in  $\mathbb{C}((z))[[z_1, z_2]]$ . Thus, we have shown that  $f(\phi(z, z_1), \phi(z, z_2))$  is a product of  $(z_1 - z_2)^k$  and an invertible element in  $\mathbb{C}((z))[[z_1, z_2]]$ . This together with (3.5) proves the proposition.  $\square$

As the main result of this section we have:

**Theorem 3.9.** *Let  $S$  be a quasi local subset of  $\mathcal{E}(W)$ . Then  $(\langle S \rangle_{\phi}, Y_{\mathcal{E}}^{\phi}, 1_W)$  is a vertex algebra with  $S$  as a generating set and with  $W$  as a faithful  $\phi$ -coordinated quasi module on which  $Y_W(a(z), z_0) = a(z_0)$  for  $a(z) \in \langle S \rangle_{\phi}$ . Furthermore, if  $S$  is local, then  $W$  is a  $\phi$ -coordinated  $\langle S \rangle_{\phi}$ -module.*

*Proof.* It suffices to prove the case that  $S$  is quasi local. Recall from [Li2] that  $\langle S \rangle_{\phi}$  is a nonlocal vertex algebra with  $W$  as a faithful  $\phi$ -coordinated quasi  $\langle S \rangle_{\phi}$ -module. On the other hand, it follows from Proposition 3.8 that

$$\{Y_{\mathcal{E}}^{\phi}(a(z), z_0) \mid a(z) \in \langle S \rangle_{\phi}\}$$

is a local subset of  $\mathcal{E}(\langle S \rangle_{\phi})$ . This implies that  $\langle S \rangle_{\phi}$  is indeed a vertex algebra.  $\square$

#### 4. $\phi$ -COORDINATED $V_{\mathcal{C}}$ -MODULES

In this section, we first recall the notion of a conformal algebra  $\mathcal{C}$  [K] (also known as a vertex Lie algebra in [DLM, P]) and then determine the  $\phi$ -coordinated modules for the universal enveloping vertex algebra of  $\mathcal{C}$  in terms of restricted modules of a Lie algebra  $\widehat{\mathcal{C}}_{\phi}$  arising from  $\mathcal{C}$  and  $\phi$ .

**Definition 4.1.** A conformal algebra  $\mathcal{C}$  is a  $\mathbb{C}[\partial]$ -module with a  $\mathbb{C}$ -bilinear product  $a_n b$  for each  $n \in \mathbb{N}$  such that the following axioms hold ( $a, b, c \in \mathcal{C}$ ,  $m, n \in \mathbb{N}$ ):

$$(C0) \quad a_n b = 0 \quad \text{for } n \gg 0,$$

$$(C1) \quad (\partial a)_n b = -n a_{n-1} b,$$

$$(C2) \quad a_n b = - \sum_{j=0}^{\infty} (-1)^{j+n} \partial^{(j)}(b_{n+j} a),$$

$$(C3) \quad a_m(b_n c) = \sum_{j=0}^m \binom{m}{j} (a_j b)_{m+n-j} c + b_n(a_m c).$$

A linear map  $\varphi : \mathcal{C} \rightarrow \mathcal{C}'$  of conformal algebras is called a *homomorphism* if

$$\varphi \partial = \partial \varphi \quad \text{and} \quad \varphi(a_n b) = \varphi(a)_n \varphi(b) \quad \text{for } a, b \in \mathcal{C}, n \in \mathbb{N}.$$

Let  $\mathcal{C}$  be a conformal algebra. Recall that  $\phi$  is an associate of  $F(x, y)$ , which is determined by a nonzero  $p(x) \in \mathbb{C}((x))$ . We define a multiplication on the formal loop space  $\mathcal{C} \otimes \mathbb{C}((x))$  by

$$(4.1) \quad [a \otimes f(x), b \otimes g(x)] = \sum_{i \geq 0} (a_i b) \otimes \left( \left( p(x) \frac{d}{dx} \right)^{(i)} f(x) \right) g(x)$$

for  $a, b \in \mathcal{C}$  and  $f(x), g(x) \in \mathbb{C}((x))$ . Form the quotient space

$$\widehat{\mathcal{C}}_\phi := \mathcal{C} \otimes \mathbb{C}((x)) / \text{Im} \left( \partial \otimes 1 + 1 \otimes p(x) \frac{d}{dx} \right).$$

It was known from [K] that the multiplication (4.1) defines a Lie algebra structure on the quotient space  $\widehat{\mathcal{C}}_\phi$ . When  $\phi = x + z$ , i.e.,  $p(x) = 1$ , we also write  $\widehat{\mathcal{C}} = \widehat{\mathcal{C}}_\phi$ . For  $a \in \mathcal{C}$  and  $f(x) \in \mathbb{C}((x))$ , we set

$$\overline{a \otimes f(x)} = \rho(a \otimes f(x)),$$

where  $\rho : \mathcal{C} \otimes \mathbb{C}((x)) \rightarrow \widehat{\mathcal{C}}_\phi$  is the quotient map.

For  $a \in \mathcal{C}$ , we set

$$a_\phi(z) = \sum_{n \in \mathbb{Z}} \overline{a \otimes x^n p(x)} z^{-n-1} \in \widehat{\mathcal{C}}_\phi[[z, z^{-1}]].$$

**Definition 4.2.** A  $\widehat{\mathcal{C}}_\phi$ -module  $W$  is called *restricted* if  $a_\phi(z) \in \mathcal{E}(W)$  for any  $a \in \mathcal{C}$ .

One notices from definition that

$$(4.2) \quad \left( p(z) \frac{d}{dz} \right) a_\phi(z) = (\partial a)_\phi(z), \quad a \in \mathcal{C}.$$

Furthermore, in view of (4.1), we have for  $a, b \in \mathcal{C}$ ,

$$(4.3) \quad [a_\phi(z), b_\phi(w)] = \sum_{i \geq 0} (a_i b)_\phi(w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \bar{p}(z) \delta \left( \frac{w}{z} \right).$$

We have the following useful lemma.

**Lemma 4.3.** *Let  $W$  be a vector space and  $\mathcal{C}$  be a conformal algebra. Assume that there is a linear map*

$$\pi : \mathcal{C} \rightarrow \mathcal{E}(W), \quad a \mapsto a(z) = \sum_{n \in \mathbb{Z}} a_{[n]} z^{-n-1},$$

such that for  $a \in \mathcal{C}$ ,

$$(4.4) \quad \left( p(z) \frac{d}{dz} \right) a(z) = (\partial a)(z),$$

and for  $a, b \in \mathcal{C}$ ,

$$(4.5) \quad [a(z), b(w)] = \sum_{i \geq 0} (a_i b)(w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \bar{p}(z) \delta \left( \frac{w}{z} \right).$$

Then there is a restricted  $\widehat{\mathcal{C}}_\phi$ -module structure on  $W$  with  $a_\phi(z) = a(z)$  for  $a \in \mathcal{C}$ .

*Proof.* Let  $a \in \mathcal{C}$  and  $f(x) \in \mathbb{C}((x))$ . Set

$$f_0(x) = p(x)^{-1} f(x) = \sum_{n \geq n_0} b_n x^n,$$

where  $n_0 \in \mathbb{Z}$ . We define an action of  $\overline{a \otimes f(x)}$  on  $W$  by

$$(4.6) \quad \overline{a \otimes f(x)} w = \sum_{n=n_0}^{n_{a,w}} b_n a_{[n]} w, \quad w \in W,$$

where  $n_{a,w}$  is a sufficiently large integer such that  $a_{[m]} w = 0$  for all  $m > n_{a,w}$ . It follows from (4.4) that this action is well-defined. Furthermore, by comparing (4.3) with (4.5), one can check that (4.6) defines a restricted  $\widehat{\mathcal{C}}_\phi$ -module structure on  $W$  with  $a_\phi(z) = a(z)$  for  $a \in \mathcal{C}$ .  $\square$

Now we recall the notion of universal enveloping vertex algebra of a conformal algebra  $\mathcal{C}$ . Set  $\widehat{\mathcal{C}}_+ = \rho(\mathcal{C} \otimes \mathbb{C}[[x]])$ , which is a Lie subalgebra of  $\widehat{\mathcal{C}}$ . Let  $\mathbb{C}$  be the one-dimensional trivial  $\widehat{\mathcal{C}}_+$ -module. Then we have the induced  $\widehat{\mathcal{C}}$ -module

$$V_{\mathcal{C}} = \mathcal{U}(\widehat{\mathcal{C}}) \otimes_{\mathcal{U}(\widehat{\mathcal{C}}_+)} \mathbb{C}.$$

Set  $\mathbf{1} = 1 \otimes 1$ . It is known that  $\mathcal{C}$  can be identified as a subspace of  $V_{\mathcal{C}}$  through the linear map

$$a \mapsto \overline{a \otimes x^{-1} \mathbf{1}}, \quad a \in \mathcal{C}.$$

Note that each vertex algebra is naturally a conformal algebra by setting  $\partial = \mathcal{D}$  and forgetting the  $n$ -products with  $n < 0$ . The following result is well-known (cf. [K, DLM, P, FB]).



**Proposition 4.4.** *There is a vertex algebra structure on  $V_{\mathcal{C}}$  such that  $\mathbf{1}$  is the vacuum vector and that*

$$Y(a, z) = a(z) = \sum_{n \in \mathbb{Z}} \overline{a \otimes x^n} z^{-n-1}$$

for  $a \in \mathcal{C}$ . Moreover, for any vertex algebra  $V$ , if  $\varphi : \mathcal{C} \rightarrow V$  is a homomorphism of conformal algebras, then  $\varphi$  extends uniquely to a vertex algebra homomorphism from  $V_{\mathcal{C}}$  to  $V$ .

The following is the main result of this section.

**Theorem 4.5.** *Let  $\mathcal{C}$  be a conformal algebra. Then any restricted  $\widehat{\mathcal{C}}_\phi$ -module  $W$  is naturally a  $\phi$ -coordinated  $V_{\mathcal{C}}$ -module with  $Y_W(a, z) = a_\phi(z)$  for  $a \in \mathcal{C}$ . On the other hand, any  $\phi$ -coordinated  $V_{\mathcal{C}}$ -module  $W$  is naturally a restricted  $\widehat{\mathcal{C}}_\phi$ -module with  $a_\phi(z) = Y_W(a, z)$  for  $a \in \mathcal{C}$ .*

*Proof.* Let  $W$  be a restricted  $\widehat{\mathcal{C}}_\phi$ -module. Then it follows from (4.3) that

$$S = \{a_\phi(z) \mid a \in \mathcal{C}\}$$

is a local subset of  $\mathcal{E}(W)$ . In view of Theorem 3.9, there is a vertex algebra  $\langle S \rangle_\phi$  with  $W$  as a faithful  $\phi$ -coordinated  $\langle S \rangle_\phi$ -module on which  $Y_W(b(z), z_0) = b(z_0)$  for  $b(z) \in \langle S \rangle_\phi$ . This together with (4.3) implies that for  $a, b \in \mathcal{C}$ ,

$$\begin{aligned} & [Y_W(a_\phi(z_0), z), Y_W(b_\phi(z_0), w)] = [a_\phi(z), b_\phi(w)] \\ &= \sum_{i \geq 0} (a_i b)_\phi(w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \bar{p}(z) \delta \left( \frac{w}{z} \right). \end{aligned}$$

As  $W$  is faithful, it follows from Proposition 3.5 that

$$(4.7) \quad a_\phi(z)_i^\phi b_\phi(z) = (a_i b)_\phi(z), \quad a, b \in \mathcal{C}, i \in \mathbb{N}.$$

We define a linear map

$$\varphi : \mathcal{C} \rightarrow \langle S \rangle_\phi, \quad a \mapsto a_\phi(z).$$

It follows from (3.3) and (4.2) that for  $a \in \mathcal{C}$ ,

$$\mathcal{D}\varphi(a) = \left( p(z) \frac{d}{dz} \right) a_\phi(z) = (\partial a)_\phi(z) = \varphi \partial(a).$$

This together with (4.7) implies that  $\varphi$  is a homomorphism of conformal algebras. Then Proposition 4.4 says that  $\varphi$  can be extended to a vertex algebra homomorphism from  $V_{\mathcal{C}}$  to  $\langle S \rangle_\phi$ . Thus,  $W$  becomes a  $\phi$ -coordinated  $V_{\mathcal{C}}$ -module with

$$Y_W(a, w) = Y_W(\varphi(a), w) = Y_W(a_\phi(z), w) = a_\phi(w) \quad \text{for } a \in \mathcal{C}.$$

On the other hand, let  $W$  be a  $\phi$ -coordinated  $V_{\mathcal{C}}$ -module. Recall from (3.3) that for  $a \in \mathcal{C}$ ,

$$Y_W(\partial a, z) = Y_W(\mathcal{D}a, z) = \left( p(z) \frac{d}{dz} \right) Y_W(a, z),$$

and recall from Proposition 3.5 that for  $a, b \in \mathcal{C}$ ,

$$[Y_W(a, z), Y_W(b, w)] = \sum_{i \geq 0} Y_W(a_i b, w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \bar{p}(z) \delta \left( \frac{w}{z} \right).$$

Combining this with Lemma 4.3, we see that  $W$  is a restricted  $\widehat{\mathcal{C}}_\phi$ -module with  $a_\phi(z) = Y_W(a, z)$  for  $a \in \mathcal{C}$ , as desired.  $\square$

**Remark 4.6.** When  $p(x) \in \mathbb{C}[x, x^{-1}]$  is a Laurent polynomial, the Lie algebra  $\widehat{\mathcal{C}}_\phi$  can be defined in the following usual way:

$$\widehat{\mathcal{C}}_\phi := \mathcal{C} \otimes \mathbb{C}[x, x^{-1}] / \text{Im}(\partial \otimes 1 + 1 \otimes p(x) \frac{d}{dx}).$$

However, for the case  $p(x) \in \mathbb{C}((x))$ , the derivation  $p(x) \frac{d}{dx}$  may be not well-defined on  $\mathbb{C}[x, x^{-1}]$ . So we need to enlarge the usual loop algebra  $\mathcal{C} \otimes \mathbb{C}[x, x^{-1}]$  to the formal loop algebra  $\mathcal{C} \otimes \mathbb{C}((x))$ .

## 5. $(G, \chi_\phi)$ -EQUIVARIANT $\phi$ -COORDINATED QUASI MODULES

Let  $G$  be a group and let  $\chi, \chi_\phi : G \rightarrow \mathbb{C}^\times$  be two linear characters on  $G$ . In this section we generalize the previous results on  $\phi$ -coordinated modules for vertex algebras to the  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for  $(G, \chi)$ -vertex algebras. We start with the notion of  $(G, \chi)$ -vertex algebra in [JKLT] (cf. [Li3, Li5]).

**Definition 5.1.** A  $(G, \chi)$ -vertex algebra is a pair  $(V, R)$  consisting of a vertex algebra  $V$  and a group homomorphism  $R : G \rightarrow \text{GL}(V)$  such that  $R_g(\mathbf{1}) = \mathbf{1}$  and

$$R_g Y(v, z) R_g^{-1} = Y(R_g(v), \chi(g)z) \quad \text{for } g \in G, v \in V.$$

For  $(G, \chi)$ -vertex algebras  $(V, R), (V', R')$ , a linear map  $\psi : V \rightarrow V'$  is called a  $(G, \chi)$ -vertex algebra homomorphism if  $\psi$  is a vertex algebra homomorphism and  $\psi \circ R_g = R'_g \circ \psi$  for  $g \in G$ .

For a subgroup  $\Gamma$  of  $\mathbb{C}^\times$ , we denote by  $\mathbb{C}_\Gamma[x]$  the set of all monic polynomials in  $\mathbb{C}[x]$  whose roots contained in  $\Gamma$ . The following notion was introduced in [JKLT].

**Definition 5.2.** Let  $(V, R)$  be a  $(G, \chi)$ -vertex algebra. A  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V$ -module is a  $\phi$ -coordinated quasi  $V$ -module  $(W, Y_W)$  such that

$$(5.1) \quad Y_W(R_g v, z) = Y_W(v, \chi_\phi(g)^{-1} z) \quad \text{for } g \in G, v \in V,$$

and that for  $u, v \in V$ , there exists  $q(z) \in \mathbb{C}_{\chi_\phi(G)}[z]$  such that

$$(5.2) \quad q(z_1/z_2) Y_W(u, z_1) Y_W(v, z_2) \in \text{Hom}(W, W((z_1, z_2))).$$

Let  $(V, R)$  be a  $(G, \chi)$ -vertex algebra and let  $(W, Y_W)$  be a  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V$ -module. If  $\frac{d}{dz}Y_W(v, z) \neq 0$  for some  $v \in V$ , then it was shown in [JKLT] that

$$(5.3) \quad \phi(x, \chi(g)z) = \chi_\phi(g)\phi(\chi_\phi(g)^{-1}x, z) \quad \text{for } g \in G.$$

Note that this relation is equivalent to

$$(5.4) \quad p(\chi_\phi(g)x) = \chi(g)^{-1}\chi_\phi(g)p(x) \quad \text{for } g \in G.$$

On the other hand, if  $\frac{d}{dz}Y_W(v, z) = 0$  for any  $v \in V$ , then the quotient vertex algebra  $V/\ker Y_W$  is simply an associative algebra and  $G$  acts trivially on  $V/\ker Y_W$ . In view of this, we always assume that the characters  $\chi$  and  $\chi_\phi$  satisfy the relation (5.3). Note that we have  $\chi = \chi_\phi$  if  $p(x) = 1$  and  $\chi = 1$  if  $p(x) = x$ .

As a generalization of Proposition 3.5, we have:

**Proposition 5.3.** *Let  $(V, R)$  be a  $(G, \chi)$ -vertex algebra and let  $(W, Y_W)$  be a  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V$ -module. Let  $\psi : \chi_\phi(G) \rightarrow G$  be a section of  $\chi_\phi$ . Then for  $u, v \in V$ ,*

$$(5.5) \quad \begin{aligned} & [Y_W(u, z_1), Y_W(v, z_2)] \\ &= \text{Res}_{z_0} \sum_{g \in \psi(\chi_\phi(G))} Y_W(Y(R_{g^{-1}}u, z_0)v, z_2) e^{z_0 p(z_2) \frac{\partial}{\partial z_2}} \left( \chi(g) \bar{p}(z_1) \delta \left( \frac{\chi_\phi(g) z_2}{z_1} \right) \right)). \end{aligned}$$

*Proof.* Let  $u, v \in V$ . From (5.2) and (3.1), it follows that there exists a  $q(z) \in \mathbb{C}_{\chi_\phi(G)}[z]$  such that

$$q(z_1/z_2)[Y_W(u, z_1), Y_W(v, z_2)] = 0.$$

Then by applying Proposition 2.6 we have

$$\begin{aligned} & [Y_W(u, z_1), Y_W(v, z_2)] \\ &= \sum_{\lambda \in \chi_\phi(G)} \sum_{j \geq 0} Y_W(u, \lambda z_2)_j^\phi Y_W(v, z_2) \left( p(z_2) \frac{\partial}{\partial z_2} \right)^{(j)} \bar{p}(\lambda^{-1} z_1) \delta \left( \frac{\lambda z_2}{z_1} \right). \end{aligned}$$

Note that (5.4) implies that for  $g \in G$ ,

$$\bar{p}(\chi_\phi(g)^{-1} z_1) = \chi_\phi(g) z_1^{-1} p(\chi_\phi(g^{-1}) z_1) = \chi(g) z_1^{-1} p(z_1) = \chi(g) \bar{p}(z_1),$$

together with  $\psi$  a section of  $\chi_\phi$ , we obtain

$$\begin{aligned} & [Y_W(u, z_1), Y_W(v, z_2)] \\ &= \text{Res}_{z_0} \sum_{g \in \psi(\chi_\phi(G))} Y_\mathcal{E}^\phi(Y_W(u, \chi_\phi(g) z_2), z_0) Y_W(v, z_2) e^{z_0 p(z_2) \frac{\partial}{\partial z_2}} \left( \chi(g) \bar{p}(z_1) \delta \left( \frac{\chi_\phi(g) z_2}{z_1} \right) \right). \end{aligned}$$

From (5.1) and (3.4), we have

$$Y_\mathcal{E}^\phi(Y_W(u, \chi_\phi(g) z), z_0) Y_W(v, z) = Y_\mathcal{E}^\phi(Y_W(R_{g^{-1}}u, z), z_0) Y_W(v, z) = Y_W(Y(R_{g^{-1}}u, z_0)v, z)$$

for  $g \in G$ . Thus (5.5) follows.  $\square$

**Definition 5.4.** Let  $W$  be a vector space and let  $\Gamma$  be a subgroup of  $\mathbb{C}^\times$ . A subset  $S$  of  $\mathcal{E}(W)$  is said to be  $\Gamma$ -quasi local if for any pair  $a(z), b(z)$  in  $S$ , there exists  $q(z) \in \mathbb{C}_\Gamma[z]$  such that

$$q(z_1/z_2)[a(z_1), b(z_2)] = 0,$$

and is said to be  $\Gamma$ -stable if for any  $a(z) \in S$  and  $\lambda \in \Gamma$ ,  $a(\lambda z) \in S$ .

For a  $\Gamma$ -quasi local and  $\Gamma$ -stable subset  $S$  of  $\mathcal{E}(W)$ ,  $\langle S \rangle_\phi$  is also  $\Gamma$ -quasi local (Theorem 3.6) and  $\Gamma$ -stable ([JKLT]). Furthermore, from [JKLT, Theorem 3.11] and Theorem 3.9, we immediately have the following result.

**Theorem 5.5.** *Let  $S$  be a  $\chi_\phi(G)$ -stable and  $\chi_\phi(G)$ -quasi local subset of  $\mathcal{E}(W)$ . Assume that the character  $\chi_\phi$  is injective. Then the vertex algebra  $\langle S \rangle_\phi$  together with the group homomorphism*

$$R : G \rightarrow \text{GL}(\langle S \rangle_\phi), \quad g \mapsto R_g$$

*is a  $(G, \chi)$ -vertex algebra and  $W$  is a faithful  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi  $\langle S \rangle_\phi$ -module with  $Y_W(a(z), z_0) = a(z_0)$  for  $a(z) \in \langle S \rangle_\phi$ , where  $R_g(a(z)) = a(\chi_\phi(g)^{-1}z)$ .*

The following notion was introduced in [Li3]:

**Definition 5.6.** A  $(G, \chi)$ -conformal algebra  $(\mathcal{C}, R)$  is a conformal algebra  $\mathcal{C}$  together with a group homomorphism

$$R : G \rightarrow \text{GL}(\mathcal{C}), \quad g \mapsto R_g,$$

such that for any  $a, b \in \mathcal{C}$ ,

$$(CR1) \quad \partial R_g = \chi(g)^{-1} R_g \partial,$$

$$(CR2) \quad R_g(a_m b) = \chi(g)^{-(m+1)} (R_g a)_m (R_g b) \quad \text{for } m \in \mathbb{N},$$

$$(CR3) \quad (R_g a)_m b = 0 \quad \text{for all but finitely many } m \in \mathbb{N}, g \in G.$$

For  $(G, \chi)$ -conformal algebras  $(\mathcal{C}, R)$ ,  $(\mathcal{C}', R')$ , a linear map  $\psi : \mathcal{C} \rightarrow \mathcal{C}'$  is called a  $(G, \chi)$ -conformal algebra homomorphism if  $\psi$  is a conformal algebra homomorphism and  $\psi \circ R_g = R'_g \circ \psi$  for  $g \in G$ .

Let  $(\mathcal{C}, R)$  be a  $(G, \chi)$ -conformal algebra. For any  $g \in G$ , we define a linear automorphism  $\hat{R}_g$  on  $\mathcal{C} \otimes \mathbb{C}((x))$  by

$$(5.6) \quad \hat{R}_g(a \otimes f(x)) = \chi(g)^{-1} R_g(a) \otimes f(\chi_\phi(g)^{-1}x), \quad a \in \mathcal{C}, f(x) \in \mathbb{C}((x)).$$

**Lemma 5.7.** *For  $g \in G$ , one has*

$$\hat{R}_g \circ (\partial \otimes 1 + 1 \otimes p(x) \frac{d}{dx}) = \chi(g) (\partial \otimes 1 + 1 \otimes p(x) \frac{d}{dx}) \circ \hat{R}_g.$$

*Proof.* For  $a \in \mathcal{C}$ ,  $f(x) \in \mathbb{C}((x))$ , we have

$$\begin{aligned} & \hat{R}_g(\partial a \otimes f(x) + a \otimes p(x) \frac{df(x)}{dx}) \\ &= \chi(g)^{-1} \left( R_g(\partial a) \otimes f(\chi_\phi(g)^{-1}x) + R_g a \otimes p(\chi_\phi(g)^{-1}x) \left( \frac{df(x)}{dx} \Big|_{x=\chi_\phi(g)^{-1}x} \right) \right) \\ &= (\partial R_g a) \otimes f(\chi_\phi(g)^{-1}x) + \chi_\phi(g)^{-1} R_g a \otimes p(x) \left( \frac{df(x)}{dx} \Big|_{x=\chi_\phi(g)^{-1}x} \right) \\ &= \chi(g) (\partial \otimes 1 + 1 \otimes p(x) \frac{d}{dx}) \hat{R}_g(a \otimes f(x)), \end{aligned}$$

where in the second equality we have used the relations (CR1) and (5.4).  $\square$

In view of Lemma 5.7,  $\hat{R}_g$  induces a linear automorphism on  $\hat{\mathcal{C}}_\phi$ , which we still denote as  $\hat{R}_g$ . Then we have:

**Lemma 5.8.** *For  $g \in G$ ,  $\hat{R}_g$  is a Lie automorphism of  $\hat{\mathcal{C}}_\phi$ . Furthermore, for any  $u, v \in \hat{\mathcal{C}}_\phi$ ,  $[\hat{R}_g(u), v] = 0$  for all but finitely many  $g \in G$ .*

*Proof.* For  $a, b \in \mathcal{C}$  and  $f(x), g(x) \in \mathbb{C}((x))$  we have

$$\begin{aligned} & [\hat{R}_g(a \otimes f(x)), \hat{R}_g(b \otimes g(x))] \\ &= \chi(g)^{-2} [R_g(a) \otimes f(\chi_\phi(g)^{-1}x), R_g(b) \otimes g(\chi_\phi(g)^{-1}x)] \\ &= \chi(g)^{-2} \sum_{i \geq 0} ((R_g a)_i (R_g b)) \otimes \left( \left( p(x) \frac{d}{dx} \right)^{(i)} f(\chi_\phi(g)^{-1}x) \right) g(\chi_\phi(g)^{-1}x) \\ &= \chi(g)^{-1+i} \sum_{i \geq 0} (R_g(a_i b)) \otimes \left( \left( p(x) \frac{d}{dx} \right)^{(i)} f(\chi_\phi(g)^{-1}x) \right) g(\chi_\phi(g)^{-1}x) \\ &= \hat{R}_g([a \otimes f(x), b \otimes g(x)]), \end{aligned}$$

where in the third and last equality we have used the relations (CR2) and (5.4), respectively. This proves that  $\hat{R}_g$  is a Lie automorphism and the second assertion follows from (CR3).  $\square$

Following [Li3], we define a multiplication on  $\hat{\mathcal{C}}_\phi$  by

$$(5.7) \quad (u, v) \mapsto \sum_{g \in G} [\hat{R}_g(u), v], \quad u, v \in \hat{\mathcal{C}}_\phi.$$

In view of Lemma 5.8, it was known that under this new multiplication, the quotient space

$$\hat{\mathcal{C}}_\phi[G] := \hat{\mathcal{C}}_\phi / \text{span}\{\hat{R}_g(u) - u \mid g \in G, u \in \hat{\mathcal{C}}_\phi\}$$

is a Lie algebra [Li3]. For convenience, for  $a \in \mathcal{C}$ ,  $f(x) \in \mathbb{C}((x))$ , we still denote the image of  $a \otimes f(x)$  in  $\hat{\mathcal{C}}_\phi[G]$  by  $\overline{a \otimes f(x)}$ .

**Remark 5.9.** When  $G$  is a finite group,  $\widehat{\mathcal{C}}_\phi[G]$  is obviously isomorphic to the subalgebra of  $\widehat{\mathcal{C}}_\phi$  fixed by  $\widehat{R}_g$  for any  $g \in G$ .

For  $a \in \mathcal{C}$ , we set

$$a_\phi^G(z) = \sum_{n \in \mathbb{Z}} \overline{a \otimes x^n p(x)} z^{-n-1} \in \widehat{\mathcal{C}}_\phi[G][[z, z^{-1}]].$$

We have

**Definition 5.10.** A  $\widehat{\mathcal{C}}_\phi[G]$ -module  $W$  is called *restricted* if  $a_\phi^G(z) \in \mathcal{E}(W)$  for any  $a \in \mathcal{C}$ .

From (5.4), we have that for  $a \in \mathcal{C}$ ,

$$(5.8) \quad (R_g a)_\phi^G(z) = a_\phi^G(\chi_\phi(g)^{-1}z).$$

Furthermore, for  $a, b \in \mathcal{C}$ , we have

$$(5.9) \quad \begin{aligned} [a_\phi^G(z), b_\phi^G(w)] &= \sum_{g \in G} [(R_g a)_\phi^G(\chi_\phi(g)z), b_\phi^G(w)] \\ &= \sum_{g \in G} \sum_{i \geq 0} ((R_{g^{-1}} a)_i b)_\phi^G(w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \left( \chi(g) \bar{p}(z) \delta \left( \frac{\chi_\phi(g)w}{z} \right) \right). \end{aligned}$$

The same proof as that of Lemma 4.3, we have

**Lemma 5.11.** *Let  $W$  be a vector space. Assume that there is a linear map*

$$\pi : \mathcal{C} \rightarrow \mathcal{E}(W), \quad a \mapsto a(z) = \sum_{n \in \mathbb{Z}} a_{[n]} z^{-n-1},$$

such that for  $a \in \mathcal{C}$ ,

$$(5.10) \quad \left( p(z) \frac{\partial}{\partial z} \right) a(z) = (\partial a)(z), \quad (R_g a)(z) = a(\chi_\phi(g)^{-1}z),$$

and for  $a, b \in \mathcal{C}$ ,

$$[a(z), b(w)] = \sum_{g \in G} \sum_{i \geq 0} ((R_{g^{-1}} a)_i b)(w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \left( \chi(g) \bar{p}(z) \delta \left( \frac{\chi_\phi(g)w}{z} \right) \right).$$

Then there is a restricted  $\widehat{\mathcal{C}}_\phi[G]$ -module structure on  $W$  with  $a_\phi^G(z) = a(z)$  for  $a \in \mathcal{C}$ .

Let  $(\mathcal{C}, R)$  be a  $(G, \chi)$ -conformal algebra and set  $H = \ker \chi_\phi$ . One notices from (5.4) that  $\ker \chi_\phi \subset \ker \chi$ , so  $\chi$  induces a linear character on  $G/H$ , denoted as  $\bar{\chi}$ . Following [Li3], we have a  $(G/H, \bar{\chi})$ -conformal algebra  $(\mathcal{C}/H, R)$  induced from  $(\mathcal{C}, R)$ . Explicitly, as a vector space,

$$\mathcal{C}/H = \mathcal{C} / \text{span}\{R_h a - a \mid h \in H, a \in \mathcal{C}\}.$$

Define  $\partial(aH) = (\partial a)H$  for  $aH \in \mathcal{C}/H$ , and the  $n$ -products on  $\mathcal{C}/H$  by

$$(uH)_n(vH) = \sum_{h \in H} ((R_h u)_n v)H \quad \text{for } u, v \in \mathcal{C}, n \in \mathbb{N},$$

we also define the  $G/H$ -action on  $\mathcal{C}/H$  by  $R_{gH}(uH) = (R_g u)H$  for  $g \in G, u \in \mathcal{C}$ . Denote by  $\bar{\chi}_\phi$  the character on  $G/H$  induced by  $\chi_\phi$ . Note that the characters  $\bar{\chi}, \bar{\chi}_\phi$  still satisfy the relation (5.4). It is known that the  $G/H$ -action  $R$  on  $\mathcal{C}/H$  can be extended uniquely to a  $G/H$ -action on  $V_{\mathcal{C}/H}$ , we still denote it by  $R$ , then  $(V_{\mathcal{C}/H}, R)$  becomes a  $(G/H, \bar{\chi})$ -vertex algebra [Li3]. In particular we have a notion of  $(G/H, \bar{\chi}_\phi)$ -equivariant  $\phi$ -coordinated quasi module for the  $(G/H, \bar{\chi})$ -vertex algebra  $V_{\mathcal{C}/H}$ . The following theorem is the main result of this section.

**Theorem 5.12.** *Let  $(\mathcal{C}, R)$  be a  $(G, \chi)$ -conformal algebra and set  $H = \ker \chi_\phi$ . Then any restricted  $\widehat{\mathcal{C}}_\phi[G]$ -module  $W$  is a  $(G/H, \bar{\chi}_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V_{\mathcal{C}/H}$ -module with  $Y_W(aH, z) = a_\phi^G(z)$  for  $a \in \mathcal{C}$ . On the other hand, any  $(G/H, \bar{\chi}_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V_{\mathcal{C}/H}$ -module  $W$  is a restricted  $\widehat{\mathcal{C}}_\phi[G]$ -module with  $a_\phi^G(z) = Y_W(aH, z)$  for  $a \in \mathcal{C}$ .*

*Proof.* Let  $W$  be a restricted  $\widehat{\mathcal{C}}_\phi[G]$ -module. Then  $S = \{a_\phi^G(z) \mid a \in \mathcal{C}\}$  is a  $\chi_\phi(G)$ -quasi local and  $\chi_\phi(G)$ -stable subset of  $\mathcal{E}(W)$ . Note that the character  $\bar{\chi}_\phi : G/H \rightarrow \mathbb{C}^\times$  is injective and its image is  $\chi_\phi(G)$ . In view of the Theorem 5.5,  $\langle S \rangle_\phi$  is a  $(G/H, \bar{\chi})$ -vertex algebra and  $W$  is a faithful  $(G/H, \bar{\chi}_\phi)$ -equivariant  $\phi$ -coordinated quasi  $\langle S \rangle_\phi$ -module with  $Y_W(a(z), z_0) = a(z_0)$  for  $a(z) \in \langle S \rangle_\phi$ . Note that for  $h \in H$  and  $a \in \mathcal{C}$ , from (5.8) we have that

$$(R_h a)_\phi^G(z) = a_\phi^G(\chi_\phi(h)^{-1}z) = a_\phi^G(z).$$

Thus the linear map

$$\varphi : \mathcal{C}/H \rightarrow \langle S \rangle_\phi, \quad aH \mapsto a_\phi^G(z)$$

is well-defined. Take a section  $\psi : \chi_\phi(G) \rightarrow G$  of  $\chi_\phi$ . For  $a, b \in \mathcal{C}$ , we have

$$\begin{aligned} & [a_\phi^G(z), b_\phi^G(w)] \\ &= \sum_{g \in G} \sum_{i \geq 0} ((R_{g^{-1}} a)_i b)_\phi^G(w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \left( \chi(g) \bar{p}(z) \delta \left( \frac{\chi_\phi(g)w}{z} \right) \right) \\ &= \sum_{g \in \psi(\chi_\phi(G))} \sum_{i \geq 0} \sum_{h \in H} ((R_{(gh)^{-1}} a)_i b)_\phi^G(w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \left( \chi(g) \bar{p}(z) \delta \left( \frac{\chi_\phi(g)w}{z} \right) \right). \end{aligned}$$

By comparing this with (5.5) we have

$$\sum_{h \in H} ((R_{(gh)^{-1}} a)_i b)_\phi^G(w) = (R_{g^{-1}} a)_\phi^G(w) \phi_i b_\phi^G(w).$$

This particular implies that

$$\varphi(aH)_i^\phi \varphi(bH) = \varphi((aH)_i(bH)), \quad a, b \in \mathcal{C}, i \in \mathbb{N}.$$

Furthermore, for any  $a \in \mathcal{C}$ , we have

$$(\mathcal{D} \circ \varphi)(aH) = \mathcal{D}(a_\phi^G(z)) = \left( p(z) \frac{\partial}{\partial z} \right) a_\phi^G(z) = (\partial a)_\phi^G(z) = (\varphi \circ \partial)(aH).$$

Thus,  $\varphi$  is a conformal algebra homomorphism. From the Proposition 4.4, there is a vertex algebra homomorphism from  $V_{\mathcal{C}/H}$  to  $\langle S \rangle_\phi$  which extends  $\varphi$ , we also denote it as  $\varphi$ . Note that for  $a \in \mathcal{C}$  and  $g \in G$ ,

$$\begin{aligned} (R_{gH} \circ \varphi)(aH) &= R_{gH} a_\phi^G(z) = a_\phi^G(\bar{\chi}_\phi(gH)^{-1}z) \\ &= a_\phi^G(\chi_\phi(g)^{-1}z) = (R_g a)_\phi^G(z) = \varphi((R_g a)H) = (\varphi \circ R_{gH})(aH). \end{aligned}$$

Then for  $a, b \in \mathcal{C}$  and  $g \in G$ , we have

$$\begin{aligned} (R_{gH} \circ \varphi)(Y(aH, z)bH) &= R_{gH}(Y_\mathcal{E}(\varphi(aH), z)\varphi(bH)) \\ &= Y_\mathcal{E}(R_{gH}(\varphi(aH)), \bar{\chi}(gH)^{-1}z)R_{gH}(\varphi(bH)) = Y_\mathcal{E}(\varphi(R_{gH}aH), \bar{\chi}(gH)^{-1}z)\varphi(R_{gH}bH) \\ &= \varphi(Y(R_{gH}(aH), \bar{\chi}(gH)^{-1}z)R_{gH}(bH)) = (\varphi \circ R_{gH})(Y(aH, z)bH). \end{aligned}$$

This together with the fact that  $\mathcal{C}/H$  is a generating set of  $V_{\mathcal{C}/H}$  proves that  $R_{gH} \circ \varphi = \varphi \circ R_{gH}$  on  $V_{\mathcal{C}/H}$ . Thus  $\varphi$  is a  $(G/H, \bar{\chi})$ -vertex algebra homomorphism from  $V_{\mathcal{C}/H}$  to  $\langle S \rangle_\phi$ . Via this homomorphism,  $W$  is naturally a  $(G/H, \bar{\chi}_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V_{\mathcal{C}/H}$ -module.

On the other hand, let  $W$  be a  $(G/H, \bar{\chi}_\phi)$ -equivariant  $\phi$ -coordinated quasi  $V_{\mathcal{C}/H}$ -module. From (3.3) and (5.8), for  $a \in \mathcal{C}$ , we have

$$\begin{aligned} Y_W(\partial(aH), z) &= Y_W(\mathcal{D}(aH), z) = \left( p(z) \frac{d}{dz} \right) Y_W(aH, z), \\ (R_g a)_\phi^G(z) &= a_\phi^G(\chi_\phi(g)^{-1}z). \end{aligned}$$

From Proposition 5.3, for  $a, b \in \mathcal{C}$ , we have

$$\begin{aligned} &[Y_W(aH, z), Y_W(bH, w)] \\ &= \sum_{g \in \psi(\chi_\phi(G))} \sum_{i \geq 0} \sum_{h \in H} Y_W(((R_{hg^{-1}}a)_i b)H, w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \left( \chi(g) \bar{p}(z) \delta \left( \frac{\chi_\phi(g)w}{z} \right) \right) \\ &= \sum_{g \in G} \sum_{i \geq 0} Y_W(((R_{g^{-1}}a)_i b)H, w) \left( p(w) \frac{\partial}{\partial w} \right)^{(i)} \left( \chi(g) \bar{p}(z) \delta \left( \frac{\chi_\phi(g)w}{z} \right) \right). \end{aligned}$$

Thus, from Lemma 5.11, we obtain that  $W$  is a restricted  $\widehat{\mathcal{C}}_\phi[G]$ -module with  $a_\phi^G(z) = Y_W(aH, z)$  for  $a \in \mathcal{C}$ .  $\square$



### 6. APPLICATIONS

In this section, as an application, a class of  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for affine vertex algebras is determined, and the  $\phi$ -coordinated modules for Virasoro vertex algebra and the vertex algebras associated to Novikov algebras are determined.

Firstly, let  $\mathfrak{g}$  be a (possibly infinite dimensional) Lie algebra, which is equipped with an invariant symmetric bilinear form  $(\cdot, \cdot)$ . Then we have the (formal) untwisted affine Lie algebra

$$\widehat{\mathfrak{g}} := \mathfrak{g} \otimes \mathbb{C}((x)) \oplus \mathbb{C}\mathbf{c},$$

where  $\mathbf{c}$  is central and for  $a, b \in \mathfrak{g}$ ,  $f(x), g(x) \in \mathbb{C}((x))$ ,

$$[a \otimes f(x), b \otimes g(x)] = [a, b] \otimes f(x)g(x) + (a, b) \operatorname{Res}_x \left( \left( \frac{d}{dx} f(x) \right) g(x) \right) \mathbf{c}.$$

On the other hand, we have the conformal algebra  $([\mathbf{K}])$

$$\mathcal{C}_{\mathfrak{g}} = (\mathbb{C}[\partial] \otimes \mathfrak{g}) \oplus \mathbb{C}\mathbf{c},$$

where for  $a, b \in \mathfrak{g}$  and  $n \in \mathbb{N}$ ,  $\partial(\partial^n \otimes a) = \partial^{n+1} \otimes a$ ,  $\partial(\mathbf{c}) = 0$ , and

$$a_n b = \delta_{n,0} [a, b] + \delta_{n,1} (a, b) \mathbf{c}, \quad a_n \mathbf{c} = 0 = \mathbf{c}_n \mathbf{c}.$$

We have

**Lemma 6.1.** *The assignment  $(a \in \mathfrak{g}, f(x) \in \mathbb{C}((x)))$*

$$\overline{a \otimes f(x)} \mapsto a \otimes f(x), \quad \overline{\mathbf{c} \otimes x^{-1}p(x)} \mapsto \mathbf{c}$$

*determines a Lie algebra isomorphism from  $\widehat{\mathcal{C}}_{\mathfrak{g}, \phi}$  to  $\widehat{\mathfrak{g}}$ .*

*Proof.* It is obvious that this assignment gives rise to a linear isomorphism, say  $\varphi$ , from  $\widehat{\mathcal{C}}_{\mathfrak{g}, \phi}$  to  $\widehat{\mathfrak{g}}$ . For any  $a, b \in \mathfrak{g}$ ,  $f(x), g(x) \in \mathbb{C}((x))$ , we have

$$\begin{aligned} & \varphi(\overline{[a \otimes f(x), b \otimes g(x)]}) \\ &= \varphi(\overline{[a, b] \otimes f(x)g(x) + (a, b) \mathbf{c} \otimes p(x) \left( \frac{d}{dx} f(x) \right) g(x)}) \\ &= [a, b] \otimes f(x)g(x) + (a, b) \operatorname{Res}_x \left( \left( \frac{d}{dx} f(x) \right) g(x) \right) \mathbf{c} \\ &= [\varphi(\overline{a \otimes f(x)}), \varphi(\overline{b \otimes g(x)})], \end{aligned}$$

noticing that

$$\overline{\mathbf{c} \otimes p(x) \left( \frac{d}{dx} f(x) \right) g(x)} = \overline{\mathbf{c} \otimes x^{-1}p(x)} \operatorname{Res}_x \left( \left( \frac{d}{dx} f(x) \right) g(x) \right).$$

Thus  $\varphi$  is a Lie algebra isomorphism. □

For any  $\ell \in \mathbb{C}$ , we denote by

$$V_{\widehat{\mathfrak{g}}}(\ell, 0) = \mathcal{U}(\widehat{\mathfrak{g}}) \otimes_{\mathcal{U}(\mathfrak{g} \otimes \mathbb{C}[[x]] \oplus \mathbb{C}\mathbf{c})} \mathbb{C}$$

the affine vertex algebra of  $\mathfrak{g}$  with the central charge  $\ell$ , where  $\mathfrak{g} \otimes \mathbb{C}[[x]]$  acts trivially on  $\mathbb{C}$  and  $\mathbf{c}$  acts as the scalar  $\ell$  on  $\mathbb{C}$ . Note that  $V_{\widehat{\mathfrak{g}}}(\ell, 0)$  is naturally isomorphic to the quotient vertex algebra of  $V_{\mathcal{C}_{\mathfrak{g}}}$  modulo the ideal generated by  $\mathbf{c} - \ell$ . We say that a  $\widehat{\mathfrak{g}}$ -module  $W$  is restricted if for any  $a \in \mathfrak{g}, w \in W, (a \otimes x^n).w = 0$  for  $n \gg 0$ , and is of level  $\ell$  if  $\mathbf{c}$  acts as the scalar  $\ell$  on  $W$ . From Lemma 6.1 and Theorem 4.5, we have the following characterization of  $\phi$ -coordinated  $V_{\widehat{\mathfrak{g}}}(\ell, 0)$ -modules.

**Proposition 6.2.**  *$\phi$ -coordinated modules for the affine vertex algebra  $V_{\widehat{\mathfrak{g}}}(\ell, 0)$  are exactly restricted  $\widehat{\mathfrak{g}}$ -modules of level  $\ell$ .*

More generally, assume that  $G$  is a group acting on  $\mathfrak{g}$  as an automorphism group preserving the bilinear form  $(\cdot, \cdot)$  such that for  $a, b \in \mathfrak{g}$ ,

$$(6.1) \quad [ga, b] = 0 \quad \text{and} \quad (ga, b) = 0 \quad \text{for all but finitely many } g \in G.$$

Let  $\chi, \chi_\phi : G \rightarrow \mathbb{C}^\times$  be any linear characters satisfy the relation (5.3). We lift the  $G$ -action from  $\mathfrak{g}$  to  $\widehat{\mathfrak{g}}$  by

$$(6.2) \quad g(a \otimes f(x) + \beta \mathbf{c}) = (ga \otimes f(\chi_\phi(g)^{-1}x)) + \beta \mathbf{c}$$

for  $g \in G, a \in \mathfrak{g}, f(x) \in \mathbb{C}((x)), \beta \in \mathbb{C}$ . From [Li3], the  $(G, \chi_\phi)$ -covariant algebra of  $\widehat{\mathfrak{g}}$  is the Lie algebra  $\widehat{\mathfrak{g}}[G]$ , where

$$(6.3) \quad \widehat{\mathfrak{g}}[G] = \widehat{\mathfrak{g}} / \text{span}\{gu - u \mid g \in G, u \in \widehat{\mathfrak{g}}\}$$

as a vector space and the Lie bracket is given by

$$[\bar{u}, \bar{v}] = \sum_{g \in G} \overline{[gu, v]} \quad \text{for } u, v \in \widehat{\mathfrak{g}},$$

where for  $u \in \widehat{\mathfrak{g}}, \bar{u}$  stands for the image of  $u$  in  $\widehat{\mathfrak{g}}[G]$  under the natural quotient map  $\widehat{\mathfrak{g}} \rightarrow \widehat{\mathfrak{g}}[G]$ . We define a  $G$ -action on  $\mathcal{C}_{\mathfrak{g}}$  by

$$R_g(\partial^n \otimes a) = \chi(g)^{n+1} \partial^n \otimes ga \quad \text{and} \quad R_g(\mathbf{c}) = \mathbf{c} \quad \text{for } g \in G, a \in \mathfrak{g}, n \in \mathbb{N}.$$

Then  $(\mathcal{C}_{\mathfrak{g}}, R)$  is a  $(G, \chi)$ -conformal algebra and so  $(V_{\mathcal{C}_{\mathfrak{g}}}, R)$  is a  $(G, \chi)$ -vertex algebra. A similar proof as that in Lemma 6.1, we have

**Lemma 6.3.** *The assignment  $(a \in \mathfrak{g}, f(x) \in \mathbb{C}((x)))$*

$$\overline{a \otimes f(x)} \mapsto \overline{a \otimes f(x)}, \quad \overline{\mathbf{c} \otimes x^{-1}p(x)} \mapsto \bar{\mathbf{c}}$$

*determines a Lie algebra isomorphism from  $\widehat{\mathcal{C}}_{\mathfrak{g}, \phi}[G]$  to  $\widehat{\mathfrak{g}}[G]$ .*

As  $R_g(\mathbf{c}) = \mathbf{c}$  for  $g \in G$ ,  $V_{\widehat{\mathfrak{g}}}(\ell, 0)$  becomes a  $(G, \chi)$ -vertex algebra with the  $G$ -action  $R$  transferred from that of  $V_{\mathcal{C}_{\mathfrak{g}}}$ . As before, we can define the notion “restricted  $\widehat{\mathfrak{g}}[G]$ -module of level  $\ell$ ”. Then from Lemma 6.3 and Theorem 5.12, we immediately have the following generalization of Proposition 6.2.

**Proposition 6.4.** *Let  $\chi, \chi_\phi$  be linear characters of  $G$  with  $\chi_\phi$  is injective and the relation (5.3) hold. Then  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for the affine  $(G, \chi)$ -vertex algebra  $V_{\widehat{\mathfrak{g}}}(\ell, 0)$  are exactly restricted  $\widehat{\mathfrak{g}}[G]$ -modules of level  $\ell$ .*

**Remark 6.5.** *We illustrate two examples for the Proposition 6.4(cf.[Li3]). First, let  $\sigma$  be an automorphism of  $\mathfrak{g}$  which has order  $M$  and preserves the form  $(\cdot, \cdot)$ , and let  $\omega$  be a primitive  $M$ -th root of unity. Set  $G = \langle \sigma \rangle$  and take  $\chi_\phi$  to be the character of  $G$  determined by  $\chi_\phi(\sigma) = \omega$ . Then  $\widehat{\mathfrak{g}}[G]$  is isomorphic to the  $\sigma$ -twisted affine Lie algebra*

$$\widehat{\mathfrak{g}}[\sigma] = \{u \in \widehat{\mathfrak{g}} \mid \hat{\sigma}(u) = u\},$$

where  $\hat{\sigma}$  is the automorphism of  $\widehat{\mathfrak{g}}$  defined by  $\hat{\sigma}(\mathbf{c}) = \mathbf{c}$  and  $\hat{\sigma}(a \otimes f(x)) = \sigma(a) \otimes f(\omega^{-1}x)$ ,  $a \in \mathfrak{g}$ ,  $f(x) \in \mathbb{C}((x))$ .

Second, let  $\mathfrak{g} = \mathfrak{gl}_\infty$  be the Lie algebra of  $\mathbb{Z} \times \mathbb{Z}$ -matrices which have only finitely many nonzero entries and set  $(x, y) = \text{Tr}(xy)$ . For any positive integer  $N$ , define an automorphism  $\sigma_N$  on  $\mathfrak{gl}_\infty$  by  $\sigma_N(E_{i,j}) = E_{i+N,j+N}$  for  $i, j \in \mathbb{Z}$ , where  $E_{i,j}$  stands for the usual unit matrix. Set  $G = \langle \sigma_N \rangle$  and take  $\chi_\phi$  to be the character of  $G$  determined by  $\chi_\phi(\sigma_N) = q^{-1}$  with  $q$  is not a root of unity. As  $G \cong \mathbb{Z}$  is an infinite group, the relation (5.3) forces that  $p(x)$  must be a homogenous polynomial in  $\mathbb{C}[x, x^{-1}]$  ([JKLT]). Thus we may define  $\widehat{\mathfrak{g}}$  to be the usual affine Lie algebra  $\mathfrak{g} \otimes \mathbb{C}[x, x^{-1}] \oplus \mathbb{C}\mathbf{c}$ . Then  $\widehat{\mathfrak{g}}[G]$  is isomorphic to a one dimensional central extension of the  $N \times N$ -matrix algebra  $\mathfrak{gl}_N(\mathbb{C}_q)$  over a 2-dimensional quantum torus  $\mathbb{C}_q$ , where  $\mathbb{C}_q = \mathbb{C}[x^{\pm 1}, y^{\pm 1}]$  as vector space with the multiplication given by  $(x^m y^n)(x^s y^t) = q^{ns} x^{m+s} y^{n+t}$  (cf.[Li3, CLTW]).

Next, we consider the  $\phi$ -coordinated modules for the Virasoro vertex algebra. Let

$$\mathcal{V} = \mathbb{C}((x)) \frac{d}{dx} \oplus \mathbb{C}\mathbf{c}$$

be (formal) Virasoro Lie algebra, where  $\mathbf{c}$  is central and for  $f(x), g(x) \in \mathbb{C}((x))$ ,

$$\begin{aligned} [f(x) \frac{d}{dx}, g(x) \frac{d}{dx}] &= f(x) \left( \frac{d}{dx} g(x) \right) \frac{d}{dx} - g(x) \left( \frac{d}{dx} f(x) \right) \frac{d}{dx} \\ &\quad + \frac{\mathbf{c}}{2} \text{Res}_x \left( \left( \frac{d}{dx} \right)^{(3)} f(x) \right) g(x). \end{aligned}$$

On the other hand, we have the *Virasoro conformal algebra* ([K])

$$\mathcal{C}_\mathcal{V} = (\mathbb{C}[\partial] \otimes \mathbb{C}L) \oplus \mathbb{C}\mathbf{c},$$

where for  $n \in \mathbb{N}$ ,  $\partial(\partial^n \otimes L) = \partial^{n+1} \otimes L$ ,  $\partial\mathbf{c} = 0$  and

$$L_n L = \delta_{n,0} \partial L + \delta_{n,1} 2L + \delta_{n,3} \frac{\mathbf{c}}{2}, \quad L_n \mathbf{c} = 0 = \mathbf{c}_n \mathbf{c}.$$

We identify  $L$  as  $1 \otimes L \in \mathcal{C}_\mathcal{V}$ . It is straightforward to see that the Lie algebra  $\widehat{\mathcal{C}}_{\mathcal{V}\phi}$  is spanned by the following elements

$$\overline{\mathbf{c} \otimes x^{-1} p(x)}, \quad \overline{L \otimes f(x)}, \quad f(x) \in \mathbb{C}((x)).$$

Furthermore, we have

**Lemma 6.6.** *The assignment*

$$\overline{L \otimes f(x)} \mapsto p(x)f(x)\frac{d}{dx} + \alpha_\phi(f(x))\mathbf{c}, \quad \overline{\mathbf{c} \otimes x^{-1}p(x)} \mapsto \mathbf{c}$$

determines a Lie algebra isomorphism from  $\widehat{\mathcal{C}}_{\mathcal{V}_\phi}$  to  $\mathcal{V}$ , where  $\alpha_\phi : \mathbb{C}((x)) \rightarrow \mathbb{C}$  is a linear map defined by

$$f(x) \mapsto \frac{1}{24}\text{Res}_x f(x) \left( 2\left(\frac{d}{dx}\right)^2 p(x) - p(x)^{-1} \left(\frac{d}{dx} p(x)\right)^2 \right).$$

*Proof.* Obviously, this assignment gives rise to a linear isomorphism  $\varphi : \widehat{\mathcal{C}}_{\mathcal{V}_\phi} \rightarrow \mathcal{V}$ . Now we prove that  $\varphi$  is a Lie algebra homomorphism. For any  $f(x), g(x) \in \mathbb{C}((x))$ , we have

$$\begin{aligned} & \varphi\left(\overline{[L \otimes f(x), L \otimes g(x)]}\right) \\ &= \varphi\left(\overline{-L \otimes p(x)\frac{d}{dx}(f(x)g(x)) + 2L \otimes p(x)\left(\frac{d}{dx}f(x)\right)g(x)}\right. \\ & \quad \left. + \frac{1}{2}\overline{\mathbf{c} \otimes \left((p(x)\frac{d}{dx})^{(3)}f(x)\right)g(x)}\right) \\ &= \varphi\left(\overline{L \otimes p(x)g(x)\frac{d}{dx}f(x) - L \otimes p(x)f(x)\frac{d}{dx}g(x)}\right. \\ & \quad \left. + \frac{1}{12}\overline{\mathbf{c} \otimes x^{-1}p(x)\text{Res}_x\left(\left(\frac{d}{dx}p(x)\right)^2\frac{d}{dx}f(x) + 3p(x)\left(\frac{d}{dx}p(x)\right)\left(\frac{d}{dx}\right)^2f(x)\right)}\right. \\ & \quad \left. + p(x)\left(\left(\frac{d}{dx}\right)^2p(x)\right)\frac{d}{dx}f(x) + p(x)^2\left(\frac{d}{dx}\right)^3f(x)\right)g(x)\Big) \\ &= p(x)^2f(x)\left(\frac{d}{dx}g(x)\right)\frac{d}{dx} - p(x)^2g(x)\left(\frac{d}{dx}f(x)\right)\frac{d}{dx} \\ & \quad + \frac{1}{12}\mathbf{c}\text{Res}_x\left(f(x)\left(\frac{d}{dx}\right)^3p(x) + 3\left(\frac{d}{dx}p(x)\right)\left(\frac{d}{dx}\right)^2f(x)\right. \\ & \quad \left. + 3\left(\left(\frac{d}{dx}\right)^2p(x)\right)\frac{d}{dx}f(x) + p(x)\left(\frac{d}{dx}\right)^3f(x)\right)p(x)g(x) \\ &= p(x)^2f(x)\left(\frac{d}{dx}g(x)\right)\frac{d}{dx} - p(x)^2g(x)\left(\frac{d}{dx}f(x)\right)\frac{d}{dx} + \frac{\mathbf{c}}{2}\text{Res}_x\left(\frac{d}{dx}\right)^{(3)}(p(x)f(x))p(x)g(x) \\ &= [p(x)f(x)\frac{d}{dx}, p(x)g(x)\frac{d}{dx}] = [\varphi(\overline{L \otimes f(x)}), \varphi(\overline{L \otimes g(x)})], \end{aligned}$$

noticing that

$$\overline{\mathbf{c} \otimes p(x)h(x)} = \overline{\mathbf{c} \otimes x^{-1}p(x)\text{Res}_x h(x)} \quad \text{for any } h(x) \in \mathbb{C}((x)).$$

Thus  $\varphi$  is a Lie algebra isomorphism.  $\square$

For any  $\ell \in \mathbb{C}$ , we denote by

$$V_{\mathcal{V}}(\ell, 0) = \mathcal{U}(\mathcal{V}) \otimes_{\mathcal{U}(\mathbb{C}[[x]] \frac{d}{dx} \oplus \mathbb{C}\mathbf{c})} \mathbb{C}$$

the Virasoro vertex algebra with central charge  $\ell$ , where  $\mathbb{C}[[x]] \frac{d}{dx}$  acts trivially on  $\mathbb{C}$  and  $\mathbf{c}$  acts as the scalar  $\ell$  on  $\mathbb{C}$ . Then  $V_{\mathcal{V}}(\ell, 0)$  is naturally isomorphic to the quotient vertex algebra of  $V_{\mathcal{C}_{\mathcal{V}}}$ , modulo the ideal generated by  $\mathbf{c} - \ell$ . We say that a  $\mathcal{V}$ -module  $W$  is restricted if for any  $w \in W$ ,  $(x^n \frac{d}{dx}).w = 0$  for  $n \gg 0$ , and is of level  $\ell$  if  $\mathbf{c}$  acts as the scalar  $\ell$  on  $W$ . From Lemma 6.6 and Theorem 4.5, we immediately have the following characterization of  $\phi$ -coordinated  $V_{\mathcal{V}}(\ell, 0)$ -modules.

**Proposition 6.7.**  *$\phi$ -coordinated modules for the Virasoro vertex algebra  $V_{\mathcal{V}}(\ell, 0)$  are exactly restricted  $\mathcal{V}$ -modules of level  $\ell$ .*

**Remark 6.8.** *When  $p(x) = x$ , Propositions 6.2 and 6.7 were proved in [CLT] by a different method.*

Finally we consider the  $\phi$ -coordinated modules of the vertex algebras associated to Novikov algebras. Let  $\mathcal{A}$  be a Novikov algebra, in the sense that  $\mathcal{A}$  is a non-associative algebra satisfying

$$(ab)c - a(bc) = (ba)c - b(ac), \quad (ab)c = (ac)b \quad \text{for } a, b, c \in \mathcal{A},$$

and let  $\langle \cdot, \cdot \rangle$  be a symmetric bilinear form of  $\mathcal{A}$  satisfying

$$(6.4) \quad \langle ab, c \rangle = \langle a, bc \rangle, \quad \langle ab, c \rangle = \langle ba, c \rangle \quad \text{for } a, b, c \in \mathcal{A}.$$

Similar to the algebra  $\widehat{\mathfrak{g}}$ , we also have an ‘‘untwisted affine Lie algebra’’

$$\widehat{\mathcal{A}} = \mathcal{A} \otimes \mathbb{C}((x)) \oplus \mathbb{C}\mathbf{c}$$

of  $\mathcal{A}$  ([BLP]), where  $\mathbf{c}$  is central and for  $a, b \in \mathcal{A}$ ,  $f(x), g(x) \in \mathbb{C}((x))$ ,

$$\begin{aligned} [a \otimes f(x), b \otimes g(x)] &= ab \otimes \left( \frac{d}{dx} f(x) \right) g(x) - ba \otimes \left( \frac{d}{dx} g(x) \right) f(x) \\ &\quad + \frac{\mathbf{c}}{2} \langle a, b \rangle \text{Res}_x \left( \left( \frac{d}{dx} \right)^{(3)} f(x) \right) g(x). \end{aligned}$$

On the other hand, following [P], we have the conformal algebra

$$\mathcal{C}_{\mathcal{A}} = \mathbb{C}[\partial] \otimes \mathcal{A} \oplus \mathbb{C}\mathbf{c},$$

where for  $a, b \in \mathcal{A}$ ,  $n \in \mathbb{N}$ ,  $\partial(\partial^n \otimes a) = \partial^{n+1} \otimes a$ ,  $\partial\mathbf{c} = 0$  and

$$a_n b = \delta_{n,0} \partial(ba) + \delta_{n,1} (ab + ba) + \delta_{n,3} \langle a, b \rangle \frac{\mathbf{c}}{2}, \quad a_n \mathbf{c} = 0 = \mathbf{c}_n \mathbf{c}.$$

Note that there is a linear isomorphism from  $\widehat{\mathcal{C}}_{\mathcal{A}\phi} \rightarrow \widehat{\mathcal{A}}$  such that  $\overline{a \otimes f(x)} \rightarrow a \otimes f(x)$  for  $a \in \mathcal{A}$ ,  $f(x) \in \mathbb{C}((x))$  and that  $\overline{\mathbf{c} \otimes x^{-1} p(x)} \mapsto \mathbf{c}$ . Via this isomorphism,

we obtain a new Lie algebra structure on  $\widehat{\mathcal{A}}$  such that

$$\begin{aligned} [a \otimes f(x), b \otimes g(x)] = & ab \otimes \left( p(x) \frac{d}{dx} f(x) \right) g(x) - ba \otimes \left( p(x) \frac{d}{dx} g(x) \right) f(x) \\ & + \frac{\mathbf{c}}{2} \langle a, b \rangle \text{Res}_x p(x)^{-1} \left( p(x) \frac{d}{dx} \right)^{(3)} f(x) g(x) \end{aligned}$$

for  $a, b \in \mathcal{A}$ ,  $f(x), g(x) \in \mathbb{C}((x))$  and that  $\mathbf{c}$  is central. We denote this Lie algebra by  $\widehat{\mathcal{A}}_\phi$ .

For any  $\ell \in \mathbb{C}$ , just as  $V_{\widehat{\mathfrak{g}}}(\ell, 0)$ , we have an affine type vertex algebra  $V_{\widehat{\mathcal{A}}}(\ell, 0)$  ([BLP]) which is isomorphic to the quotient vertex algebra  $V_{\mathcal{C}_{\mathcal{A}}}/\langle \mathbf{c} - \ell \rangle$ . We say that a  $\widehat{\mathcal{A}}_\phi$ -module  $W$  is restricted if for any  $a \in \mathcal{A}, w \in W$ ,  $(a \otimes x^n).w = 0$  for  $n \gg 0$ , and is of level  $\ell$  if  $\mathbf{c}$  acts as the scalar  $\ell$  on  $W$ . In view of Theorem 4.5, we have the following slight generalization of a result in [BLP]:

**Proposition 6.9.**  *$\phi$ -coordinated  $V_{\widehat{\mathcal{A}}}(\ell, 0)$ -modules are exactly restricted  $\widehat{\mathcal{A}}_\phi$ -modules of level  $\ell$ .*

**Remark 6.10.** *Equip  $\mathcal{A} = \mathbb{C}[y, y^{-1}]$  with a bilinear operator*

$$\circ : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}, \quad (y^i, y^j) \mapsto jy^{i+j}.$$

*Then  $(\mathcal{A}, \circ)$  is a Novikov algebra. Unlike the usual affine Lie algebra case (cf. Lemma 6.1), it follows from [DZ] that the Lie algebra  $\widehat{\mathcal{A}}$  is not isomorphic to  $\widehat{\mathcal{A}}_\phi$  with  $\phi = xe^z$ .*

## REFERENCES

- [BLP] C. Bai, H.-S. Li, Y. Pei,  $\phi_\epsilon$ -Coordinated modules for vertex algebras, *J. Algebra* **426** (2015), 211-242.
- [CLT] F. Chen, H.-S. Li, S. Tan, Toroidal extended affine Lie algebras and vertex algebras, arXiv:2102.10968.
- [CLTW] F. Chen, H.-S. Li, S. Tan, Q. Wang, Extended affine Lie algebras, vertex algebras, and reductive groups, arXiv:2004.02821.
- [DLM] C. Dong, H.-S. Li, G. Mason, Vertex Lie algebras, vertex Poisson algebras and vertex algebras, in: Recent Developments in Infinite-Dimensional Lie Algebras and Conformal Field Theory, Charlottesville, VA, 2000, in: Contemp. Math., Vol. **297**, Amer. Math. Soc., Providence, RI, 2002, pp. 69-96.
- [DZ] D. Dokovic, K. Zhao, Derivations, isomorphisms and second cohomology of generalized Block algebras, *Algebra Colloq.* **3** (1996), 245-272.
- [FB] E. Frenkel, D. Ben-Zvi, *Vertex Algebras and Algebraic Curves*, Mathematical Surveys and Monographs, Vol. **88**, Amer. Math. Soc., Providence, 2004.
- [FLM] I. B. Frenkel, J. Lepowsky, A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Applied Math., Vol. **134**, Academic Press, Boston, 1988.
- [FZ] I. B. Frenkel, Y.-C. Zhu, Vertex operator algebras associated to representations of affine and Virasoro algebras, *Duke Math. J.* **66** (1992), 123-168.
- [G-K-K] M. Golenishcheva-Kutuzova, V. Kac,  $\Gamma$ -conformal algebras, *J. Math. Phys.* **39** (1998), 2290-2305.

- [JKLT] N. Jing, F. Kong, H.-S. Li, S. Tan,  $(G, \chi_\phi)$ -equivariant  $\phi$ -coordinated quasi modules for nonlocal vertex algebras, *J. Algebra* **570** (2021), 24-74.
- [Li1] H.-S. Li, Local systems of vertex operators, vertex superalgebras and modules, *J. Pure Appl. Algebra* **109** (1996), 143-195.
- [Li2] H.-S. Li,  $\phi$ -Coordinated Quasi-Modules for Quantum Vertex Algebras, *Comm. Math. Phys.* **308** (2011), 703-741.
- [Li3] H.-S. Li, On certain generalizations of twisted affine Lie algebras and quasimodules for  $\Gamma$ -vertex algebras, *J. Pure Appl. Algebra* **209** (2007), 853-871.
- [Li4] H.-S. Li,  $G$ -equivariant  $\phi$ -coordinated quasi modules for quantum vertex algebras, *J. Math. Phys.* **54** (2013), 1-26.
- [Li5] H.-S. Li, A new construction of vertex algebras and quasi-modules for vertex algebras, *Adv. Math.* **202** (2006), 232-286.
- [Li6] H.-S. Li, Local systems of twisted vertex operators, vertex superalgebras and twisted modules, *Contemporary Math.*, Vol. **193**, Amer. Math. Soc., Providence, 1996, 203-236.
- [K] V. G. Kac, *Vertex Algebras for Beginners*, Second edition, University Lecture Series, Vol. **10**, Amer. Math. Soc., Providence, 1998.
- [P] M. Primc, Vertex algebras generated by Lie algebras, *J. Pure Appl. Algebra* **135** (1999), 253-293.

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN, CHINA 361005  
Email address: chenfxmu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN, CHINA 361005  
Email address: xiaoling@stu.xmu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN, CHINA 361005  
Email address: tans@xmu.edu.cn

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN, CHINA 361005  
Email address: qingwang@xmu.edu.cn