

VERSALITY OF BRILL-NOETHER FLAGS AND DEGENERACY LOCI OF TWICE-MARKED CURVES

NATHAN PFLUEGER

ABSTRACT. A line bundle on a smooth curve C with two marked points determines a rank function $r(a, b) = h^0(C, L(-ap - bq))$. This paper studies Brill-Noether degeneracy loci; such a locus is defined to be the closure in $\text{Pic}^d(C)$ of the locus of line bundles with a specified rank function. These loci generalize the classical Brill-Noether loci $W_d^r(C)$ as well as Brill-Noether loci with imposed ramification. For general (C, p, q) we determine the dimension, singular locus, and intersection class of Brill-Noether degeneracy loci, generalizing classical results about $W_d^r(C)$. The intersection class has a combinatorial interpretation in terms of the number of reduced words for a permutation associated to the rank function, or alternatively the number of saturated chains in the Bruhat order. The essential tool is a versality theorem for a certain pair of flags on $\text{Pic}^d(C)$, conjectured by Melody Chan and the author. We prove this versality theorem by showing the injectivity of a generalized Petri map, along the lines of Eisenbud and Harris’s proof of the Gieseker-Petri theorem.

1. INTRODUCTION

Classical Brill-Noether theory concerns the varieties $W_d^r(C)$, where C is a smooth curve of genus g , parameterizing line bundles L with at least $r + 1$ linearly independent global sections. One way to understand such loci is to mark two points p, q on C , choose a large integer N , and view $H^0(C, L)$ as the intersection of two subspaces $P, Q \subseteq H = H^0(C, L(Np + Nq))$, where P, Q consist of sections of $L(Np + Nq)$ vanishing to order at least N at p and q , respectively. Provided that $d + N \geq 2g - 1$, the dimensions of P, Q, H do not depend on L , and $W_d^r(C)$ may be described as the locus where P, Q are sufficiently non-transverse. Globalizing this construction, we obtain a vector bundle \mathcal{H} on $\text{Pic}^{d+2N}(C)$ with two subbundles \mathcal{P}, \mathcal{Q} , and $W_d^r(C)$ is isomorphic (via a twist by $Np + Nq$) to the degeneracy locus where the fiber of $\mathcal{P} \cap \mathcal{Q}$ jumps dimension by $g - d + r$.

The Brill-Noether and Gieseker-Petri theorems amount to the claim that, for a general curve C , $W_d^r(C)$ is nonempty if and only if the expected codimension $(r + 1)(g - d + r)$ is at most g , has the expected codimension if so, and has singular locus equal to $W_d^{r+1}(C)$. I hope to convince the reader that it is fruitful to view these facts through the following lens: as the line bundle L varies in $\text{Pic}^d(C)$, the *relative position* of the subbundles P, Q deforms versally. This amounts to saying that, up to pullback along smooth morphisms (equivalently, étale-locally and up to products with affine space), $\text{Pic}^d(C)$ with this choice of two subbundles is the same as a product of Grassmannians, where dimension and singularity of the relevant degeneracy locus can be found by elementary means.

Melody Chan and the author conjectured in [CP19] that the same phenomenon occurs for flags recording vanishing order at the points p, q .

1.1. The versality theorem. Fix a smooth twice-marked curve (C, p, q) over an algebraically closed field k , and an integer $d \geq 2g - 1$ (for notation conventions about what we mean by “smooth curve” and other matters, consult Section 1.6). Pushing down a Poincaré line bundle on $C \times \text{Pic}^d(C)$ gives a rank $n = d + 1 - g$ vector bundle \mathcal{H}_d on $\text{Pic}^d(C)$, with fibers $(\mathcal{H}_d)_{[L]} \cong H^0(C, L)$. For any

divisor D on C , denote by $\text{tw}_D : \text{Pic}^d(C) \rightarrow \text{Pic}^{d+\deg D}(C)$ the “twist” isomorphism given pointwise by $[L] \mapsto [L(D)]$, and define for all $0 \leq a, b \leq d - 2g + 1$,

$$\begin{aligned}\mathcal{P}_d^a &= \text{tw}_{ap} \mathcal{H}_{d-a} \\ \mathcal{Q}_d^b &= \text{tw}_{bq} \mathcal{H}_{d-b}.\end{aligned}$$

Regarding these as subbundles of \mathcal{H}_d , we obtain two flags $\mathcal{P}_d^\bullet, \mathcal{Q}_d^\bullet$ in \mathcal{H}_d , both of coranks $[0, d - 2g + 1] \cap \mathbb{Z}$. We call these the *degree- d Brill-Noether flags* of (C, p, q) .

We discuss in Section 4 the notion of a *versal pair of flags*; for now we simply remark that roughly speaking we call a pair of flags *versal* if, up to pullback along smooth morphisms, it is isomorphic to a pair of tautological flags on a product of two flag varieties, and therefore the relative positions of the two flags in any fiber deform “as generally as possible” around that fiber. We prove [CP19, Conjecture 6.3], namely

Theorem 1.1. *Fix integers d, g , and an algebraically closed field k of any characteristic. If (C, p, q) is a general twice-marked curve of genus g over k , then the pair of degree- d Brill-Noether flags $\mathcal{P}_d^\bullet, \mathcal{Q}_d^\bullet$ of (C, p, q) is versal.*

Our proof of Theorem 1.1 is a degeneration argument, which can be summarized in the slogan: “twice-marked curves with versal Brill-Noether flags can be chained together.” We will show that if two twice-marked curves (C_1, p_1, q_1) and (C_2, p_2, q_2) both have versal Brill-Noether flags in *all* degrees, then if q_1 is glued to p_2 to form a twice-marked stable curve (X, p_1, q_2) , a general member of any smoothing of (X, p_1, q_2) has versal Brill-Noether flags in degree d (so a *very general* member has versal Brill-Noether flags in all degrees). More generally, the same principle holds for a chain of twice-marked curves. The base case $g = 1$ is provided by twice-marked elliptic curves with marked points not differing by torsion. This strategy is similar in spirit to the proof in [Oss14] of the Brill-Noether theorem, where the Brill-Noether condition is strengthened to a property of twice-marked curves that is preserved by chaining two such curves together and smoothing.

Remark 1.2. One might hope to generalize Theorem 1.1 to three or more marked points, but this is impossible, even for $g = 0$. A tuple of three or more fixed flags in a vector space of dimension at least 3 is never versal [CP19, Example 3.4].

Remark 1.3. The use of chains of elliptic curves in Brill-Noether theory has a long history; such chains possess a curious ability to behave like “general” curves, provided that the attachment points on each elliptic curve do not differ by torsion. This remark gives a few examples of that history, but is in no way comprehensive. In contrast to the flag curves used e.g. in [EH83], elliptic chains are often useful without need for a characteristic 0 hypothesis. The earliest use of elliptic chains in Brill-Noether theory was by Welters [Wel85], and they have been used for many applications to Brill-Noether theory of vector bundles on curves, e.g. [TiB92, TiB08, TiB14, CLMTiB18], and to study other aspects of the geometry of special divisors on curves, e.g. [CTiB08, CLMPTiB18, CP21]. The tropical analog of an elliptic chain is a chain of loops; these have found many applications in tropical Brill-Noether theory, e.g. [CDPR12, JP14, Pfl17, CLRW20, Man20]. An interesting perspective on why such chains are so applicable to Brill-Noether theory, and what generalizations of them may have similar properties, is in [Oss16, §5]; see especially Questions 5.4 and 5.5.

1.2. The coupled Petri map. The basic tool used to prove the versality theorem is a generalization of the Petri map. First define, for any line bundle L on a smooth twice-marked curve (C, p, q) and any subset $S \subseteq \mathbb{Z} \times \mathbb{Z}$,

$$(1) \quad T_{p,q}^L(S) = \sum_{(a,b) \in S} H^0(C, L(-ap - bq)) \otimes H^0(C, \omega_C \otimes L^\vee(ap + bq)).$$

We write $T_{p,q}^L$ as shorthand for $T_{p,q}^L(\mathbb{Z} \times \mathbb{Z})$. To define this sum, regard all terms as subspaces of $H^0(C^*, L) \otimes H^0(C^*, \omega_C \otimes L^\vee)$, where $C^* = C \setminus \{p, q\}$. Call the elements of $T_{p,q}^L$ *coupled tensors*.

Definition 1.4. The *(fully) coupled Petri map* of L on (C, p, q) is the map

$$\mu_{p,q}^L : T_{p,q}^L \rightarrow H^0(C, \omega_C)$$

given in each summand by the cup product. The restriction $\mu_{p,q}^L|_{T_{p,q}^L(S)}$ will be abbreviated $\mu_{p,q}^L|_S$ and called the *S-coupled Petri map*.

If $\mu_{p,q}^L$ is injective for every line bundle $L \in \text{Pic}(C)$, then we say that (C, p, q) satisfies the *(fully) coupled Petri condition*; if $\mu_{p,q}^L|_S$ is injective for all $L \in \text{Pic}(C)$ we say that (C, p, q) satisfies the *S-coupled Petri condition*.

The $\{(0,0)\}$ -coupled Petri map is the usual Petri map; the fully coupled Petri map may be viewed as gluing together information about the local geometry of Brill-Noether loci around not only L but all of its twists by multiples of p and q . The transpose of a specific type of coupled Petri map, in which only twists by a single point p are used, is studied in [CHTiB92, Lemma 3.1], where it is used to analyze the geometry of Brill-Noether varieties of once-marked curves.

Example 1.5. (Genus 0 curves) If (C, p, q) is a twice-marked smooth curve of genus 0, then every line bundle on C is nonspecial, and $T_{p,q}^L$ is trivial, so (C, p, q) satisfies the coupled Petri condition.

Example 1.6. (Genus 1 curves) A genus 1 twice-marked smooth curve (C, p, q) satisfies the coupled Petri condition if and only if $p - q$ is a non-torsion element of the Jacobian.

If (C, p, q) is a twice-marked smooth curve of genus 1, then $\omega_C \cong \mathcal{O}_C$ and the only line bundles L for which $H^0(C, L) \otimes H^0(C, \omega_C \otimes L^\vee) \neq \{0\}$ are those isomorphic to \mathcal{O}_C . If $p - q$ is non-torsion, then there is at most one $(a, b) \in \mathbb{Z}^2$ such that $L \cong \mathcal{O}_C(ap + bq)$, and it follows that $T_{p,q}^L$ is either trivial or one-dimensional. Since the image $\mu_{p,q}^L(\sigma \otimes \tau)$ of a simple tensor with $\sigma, \tau \neq 0$ is nonzero, $\mu_{p,q}^L$ is injective for all L in this case. So twice-marked elliptic curves with $p - q$ nontorsion satisfy the coupled Petri condition. On the other hand, if $p - q$ is torsion, then for any line bundle $L \cong \mathcal{O}_C(ap + bq)$, $T_{p,q}^L$ is infinite-dimensional and $\mu_{p,q}^L$ fails spectacularly to be injective.

We will prove (Corollary 4.10) that $\mathcal{P}_d^*, \mathcal{Q}_d^*$ is a versal pair at $[L] \in \text{Pic}^d(C)$ if and only if the $[0, d - 2g + 1]^2$ -coupled Petri map of L is injective. Indeed, the transpose of that map is equal (up to isomorphism of the domain and codomain) to the map from the tangent space $T_{[L]} \text{Pic}^d(C)$ to the space of first-order deformations of pairs of flags. This identification is the principal goal of Section 4. Our main result about the coupled Petri map is the following; it is proved in Section 5.

Theorem 1.7. *Let $S \subseteq \mathbb{Z} \times \mathbb{Z}$ be finite, and fix an algebraically closed field k . A general twice-marked genus g curve (C, p, q) over k satisfies the S-coupled Petri condition.*

Remark 1.8. By taking a countable intersection in $\mathcal{M}_{g,2}$, Theorem 1.7 implies that a *very general* twice-marked curve satisfies the fully coupled Petri condition. In fact, our strategy is to prove the existence of such a curve over some field in every characteristic. It is not possible to replace “very general” with “general” in this statement, however, as Example 1.6 shows for $g = 1$. Over \mathbb{C} the locus of genus-1 twice-marked curves (C, p, q) with $p - q$ torsion is analytically dense, and over the algebraic closure of a finite field this locus is all of $\mathcal{M}_{g,2}$.

Remark 1.9. It is not clear at first that one should even expect $T_{p,q}^L$ to be finite-dimensional in general, and Example 1.6 shows that it need not be for special twice-marked curves. There is a combinatorially interesting way to describe its dimension; we sketch it here without proof since it is not needed elsewhere in this paper. There is a unique permutation π of \mathbb{Z} such that for all $a, b \in \mathbb{Z}$, $h^0(C, L(-ap - bq)) = r^\pi(a, b) = \#\{a' \geq a : \pi(a') \geq b\}$. The “typical” case is that none of the line

bundles $L(-ap - bq)$ are special; in that case $\pi = \omega_{d-g}$, the descending permutation $n \mapsto d - g - n$. The dimension of $T_{p,q}^L$ is equal to the length $\ell(\omega_{d-g}\pi)$, which may be finite or infinite. When $g = 1$, the inversions of $\omega_{d-g}\pi$ are in bijection with pairs a, b such that $L \cong \mathcal{O}_C(ap + bq)$. Of course, Theorem 1.7 implies that for (C, p, q) very general, $\ell(\omega_{d-g}\pi) \leq g$.

1.3. Background on Brill-Noether theory of marked curves. Eisenbud and Harris [EH86] proved an extended form of the classical Brill-Noether theorem to curves with marked points in their development of the theory of limit linear series. This extended Brill-Noether theorem considers a general n -marked smooth curve (C, p_1, \dots, p_n) . Fix integers d, r and sequences $a_\bullet^1, \dots, a_\bullet^n$, called vanishing sequences, where each is a strictly increasing sequence $a_\bullet^i = (a_0^i, \dots, a_r^i)$ of nonnegative integers. Define the variety $G_d^r(C, (p_1, a_\bullet^1), \dots, (p_n, a_\bullet^n))$, parameterizing linear series (L, V) on C with $\deg L = d, \dim V = r + 1$ such that for all i, j , $\dim V(-a_j^i p_i) \geq r + 1 - j$. These varieties are analogs of the Brill-Noether varieties $G_d^r(C)$ for curves with n marked points, which allow for imposed ramification at the marked points. Such varieties are crucial in the theory of limit linear series, because when studying the limit of a linear series as a smooth curve degenerates to a nodal curve, one must control ramification conditions at the nodes. Then the extended Brill-Noether theorem [EH86, Theorem 4.5] states that, in characteristic 0, for (C, p_1, \dots, p_n) general in $\mathcal{M}_{g,n}$,

$$\dim G_d^r(C, (p_1, a_\bullet^1), \dots, (p_n, a_\bullet^n)) = g - (r + 1)(g - d + r) - \sum_{i=1}^n \sum_{j=0}^r (a_j^i - j),$$

when the variety is nonempty. The characteristic 0 hypothesis is used in the base case $g = 0$, and indeed this extended Brill-Noether theorem is false in positive characteristic when $n > 2$. However, the theorem is true in all characteristics when $n \leq 2$; a simple proof appears in [Oss14].

Eisenbud and Harris's proof does not consider the singularities of these varieties. In the case $n = 1$, this question is considered in [CHTiB92], where the singular locus is described using a map that is essentially a one-marked-point version of the coupled Petri map described above. In the case $n = 2$ and in all characteristics, an analog of the Gieseker-Petri theorem for $G_d^r(C, (p_1, a_\bullet^1), (p_2, a_\bullet^2))$ was proved in [COP19]. Although the varieties $G_d^r(C, (p_1, a_\bullet^1), (p_2, a_\bullet^2))$ are not smooth in general, their singular loci can be completely described in terms of singularities of Schubert varieties. The approach in [COP19] does not make use of any version of the Petri map, instead analyzing the geometry of the space of limit linear series on a nodal curve.

Brill-Noether varieties, for both marked and unmarked curves, have also been studied from the point of view of enumerative geometry. This study begins in Castelnuovo's calculation in 1889 of the number of points in $W_d^r(C)$ when it is 0-dimensional and C is general, under the hypothesis that the Brill-Noether theorem was true. Castelnuovo's approach used a degeneration argument; a nice summary of this argument and how it has been adapted to other situations is in [HM06, §5A]. Modern approaches have typically used techniques from the theory of degeneracy loci, beginning with the work of Kempf [Kem71] and Kleiman-Laksov [KL72, KL74b], who computed the intersection theoretic class $[W_d^r(C)]$ via the Porteous formula. An analogous computation for curves with one marked point may be deduced from the degeneracy locus formulas in [KL74a]; this computation may be found, for example, in [Pfl18, §4], and an analogous theorem for curves with a *moving* marked point can be found in [FT16]. The case of two marked points with imposed ramification was studied in detail by Anderson, Chen, and Tarasca [ACT21], who used degeneracy locus formulas to obtain not just the intersection class of the Brill-Noether loci but classes in K-theory as well. A recent paper [ACT20] also considers the case of one marked point from a motivic point of view.

The present paper focuses, as in [COP19], on the case $n = 2$, where results are possible in all characteristics. Some of the works mentioned above focus on parameter spaces (generalizing

$G_d^r(C)$), while others focus on loci in $\text{Pic}^d(C)$ (generalizing $W_d^r(C)$) or consider both points of view. We focus entirely in loci in $\text{Pic}^d(C)$ in this paper. Most significantly, the present work goes beyond imposing vanishing conditions at the individual marked points, and allows conditions to be imposed on combinations of the marked points. The enumerative aspects of this paper, developed in Section 6, follow similar methods to those of [ACT21]. The principle difference in this paper is that while [ACT21] considers degeneracy loci indexed by 321-avoiding permutations, imposing conditions on combinations of marked points requires us to consider degeneracy loci indexed by arbitrary permutations (of finite length). See also Remark 1.13.

For example, while vanishing conditions may impose a cusp, flex point, or higher ramification condition at a single marked point, the approach in this paper may also impose nodes, bitangents, and other conditions concerning the interaction of the two marked points.

1.4. Brill-Noether degeneracy loci. We will apply the versality theorem to prove a generalization of the basic theorems about $W_d^r(C)$ to degeneracy loci that we call *Brill-Noether degeneracy loci*. Given a twice-marked smooth curve (C, p, q) , integer d , and function $r : \mathbb{N}^2 \rightarrow \mathbb{N}$, we consider the locus

$$(2) \quad \{[L] \in \text{Pic}^d(C) : h^0(C, L(-ap - bq)) \geq r(a, b) \text{ for all } a, b \in \mathbb{N}^2\}.$$

For example, by choosing $r(a, b) = \max(0, r_0 + 1 - a - b)$ for some fixed number r_0 , this locus is the usual Brill-Noether locus $W_d^{r_0}(C)$; other choices of function $r(a, b)$ encode information about the geometry of the map $C \rightarrow \mathbb{P}H^0(C, L)^\vee$.

A warning about an unfortunately confusing aspect of our notation: traditionally in Brill-Noether theory, the number r is used for the *projective* dimension of a linear series, and called the “rank,” hence $W_d^r(C)$ is defined by the inequality $h^0(C, L) \geq r + 1$. But in (2) above, we use $r(a, b)$ to bound the vector space dimension, to be consistent with usage when speaking about degeneracy loci, and also use the word “rank” for $r(a, b)$. There does not seem to be a good way to avoid this.

Only certain functions $r(a, b)$ are useful the construction (2). In fact, the only functions where equality $r(a, b) = h^0(C, L(-ap - bq))$ is possible are *rank functions of dot arrays*, defined as follows.

Definition 1.10. A *dot array* is a finite subset of $\mathbb{N} \times \mathbb{N}$ such that no two distinct elements of Π are in the same row or the same column. When drawing a dot array, we index positions like entries in a matrix: (a, b) is placed in row a and column b , with $(0, 0)$ in the upper-left corner.

The *rank function* of a dot array is the function $\mathbb{N}^2 \rightarrow \mathbb{N}$ defined by

$$(3) \quad r^\Pi(a, b) = \#\{(a', b') \in \Pi : a' \geq a \text{ and } b' \geq b\}.$$

We call a function $r : \mathbb{N}^2 \rightarrow \mathbb{N}$ a *rank function* if it is the rank function of some dot array.

Given a dot array Π , the *row sequence* is the set of first coordinates of elements of Π sorted in increasing order and usually denoted (a_0, \dots, a_r) , where $r = |\Pi| - 1$, and the *column sequence* is the sorted set of second coordinates, usually denoted (b_0, \dots, b_r) .

See [FP98, §1.2] for a nice discussion and many examples. However, note that we use slightly different index conventions since we think of a, b as codimensions rather than dimensions. This better matches our intended applications to Brill-Noether theory, where a, b will be vanishing orders at a point. For example, we index beginning at 0 and count “towards the lower right” rather than “towards the upper left.”

Definition 1.11. Let (C, p, q) be a twice-marked curve of genus g , d a positive integer, and Π a dot array such that $|\Pi| \geq d + 1 - g$. Define set-theoretically

$$W_d^\Pi(C, p, q) = \{[L] \in \text{Pic}^d(C) : h^0(C, L(-ap - bq)) \geq r^\Pi(a, b) \text{ for all } a, b \in \mathbb{N}^2\}.$$




FIGURE 1. Examples of Brill-Noether degeneracy loci, with $|\Pi| = 3$. The shaded boxes indicate $\text{Ess}(\Pi)$.

Give this locus a scheme structure by interpreting each inequality with a Fitting ideal; see Section 6. We call $W_d^\Pi(C, p, q)$ a *Brill-Noether degeneracy locus*.

In fact, most of the inequalities in Definition 1.11 are redundant. Define

$$(4) \quad \text{Ess}(\Pi) = \{(a, b) \in \mathbb{N}^2 : r^\Pi(a-1, b) = r^\Pi(a, b-1) = r^\Pi(a, b) > r^\Pi(a+1, b) = r^\Pi(a, b+1)\},$$

and call this the *essential set* of Π . To interpret this definition when $a = 0$ or $b = 0$, define $r^\Pi(-1, b) = r^\Pi(0, b)$ and $r^\Pi(a, -1) = r^\Pi(a, 0)$ (this accords with Equation 3 by simply allowing $a, b \in \mathbb{Z}$). One may replace “for all $a, b \in \mathbb{N}^2$ ” with “for all $(a, b) \in \text{Ess}(\Pi)$ ” in Definition 1.11 without changing the scheme structure. See Section 3.2. It will also be convenient to take a slightly larger set than $\text{Ess}(\Pi)$ for some arguments; define the *essential rows* $\text{EssR}(\Pi)$ and *essential columns* $\text{EssC}(\Pi)$ to be the images of $\text{Ess}(\Pi)$ along the first and second projections to \mathbb{N} , respectively. With these definitions, define

$$\widetilde{W}_d^\Pi(C, p, q) = \{[L] \in \text{Pic}^d(C) : h^0(C, L(-ap - bq)) \geq r^\Pi(a, b) \text{ for all } (a, b) \in \text{EssR}(\Pi) \times \text{EssC}(\Pi)\}.$$

We illustrate the geometric significance of dot arrays and degeneracy loci in Figure 1. This figure shows various examples of dot arrays with 3 dots. The sketch shows a cartoon of a twice-marked curve with a line bundle $[L] \in \widetilde{W}_d^\Pi(C, p, q)$, immersed in \mathbb{P}^2 by the complete linear series of L (which we assume to be birational in these sketches). The boxes of the set $\text{Ess}(\Pi)$ are shaded. Note for example that for the “generic” dot array of size $r+1$, $\Pi = \{(0, r), (1, r-1), \dots, (r, 0)\}$, $W_d^\Pi(C, p, q) = W_d^r(C)$, $\text{Ess}(\Pi) = \{(0, 0)\}$, and $\widetilde{W}_d^\Pi(C, p, q) = W_d^r(C) \setminus W_d^{r+1}(C)$ (the choice of marked points is immaterial in this case).

The expected dimension of $W_d^\Pi(C, p, q)$ is the following number. Here a_0, \dots, a_r and b_0, \dots, b_r denote the row and column sequences, and $r = |\Pi| - 1$.

$$\begin{aligned} \rho_g(d, \Pi) &= g - (r+1)(g-d+r) - \sum_{i=0}^r (a_i - i) - \sum_{i=0}^r (b_i - i) \\ &\quad - \#\{(a, b), (a', b') \in \Pi : a < a' \text{ and } b < b'\}. \end{aligned}$$

This generalizes the classical Brill-Noether number $\rho_g(d, r) = g - (r+1)(g-d+r)$ that gives the expected dimension of $W_d^r(C)$; the remaining terms count conditions imposed by the required vanishing sequences at p and q and the interaction between them.

Dot patterns $\Pi \subset \mathbb{N}^2$ with $|\Pi| \geq d+1-g$ are in bijection with a type of permutation of \mathbb{Z} that we call (d, g) -*confined permutations* and describe in Section 6.1. The (d, g) -confined permutation π associated to Π has the feature that $\omega_{d-g}\pi$ has finite length, and in fact this length is the expected codimension: $\rho_g(d, \Pi) = g - \ell(\omega_{d-g}\pi)$.

Theorem 1.12. *Fix positive integers g, d and an algebraically closed field k . Let (C, p, q) be a general twice-marked smooth curve of genus g . For any dot array Π such that $|\Pi| \geq d + 1 - g$, $W_d^\Pi(C, p, q)$ is nonempty if and only if $\rho_g(d, \Pi) \geq 0$. If it is nonempty, it has pure dimension $\rho_g(d, \Pi)$, and the open subscheme $\widetilde{W}_d^\Pi(C, p, q)$ is smooth and dense.*

Let π be the (d, g) -confined permutation associated to Π , and let $R(\omega_{d-g}\pi)$ denote the set of reduced words for the permutation $\omega_{d-g}\pi$ (or equivalently, saturated chains from the identity to $\omega_{d-g}\pi$ in the Bruhat order). Then the Chow class of $W_d^\Pi(C, p, q)$ is

$$[W_d^\Pi(C, p, q)] = \frac{|R(\omega_{d-g}\pi)|}{\ell(\omega_{d-g}\pi)!} \Theta^{\ell(\omega_{d-g}\pi)}.$$

In fact, we prove a stronger version of the smoothness statement in Theorem 6.10, which requires a bit more terminology.

Since $\deg \Theta^g = g!$, Theorem 1.12 implies in part that, in the case $\rho_g(d, \Pi) = 0$, $W_d^\Pi(C, p, q)$ is a set of reduced points, where the number of points is the number of reduced words for a permutation.

Remark 1.13. When Π has no ascents, i.e. no pairs $(a, b), (a', b') \in \Pi$ with $a' \leq a$ and $b' \leq b$, the degeneracy locus $W_d^\Pi(C, p, q)$ is the subscheme of $W_d^{|\Pi|-1}(C)$ where a certain pair of vanishing conditions is imposed on the marked points individually but no conditions are imposed on combinations of the marked points. In this case, a stronger form of the Chow class formula above is proved Proposition 5.1 of [ACT21] (see also Theorem B, in the case $\beta = 0$); namely a formula is obtained for the class in K-theory. The condition that Π has no ascents is equivalent to the condition that $\omega_{d-g}\pi$ is 321-avoiding. Also, the smoothness theorem in [COP19] may be used to deduce the smoothness statement in Theorem 1.12, in the case where Π has no ascents.

Remark 1.14. The number of points in a classical Brill-Noether locus of dimension 0 is the number of standard Young tableaux on a rectangular partition; it was observed in [CDPR12, p. 3] that this statement can be seen by a bijection in an analogous tropical situation using a chain of loops. In [Pfl17, Theorem 1.4] this bijection was extended to non-rectangular tableaux by considering divisors with prescribed ramification at a point.

Partitions may be associated with vexillary permutations, and standard young tableaux are then in bijection with reduced words of these permutations. It seems plausible that, upon formulating a tropical analog of the degeneracy loci $W_d^\Pi(C, p, q)$, there should be a nice bijection between the set of reduced words of a given permutation and a 0-dimensional tropical degeneracy locus on a chain of loops.

Question 1.15. Under what circumstances is the degeneracy locus $W_d^\Pi(C, p, q)$ connected? The connectedness of $W_d^r(C)$ for general curves is usually established using the main theorem of [FL81], which does not apply in this situation. It is possible that a degeneration proof, along the lines of Osserman's proof of the connectedness of $W_d^r(C)$ [Oss19], may work.

We will use the versality theorem to prove Theorem 1.12, by demonstrating that the desired statements can all be deduced from the assumption that the degree- $(d + 4g - 2)$ Brill-Noether flags are versal. This analysis occupies Section 6.

1.5. Generalizations of $G_d^r(C)$. Although the details are not developed in this paper, the versality theorem should also be able to provide new proofs of the main results of [COP19] about the geometry of the varieties $G_d^r(C, (p, a.), (q, b.))$ parameterizing linear series on C with prescribed vanishing orders at two marked points p and q , as described in Section 1.3. This potential application is also discussed in [CP19, §6] and [Li20, §4]. By choosing sufficiently large integers M and N and letting

$d' = d + M + N$, this variety can be described set-theoretically as

$$(5) \quad \{(L, V) \in G_{d'}^r(C) : \dim V \cap (\mathcal{P}_{d'}^{M+a_i})_{[L(Mp+Nq)]} \geq r+1-i \\ \text{and } \dim V \cap (\mathcal{Q}_{d'}^{N+b_i})_{[L(Mp+Nq)]} \geq r+1-i \text{ for all } 0 \leq i \leq r\}.$$

This has the form of a *relative Richardson variety*, in the sense of [CP19]. However, the results of [CP19] on such varieties work with pairs of *complete flags*, so they do not immediately apply in this situation.

Assuming that the results of [CP19] generalize to partial flags, they would quickly imply the results of [COP19] about the smooth locus of a twice-pointed Brill-Noether variety, and also imply that the image of the projection

$$G_d^r(C, (p, a_\bullet), (q, b_\bullet)) \rightarrow \text{Pic}^d(C)$$

is equal to a Brill-Noether degeneracy locus of the type discussed in this paper. Furthermore, the techniques of [Li20] provide natural resolutions of singularities for twice-pointed Brill-Noether varieties, via *relative Bott-Samelson varieties*; see [Li20, Conjecture 4.4].

1.6. Conventions. By a *smooth curve* we mean a smooth proper geometrically connected curve over a field k of any characteristic (we will usually, but not always, take k to be algebraically closed). A *point* of a smooth curve will refer to a k -valued point unless otherwise stated. A *twice-marked smooth curve* (C, p, q) is a smooth curve with two distinct marked points. For a twice-marked curve (C, p, q) , we denote the punctured curve $C \setminus \{p, q\}$ by C^* , where the marked points p, q will be clear from context.

To simplify our notation, intervals $[a, b]$, $(-\infty, b]$, $[a, \infty)$ will always refer to sets of integers $[a, b] \cap \mathbb{Z}$, $(-\infty, b] \cap \mathbb{Z}$, $[a, \infty) \cap \mathbb{Z}$.

A (*partial*) *flag* in a vector bundle \mathcal{H} (or vector space) is a sequence of nested subbundles (or subspaces) $\mathcal{P}^{a_0} \supset \dots \supset \mathcal{P}^{a_\ell}$, where the subscript always denotes corank (i.e. the rank of $\mathcal{H}/\mathcal{P}^a$ is a). The set $A = \{a_0, \dots, a_\ell\}$ is called the set of *coranks* of the flag. A *complete flag* is a flag with set of coranks $[0, n]$, where n is the rank of \mathcal{H} .

If $\alpha : \mathbb{Z} \xrightarrow{\sim} \mathbb{Z}$ is a permutation fixing all but finitely many $n \in \mathbb{Z}$, then the *length* of α is $\ell(\alpha) = \#\{(a, b) \in \mathbb{Z} : a < b \text{ and } \alpha(a) > \alpha(b)\}$, i.e. the number of inversions in α . The length is also equal to the minimum length of a factorization of α into a product of simple transpositions; such a factorization is called a *reduced word* for α . If a permutation of \mathbb{Z} moves infinitely many points, we say that it has infinite length.

Denote by ω_m the descending permutation of \mathbb{Z} given by $\omega_m(n) = m - n$, which we will sometimes also regard as a permutation of $[0, m - 1]$ rather than all of \mathbb{Z} . We also use ω_C and $\omega_{C/B}$ to denote the canonical sheaf of a smooth curve and the relative dualizing sheaf of a family of curves, and hope this leads to no confusion.

The symbol \mathbb{N} will denote the set of *nonnegative* integers.

1.7. Outline of the paper. Section 2 carries out the degeneration argument needed to prove Theorem 1.7. Section 3 gives some preliminary material about relative positions of flags. Section 4 is concerned with first-order deformations of a pair of flags, and builds the link between versality of Brill-Noether flags and the coupled Petri map. Section 5 completes the proofs of Theorems 1.1 and 1.7. Section 6 applies the versality theorem to the analysis of Brill-Noether degeneracy loci and proves Theorem 1.12.

ACKNOWLEDGEMENTS

This work was supported by a Miner D. Crary Sabbatical Fellowship from Amherst College. The work and manuscript were completed during the special semester on Combinatorial Algebraic




FIGURE 2. The arithmetic surface $\mathcal{C} \rightarrow \text{Spec } R$ in Situation 2.1.

Geometry at ICERM. I am also grateful to Melody Chan, Montserrat Teixidor i Bigas, and David Anderson for helpful conversations and suggestions.

2. DEGENERATION TO CHAINS TWICE-MARKED CURVES

This section is largely independent of the rest of the paper; its object may be summarized by the slogan: “twice-marked curves satisfying the coupled Petri condition can be chained together.” The approach here is based on Eisenbud and Harris’s proof of the Gieseker-Petri theorem [EH83], with two main adaptations. First, we degenerate to a chain of elliptic curves, rather than a flag curve; this is similar to the approach taken in [Wel85], and shares some features with the tropical proof of the Gieseker-Petri theorem in [JP14]. In fact, we formulate our degeneration in a more flexible way. Any chain of smooth curves, all satisfying the coupled Petri condition when the attachment points are marked, will do; elliptic curves are simply a convenient base case. Second, we work throughout with sections of line bundles on twice-punctured curves (i.e. rational sections); this adds the flexibility to consider the fully coupled Petri condition all at once, and also simplifies the argument somewhat.

Our aim is to show that, if a chain of twice-marked curves satisfying the coupled Petri condition is smoothed in a family $\mathcal{C} \rightarrow B$, then for any finite set S a general member of \mathcal{C} satisfies the S -coupled Petri condition (and therefore a very general member satisfies the fully coupled Petri condition). As in [EH83], we demonstrate this by working over a discrete valuation ring, and studying the geometric general fiber.

An *arithmetic surface* is a proper flat scheme over a discrete valuation ring whose generic fiber is a smooth curve. Throughout this section, we work with an arithmetic surface as described in the following Situation, and illustrated in Figure 2.

Situation 2.1. Let R be a discrete valuation ring, with residue field k , uniformizer t , and fraction field K . Assume that k is algebraically closed. Let $\pi : \mathcal{C} \rightarrow \text{Spec } R$ be a regular arithmetic surface, with two disjoint sections P, Q . We sometimes also denote P by A_0 and Q by A_{l+1} . Assume that

the special fiber \mathcal{C}_k is a chain of ℓ smooth curves $A_1 \cup \cdots \cup A_\ell$, where A_n and A_{n+1} meet nodally at a point that we denote by both q_n and p_{n+1} ¹. Furthermore, assume that the section P meets the special fiber at a point $p_1 \in A_1$ distinct from q_1 , and Q meets the special fiber at a point $q_\ell \in A_\ell$ distinct from p_ℓ .

Denote by \mathcal{C}^* the open subscheme given by the complement of $P \cup A_1 \cup \cdots \cup A_\ell \cup Q = A_0 \cup \cdots \cup A_{\ell+1}$, and by \mathcal{U}_n the complement of $A_0 \cup \cdots \cup A_{n-1} \cup A_{n+1} \cup \cdots \cup A_{\ell+1}$. Also denote by A_n^* the punctured smooth curve $A_n \setminus \{p_n, q_n\}$ (for $1 \leq n \leq \ell$). The generic fiber $\mathcal{C} \times_R \text{Spec } K$ will be denoted \mathcal{C}_η , and the geometric generic fiber $\mathcal{C} \times_R \text{Spec } \overline{K}$ will be denoted by $\mathcal{C}_{\overline{\eta}}$.

Throughout this section, we will follow the notation convention that if \mathcal{L}, \mathcal{M} are line bundles on \mathcal{C} , then L_n, M_n denote the restrictions of these line bundles to A_n .

Our objective in this section is to prove the following.

Theorem 2.2. *In Situation 2.1, suppose that each twice-marked curve (A_n, p_n, q_n) satisfies the coupled Petri condition. Then $(\mathcal{C}_{\overline{\eta}}, P_{\overline{\eta}}, Q_{\overline{\eta}})$ satisfies the coupled Petri condition.*

It will simplify notation slightly and add no additional difficulty to work in a slightly more general situation. If \mathcal{L}, \mathcal{M} are line bundles on \mathcal{C} , denote by μ the cup product map

$$\mu : H^0(\mathcal{C}^*, \mathcal{L}) \otimes_R H^0(\mathcal{C}^*, \mathcal{M}) \rightarrow H^0(\mathcal{C}^*, \mathcal{L} \otimes \mathcal{M}).$$

We will define a notion of coupled tensors for this multiplication map and prove in Theorem 2.10 an analog of Theorem 2.2 in this setting, from which we will deduce Theorem 2.2.

2.1. Valuation of tensors. In Situation 2.1, the divisors $P = A_0, A_1, \dots, A_\ell, A_{\ell+1} = Q$ determine valuations $\nu_0, \dots, \nu_{\ell+1}$ on the function field of \mathcal{C} . If \mathcal{L} is a line bundle on \mathcal{C} , then ν_n also naturally determines a map $\nu_n : H^0(\mathcal{C}^*, \mathcal{L}) \rightarrow \mathbb{Z} \cup \{\infty\}$ giving the vanishing order of a rational section along A_n . Following the strategy of Eisenbud and Harris [EH83], we further extend ν_n to tensors, as follows. If \mathcal{L}, \mathcal{M} are two line bundles on \mathcal{C} , then we define a function

$$\nu_n : H^0(\mathcal{C}^*, \mathcal{L}) \otimes_R H^0(\mathcal{C}^*, \mathcal{M}) \rightarrow \mathbb{Z} \cup \{\infty\}$$

by setting $\nu_n(\rho)$ to the maximum, taken over all expansions $\rho = \sum \sigma_i \otimes \tau_i$ into rank-1 tensors, of $\min_i \{\nu_n(\sigma_i) + \nu_n(\tau_i)\}$. Importantly, $\nu_n(\rho)$ is *not* always equal to $\nu_n(\mu(\rho))$, although it does provide a lower bound. In the same fashion, for each $n \in [1, \ell]$ we extend the valuations ν_{p_n}, ν_{q_n} on the function field of A_n to maps $H^0(A_n^*, L_n) \otimes_k H^0(A_n^*, M_n) \rightarrow \mathbb{Z} \cup \{\infty\}$.

For $n \in [1, \ell]$, $\nu_n|_R$ is the discrete valuation on R , so any rank-1 tensor $\sigma \otimes \tau$ may be written $t^a \sigma' \otimes \tau'$, where $a \in \mathbb{Z}$ and $\nu_n(\sigma) = \nu_n(\tau) = 0$. It follows that for all $a \in \mathbb{Z}$,

$$\{\rho \in H^0(\mathcal{C}^*, \mathcal{L}) \otimes_R H^0(\mathcal{C}^*, \mathcal{M}) : \nu_n(\rho) \geq a\} = t^a H^0(\mathcal{U}_n, \mathcal{L}) \otimes_R H^0(\mathcal{U}_n, \mathcal{M}).$$

On the other hand, if $n \in \{0, \ell + 1\}$, then ν_n induces the trivial valuation on R . In general,

$$(6) \quad \nu_n(t^a \rho) = \begin{cases} a + \nu_n(\rho) & \text{if } 1 \leq n \leq \ell \\ \nu_n(\rho) & \text{if } n \in \{0, \ell + 1\}. \end{cases}$$

The exact sequence $0 \rightarrow \mathcal{L}(-A_n) \rightarrow \mathcal{L} \rightarrow L_n \rightarrow 0$ induces an exact sequence $0 \rightarrow H^0(\mathcal{U}_n, \mathcal{L}(-A_n)) \rightarrow H^0(\mathcal{U}_n, \mathcal{L}) \rightarrow H^0(A_n^*, L_n)$, and the image of the first map may be identified with $t \cdot H^0(\mathcal{U}_n, \mathcal{L})$, hence the second map induces an injection $H^0(\mathcal{U}_n, \mathcal{L}) \otimes_R k \hookrightarrow H^0(A_n^*, L_n)$. The same remarks apply to \mathcal{M} , and we obtain a restriction homomorphism

$$H^0(\mathcal{U}_n, \mathcal{L}) \otimes_R H^0(\mathcal{U}_n, \mathcal{M}) \rightarrow H^0(A_n^*, L_n) \otimes_k H^0(A_n^*, M_n),$$

¹Why give the same point two names? I have found it psychologically helpful to use q_n when treating it as a point of A_n , and p_{n+1} when treating it as a point of A_{n+1} , so that the central fiber consists of twice-marked curves (A_n, p_n, q_n) .

the kernel of which consists of all tensors divisible by t , i.e. all ρ with $\nu_n(\rho) \geq 1$. We denote the image of a tensor ρ under this homomorphism by $\rho|_{A_n^*}$; this is nonzero if $\nu_n(\rho) = 0$. The basic fact we need about valuations of tensors is the following.

Proposition 2.3. *Let $\rho \in H^0(\mathcal{C}^*, \mathcal{L}) \otimes_R H^0(\mathcal{C}^*, \mathcal{M})$, and let $n \in [1, \ell]$. If $\nu_n(\rho) = 0$, then the restriction $\rho|_{A_n^*}$ is nonzero, and satisfies*

$$\nu_{p_n}(\rho|_{A_n^*}) \geq \nu_{n-1}(\rho), \text{ and } \nu_{q_n}(\rho|_{A_n^*}) \geq \nu_{n+1}(\rho).$$

Proposition 2.3 is plausible enough, but it requires a bit of algebra to prove; we briefly develop the necessary tools in the next subsection.

2.2. Adapted bases. To prove Proposition 2.3, we make use of a special type of basis for finite-dimensional subspaces of sections of line bundles on \mathcal{C}^* . These bases play the role of the ‘‘compatible bases’’ used in [EH83, Lemma 1.2].

Definition 2.4. In Situation 2.1, let \mathcal{L} be a line bundle on \mathcal{C} , and suppose $n \in [0, \ell + 1]$. Let $S \subseteq H^0(\mathcal{C}, \mathcal{L})$, and let $V = \text{span } S$. The set S is called *adapted to A_n* if it is linearly independent (over K), and for all $a \in \mathbb{Z}$, the set $\{c\sigma : c \in K, \sigma \in S, \nu_n(c\sigma) \geq a\}$ generates the R -module $V_a = \{v \in V : \nu_n(v) \geq a\}$.

Adapted bases are convenient for evaluating valuations of sections: if $\{\sigma_i\}$ is adapted to A_n , then for any choice of constants $\{c_i\}$,

$$(7) \quad \nu_n\left(\sum c_i \sigma_i\right) = \min\{\nu_n(c_i \sigma_i)\}.$$

Lemma 2.5. *Let $V \subseteq H^0(\mathcal{C}^*, \mathcal{L})$ be a finite-dimensional subspace, and $n \in [0, \ell + 1]$. There exists a basis of V that is adapted to A_n .*

Proof. Observe that if $n \in \{0, \ell + 1\}$, then each V_a is a K -vector space, while if $n \in [1, \ell]$ then $V_a = t^a V_0$ for all $a \in \mathbb{Z}$, $V_0 \otimes_R K \cong V$, and V_0 is a free module since it is torsion-free. In these two cases, we may construct bases adapted to A_n as follows.

- (1) If $n \in \{0, \ell + 1\}$: choose $\sigma_1, \dots, \sigma_{\dim_K V}$ inductively so that for each $a \in \mathbb{Z}$, $\{\sigma_1, \dots, \sigma_{\dim_K V_a}\}$ is a basis for V_a .
- (2) If $n \in [1, \ell]$: choose an R -basis $\sigma_1, \dots, \sigma_{\dim_K V}$ for V_0 .

In either case, $\{\sigma_i\}$ is a K -basis for V adapted to A_n . \square

If adapted bases are constructed as in items (1) and (2) above, they lend themselves to modification by the operations of Gaussian elimination. Indeed, if we modify exactly one element of the basis by replacing σ_i by $\sigma_i + c\sigma_j$, then we obtain a new adapted basis of the same form provided that

- (1) if $n \in \{0, \ell + 1\}$, we choose $c \in K$ and $i > j$, and
- (2) if $n \in [1, \ell]$ we choose $c \in R$ and $i \neq j$.

Lemma 2.6. *Let $m, n \in [0, \ell + 1]$. There exists a basis of V that is adapted to both A_m and A_n .*

Proof. Let $\{\sigma_i\}$ be a basis adapted to A_m , and $\{\tau_i\}$ a basis adapted to A_n , constructed as in the proof of Lemma 2.5. Let T be the change of basis matrix from basis $\{\sigma_i\}$ to basis $\{\tau_i\}$. We may modify the bases, according to the operations described above, in such a way that T has exactly one nonzero entry in each row and in each column; to do so is an exercise in Gaussian elimination. This proves the lemma, since upon permuting one of the two bases we have that $\sigma_i = c_i \tau_i$ for nonzero constants $c_i \in K$, and it follows that $\{\sigma_i\}$ is now adapted to both A_m and A_n . \square

Lemma 2.7. *Suppose $\rho \in H^0(\mathcal{C}^*, \mathcal{L}) \otimes_R H^0(\mathcal{C}^*, \mathcal{M})$. If $\nu_m(\rho) \geq a$ and $\nu_n(\rho) \geq b$, then there exist $\sigma_1, \dots, \sigma_N \in H^0(\mathcal{C}^*, \mathcal{L})$ and $\tau_1, \dots, \tau_N \in H^0(\mathcal{C}^*, \mathcal{M})$ such that*

- (1) $\rho = \sum \sigma_i \otimes \tau_i$,
- (2) $\nu_m(\sigma_i) + \nu_m(\tau_i) \geq a$ for all i , and
- (3) $\nu_n(\sigma_i) + \nu_n(\tau_i) \geq b$ for all i .

Proof. By definition of $\nu_m(\rho), \nu_n(\rho)$, there exist two expansions $\rho = \sum_{k=1}^{N_1} \alpha_k \otimes \beta_k = \sum_{k=1}^{N_2} \gamma_k \otimes \delta_k$ such that $\nu_m(\alpha_i) + \nu_m(\beta_i) \geq a$ for all $i \in [1, N_1]$ and $\nu_n(\gamma_i) + \nu_n(\delta_i) \geq b$ for all $i \in [1, N_2]$. Let V be the span of all α_i and γ_i , and W be the span of all β_i and δ_i . By Lemma 2.6, we may choose bases $\{\sigma_i\}$ of V and $\{\tau_j\}$ of W adapted to both A_m and A_n , and $\{\sigma_i \otimes \tau_j\}$ is a basis for $V \otimes W$. Expanding an individual term $\alpha_k \otimes \beta_k$ in this basis results in a sum $\sum f_{i,j}^k \sigma_i \otimes \tau_j$, where, by Equation 7, $\nu_m(f_{i,j}^k) + \nu_m(\sigma_i) + \nu_m(\tau_j) \geq a$ for all i, j . Thus when ρ is expanded in this basis we obtain an expression $\rho = \sum f_{i,j} \sigma_i \otimes \tau_j$ where $\nu_m(f_{i,j}) + \nu_m(\sigma_i) + \nu_m(\tau_j) \geq a$ for all i, j . By linear independence, the same coefficients $f_{i,j}$ are obtained from expanding $\sum_{k=1}^{N_2} \gamma_k \otimes \delta_k$, so a similar argument shows that for all i, j , $\nu_n(f_{i,j}) + \nu_n(\sigma_i) + \nu_n(\tau_j) \geq b$, which gives the lemma. \square

We now have the tools to prove Proposition 2.3.

Proof of Proposition 2.3. Suppose that $1 \leq n \leq \ell$ and $\rho \in H^0(\mathcal{C}^*, \mathcal{L}) \otimes_R H^0(\mathcal{C}^*, \mathcal{M})$ satisfies $\nu_n(\rho) = 0$. The claim that $\rho|_{A_n^*}$ is nonzero was verified in the paragraph before the proposition statement.

By Lemma 2.7, there exists an expansion $\rho = \sum \sigma_i \otimes \tau_i$ such that $\nu_{n-1}(\sigma_i) + \nu_{n-1}(\tau_i) \geq \nu_{n-1}(\rho)$ and $\nu_n(\sigma_i) + \nu_n(\tau_i) \geq 0$ for all i . Since $\nu_n(t) = 1$, we may move a power of t from one factor to the other in each term so that $\nu_n(\sigma_i), \nu_n(\tau_i) \geq 0$ for all i . Thus we have well-defined restrictions $\sigma_i|_{A_n^*} \in H^0(A_n^*, L_n)$ and $\tau_i|_{A_n^*} \in H^0(A_n^*, M_n)$, and $\rho|_{A_n^*} = \sum (\sigma_i|_{A_n^*}) \otimes (\tau_i|_{A_n^*})$.

The restriction of $\mathcal{O}_{\mathcal{C}}(A_{n-1})$ to A_n is equal to $\mathcal{O}_{A_n}(p_n)$, and thus $\nu_{p_n}(\sigma_i|_{A_n^*}) \geq \nu_{n-1}(\sigma_i)$ and $\nu_{p_n}(\tau_i|_{A_n^*}) \geq \nu_{n-1}(\tau_i)$. Therefore $\nu_{p_n}(\rho|_{A_n^*}) \geq \min_i (\nu_{n-1}(\sigma_i) + \nu_{n-1}(\tau_i)) = \nu_{n-1}(\rho)$, as desired. The bound $\nu_{q_n}(\rho|_{A_n^*}) \geq \nu_{n+1}(\rho)$ follows by replacing ν_{n-1} with ν_{n+1} in this argument. \square

Although we have been working with an arithmetic surface $\mathcal{C} \rightarrow \text{Spec } R$ throughout this section, the reader may verify that all arguments regarding the valuations ν_0 and $\nu_{\ell+1}$ work with no modification if we instead consider a smooth curve $C \rightarrow \text{Spec } k$ over a field, with valuations ν_p, ν_q coming from two distinct rational points. The existence of an adapted basis in that setting is also given by Fact 3.2. Adapted to this context, we obtain the following analog of Lemma 2.7, which will be useful in the next section.

Lemma 2.8. *Let $C \rightarrow \text{Spec } k$ be a smooth curve over a field (not necessarily algebraically closed), with distinct rational points p, q . Let L, M be two line bundles on C , and $\rho \in H^0(\mathcal{C}^*, L) \otimes_k H^0(\mathcal{C}^*, M)$. If $A, B \in \mathbb{Z}$ satisfy $\nu_p(\rho) \geq A$ and $\nu_q(\rho) \geq B$, then*

$$\rho \in \sum_{a,b \in \mathbb{Z}} H^0(C, L((-a-A)p + (-b-B)q)) \otimes_k H^0(C, M(ap + bq)).$$

2.3. Injective coupling. We prove in this subsection our basic inductive result. Suppose that $C \rightarrow \text{Spec } k$ is a smooth curve over a field (not necessarily algebraically closed), p, q are distinct rational points, and L, M are two line bundles on C . Define

$$T_{p,q}^{L,M} = \sum_{a,b \in \mathbb{Z}} H^0(C, L(-ap - bq)) \otimes_k H^0(C, M(ap + bq)),$$

and define the *coupled multiplication map* to be the linear map

$$(8) \quad \mu_{p,q}^{L,M} : T_{p,q}^{L,M} \rightarrow H^0(C, L \otimes M),$$

given in each summand by the cup product. This generalizes the coupled Petri map.

Remark 2.9. The set $T_{p,q}^{L,M}$ also has the following two equivalent descriptions.

- (1) We may add the restrictions $-\deg M \leq a + b \leq \deg L$ to the sum defining $T_{p,q}^{L,M}$, since in all other summands one of the line bundles has negative degree.
- (2) By Lemma 2.8, $T_{p,q}^{L,M}$ is equal to the set of all $\rho \in H^0(C^*, L) \otimes_k H^0(C^*, M)$ such that both $\nu_p(\rho) \geq 0$ and $\nu_q(\rho) \geq 0$.

We say that two line bundles L, M on a smooth twice-marked curve (C, p, q) over a field have *injective coupling* if $\mu_{p,q}^{L,M}$ is injective.

Theorem 2.10. *In Situation 2.1, let \mathcal{L}, \mathcal{M} be line bundles on \mathcal{C} . Suppose that for all $1 \leq n \leq \ell$, $L_n(-p_n)$ and M_n have injective coupling on (A_n, p_n, q_n) . Then $L_\eta(-P_\eta), M_\eta$ have injective coupling on (C_η, P_η, Q_η) .*

Proof. Suppose that $\rho \in H^0(\mathcal{C}^*, \mathcal{L}) \otimes_R H^0(\mathcal{C}^*, \mathcal{M})$ is nonzero and satisfies $\nu_P(\rho) \geq 1$. We will demonstrate that $\nu_Q(\rho) < 0$.

For each n between 1 and ℓ inclusive, let $\rho_n = t^{-\nu_n(\rho)}\rho$. Proposition 2.3 implies that $\rho_n|_{A_n^*}$ is nonzero and satisfies $\nu_{p_n}(\rho_n|_{A_n^*}) \geq \nu_{n-1}(\rho_n)$ and $\nu_{q_n}(\rho_n|_{A_n^*}) \geq \nu_{n+1}(\rho_n)$. Injective coupling on A_n implies that either $\nu_{n-1}(\rho_n) < 1$ or $\nu_{n+1}(\rho_n) < 0$. By Equation 6,

$$\nu_{n-1}(\rho_n) = \begin{cases} \nu_P(\rho) & \text{if } n = 1 \\ \nu_{n-1}(\rho) - \nu_n(\rho) & \text{if } 2 \leq n \leq \ell \end{cases},$$

and

$$\nu_{n+1}(\rho_n) = \begin{cases} \nu_Q(\rho) & \text{if } n = \ell \\ \nu_{n+1}(\rho) - \nu_n(\rho) & \text{if } 1 \leq n \leq \ell - 1 \end{cases}.$$

A straightforward induction shows that if $\nu_P(\rho) \geq 1$, then $\nu_{n+1}(\rho) < \nu_n(\rho)$ for $1 \leq n \leq \ell - 1$, and $\nu_Q(\rho) < 0$. This establishes the theorem. \square

The following lemma is needed to apply Theorem 2.10 to the coupled Petri map.

Lemma 2.11. *Let (C, p, q) be a twice-marked smooth curve of any genus, and L any line bundle on C . The line bundles L and $\omega_C \otimes L^\vee$ have injective coupling if and only if $L(-p)$ and $\omega_C(p+q) \otimes L^\vee$ have injective coupling.*

Proof. By the Riemann-Roch formula, for all a, b either $H^0(C, L(-ap - (b-1)q)) = H^0(C, L(-ap - bq))$ or $H^0(C, \omega_C \otimes L^\vee(ap + bq)) = H^0(C, \omega_C \otimes L^\vee(ap + (b-1)q))$. This shows that $T_{p,q}^{L, \omega_C \otimes L^\vee} = T_{p,q}^{L(q), \omega_C \otimes L^\vee}$. Next, observe that $T_{p,q}^{L(q), \omega_C \otimes L^\vee} = T_{p,q}^{L(-p), \omega_C(p+q) \otimes L^\vee}$, simply by replacing a, b with $a+1, b+1$ in the summation. So despite appearances, the coupled multiplication maps for the pair $L, \omega_C \otimes L^\vee$ and the pair $L(-p), \omega_C(p+q) \otimes L^\vee$ are precisely the same map. \square

Proof of Theorem 2.2. In Situation 2.1, assume that each (A_n, p_n, q_n) satisfies the coupled Petri condition. Let L be any line bundle on \mathcal{C}_η ; we will prove that μ_{P_η, Q_η}^L is injective.

First consider the case where L is defined over K , so that L may be regarded as a line bundle on C_η . Since \mathcal{C} is regular, this line bundle may be extended (non-uniquely) to a line bundle \mathcal{L} on \mathcal{C} (e.g. choose a Cartier divisor D with $L \cong \mathcal{O}_{C_\eta}(D)$, and take its closure to obtain a Cartier divisor in \mathcal{C}). Let $\omega_{\mathcal{C}/R}$ be the relative dualizing sheaf. By the adjunction formula [Liu02, Theorem 9.1.37], together with the fact that $\mathcal{O}_{\mathcal{C}}(A_n)|_{A_n} \cong \mathcal{O}_{A_n}(-p_n - q_n)$, we have $\omega_{\mathcal{C}}(P+Q)|_{A_n} \cong \omega_{A_n}(p_n + q_n)$. Let $\mathcal{M} = \omega_{\mathcal{C}}(P+Q) \otimes \mathcal{L}^\vee$. Then $M_n = \omega_{A_n}(p_n + q_n) \otimes L_n^\vee$. Since (A_n, p_n, q_n) satisfies the coupled Petri condition, the bundles L_n and $\omega_{A_n} \otimes L_n^\vee$ have injective coupling; by Lemma 2.11 so do $L_n(-p_n)$ and M_n . Now Theorem 2.10 implies that $L(-P_\eta)$ and $M = \omega_{C_\eta}(P_\eta + Q_\eta) \otimes L^\vee$ have injective coupling on C_η , and Lemma 2.11 shows that L and $\omega_{C_\eta} \otimes L^\vee$ have injective coupling. Since the

global sections functor commutes with flat base extension, the same is true after L is pulled back from C_η to $C_{\overline{\eta}}$. This settles the case of line bundles on $C_{\overline{\eta}}$ defined over K .

For the general case, note that any line bundle L on $C_{\overline{\eta}}$ is defined over some finite extension $K' \supseteq K$. We may extend R to a discrete valuation ring R' with fraction field K' . Upon forming the base extension $\mathcal{C} \times_R \text{Spec } R'$ and resolving singularities, we obtain an arithmetic surface $\mathcal{C}' \rightarrow \text{Spec } R'$ with general fiber $C_\eta \times_K \text{Spec } K'$ and special fiber given by replacing each node p_n ($2 \leq n \leq \ell$) by a chain of rational curves (see e.g. [Liu02, Corollary 3.25]). By assumption on A_1, \dots, A_ℓ and Example 1.5, each component of the central fiber of \mathcal{C}' satisfies the coupled Petri condition. Therefore we may invoke the first case to conclude that L has injective coupling on $C_{\overline{\eta}}$. \square

3. FLAGS AND PERMUTATIONS

This section collects, without proofs, some basic facts about how permutations are used to record the relative position of a pair of flags in a fixed vector space and to define degeneracy loci of flags within vector bundles. There are a variety of conventions in the literature, depending on whether flags are indexed by dimension or codimension, whether indexing begins at 0 or 1, and other choices, so it is useful to consolidate the facts that we need in the notation used in this paper.

3.1. The permutation associated to a pair of flags.

Fact 3.1. *Let P^\bullet, Q^\bullet be two flags in a vector space H , with sets of coranks A and B respectively, and let $n = \dim H$. There exists a unique permutation σ of $[0, n-1]$ such that*

- (1) *For all $a \in A, b \in B$, $\dim P^a \cap Q^b = \#\{i \in [0, n-1] : i \geq a \text{ and } \sigma(i) \geq b\}$,*
- (2) *for all $a \in [0, n-1]$ with $a > 0$ and $a \notin A$, $\sigma(a) < \sigma(a-1)$, and*
- (3) *for all $b \in [0, n-1]$ with $b > 0$ and $b \notin B$, $\sigma^{-1}(b) < \sigma^{-1}(b-1)$.*

The permutation σ is called the *permutation associated to P^\bullet, Q^\bullet* . We call any permutation of $[0, n-1]$ satisfying conditions (2) and (3) above *compatible with coranks A, B* .

Fact 3.2. *Let P^\bullet, Q^\bullet be a pair of flags with associated permutation σ . There exists a basis $\{v_0, \dots, v_{n-1}\}$ of H such that for all $a \in A, b \in B$, $\{v_i : i \geq a\}$ is a basis for P^a and $\{v_i : \sigma(i) \geq b\}$ is a basis for Q^b . Therefore $\{v_i : i \geq a, \sigma(i) \geq b\}$ is a basis for $P^a \cap Q^b$.*

Although we will not prove these facts, we sketch a strategy of proof, for the reader who is new to these notions. First, construct a basis \mathcal{B} of H that contains a basis for all subspaces $P^a \cap Q^b$; this can be done inductively, ordering the intersections by dimension. Now sort this basis in two ways: one in order of stratum from P^\bullet , and one in order of stratum from Q^\bullet . Furthermore, ensure that any two elements $v, w \in \mathcal{B}$ are placed in the same order in both orderings whenever possible without violating the previous sentence. Comparing these two orderings gives σ .

We will call a basis as in Fact 3.2 *adapted to P^\bullet, Q^\bullet* .

3.2. Degeneracy loci and the essential set. The association of a permutation to a pair of flags can also be reversed, to define *degeneracy loci* for a pair of flags in a vector bundle. We follow the conventions of [CP19, §2] here. Given a permutation σ of $[0, n-1]$, define the *rank function* of σ to be the function $r^\sigma : \mathbb{N}^2 \rightarrow \mathbb{N}$ given by

$$(9) \quad r^\sigma(a, b) = \#\{a' \in [0, n-1] : a' \geq a \text{ and } \sigma(a') \geq b\}.$$

If S is a scheme with a rank n vector bundle \mathcal{H} and two complete flags $\mathcal{P}^\bullet, \mathcal{Q}^\bullet$ in \mathcal{H} , we define a degeneracy locus

$$(10) \quad D_\sigma(\mathcal{P}^\bullet; \mathcal{Q}^\bullet) = \{x \in S : \dim(\mathcal{P}^a)_x \cap (\mathcal{Q}^b)_x \geq r^\sigma(a, b) \text{ for all } a, b \in [0, n-1]\}.$$

Of course, the equation above is only a set-theoretic definition. To give a scheme-theoretic definition, translate each inequality into a bound on the rank of the bundle map $\mathcal{Q}^b \rightarrow \mathcal{H}/\mathcal{P}^a$, trivialize

this bundle map locally, and consider the scheme defined by all the minors of appropriate size in the matrix trivializing the bundle map, which is independent of the choice of trivialization. Alternatively, one can bound the rank of the difference map $\mathcal{P}^a \oplus \mathcal{Q}^b \rightarrow \mathcal{H}$, which results in the same ideal (this can be seen by representing this map in block form and comparing its minors to those of $\mathcal{Q}^b \rightarrow \mathcal{H}/\mathcal{P}^a$), and makes the symmetry evident. The scheme structure on D_σ is the intersection of these schemes.

Although there are a priori d^2 inequalities imposed in Equation 10, many of them are redundant. For example, if $r^\sigma(a, b) = r^\sigma(a + 1, b)$ or $r^\sigma(a, b) = r^\sigma(a - 1, b) - 1$, then the bound on $\dim(\mathcal{P}^a)_x \cap (\mathcal{Q}^b)_x$ is redundant. A minimal set of rank conditions is provided by the *essential set*, defined in [Ful92]. Translated into our notation, this set is

$$(11) \quad \text{Ess}(\sigma) = \{(a, b) : 1 \leq a, b < n, \sigma(a - 1) < b \leq \sigma(a) \text{ and } \sigma^{-1}(b - 1) < a \leq \sigma^{-1}(b)\}.$$

The inequalities in Equation 11 are equivalent to the equations

$$(12) \quad r^\sigma(a - 1, b) = r^\sigma(a, b - 1) = r^\sigma(a, b) > r^\sigma(a + 1, b) = r^\sigma(a, b + 1),$$

which make the redundancy more clear (cf. Equation 4). By [Ful92, Lemma 3.10], the scheme defined in Equation 10 is the same if we consider only $(a, b) \in \text{Ess}(\sigma)$, rather than all $(a, b) \in [0, d - 1]^2$ (this is not hard to see set-theoretically; importantly, it is also true scheme-theoretically). A pleasant consequence of this is that it is also possible to define D_σ for *partial* flags, provided that the flags have strata of all coranks mentioned in $\text{Ess}(\sigma)$.

Definition 3.3. Let $\mathcal{P}^\bullet, \mathcal{Q}^\bullet$ be flags with coranks A, B respectively, and let σ be a permutation of $[0, n - 1]$ such that $\text{Ess}(\sigma) \subseteq A \times B$. Define

$$D_\sigma(\mathcal{P}^\bullet; \mathcal{Q}^\bullet) = \{x \in S : \dim(\mathcal{P}^a)_x \cap (\mathcal{Q}^b)_x \geq r^\sigma(a, b) \text{ for all } (a, b) \in \text{Ess}(\sigma)\},$$

where the scheme structure is defined in the manner described above.

Permutations of $[0, n - 1]$ have a partial order, the *Bruhat order*, characterized by $\sigma \leq \tau$ if and only if $r^\sigma \geq r^\tau$. Therefore $\tau \leq \sigma$ implies $D_\tau(\mathcal{P}^\bullet; \mathcal{Q}^\bullet) \subseteq D_\sigma(\mathcal{P}^\bullet; \mathcal{Q}^\bullet)$, and $D_\sigma(\mathcal{P}^\bullet; \mathcal{Q}^\bullet)$ can also be described (set theoretically) as the locus of points $x \in S$ where the permutation τ associated to $\mathcal{P}_x, \mathcal{Q}_x$ satisfies $\tau \leq \sigma$ in Bruhat order.

Denote by $\text{Fl}(A; n)$ the flag variety parameterizing flags V^\bullet of coranks A in k^n . Taking \mathcal{V}^\bullet to be the tautological flag of coranks A on $\text{Fl}(A; n)$, and F^\bullet to be a fixed flag in k^n of coranks B , we denote

$$X_\sigma(F^\bullet) = D_\sigma(\mathcal{V}^\bullet; F^\bullet) \subseteq \text{Fl}(A; n),$$

where σ is any permutation compatible with coranks A, B . These are *Schubert varieties*, and they are prototypical of all degeneracy loci of pairs of flags. The literature on Schubert varieties is vast; we mention here only a few facts that we need in this paper.

Fact 3.4. Fix a permutation σ compatible with coranks A, B , and a fixed flag F^\bullet of coranks B in k^n . The Schubert variety $X_\sigma(F^\bullet)$ has codimension $\ell(\omega_{n-1}\sigma)$ in $\text{Fl}(A; n)$. Its singular locus is a union of Schubert varieties $X_\tau(F^\bullet)$ for certain permutations $\tau \leq \sigma$. All such τ have $\ell(\omega_{n-1}\tau) \geq \ell(\omega_{n-1}\sigma) + 2$, so Schubert varieties are smooth in codimension 1.

If $X_\tau(F^\bullet) \subseteq X_\sigma(F^\bullet)_{\text{sing}}$, we will say that τ is *singular* in X_σ . A combinatorial description of those σ, τ for which τ is singular in X_σ was conjectured in [LS90] and proved independently by several groups [BW03, Cor01, KLR03, Man01].

4. VERSALITY OF A PAIR OF FLAGS

This section gives a definition and some preliminary results about versality of flags, following the point of view of [CP19], and applies these results to relate versality of Brill-Noether flags to the coupled Petri map. Unlike [CP19], we do not assume the flags are complete, and we restrict our attention to pairs of *two* versal flags.

Throughout this section, let S be a finite-type scheme over an algebraically closed field k of any characteristic, with a rank n vector bundle \mathcal{H} . Suppose further that two (partial or complete) flags $\mathcal{P}^\bullet, \mathcal{Q}^\bullet$ are chosen in \mathcal{H} , with sets of coranks A, B respectively. We will reserve the letters a, b to denote elements of A or B respectively.

Denote by $f : \text{Fr}(\mathcal{H}) \rightarrow S$ the frame bundle of \mathcal{H} and by $\text{Fl}(A; n)$ the flag variety parameterizing flags F^\bullet of coranks A in k^n . The flags $\mathcal{P}^\bullet, \mathcal{Q}^\bullet$ induce a morphism $p : \text{Fr}(\mathcal{H}) \rightarrow \text{Fl}(A; n) \times \text{Fl}(B; n)$.

Definition 4.1. The pair of flags $\mathcal{P}^\bullet, \mathcal{Q}^\bullet$ is called *versal* if morphism $p : \text{Fr}(\mathcal{H}) \rightarrow \text{Fl}(A; n) \times \text{Fl}(B; n)$ described above is a smooth morphism.

Remark 4.2. Definition 4.1 may be stated stack-theoretically: the pair of flags $\mathcal{P}^\bullet, \mathcal{Q}^\bullet$ is versal if it induces a smooth map from S to the stack quotient $[\text{Fl}(A; d) \times \text{Fl}(B; d) / \text{GL}_n(k)]$. There is a moduli problem for pairs of flags, which is represented by this stack, and this explains our use of the term “versal.” The details are omitted since we do not need this point of view.

4.1. First-order deformations of a pair of flags. Fix a k -point $x \in S$. For simplicity, denote by H, P^\bullet, Q^\bullet the fibers of $\mathcal{H}, \mathcal{P}^\bullet, \mathcal{Q}^\bullet$ over x . Define

$$(13) \quad M_x = \text{coker} \left(\text{End } H \xrightarrow{\Delta} \text{End } H / \text{Fix } P^\bullet \oplus \text{End } H / \text{Fix } Q^\bullet \right).$$

where $\text{Fix } P^\bullet$ denotes $\{\phi \in \text{End } H : \phi(P^a) \subseteq P^a \text{ for all } a \in A\}$, $\text{Fix } Q^\bullet$ is defined similarly, and Δ is the diagonal map.

Fix a point $y \in f^{-1}(x)$. This amounts to an isomorphism $k^n \xrightarrow{\sim} H$, which induces an isomorphism

$$T_{p(y)} \text{Fl}(A; n) \times \text{Fl}(B; n) \cong \text{End } H / \text{Fix } P^\bullet \oplus \text{End } H / \text{Fix } Q^\bullet.$$

Using this identification we define a quotient map $q_y : T_{p(y)} \text{Fl}(A; n) \times \text{Fl}(B; n) \rightarrow M_x$. The kernel of q_y is isomorphic to the image of the diagonal map from $\text{End } H$, and may be identified with the tangent space to the $\text{GL}_n(k)$ -orbit of $p(y)$.

Lemma 4.3. *There is a linear map $\delta_x : T_x S \rightarrow M_x$, such that the following diagram commutes. In this diagram, the rows are exact, $\text{GL}_n(k)$ is identified with the fiber $f^{-1}(x)$ and the map to $T_y \text{Fr}(\mathcal{H})$ is the differential of the inclusion.*

$$\begin{array}{ccccccc} 0 & \longrightarrow & T_{\text{id}} \text{GL}_n(k) & \longrightarrow & T_y \text{Fr}(\mathcal{H}) & \xrightarrow{df_y} & T_x S \longrightarrow 0 \\ & & \parallel & & \downarrow dp_y & & \downarrow \delta_x \\ 0 & \longrightarrow & \text{End } H & \xrightarrow{\Delta} & T_{p(y)} \text{Fl}(A; d) \times \text{Fl}(B; d) & \xrightarrow{q_y} & M_x \longrightarrow 0 \end{array}$$

The map δ_x does not depend on the choice of $y \in f^{-1}(x)$.

We omit the proof of Lemma 4.3; see the discussion in [CP19, §3], which generalizes with minor notation changes to the case of incomplete flags.

Therefore the linear map δ_x describes the essential content of the differential dp_y , discarding any information that depends on the choice of frame. The utility of δ_x stems from the following Lemma, which boils down to the first-order criterion for smoothness of morphisms. We omit the proof; see [CP19, Proposition 3.2], which can be adapted to partial flags by minor notational changes.

Lemma 4.4. *The flag pair $\mathcal{P}^\bullet, \mathcal{Q}^\bullet$ is versal on S if and only if S is smooth over k and δ_x is surjective for all $x \in S$.*

We require a more concrete description of the map δ_x suitable to study Brill-Noether flags. Let $v \in T_x S$, and regard v as a morphism $v : \text{Spec } k[\varepsilon] \rightarrow S$, where $k[\varepsilon]$ is the ring of dual numbers. This gives an extension $0 \rightarrow H \rightarrow v^* \mathcal{H} \rightarrow H \rightarrow 0$, where the first map is multiplication by ε and the second is quotient modulo ε . A choice of $y \in f^{-1}(x)$ and $w \in df_y^{-1}(v)$ determines a trivialization $v^* \mathcal{H} \xrightarrow{\sim} k[\varepsilon]^n$, which determines a splitting $s : v^* \mathcal{H} \rightarrow H$. It is possible to inductively construct two splittings $\phi_P, \phi_Q : H \rightarrow v^* \mathcal{H}$ of the quotient map $v^* \mathcal{H} \rightarrow H$, such that $\phi_P(P^a) \subseteq v^* \mathcal{P}^a$ and $\phi_Q(Q^b) \subseteq v^* \mathcal{Q}^b$ for all $a \in A, b \in B$. Although these are not unique, $s \circ \phi_P$ is unique modulo $\text{Fix } P^\bullet$ and $s \circ \phi_Q$ is unique modulo $\text{Fix } Q^\bullet$. In fact, these two maps precisely give the differential of p :

$$(14) \quad dp_y(w) = (s \circ \phi_P + \text{Fix } P^\bullet, s \circ \phi_Q + \text{Fix } Q^\bullet) \in \text{End } H / \text{Fix } P^\bullet \oplus \text{End } H / \text{Fix } Q^\bullet.$$

Now, applying the quotient q_y to this formula, we obtain

Lemma 4.5. *For any $x \in S$ and $v \in T_x S$, let $s : v^* \mathcal{H} \rightarrow H$ be any k -linear splitting, and $\phi_P, \phi_Q : H \rightarrow v^* \mathcal{H}$ be any k -linear splittings such that $\phi_P(P^a) \subseteq v^* \mathcal{P}^a$ and $\phi_Q(Q^b) \subseteq v^* \mathcal{Q}^b$ for all $a \in A, b \in B$. Then*

$$\delta_x(v) = [s \circ \phi_P, s \circ \phi_Q],$$

where the square brackets indicate the coset in M_x .

4.2. A description of M_x^\vee . The dual vector space M_x^\vee has a convenient description that will be well-suited to relating it to the coupled Petri map. This description is based on the observation that elements of M_x encode first-order deformations of the relative positions of each pair P^a, Q^b of strata, via the following maps.

Definition 4.6. For $a \in A, b \in B$, define $r_{a,b} : M_x \rightarrow \text{Hom}(P^a \cap Q^b, H/(P^a + Q^b))$ by

$$r_{a,b}([\psi_P, \psi_Q]) = q \circ (\psi_P - \psi_Q) |_{P^a \cap Q^b},$$

where q denotes the quotient map $H \rightarrow H/(P^a + Q^b)$.

In the following statement, \perp denotes the annihilator subspace in H^\vee , all summands are regarded as subspaces of $H^\vee \otimes H$, and we implicitly identify $(H/(P^a + Q^b))^\vee$ with $(P^a + Q^b)^\perp$ and $(P^a + Q^b)^\perp \otimes (P^a \cap Q^b)$ with $\text{Hom}(P^a \cap Q^b, H/(P^a + Q^b))^\vee$.

Proposition 4.7. *There is an isomorphism*

$$\zeta : \sum_{a \in A, b \in B} (P^a + Q^b)^\perp \otimes (P^a \cap Q^b) \rightarrow M_x^\vee$$

such that for all a, b the induced map $(P^a + Q^b)^\perp \otimes (P^a \cap Q^b) \rightarrow M_x^\vee$ is equal to ${}^t r_{a,b}$.

Proof. First, we claim that the product map

$$r = \prod_{a,b} r_{a,b} : M_x \rightarrow \prod_{a \in A, b \in B} \text{Hom}(P^a \cap Q^b, H/(P^a + Q^b))$$

is injective. To see this, first choose a basis \mathcal{B} that is adapted to P^\bullet, Q^\bullet (see Fact 3.2). Any element of M_x is equal to $[\psi, 0]$ for some $\psi \in \text{End } H$, since $[\alpha, \beta] = [\alpha - \beta, 0]$; let $[\psi, 0] \in M_x$ be any element of $\ker r$. We will show that $[\psi, 0] = 0$, by constructing $\psi_1 \in \text{Fix } P^\bullet, \psi_2 \in \text{Fix } Q^\bullet$ such that $\psi = \psi_1 - \psi_2$. We will construct ψ_1, ψ_2 , by specifying their values on each element of \mathcal{B} . For each $v \in \mathcal{B}$, choose $a \in A, b \in B$ maximal such that $v \in P^a \cap Q^b$; the assumption $[\psi, 0] \in \ker r_{a,b}$ implies that $\psi(v) \in P^a + Q^b$, so we may write $v = v_1 - v_2$ for some $v_1 \in P^a, v_2 \in Q^b$. Define

$$\begin{array}{ccccccc}
& & & & & & 0 \\
& & & & & & \downarrow \\
& & & & & & P^a \cap Q^b \\
& & & & & & \downarrow \iota \\
& & & & & & \downarrow \phi_P \oplus \phi_Q \\
0 & \longrightarrow & P^a \oplus Q^b & \longrightarrow & v^*P^a \oplus v^*Q^b & \longrightarrow & P^a \oplus Q^b \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
& & & & & & \downarrow \\
0 & \longrightarrow & H & \xleftarrow{s} & v^*H & \longrightarrow & H \longrightarrow 0 \\
& & \downarrow q & & & & \downarrow \\
& & & & & & H/(P^a + Q^b) \\
& & & & & & \downarrow \\
& & & & & & 0
\end{array}$$

FIGURE 3. The composition $r_{a,b} \circ \delta_x(v)$ as a snake map.

$\psi_1(v) = v_1$, $\psi_2(v) = v_2$. If ψ_1, ψ_2 are constructed in this way, then $[\psi, 0] = [\psi_1, \psi_2] = 0 \in M_x$. This establishes that $\prod_{a,b} r_{a,b}$ is injective.

Now, define $\Delta : \text{End } H \rightarrow \prod_{a,b} \text{Hom}(P^a \cap Q^b, H/(P^a + Q^b))$ to be the diagonal map, and $\pi : \text{End } H \rightarrow M$ to be the map $\psi \mapsto [\psi, 0]$. Then $r \circ \pi = \Delta$. Dualizing, and using the fact that r is injective and π is surjective, we obtain an isomorphism $\zeta : \text{im } {}^t\Delta \xrightarrow{\sim} M^\vee$. Identifying $\text{im } {}^t\Delta$ with $\sum_{a,b} (P^a + Q^b)^\perp \otimes (P^a \cap Q^b)$ gives the desired isomorphism. It follows from the construction that $\zeta_{a,b} = {}^t r_{a,b}$. \square

The isomorphism ζ demonstrates that the map δ_x may be conveniently studied via the compositions $r_{a,b} \circ \delta_x$, which record first-order deformations of the pair P^a, Q^b of subspaces. These maps have a convenient cohomological description, which is shown in Figure 3. Consider the difference morphism $\mathcal{P}^a \oplus \mathcal{Q}^b \rightarrow \mathcal{H}$ given by $(\alpha, \beta) \mapsto \alpha - \beta$. Given a tangent vector $v \in T_x S$, we obtain a map of short exact sequences from $0 \rightarrow P^a \oplus Q^b \rightarrow v^*P^a \oplus v^*Q^b \rightarrow P^a \oplus Q^b \rightarrow 0$ to $0 \rightarrow H \rightarrow v^*H \rightarrow H \rightarrow 0$. Now, identifying the kernel of $P^a \oplus Q^b \rightarrow H$ with $P^a \cap Q^b$, this map of short exact sequences gives a snake map $P^a \cap Q^b \rightarrow H/(P^a + Q^b)$. This snake map can be described explicitly as $q \circ s \circ (\phi_P - \phi_Q) \circ \iota$, where $q : H \rightarrow H/(P^a + Q^b)$ is the quotient map, s, ϕ_P, ϕ_Q are splitting as in Lemma 4.5, and ι is the inclusion $P^a \cap Q^b \rightarrow H$. But $q \circ s \circ (\phi_P - \phi_Q) \circ \iota$ is also equal to $r_{a,b} \circ \delta_x$. This proves

Lemma 4.8. *Let $v \in T_x S$ be a tangent vector. For all $a \in A, b \in B$, the map $r_{a,b} \circ \delta_x(v)$ is equal to snake map described above and shown in Figure 3.*

4.3. First-order deformation of Brill-Noether flags. We now specialize to our intended application: let (C, p, q) be a twice-marked smooth curve over an algebraically closed field k , $S = \text{Pic}^d(C)$ for some integer $d \geq 2g - 1$, and consider the degree- d Brill-Noether flags $\mathcal{P}_d^\bullet, \mathcal{Q}_d^\bullet$ in the vector bundle \mathcal{H}_d . For convenience, let $n = d + 1 - g = \text{rank}(\mathcal{H}_d)$.

Fix a point $[L] \in \text{Pic}^d(C)$. In the notation this section, $H = H^0(C, L)$, $P^a = H^0(C, L(-ap))$, $Q^b = H^0(C, L(-bq))$, $P^a \cap Q^b = H^0(C, L(-ap - bq))$, and the sets of coranks are $A = B = [0, d - 2g + 1]$. For all $a \in A, b \in B$, there is an exact sequence of \mathcal{O}_C -modules

$$(15) \quad 0 \rightarrow L(-ap - bq) \rightarrow L(-ap) \oplus L(-bq) \rightarrow L \rightarrow 0,$$

where the second map is given by $(s, t) \mapsto s - t$ on sections. Taking cohomology, and using the fact that $L, L(-ap)$, and $L(-bq)$ are nonspecial line bundles, gives an isomorphism

$$h_{a,b} : H/(P^a + Q^b) \xrightarrow{\sim} H^1(C, L(-ap - bq)).$$

Dualizing these maps and using functoriality of the long exact sequence gives an isomorphism

$$(16) \quad \theta : \sum_{0 \leq a, b \leq n} H^1(C, L(-ap - bq))^\vee \otimes H^0(L(-ap - bq)) \xrightarrow{\sim} \sum_{0 \leq a, b \leq n} (P^a + Q^b)^\perp \otimes (P^a \cap Q^b).$$

Using Serre duality and identifying the domain of θ with $T_{p,q}^L|_{[0,n]^2}$ gives an isomorphism

$$\zeta \circ \theta : T_{p,q}^L|_{[0,n]^2} \xrightarrow{\sim} M_{[L]}^\vee.$$

Identifying the cotangent space $T_x^\vee S \cong H^1(C, \mathcal{O}_C)^\vee$ with $H^0(C, \omega_C)$, regard ${}^t\delta_{[L]}$ as a map

$${}^t\delta_{[L]} : M_{[L]}^\vee \rightarrow H^0(C, \omega_C).$$

We may finally link versality of Brill-Noether flags to the coupled Petri map.

Proposition 4.9. *The map $\eta = {}^t\delta_{[L]} \circ \zeta \circ \theta$ is equal to the $[0, n]^2$ -coupled Petri map $\mu_{p,q}^L|_{[0,n]^2}$.*

Proof. Let $\eta_{a,b} = \eta \circ \iota$ denote the restriction of η to $H^0(C, L(-ap - bq)) \otimes H^0(C, \omega_C \otimes L^\vee(ap + bq))$. Then ${}^t\eta_{a,b} = {}^t\iota \circ {}^t\theta \circ {}^t\zeta \circ \delta_{[L]} = h_{a,b} \circ r_{a,b} \circ \delta_{[L]} = h_{a,b} \circ s$, where s is examined in Lemma 4.8: it is the map

$$s : T_{[L]} \text{Pic}^d(C) \rightarrow \text{Hom}(H^0(C, L(-ap - bq)), H^1(C, L(-ap - bq)))$$

such that for all $v \in T_{[L]} \text{Pic}^d(C)$, the map $s(v)$ is equal to the snake map in Lemma 4.8. That snake diagram has a cohomological interpretation. Fix $v \in T_{[L]} \text{Pic}^d(C)$, corresponding to an extension $0 \rightarrow L \rightarrow \mathcal{L} \rightarrow L \rightarrow 0$ of L to a line bundle on $C \times_k \text{Spec } k[\varepsilon]$, and consider the following map of short exact sequences.

$$\begin{array}{ccccccc} 0 & \longrightarrow & L(-ap) \oplus L(-bq) & \longrightarrow & \mathcal{L}(-ap) \oplus \mathcal{L}(-bq) & \longrightarrow & L(-ap) \oplus L(-bq) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & L & \longrightarrow & \mathcal{L} & \longrightarrow & L \longrightarrow 0 \end{array}$$

Observe that none of these sheaves have higher cohomology, and the kernels of the vertical maps are $L(-ap - bq), \mathcal{L}(-ap - bq), L(-ap - bq)$, respectively. In other words, this diagram gives an acyclic resolution of the exact sequence

$$(17) \quad 0 \rightarrow L(-ap - bq) \rightarrow \mathcal{L}(-ap - bq) \rightarrow L(-ap - bq) \rightarrow 0.$$

Therefore, applying the global sections functor and forming the snake diagram gives the long exact cohomology sequence of (17), and the snake map of Figure 3 is equal, up to the isomorphism $h_{a,b}$, to the cohomology boundary map

$$(18) \quad \partial : H^0(C, L(-ap - bq)) \rightarrow H^1(C, L(-ap - bq)).$$

In other words, $\partial = h_{a,b} \circ r_{a,b} \circ \delta_{[L]}(v)$. On the other hand, ∂ is also given by taking the cup product with v , where we now regard v as an element of $H^1(C, \mathcal{O}_C)^2$. In other words, the map

$${}^t\eta_{a,b} : T_{[L]} \text{Pic}^d(C) \cong H^1(C, \mathcal{O}_C) \rightarrow \text{Hom}(H^0(C, L(-ap - bq)), H^1(C, L(-ap - bq)))$$

²This can be seen explicitly by representing the extension \mathcal{L} by a Čech cocycle, or more abstractly by noting that in the long exact sequence for $\text{Hom}_{\mathcal{O}_C}(\mathcal{O}_C, \bullet)$, the boundary map $\text{Hom}_{\mathcal{O}_C}(\mathcal{O}_C, L) \rightarrow \text{Ext}_{\mathcal{O}_C}^1(\mathcal{O}_C, L)$ is given by taking the Yoneda product with the class in $\text{Ext}_{\mathcal{O}_C}^1(L, L)$ of the extension; the latter is identified with $v \in H^1(C, \mathcal{O}_C)$.

is given by ${}^t\eta_{a,b}(v)(\sigma) = v \cup \sigma$. Dualizing, and unwinding definitions (including the description of the Serre duality isomorphism as cup product followed by $H^1(C, \omega_C) \xrightarrow{\sim} k$), it follows that

$$\eta_{a,b} : H^0(C, \omega_C \otimes L^\vee(ap + bq)) \otimes H^0(C, L(-ap - bq)) \rightarrow H^0(C, \omega_C)$$

is the cup product map. Considering all the $\eta_{a,b}$ together shows that $\eta = \mu_{p,q}^L|_{[0,n]^2}$. \square

Corollary 4.10. *Let (C, p, q) be a twice-marked smooth curve over an algebraically closed field k . The degree- d Brill-Noether flags of (C, p, q) are versal at $[L] \in \text{Pic}^d(C)$ if and only if the $[0, d - 2g + 1]^2$ -coupled Petri map $\mu_{p,q}^L|_{[0, d - 2g + 1]^2}$ is injective.*

5. PROOF OF THE VERSALITY THEOREM

We now have all the pieces in place to prove Theorem 1.7 on the S -coupled Petri condition, and the versality Theorem 1.1. All that remains is to use some properties of the moduli space of curves to obtain the desired results for all algebraically closed fields.

Proof of Theorem 1.1. If $\pi : \mathcal{C} \rightarrow B$ is a proper flat family of smooth curves with two disjoint sections P, Q , the construction of degree- d Brill-Noether flags globalizes to give flags $\mathcal{P}_d^\bullet, \mathcal{Q}_d^\bullet$ in a vector bundle \mathcal{H}_d on the relative Picard variety $\text{Pic}^d(\pi) \rightarrow B$. Define a morphism $\text{Fr}(\mathcal{H}_d) \rightarrow \text{Fl}(A; n) \times \text{Fl}(B; n)$ as in Definition 4.1. The locus where this map fails to be smooth is a closed subscheme, as is its image in B . So the locus of $b \in B$ where the degree- d Brill-Noether flags are versal is open. Since $\mathcal{M}_{g,2}$ is irreducible, we need only verify the existence of a twice-marked curve of genus g with versal degree- d Brill-Noether flags over some field in every characteristic. This can be done as follows: find a twice-marked elliptic curve (C, p, q) with $p - q$ non-torsion, chain g such curves together, and deform to an arithmetic surface over a discrete valuation ring with fraction field of the desired characteristic. By Theorem 2.2, the geometric general fiber satisfies the fully coupled Petri condition, so by Corollary 4.10 its Brill-Noether flags (in every degree) are versal. \square

Proof of Theorem 1.7. Fix a genus g , an algebraically closed field k , and a finite set $S \subseteq \mathbb{Z} \times \mathbb{Z}$. Let $\mathcal{V} \subseteq \mathcal{M}_{g,2}$ denote the locus of twice-marked curves satisfying the S -coupled Petri condition, and for all $d \in \mathbb{Z}$ let \mathcal{V}_d denote the locus of twice-marked curves for which every degree- d line bundle has injective S -coupled Petri map. So $\mathcal{V} = \bigcap_{d \in \mathbb{Z}} \mathcal{V}_d$. For all but finitely many $d \in \mathbb{Z}$, all elements of $\{d + a + b : (a, b) \in S\}$ are either less than 0 or greater than $2g - 2$; for line bundles L of these degrees the space $T_{p,q}^L(S)$ is trivial, so the S -coupled Petri map is injective. Therefore all but finitely many \mathcal{V}_d are equal to all of $\mathcal{M}_{g,2}$, so it suffices to fix a single integer d and verify that \mathcal{V}_d is Zariski-dense.

For all $N \in \mathbb{Z}$, the S -coupled Petri map of a degree- d line bundle L is equal to the S' -coupled Petri map of the degree- $(d + 2N)$ line bundle $L(Np + Nq)$, where $S' = \{(a + N, b + N) : (a, b) \in S\}$. For N sufficiently large, $S' \subseteq [0, d + 2N - 2g + 1]^2$. By Corollary 4.10, \mathcal{V}_d contains the locus of twice-marked curves with versal degree- $(d + 2N)$ Brill-Noether flags. This is dense by Theorem 1.1, hence \mathcal{V}_d is dense for all $d \in \mathbb{Z}$. \square

6. BRILL-NOETHER DEGENERACY LOCI

This section analyzes the geometry of Brill-Noether degeneracy loci $W_d^\Pi(C, p, q)$ using the versality of Brill-Noether flags, and proves Theorem 1.12. Assume we are in the following situation.

Situation 6.1. Let g, d be positive integers, (C, p, q) a twice-marked smooth curve of genus g , and Π a dot array of size $r + 1$ with $g - d + r \geq 0$. Also assume that $\Pi \subseteq [0, d]^2$.




FIGURE 4. Extension of a dot pattern with $r + 1 = |\Pi| = 3$ to a (d, g) -confined permutation, for various values of $d - g$. The dot pattern in the last example is not (d, g) -confined. The dashed squares are explained in Lemma 6.7.

The simplifying assumption $\Pi \subseteq [0, d]^2$ is harmless; see the proof of Theorem 1.12.

We begin by specifying the scheme structure on $W_d^\Pi(C, p, q)$ and interpreting it as a degeneracy locus of Brill-Noether flags. First, we recall the construction of the scheme structure of $W_d^r(C)$ (see e.g. [ACGH85, §IV.3]): the set-theoretic locus $\{[L] \in \text{Pic}^d(C) : h^0(C, L) \geq r + 1\} = \{[L] \in \text{Pic}^d(C) : h^1(C, L) \geq g - d + r\}$ has a natural scheme structure given by the $(g - d + r)$ th Fitting ideal of $R^1\nu_*\mathcal{L}$, where $\nu : C \times \text{Pic}^d(C) \rightarrow \text{Pic}^d(C)$ is the projection and \mathcal{L} is a Poincaré line bundle. The Fitting ideal may be computed as a determinantal locus using any resolution by vector bundles. In particular, if we choose any two integers $a, b \geq 2g - 1 - d$ and let $d' = d + a + b$, we have a resolution $\mathcal{P}_{d'}^a \oplus \mathcal{Q}_{d'}^b \rightarrow \mathcal{H}_{d'} \rightarrow R^1\nu_*\mathcal{L} \rightarrow 0$, and the Fitting ideal is therefore equal to the determinantal ideal defining the standard scheme structure of $\{x \in \text{Pic}^{d'}(C) : \dim(\mathcal{P}_{d'}^a)_x \cap (\mathcal{Q}_{d'}^b)_x \geq r + 1\}$, as described in Section 3.2.

Now, we define the scheme structure on $W_d^\Pi(C, p, q)$ by a scheme-theoretic intersection

$$(19) \quad W_d^\Pi(C, p, q) = \bigcap_{(a,b) \in \mathbb{N}^2} \text{tw}_{ap+bq} \left(W_{d-a-b}^{r(a,b)-1}(C) \right).$$

This in turn may be rephrased as a degeneracy locus of Brill-Noether flags of larger degree. For any two integers $M, N \geq 2g - 1$, we may define $d' = d + M + N$ and write equivalently

$$(20) \quad W_d^\Pi(C, p, q) = \text{tw}_{-Mp-Nq} \left(\{x \in \text{Pic}^{d'}(C) : \dim(\mathcal{P}_{d'}^{M+a})_x \cap (\mathcal{Q}_{d'}^{N+b})_x \geq r^\Pi(a, b) \text{ for all } a, b \in [0, d]\} \right).$$

The bounds $M, N \geq 2g - 1$ ensure that $M + a, N + b \leq d' - 2g + 1$ for all $a, b \in [0, d]$, so that the flag elements $\mathcal{P}_{d'}^{M+a}, \mathcal{Q}_{d'}^{N+b}$ exist.

Now, we wish to write this locus as a degeneracy locus $D_\sigma(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet)$ of the form discussed in Section 3.2. To do so, we must convert the dot pattern Π to a permutation. This requires a few combinatorial preliminaries.

6.1. (d, g) -confined permutations. We wish to extend a dot pattern $\Pi \subseteq \mathbb{N} \times \mathbb{N}$ to a dot pattern in $\mathbb{Z} \times \mathbb{Z}$ in the “most efficient way,” so that a rank function for L may be translated into a rank function for $L(Mp + Nq)$ that results in the same degeneracy locus.

Definition 6.2. A permutation $\pi : \mathbb{Z} \rightarrow \mathbb{Z}$ is called (d, g) -confined if both π, π^{-1} are decreasing on $(-\infty, -1]$ and $\omega_{d-g}\pi$ has finite length.

Note that the definition of (d, g) -confined actually depends only on the difference $d - g$. We may extend Equation 9 to (d, g) -confined permutations to define, for all $a, b \in \mathbb{Z}$,

$$(21) \quad r^\pi(a, b) = \#\{a' \geq a : \pi(a') \geq b\},$$

and the requirement that $\omega_{d-g}\pi$ has finite length ensures that $r^\pi(-a, -b) = d + a + b + 1 - g$ for sufficiently large a, b , in accordance with Riemann-Roch.

Lemma 6.3. *Fix positive integers d, g . For every dot pattern Π with $|\Pi| \geq d + 1 - g$, there is a unique (d, g) -confined permutation π such that $\Pi = \{(a, \pi(a)) : a \geq 0 \text{ and } \pi(a) \geq 0\}$.*

Proof. This is analogous to an infinite-grid version of the process described in [FP98, p. 9]. Let $r = |\Pi| - 1$ and $A = \{a_0, \dots, a_r\}$ and $B = \{b_0, \dots, b_r\}$ be the set of rows and set of columns of Π . A permutation π of \mathbb{Z} satisfies the desired conditions if and only if

- (1) $\pi|_A$ is the bijection $A \rightarrow B$ with graph Π ,
- (2) $\pi|_{\mathbb{Z} \setminus A}$ is a *decreasing* bijection $\mathbb{Z} \setminus A \rightarrow \mathbb{Z} \setminus B$ such that $\pi(n) = d - g - n$ for all but finitely many n , and
- (3) $\pi(\mathbb{N} \setminus A)$ is disjoint from $\mathbb{N} \setminus B$.

For any dot pattern $\Pi \subset \mathbb{N} \times \mathbb{N}$ (regardless of $|\Pi|$), there is a unique π satisfying conditions (1) and (2). For N sufficiently large, this permutation induces a bijection $[-N, (d - g) + N] \setminus A \rightarrow [-N, (d - g) + N] \setminus B$, and condition (3) is satisfied if and only if

$$|[0, (d - g) + N] \setminus A| \leq |[-N, -1]|,$$

i.e. $d - g - r + N \leq N$. So a permutation π satisfying (1-3) above exists if and only if $g - d + r \geq 0$, i.e. $|\Pi| \geq d + 1 - g$. \square

The rightmost grid in Figure 4 illustrates where Lemma 6.3 breaks down when Π has too few dots: there is not enough space outside $\mathbb{N} \times \mathbb{N}$ to complete the dot pattern (hence a white dot must be added within $\mathbb{N} \times \mathbb{N}$).

Remark 6.4. Observe that for all $a, b \in \mathbb{N}$, $r^\Pi(a, b) = r^\pi(a, b)$. So r^π may be viewed as the “most efficient” extension of r^Π to all of $\mathbb{Z} \times \mathbb{Z}$. However, this extension depends on the choice of (d, g) (rather, on $d - g$), which is why we made the ad hoc choice to define $r^\Pi(a, -1) = r^\Pi(a, 0)$, $r^\Pi(-1, b) = r^\Pi(0, b)$ when defining $\text{Ess}(\Pi)$ in Equation 4. The following lemma shows that this ad hoc choice makes almost no difference anyway.

Lemma 6.5. *Let Π be a dot pattern of size $r + 1$, and let a_0, b_0 be the minimum row and column occurring in Π . Let π be its (d, g) -confined permutation, where $g - d + r \geq 0$. Then $\text{Ess}(\Pi) = \text{Ess}(\pi)$ unless $g - d + r = 0$ and $a_0 = b_0 = 0$. In the case $g - d + r = 0, a_0 = b_0 = 0$, we have instead $(0, 0) \notin \text{Ess}(\pi)$ and $\text{Ess}(\Pi) = \text{Ess}(\pi) \cup \{(0, 0)\}$.*

In what follows, we will only actually need the inclusion $\text{Ess}(\pi) \subseteq \text{Ess}(\Pi)$, but we prove the sharper statement here partly to justify that the definition of $\text{Ess}(\Pi)$ was sensible.

Proof. The monotonicity of π, π^{-1} on $(-\infty, -1]$ implies that $\text{Ess}(\pi) \subseteq \mathbb{N}^2$. Comparing Equations 4 and 12 and the definitions of r^Π and r^π shows that $\text{Ess}(\pi) \subseteq \text{Ess}(\Pi)$. Suppose that $(a, b) \in \text{Ess}(\Pi) \setminus \text{Ess}(\pi)$. Since r^π and r^Π match on \mathbb{N}^2 , either $a = 0$ or $b = 0$. Consider first the case where $a > 0, b = 0$. Then $(a, 0) \in \text{Ess}(\Pi)$ implies that $\pi(a - 1) < 0 \leq \pi(a)$ and $a \leq \pi^{-1}(0)$; $(a, 0) \notin \text{Ess}(\pi)$ implies that $a \leq \pi^{-1}(-1)$ as well. The construction of π in the proof of Lemma 6.3 shows that $\pi^{-1}(-1)$ is the minimum element of $[-(g - d + r), \infty) \setminus A$, where A is the set of rows occurring in Π . So $a \leq \pi^{-1}(-1)$ implies that $g - d + r = 0$ and that $0, 1, 2, \dots, a$ are all rows occurring in Π . But that implies that $\pi^{-1}(a - 1) \geq 0$, a contradiction. So this case is impossible. Switching rows with columns shows that the case $a = 0, b > 0$ is also impossible. So the only possible element of




FIGURE 5. The number of non-inversions of π is $g - \rho_g(d, \Pi)$.

$\text{Ess}(\Pi) \setminus \text{Ess}(\pi)$ is $(0, 0)$. Now, observe that $(0, 0) \in \text{Ess}(\Pi)$ if and only if $a_0 = b_0 = 0$. On the other hand, $(0, 0) \in \text{Ess}(\pi)$ if and only if $\pi(-1) < 0 \leq \pi(0)$ and $\pi^{-1}(-1) < 0 \leq \pi^{-1}(0)$. Now, $\pi(-1) < 0$ and $\pi^{-1}(-1) < 0$ are both equivalent to $g - d + r > 0$, while $\pi(0) \geq 0$ and $\pi^{-1}(0) \geq 0$ are equivalent to $b_0 = 0$ and $a_0 = 0$, respectively. So $(0, 0) \in \text{Ess}(\Pi) \setminus \text{Ess}(\pi)$ if and only if $g - d + r = 0$ and $a_0 = b_0 = 0$. \square

The (d, g) -confined permutation associated to Π also provides a convenient and concise reformulation of the expected dimension $\rho_g(d, \Pi)$.

Lemma 6.6. *If the dot pattern Π has $|\Pi| \geq d + 1 - g$ and associated (d, g) -confined permutation π , then $\rho_g(d, \Pi) = g - \ell(\omega_{d-g}\pi)$.*

We omit the straightforward proof of Lemma 6.6, but instead illustrate it visually in Figure 5.

In the proof of Theorem 1.12, the replacement of a line bundle L by $L(ap + bq)$ has the effect of sliding the graph of π down a steps and to the right b steps. We will want to do this in such a way that the portion in $\mathbb{N} \times \mathbb{N}$ is the graph of a permutation of $[0, n]$ for some integer n . The dashed squares in Figure 4 show what we are looking for: squares that enclose the graph of a permutation. The following Lemma formalizes how such squares can be found.

Lemma 6.7. *Let π be the (d, g) -confined permutation associated to a dot pattern Π of size $r + 1$, and let a_r, b_r be the largest row and largest column occurring in Π , respectively. Then π restricts to a bijection $[d - g - b_r, a_r] \rightarrow [d - g - a_r, b_r]$, and $\pi(n) = d - g - n$ for all n such that $n < d - g - b_r$ or $n > a_r$. Also, π is decreasing on $[a_r, \infty)$, and π^{-1} is decreasing on $[b_r, \infty)$.*

Proof. By the construction in the proof of Lemma 6.3, π restricts to a decreasing bijection between $(-\infty, d - g - b_r - 1] \cup [a_r + 1, \infty)$ and $(\infty, d - g - a_r - 1] \cup [b_r + 1, \infty)$. Such a bijection is determined by its value on a single point, and the fact that $\omega_{d-g}\pi$ has finite length forces $\pi(n) = d - g - n$ for all n in this restricted domain. This implies that π also restricts to a bijection between the complements $[d - g - b_r, a_r]$ and $[d - g - a_r, b_r]$. The fact that π is decreasing on $[a_r + 1, \infty)$ is clear; note finally that $\pi(a_r) \in [d - g - a_r, b_r]$ and $\pi(a_r + 1) = d - g - a_r - 1$, so $\pi(a_r) > \pi(a_r + 1)$ and π is decreasing on all of $[a_r, \infty)$. By the same argument applied to π^{-1} , π^{-1} is decreasing on $[b_r, \infty)$. \square

For fixed $m \in \mathbb{Z}$, denote by α_m the “add m ” permutation $\alpha_m(n) = m + n$. The following corollary will be useful in converting Brill-Noether degeneracy loci to degeneracy loci of Brill-Noether flags.

Corollary 6.8. *If $M \geq b_r - (d - g)$ and $N \geq a_r - (d - g)$, then the permutation $\alpha_N \pi \alpha_M^{-1}$ restricts to a bijection from $[0, M + N + d - g]$ to itself that is decreasing on $[a_r + M, \infty)$, and its inverse is decreasing on $[b_r + N, \infty)$.*

6.2. Geometry of Brill-Noether degeneracy loci. We now have what we need to study Brill-Noether degeneracy loci as degeneracy loci of Brill-Noether flags. With an eye on Equation 20, we add a bit more notation to Situation 6.1.

Situation 6.9. In Situation 6.1, also fix integers $M, N \geq 2g - 1$, define $d' = d + M + N$, and define $\pi' = \alpha_N \pi \alpha_M^{-1}$. For convenience, define $n = d' + 1 - g$ (the rank of $\mathcal{H}_{d'}$).

If a_r, b_r are the largest row and column occurring in Π , the assumption $\Pi \subseteq [0, d]$ (from Situation 6.1) ensures $a_r, b_r \leq d$. So Corollary 6.8 implies that π' restricts to a permutation of $[0, n - 1]$, and that π' and its inverse are both decreasing on $[d' - (2g - 1), n - 1]$. Both flags $\mathcal{P}_{d'}, \mathcal{Q}_{d'}$ have corank sets $[0, d' - (2g - 1)]$ so π' meets the monotonicity requirements of Definition 3.3, and there is a well-defined degeneracy locus $D_{\pi'}(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet)$. Furthermore, adding degeneracy conditions for pairs (a, b) not in the essential set does not change the scheme structure, so Equation 20, Lemma 6.5 and the observation that $\text{Ess}(\pi') = \text{Ess}(\pi) + (M, N)$ imply that

$$(22) \quad W_d^\Pi(C, p, q) = \text{tw}_{-Mp-Nq}(D_{\pi'}(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet)).$$

So Brill-Noether degeneracy loci are just degeneracy loci of Brill-Noether flags, hiding behind a twist. We can now use the versality theorem to prove the local statements from Theorem 1.12, including a stronger form of the smoothness statement.

Theorem 6.10. *In Situation 6.9, assume that (C, p, q) has versal degree- d' Brill-Noether flags. Let $[L]$ be any point of $W_d^\Pi(C, p, q)$. The local dimension of $W_d^\Pi(C, p, q)$ is $\rho_g(d, \Pi)$.*

Furthermore, let $L' = L(Mp + Nq)$, and let $\sigma \leq \pi'$ be the permutation associated to the flags $\mathcal{P}_{[L']}^\bullet, \mathcal{Q}_{[L']}^\bullet$. Then $W_d^\Pi(C, p, q)$ is singular at $[L]$ if and only if σ is singular in $X_{\pi'}$. If $[L] \in \widetilde{W}_d^\Pi(C, p, q)$ then $W_d^\Pi(C, p, q)$ is smooth at $[L]$.

Proof. By definition of versality and basic properties of smooth morphisms, the local codimension at $[L']$ of $D_{\pi'}(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet) \subset \text{Pic}^{d'}(C)$ is equal to the local codimension at a point $y \in \text{Fl}(A; n) \times \text{Fl}(B; n)$ of $D_{\pi'}(\mathcal{V}^\bullet; \mathcal{W}^\bullet)$, where $\mathcal{V}^\bullet, \mathcal{W}^\bullet$ are the two (pull-backs of) tautological flags, and y is a point where the permutation associated to $\mathcal{V}_y^\bullet, \mathcal{W}_y^\bullet$ is equal to σ . Using the fact that the tautological flag of $\text{Fl}(A; n)$ is versal to any fixed flag F^\bullet , we can apply the same reasoning to deduce that the codimension at y of $D_{\pi'}(\mathcal{V}^\bullet; \mathcal{W}^\bullet)$ is equal to the codimension at $z \in \text{Fl}(A; d)$ of $X_{\pi'}(F^\bullet)$, where z is a point corresponding to a flag V^\bullet such that the permutation associated to V^\bullet, F^\bullet is σ . The Schubert variety $X_{\pi'}$ has pure codimension $\ell(\omega_{n-1}\pi') = g - \rho_g(d, \Pi)$, so we obtain the desired local dimension statement. Furthermore, since smooth morphisms preserve smoothness of subschemes under inverse image, $[L]$ is a smooth point of $W_d^\Pi(C, p, q)$ if and only if z is a smooth point of $X_{\pi'}(F^\bullet)$, which holds if and only if σ is singular in $X_{\pi'}$, as desired.

There is one final wrinkle in proving that $\widetilde{W}_d^\Pi(C, p, q)$ is smooth: we must show that if $[L] \in \widetilde{W}_d^\Pi(C, p, q)$ then σ is in the smooth locus of $X_{\pi'}$. It is tempting to say that $\pi' = \sigma$, but unfortunately this need not be true if the strata $\mathcal{P}_{d'}^a, \mathcal{Q}_{d'}^b$ with $(a + M, b + N) \notin \text{EssR}(\Pi) \times \text{EssC}(\Pi)$ meet in dimension larger than $r^{\pi'}(a, b)$. One can resolve this by invoking a combinatorial description of the singular locus of $X_{\pi'}$, but we can also accomplish it with a cheap trick: simply replace $\mathcal{P}_{d'}^\bullet, \mathcal{Q}_{d'}^\bullet$ by subflags $\hat{\mathcal{P}}^\bullet, \hat{\mathcal{Q}}^\bullet$ in which we only retain the strata of coranks $A = \{a : \pi'(a) > \pi'(a - 1)\}$, $B = \{b : \pi'^{-1}(b) > \pi'^{-1}(b - 1)\}$. Observe that $A \times B \supseteq \text{Ess}(\pi')$, so the degeneracy locus $D_{\pi'}(\hat{\mathcal{P}}^\bullet; \hat{\mathcal{Q}}^\bullet)$ is well-defined and equal to $D_{\pi'}(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet)$. Also, Lemma 6.5 implies that $A \subseteq \text{EssR}(\Pi) + M$ and $B \subseteq \text{EssC}(\Pi) + N$, so $[L] \in \widetilde{W}_d^\Pi(C, p, q)$ implies that the permutation associated to the subflags $(\hat{\mathcal{P}}^\bullet; \hat{\mathcal{Q}}^\bullet)$ is exactly π' ; thus $[L']$ is a smooth point of $D_{\pi'}(\hat{\mathcal{P}}^\bullet; \hat{\mathcal{Q}}^\bullet)$ and $[L]$ is a smooth point of $W_d^\Pi(C, p, q)$. \square

Remark 6.11. The proof above follows the techniques of [CP19, §4], although we work here with possibly partial flags. Using [CP19, Theorem 4.4] (with mild modifications for partial flags) one can deduce much stronger results about the nature of the singularities of $W_d^\Pi(C, p, q)$; roughly speaking, they are isomorphic, étale-locally and up to products with affine space, to the singularities of Schubert varieties. Together with the results of [BW03, Cor01, KLR03, Man01], this gives essentially a complete description of the singularities of $W_d^\Pi(C, p, q)$.

6.3. The Chow class of $W_d^\Pi(C, p, q)$. To obtain the existence and intersection-theoretic part of Theorem 1.12, we use the results of [Ful92] about the intersection theory of degeneracy loci (see also the exposition in [FP98, §1-2]). We assume throughout this subsection that we are in Situation 6.9, and furthermore that $W_d^\Pi(C, p, q)$ has the expected dimension $\rho_g(d, \Pi)$ (e.g. it suffices to assume that the degree- d' Brill-Noether flags are versal, by Theorem 6.10).

Translating our notation (indexed by codimension, starting at 0) into the notation of [Ful92] (indexed by dimension, starting at 1), our permutation π' gives rise to a permutation w of $[1, n]$ via the formula

$$w(i) = d' + 1 - g - \pi(i - 1) \text{ for } 1 \leq i \leq n.$$

Note that $\ell(w) = \ell(\omega_{d'-g}\pi')$ is the codimension of $D_{\pi'}(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet)$.

Let x_1, \dots, x_n be Chern roots of the bundle $\mathcal{H}/\mathcal{P}^n$, ordered such that prefixes of this sequence give Chern roots of $\mathcal{H}/\mathcal{P}^a$ for each a . Let y_1, \dots, y_n be Chern roots of \mathcal{H} such that prefixes of this sequence give Chern roots of \mathcal{Q}^b for each b . As usual, these Chern roots cannot be taken literally as elements of the Chow ring of $\text{Pic}^d(C)$, but any symmetric polynomial in the first ℓ of them, for any $\ell \geq g$, is identified with a polynomial in the Chern classes of the bundles involved. Finally, let \mathfrak{S}_w be the double Schubert polynomial of Lascoux and Schützenberger. Since $\text{Pic}^d(C)$ is smooth and $D_{\pi'}(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet)$ has the expected dimension, [Ful92, Theorem 8.2] implies that in the Chow ring,

$$(23) \quad [D_{\pi'}(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet)] = \mathfrak{S}_w(x_1, \dots, x_n; y_1, \dots, y_n).$$

The calculation in [ACGH85, §VII.4] shows that all of the bundles $\mathcal{H}_{d'}, \mathcal{P}_{d'}^a, \mathcal{Q}_{d'}^b$ have the same total Chern class $e^{-\Theta}$, where Θ is the theta class in $\text{Pic}^d(C)$. It follows from this that the Chern roots in Equation 23 simplify considerably. All the quotients $\mathcal{H}/\mathcal{P}^a$ have trivial Chern classes, so $x_1 = \dots = x_n = 0$. The first g roots y_1, \dots, y_g are Chern roots of the rank- g bundle \mathcal{Q}^{n-g} , which has Chern class $e^{-\Theta}$. If e_k denotes the degree- k elementary symmetric polynomial, this means that $e_k(y_1, \dots, y_g) = (-1)^k \frac{\Theta^k}{k!}$. Each quotient $\mathcal{Q}^b/\mathcal{Q}^{b+1}$ has trivial total Chern class, so $y_{g+1} = \dots = y_n = 0$. Using the formula $\mathfrak{S}_w(x, y) = (-1)^{\ell(w)} \mathfrak{S}_{w^{-1}}(y, x)$, and the fact that \mathfrak{S}_w is a polynomial of degree $\ell(w)$ (e.g. [FP98, §1.3]), we may rewrite Equation 23 as

$$(24) \quad \begin{aligned} [D_{\pi'}(\mathcal{P}_{d'}^\bullet; \mathcal{Q}_{d'}^\bullet)] &= (-1)^{\ell(w)} \mathfrak{S}_{w^{-1}}(y_1, \dots, y_g, 0, \dots, 0; 0, \dots, 0) \\ &= \mathfrak{S}_{w^{-1}}(-y_1, \dots, -y_g, 0, \dots, 0; 0, \dots, 0). \end{aligned}$$

When all of the second set of variables in a double Schubert polynomial are set to 0, one obtains an ordinary Schubert polynomial. So we may write this class more simply as $\mathfrak{S}_{w^{-1}}(-y_1, \dots, -y_g)$. To finish our computation, we need the following fact about Schubert polynomials.

Lemma 6.12. *Let w be a permutation of $[1, n]$, and g be an integer such that $w(1) < \dots < w(g)$ and $g \leq \ell(w)$. Let e_k denote the elementary symmetric polynomial of degree k in g variables, and let $\alpha_1, \dots, \alpha_g, \Theta$ be elements of a ring such that $e_k(\alpha_1, \dots, \alpha_g) = \frac{\Theta^k}{k!}$. Then the Schubert polynomial \mathfrak{S}_w satisfies $\mathfrak{S}_w(\alpha_1, \dots, \alpha_g, 0, \dots, 0) = \frac{\Theta^{\ell(w)}}{\ell(w)!} |R(w)|$, where $R(w)$ denotes the set of reduced words for w .*

Proof. We use a theorem of Billey, Jockusch, and Stanley. Let $\ell = \ell(w)$, denote by s_a the transposition of a and $a + 1$, and identify the elements $a \in R(w)$ by ℓ -tuples $a = (a_1, a_2, \dots, a_\ell)$

where $w = s_{a_1} s_{a_2} \cdots s_{a_\ell}$. For each $a \in R(w)$, define the set $K(a)$ to be the set of all sequences $i = (i_1, \dots, i_\ell)$ of positive integers such that

$$(25) \quad i_1 \leq i_2 \leq \dots \leq i_\ell,$$

$$(26) \quad i_j \leq a_j \text{ for } 1 \leq j \leq \ell, \text{ and,}$$

$$(27) \quad i_j < i_{j+1} \text{ if } a_j < a_{j+1}.$$

With these definitions, [BJS93, Theorem 1.1] states that $\mathfrak{S}_w = \sum_{a \in R(w)} \sum_{i \in K(a)} x_{i_1} \cdots x_{i_\ell}$. Split \mathfrak{S}_w into a sum $\mathfrak{S}_{w,1} + \mathfrak{S}_{w,2} + \mathfrak{S}_{w,3}$ by partitioning $K(a)$ into three sets as follows. Let $K_1(a)$ consist of those sequences $i \in K(a)$ such that $i_1 < \dots < i_\ell \leq g$. Let $K_2(a)$ consist of those sequences $i \in K(a)$ with $i_\ell \leq g$ but with at least one index repeated. Let $K_3(a)$ consist of all $i \in K(a)$ such that $i_\ell > g$. The assumption that w is increasing on $1, 2, \dots, g$ implies that \mathfrak{S}_w is symmetric in the variables x_1, \dots, x_g (see e.g. [Mac91, 4.3(iii)] or [Ful92, Corollary 2.11]). The definitions of the three summands $\mathfrak{S}_{w,1}, \mathfrak{S}_{w,2}, \mathfrak{S}_{w,3}$ imply that each one is symmetric in the variables x_1, \dots, x_g as well; in particular, $\mathfrak{S}_{w,1}$ is an integer multiple of the elementary symmetric function e_ℓ and $\mathfrak{S}_{w,2}$ is a linear combination of monomial symmetric polynomials with at least one exponent greater than 1.

Clearly $\mathfrak{S}_{w,3}(\alpha_1, \dots, \alpha_g, 0, \dots, 0) = 0$. The expression $\mathfrak{S}_{w,2}(\alpha_1, \dots, \alpha_g, 0, \dots, 0)$ may be computed using the *exponential substitution*, discussed in [Sta99, §7.8]. The fact that $e_k(\alpha_1, \dots, \alpha_g) = \frac{\Theta^k}{k!}$ uniquely determines the value of all symmetric polynomials evaluated on $\alpha_1, \dots, \alpha_g$. By [Sta99, Proposition 7.8.4(b)], all monomial symmetric polynomials except the e_k evaluate to 0. Therefore $\mathfrak{S}_{w,2}(\alpha_1, \dots, \alpha_g, 0, \dots, 0) = 0$.

Finally, we claim that $\mathfrak{S}_{w,1} = |R(w)| \cdot e_\ell$. It suffices to show that for all $a \in R(w)$, $K(a)$ contains all ℓ -element increasing subsequences of $\{1, 2, \dots, g\}$. Fix a reduced word $a \in R(w)$. For all $j \in \{1, \dots, \ell\}$, define $w_j = s_{a_1} s_{a_2} \cdots s_{a_j}$. Let $x_j = w_j(a_j)$ and $y_j = w_j(a_j + 1)$. Since a is a reduced word, $x_j > y_j$, and x_j occurs before y_j in each of $w_j, w_{j+1}, \dots, w_\ell$. The position $w_k^{-1}(y_j)$ may increase by at most 1 at a time when k increases, so $w^{-1}(y_j) \leq w_j^{-1}(y_j) + (\ell - j) = a_j + 1 - \ell - j$. Since w is assumed to be increasing on $\{1, \dots, g\}$, it follows that

$$(28) \quad a_j \geq g + \ell + j.$$

Now, let $i = (i_1, \dots, i_\ell)$ be any sequence of positive integers such that $1 \leq i_1 < \dots < i_\ell \leq g$. Then i automatically satisfies conditions (25) and (26). Also, for all j the fact that i is strictly increasing implies that $i_j \leq g - \ell + j$. Equation (28) implies that $i_j \leq a_j$, so condition (26) holds as well. Therefore $K_1(a)$ includes all ℓ -element increasing subsequences of $\{1, \dots, g\}$. The reverse inclusion is clear from definitions; it follows that $\mathfrak{S}_{w,1} = |R(w)| \cdot e_\ell$ as claimed, and the lemma follows. \square

We can now prove the existence and intersection-theoretic part of Theorem 1.12.

Theorem 6.13. *In Situation 6.9, suppose that the degree- d' Brill-Noether flags of (C, p, q) are versal. Then in the Chow ring, $[W_d^\Pi(C, p, q)] = \frac{|R(\omega_{d-g}\pi)|}{\ell(\omega_{d-g}\pi)!} \Theta^{\ell(\omega_{d-g}\pi)}$.*

Proof. In the notation above, we have $\ell(\omega_{d-g}\pi) = \ell(w)$. Consider first the case $\ell(w) > g$. Then by Theorem 6.10, $W_d^\Pi(C, p, q)$ is empty, so the formula holds in this case. So assume $\ell(w) \leq g$.

Corollary 6.8 implies that π'^{-1} is decreasing on $[d' - (2g - 1), n - 1]$, so w^{-1} is increasing on $[1, g]$. So we have verified the hypotheses of Lemma 6.12, and it follows via Equation 22 that

$$[W_d^\Pi(C, p, q)] = \frac{\Theta^{\ell(w^{-1})}}{\ell(w^{-1})!} |R(w^{-1})|.$$

Since w and w^{-1} have the same length and there is a bijection between their sets of reduced words, we obtain the desired formula. \square

Proof of Theorem 1.12. Suppose that g, d are positive integers, k is an algebraically closed field, and Π is a dot array of size at least $d + 1 - g$. By Theorem 1.1, a general twice-marked smooth curve (C, p, q) has versal degree- $(d + 4g - 2)$ Brill-Noether flags, so assume that (C, p, q) is such a curve. First, consider the case where $\Pi \not\subseteq [0, d]^2$. Then either $r^\Pi(d + 1, 0) > 0$ or $r^\Pi(0, d + 1) > 0$, so $W_d^\Pi(C, p, q)$ is empty. But either the largest row a_r or largest column b_r in Π exceeds d , so $\rho_g(d, \Pi) < 0$ and the theorem statement is correct in this case.

Now assume that $\Pi \subseteq [0, d]^2$, so that we are in Situation 6.1. Let $M = N = 2g - 1$ in Situation 6.9. If $\rho_g(d, \Pi) < 0$ then $\ell(\omega_{d-g}\pi) > g$ and Theorem 6.13 implies that $W_d^\Pi(C, p, q)$ is empty, as desired. On the other hand, if $\rho_g(d, \Pi) \geq 0$ then Theorem 6.13 shows that $W_d^\Pi(C, p, q)$ is nonempty of the claimed class. Theorem 6.10 shows that $W_d^\Pi(C, p, q)$ has pure dimension $\rho_g(d, \Pi)$ and that $\widetilde{W}_d^\Pi(C, p, q)$ is smooth. Since the complement of $\widetilde{W}_d^\Pi(C, p, q)$ is a union of loci $\text{tw}_{-Mp-Nq}(D_\sigma(\mathcal{P}^\bullet, \mathcal{Q}^\bullet))$ of larger codimension, $\widetilde{W}_d^\Pi(C, p, q)$ is dense. \square

REFERENCES

- [ACGH85] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris, *Geometry of algebraic curves. Vol. I*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 267, Springer-Verlag, New York, 1985. MR 770932 21, 25
- [ACT20] Dave Anderson, Linda Chen, and Nicola Tarasca, *Motivic classes of degeneracy loci and pointed brill-noether varieties*, arXiv:2011.04814, 2020. 4
- [ACT21] Dave Anderson, Linda Chen, and Nicola Tarasca, *K-classes of Brill–Noether Loci and a Determinantal Formula*, International Mathematics Research Notices (2021), rnab025. 4, 5, 7
- [BJS93] Sara C. Billey, William Jockusch, and Richard P. Stanley, *Some combinatorial properties of Schubert polynomials*, J. Algebraic Combin. **2** (1993), no. 4, 345–374. MR 1241505 26
- [BW03] Sara C. Billey and Gregory S. Warrington, *Maximal singular loci of Schubert varieties in $\text{SL}(n)/B$* , Trans. Amer. Math. Soc. **355** (2003), no. 10, 3915–3945. MR 1990570 15, 25
- [CDPR12] Filip Cools, Jan Draisma, Sam Payne, and Elina Robeva, *A tropical proof of the Brill-Noether theorem*, Adv. Math. **230** (2012), no. 2, 759–776. MR 2914965 2, 7
- [CHTiB92] Ciro Ciliberto, Joe Harris, and Montserrat Teixidor i Bigas, *On the endomorphisms of $\text{Jac}(W_d^1(C))$ when $\rho = 1$ and C has general moduli*, Classification of irregular varieties (Trento, 1990), Lecture Notes in Math., vol. 1515, Springer, Berlin, 1992, pp. 41–67. MR 1180337 3, 4
- [CLMPTiB18] Melody Chan, Alberto López Martín, Nathan Pflueger, and Montserrat Teixidor i Bigas, *Genera of Brill-Noether curves and staircase paths in Young tableaux*, Trans. Amer. Math. Soc. **370** (2018), no. 5, 3405–3439. MR 3766853 2
- [CLMTiB18] Abel Castorena, Alberto López Martín, and Montserrat Teixidor i Bigas, *Petri map for vector bundles near good bundles*, J. Pure Appl. Algebra **222** (2018), no. 7, 1692–1703. MR 3763277 2
- [CLRW20] Steven Creech, Yoav Len, Caelan Ritter, and Derek Wu, *Prym–Brill–Noether Loci of Special Curves*, International Mathematics Research Notices (2020), rnaa207. 2
- [COP19] Melody Chan, Brian Osserman, and Nathan Pflueger, *The Gieseker–Petri theorem and imposed ramification*, Bull. Lond. Math. Soc. **51** (2019), no. 6, 945–960. MR 4041002 4, 7, 8
- [Cor01] Aurélie Cortez, *Singularités génériques et quasi-résolutions des variétés de Schubert pour le groupe linéaire*, C. R. Acad. Sci. Paris Sér. I Math. **333** (2001), no. 6, 561–566. MR 1860930 15, 25
- [CP19] Melody Chan and Nathan Pflueger, *Relative richardson varieties*, arXiv:1909.12414, 2019. 1, 2, 7, 8, 14, 16, 25
- [CP21] Melody Chan and Nathan Pflueger, *Euler characteristics of Brill-Noether varieties*, Trans. Amer. Math. Soc. **374** (2021), no. 3, 1513–1533. MR 4216716 2
- [CTiB08] Abel Castorena and Montserrat Teixidor i Bigas, *Divisorial components of the Petri locus for pencils*, J. Pure Appl. Algebra **212** (2008), no. 6, 1500–1508. MR 2391662 2
- [EH83] D. Eisenbud and J. Harris, *A simpler proof of the Gieseker–Petri theorem on special divisors*, Invent. Math. **74** (1983), no. 2, 269–280. MR 723217 2, 9, 10, 11
- [EH86] David Eisenbud and Joe Harris, *Limit linear series: basic theory*, Invent. Math. **85** (1986), no. 2, 337–371. MR 846932 4
- [FL81] W. Fulton and R. Lazarsfeld, *On the connectedness of degeneracy loci and special divisors*, Acta Math. **146** (1981), no. 3-4, 271–283. MR 611386 7

- [FP98] William Fulton and Piotr Pragacz, *Schubert varieties and degeneracy loci*, Lecture Notes in Mathematics, vol. 1689, Springer-Verlag, Berlin, 1998, Appendix J by the authors in collaboration with I. Ciocan-Fontanine. MR 1639468 5, 22, 25
- [FT16] Gavril Farkas and Nicola Tarasca, *Pointed Castelnuovo numbers*, Math. Res. Lett. **23** (2016), no. 2, 389–404. MR 3512891 4
- [Ful92] William Fulton, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), no. 3, 381–420. MR 1154177 15, 25, 26
- [HM06] Joe Harris and Ian Morrison, *Moduli of curves*, Springer Science & Business Media, 2006. 4
- [JP14] David Jensen and Sam Payne, *Tropical independence I: Shapes of divisors and a proof of the Gieseker-Petri theorem*, Algebra Number Theory **8** (2014), no. 9, 2043–2066. MR 3294386 2, 9
- [Kem71] George Kempf, *Schubert methods with an application to algebraic curves*, Stichting Mathematisch Centrum. Zuivere Wiskunde (1971), no. ZW 6/71. 4
- [KL72] Steven L. Kleiman and Dan Laksov, *On the existence of special divisors*, Amer. J. Math. **94** (1972), 431–436. MR 323792 4
- [KL74a] George Kempf and Dan Laksov, *The determinantal formula of schubert calculus*, Acta mathematica **132** (1974), 153–162. 4
- [KL74b] Steven L. Kleiman and Dan Laksov, *Another proof of the existence of special divisors*, Acta Math. **132** (1974), 163–176. MR 357398 4
- [KLR03] Christian Kassel, Alain Lascoux, and Christophe Reutenauer, *The singular locus of a Schubert variety*, J. Algebra **269** (2003), no. 1, 74–108. MR 2015302 15, 25
- [Li20] Shiyue Li, *Relative bott-samelson varieties*, arXiv:2011.04814, 2020. 7, 8
- [Liu02] Qing Liu, *Algebraic geometry and arithmetic curves*, Oxford Graduate Texts in Mathematics, vol. 6, Oxford University Press, Oxford, 2002, Translated from the French by Reinie Ern e, Oxford Science Publications. MR 1917232 13, 14
- [LS90] V. Lakshmibai and B. Sandhya, *Criterion for smoothness of Schubert varieties in $Sl(n)/B$* , Proc. Indian Acad. Sci. Math. Sci. **100** (1990), no. 1, 45–52. MR 1051089 15
- [Mac91] Ian Grant Macdonald, *Notes on schubert polynomials*, vol. 6, D ep. de math ematique et d’informatique, Universit e du Qu ebec  a Montr eal, 1991. 26
- [Man01] L. Manivel, *Le lieu singulier des vari et es de Schubert*, Internat. Math. Res. Notices (2001), no. 16, 849–871. MR 1853139 15, 25
- [Man20] Madhusudan Manjunath, *Poincar e series of divisors on graphs and chains of loops*, arXiv:2011.11910, 2020. 2
- [Oss14] Brian Osserman, *A simple characteristic-free proof of the Brill-Noether theorem*, Bull. Braz. Math. Soc. (N.S.) **45** (2014), no. 4, 807–818. MR 3296194 2, 4
- [Oss16] ———, *Dimension counts for limit linear series on curves not of compact type*, Math. Z. **284** (2016), no. 1-2, 69–93. MR 3545485 2
- [Oss19] ———, *Connectedness of Brill-Noether loci via degenerations*, Int. Math. Res. Not. IMRN (2019), no. 19, 6162–6178. MR 4016895 7
- [Pfl17] Nathan Pflueger, *Special divisors on marked chains of cycles*, J. Combin. Theory Ser. A **150** (2017), 182–207. MR 3645573 2, 7
- [Pfl18] ———, *On nonprimitive Weierstrass points*, Algebra Number Theory **12** (2018), no. 8, 1923–1947. MR 3892968 4
- [Sta99] Richard P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999, With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin. MR 1676282 26
- [TiB92] Montserrat Teixidor i Bigas, *On the Gieseker-Petri map for rank 2 vector bundles*, Manuscripta Math. **75** (1992), no. 4, 375–382. MR 1168175 2
- [TiB08] ———, *Petri map for rank two bundles with canonical determinant*, Compos. Math. **144** (2008), no. 3, 705–720. MR 2422346 2
- [TiB14] ———, *Injectivity of the Petri map for twisted Brill-Noether loci*, Manuscripta Math. **145** (2014), no. 3-4, 389–397. MR 3268854 2
- [Wel85] Gerald E. Welters, *A theorem of Gieseker-Petri type for Prym varieties*, Ann. Sci.  cole Norm. Sup. (4) **18** (1985), no. 4, 671–683. MR 839690 2, 9