

# Stochastic comparisons, differential entropy and varentropy for distributions induced by probability density functions\*

Antonio Di Crescenzo<sup>†</sup>    Luca Paolillo<sup>‡</sup>    Alfonso Suárez-Llorens<sup>§</sup>

## Abstract

Stimulated by the need of describing useful notions related to information measures, we introduce the ‘pdf-related distributions’. These are defined in terms of transformation of absolutely continuous random variables through their own probability density functions. We investigate their main characteristics, with reference to the general form of the distribution, the quantiles, and some related notions of reliability theory. This allows us to obtain a characterization of the uniform distribution based on pdf-related distributions of exponential and Laplace type as well. We also face the problem of stochastic comparing the pdf-related distributions by resorting to suitable stochastic orders. Finally, the given results are used to analyse properties and to compare some useful information measures, such as the differential entropy and the varentropy.

*Keywords:* Random lifetimes; Quantiles; Unimodal distributions; Stochastic orders.

*Mathematics Subject Classification:* 60E05, 60E15, 62E99, 94A17

## 1 Introduction

It is well known that various information measures for absolutely continuous random variables are given by expectations of functions of probability densities. Typical examples in this area involve the differential entropy and the related varentropy. These notions have been largely used in several contexts in order to describe the information content and the variability of stochastic systems. However, in some cases the results dealing with these measures are not easily manageable, and require non trivial efforts in order to appropriately compare the quantities under consideration. Stimulated by the need of dealing with effective probabilistic and statistical tools for assessing the above mentioned information measures, in this paper we introduce the so-called ‘pdf-related distributions’. These are constructed by means of transformation of absolutely continuous random variables through their own probability density functions (pdf’s).

---

\*Submitted for publication on March 11, 2021.

<sup>†</sup>Dipartimento di Matematica, Università degli Studi di Salerno, Via Giovanni Paolo II, 132, 84084 Fisciano (SA), Italy, Email: adicrescenzo@unisa.it – ORCID: 0000-0003-4751-7341

<sup>‡</sup>Dipartimento di Matematica, Università degli Studi di Salerno, Via Giovanni Paolo II, 132, 84084 Fisciano (SA), Italy, Email: lpaolillo@unisa.it – ORCID: 0000-0001-7146-4863

<sup>§</sup>Dpto. Estadística e Investigación Operativa, Universidad de Cádiz, Facultad de Ciencias, Campus Universitario, Río San Pedro s/n, 11510 Puerto Real, Cádiz, Spain, Email: alfonso.suarez@uca.es – ORCID: 0000-0002-7679-9328

We investigate the main properties of the pdf-related distributions, with special reference to (i) the general form of their distribution functions, (ii) the connections with various notions of interest in reliability theory (such as the residual lifetime), (iii) the determination of quantiles. We analyse pdf-related distributions generated by general distributions. However, special attention is devoted to the case when the underlying distributions possess monotone or unimodal pdf's. Here, unimodality refers to strictly decreasing pdf's or to pdf's that are first strictly increasing and then strictly decreasing.

The special form that is involved in their definition, allows to tackle the problem of comparing the pdf-related distributions by resorting to suitable stochastic orders, such as the usual stochastic order, the dispersive order, the convex transform order, the star order and the kurtosis order. Hence, in this framework the main results involve both location and variability orderings, and are obtained by techniques based on mapping of quantiles, convexity and rearrangement of pdf's.

As main application, the pdf-related distributions are finalized to construct a stochastic framework aimed to compare the basic information measures for absolutely continuous random variables. This allows to come to new results on the differential entropy, on the varentropy, and on the varentropy of residual lifetimes.

This is the plan of the paper. In Section 2 we recall some useful notions on random lifetimes and residual lifetimes, also with reference to the related information content, the differential entropy and the differential varentropy. Then, necessary results on quantiles for unimodal lifetime distributions and residual lifetime distributions are provided. In Section 3 we introduce the notion of distributions induced by probability density functions, namely pdf-related distributions. Under suitable monotonicity or unimodality assumptions we investigate their distributions and the related residual lifetime distributions, expressed in terms of quantiles. We also obtain a characterization of the  $(0,1)$ -uniform distribution based on pdf-related distributions of exponential and Laplace type. Section 4 focus on results based on stochastic orders, mainly finalized to perform stochastic comparisons between pdf-related distributions, also with reference to rearrangements of distributions. In Section 5 further stochastic comparisons are performed for the pdf-related distributions, and are also finalized to compare the differential entropy and varentropy of random lifetimes and of residual lifetimes.

Throughout the paper we denote by  $[X|B]$  a random variable having the same distribution of  $X$  conditional on  $B$ . Moreover, we write  $Y =_{st} X$  when  $X$  and  $Y$  are identically distributed. Furthermore,  $\phi'(x)$  denotes the derivative of any differentiable function  $\phi(x)$ .

## 2 Background

Let  $X$  be an absolutely continuous random variable defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , and let the interval  $S_X \subseteq \mathbb{R}$  be its support. Let  $F(x) = \mathbb{P}(X \leq x)$ ,  $x \in \mathbb{R}$ , be the cumulative distribution function (cdf), and  $\bar{F}(x) = 1 - F(x)$  the complementary distribution function, also known as survival function. We assume that the pdf  $f(x)$  of  $X$  satisfies one of the following assumptions:

- (i)  $f(x)$  is strictly decreasing for all  $x \in S_X$ , with  $S_X = (m, b)$ , with  $m < b \leq \infty$ ,
- (ii)  $f(x)$  is strictly increasing for  $x \leq m$  and strictly decreasing for  $x \geq m$ , with  $m \in S_X$ .

Henceforth,  $X$  is unimodal with mode  $m \in S_X$ .

If  $X$  is a nonnegative random lifetime, with support  $S_X = (0, b)$ , for  $0 < b \leq \infty$ , then

$$X_t := [X - t | X > t], \quad t \in S_X \tag{1}$$

describes the residual lifetime of  $X$  at age  $t$ , this being of interest in reliability or lifetime analysis. Then, the cdf, survival function and pdf of  $X_t$ , for  $x \in (0, b - t)$  and  $t \in S_X$  are given respectively by:

$$F_t(x) = \mathbb{P}(X_t \leq x) = \frac{F(x+t) - F(t)}{\bar{F}(t)},$$

$$\bar{F}_t(x) = 1 - F_t(x) = \frac{\bar{F}(x+t)}{\bar{F}(t)}, \quad (2)$$

$$f_t(x) = \frac{dF_t(x)}{dx} = \frac{f(x+t)}{\bar{F}(t)}. \quad (3)$$

The conventional approach used to characterize the failure distribution of  $X$  is either by its (instantaneous) hazard rate function

$$\lambda(t) = \frac{f(t)}{\bar{F}(t)} = \lim_{h \rightarrow 0^+} \frac{1}{h} \mathbb{P}[X \leq t+h | X > t], \quad t \in S_X, \quad (4)$$

or by its mean residual lifetime function, defined as

$$m(t) = \mathbb{E}(X_t) = \mathbb{E}[X - t | X > t] = \frac{1}{\bar{F}(t)} \int_t^\infty \bar{F}(x) dx, \quad t \in S_X.$$

We recall also the cumulative hazard rate function of  $X$ ,

$$\Lambda(t) = -\log \bar{F}(t) = \int_0^t \lambda(x) dx, \quad t \in S_X, \quad (5)$$

which plays a relevant role in numerous contexts.

For an absolutely continuous random variable  $X$  with pdf  $f(t)$ , the *information content* of  $X$  is defined as

$$IC(X) := -\log f(X). \quad (6)$$

This is a random variable that plays a relevant role in Information Theory, since it describes the amount of uncertainty contained in the outcome of  $X$ . Indeed, it is well-known that the *differential entropy* of  $X$  is given by the expectation of  $IC(X)$ , i.e. (cf. Cover and Thomas [4])

$$H(X) := \mathbb{E}[IC(X)] = -\mathbb{E}[\log f(X)] = -\int_{-\infty}^{\infty} f(x) \log f(x) dx. \quad (7)$$

In this framework, a relevant notion that measures the variability of the information content is the *differential varentropy* of  $X$ , given by (cf. Fradelizi *et al.* [11])

$$\begin{aligned} V(X) &:= \text{Var}[IC(X)] = \text{Var}[-\log f(X)] = \mathbb{E}[(IC(X))^2] - [H(X)]^2 \\ &= \int_{-\infty}^{\infty} f(x) [\log f(x)]^2 dx - \left[ \int_{-\infty}^{\infty} f(x) \log f(x) dx \right]^2. \end{aligned} \quad (8)$$

It is well known that the residual entropy is given by (cf. Ebrahimi [7] and Ebrahimi and Kirmani [8])

$$H(X_t) = \mathbb{E}[IC(X_t)] = -\int_t^\infty \frac{f(x)}{\bar{F}(t)} \log \frac{f(x)}{\bar{F}(t)} dx, \quad t \in S_X. \quad (9)$$

Equivalently, the following alternative expressions hold:

$$H(X_t) = -\Lambda(t) - \frac{1}{\bar{F}(t)} \int_t^\infty f(x) \log f(x) dx, \quad (10a)$$

$$H(X_t) = 1 - \frac{1}{\bar{F}(t)} \int_t^\infty f(x) \log \lambda(x) dx, \quad (10b)$$

for  $t \in D$ . The residual varentropy is (cf. Di Crescenzo and Paolillo [6])

$$\begin{aligned} V(X_t) &= \text{Var}[IC(X_t)] = \int_t^\infty \frac{f(x)}{\bar{F}(t)} \left( \log \frac{f(x)}{\bar{F}(t)} \right)^2 dx - [H(X_t)]^2 \\ &= \frac{1}{\bar{F}(t)} \int_t^\infty f(x) [\log f(x)]^2 dx - [\Lambda(t) + H(X_t)]^2, \end{aligned} \quad (11)$$

where  $\Lambda(t)$  can be found in (5), and  $H(X_t)$  is provided in (10a) and (10b). See, also, Goodarzi *et al.* [12] and [13] for various bounds to the variance of measures in dynamic settings.

## 2.1 Quantiles for unimodal distributions

In the following, for all  $u \in (0, 1)$  we shall denote by  $x_u$  the lower quantile of a random variable  $X$ , such that  $F(x_u) = u$ , i.e.

$$x_u = F^{-1}(u), \quad u \in (0, 1),$$

where

$$F^{-1}(\alpha) := \sup\{x \in S_X : F(x) \leq \alpha\}, \quad \alpha \in (0, 1) \quad (12)$$

denotes the right-continuous inverse of the distribution function  $F$ . In the following the function (12) will be said quantile function of  $X$ . Clearly,  $\bar{x}_u := x_{1-u}$  is the upper quantile, and  $x_{1/2}$  is the median of the distribution of  $X$ .

We recall that the distribution of  $X$  is symmetric if there exists a constant  $m$  such that  $F(m-x) = \bar{F}(m+x)$  for all  $x \in S_X$ , equivalently if  $F(x_u) = \bar{F}(\bar{x}_u)$  for all  $u \in (0, 1)$ . Moreover, if  $X$  is absolutely continuous, a sufficient condition for the symmetry is  $f(m-x) = f(m+x)$  for all  $x \in S_X$ , and an equivalent condition is (cf. Lemma 1 of Fashandi and Ahmadi [9])  $f(x_u) = f(\bar{x}_u)$ , for almost all  $u \in (0, 1)$ .

In the following, given a pdf  $f$  having support  $S_X$ , we define

$$\text{Im}^+(f) := \{y > 0 : \exists x \in S_X \text{ s.t. } f(x) = y\}.$$

**Definition 2.1** Let  $X$  be an absolutely continuous random variable having support  $S_X = (a, b)$  and a unimodal pdf  $f$  with mode  $m$ . Let  $y \in \text{Im}^+(f)$ .

- (i) If  $f(x)$  is continuous and strictly monotone on  $(a, b)$ , then the *inverse* of  $f$  at  $y$  is denoted by

$$l_y := f^{-1}(y). \quad (13)$$

- (ii) If  $f(x)$  is continuous in all the support  $S_X$  and is strictly increasing for  $x \leq m$  and strictly decreasing for  $x \geq m$ , then the *lower inverse* of  $f$  at  $y$  is given in (13), where in this case  $f^{-1}$  denotes the inverse of the restriction of  $f$  to  $S_X \cap (-\infty, m]$ . Moreover, the *upper inverse* of  $f$  at  $y$  is denoted by

$$u_y := f_2^{-1}(y) \quad (14)$$

where in this case  $f_2^{-1}$  denotes the inverse of the restriction of  $f$  to  $S_X \cap [m, \infty)$ .

**Remark 2.1** *If, in the case (ii) of Definition 2.1,  $f$  is symmetric in  $m$  then*

$$u_y = 2m - f^{-1}(y) = 2m - l_y.$$

**Remark 2.2** *Under the assumptions specified in the case (ii) of Definition 2.1 we have*

$$f(u_y) = f(l_y) \quad \text{and} \quad F(l_y) = \bar{F}(u_y).$$

## 2.2 Unimodal residual lifetime distributions and quantiles

In this section we generalize the above concepts to the case of residual lifetimes.

We first consider some properties of the residual lifetime when the underlying pdf is unimodal.

**Proposition 2.1** *Let  $X$  be an absolutely continuous random variable with support  $S_X = (0, b)$ , for  $0 < b \leq \infty$ , and having a unimodal pdf  $f(x)$  with mode  $m$ . Let  $X_t$  be defined as in (1), for  $t \in S_X$ , and with pdf (3).*

(i) *If  $t < m$ , then  $f_t(x)$  is unimodal with mode  $m_t = m - t$ .*

(ii) *If  $t \geq m$ , then  $f_t(x)$  is decreasing for all  $x \in (0, b - t)$ .*

The proof is straightforward and thus is omitted.

We can now recover the expression of the quantile function of the residual lifetime.

**Proposition 2.2** *If  $X$  is an absolutely continuous random variable having distribution function  $F(x)$  and support  $S_X = (0, b)$ , for  $0 < b \leq \infty$ , then the quantile function associated with the residual lifetime  $X_t$ , for  $t \in S_X$  can be expressed as*

$$F_t^{-1}(p) = F^{-1}(1 - (1 - p)\bar{F}(t)) - t, \quad p \in (0, 1).$$

**Proof.** From Equation (2), imposing  $F_t(x) = p \in (0, 1)$  we have  $\bar{F}(x + t) = (1 - p)\bar{F}(t)$ , for  $x \in (0, b - t)$ . Hence, one has  $F(x + t) = 1 - \bar{F}(x + t) = 1 - (1 - p)\bar{F}(t)$ , which gives the thesis.

If the pdf  $f(x)$  is symmetrical and unimodal, then the median and the mode of  $X$  coincide. However, even in this case the median and the mode of  $X_t$  are not generally coincident.

## 3 Pdf-related distributions

Let us now define a special type of distribution, named “pdf-related distribution”. This is defined by means of a transformation expressed by the pdf of a given baseline random variable.

**Definition 3.1** Let  $X$  be an absolutely continuous random variable with support  $S_X = (a, b)$ , and having pdf  $f(x)$ . Then,  $f(X)$  is named as the pdf-related random variable of  $X$ . For any  $y \in \text{Im}^+(f)$  the distribution function of  $f(X)$  is defined as

$$K(y) := \mathbb{P}(f(X) \leq y). \tag{15}$$

In the following proposition, which refers to the distribution function of pdf-related distributions, we use the notation adopted in Eqs. (13) and (14).

**Proposition 3.1** *Let  $X$  be an absolutely continuous random variable with support  $S_X = (a, b)$ , and having pdf  $f(x)$  that is continuous in all the support  $S_X$ .*

(a) *If  $f(x)$  is strictly decreasing, then*

$$K(y) = \bar{F}(l_y) = \bar{F}(f^{-1}(y)), \quad y \in \text{Im}^+(f).$$

(b) *If  $f(x)$  is strictly increasing, then*

$$K(y) = F(l_y) = F(f^{-1}(y)), \quad y \in \text{Im}^+(f).$$

(c) *If  $f(x)$  is strictly increasing for  $x \leq m$  and strictly decreasing for  $x \geq m$ , then*

$$K(y) = F(l_y) + \bar{F}(u_y) = F(f^{-1}(y)) + \bar{F}(f_2^{-1}(y)), \quad y \in \text{Im}^+(f).$$

The proof easily follows from Definition 2.1 and Eq. (15). Clearly, in the case (c) of Proposition 3.1 if the pdf  $f(x)$  is symmetric then

$$K(y) = 2F(l_y) = 2\bar{F}(u_y), \quad y \in \text{Im}^+(f).$$

It is worth pointing out that a pdf-related distribution is not necessarily continuous. Indeed, for instance, if  $X$  is uniformly distributed on  $(a, b)$ , then clearly  $f(X)$  is degenerate in  $(b-a)^{-1}$ .

Hereafter we obtain a characterization of the  $(0, 1)$ -uniform distribution based on pdf-related distributions of exponential and Laplace type. The proof follows easily from Proposition 3.1 and thus is omitted.

**Proposition 3.2** *Under the assumptions of Proposition 3.1, we have that the pdf-related random variable  $f(X)$  is uniform on  $(0, 1)$  if and only if*

(a)  *$f(x) = e^{a-x}$  for  $x \in (a, \infty)$  and  $f(x) = 0$  otherwise, when  $f$  is strictly decreasing in  $(a, \infty)$ , for  $a \in \mathbb{R}$ ;*

(b)  *$f(x) = e^{x-b}$  for  $x \in (-\infty, b)$  and  $f(x) = 0$  otherwise, when  $f$  is strictly increasing in  $(-\infty, b)$ , for  $b \in \mathbb{R}$ ;*

(c)  *$f(x) = e^{2(x-m)}$  for  $x \in (-\infty, m]$  and  $f(x) = e^{2(m-x)}$  for  $x \in [m, \infty)$ , when  $f$  is strictly increasing in  $(-\infty, m]$  and is strictly decreasing in  $[m, \infty)$ , for  $m \in \mathbb{R}$ .*

In the following, when the pdf  $f$  has support  $S_X = (a, b)$ , we adopt the following notation:  $f(a) = \lim_{x \rightarrow a^+} f(x)$  and  $f(b) = \lim_{x \rightarrow b^-} f(x)$ .

Hereafter we determine the cdf of the pdf-related random variable of the residual lifetime defined in (1), denoted as

$$K_t(y) := \mathbb{P}(f_t(X_t) \leq y) = \mathbb{P}(f(X) \leq y\bar{F}(t) \mid X > t), \quad (16)$$

with the last identity due to (3).

**Proposition 3.3** *Let  $X$  be an absolutely continuous random variable defined on a support  $S_X = (0, b)$ .*

(a) *If the pdf  $f$  is continuous and strictly decreasing in  $(t_0, b)$  for a given  $t_0 \in (0, b)$ , then for all  $t \in (t_0, b)$  one has*

$$K_t(y) = \frac{\bar{F}(f^{-1}(y\bar{F}(t)))}{\bar{F}(t)}, \quad y \in \text{Im}^+(f_t) = \left( \frac{f(b)}{\bar{F}(t)}, \lambda(t) \right), \quad (17)$$

where  $f^{-1}$  denotes the inverse of the restriction to  $(t_0, b)$  of  $f$ .

(b) If the pdf  $f$  is continuous and strictly increasing in  $(t_0, b)$  for a given  $t_0 \in (0, b)$ , then for all  $t \in (t_0, b)$  one has

$$K_t(y) = 1 - \frac{\bar{F}(f^{-1}(y\bar{F}(t)))}{\bar{F}(t)}, \quad y \in \text{Im}^+(f_t) = \left( \lambda(t), \frac{f(b)}{\bar{F}(t)} \right), \quad (18)$$

where  $f^{-1}$  denotes the inverse of the restriction to  $(t_0, b)$  of  $f$ .

(c) Let the pdf  $f$  be continuous in  $S_X$ , unimodal and symmetric, strictly increasing for  $x \in (0, m]$  and strictly decreasing for  $x \in [m, b)$ , with  $m = b/2$ . Then, for all  $t \in (0, m]$  one has

$$K_t(y) = \begin{cases} \frac{F(f^{-1}(y\bar{F}(t)))}{\bar{F}(t)}, & y \in \left( \frac{f(0)}{\bar{F}(t)}, \lambda(t) \right) \\ \frac{2F(f^{-1}(y\bar{F}(t))) - F(t)}{\bar{F}(t)}, & y \in \left[ \lambda(t), \frac{f(m)}{\bar{F}(t)} \right) \end{cases} \quad (19)$$

where  $f^{-1}$  is the inverse of the restriction to  $(0, m]$  of  $f$ .

**Proof.** In the case (a), from (16) for all  $t \in (t_0, b)$  we have

$$K_t(y) = \mathbb{P}(X \geq f^{-1}(y\bar{F}(t)) | X > t), \quad y \in \text{Im}^+(f_t).$$

Eq. (17) then follows from the assumption that  $f$  is strictly decreasing in  $(t_0, b)$ . In the case (b), from (16) one similarly has, for all  $t \in (t_0, b)$ ,

$$K_t(y) = \frac{\mathbb{P}(t < X \leq f^{-1}(y\bar{F}(t)))}{\bar{F}(t)}, \quad y \in \text{Im}^+(f_t),$$

this giving Eq. (18) due to the assumption that  $f$  is strictly increasing in  $(t_0, b)$ . In the case (c), when  $y \in \left( \frac{f(0)}{\bar{F}(t)}, \lambda(t) \right)$  the function  $K_t(y)$  can be obtained similarly as in case (a), whereas when  $y \in \left[ \lambda(t), \frac{f(m)}{\bar{F}(t)} \right)$  we have

$$K_t(y) = \mathbb{P}(t < X \leq f^{-1}(y\bar{F}(t)) | X > t) + \mathbb{P}(b - f^{-1}(y\bar{F}(t)) \leq X \leq b | X > t).$$

The expression given in (19) thus follows by noting that  $\bar{F}(b - x) = F(x)$  for all  $0 \leq x \leq b$ .

Clearly, the cdf given in (19) is continuous in  $y = \lambda(t)$ , with

$$K_t(\lambda(t)) = \frac{F(t)}{\bar{F}(t)}, \quad t \in (0, m],$$

where the right-hand-side is the odds function of  $X$  evaluated at  $t \leq m$ .

**Example 3.1** Let  $X$  be a Pareto-type random variable having pdf and cdf given respectively by

$$f(x) = \frac{1}{(1+x)^2}, \quad x \in (0, +\infty), \quad F(x) = \frac{x}{1+x}, \quad x \in [0, +\infty).$$

Since  $f$  is strictly decreasing for all  $x \in (0, +\infty)$ , with inverse  $f^{-1}(y) = y^{-1/2} - 1$ ,  $y \in (0, 1)$ , from (17) we have that, for all  $t > 0$

$$K_t(y) = (y(1+t))^{1/2}, \quad y \in \left( 0, \frac{1}{1+t} \right).$$




Figure 1: The cdf  $K_t(y)$  of Example 3.2 for  $t = 0.25$  (left) and for  $t = 0.5$  (right), with  $b = 0.25$  (dotted),  $b = 0.5$  (dashed) and  $b = 0.75$  (full).

**Example 3.2** Let  $X$  be a random variable having support  $S_X = (0, 2)$ , with pdf and cdf given respectively by

$$f(x) = \frac{b}{3} + \frac{1}{2} - b(x-1)^2, \quad F(x) = \left(\frac{b}{3} + \frac{1}{2}\right)x - \frac{b}{3}(x-1)^3 - \frac{b}{3}, \quad x \in S_X,$$

with  $b \in (0, \frac{3}{4}]$ . It satisfies the assumptions of Case (c) of Proposition 3.3, with mode  $m = 1$  and inverse

$$f^{-1}(y) = 1 - \sqrt{\frac{1}{b} \left( \frac{b}{3} + \frac{1}{2} - y \right)}, \quad y \in \left( \frac{1}{2} - \frac{2b}{3}, \frac{1}{2} + \frac{b}{3} \right).$$

The cdf  $K_t(y)$  can be easily obtained from (19). We omit the details for brevity, and show some instances in Figure 1.

A further case of interest is considered hereafter. For a given absolutely continuous random variable  $X$  with support  $S_X = (a, b)$ , and having pdf  $f$ , let us now analyze the survival function of  $f(t + X_t)$ . For all  $y \in \text{Im}^+(f)$  and  $t \in (a, b)$ , recalling Eq. (1) we have

$$\bar{G}_t(y) := \mathbb{P}(f(t + X_t) > y) = \mathbb{P}(f(X) > y \mid X > t). \quad (20)$$

**Proposition 3.4** Let  $X$  be an absolutely continuous random variable with support  $S_X = (a, b)$ , and let its pdf  $f$  be continuous and unimodal with mode  $m$ , such that  $f(x)$  is strictly increasing for  $x \leq m$  and strictly decreasing for  $x \geq m$ . Let  $y \in \text{Im}^+(f)$ ; denoting by  $F$  the cdf of  $X$  we have

(a) for  $t \in (a, m]$  and  $f(a) < f(b) < f(t)$

$$\bar{G}_t(y) = \begin{cases} 1 & \text{if } f(a) < y < f(b) \\ \frac{F(u_y) - F(t)}{\bar{F}(t)} & \text{if } f(b) < y \leq f(t) \\ \frac{F(u_y) - F(l_y)}{\bar{F}(t)} & \text{if } f(t) \leq y \leq f(m) \end{cases}$$



(b) for  $t \in (a, m]$  and  $f(a) < f(t) < f(b)$

$$\bar{G}_t(y) = \begin{cases} 1 & \text{if } f(a) < y \leq f(t) \\ \frac{1 - F(l_y)}{\bar{F}(t)} & \text{if } f(t) \leq y < f(b) \\ \frac{F(u_y) - F(t)}{\bar{F}(t)} & \text{if } f(b) < y \leq f(m); \end{cases}$$

(c) for  $t \in (a, m]$  and  $f(b) \leq f(a)$

$$\bar{G}_t(y) = \begin{cases} \frac{F(u_y) - F(t)}{\bar{F}(t)} & \text{if } f(b) < y < f(a) \text{ or } f(a) < y \leq f(t) \\ \frac{F(u_y) - F(l_y)}{\bar{F}(t)} & \text{if } f(t) \leq y \leq f(m); \end{cases}$$

(d) for  $t \in [m, b)$

$$\bar{G}_t(y) = \frac{F(u_y) - F(l_y)}{\bar{F}(t)} \quad \text{if } f(b) < y \leq f(t).$$

**Proof.** According to Eq. (20) we can consider the following cases.

(a) for  $t \in (a, m]$  and  $f(a) < f(b) < f(t)$ :

$$\begin{aligned} \bar{G}_t(y) &= \mathbb{P}(t \leq X < b | X > t) = 1, & y \in (f(a), f(b)), \\ \bar{G}_t(y) &= \mathbb{P}(t \leq X \leq u_y | X > t) = \frac{F(u_y) - F(t)}{\bar{F}(t)}, & y \in (f(b), f(t)], \\ \bar{G}_t(y) &= \mathbb{P}(l_y \leq X \leq u_y | X > t) = \frac{F(u_y) - F(l_y)}{\bar{F}(t)}, & y \in [f(t), f(m)]; \end{aligned}$$

(b) for  $t \in (a, m]$  and  $f(a) < f(t) < f(b)$ :

$$\begin{aligned} \bar{G}_t(y) &= \mathbb{P}(t \leq X < b | X > t) = 1, & y \in (f(a), f(b)), \\ \bar{G}_t(y) &= \mathbb{P}(l_y \leq X < b | X > t) = \frac{\bar{F}(l_y)}{\bar{F}(t)}, & y \in (f(b), f(t)], \\ \bar{G}_t(y) &= \mathbb{P}(l_y \leq X \leq u_y | X > t) = \frac{F(u_y) - F(l_y)}{\bar{F}(t)}, & y \in [f(t), f(m)]; \end{aligned}$$

(c) for  $t \in (a, m]$  and  $f(b) \leq f(a)$ :

$$\bar{G}_t(y) = \mathbb{P}(t \leq X \leq u_y | X > t) = \frac{F(u_y) - F(t)}{\bar{F}(t)}, \quad y \in (f(b), f(a)) \cup (f(a), f(t));$$

$$\bar{G}_t(y) = \mathbb{P}(l_y \leq X \leq u_y | X > t) = \frac{F(u_y) - F(l_y)}{\bar{F}(t)}, \quad y \in [f(t), f(m)];$$

(d) for  $t \in [m, b)$ :

$$\bar{G}_t(y) = \mathbb{P}(t \leq X \leq u_y | X > t) = \frac{F(u_y) - F(t)}{\bar{F}(t)}, \quad y \in (f(b), f(t)].$$

The proof is thus completed.

**Remark 3.1** With reference to Proposition 3.4, recalling Definition 2.1 we have

$$\frac{d}{dy}F(u_y) = \frac{y}{f'(u_y)}, \quad \frac{d}{dy}F(l_y) = \frac{y}{f'(l_y)},$$

so that we can obtain the pdf of  $f(t + X_t)$ , denoted by  $g_t(y) := -\frac{\partial \bar{G}_t(y)}{\partial y}$ , as follows:

- for  $t < m$  and  $f(a) \leq f(b) < f(t)$

$$g_t(y) = \begin{cases} -\frac{y}{\bar{F}(t) f'(u_y)} & \text{if } f(b) < y \leq f(t) \\ -\frac{y}{\bar{F}(t)} \left( \frac{1}{f'(u_y)} - \frac{1}{f'(l_y)} \right) & \text{if } f(t) < y \leq f(m), \end{cases} \quad (21)$$

- for  $t < m$  and  $f(a) < f(t) < f(b)$

$$g_t(y) = \begin{cases} \frac{y}{\bar{F}(t) f'(l_y)} & \text{if } f(b) < y \leq f(t) \\ -\frac{y}{\bar{F}(t)} \left( \frac{1}{f'(u_y)} - \frac{1}{f'(l_y)} \right) & \text{if } f(t) \leq y \leq f(m), \end{cases} \quad (22)$$

- for  $t < m$  and  $f(b) \leq f(a)$

$$g_t(y) = \begin{cases} -\frac{y}{\bar{F}(t) f'(u_y)} & \text{if } f(b) < y < f(a) \vee f(a) < y \leq f(t) \\ -\frac{y}{\bar{F}(t)} \left( \frac{1}{f'(u_y)} - \frac{1}{f'(l_y)} \right) & \text{if } f(t) \leq y \leq f(m), \end{cases} \quad (23)$$

- for  $t \geq m$

$$g_t(y) = -\frac{y}{\bar{F}(t) f'(u_y)} \quad \text{if } f(b) < y \leq f(t).$$

We note that  $f'(u_y) < 0 < f'(l_y)$ . Moreover, if  $X$  has a symmetric pdf, i.e.  $f(u_y) = f(l_y)$  then  $f'(u_y) = -f'(l_y)$ , so that the last expressions in the right-hand-sides of Eqs. (21)-(23) are simplified.

Hereafter we discuss the expression of the survival function (20) and the corresponding pdf as a function of  $t$ .

**Remark 3.2** If  $f(a) < f(b)$  then we have the following cases:

(a) for  $y \in (f(a), f(b))$

$$\bar{G}_t(y) = \frac{\bar{F}(l_y)}{\bar{F}(t)}, \quad g_t(y) = \frac{y}{f'(l_y)\bar{F}(t)}, \quad a < t \leq l_y$$

(b) for  $y \in (f(b), f(m))$

$$\bar{G}_t(y) = \begin{cases} \frac{F(u_y) - F(l_y)}{\bar{F}(t)}, & a < t \leq l_y \\ \frac{F(u_y) - F(t)}{\bar{F}(t)}, & l_y \leq t \leq u_y \end{cases} \quad g_t(y) = \begin{cases} -\frac{y}{\bar{F}(t)} \left( \frac{1}{f'(u_y)} - \frac{1}{f'(l_y)} \right), & a < t \leq l_y \\ -\frac{y}{f'(l_y)\bar{F}(t)}, & l_y \leq t \leq u_y. \end{cases}$$

If  $f(b) \leq f(a)$  then we have:

(a) when  $f(b) < f(a)$ , for  $y \in (f(b), f(a))$

$$\bar{G}_t(y) = \frac{F(u_y) - F(t)}{\bar{F}(t)}, \quad g_t(y) := -\frac{y}{f'(u_y)\bar{F}(t)}, \quad a < t \leq u_y$$

(b) for  $y \in (f(a), f(m)]$

$$\bar{G}_t(y) = \begin{cases} \frac{F(u_y) - F(l_y)}{\bar{F}(t)}, & a < t \leq l_y \\ \frac{F(u_y) - F(t)}{\bar{F}(t)}, & l_y \leq t \leq u_y \end{cases} \quad g_t(y) = \begin{cases} -\frac{y}{\bar{F}(t)} \left( \frac{1}{f'(u_y)} - \frac{1}{f'(l_y)} \right), & a < t \leq l_y \\ -\frac{y}{f'(u_y)\bar{F}(t)}, & l_y \leq t \leq u_y. \end{cases}$$

As in Remark 3.1, some expressions are simplified in the symmetric case for  $f'(u_y) = -f'(l_y)$ .

### 3.1 Determination of quantiles

Generally the determination of quantiles for a pdf-related distribution is not an easy task. Nevertheless, in some cases this problem can be solved, as reported hereafter.

**Proposition 3.5** *Let  $X$  be an absolutely continuous random variable with support  $S_X = (0, b)$  and having a symmetric and unimodal pdf  $f(x)$ . Then the lower quantile and the upper quantile of  $f(X)$  are given respectively by*

$$\xi_u = f \left( F^{-1} \left( \frac{u}{2} \right) \right), \quad \xi_{1-u} = f \left( F^{-1} \left( 1 - \frac{u}{2} \right) \right), \quad u \in (0, 1). \quad (24)$$

**Proof.** Recalling (15), from Proposition 3.1 we have

$$K(y) = F(l_y) + \bar{F}(u_y), \quad y \in \text{Im}^+(f).$$

For the symmetry, we can apply Remark 2.2 so that

$$K(y) = 2F(l_y) = 2\bar{F}(u_y).$$

Hence, applying Eqs. (13) and (14) we obtain the lower and upper quantiles of  $f(X)$  given in (24).

The following result refers to the cdf of the the pdf-related random variable of the residual lifetime (1), that is given in (16).

**Proposition 3.6** *Let  $X$  be an absolutely continuous random variable with support  $S_X = (0, b)$  and having a strictly monotone pdf  $f(x)$ .*

(a) *If  $f(x)$  is strictly decreasing, then for all  $t \in S_X$  and  $0 < p < 1$ ,*

$$K_t(y) = 1 - p, \quad y \in \text{Im}^+(f_t)$$

*if and only if*

$$y = f_t(F_t^{-1}(p)) = \frac{1}{\bar{F}(t)} f(F^{-1}(1 - (1 - p)\bar{F}(t))).$$

(b) *If  $f(x)$  is strictly increasing, then for all  $t \in S_X$  and  $0 < q < 1$ ,*

$$K_t(y) = q, \quad y \in \text{Im}^+(f_t)$$

*if and only if*

$$y = f_t(F_t^{-1}(q)) = \frac{1}{\bar{F}(t)} f(F^{-1}(1 - (1 - q)\bar{F}(t))).$$

**Proof.** After straightforward calculations, in both cases the result follows making use of Proposition 3.3 and Proposition 2.2.

With reference to the cdf (15), a straightforward consequence of Proposition 3.6 is given hereafter.

**Corollary 3.1** *Under the assumptions of Proposition 3.6,*

(a) *if  $f(x)$  is strictly decreasing, then for  $0 < p < 1$ ,*

$$K(y) = 1 - p \quad \text{if and only if} \quad y = f(F^{-1}(p));$$

(b) *if  $f(x)$  is strictly increasing, then for  $0 < q < 1$ ,*

$$K(y) = q \quad \text{if and only if} \quad y = f(F^{-1}(q)).$$

## 4 Results based on stochastic orders

In this section we provide some results involving stochastic orders and quantiles, for a comprehensive review on the background we refer to Shaked and Shantikumar [20]. We first recall notions that will be used in the following to establish an ordering between the varentropy of different random variables. Hereafter, the notation  $F^{-1}$  refers to the right-continuous inverse introduced in (12). Moreover,  $Me = x_{1/2}$  is the median of the distribution of  $X$ .

**Definition 4.1** *Let  $X$  and  $Y$  be random variables having cdfs  $F(x)$  and  $G(x)$ , respectively. Then,*

(i)  *$X$  is stochastically smaller than  $Y$ , and write  $X \leq_{st} Y$ , if  $F(x) \geq G(x)$  for all  $x \in \mathbb{R}$ , or equivalently if*

$$F^{-1}(u) \leq G^{-1}(u), \quad \text{for all } u \in (0, 1);$$

(ii)  $X$  is smaller than  $Y$  in the dispersive order, and write  $X \leq_{disp} Y$ , if

$$F^{-1}(\beta) - F^{-1}(\alpha) \leq G^{-1}(\beta) - G^{-1}(\alpha), \quad \text{for all } 0 < \alpha \leq \beta < 1.$$

(iii)  $X$  is smaller than  $Y$  in the convex transform order, with  $X$  and  $Y$  nonnegative, and write  $X \leq_c Y$ , if  $G^{-1}(F(x))$  is convex in  $x$  on the support of  $F(x)$ ;

(iv)  $X$  is smaller than  $Y$  in the star order, with  $X$  and  $Y$  nonnegative, and write  $X \leq_* Y$ , if  $G^{-1}$  is starshaped in  $x$ , or equivalently

$$\frac{G^{-1}(F(x))}{x} \quad \text{is increasing in } x \text{ on the support of } F(x);$$

(v)  $X$  is smaller than  $Y$  in kurtosis order, and write  $X \leq_k Y$ , if  $X$  and  $Y$  symmetric and  $G^{-1}(F(x))$  is concave for all  $x < Me$ , or, equivalently,  $G^{-1}(F(x))$  is convex for all  $x > Me$  (see, for example, Arriaza et al. [2]).

## 4.1 Mapping of quantiles

Given two random variables  $X$  and  $Y$  having cdfs  $F(x)$  and  $G(x)$ , respectively, let us define the mapping function

$$\phi(x) := G^{-1}(F(x)). \quad (25)$$

Clearly, this well-known function maps the quantiles of  $X$  to the quantile of  $Y$ , since  $Y =_{st} \phi(X)$ . The following result follows from Theorem 1.A.17 of [20].

**Remark 4.1** *The random variables  $X$  and  $Y$  satisfy  $X \leq_{st} Y$  if and only if, for all  $x$  in the support of  $X$ , we have*

$$\phi(x) \geq x.$$

The following result is related to the dispersive order, and involves the function (25).

**Remark 4.2** *For continuous random variables  $X$  and  $Y$  we have that  $X \leq_{disp} Y$  if and only if  $Y =_{st} \psi(X)$  for some function  $\psi$  which satisfies (cf. Section 3.B.1 of [20])*

$$\psi(x_2) - \psi(x_1) \geq x_2 - x_1 \quad \text{whenever } x_1 \leq x_2. \quad (26)$$

*It is easy to verify that if  $\psi(x)$  is equal to the function given in (25), then condition (26) is satisfied.*

## 4.2 Stochastic orders and convexity

Let us now recall a characterization of the convex transform order.

**Remark 4.3** *Let  $X$  and  $Y$  be nonnegative random variables with cdfs  $F(x)$  and  $G(x)$ , and pdfs  $f(x)$  and  $g(x)$ , respectively. Then, (cf. Section 4.B.2 of [20])*

$$X \leq_c Y \quad \iff \quad \frac{f(F^{-1}(p))}{g(G^{-1}(p))} \quad \text{is increasing for all } p \in (0, 1).$$

We also recall two results involving the star order.

**Remark 4.4** Let  $X$  and  $Y$  be nonnegative random variables with cdfs  $F(x)$  and  $G(x)$ , respectively. Then, (cf. Section 4.B.1 of [20])

$$X \leq_* Y \iff \frac{G^{-1}(p)}{F^{-1}(p)} \text{ is increasing in } p \in (0, 1).$$

**Remark 4.5** Let  $X$  and  $Y$  be nonnegative random variables. Then, (cf. Theorem 4.B.1 of [20])

$$X \leq_* Y \iff \log X \leq_{disp} \log Y.$$

We are now able to perform the comparison of pdf-related distributions as defined in Section 3 in terms of the star order.

**Theorem 4.1** Let  $X$  and  $Y$  be absolutely continuous random variables having pdfs  $f(x)$  and  $g(x)$ , respectively. Then,

$$f(X) \leq_* g(Y)$$

if and only if

$$\frac{\phi(x)}{x} \text{ is increasing in } x \text{ for all } x \text{ in the support of } f(X),$$

where

$$\phi(x) := S^{-1}(K(x)),$$

for

$$K(x) = \mathbb{P}(f(X) \leq x) \quad \text{and} \quad S(x) = \mathbb{P}(g(Y) \leq x). \quad (27)$$

**Proof.** The proof is a consequence of the point (iv) of Definition 4.1 applied to the random variables  $f(X)$  and  $g(Y)$ .

From the point (iv) of Definition 4.1 one has that both relations  $U \leq_* V$  and  $V \leq_* U$  hold simultaneously, and in the following we shall write  $U =_* V$ , if the random variables  $U$  and  $V$  have proportional quantile functions. In this case they belong to the same scale family of distributions (see, for instance, Section 4.1 of Di Crescenzo *et al.* [5]). Let us now investigate this relation for the pdf-related distributions connected by affine transformations.

**Theorem 4.2** Let  $X$  be an absolutely continuous random variable with pdf  $f(x)$ , and let  $Y = aX + b$ ,  $a > 0$ , have pdf  $g(x)$ . Then, one has

$$f(X) =_* g(Y). \quad (28)$$

**Proof.** For the cdfs (27), recalling that  $g(x) = \frac{1}{a} f\left(\frac{x-b}{a}\right)$ , for all  $x$  in the support of  $Y$  we have

$$S(x) = \mathbb{P}\left(\frac{1}{a} f\left(\frac{Y-b}{a}\right) \leq x\right) = K(ax).$$

Hence, one has

$$S^{-1}(p) = \frac{1}{a} K^{-1}(p), \quad \text{for all } p \in (0, 1),$$

so that

$$\frac{\phi(x)}{x} = \frac{S^{-1}(K(x))}{x} = \frac{1}{a} \quad \text{for all } x \text{ in the support of } f(X).$$

This implies relation (28) by Theorem 4.1.

The notion of rearrangement of a function has played a key role in many inequalities in literature. It is described in the seminal book of Hardy *et al.* [15] and has been studied in the context of entropy, randomness, majorization and dispersion in Hickey [16], [17] and Fernández-Ponce and Suárez-Llorens [10], among other authors. We next recall the decreasing rearrangement of a density function and some of its important properties.

**Definition 4.2** *Let  $X$  be an absolutely continuous random variable having a pdf  $f(x)$ . The decreasing rearrangement of  $f(x)$ , denoted by  $f^*(x)$ , is given by*

$$f^*(x) = \sup\{c : m(c) > x\}, \quad x > 0,$$

where  $m(c) = \mu\{t : f(t) > c\}$ ,  $\mu$  denoting the Lebesgue measure.

It is well known, see [14], that the decreasing rearrangement satisfies the following identity (see, Hardy *et al.* [14])

$$\int_0^\infty f^*(x) dx = \int_{-\infty}^\infty f(x) dx = 1.$$

Then  $f^*$  can be considered as a pdf of a particular random variable that we will denote by  $X^*$ .

The following implication is well-known (see Hardy *et al.* [14], Theorems 9 and 10). Let  $X$  and  $Y$  be absolutely continuous random variables having pdf  $f$  and  $g$ , respectively. Then

$$\int_0^t f^*(x) dx \geq \int_0^t g^*(x) dx, \quad \forall t > 0 \iff \int_{-\infty}^\infty u(f(x)) dx \leq \int_{-\infty}^\infty u(g(x)) dx, \quad (29)$$

for all continuous and concave functions  $u$ . From Definition 4.1, the expressions in (29) are also equivalent to  $X^* \leq_{st} Y^*$ . It is worth mentioning that  $X^* \leq_{st} Y^*$  is interpreted as continuous majorisation in Hickey [16].

It seems that  $X^* \leq_{st} Y^*$  implies a kind of stochastic comparison between  $f(X)$  and  $g(Y)$ . Rewriting the right side of expression (29), we obtain that  $X^* \leq_{st} Y^*$  is equivalent to

$$\mathbb{E} \left[ \frac{u(f(X))}{f(X)} \right] \leq \mathbb{E} \left[ \frac{u(g(Y))}{g(Y)} \right],$$

for all continuous and concave functions  $u$ . The latter relation provides a class of measures of entropy. Indeed, as it is interpreted in Hickey [16], just taking  $u(x) = -x \log(x)$ , one has that if  $X^* \leq_{st} Y^*$  then  $H(X) \leq H(Y)$ , due to (7).

Hence, it is worth mentioning that the rearrangement captures in some sense the degree of randomness in a density function. Now we wonder if the rearrangement is also related with the concept of varentropy. From now on, we will assume that the density functions have no flat zones or, equivalently, the rearrangements are strictly decreasing. Next we provide a result that we will use later on.

**Theorem 4.3** *Let  $X$  and  $Y$  be absolutely continuous random variables having pdf  $f$  and  $g$ , respectively, with no flat zones. Then*

$$X^* \leq_c Y^* \iff f(X) \leq_* g(Y).$$

**Proof.** From Eq. (2.1) of Hickey [16] we have that the distribution function of  $X^*$  can be expressed as

$$F_{X^*}(t) = \mathbb{P}(X^* \leq t) = \int_0^t f^*(x) dx = \int_{\{x \in \mathbb{R}: f(x) > f^*(t)\}} f(x) dx = \mathbb{P}(f(X) > f^*(t)).$$

The use of strict inequality in the set  $\{x \in \mathbb{R} : f(x) > f^*(x)\}$  follows from the assumption of  $f$  having no flat zones. Therefore, with obvious notation, one has

$$F_{f(X)}(f^*(t)) = 1 - F_{X^*}(t), \quad t \in \mathbb{R}.$$

Because  $f^*$  is strictly decreasing and thus invertible, we have

$$F_{f(X)}^{-1}(p) = f^*(F_{X^*}^{-1}(1-p)), \quad 0 < p < 1, \quad (30)$$

where  $F_{X^*}^{-1}$  denotes the right-continuous inverse of  $F_{X^*}$ . Then, recalling Remark 4.4 the stochastic relation

$$f(X) \leq_* g(Y)$$

is satisfied if and only if

$$\frac{F_{g(Y)}^{-1}(p)}{F_{f(X)}^{-1}(p)} = \frac{g^*(G_{Y^*}^{-1}(1-p))}{f^*(F_{X^*}^{-1}(1-p))}, \quad \text{is increasing in } p \in (0, 1).$$

Finally, this is equivalent to  $X^* \leq_c Y^*$  by virtue of Remark 4.3.

**Example 4.1** From expression (30) and taking in account the inverse probability integral transform, we obtain that  $f(X) =_{st} f^*(X^*)$ . Therefore, it is clear that random variables having the same rearrangements have also the same differential entropy and varentropy. Here we provide an illustrative example. Let  $X$  and  $Y$  be random variables taking values in  $[-1, 1]$ , with density functions  $f$  and  $g$  given by

$$f(x) = 1 - |x|, \quad -1 \leq x \leq 1, \quad g(x) = |x|, \quad -1 \leq x \leq 1,$$

respectively. It is easy to compute that

$$m_f(c) = m_g(c) = \mu \{x : g(x) > c\} = 2(1-c)I_{[0,1]}(c).$$

Then  $f^*(x) = g^*(x) = (1-x/2)I_{[0,2]}(x)$ . Therefore, we have that  $f^*(X^*) =_{st} g^*(Y^*)$  and thus we can conclude that  $H(X) = H(Y)$  and  $V(X) = V(Y)$ . It is worth mentioning that  $g$  is not unimodal.

**Remark 4.6** Let  $X$  and  $Y$  be nonnegative random variables having common support  $S = (a, b)$ , and having pdfs  $f(x)$  and  $g(x)$ , respectively. Then, as a direct consequence of Theorem 4.3 we have

$$X \leq_c Y \quad \iff \quad \begin{cases} f(X) \leq_* g(Y), & \text{if } f \text{ and } g \text{ are strictly decreasing} \\ f(X) \geq_* g(Y), & \text{if } f \text{ and } g \text{ are strictly increasing.} \end{cases}$$

The following remarks are straightforward and they will be useful later on.



**Remark 4.7** *If  $X$  has a symmetric and unimodal pdf  $f(x)$ , then  $X^* =_{st} 2|X - Me|$  where  $Me$  is the median of  $X$ .*

**Remark 4.8** *If  $X$  has a strictly decreasing pdf in the interval  $(0, +\infty)$ , then  $f^*(x) = f(x)$  for all  $x \in (0, +\infty)$ , i.e.,  $X^* =_{st} X$ .*

A possible comparison in terms of the kurtosis order is provided by Remark 4.7 through the following theorem (see Oja [18]).

**Theorem 4.4** *Let  $X$  and  $Y$  be symmetric unimodal random variables having median zero.  $X \leq_k Y$  if and only if  $|X| \leq_c |Y|$ .*

### 4.3 Pdf-related distributions and variability orders

Starting from the results illustrated in Section 4 we apply those concepts to investigate relations between pdf-related distributions.

**Theorem 4.5** *Let  $X$  and  $Y$  be random variables having symmetric and unimodal pdfs  $f$  and  $g$ , respectively. Then,  $X \leq_k Y$  if and only if  $f(X) \leq_* g(Y)$ .*

**Proof.** Since changes in location do not affect the kurtosis order we can assume that  $Me(X) = Me(Y) = 0$ . Therefore from Theorem 4.4 and using Remark 4.7 we obtain that  $X \leq_k Y$  if and only if  $X^* \leq_c Y^*$ . The result follows easily just applying Theorem 4.3.

## 5 Differential entropy, varentropy and stochastic orders

In this section we will use concepts related to stochastic orders to perform some comparisons for the differential entropy and the differential varentropy. We first show the effect of the usual stochastic ordering between pdf-related random variables on the differential entropy of the underlying random variables.

**Theorem 5.1** *Let  $X$  and  $Y$  be absolutely continuous random variables having pdfs  $f(x)$  and  $g(x)$ , respectively, and having finite differential entropy. If  $f(X) \leq_{st} g(Y)$ , then*

$$H(X) \geq H(Y).$$

**Proof.** From the hypothesis  $f(X) \leq_{st} g(Y)$  it follows that  $-\log f(X) \geq_{st} -\log g(Y)$ , so that  $\mathbb{E}[IC(X)] \geq \mathbb{E}[IC(Y)]$ . The thesis then follows from Eq. (7).

**Remark 5.1** *When  $X$  and  $Y$  have cdf's  $F$  and  $G$ , and pdf's  $f$  and  $g$ , respectively, then  $X \leq_{disp} Y$  if and only if  $f(F^{-1}(u)) \geq g(G^{-1}(u))$  for all  $u \in (0, 1)$ , see (3.B.11) in [20]. Just considering the inverse probability integral transformation and Theorem 1.A.1 in [20], if  $X \leq_{disp} Y$  holds then  $f(X) \geq_{st} g(Y)$  and using Theorem 5.1 we obtain that  $H(X) \leq H(Y)$ . On the other hand, it is well known that if  $X$  is IFR then  $X_{t_2} \leq_{disp} X_{t_1}$  for all  $t_1 \leq t_2$ , see Belzunce et al. [3] and Pellerey and Shaked [19]. From the previous arguments it is apparent that if  $X$  is IFR then  $H(X_t)$  is decreasing when  $t$  increases. This result is well known in the literature, see Ebrahimi [7]. A similar result holds for DFR distributions by exchanging all inequalities and replacing decreasing for increasing in the residual differential entropy.*

A similar result can be stated for the differential varentropy when the pdf-related random variables are compared in the star order.

**Theorem 5.2** *Let  $X$  and  $Y$  be absolutely continuous random variables having pdf  $f(x)$  and  $g(x)$ , respectively, and having finite differential varentropy. If  $f(X) \leq_* g(Y)$ , then*

$$V(X) \leq V(Y).$$

**Proof.** Recalling Remark 4.5 one has that the assumption  $f(X) \leq_* g(Y)$  is equivalent to  $\log f(X) \leq_{disp} \log g(Y)$ . The latter relation implies that  $\text{Var}[\log f(X)] \leq \text{Var}[\log g(Y)]$  (see, for instance, Section 3.B.2 of [20]). The thesis then follows from (8).

With reference to equation (6), we denote by

$$L(x) = \mathbb{P}(IC(X) \leq x) \quad \text{and} \quad M(x) = \mathbb{P}(IC(Y) \leq x) \quad (31)$$

the cdfs of the information contents  $IC(X) = -\log f(X)$  and  $IC(Y) = -\log g(Y)$ , respectively. Then, in the following proposition we show some stochastic comparisons between the information contents of  $X$  and  $Y$ , where a role is played by the mapping function related to these cdfs.

**Remark 5.2** *Let  $X$  and  $Y$  be absolutely continuous random variables having pdf  $f(x)$  and  $g(x)$ , respectively, and let  $\varphi(x) := M^{-1}(L(x))$  be the mapping function related to the cdfs (31).*

*(i) From Remark 4.1 we have that  $IC(X) \leq_{st} IC(Y)$  if and only if  $\varphi(x) \geq x$  for all  $x$  in the support of  $X$ . Moreover, due to Theorem 1.A.3(a) of [20] these conditions are equivalent to  $f(X) \geq_{st} g(Y)$ .*

*(ii) From Remark 4.2 one has  $IC(X) \leq_{disp} IC(Y)$  if and only if  $\varphi'(x) \geq 1$  for all  $x$  in the support of  $X$ .*

*(iii) Due to Theorem 3.4 of Alimohammadi et al. [1], if  $IC(X) \geq_{st} IC(Y)$  and  $IC(X) \leq_{disp} IC(Y)$ , then  $f(X) \leq_{disp} g(Y)$ .*

Hereafter we show further comparison results for the differential varentropy.

**Theorem 5.3** *Let  $X$  and  $Y$  be symmetric and unimodal random variables. If  $X \leq_k Y$  then*

$$V(X) \leq V(Y).$$

**Proof.** The result follows easily from Theorem 4.4 and 4.5.

**Remark 5.3** *Theorem 5.3 provides many possible comparisons. We find in Arriaza et al. [2] a compilation of many unimodal symmetric distributions that are ordered in the kurtosis sense. For example, we know that the classical normal, logistic and Cauchy distributions satisfy that  $\text{normal} \leq_k \text{logistic} \leq_k \text{Cauchy}$ . Therefore, just applying Theorem 5.3 we obtain that  $V(\text{normal}) \leq V(\text{logistic}) \leq V(\text{Cauchy})$ .*

**Proposition 5.1** *If  $X$  and  $Y$  are absolutely continuous random variables having strictly decreasing pdfs in the interval  $(a, +\infty)$  and if  $X \leq_c Y$ , then*

$$V(X) \leq V(Y).$$

**Proof.** Without lack of generality we assume that  $a = 0$ . By Definition 4.2 we have that  $X^*$  is identically distributed as  $X$ . Hence, by Theorem 4.3 the assumption  $X^* \leq_c Y^*$  is equivalent to  $f(X) \leq_* g(Y)$ , where  $f(x)$  and  $g(x)$  denote respectively the pdfs of  $X$  and  $Y$ . We thus obtain  $V(X) \leq V(Y)$  from Theorem 5.2.

**Theorem 5.4** *Let  $X$  be a nonnegative absolutely continuous random lifetime with support  $S_X$  having a cdf  $F$  and a pdf  $f(x)$ . Let  $X_t$  be its residual lifetime at time  $t$ , as defined in (1). Let us suppose that exists  $t_0 \in S_X$  such the pdf of  $X_t$ , defined in (3), is strictly decreasing  $\forall t \geq t_0$ . If the ratio*

$$\frac{f(F^{-1}(1 - (1 - p)u))}{f(F^{-1}(1 - (1 - p)v))}$$

*increases (decreases) in  $p \in (0, 1)$ ,  $\forall 0 < v < u < 1$ . Then  $V(X_t)$  increases (decreases) in  $t \geq t_0$ .*

**Proof.** Given  $t_1$  and  $t_2$  such that  $t_0 < t_1 < t_2$  and using Proposition 5.1 we just need to check that  $X_{t_1} \leq_c (\geq_c) X_{t_2}$ . From the characterization of the convex transfor order given in Remark 4.3, the expression of the PDF of the residual life given in (3) and the expression of the inverse of the distribution function of the residual life given in Proposition 2.2 we obtain that

$$X_{t_1} \leq_c (\geq_c) X_{t_2} \iff \frac{f(F^{-1}(1 - (1 - p)\bar{F}(t_1)))}{f(F^{-1}(1 - (1 - p)\bar{F}(t_2)))} \text{ is increasing (decreasing) in } p \in (0, 1).$$

The result follows easily from the hypothesis assumption just considering  $v = \bar{F}(t_2)$  and  $u = \bar{F}(t_1)$ .

**Example 5.1** *Let  $X \sim \text{Weibull}(k, \lambda)$  be a Weibull distribution with shape parameter  $k$  and scale parameter  $\lambda$ . From the expression of the cdf of  $X$  given by  $F(x) = 1 - \exp(-(x/\lambda)^k)$ , it is a straightforward matter to compute the ratio*

$$\frac{f(F^{-1}(1 - (1 - p)u))}{f(F^{-1}(1 - (1 - p)v))} = \frac{u}{v} \left( \frac{\ln((1 - p)u)}{\ln((1 - p)v)} \right)^{\frac{k-1}{k}}$$

*for all  $0 < v < u < 1$ . It follows easily that the above ratio is increasing (decreasing) when  $k > 1$  ( $k < 1$ ) and it is constant for  $k = 1$ . It is well-known that Weibull distributions are always unimodal having decreasing density functions after the mode. Then, the pdfs of the residual lives after the mode satisfy the conditions of Theorem 5.4, see Proposition 2.1. Then,  $V(X_t)$  increases (decreases)  $\forall t \geq \text{mode}$ , for  $k > 1$  ( $k < 1$ ). On the other hand, the Weibull distribution is IFR (DFR) for  $k > 1$  ( $k < 1$ ). Then using Remark 5.1 we obtain that the differential entropy and the varentropy of the residual lives have opposite directions of growth depending on the shape parameter  $k$ . Of course, the case  $k = 1$  is just the exponential distribution where the residual lives have both differential entropy and varentropy constant. Some examples of the varentropy of the residual lives in the Weibull distributions are shown computationally in [6].*

We recall that in Theorem 3.4 of [6] it has been proved that if a random lifetime  $X$  is ILR, i.e. its pdf is log-concave, then the residual varentropy (11) is such that  $V(X_t) \leq 1$  for all  $t$  in the support of  $X$ . The condition of log-concave density means that the pdf belongs to the class of strong unimodal densities. Hereafter we show a similar result making use of some of the above results. To this aim, recall that a random lifetime  $X$  is said IFR if its hazard rate function (4) is increasing.

**Theorem 5.5** *Let  $X$  be a nonnegative absolutely continuous random lifetime with support  $S_X$ , and let  $X_t$  be its residual lifetime at time  $t$  as defined in (1). If, for a given  $t \in S_X$ ,  $X_t$  is IFR and its pdf (3) is strictly decreasing, then*

$$V(X_t) \leq 1.$$

**Proof.** Since, for a given  $t \in S_X$ , the residual lifetime  $X_t$  is IFR, then for Theorem 4.B.11 of [20] one has that  $X_t \leq_c \text{Exp}$ , where  $\text{Exp}$  denotes an exponential random variable. Moreover, by assumption  $X_t$  has a strictly decreasing pdf  $f_t(x)$  for all  $x \geq 0$ , so that  $X_t \leq_c \text{Exp}$  if and only if  $X_t^* \leq_c \text{Exp}^*$  for all  $t \in S_X$ , where  $X_t^*$  and  $\text{Exp}^*$  are the rearrangement of  $X_t$  and  $\text{Exp}$ , respectively. Due to Theorem 4.3, the last stochastic inequality is satisfied if and only if

$$f_{X_t}(X_t) \leq_* g(\text{Exp}),$$

where  $g$  is the pdf of  $\text{Exp}$ . Hence, making use of Theorem 5.2 we find

$$V(X_t) \leq V(\text{Exp}).$$

Finally, the thesis follows recalling that  $V(\text{Exp}) = 1$ .

**Remark 5.4** *From Proposition 2.1, we have that the conditions of Theorem 5.5 easily hold for all unimodal IFR distributions when  $t \geq m$ , where  $m$  is the mode of  $X$ . For example, this is the case of the residual lives of the Weibull distribution for  $k > 1$ , as it is described in Example 5.1. After the mode the varentropy increases but it has an upper bound equal to 1.*

## Acknowledgements

The authors acknowledge support received from the Ministerio de Economía y Competitividad (Spain) under grant no. MTM2017-89577-P, by the 2014-2020 ERDF Operational Programme and by the Consejería de Economía, Conocimiento, Empresas y Universidad (Junta de Andalucía, Spain), under grant FEDER-UCA18-107519, and from the Italian MIUR-PRIN 2017, project ‘Stochastic Models for Complex Systems’, No. 2017JFFHSH.

A. Di Crescenzo and L. Paolillo are members of the research group GNCS of Istituto Nazionale di Alta Matematica (INdAM).

L. Paolillo expresses his warmest thanks to the Departamento de Estadística e Investigación Operativa of Universidad de Cádiz for the hospitality during the visit carried out in 2020.

## References

- [1] Alimohammadi M, Esna-Ashari M, Cramer E (2021) On dispersive and star orderings of random variables and order statistics. *Stat Prob Lett* 170:109014. <https://doi.org/10.1016/j.spl.2020.109014>
- [2] Arriaza A, Di Crescenzo A, Sordo M A, Suárez Llorens A (2019) Shape measures based on the convex transform order. *Metrika* 82:99-124. <https://doi.org/10.1007/s00184-018-0667-y>
- [3] Belzunce F, Candel J, Ruiz J M (1996), Dispersive Ordering and Characterizations of Aging Classes. *Stat Prob Lett* 28:321-327. [https://doi.org/10.1016/0167-7152\(95\)00141-7](https://doi.org/10.1016/0167-7152(95)00141-7)
- [4] Cover T M, Thomas J A (1991) *Elements of Information Theory*. J Wiley & Sons, New York.

- [5] Di Crescenzo A, Martinucci B, Mulero J (2016) A quantile-based probabilistic mean value theorem. *Prob Eng Inf Sci* 30:261-280. <https://doi.org/10.1017/S0269964815000376>
- [6] Di Crescenzo A, Paolillo L (2020) Analysis and applications of the residual varentropy of random lifetimes. *Prob Eng Inf Sci* (online first) <https://doi.org/10.1017/S0269964820000133>
- [7] Ebrahimi N (1996) How to measure uncertainty in the residual life time distribution. *Sankhyā. The Indian Journal of Statistics. Series A* 58:48-56.
- [8] Ebrahimi N., Kirmani S.N.U.A. (1996) Some results on ordering of survival functions through uncertainty. *Stat Prob Lett* 29:167-176. <https://doi.org/10.1007/BF00054789>
- [9] Fashandi M, Ahmadi J (2012) Characterizations of symmetric distributions based on Rényi entropy. *Stat Prob Lett* 82:798-804. <https://doi.org/10.1016/j.spl.2012.01.004>
- [10] Fernández-Ponce J M, Suárez-Llorens A (2003) An aging concept based on majorization. *Prob Eng Inf Sci* 17:107-117. <https://doi.org/10.1017/S0269964803171069>
- [11] Fradelizi M, Madiman M, Wang L (2016) Optimal concentration of information content for log-concave densities. In: Houdré C, Mason D, Reynaud-Bouret P, Rosiński J (eds.) *High Dimensional Probability VII. Progress in Probability*, Cham, Springer, 71:45-60.
- [12] Goodarzi F, Amini M, Mohtashami Borzadaran G R (2016) On upper bounds for the variance of functions of the inactivity time. *Stat Prob Lett* 117:62-71. <https://doi.org/10.21136/AM.2017.0182-16>
- [13] Goodarzi F, Amini M, Mohtashami Borzadaran G R (2017) Some results on upper bounds for the variance of functions of the residual life random variables. *J Comp Appl Math* 320:30-42. <https://doi.org/10.1016/j.cam.2017.01.001>
- [14] Hardy G H, Littlewood J E, Polya G (1929) Some simple inequalities satisfied by convex functions. *Messenger of Math* 58:145-152.
- [15] Hardy G H, Littlewood J E, Polya G (1934) *Inequalities*. Cambridge University Press, Cambridge.
- [16] Hickey R J (1984) Continuous majorisation and randomness. *J Appl Prob* 21:924-929.
- [17] Hickey R J (1986) Concepts of dispersion in distributions: a comparative note. *J Appl Prob* 23:914-921.
- [18] Oja H (1983) Descriptive statistics for multivariate distributions. *Stat Prob Lett* 1:327-332.
- [19] Pellerey F, Shaked M (1997) Characterizations of the IFR and DFR Aging Notions by Means of the Dispersive Order. *Stat Prob Lett* 33:389-393. [https://doi.org/10.1016/S0167-7152\(96\)00152-6](https://doi.org/10.1016/S0167-7152(96)00152-6)
- [20] Shaked M, Shantikumar J G (2007) *Stochastic Orders*. Springer.