

CURVES CONTRACTED BY THE GAUSS MAP

LEI SONG

ABSTRACT. Given a singular projective variety in some projective space, we characterize the smooth curves contracted by the Gauss map in terms of normal bundles. As a consequence, we show that if the variety is normal, then a contracted line always has local obstruction for the embedded deformation and each component of the Hilbert scheme where the line lies is non-reduced everywhere.

1. INTRODUCTION

Let $X \subseteq \mathbb{P}^N$ be a subvariety of dimension n . The Gauss map $\gamma : X \dashrightarrow \mathbb{G}(n, N)$ sends a smooth point $x \in X$ to the embedded tangent space $\widehat{T}_x X$ of X at x .

If X is non-degenerate and smooth, then the map is a finite morphism by F. Zak [Zak93]. For singular varieties, Griffiths-Harris [GH79] established a structure theorem for Gauss map. Specifically it states that if the dimension of the image $\dim(\text{Im}(\gamma))$ of γ is less than n (in this case, the Gauss map is called to be degenerate), then for a general point $[T] \in \text{Im}(\gamma)$, the closure of the fibre $\gamma^{-1}([T])$ is a projective space of dimension $n - \dim(\text{Im}(\gamma))$. Roughly speaking, a projective subvariety with the degenerate Gauss map can be built up by some basic blocks, such as cones and developables, see [GH79], [AG01] for more precise statements, [AG01] for a historical account, and [Ran19] for a recent development.

In general, it may happen that γ is generically finite but has some positive-dimensional fibres, i.e. it contacts some positive-dimensional subvarieties. Here for a given subvariety $V \subseteq X$, we say V is *contracted* by the Gauss map γ or say γ is *constant* on V if V is not contained in the singular locus X_{Sing} and if the restricted map $\gamma|_V : V \dashrightarrow \mathbb{G}(n, N)$ is constant.

Projective subvarieties whose Gauss maps contract positive-dimensional subvarieties naturally appear in several contexts, notably in A. Noma's classification for projective subvarieties (see [Nom18] and the references therein). The usefulness of the classification is manifested in [Nom14], where a sharp bound for the Castelnuovo-Mumford regularity of \mathcal{O}_X was established for all non-degenerate, smooth varieties, and subsequently in [MPS20] for an extension to isolated singularities. In the least specific class of Noma's classification, a projective subvariety has positive dimensional set of non-birational centers, which is a union of linear spaces in characteristic 0 by B. Segre, and the Gauss map is constant on each of components. It is worth further understanding of the geometry of such class of varieties.

2010 *Mathematics Subject Classification.* 14N05, 14M99, 14C05.

Key words and phrases. Gauss map, Normal bundle, Hilbert scheme.

In this article, we study smooth curves that are contracted by the Gauss map. Our first main result is a characterization of contracted curves via the normal bundle.

Theorem 1.1. *Let $X \subseteq \mathbb{P}^N$ be a projective variety of dimension n and $C \hookrightarrow X$ be a smooth projective curve not contained in X_{Sing} . Then the Gauss map is constant on C if and only if the normal bundle $N_{C/X}$ of C fits into the exact sequence*

$$(1.1) \quad 0 \rightarrow P^1(\mathcal{O}_C(1))^* \otimes \mathcal{O}_C(1) \rightarrow \oplus^{n+1} \mathcal{O}_C(1) \rightarrow N_{C/X} \rightarrow 0.$$

In particular when $C = L$ is a line, the Gauss map is constant on L if and only if

$$(1.2) \quad N_{L/X} \simeq \oplus^{n-1} \mathcal{O}_L(1).$$

The sequence (1.1) is well-known when X is a projective space, and one direction of the result may be viewed as a generalization.

Next let $L \subseteq X$ be a line contracted by the Gauss map. In view of the normal bundle formula (1.2), one might expect at first glance that L deforms in a family whose members cover X . Somewhat surprisingly, we show as long as X is normal, $[L]$ *always* have local obstruction for the embedded deformation in X (except for trivial cases), by dimension counting and a deep result of Cho, Miyaoka, and Shepherd-Barron [CMSB02]. The details are presented in §4. As a consequence, we obtain

Theorem 1.2. *Assume $X \subseteq \mathbb{P}^N$ is non-degenerate and normal. Let $L \subseteq X$ be a line contracted by the Gauss map. Then any irreducible component W of Hilb^o containing $[L]$ is non-reduced everywhere, where Hilb^o is the Hilbert scheme of lines on X which are not contained in X_{Sing} .*

In [Mum62], a component of the Hilbert scheme of space curves of degree 14 and genus 24 was proved to be nowhere reduced. Since then, more pathologies of Hilbert schemes have been found. Theorem 1.2 however presents pathologies of Hilbert schemes in a different flavor; here the parameterized objects are simply lines while the ambient space could be complicated. We remark that the result readily extends to rational curves of minimal degree by the nature of the proof.

Finally we in §5 investigate upper bounds for degree of curves contracted by the Gauss map.

Conventions and notations: we work throughout over a base field k which is algebraically closed and of characteristic 0. By a variety, we mean an integral, separated scheme, finite type over k . Given k -schemes Z and T , we write Z_T for $Z \times_{\text{Spec} k} T$. Given an embedding $i : X \hookrightarrow \mathbb{P}^N$, we put $V = H^0(\mathbb{P}^N, \mathcal{O}_{\mathbb{P}^N}(1))$ and $\mathcal{O}_X(1) = i^* \mathcal{O}_{\mathbb{P}^N}(1)$.

Acknowledgements. I wish to take the opportunity to thank Xiaowen Hu, Joaquín Moraga and Jinhyung Park for stimulating discussions.

2. PRELIMINARIES

2.1. Normal bundles of smooth curves. We recall some general facts on the normal bundle of a smooth curve on a possibly singular variety.

Let C be a smooth projective curve and \mathcal{E} a coherent sheaf on C . Let τ be the torsion subsheaf of \mathcal{E} . Then \mathcal{E}/τ is torsion free, and hence locally free. Consider the exact sequence

$$0 \rightarrow \tau \rightarrow \mathcal{E} \rightarrow \mathcal{E}/\tau \rightarrow 0.$$

Since $\text{Ext}^1(\mathcal{E}/\tau, \tau) \simeq H^1(C, \tau \otimes (\mathcal{E}/\tau)^\vee) = 0$ by the Grothendieck vanishing, the sequence splits

$$(2.1) \quad \mathcal{E} \simeq \tau \oplus \mathcal{E}/\tau.$$

Let X be a projective scheme (possibly singular) and $C \hookrightarrow X$. The normal sheaf

$$(2.2) \quad N_{C/X} \simeq \mathcal{H}om_{\mathcal{O}_C}(\mathcal{I}_C/\mathcal{I}_C^2, \mathcal{O}_C) \simeq \mathcal{H}om_{\mathcal{O}_C}((\mathcal{I}_C/\mathcal{I}_C^2)/\text{tor}, \mathcal{O}_C)$$

is locally free by (2.1). This justifies the terminology ‘‘normal bundle’’.

By the theory of T^i functors (see [Har09, Thm 3.5, Ex. 3.5, 3.6]), one has the exact sequence

$$0 \rightarrow T_C \rightarrow \mathcal{H}om(\Omega_X^1|_C, \mathcal{O}_C) \rightarrow N_{C/X} \rightarrow \mathcal{T}^1(C, \mathcal{O}_C) \rightarrow \cdots,$$

where T_C is the tangent bundle of C . Since C is smooth, $\mathcal{T}^1(C, \mathcal{O}_C) = 0$ cf. [Har09, Ex. 4.3]. Thus $N_{C/X}$ fits into the short exact sequence

$$(2.3) \quad 0 \rightarrow T_C \rightarrow \mathcal{H}om(\Omega_X^1|_C, \mathcal{O}_C) \rightarrow N_{C/X} \rightarrow 0.$$

2.2. First order principle sheaf. Let \mathcal{F} be a quasi-coherent sheaf on X . We briefly recall $P^1(\mathcal{F})$, the first order principle sheaf of \mathcal{F} , with details referred to the excellent exposition [Kle77, Chap. IV, Sec. A].

In general, one has the short exact sequence

$$(2.4) \quad 0 \rightarrow \Omega_X^1 \otimes \mathcal{F} \rightarrow P^1(\mathcal{F}) \rightarrow \mathcal{F} \rightarrow 0.$$

In complex analytic context, Atiyah [Ati57, Prop. 12] shows that if \mathcal{F} is invertible, the extension class of (2.4) is indeed given by $-2\pi ic_1(\mathcal{F}) \in H^1(X, \Omega_X^1)$. By Lefschetz principle and Serre’s GAGA, this result carries over to the algebraic setting immediately.

When $X = \mathbb{P}^1$, an extension sheaf of $\mathcal{O}_{\mathbb{P}^1}$ by $\Omega_{\mathbb{P}^1}^1$ is either trivial or isomorphic to $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$. Thus for any $d \in \mathbb{Z}$, it holds

$$(2.5) \quad P^1(\mathcal{O}_{\mathbb{P}^1}(d)) \simeq \begin{cases} \mathcal{O}_{\mathbb{P}^1}(d-1) \oplus \mathcal{O}_{\mathbb{P}^1}(d-1) & d \neq 0 \\ \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-2) & d = 0. \end{cases}$$

Lemma 2.1. [[Kle77, (IV, 19)]] *For any closed embedding $X \hookrightarrow \mathbb{P}(V)$, the natural map*

$$V \otimes \mathcal{O}_X \twoheadrightarrow P^1(\mathcal{O}_X(1))$$

is surjective.

This surjection induces an embedding $\mathbb{P}(P^1(\mathcal{O}_X(1))) \hookrightarrow \mathbb{P}(V) \times X$ over X . In fact, over a smooth closed point $x \in X$, $\mathbb{P}(P^1(\mathcal{O}_X(1)) \otimes k(x))$ gives the embedded tangent space of X at x .

3. NORMAL BUNDLE AND CHARACTERIZATION

In the section, we study the normal bundle $N_{C/X}$ of a smooth curve C in a possibly singular subvariety $X \subseteq \mathbb{P}^N$. Put $\mathcal{O}_C(1) = \mathcal{O}_X(1)|_C$, so the degree of a curve equals the degree of $\mathcal{O}_C(1)$. As usual, by a line, we mean a smooth rational curve of degree one.

Proposition 3.1. *Let C be a smooth curve in X of degree d . Then $\mathcal{N}_{C/X}^*/\text{tor} \otimes \mathcal{O}_C(d)$ is globally generated. Consequently when $C = L$ is a line,*

$$N_{L/X} \simeq \bigoplus_i \mathcal{O}_{\mathbb{P}^1}(a_i),$$

for some integers $a_i \leq 1$.

Proof. Note to begin with that $\mathcal{I}_{C/\mathbb{P}^N}(d)$ is globally generated, so the evaluation morphism

$$U \otimes_k \mathcal{O}_{\mathbb{P}^N}(-d) \rightarrow \mathcal{I}_{C/\mathbb{P}^N}$$

is surjective, where $U = H^0(\mathbb{P}^N, \mathcal{I}_{C/\mathbb{P}^N}(d))$. Pulling back to X , we have

$$U \otimes_k \mathcal{O}_X(-d) \twoheadrightarrow \mathcal{I}_{C/\mathbb{P}^N} \otimes_{\mathcal{O}_{\mathbb{P}^N}} \mathcal{O}_X \simeq \mathcal{I}_{C/\mathbb{P}^N} / (\mathcal{I}_{X/\mathbb{P}^N} \cdot \mathcal{I}_{C/\mathbb{P}^N}) \twoheadrightarrow \mathcal{I}_{C/\mathbb{P}^N} / \mathcal{I}_{X/\mathbb{P}^N} \simeq \mathcal{I}_{C/X},$$

which gives the surjection

$$U \otimes_k \mathcal{O}_X(-d) \twoheadrightarrow \mathcal{I}_{C/X}.$$

Then pulling back to C , we get

$$U \otimes_k \mathcal{O}_C(-d) \twoheadrightarrow \mathcal{N}_{C/X}^*.$$

Therefore

$$\mathcal{N}_{C/X}^*/\text{tor} \otimes \mathcal{O}_C(d),$$

as a quotient of $U \otimes_k \mathcal{O}_C$ is globally generated. When $C = L$ is a line, by taking dual and applying Lemma 2.1, we get the assertion that $a_i \leq 1$ for i . \square

Proposition 3.2. *Let C be a smooth curve not contained in X_{Sing} . Then there exists an exact sequence*

$$\mathcal{I}_{C/X} / \mathcal{I}_{C/X}^2 \otimes \mathcal{O}_C(1) \rightarrow P^1(\mathcal{O}_X(1))|_C \rightarrow P^1(\mathcal{O}_C(1)) \rightarrow 0.$$

Consequently one has the short exact sequence

$$(3.1) \quad 0 \rightarrow (\mathcal{I}_{C/X} / \mathcal{I}_{C/X}^2 / \text{tor}) \otimes \mathcal{O}_C(1) \rightarrow P^1(\mathcal{O}_X(1))|_C / \text{tor} \rightarrow P^1(\mathcal{O}_C(1)) \rightarrow 0.$$

Proof. We have the commutative diagram with exact rows, see [Kle77, p. 343, (IV, 11)]

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega_X^1|_C \otimes \mathcal{O}_C(1) & \longrightarrow & P^1(\mathcal{O}_X(1))|_C & \longrightarrow & \mathcal{O}_C(1) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow = \\ 0 & \longrightarrow & \Omega_C^1 \otimes \mathcal{O}_C(1) & \longrightarrow & P^1(\mathcal{O}_C(1)) & \longrightarrow & \mathcal{O}_C(1) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array} .$$

Moreover we have the natural exact sequence

$$\mathcal{I}_{C/X}/\mathcal{I}_{C/X}^2 \rightarrow \Omega_X^1|_C \rightarrow \Omega_C^1 \rightarrow 0.$$

Putting these together with the snake lemma, we obtain the exact sequence

$$\mathcal{I}_{C/X}/\mathcal{I}_{C/X}^2 \otimes \mathcal{O}_C(1) \rightarrow P^1(\mathcal{O}_X(1))|_C \rightarrow P^1(\mathcal{O}_C(1)) \rightarrow 0.$$

This yields the surjection of locally free sheaves

$$\left(\mathcal{I}_{C/X}/\mathcal{I}_{C/X}^2/\text{tor}\right) \otimes \mathcal{O}_C(1) \rightarrow \ker\left(P^1(\mathcal{O}_X(1))|_C/\text{tor} \rightarrow P^1(\mathcal{O}_C(1))\right).$$

By the assumption that C is not contained in the singular locus, the two bundles are of the same rank $n - 1$, and hence are isomorphic. \square

We need the lemmas in the sequel.

Lemma 3.3. *Let C be a smooth curve not contained in X_{Sing} . Then the vector bundle $P^1(\mathcal{O}_X(1))|_C/\text{tor}$ is globally generated.*

Proof. In view of Lemma 2.1, we have the surjections

$$V \otimes \mathcal{O}_C \twoheadrightarrow P^1(\mathcal{O}_X(1))|_C \twoheadrightarrow P^1(\mathcal{O}_X(1))|_C/\text{tor}.$$

Since $V \otimes \mathcal{O}_C$ is globally generated, so is $P^1(\mathcal{O}_X(1))|_C/\text{tor}$. \square

Lemma 3.4. *Let C be a smooth projective curve and E is a globally generated vector bundle of rank r . Then E is trivial provided that either $h^0(C, E) = r$ or $\deg E = 0$.*

Proof. By definition of globally generation, the evaluation map

$$\text{ev} : H^0(C, E) \otimes \mathcal{O}_C \rightarrow E$$

is surjective. If $h^0(C, E) = r$, then the two bundles have the same rank. So ev is an isomorphism by Nakayama's lemma.

Now assume $\deg E = 0$. We shall proceed by induction on r . If $r = 1$, a degree 0 line bundle with sections has to be trivial. For $r \geq 2$, take a nonzero section s of E and consider the exact sequence

$$0 \rightarrow \mathcal{O}_C \xrightarrow{s} E \rightarrow E' \rightarrow 0.$$

Suppose E' has the torsion part τ . Then by the snake lemma, we arrive at the commutative diagram with exact rows and columns

$$\begin{array}{ccccccc} & & 0 & & \tau & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{O}_C & \longrightarrow & E & \longrightarrow & E' \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \longrightarrow & L & \longrightarrow & E & \longrightarrow & E'/\tau \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \tau & & 0 & & \end{array}$$

Then $\deg L > 0$, and hence $\deg(E'/\tau) = -\deg L < 0$. However this is impossible as E'/τ is globally generated.

So E' is torsion free, and hence locally free. As a quotient of E , E' is globally generated. By the induction hypothesis, $E' \simeq \oplus^{r-1} \mathcal{O}_C$. It follows that

$$r \leq h^0(E) \leq h^0(\mathcal{O}_C) + h^0(E') = r.$$

This reduces to the case $h^0(C, E) = r$ and we finish the proof. \square

Proposition 3.5. *Let $C \hookrightarrow X$ be a smooth curve not contained in X_{Sing} . Then the Gauss map is constant on C if and only if*

$$P^1(\mathcal{O}_X(1))|_C/\text{tor} \simeq \oplus^{n+1} \mathcal{O}_C.$$

Proof. Put $E = P^1(\mathcal{O}_X(1))|_C/\text{tor}$. Consider the commutative diagram

$$\begin{array}{ccccccc} & & & & f & & \\ & & & & \curvearrowright & & \\ \mathbb{P}(E) & \hookrightarrow & \mathbb{P}(P^1(\mathcal{O}_X(1))|_C) & \hookrightarrow & \mathbb{P}(P^1(\mathcal{O}_X(1))) & \hookrightarrow & X \times \mathbb{P}(V) \xrightarrow{p_2} \mathbb{P}(V). \\ & \searrow & \downarrow & & \downarrow & \swarrow p_1 & \\ & & C & \hookrightarrow & X & & \end{array}$$

The first horizontal morphism on the top row is an injection because of the surjection $P^1(\mathcal{O}_X(1))|_C \rightarrow E$, the square in the middle is cartesian, and the third horizontal morphism is an injection by Lemma 2.1.

Denote by f the composition of morphisms on the top row. The Gauss map on C can be extended to a morphism from $C \rightarrow \mathbb{G}(n, N)$ in a unique way (see [Har77, I.6.8]), which corresponds to $f : \mathbb{P}(E) \rightarrow \mathbb{P}(V)$. In view of the diagram, f is induced by a sub-linear system of $|\mathcal{O}_{\mathbb{P}(E)}(1)|$.

Suppose the Gauss map is constant on C . Then the image of f is the common embedded tangent space $\Lambda \simeq \mathbb{P}^n$ of X along C . Since $\mathcal{O}_{\mathbb{P}(E)}(1) = f^* \mathcal{O}_{\mathbb{P}(V)}(1)$ (or by Lemma 3.3), E is globally generated. Moreover we have

$$h^0(C, E) = h^0(\mathbb{P}(E), \mathcal{O}_{\mathbb{P}(E)}(1)) = h^0(\Lambda, \mathcal{O}_\Lambda(1)) = n + 1.$$

So it follows from Lemma 3.4 that E must be trivial.

Conversely if E is trivial, then $h^0(\mathbb{P}(E), \mathcal{O}_{\mathbb{P}(E)}(1)) = h^0(C, E) = n + 1$. So f factors through an n -dimensional linear space $\Lambda \hookrightarrow \mathbb{P}(V)$, and hence the Gauss map is constant on L . This completes the proof. \square

Proof of Theorem 1.1. Suppose the Gauss map γ is constant on C . The claimed exact sequence is obtained by taking dual of (3.1) and using Proposition 3.5.

Conversely, by comparison between sequence (1.1) and (3.1), we deduce that the degree of the vector bundle $E = P^1(\mathcal{O}_X(1))|_C/\text{tor}$ is 0. On the other hand, E is globally generated by Lemma 3.3. So it follows from Lemma 3.4 that E is trivial. Consequently γ is constant on C by Proposition 3.5.

Consider the case $C = L$ be a line. In view of Proposition 3.1 and the fact $P^1(\mathcal{O}_L(1)) \simeq \oplus^2 \mathcal{O}_L$, one sees that the sequence (3.1) splits. Therefore $P^1(\mathcal{O}_X(1))|_L/\text{tor}$ is trivial if and only if $N_{L/X} \simeq \oplus^{n-1} \mathcal{O}_L(1)$. This completes the proof. \square

Observe that the normal bundle $N_{C/X}$ only depends on the embedding $C \hookrightarrow X$, but not on the embedding $X \hookrightarrow \mathbb{P}^N$, so the exact sequence (1.1) imposes strong restrictions on the possible situations where C is contracted by γ . As an example, we have

Corollary 3.6. *Given $X \subseteq \mathbb{P}^N$ and $C \subseteq X$ be a smooth projective curve contracted by γ . Then for any ample line bundle L on X and a very ample linear system $U \subseteq H^0(X, \mathcal{O}_X(1) \otimes L)$, the Gauss map γ' associated to $X \subseteq \mathbb{P}(U)$ does not contract C . In particular, any d -uple embedding does not contract C .*

Proof. Keep the notations as before. By Theorem 1.1,

$$\deg N_{C/X} = (n + 1)d_0 + 2g - 2,$$

where $d_0 = \deg \mathcal{O}_X(1)|_C$. But by the ampleness of L , $\deg(\mathcal{O}_X(1) \otimes L)|_C > d_0$, so $N_{C/X}$ cannot fit into (1.1). This shows γ' does not contract C by Theorem 1.1. \square

4. LOCAL OBSTRUCTIONS AND NON-REDUCEDNESS OF HILBERT SCHEME

In this section, we study families of lines on a projective variety X contracted by the Gauss map.

Fix a very ample line bundle $\mathcal{O}_X(1)$ and a polynomial $P \in \mathbb{Q}[t]$. By Grothendieck [Gro61], the Hilbert scheme $\text{Hilb}_{X/k}^{P, \mathcal{O}_X(1)}$ of X , whose closed points parameterize the closed subschemes of X with the Hilbert polynomial P , exists and is projective. In particular for $P(t) = t + 1$, we have

$$\text{Hilb}_{X/k}^{t+1, \mathcal{O}_X(1)} \hookrightarrow \text{Hilb}_{\mathbb{P}^N/k}^{t+1, \mathcal{O}_{\mathbb{P}^N}(1)} \simeq \mathbb{G}(1, N)$$

parameterizes lines on X .

Lemma 4.1. *Let $Y \subset \mathbb{P}^N$ be a projective variety of dimension n and $y_0 \in Y$ a closed point. Then there exists no n -dimensional family of lines on Y which pass y_0 and cover Y .*

Proof. Suppose not, then there exists an irreducible projective scheme Z of dim n and a flat family \mathcal{U} of lines passing through y_0 over Z such that the morphism p

$$\begin{array}{ccc} & & p \\ & \curvearrowright & \\ \mathcal{U} & \xrightarrow{\quad} & Z \times Y \xrightarrow{p_Y} Y, \\ & \searrow p_Z & \\ & & Z \end{array}$$

is surjective. Thus for general $y \in Y$, it holds that $\dim p^{-1}(y) \geq 1$. This contradicts the fact that there exists at most one line passing through both y_0 and y . \square

Let Y be a k -scheme and $Z \hookrightarrow Y$ be a closed subscheme with defining ideal \mathcal{I}_Z . Recall (cf. [Kol96]) the *obstruction space* $\text{Obs}(Z) \subseteq \text{Ext}_{\mathcal{O}_Y}^1(\mathcal{I}_Z, \mathcal{O}_Z)$ is the smallest k -subspace such that for each sequence $0 \rightarrow J \rightarrow B \rightarrow A \rightarrow 0$, where A, B are local Artin k -algebras with residue field k and $J\mathfrak{m}_B = 0$, the obstruction class for extending a deformation of Z over A to one over B lies in $\text{Obs}(Z) \otimes_k J$.

Now let $L \subset X$ be a line contracted by the Gauss map. The following result shows that in general, the corresponding closed point $[L]$ is a singular point of $\mathrm{Hilb}_{X/k}^{1+t, \mathcal{O}_X(1)}$.

Theorem 4.2. *Assume $X \subseteq \mathbb{P}^N$ is non-degenerate and normal. Then $\mathrm{Obs}(L) \neq (0)$.*

Proof. Let W be an irreducible component containing $[L]$ with the universal family \mathcal{U} above. Once again we have the following diagram

$$\begin{array}{ccc} \mathcal{U} & \xrightarrow{\quad} & W \times X \xrightarrow{p_X} X, \\ \pi \downarrow & \swarrow p_W & \\ W & & \end{array}$$

By [Kol96, (2.10.3)] and Theorem 1.1,

$$\dim W \geq h^0(L, N_{L/X}) - \dim_k \mathrm{Obs}(L) = 2(n-1) - \dim_k \mathrm{Obs}(L).$$

Since W is proper, so is p . Thus $p(\mathcal{U})$ is closed in X .

Suppose to the contrary that $\mathrm{Obs}(L) = (0)$. Then p has to be surjective, for otherwise there would exist $x_0 \in X$, and irreducible $T \subseteq p^{-1}(x_0)$ with $\dim T \geq n$. Then $\pi(T)$ parameterizes a family of lines on $p(\mathcal{U})$ passing through x_0 with $\dim T \geq n$. This contradicts Lemma 4.1.

Then for each $x \in X$, $\dim p^{-1}(x) \geq n-1$. In view of Lemma 4.1, we deduce that for each $x \in X$, $\dim p^{-1}(x) = n-1$. Put $W_x = \pi(p^{-1}(x))$, which gives an $(n-1)$ -dimensional family of lines passing through x . Consider the induced morphism $p_x : \mathcal{U}|_{W_x} \rightarrow X$. Because $\dim \mathcal{U}|_{W_x} = \dim X$ and there exists at most one line l connecting x and any other x' , we deduce that p_x is surjective. This shows that for any $x \neq x' \in X$, there exists a line l connecting x and x' . Now by [CMSB02, Theorem 1], it holds that $X \simeq \mathbb{P}^n$. But this contradicts Zak's finiteness of Gauss maps for smooth varieties. \square

As a corollary we obtain:

Corollary 4.3. *Assumptions as above. Then L is locally obstructed.*

Proof. Suppose to the contrary that $L \subset X$ is locally unobstructed, then by [Kol96, I, Prop. 2.14], it holds that $\mathrm{Obs}(L) \subseteq H^1(L, N_{L/X})$. By Theorem 1.1, $H^1(L, N_{L/X}) = 0$, which implies that $\mathrm{Obs}(L) = (0)$, a contradiction to Theorem 4.2. \square

Lemma 4.4. *Given a flat family of smooth curves on X*

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & X_T \dashrightarrow \mathbb{G}(n, N)_T \\ & \searrow \pi & \downarrow \\ & & T \end{array}$$

with the property that for each $t \in T$, $\mathcal{C}_t \not\subseteq (X_{\mathrm{Sing}})_t$, where X_{Sing} has the induced reduced scheme structure. For each $t \in T$, one has the Gauss map $\gamma_t : \mathcal{C}_t \rightarrow \mathbb{G}(n, N)_t$. Assume the base scheme T is irreducible and of finite type over k , and

suppose that for some closed point t_0 , γ_{t_0} is constant. Then for general $t \in T$, γ_t is constant.

Proof. Note to begin with that given a curve $C \subseteq X$ and a change of the base field $\text{spec}K \rightarrow \text{spec}k$, the Gauss map $\gamma_K : C_K \dashrightarrow \mathbb{G}(n, N)_K$ is constant on C_K if and only if γ is constant on C . Thus we can assume T is reduced. Moreover, by taking a resolution of singularities $T' \rightarrow T$ and considering the family $\mathcal{C}_{T'} \rightarrow T'$, we can assume that T and hence \mathcal{C}_T are smooth. By the valuative criterion for properness, one can extend the rational map $\mathcal{C}_T \dashrightarrow \mathbb{G}(n, N)_T$ over T to a map defined over all codimension one points of \mathcal{C}_T . Therefore there exists a nonempty open subset $U \subseteq T$ over which we have the morphism $f : \mathcal{C}_U \rightarrow \mathbb{G}(n, N)_U$.

Now let $\mathcal{O}_{\mathbb{G}_U}(1)$ be the tautological line bundle on the Grassmannian $\mathbb{G}(n, N)_U$. It is evident that for each $t \in U$, γ_t is constant if and only if $f^*\mathcal{O}_{\mathbb{G}_U}(1)|_{\mathcal{C}_t}$ is trivial. Since $f^*\mathcal{O}_{\mathbb{G}_U}(1)|_{\mathcal{C}_t}$ is globally generated, $f^*\mathcal{O}_{\mathbb{G}_U}(1)|_{\mathcal{C}_t}$ is trivial if and only if its degree is zero. Then the assertion follows as the Hilbert polynomial is locally constant for fibres of a coherent sheaf flat over the base. \square

Since $Z = X_{\text{Sing}} \hookrightarrow X$, we have the closed embedding $\text{Hilb}_{Z/k}^{1+t, \mathcal{O}_X(1)} \hookrightarrow \text{Hilb}_{X/k}^{1+t, \mathcal{O}_X(1)}$. Put $\text{Hilb}^o = \text{Hilb}_{X/k}^{1+t, \mathcal{O}_X(1)} \setminus \text{Hilb}_{Z/k}^{1+t, \mathcal{O}_X(1)}$ with the induced scheme structure, and let W be an irreducible component of Hilb^o .

Proof of Theorem 1.2. By Lemma 4.4, for a general point $x : \text{spec}k \rightarrow W$, the Gauss map on the corresponding line L_x is constant, and hence x is singular by Theorem 4.2. Suppose to the contrary that W is reduced at some point x' , then there exists an open neighborhood $U \subseteq W$ of x' such that U is reduced with the induced scheme structure. Then by the generic smoothness, there exists a nonempty open $V \subseteq U$ that is smooth over k . This gives a contradiction. \square

5. BOUND OF DEGREE

Gauss maps which are generically finite but not quasi-finite seem much less understood than the degenerate ones. Mezzetti-Tommasi give an example (c.f. [Ran19, Remark 2]) where a singular quartic surface has a generically finite Gauss map which contracts a smooth plane quadric.

In surface case, we give an upper bound of the degree of the contracted curves. Note in higher dimensions, such a bound cannot exist because the closure of a fibre of the Gauss map can be \mathbb{P}^m with $m \geq 2$.

Proposition 5.1. *Let X be a non-degenerate singular surface in \mathbb{P}^N . Suppose γ is constant on a smooth projective curve C . Then C is a plane curve and*

$$\deg X \geq 2 \deg C + (N - 3).$$

Proof. It is clear that C is a plane curve contained in the embedded tangent space $\widehat{T} = \widehat{T}_x X$ for general $x \in C$. When $N = 3$, since X and \widehat{T} intersect at C doubly, by Bézout's theorem, we have $\deg X \geq 2 \deg C$. For $N > 3$, note the secant variety $\text{Sec}(C) \subseteq \widehat{T}$. Let $\overline{C}(X)$ be the closure of the non-birational centers of X , which is a proper linear space in X . Then for any smooth point $x_0 \in X \setminus (\text{Sec}(C) \cup \overline{C}(X))$,

the inner projection $\pi : X \setminus \{x_0\} \rightarrow \mathbb{P}^{N-1}$ gives rise to a birational map $X \dashrightarrow X' := \pi(X \setminus \{x_0\})$ such that the restriction $\pi|_C : C \rightarrow C' := \pi(C)$ is an isomorphism. So $X' \subseteq \mathbb{P}^{N-1}$ is a surface with a smooth rational curve C' on it, with $\deg X' = \deg X - 1$ and $\deg C' = \deg C$. Fix a closed point $y_0 \in C \setminus X_{\text{Sing}}$. By taking the center x_0 general, we can guarantee that π is isomorphic at y_0 , and hence is isomorphic at a neighborhood U of y_0 . Therefore for $y \in U$, $\widehat{T}_{\pi(y)}(X') = \pi(\widehat{T}_y(X)) = \pi(\widehat{T})$, which implies that $\gamma_{X'}$ is constant on C' provided x_0 is taken general. So we can proceed by induction on N . \square

In a different direction, we establish an upper bound for the degree of contracted curves by the Gauss map with respect to an anticanonical divisor.

Theorem 5.2. *Let $X \subseteq \mathbb{P}^N$ be a normal \mathbb{Q} -Gorenstein variety of dimension n . Let $C \subseteq X$ be a smooth projective curve of degree d . Suppose the Gauss map is constant on C . Then $-K_X \cdot C \leq (n+1)d$.*

Proof. Suppose rK_X is Cartier for some integer $r > 0$. Put $\Omega_X^n = \wedge^n \Omega_X$. We have the natural morphism $\eta : \Omega_X^n \rightarrow \omega_X$ (cf. [EM09, Appendix]), where ω_X is the canonical sheaf. The morphism η induces $(\Omega_X^n)^{[r]} \rightarrow \omega_X^{[r]} \simeq \mathcal{O}_X(rK_X)$, where $\star^{[r]}$ denotes the double dual of $\star^{\otimes r}$. Consider the composite of the natural morphisms

$$\xi : ((\Omega_X^n)|_C)^{\otimes r} \simeq (\Omega_X^n)^{\otimes r}|_C \rightarrow (\Omega_X^n)^{[r]}|_C \rightarrow \mathcal{O}_X(rK_X)|_C.$$

Since ξ is generically injective, the induced morphism

$$((\Omega_X^n)|_C)^{\otimes r} / \text{tor} \rightarrow \mathcal{O}_X(rK_X)|_C$$

is injective. Consequently

$$rK_X \cdot C = \deg \mathcal{O}_X(rK_X)|_C \geq \deg((\Omega_X^n)|_C)^{\otimes r} / \text{tor}.$$

To finish the proof, note by (2.3) and Theorem 1.1, $\mathcal{H}om(\Omega_X^1|_C, \mathcal{O}_C)$ is a vector bundle of degree $(n+1)d$, so $(\Omega_X^n|_C)/\text{tor}$ is a line bundle of degree $-(n+1)d$. \square

REFERENCES

- [AG01] M. A. Akivis and V. V. Goldberg. On the structure of submanifolds with degenerate Gauss maps. *Geometriae Dedicata*, 86(1):205–226, 2001.
- [Ati57] M. F. Atiyah. Complex analytic connections in fibre bundles. *Trans. Amer. Math. Soc.*, 85(1):181–207, 1957.
- [CMSB02] K. Cho, Y. Miyaoka, and N. I. Shepherd-Barron. Characterizations of projective space and applications to complex symplectic manifolds. In *Higher dimensional birational geometry*, pages 1–88. Mathematical Society of Japan, 2002.
- [EM09] L. Ein and M. Mustață. Jet schemes and singularities. *Proc. Sympos. Pure Math., part 2*, 80:505–546, 2009.
- [GH79] P. Griffiths and J. Harris. Algebraic geometry and local differential geometry. *Ann. Sci. École Norm. Sup. (4)*, 12(3):355–452, 1979.
- [Gro61] A. Grothendieck. Techniques de construction et théorèmes d'existence en géométrie algébrique IV: Les schémas de Hilbert. *Séminaire Bourbaki*, 221, 1961.
- [Har77] R. Hartshorne. *Algebraic Geometry*, volume 52. Springer-Verlag, New York, 1977.
- [Har09] R. Hartshorne. *Deformation Theory*, volume 257. Springer Science & Business Media, 2009.

- [Kle77] S. L. Kleiman. The enumerative theory of singularities. *Real and complex singularities (Proc. Ninth Nordic Summer School, Oslo, 1976)*, pages 297–396, 1977.
- [Kol96] J. Kollár. *Rational Curves on Algebraic Varieties*, volume 32. Springer-Verlag, Berlin, 1996.
- [MPS20] J. Moraga, J. Park, and L. Song. Regularity of structure sheaves of varieties with isolated singularities. *Commun. Contemp. Math.*, page 2050039, 2020.
- [Mum62] D. Mumford. Further pathologies in algebraic geometry. *Amer. J. Math.*, 84(4):642–648, 1962.
- [Nom14] A. Noma. Generic inner projections of projective varieties and an application to the positivity of double point divisors. *Trans. Amer. Math. Soc.*, 366(9):4603–4623, 2014.
- [Nom18] A. Noma. Projective varieties with nonbirational linear projections and applications. *Trans. Amer. Math. Soc.*, 370(4):2299–2320, 2018.
- [Ran19] Z. Ran. Stationary flags, Gaussian scrolls and duality. *arXiv preprint arXiv:1909.03307*, 2019.
- [Zak93] F. L. Zak. Tangents and secants of algebraic varieties, Translations of Mathematical Monographs, vol. 127. *American Mathematical Society, Providence, RI*, 1993.

SCHOOL OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, W. 135 XINGANG ROAD, GUANGZHOU, GUANGDONG 510275, CHINA

Email address: songlei3@mail.sysu.edu.cn