

A THEOREM ON MEROMORPHIC DESCENT AND THE SPECIALIZATION OF THE PRO-ÉTALE FUNDAMENTAL GROUP

MARCIN LARA, JIU-KANG YU, LEI ZHANG

ABSTRACT. Given a Noetherian formal scheme \hat{X} over $\mathrm{Spf}(R)$, where R is a complete DVR, we first prove a theorem of meromorphic descent along a possibly infinite cover of \hat{X} . Using this we construct a specialization functor from the category of continuous representations of the pro-étale fundamental group of the special fiber to the category of F -divided sheaves on the generic fiber. This specialization functor partially recovers the specialization functor of the étale fundamental groups. We also express the pro-étale fundamental group of a connected scheme X of finite type over a field as coproducts and quotients of the free group and the étale fundamental groups of the normalizations of the irreducible components of X and those of its singular loci.

INTRODUCTION

Let R be a complete DVR with function field K , and let X be a proper scheme over $\mathrm{Spec}(R)$. Let X_0 be the special fiber and X_K the generic fiber. Thanks to Grothendieck's existence theorem the pullback along $X_0 \subseteq X$ induces an equivalence from the category $\mathrm{ECov}(X)$ of finite étale covers of X to the category $\mathrm{ECov}(X_0)$ of finite étale covers of X_0 . In this way, we get the specialization functor

$$(1) \quad \mathrm{ECov}(X_0) \xrightarrow{\simeq} \mathrm{ECov}(X) \longrightarrow \mathrm{ECov}(X_K)$$

If X, X_0, X_K are moreover connected, and if $x_0 \in X_0, x_K \in X_K$ are two geometric points, then up to a choice of a path between x_0 and x_K , there is the specialization map

$$(2) \quad \pi_1^{\acute{e}t}(X_K, x_K) \longrightarrow \pi_1^{\acute{e}t}(X, x_K) \simeq \pi_1^{\acute{e}t}(X, x_0) \xrightarrow{\simeq} \pi_1^{\acute{e}t}(X_0, x_0)$$

In [3], B. Bhatt and P. Scholze introduced the notion of the pro-étale fundamental group. This is a topological group which classifies geometric covers $\mathrm{Cov}(-)$ which are étale but possibly infinite. If we consider $\mathrm{Cov}(-)$ instead of $\mathrm{ECov}(-)$, then we quickly run into a problem: Grothendieck's existence theorem is not at our disposal anymore, because it can only be applied to the algebraization of *finite* covers.

Indeed, it is known to E. Lavanda [10, Remark 3.2, p. 39] that the naive specialization map of the pro-étale fundamental group does not exist. However, she managed to construct a version of the specialization functor ([10, Proposition 4.15, p. 52])

$$\mathrm{sp}_K : \mathrm{Rep}_K^{\mathrm{cts}}(\pi_1^{\mathrm{pro}\acute{e}t}(X_0, x_0)) \longrightarrow \mathrm{Fdiv}(X_K)$$

where R is a complete DVR of equal characteristic p , X is a semi-stable curve over R , and $\mathrm{Rep}_K^{\mathrm{cts}}$ denotes the category of finite dimensional continuous representations viewing K as a discrete field. This specialization functor also partially recovers the one of the étale fundamental groups in the following sense: there is a 2-commutative diagram

$$\begin{array}{ccc} \mathrm{Rep}_K^{\mathrm{cts}}(\pi_1^{\acute{e}t}(X_0, x_0)) & \xrightarrow{\mathrm{sp}^{\acute{e}t}} & \mathrm{Fdiv}(X_K) \\ \subseteq \downarrow & & \downarrow = \\ \mathrm{Rep}_K^{\mathrm{cts}}(\pi_1^{\mathrm{pro}\acute{e}t}(X_0, x_0)) & \xrightarrow{\mathrm{sp}_K} & \mathrm{Fdiv}(X_K) \end{array}$$

where $\text{Rep}_K^{\text{cts}}(\pi_1^{\acute{e}t}(X_0, x_0))$ is seen as the full subcategory of $\text{Rep}_K^{\text{cts}}(\pi_1^{\text{pro}\acute{e}t}(X_0, x_0))$ consisting of those representations whose monodromy groups are finite ([3, Theorem 1.10, p. 104]), and $\text{sp}_{\acute{e}t}$ is the composition:

$$(3) \quad \text{Rep}_K^{\text{cts}}(\pi_1^{\acute{e}t}(X_0, x_0)) \longrightarrow \text{Rep}_K^{\text{cts}}(\pi_1^{\acute{e}t}(X_K, x_K)) \longrightarrow \text{Fdiv}(X_K)$$

where the second map is recalled in 1.15.

In this paper, we first generalize E. Lavanda's specialization map ([10, Theorem 4.16, p. 54]) to higher dimensional schemes (cf. 1.14, 1.16). Another approach of considering the de Jong fundamental group of the rigid generic fiber can be found in [1].

If π_1 is a topological group, then we denote by π_1^{cts} the algebraic hull of π_1 as in [10, Definition 2.2, p. 36]. Here is our first main result:

Theorem I. *Let X be a proper scheme over a complete discrete valuation ring R with a connected special fiber. Then the functor sp_K exists and recovers the functor $\text{sp}_{\acute{e}t}$. If the residue field k of R is algebraically closed, and if X is flat geometrically connected over R , then there is a finite extension L of the function field K of R and a homomorphism of L -group schemes:*

$$\pi_1^{\text{Fdiv}}(X \otimes_R L, x_L) \longrightarrow \pi_1^{\text{pro}\acute{e}t}(X_0, x_0)^{\text{cts}}$$

which recovers the one of the étale fundamental groups (cf. 1.20).

There are two main ingredients in E. Lavanda's construction of sp_K : (1) A formula for the pro-étale fundamental group of a semi-stable curve [10, Theorem 1.17, p. 33]; (2) The theory of meromorphic descent along a torsor under a torsion-free discrete group developed by D. Gieseker in [5, Lemma 1, p. 97]. Her proof of (2) is only for curves, partly because it relies on (1). Here we generalized (2) to arbitrary discrete groups and any Noetherian base scheme (cf. 1.11). Using this we can construct sp_K directly.

Theorem II. *If q is a G -torsor, where G is any discrete group, and if \hat{X} is Noetherian, then the pullback along q induces an equivalence between $\text{Coh}(\hat{X}) \otimes_R K$ and the category of sheaves in $\text{Coh}(\hat{Y}) \otimes_R K$ equipped with meromorphic descent data.*

On the other hand, we also generalize (1) to very general schemes, even though it is not needed in our generalization of (2). We consider the case when X is a connected Nagata Noetherian J-2 scheme (e.g. a connected scheme of finite type over a field). The idea is to express the pro-étale fundamental group of X in terms of the étale fundamental groups of the normalization of its irreducible components and the pro-étale fundamental group of its singular locus. Then using Noetherian induction one may understand the pro-étale fundamental group based on one's knowledge on the étale fundamental groups.

Theorem III. *Let X be a connected Nagata Noetherian J-2 scheme. If X is irreducible, and if the singular locus Z of the reduced induced structure of X is connected, then there is an isomorphism of topological groups*

$$\pi_1^{\text{pro}\acute{e}t}(X, x) \simeq \left[\frac{\left(\pi_1^{\acute{e}t}(\tilde{X}, x_1) \coprod \pi_1^{\text{pro}\acute{e}t}(Z_1, x_1) \pi_1^{\text{pro}\acute{e}t}(Z, x) \right) \coprod \coprod_{2 \leq j \leq n} \mathbb{Z}_j}{\langle \{ \phi_j(a) \cdot 1_j = 1_j \cdot \psi_j(a) \mid 2 \leq j \leq n, a \in \pi_1^{\text{pro}\acute{e}t}(Z_j, x_j) \} \rangle} \right]^{\text{Noohi}}$$

where $\mathbb{Z}_j := \mathbb{Z}$, \tilde{X} is the normalization of X , $\{Z_j\}_{1 \leq j \leq n}$ are connected components of the inverse image of Z under $f: \tilde{X} \rightarrow X$, $* \coprod^* *$ is the coproduct in the category of Noohi groups (2.5), $[\frac{*}{*}]^{\text{Noohi}}$ is the Noohi quotient (2.4), and

$$\phi_j: \pi_1^{\text{pro}\acute{e}t}(Z_j, x_j) \rightarrow \pi_1^{\text{pro}\acute{e}t}(Z, x)$$

$$\psi_j: \pi_1^{\text{pro}\acute{e}t}(Z_j, x_j) \rightarrow \pi_1^{\text{pro}\acute{e}t}(\tilde{X}, x_1)$$

are obtained by choosing paths to identify fundamental groups with different chosen points.

If X has more irreducible components, then we “glue” them together along Z (cf. 2.20). If Z has more connected components, then we compute $\pi_1^{\text{proét}}(X, x)$ by induction on the number of connected components of Z (cf. 2.24).

As an application of Theorem III, we have

Theorem IV. *If X is a connected scheme of finite type over a separably closed field k and if the singular locus Z is 0-dimensional, then we have*

$$\pi_1^{\text{proét}}(X, x) \simeq \left(\coprod_{1 \leq i \leq n} \pi_1^{\text{ét}}(\tilde{X}_i, x_i) \right) \amalg \left(\coprod_{1 \leq j \leq |f^{-1}(Z)| - |Z| - n + 1} \mathbb{Z}_j \right)$$

where $|\cdot|$ denote the number of topological points of the underlying scheme (cf. 2.27).

As another application of Theorem III we prove that a continuous representation of $\pi_1^{\text{proét}}(X, x)$ factors through a discrete quotient iff its restrictions to the étale fundamental groups of the normalizations of the irreducible components of X do so (cf. 2.31).

1. MEROMORPHIC DESCENT

1.1. Noetherian meromorphic descent.

Conventions 1.1. Let R be an adic Noetherian ring, and let $I \subseteq R$ be an ideal of definition. Suppose that $I = (f)$. Let K be a localization of R at the element f . Let \hat{X} be an *adic formal scheme* ([6, 10.4.2, p. 407]) equipped with an *adic morphism* $\hat{X} \rightarrow \text{Spf}(R)$ ([6, 10.12.1, p. 436]). In this case, we simply call \hat{X} a formal scheme over $\text{Spf}(R)$ or over R . We denote by $\text{Coh}(\hat{X})$ the category of sheaves of $\mathcal{O}_{\hat{X}}$ -modules on \hat{X} whose pullbacks to the scheme $X_n := (\hat{X}, \mathcal{O}_{\hat{X}}/I^{n+1})$ are finitely presented for each $n \in \mathbb{N}$. If \hat{X} is a locally Noetherian formal scheme ([6, 10.4.2, p. 407]), then $\text{Coh}(\hat{X})$ is just the category of coherent sheaves on \hat{X} ([15, 01XZ]).

Definition 1.2. We denote by $\text{Coh}^m(\hat{X})$ the category whose objects are exactly the same as those in $\text{Coh}(\hat{X})$, whose set of morphisms between two objects \mathcal{F}, \mathcal{G} is

$$\Gamma(\hat{X}, \underline{\text{Hom}}_{\mathcal{O}_{\hat{X}}}(\mathcal{F}, \mathcal{G}) \otimes_R K)$$

where $\underline{\text{Hom}}$ denotes the internal sheaf Hom.

Definition 1.3. Let \mathcal{C} be an R -linear category. We define the category $\mathcal{C} \otimes_R K$ to the category whose objects are exactly the same as those in \mathcal{C} , whose set of morphisms between two objects X, Y is $\text{Hom}_{\mathcal{C}}(X, Y) \otimes_R K$. If \mathcal{C} is an Abelian category, then so is $\mathcal{C} \otimes_R K$ ([13, Lemma 1.5, p. 7]).

Definition 1.4. Let $q: \hat{Y} \rightarrow \hat{X}$ be an adic map of locally Noetherian formal schemes over $\text{Spf}(R)$. We define the category $\text{Coh}^m(\hat{Y}/\hat{X})$ of coherent sheaves with *meromorphic descent data* (MDD for short) to be the category whose objects are pairs (\mathcal{F}, ϕ) , where \mathcal{F} belongs to $\text{Coh}^m(\hat{Y})$ and $\phi: p_1^* \mathcal{F} \rightarrow p_2^* \mathcal{F}$ is a map in $\text{Coh}^m(\hat{Y} \times_{\hat{X}} \hat{Y})$ which satisfies

- the identity condition: the pullback of ϕ along the diagonal map $\hat{Y} \rightarrow \hat{Y} \times_{\hat{X}} \hat{Y}$ is the identity map in $\text{Coh}^m(\hat{Y})$;
- the cocycle condition: we have an equality

$$p_{23}^* \phi \circ p_{12}^* \phi = p_{13}^* \phi$$

in $\text{Coh}^m(\hat{Y} \times_{\hat{X}} \hat{Y} \times_{\hat{X}} \hat{Y})$,

whose morphisms between two objects (\mathcal{F}, ϕ) and (\mathcal{G}, φ) are those morphisms in

$$\text{Hom}_{\text{Coh}^m(\hat{Y})}(\mathcal{F}, \mathcal{G})$$

which are compactifiable with ϕ and φ .

Theorem 1.5. *Let $q: \hat{Y} \rightarrow \hat{X}$ be an adic map of Noetherian formal schemes over $\mathrm{Spf}(R)$. Then there is a canonical functor*

$$\mathrm{Coh}(\hat{X}) \otimes_R K \longrightarrow \mathrm{Coh}^m(\hat{Y}/\hat{X})$$

which is an equivalence if q is faithfully flat.

Proof. The functor is simply the formation of the *canonical descent datum* ([15, 023D]). A proof of this theorem, which is due to O. Gabber, can be found in [12, (1.9), p. 774] (the result was stated there when R is a complete DVR and $K = \mathrm{Frac}(R)$, but the proof works in our setting). \square

Example 1.6. Theorem 1.5 would not be true if one replaces “Noetherian” by “locally Noetherian”. Let R be a complete DVR with maximal ideal $I = (\pi)$ and quotient field K . Let T (e.g. $R[x, y]/(xy)$) be a scheme over R equipped with two disjoint opens U_x, U_y (e.g. $U_x = (x \neq 0)$, $U_y = (y \neq 0)$) whose special fibers are not empty and $U_x \simeq U_y$ as R -schemes. Let $\{T_n | n \in \mathbb{Z}\}$ be \mathbb{Z} -copies of T . We glue $U_{y_n} \subseteq T_n$ with $U_{x_{n+1}} \subseteq T_{n+1}$ for all $n \in \mathbb{Z}$ in the sense of [8, Chapter II, Exercise 2.12, p. 80]. Then we get a scheme X with a Zariski-covering $\{T_i\}_{i \in \mathbb{Z}}$ such that

$$T_i \cap T_j = \begin{cases} \emptyset, & \text{if } i - j \neq \pm 1; \\ T_i = T_j, & \text{if } i = j; \\ U_{y_i} = U_{x_j}, & \text{if } j = i + 1; \end{cases}$$

Set $Y = \coprod_{n \in \mathbb{Z}} T_n$, and let $q: \hat{Y} \rightarrow \hat{X}$ be the formal completion of the map $Y \rightarrow X$ along the special fiber. Now if we take $\mathcal{O}_{\hat{Y}}$ on \hat{Y} and glue the restrictions on \hat{U}_{x_i} by the uniformizer π . The so obtained meromorphic descent data can not descent to \hat{X} .

This shows that meromorphic descent may not be effective even for a Zariski covering with by infinitely many Zariski open sets. So the next result is somewhat surprising.

1.2. Meromorphic descent along G -torsors.

Conventions 1.7. Let R is a complete DVR, and let k be the residue field of R . Let K be the quotient field. Let π be a uniformizer of R , $S := \mathrm{Spec}(R)$, $\hat{S} := \mathrm{Spf}(R)$. Let $\hat{X} \rightarrow \hat{S}$ be an adic morphism of Noetherian formal schemes. We denote X_n the scheme given by the ringed space $(\hat{X}, \mathcal{O}_{\hat{X}}/(\pi)^{n+1})$. Let $q: \hat{Y} \rightarrow \hat{X}$ be a torsor under a discrete group G , i.e. q is a faithfully flat adic map of formal schemes over \hat{S} and there is a group homomorphism $\rho: G \rightarrow \mathrm{Aut}_{\hat{X}}(\hat{Y})$ such that the induced map

$$\begin{aligned} \hat{Y} \times G &\longrightarrow \hat{Y} \times_{\hat{X}} \hat{Y} \\ (y, w) &\longmapsto (y, \rho(w)(y)) \end{aligned}$$

is an isomorphism, where $\hat{Y} \times G := \coprod_{w \in G} \hat{Y}$. Note that \hat{Y} is a locally Noetherian formal scheme, but not Noetherian when G is infinite. We denote by Y_n the scheme given by the ringed space $(\hat{Y}, \mathcal{O}_{\hat{Y}}/(\pi)^{n+1})$.

By using the isomorphism $\hat{Y} \times_{\hat{X}} \hat{Y} \simeq \hat{Y} \times G$, it is evident that $\mathrm{Coh}^m(\hat{Y}/\hat{X})$ is equivalent to the category $\mathrm{Coh}^m(\hat{Y}/\hat{X}, G)$, which we define now:

Definition 1.8. The category $\mathrm{Coh}^m(\hat{Y}/\hat{X}, G)$ of *coherent sheaves with MDD* is the category whose objects are pair $(\mathcal{F}, \{h_w\}_{w \in G})$ where \mathcal{F} is an object in $\mathrm{Coh}^m(\hat{Y})$ and $\{h_w\}_{w \in W}$ is a collection of elements such that

$$h_w \in \mathrm{Hom}_{\mathrm{Coh}^m(\hat{Y})}(\mathcal{F}, w^* \mathcal{F}) = \Gamma(\hat{Y}, \underline{\mathrm{Hom}}_{\mathcal{O}_{\hat{Y}}}(\mathcal{F}, w^* \mathcal{F}) \otimes_R K)$$

satisfying the “identity condition” and the “cocycle condition”:

- the identity condition: $h_e = \mathrm{id}$ for the unit $e \in G$;
- the cocycle condition: we have an equality

$$w^*(h_{w'}) \circ h_w = h_{w'w}$$

for all $w, w' \in G$.

A map

$$(\mathcal{F}, \{h_w\}_{w \in G}) \longrightarrow (\mathcal{F}', \{h'_w\}_{w \in G})$$

between two coherent sheaves with MDD is an element in

$$\mathrm{Hom}_{\mathrm{Coh}^m(\hat{Y})}(\mathcal{F}, \mathcal{F}') = \Gamma(\hat{Y}, \underline{\mathrm{Hom}}_{\mathcal{O}_{\hat{Y}}}(\mathcal{F}, \mathcal{F}') \otimes_R K)$$

which is compatible with $\{h_w\}_{w \in G}$ and $\{h'_w\}_{w \in G}$ in the obvious sense.

Definition 1.9. Let $(\mathcal{F}, \{h_w\}_{w \in G})$ be a coherent sheaf with MDD on \hat{Y} . We say that it is *genuine* if for all $w \in G$, the element h_w lies in the subset $\mathrm{Hom}_{\mathrm{Coh}(\hat{Y})}(\mathcal{F}, w^*\mathcal{F})$ of $\mathrm{Hom}_{\mathrm{Coh}^m(\hat{Y})}(\mathcal{F}, w^*\mathcal{F})$.

Lemma 1.10. *Any object in $\mathrm{Coh}^m(\hat{Y}/\hat{X}, G)$ is isomorphic to a genuine one.*

Proof. Suppose that $(\mathcal{F}, \{h_w\}_{w \in G})$ is an object in $\mathrm{Coh}^m(\hat{Y}/\hat{X}, G)$. Let \mathcal{G} be the image of \mathcal{F} in $\mathcal{F} \otimes_R K$. Then we have $\mathcal{F} \otimes_R K \simeq \mathcal{G} \otimes_R K$. Thus replacing \mathcal{F} by \mathcal{G} we may assume that \mathcal{F} is R -torsion free. Now we claim that there is a coherent subsheaf $\mathcal{F}' \subseteq \mathcal{F} \otimes_R K$ such that $\mathcal{F}' \otimes_R K = \mathcal{F} \otimes_R K$ and h_w sends $\mathcal{F}' \subseteq \mathcal{F} \otimes_R K$ to $w^*\mathcal{F}' \subseteq w^*\mathcal{F}' \otimes_R K = w^*\mathcal{F} \otimes_R K$. Evidently, the lemma follows from this claim.

We will find \mathcal{F}' using Noetherian induction on \hat{X} . The key is the induction step: if the claim is true when \hat{Y}/\hat{X} is replaced by $q^{-1}(U)/U$ for some open U of \hat{X} , and $U \neq \hat{X}$, then there exists an open U' such that $U \subsetneq U' \subset \hat{X}$ and the claim is true when \hat{Y}/\hat{X} is replaced by $q^{-1}(U')/U'$.

Let $Z = \hat{X} \setminus U$. We regard Z as a closed subscheme of X_0 with the reduced induced structure. Let $\eta \in Z$ be a generic point of Z . Since

$$\pi_1^{\mathrm{pro\acute{e}t}}(\mathrm{Spec}(\kappa(\eta))) = \pi_1^{\acute{e}t}(\mathrm{Spec}(\kappa(\eta)))$$

is profinite, the G -torsor $\hat{Y}|_{\mathrm{Spec}(\kappa(\eta))}$ obtained by pulling back \hat{Y} through

$$\mathrm{Spec}(\kappa(\eta)) \hookrightarrow Z \hookrightarrow X_0 \hookrightarrow \hat{X}$$

is induced from an H -torsor over $\mathrm{Spec}(\kappa(\eta))$ with $H \subseteq G$ finite. Spreading it out we find $V \subseteq Z$, an open neighborhood of x , and an H -torsor $\tilde{Z} \rightarrow V$ which induces $\hat{Y}|_V$. Thus \tilde{Z} is an open in $\hat{Y}|_V = q^{-1}(V)$ and $\hat{Y}|_V$ is the disjoint union of $w\tilde{Z}$ over $w \in G/H$.

We can find an H -invariant quasi-compact open subset W of Y_0 such that $W \cap q^{-1}(Z) = \tilde{Z}$. Consider W as an open formal subscheme of \hat{Y} . We can convert the MDD on \mathcal{F} in the sense of 1.8 to an MDD ϕ in the sense of 1.4, and restrict it to get an MDD for $W \rightarrow q(W)$. By Gabber's theorem 1.5, there is a coherent sheaf \mathcal{E} on $q(W)$ such that $(q|_W)^*\mathcal{E}$ is isomorphic to $(\mathcal{F}|_W, \phi)$ in $\mathrm{Coh}^m(W/q(W))$. We may and do assume that the isomorphism is given by a morphism in $\mathrm{Hom}_{\mathrm{Coh}(W)}((q|_W)^*\mathcal{E}, \mathcal{F}|_W)$. Let \mathcal{E}' be the image of this morphism.

Let \mathcal{F}' be the coherent sheaf on $q^{-1}(U)$ asserted by the induction hypothesis. We now use \mathcal{F}' on $q^{-1}(U)$ and \mathcal{E}' on W to form a sheaf \mathcal{F}'' on $q^{-1}(U')$ extending \mathcal{F}' , where $U' := U \cup q(W) = U \cup q(\tilde{Z})$, using a method of Gieseker and Raynaud, as follows. We first try to extend it to $q^{-1}(U) \cup W = q^{-1}(U) \cup \tilde{Z}$. On the open set $q^{-1}(U) \cap W$, the restriction of $\mathcal{F}, \mathcal{F}', \mathcal{E}'$ become the same upon tensoring with K . Therefore, for n large enough we have

$$\pi^n \mathcal{E}' \subset \mathcal{F}' \subset \pi^{-n} \mathcal{E}' \quad \text{on } q^{-1}(U) \cap W.$$

This implies

$$\mathcal{F}'/\pi^n \mathcal{E}' \subset \pi^{-n} \mathcal{E}'/\pi^n \mathcal{E}' \quad \text{on } q^{-1}(U) \cap W.$$

On the other hand, the right-hand side is obviously the restriction of a coherent sheaf on W_{2n} (by which we mean W regarded as an open subscheme of Y_{2n}). By [15, 01PD], the left-hand side extends to a coherent sheaf \mathcal{G} on W such that \mathcal{G} is a subsheaf of $\pi^{-n} \mathcal{E}'/\pi^n \mathcal{E}'$. Let \mathcal{E}'' be the subsheaf of $\pi^{-n} \mathcal{E}'$ such that $\mathcal{E}''/\pi^n \mathcal{E}' = \mathcal{G}$. By construction, $\mathcal{E}'' = \mathcal{F}'$ on $q^{-1}(U) \cap W$. Since both $\mathcal{E}', \mathcal{F}'$ are H -invariant and the extension in [15, 01PD] is functorial, \mathcal{E}'' is also H -invariant.

We now define a sheaf \mathcal{F}'' on

$$q^{-1}(U') = q^{-1}(U) \cup \bigcup_{w \in G} w^{-1}W$$

as follows: it is the subsheaf of $\mathcal{F} \otimes_R K$ generated by sections of \mathcal{F}' and sections of $h_w^{-1}(w^*\mathcal{E}'')$ (a sheaf on $w^{-1}W$) for all $w \in G$. Thus a local section of $\mathcal{F} \otimes_R K$ is in \mathcal{F}'' iff its stalk at each point is a sum of stalks of some local sections of \mathcal{F}' and $h_w^{-1}(w^*\mathcal{E}'')$. We claim that \mathcal{F}'' fulfills all the conditions required by the induction step.

From the above definition and the induction hypothesis, it is easy to see that $\mathcal{F}_y'' = \mathcal{F}_y'$ for all $y \in q^{-1}(U)$. For $y \in q^{-1}(U') \setminus q^{-1}(U)$, $\mathcal{F}_y'' = \sum h_w^{-1}(\mathcal{E}_z'')$, where the sum is over $\{(w, z) : w \in G, z \in \tilde{Z}, w^{-1}z = y\}$, a transitive H -set. It follows that for any $w \in G$,

$$\mathcal{F}''|_{w^{-1}W} = \sum_{j \in H} h_{jw}^{-1}((jw)^*\mathcal{E}'') = h_w^{-1}(w^*\mathcal{E}'')$$

where the second equality follows from the fact that both W and \mathcal{E}'' are H -invariant. We conclude that \mathcal{F}'' is coherent on $q^{-1}(U')$.

It remains to verify that h_u maps \mathcal{F}'' to $u^*\mathcal{F}''$ for each $u \in G$. It suffices to check this over $q^{-1}(U)$ and $w^{-1}W$ for all $w \in G$. The case of $q^{-1}(U)$ is clear. Over $w^{-1}W$, we have the following commutative diagrams:

$$\begin{array}{ccc} & uw^{-1}W & \\ u \nearrow & & \searrow wu^{-1} \\ w^{-1}W & \xrightarrow{w} & W \end{array} \quad \begin{array}{ccc} & u^*h_{wu^{-1}}^{-1}((wu^{-1})^*\mathcal{E}'') & \\ h_u \nearrow & & \searrow u^*h_{wu^{-1}} \\ h_w^{-1}(w^*\mathcal{E}'') & \xrightarrow{h_w} & w^*\mathcal{E}'' \end{array}$$

due to the cocycle condition $h_w = (u^*h_{wu^{-1}}) \circ h_u$. Thus h_u maps $\mathcal{F}''|_{w^{-1}W}$ to $u^*(\mathcal{F}''|_{uw^{-1}W})$ which is exactly $(u^*\mathcal{F}'')|_{w^{-1}W}$. The lemma is proved completely. \square

Theorem 1.11. *The pullback functor*

$$\mathrm{Coh}(\hat{X}) \otimes_R K \longrightarrow \mathrm{Coh}^m(q, G)$$

is an equivalence.

Proof. It is enough to show that the functor is fully faithful. Then the last statement follows readily from 1.10. By fpqc-descent of quasi-coherent sheaves we have for any two sheaves $\mathcal{F}, \mathcal{G} \in \mathrm{Coh}(\hat{X})$ an exact sequence of R -modules:

$$0 \longrightarrow \mathrm{Hom}_{\mathcal{O}_{\hat{X}}}(\mathcal{F}, \mathcal{G}) \longrightarrow \mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \longrightarrow \prod_{w \in G} \mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G})$$

where the second map is the pullback, and the third map is $w^* - \mathrm{id}^*$. Since we work in $\mathrm{Coh}(\hat{X}) \otimes_R K$, we may assume that \mathcal{F}, \mathcal{G} are R -torsion free. Note that the natural map

$$\left(\prod_{w \in G} \mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \right) \otimes_R K \longrightarrow \prod_{w \in G} (\mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \otimes_R K)$$

is injective, so we get an exact sequence:

$$0 \longrightarrow \mathrm{Hom}_{\mathcal{O}_{\hat{X}}}(\mathcal{F}, \mathcal{G}) \otimes_R K \longrightarrow \mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \otimes_R K \longrightarrow \prod_{w \in G} (\mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \otimes_R K)$$

On the other hand, we have an exact sequence

$$\begin{aligned} 0 \longrightarrow \mathrm{Hom}_{\mathrm{Coh}^m(q, G)}(q^*\mathcal{F}, q^*\mathcal{G}) &\longrightarrow \Gamma(\hat{Y}, \underline{\mathrm{Hom}}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \otimes_R K) \\ &\longrightarrow \prod_{w \in G} \Gamma(\hat{Y}, \underline{\mathrm{Hom}}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \otimes_R K) \end{aligned}$$

where the last map is again given by $w^* - \mathrm{id}^*$. Since

$$\mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \otimes_R K \subseteq \Gamma(\hat{Y}, \underline{\mathrm{Hom}}_{\mathcal{O}_{\hat{Y}}}(q^*\mathcal{F}, q^*\mathcal{G}) \otimes_R K)$$

is the subspace consisting of elements s such that $\pi^n s$ can be lifted to $\mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^* \mathcal{F}, q^* G)$ for some $n \in \mathbb{N}$, we just have to show that if

$$s \in \Gamma(\hat{Y}, \underline{\mathrm{Hom}}_{\mathcal{O}_{\hat{Y}}}(q^* \mathcal{F}, q^* G) \otimes_R K)$$

such that $w^* s = s$ for all $w \in G$, then there exist n large such that $\pi^n s$ can be lifted to $\mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^* \mathcal{F}, q^* G)$.

Let $U \rightarrow X_0$ be a quasi-compact étale map which trivializes the G -torsor $Y_0 \rightarrow X_0$. Then the image V of $U \rightarrow Y_0$ is a quasi-compact open whose G -translates $\{V_w\}_{w \in G}$ cover Y_0 . Since V is quasi-compact, $\pi^n s|_V$ can be lifted to $\mathrm{Hom}_{\mathcal{O}_{\hat{Y}|_V}}(q^* \mathcal{F}|_V, q^* \mathcal{G}|_V)$. Now the condition $w^* s = s$ implies that $\pi^n s|_{V_w}$ can be lifted to $\mathrm{Hom}_{\mathcal{O}_{\hat{Y}|_{V_w}}}(q^* \mathcal{F}|_{V_w}, q^* \mathcal{G}|_{V_w})$. Thus $\pi^n s$ can be lifted to $\mathrm{Hom}_{\mathcal{O}_{\hat{Y}}}(q^* \mathcal{F}, q^* G)$. \square

Remark 1.12. The above pullback functor is functorial in the following sense. If there is a Cartesian diagram of locally Noetherian formal schemes over $\mathrm{Spf}(R)$

$$\begin{array}{ccc} \hat{Y}' & \xrightarrow{\beta} & \hat{Y} \\ q' \downarrow & \square & \downarrow q \\ \hat{X}' & \xrightarrow{\alpha} & \hat{X} \end{array}$$

where q, q' are G -torsors, and if \mathcal{F} is a coherent sheaf on \hat{X} and $(q^* \mathcal{F}, \{h_w\}_{w \in G})$ is the corresponding pair, then the pair corresponding to $\alpha^* \mathcal{F}$ is $(\beta^* q^* \mathcal{F}, \{\beta^* h_w\}_{w \in G})$.

1.3. The specialization map. We are in Setting 1.7 except that here we assume that R is of characteristic p . Let X be a proper scheme over $\mathrm{Spec}(R)$, and let \hat{X} be the formal scheme associated with the special fiber of $X \rightarrow \mathrm{Spec}(R)$. Suppose that the special fiber X_0 is connected and $\xi \in X_0$ is a geometric point. Set $X_K := X \times_{\mathrm{Spec}(R)} \mathrm{Spec}(K)$.

Theorem 1.13. *There is a natural equivalence of categories:*

$$\mathrm{Coh}(\hat{X}) \otimes_R K \rightarrow \mathrm{Coh}(X_K)$$

which is additive, monoidal and exact.

Proof. By Grothendieck's existence theorem we have an additive, monoidal and exact equivalence

$$\mathrm{Coh}(X) \otimes_R K \xrightarrow{\cong} \mathrm{Coh}(\hat{X}) \otimes_R K$$

On the other hand, since X is Noetherian the natural pullback functor

$$\mathrm{Coh}(X) \otimes_R K \longrightarrow \mathrm{Coh}(X_K)$$

is an equivalence. This completes the proof. \square

Construction 1.14. Let $\hat{X}^{(i)}$ denote the formal scheme obtained by the following Cartesian diagram

$$\begin{array}{ccc} \hat{X}^{(1)} & \longrightarrow & \hat{X} \\ \downarrow & \square & \downarrow \\ \mathrm{Spf}(R) & \xrightarrow{F_{\mathrm{Spf}(R)}} & \mathrm{Spf}(R) \end{array}$$

where $F_{\mathrm{Spf}(R)}$ denotes the absolute Frobenius of $\mathrm{Spf}(R)$. The universal property of pullback diagrams provides a map $\hat{X} \rightarrow \hat{X}^{(1)}$ over $\mathrm{Spf}(R)$. In other words, $\hat{X} \rightarrow \hat{X}^{(1)}$ is the *relative Frobenius* of $\hat{X}/\mathrm{Spf}(R)$. Then we can define the relative Frobenius maps $\hat{X}^{(i)} \rightarrow \hat{X}^{(i+1)}$ of $\hat{X}^{(i)}/\mathrm{Spf}(R)$ inductively for each $i \in \mathbb{N}$. Moreover, we have commutative diagrams

$$\begin{array}{ccc} \hat{Y}^{(i)} & \longrightarrow & \hat{Y}^{(i+1)} \\ q^{(i)} \downarrow & \square & \downarrow q^{(i+1)} \\ \hat{X}^{(i)} & \longrightarrow & \hat{X}^{(i+1)} \end{array}$$

which are Cartesian because $\hat{Y}^{(i)} \rightarrow \hat{X}^{(i)}$ are étale.

We want to construct the specialization functor

$$\mathrm{sp}_K: \mathrm{Rep}_K^{\mathrm{cts}}(\pi_1^{\mathrm{pro\acute{e}t}}(X_0, \xi)) \longrightarrow \mathrm{Fdiv}(X_K)$$

First we choose an equivalence from $\mathrm{Rep}_K^{\mathrm{cts}}(\pi_1^{\mathrm{pro\acute{e}t}}(X_0, \xi))$ to the category of finite dimensional representations on the standard vector spaces $K^{\oplus n}$ ($n \in \mathbb{N}$), so it is enough to construct from there. Given a continuous representation

$$\rho: \mathrm{Rep}_K^{\mathrm{cts}}(\pi_1^{\mathrm{pro\acute{e}t}}(X_0, \xi)) \rightarrow \mathrm{GL}(K^{\oplus n})$$

we can take the image $G \subseteq \mathrm{GL}(K^{\oplus n})$ of ρ , which is a discrete group. Let $w \in G$, then the action of w on $K^{\oplus n}$ defines an element h_w in

$$\mathrm{Hom}_{\mathcal{O}_{\hat{Y}^{(i)}}}(\mathcal{O}_{\hat{Y}^{(i)}} \otimes_R K^{\oplus n}, \mathcal{O}_{\hat{Y}^{(i)}} \otimes_R K^{\oplus n}) = \mathrm{Hom}_{\mathcal{O}_{\hat{Y}^{(i)}}}(\mathcal{O}_{\hat{Y}^{(i)}}^{\oplus n} \otimes_R K, \mathcal{O}_{\hat{Y}^{(i)}}^{\oplus n} \otimes_R K)$$

We can identify $\mathcal{O}_{\hat{Y}^{(i)}}^{\oplus n} \otimes_R K$ with $w^* \mathcal{O}_{\hat{Y}^{(i)}}^{\oplus n} \otimes_R K$, so h_w is indeed in

$$\mathrm{Hom}_{\mathcal{O}_{\hat{Y}^{(i)}}}(\mathcal{O}_{\hat{Y}^{(i)}}^{\oplus n} \otimes_R K, w^* \mathcal{O}_{\hat{Y}^{(i)}}^{\oplus n} \otimes_R K)$$

In this way, the pair $(\mathcal{O}_{\hat{Y}^{(i)}}^{\oplus n}, \{h_w\}_{w \in G})$ becomes a pair in $\mathrm{Coh}^m(q^{(i)}, G)$ which descends, via 1.11, to a coherent sheaf \mathcal{E}_i on $\hat{X}^{(i)}$, hence a coherent sheaf on $X_K^{(i)}$ by 1.13. By the functoriality of the descent 1.12 we get a natural isomorphism $\sigma_i: F_{X^{(i)K}/K}^* \mathcal{E}_{i+1} \xrightarrow{\cong} \mathcal{E}_i$ for each i . Thus $(\mathcal{E}_i, \sigma_i)_{i \in \mathbb{N}}$ form an object in $\mathrm{Fdiv}(X_K)$.

Remark 1.15. If X and the generic fiber X_K are connected schemes, and if $\xi \in X_0$ and $\eta \in X_K$, then we have a specialization map of the étale fundamental groups

$$(4) \quad \pi_1^{\acute{e}t}(X_K, \eta) \longrightarrow \pi_1^{\acute{e}t}(X, \eta) \simeq \pi_1^{\acute{e}t}(X, \xi) \longleftarrow \cong \pi_1^{\acute{e}t}(X_0, \xi)$$

which is defined up to a choice of a path from η to ξ . The specialization map induces a functor on the representations

$$\mathrm{Rep}_K^{\mathrm{cts}}(\pi_1^{\acute{e}t}(X_0, \xi)) \longrightarrow \mathrm{Rep}_K^{\mathrm{cts}}(\pi_1^{\acute{e}t}(X_K, \eta))$$

Let's make the functor more explicit. Given a finite dimensional continuous representation $\rho: \pi_1^{\acute{e}t}(X_0, \xi) \rightarrow \mathrm{GL}(K^{\oplus n})$. Let $G \subseteq \mathrm{GL}(K^{\oplus n})$ be the image of ρ . Then the surjection

$$\pi_1^{\acute{e}t}(X_0, \xi) \twoheadrightarrow G$$

corresponds to a pointed G -torsor

$$\begin{array}{ccc} & & Y_0 \\ & \nearrow & \downarrow \\ \xi & \longrightarrow & X_0 \end{array}$$

which corresponds, via Grothendieck's existence theorem, to the pointed G -torsor:

$$\begin{array}{ccc} & & Y \\ & \nearrow & \downarrow \\ \xi & \longrightarrow & X \end{array}$$

and then after a (functorial) choice of the dotted arrow:

$$\begin{array}{ccc} & & Y_K \\ & \nearrow \cdots & \downarrow \\ \eta & \longrightarrow & X_K \end{array}$$

The above pointed torsor defines a map

$$\pi_1^{\acute{e}t}(X_K, \eta) \longrightarrow G$$

which in turn gives a representation $\rho_K: \pi_1^{\acute{e}t}(X_K, \eta) \rightarrow \mathrm{GL}(K^{\oplus n})$.

Given a representation $\rho_K: \pi_1^{\text{ét}}(X_K, \eta) \rightarrow \text{GL}(K^{\oplus n})$, we can take the image G as before. Then the surjection $\pi_1^{\text{ét}}(X_K, \eta) \rightarrow G$ gives a G -torsor $q_K: Y_K \rightarrow X_K$ which extends to Cartesian diagrams:

$$(5) \quad \begin{array}{ccc} Y_K^{(i)} & \longrightarrow & Y_K^{(i+1)} \\ \downarrow & \square & \downarrow \\ X_K^{(i)} & \longrightarrow & X_K^{(i+1)} \end{array}$$

The representation of the finite group G on $K^{\oplus n}$ provides descent data of $\mathcal{O}_{Y_K}^{\oplus n}$ with respect to $Y_K^{(i)} \rightarrow X_K^{(i)}$ for each $i \in \mathbb{N}$ and they are compatible along the diagrams (5). This defines, by the functoriality of descent, an object in $\text{Fdiv}(X_K)$. Thus we have constructed a natural functor

$$\text{Rep}_K^{\text{cts}}(\pi_1^{\text{ét}}(X_K, \eta)) \longrightarrow \text{Fdiv}(X_K)$$

In conclusion, putting everything together with get a functor

$$\text{sp}_{\text{ét}}: \text{Rep}_K^{\text{cts}}(\pi_1^{\text{ét}}(X_0, \xi)) \longrightarrow \text{Fdiv}(X_K)$$

by the specialization of the étale fundamental group. Note that the assumption that X and X_K are connected and the choice of the point η are only to make sense of the corresponding fundamental groups, but if one directly start with a representation ρ in $\text{Rep}_K^{\text{cts}}(\pi_1^{\text{ét}}(X_0, \xi))$ and take its image then one can go through the whole construction of $\text{sp}_{\text{ét}}$ without referring to the étale fundamental group of X or X_K , not to say the choice of the path from η to ξ .

Theorem 1.16. *The following natural diagram of categories*

$$\begin{array}{ccc} \text{Rep}_K^{\text{cts}}(\pi_1^{\text{ét}}(X_0, \xi)) & \xrightarrow{\text{sp}_{\text{ét}}} & \text{Fdiv}(X_K) \\ \downarrow & & \downarrow = \\ \text{Rep}_K^{\text{cts}}(\pi_1^{\text{proét}}(X_0, \xi)) & \xrightarrow{\text{sp}_K} & \text{Fdiv}(X_K) \end{array}$$

is 2-commutative.

Proof. Indeed $\text{Rep}_K^{\text{cts}}(\pi_1^{\text{ét}}(X_0, \xi))$ is the full subcategory of $\text{Rep}_K^{\text{cts}}(\pi_1^{\text{proét}}(X_0, \xi))$ consisting of representations whose monodromy groups are finite. Now let's come back to 1.14 and we assume that G is finite. In this case we have a G -torsor $q_K: Y_K \rightarrow X_K$ at our disposal. Indeed, thanks to Grothendieck existence theorem, there is a G -torsor $q: Y \rightarrow X$ whose formal completion along the special fiber is $q: \hat{Y} \rightarrow \hat{X}$ and whose generic fiber is q_K . Applying 1.13 to $X_K, \hat{X}, Y_K, \hat{Y}$ and their fibred products we get an equivalence between $\text{Coh}^m(q, G)$ and the category of fpqc descent data of coherent sheaves along q_K , and it is clear that the meromorphic descent data along q and the fpqc descent data along q_K induced by the representation $G \subseteq \text{GL}(K^{\oplus n})$ correspond to each other under this equivalence. Thus they produce the same F -divided sheaves, whence the 2-commutativity of the diagram. \square

Lemma 1.17. *Let A be a flat finitely generated R -algebra. Suppose that A is a integral domain, and P is a maximal ideal of A containing the uniformizer π of R . If $A_P[\frac{1}{\pi}]$ is a field, then $A[\frac{1}{\pi}]$ is also a field.*

Proof. The conditions implies that if $Q \subseteq P$ is a non-zero prime ideal of A , then $\pi \in Q$. Consider the minimal prime ideals of the ideal (π) in A_P . If PA_P was not one of them, then there would be $a \in PA_P$ such that a is not contained in any of those minimal prime ideals of (π) . By Krull principal ideal theorem each minimal prime ideal I of (a) has height 1, but since it is non-zero, $\pi \in I$. Thus I has to be also a minimal prime ideal of (π) - a contradiction! So PA_P is a minimal prime ideal of (π) . This implies that $\text{ht}(P) = 1$, so by [11, 14.C, Theorem 23, p. 84] and the fact that R is universally catenary, the function field of $A[\frac{1}{\pi}]$ is finite over K , hence $A[\frac{1}{\pi}]$ a field. \square

Lemma 1.18. *If the residue field k of R is algebraically closed, and if X is flat over $\text{Spec}(R)$, then for any $\xi \in X_0(k)$ we can find a complete DVR R' which is finite flat over R and a R -map $\text{Spec}(R') \rightarrow X$ whose special point goes to ξ .*

Proof. We can find $\eta \in X_K$ which is a generalization of ξ (going down). Set $P_\eta \subseteq \mathcal{O}_{X,\xi}$ the prime ideal corresponding to η . Let $\eta' \in X_K$ be the point corresponding to a maximal ideal of $(\mathcal{O}_{X,\xi}/P_\eta)[\frac{1}{\pi}]$. Then by 1.17, $\eta' \in X_K$ is a closed point. Replacing η by η' we may assume that η is a closed point of X_K , then its residue field $\kappa(\eta)$ is a finite extension of K . By [15, 054F] there exists a DVR R' and a map $\text{Spec}(R') \rightarrow X$, whose special point goes to ξ and whose generic point goes to η . Moreover, R' can be chosen so that its function field is equal to $\kappa(\eta)$. Thus $R \rightarrow R'$ is a finite ring extension. \square

Let $\eta \in X_K(K)$ be a rational point, and let $\bar{\eta}$ be the composition:

$$\text{Spec}(\bar{K}) \rightarrow \text{Spec}(K) \rightarrow X_K$$

Recall that the pro-constant quotient $\pi^G(X_K/K, \eta)$ of the Nori-étale fundamental K -group scheme $\pi^E(X_K/K, \eta)$ ([18, Definition 4.5, p. 1361] or the gerbe version [17, Definition 4.1, p. 866]) is the quotient of the étale fundamental group $\pi_1^{\text{ét}}(X_K, \bar{\eta}) = \pi^G(X_K/K, \bar{\eta})$ classifying covers equipped with a K -rational point over η . We have fully faithful inclusions:

$$\text{Rep}_K^{\text{cts}}(\pi^G(X_K/K, x_K)) \subseteq \text{Rep}_K(\pi^E(X_K/K, x_K)) \subseteq \text{Fdiv}(X_K)$$

where the first inclusion is obtained by taking the finite dimensional representations of the K -group scheme $\pi^E(X_K/K, x_K)$ whose monodromy groups are constant K -group schemes, and the second inclusion is obtained by taking the essentially finite objects of $\text{Fdiv}(X_K)$ ([17, Theorem 5.8, p. 869, Theorem 6.23, p. 885]). The category $\text{Fdiv}(X_K)$ is Tannakian (see [17, Theorem I (3), p. 856]). If $\eta \in X_K(K)$ is a K -rational point, then we denote $\pi_1^{\text{Fdiv}}(X_K, \eta)$ the affine K -group scheme corresponding to $\text{Fdiv}(X_K)$ and the fiber functor induced by η .

Proposition 1.19. *Suppose that the residue field k of R is algebraically closed, X is flat over $\text{Spec}(R)$, and that X_K is connected. If there is a R -rational point $(\xi, \eta) \in X(R)$, then this point defines a path between ξ and $\bar{\eta}$ (indeed also η) and the map (4) factors as*

$$\pi_1^{\text{ét}}(X_K, \bar{\eta}) \twoheadrightarrow \pi^G(X_K/K, \eta) \longrightarrow \pi_1^{\text{ét}}(X_0, \xi)$$

Therefore, the functor $\text{sp}_{\text{ét}}$ factors as

$$\text{Rep}_K^{\text{cts}}(\pi_1^{\text{ét}}(X_0, \xi)) \longrightarrow \text{Rep}_K^{\text{cts}}(\pi^G(X_K/K, \eta)) \subseteq \text{Fdiv}(X_K)$$

By taking Tannakian dual we get a homomorphism of K -group schemes:

$$\pi_1^{\text{Fdiv}}(X_K, \eta) \longrightarrow \pi_1^{\text{ét}}(X_0, \xi)$$

Proof. In Remark 1.15 the pointed G -torsor on X_0

$$\begin{array}{ccc} & & Y_0 \\ & \nearrow & \downarrow \\ \text{Spec}(k) & \xrightarrow{\xi} & X_0 \end{array}$$

indeed corresponds to the pointed G -torsor on X

$$\begin{array}{ccc} & & Y \\ & \nearrow & \downarrow \\ \text{Spec}(R) & \xrightarrow{(\xi, \eta)} & X \end{array}$$

because R is now strictly Henselian so that all the finite étale covers splits completely. This then provides the pointed torsor over X_K

$$(6) \quad \begin{array}{ccc} & & Y_K \\ & \nearrow & \downarrow \\ \text{Spec}(K) & \xrightarrow{\eta} & X_K \end{array}$$

Therefore, the map $\pi_1^{\text{ét}}(X_K, \bar{\eta}) \rightarrow G$ corresponding to the pointed torsor (6) factors as

$$\pi_1^{\text{ét}}(X_K, \bar{\eta}) \rightarrow \pi^G(X_K/K, \eta) \rightarrow G$$

This provides the desired factorization. \square

If the residue field k of R is algebraically closed, and X is flat over R , then any point $\xi \in X_0$ generalizes to a closed point $\eta \in X_K$ (1.18). Set $L := \kappa(\eta)$. Then L/K is finite. Let R' be the integral closure of R in L . Then R' is a complete DVR with residue field k . The scheme $X' := X \otimes_R R'$ has an R' -rational point by [15, 054F]. Its special fiber is still X_0 , its general fiber is $X'_L = X_K \otimes_K L$, and $\eta_L := \eta \otimes_K L$ is an L -rational point.

Corollary 1.20. *If the residue field k of R is algebraically closed, and if X is flat geometrically connected over R , then there is a finite extension L of the function field K of R and a homomorphism of L -group schemes:*

$$\pi_1^{\text{Fdiv}}(X \otimes_R L, \eta_L) \rightarrow \pi_1^{\text{ét}}(X_0, \xi)$$

Proof. We apply 1.19 to the R' -scheme $X' := X \otimes_R R'$ discussed above. \square

2. THE PRO-ÉTALE FUNDAMENTAL GROUP

In this section, we generalize E. Lavanda's theorem [10, Theorem 1.17, p. 33] to more general schemes (see Theorem 2.20, Theorem 2.24 and Theorem 2.27). The core techniques we use here are similar to those in E. Lavanda's proof: (1) the technique of proper descent of étale morphisms [10, Proposition 1.16, p. 33] due to A. Grothendieck and D. Rydh; (2) a combinatorial method which turns out to be a folklore (see e.g. [4, 8.4.1, p. 333]).

2.1. Results about Noohi groups.

Lemma 2.1. *A Noohi group π is Hausdorff and has a basis of open neighborhoods of $e \in \pi$ given by subgroups.*

Proof. Let $\pi\text{-(Sets)}$ be the category of discrete sets with π -actions, and let $F_\pi : \pi\text{-(Sets)} \rightarrow (\text{Sets})$ be the forgetful functor. Since π is Noohi, the continuous map $\pi \rightarrow \text{Aut}(F_\pi)$ is an isomorphism. Let T denote the set of open subgroups of π . Then for each $U \subseteq \pi$ in T the coset π/U is a discrete set equipped with a continuous π -action. Let $\text{Aut}(\pi/U)$ be the automorphism group (with the compact-open topology) of π/U . Then $\text{Aut}(F_\pi)$ is a subgroup of $\prod_{U \in T} \text{Aut}(\pi/U)$ equipped with the subspace topology. Since each $\text{Aut}(\pi/U)$ is Hausdorff and has a basis of open neighborhoods of the unit given by subgroups, so is the product $\prod_{U \in T} \text{Aut}(\pi/U)$ and its topological subgroup $\text{Aut}(F_\pi) \cong \pi$. \square

Lemma 2.2. *Let $f, g : G \rightarrow \pi$ be two continuous maps of topological groups, where π is Hausdorff and has a basis of open neighborhoods of $e \in \pi$. If the induced functors $f^*, g^* : \pi\text{-(Sets)} \rightarrow G\text{-(Sets)}$ coincide, then $f = g$. In other words, $\text{Hom}(G, \pi) \rightarrow \text{Fun}(\pi\text{-(Sets)}, G\text{-(Sets)})$ is injective.*

Proof. Since π is Hausdorff and has a basis of open neighborhoods of $e \in \pi$, the intersection of all open subgroups of π is e . Therefore, it is enough to show that f is equal to g after composing with $\pi \rightarrow \pi/U$ for all open subgroups $U \subseteq \pi$. But $\pi/U \in \pi\text{-(Sets)}$, so its restrictions to $G\text{-(Sets)}$ via f and g must coincide. This implies exactly that $\pi \rightarrow \pi/U$ equalizes f and g . \square

The following is a generalization of [9, Lemma 2.51, p. 18] to not necessarily Hausdorff topological groups.

Lemma 2.3. *The natural inclusion*

$$(\text{Noohi groups}) \subseteq (\text{Topological groups})$$

has a left adjoint, which we will denote by $(-)^{\text{Noohi}}$.

Proof. Given $G \in (\text{Topological groups})$ the category $G\text{-(Sets)}$ is a tame infinite Galois category. Let $F: G\text{-(Sets)} \rightarrow (\text{Sets})$ be the forgetful functor, then $\text{Aut}(F)$ equipped with the compact-open topology is a Noohi group by [3, Theorem 7.2.5.1, p. 191]. By [3, Theorem 7.2.5.3, p. 191] the continuous map $\alpha: G \rightarrow \text{Aut}(F)$ identifies $\text{Aut}(F)\text{-(Sets)}$ and $G\text{-(Sets)}$. We put $G^{\text{Noohi}} := \text{Aut}(F)$. If $\gamma: G \rightarrow G'$ is a continuous homomorphism, where G' is Noohi, then by [3, Theorem 7.2.5.2, p. 191] the functor $G'\text{-(Sets)} \rightarrow G\text{-(Sets)}$ corresponds to a unique map $\beta: \text{Aut}(F) \rightarrow G'$. Thus by 2.2 we have $\beta \circ \alpha = \gamma$. This concludes the proof. \square

Definition 2.4. Let π be a Noohi group, and let $H \subseteq \pi$ be a subgroup. Denote $\langle H \rangle$ the smallest normal subgroup of π containing H . We define the *Noohi quotient of π by H* to be $(\pi/\langle H \rangle)^{\text{Noohi}}$. Let $\langle H \rangle^c$ be the closure of $\langle H \rangle$ in π . By [3, Proposition 7.1.5, p. 188] and 2.1, $(\pi/\langle H \rangle)^{\text{Noohi}}$ is the Raïkov completion of $\pi/\langle H \rangle^c$.

If $\{f_i, g_i\}_{i \in I}$ are elements of π , we often form the Noohi quotient $(\pi/\langle H \rangle)^{\text{Noohi}}$ with H being the subgroup generated by $\{f_i^{-1}g_i\}_{i \in I}$. We refer to this as the *Noohi quotient of π by the relations $\{f_i = g_i\}_{i \in I}$* .

Lemma 2.5. *Finite coproducts exist in the category of Noohi groups.*

Proof. It suffices to show that $\pi_1 \coprod^{\pi} \pi_2$ exists for any diagram $\pi_1 \xleftarrow{p} \pi \xrightarrow{q} \pi_2$ of Noohi groups. Indeed when π is trivial, $\pi_1 \coprod \pi_2$ is just the Noohi group corresponding to the tame infinite Galois category of discrete sets equipped with two independent actions from π_1 and π_2 . See [3, Example 7.2.6].

In general, $\pi_1 \coprod^{\pi} \pi_2$ is nothing but the Noohi quotient of $\pi_1 \coprod \pi_2$ by the relations $\{p(x) = q(x)\}_{x \in \pi}$. Alternatively, it can also be defined by the Noohi group corresponding to the tame infinite Galois category of discrete sets equipped with two actions from π_1 and π_2 which agree on π . \square

2.2. A construction of Van Kampen. The construction we are about to discuss, in the case of discrete groups, can be found in [4, 8.4.1, p. 333], where it is attributed to Van Kampen.

Let $s \geq 1$ and let $\pi, \pi', \pi''_1, \dots, \pi''_s$ be Noohi groups. Let $\psi_i: \pi''_i \rightarrow \pi, \phi_i: \pi''_i \rightarrow \pi'$ be continuous homomorphisms. We define a (discrete) group F by generators $\{u_{ij}: 1 \leq i, j \leq s\}$ and relations $u_{ii} = e, u_{ij}u_{jk} = u_{ik}$. Evidently, F is free of rank $s-1$ on the generators v_2, \dots, v_s , where $v_i := u_{1i}$ for $i = 2, \dots, s$. We put $v_1 = e$.

Lemma 2.6. *The following Noohi groups are isomorphic:*

- (i) *The Noohi coproduct of π' and F .*
- (ii) *The Noohi coproduct of s copies of π' , and one copy of F , then Noohi quotient by the relations $u_{ij}^{-1}[y]_i u_{ij} = [y]_j, 1 \leq i, j \leq s, y \in \pi'$. Here $[y]_i$ denote y regarded as an element in the i -th copy of π' .*

Proof. The isomorphism is characterized by $y \leftrightarrow [y]_1$ for $y \in \pi'$ and $g \leftrightarrow g$ for $g \in F$. The universal property of coproduct implies that there are homomorphisms in both directions matching these elements, and they are inverse to each other. \square

Lemma 2.7. *The following Noohi groups are isomorphic:*

- (i) *The Noohi coproduct of π and the group in 2.6 (i), Noohi quotient by the relations $\psi_i(a) = v_i^{-1}\phi_i(a)v_i, a \in \pi''_i, i = 1, \dots, s$.*
- (ii) *The Noohi coproduct of π and the group in 2.6 (ii), Noohi quotient by the relations $\psi_i(a) = [\phi_i(a)]_i, a \in \pi''_i, i = 1, \dots, s$.*
- (iii) *The Noohi coproduct of F with $\pi \coprod^{\pi''_1, \phi_1, \psi_1} \pi'$, Noohi quotient by the relations $\psi_i(a) = v_i^{-1}\phi_i(a)v_i, a \in \pi''_i, i = 2, \dots, s$.*

- (iv) *The Noohi coproduct of F and (the Noohi fiber coproduct of $\pi \rightarrow \pi \coprod_{\pi_1''^{\phi_1, \psi_1}} \pi', \dots, \pi \rightarrow \pi \coprod_{\pi_s''^{\phi_s, \psi_s}} \pi')$, Noohi quotient by the relations $u_{ij}^{-1}(e *_i y)u_{ij} = e *_j y$ for all $y \in \pi'$, $i, j = 1, \dots, s$. Here $e *_i y$ denote the product of $e \in \pi$ and $y \in \pi'$ in $\pi \coprod_{\pi_i''^{\phi_i, \psi_i}} \pi'$.*

Proof. The isomorphism between (i) and (ii) is induced by that of 2.6. The isomorphism between (i) and (iii) is rather obvious. The isomorphism between (iii) and (iv) is constructed in a way similar to that of 2.6. \square

Remark 2.8. We will denote the group in 2.7 as $\mathbf{VK}(\pi, \pi'; \pi_1'', \dots, \pi_s'')_{(\psi_1, \dots, \psi_s), (\phi_1, \dots, \phi_s)}$ and omit the homomorphisms in subscripts when there is no confusion. The description (i) is the one usually found in the literature, e.g. [4, 8.4.1, p. 333]. It relies on single out the index 1 and so does (iii). The descriptions (ii) and (iv) make it clear that the construction actually treats all indices in equal footing. This construction is very useful due to the next result.

Setting 2.9. To state the result, we work in the following situation with $s \geq 1$:

- Let (\mathcal{C}, F) , (\mathcal{D}, G) , $(\mathcal{E}_1, H_1), \dots, (\mathcal{E}_s, H_s)$ be tame infinite Galois categories.
- For $j = 1, \dots, s$, let $u_j : \mathcal{C} \rightarrow \mathcal{E}_j$ (resp. $v_j : \mathcal{D} \rightarrow \mathcal{E}_j$) be a functor such that $(\mathcal{C}, H_j \circ u_j)$ (resp. $(\mathcal{D}, H_j \circ v_j)$) is a tame infinite Galois category and $H_j \circ u_j$ is compatible with F (resp. $H_j \circ v_j$ is compatible with G) in the sense of [9, Definition 3.14].
- Recall that an object of the 2-fibre product ([15, 003R])

$$\mathcal{C} \times_{\mathcal{E}_1 \times \dots \times \mathcal{E}_s} \mathcal{D}$$

is a triple (X, Y, ϕ) , where $X \in \mathcal{C}$, $Y \in \mathcal{D}$, and ϕ is an isomorphism

$$(u_j(X))_{1 \leq j \leq s} \rightarrow (v_j(Y))_{1 \leq j \leq s}$$

in $\mathcal{E}_1 \times \dots \times \mathcal{E}_s$. We define a functor F' from this 2-fibre product to (Sets) by setting $F'(X, Y, \phi) = F(X)$.

- For $j = 1, \dots, s$, choose an isomorphism of fiber functors on \mathcal{C} : $H_j \circ u_j \xrightarrow{\cong} F$ and denote the composition $\pi_1(\mathcal{E}_j, H_j) \rightarrow \pi_1(\mathcal{C}, H_j \circ u_j) \rightarrow \pi_1(\mathcal{C}, F)$ by ψ_j . We also choose an isomorphism of fiber functors on \mathcal{D} : $H_j \circ v_j \xrightarrow{\cong} G$ and denote the composition $\pi_1(\mathcal{E}_j, H_j) \rightarrow \pi_1(\mathcal{D}, H_j \circ v_j) \rightarrow \pi_1(\mathcal{D}, G)$ by ϕ_j .

Proposition 2.10. *With the above setting, the pair*

$$(\mathcal{C} \times_{\mathcal{E}_1 \times \dots \times \mathcal{E}_s} \mathcal{D}, F')$$

is a tame infinite Galois category, whose fundamental group is isomorphic to

$$\mathbf{VK}(\pi_1(\mathcal{C}, F), \pi_1(\mathcal{D}, G); \pi_1(\mathcal{E}_1, H_1), \dots, \pi_1(\mathcal{E}_s, H_s))_{(\psi_1, \dots, \psi_s), (\phi_1, \dots, \phi_s)}.$$

Proof. Let us denote the pair in the proposition by (\mathcal{C}', F') . An object in \mathcal{C}' is an $(s+2)$ -tuple $(A, B, \lambda_1, \dots, \lambda_s)$, where $A \in \mathcal{C}$, $B \in \mathcal{D}$, and $\lambda_j : u_j(A) \xrightarrow{\cong} v_j(B)$. We will view this as an object (A, B, λ_1) of $\mathcal{C} \times_{\mathcal{E}_1} \mathcal{D}$, together with isomorphisms $(\lambda_2, \dots, \lambda_s)$ in $\mathcal{E}_2 \times \dots \times \mathcal{E}_s$.

By [3, Theorem 7.2.5], we may further describe a category equivalent to \mathcal{C}' using Noohi group actions. Indeed, \mathcal{C}' is equivalent to the category of s -tuples $(S, \lambda'_2, \dots, \lambda'_s)$, where S is a discrete Π_1 -set with $\Pi_1 := \pi_1(\mathcal{C}, F) \coprod_{\pi_1(\mathcal{E}_1, H_1), \psi_1, \phi_1} \pi_1(\mathcal{D}, G)$, and for $s = 2, \dots, s$, $\lambda'_s : S \rightarrow S$ is a bijection of $\pi_1(\mathcal{E}_j, H_j)$ -sets where the source (resp. target) has the structure of a $\pi_1(\mathcal{E}_j, H_j)$ -set via $u_j(A)$ (resp. $v_j(B)$). This exactly means: $\phi_j(a) \cdot \lambda'_j(s) = \lambda'_j(\psi_j(a) \cdot s)$ for all $a \in \pi_1(\mathcal{E}_j, H_j)$, $s \in S$. Therefore, \mathcal{C}' is exactly equivalently the category of discrete π -set, where π is the Noohi group given by the proposition, by 2.7 (iii). This completes the proof by [3, Example 7.2.2]. \square

Remark 2.11. With the notation of the proof, it is obvious that under the identification $\pi_1(\mathcal{C}', F') \simeq \pi$, the maps $\pi_1(\mathcal{C}, F) \rightarrow \pi_1(\mathcal{C}', F')$, $\pi_1(\mathcal{D}, G) \rightarrow \pi_1(\mathcal{C}', F')$ correspond to the canonical maps $\pi_1(\mathcal{C}, F) \rightarrow \pi$, $\pi_1(\mathcal{D}, G) \rightarrow \pi$.

Remark 2.12. More sophisticated forms of this sort of result can be found in [16, Corollary 5.3, p. 19] and [9, Corollary 3.18, p. 32]. However, the form given here is more readily applicable and is enough for almost all known applications, such as [10, Theorem 1.17, p. 33], which we generalize in 2.27. More applications are given in the next few subsections.

2.3. Van Kampen theorems.

Remark 2.13. From now on we assume that X is a locally topologically Noetherian scheme. Let $\text{Cov}(X)$ denote the category of *geometric covers of X* in the sense of [3, Definition 7.3.1, p. 192]. This construction gives a category fibered over the category of locally topological Noetherian schemes in an obvious way.

Let X_{red} denote the reduced induced scheme structure of X . Then by [2, Exposé VIII, Théorème 1.1, p. 247] or [7, Exposé IX, 4.10, p. 186] we can conclude that $\text{Cov}(X) \xrightarrow{\cong} \text{Cov}(X_{\text{red}})$ just as in [10, Lemma 1.15, p. 32]. Thus if X is a locally Noetherian connected scheme and $x \in X$ is a geometric point, then $\pi_1^{\text{proét}}(X_{\text{red}}, x) \rightarrow \pi_1^{\text{proét}}(X, x)$ is an isomorphism.

Lemma 2.14. *Let $f_1 : Z_1 \rightarrow X, f_2 : Z_2 \rightarrow X$ be monomorphisms of schemes. Assume that the induced map $Z_1 \amalg Z_2 \rightarrow X$ is a morphism of effective descent for $\text{Cov}(-)$. Then we have an equivalence given by the pullback functor:*

$$\text{Cov}(X) \simeq \text{Cov}(Z_1) \times_{\text{Cov}(Z)} \text{Cov}(Z_2)$$

where $Z = Z_1 \times_X Z_2$.

Proof. By the effectiveness of descent, an object of $\text{Cov}(X)$ amounts to a descent datum, which consists of $Y_i \in \text{Cov}(Z_i)$, $i = 1, 2$ together with $\varphi_{ij} : p_1^* Y_i \rightarrow p_2^* Y_j$ for $i, j \in \{1, 2\}$ satisfying the cocycle condition. The cocycle condition says that φ_{11} and φ_{22} are just identities (by the monomorphism assumption) and $\varphi_{12}, \varphi_{21}$ give isomorphisms $Y_1|_Z \simeq Y_2|_Z$, inverse to each other. Thus we get an object in the $\text{Cov}(Z_1) \times_{\text{Cov}(Z)} \text{Cov}(Z_2)$. It is routine to verify that this construction gives an equivalence. \square

Lemma 2.15. *The assumption of Lemma 2.14 is satisfied when Z_1, Z_2 are open subschemes of X such that $X = Z_1 \cup Z_2$.*

Proof. This is essentially the classical van Kampen theorem and an easy case of fpqc descent. Notice that here $Z = Z_1 \cap Z_2$. \square

Lemma 2.16. *The assumption of Lemma 2.14 is satisfied when X is locally Noetherian and Z_1, Z_2 are closed subschemes of X such that $X = Z_1 \cup Z_2$ set-theoretically.*

Proof. This follows from [10, Proposition 1.16, p. 33] applied to the map $\phi : Z_1 \amalg Z_2 \rightarrow X$, which relies on Rydh's work [14] and generalizes [7, IX Théorème 4.12]. \square

Lemma 2.17. *Let X be a locally Noetherian scheme. Let $Z \subseteq X$ be a closed subscheme. Consider a proper surjective map*

$$\tilde{X} \xrightarrow{f} X$$

Denote $Z \times_X \tilde{X}$ by \tilde{Z} . Suppose that the union of the image of the two closed immersions

$$\Delta_f : \tilde{X} \rightarrow \tilde{X} \times_X \tilde{X}, \quad \tilde{Z} \times_Z \tilde{Z} \rightarrow \tilde{X} \times_X \tilde{X}$$

is $\tilde{X} \times_X \tilde{X}$ set-theoretically. Then the pullback functor induces an equivalence of categories:

$$\text{Cov}(X) \xrightarrow{\cong} \text{Cov}(\tilde{X}) \times_{\text{Cov}(\tilde{Z})} \text{Cov}(Z).$$

Proof. Let us construct a quasi-inverse of the pullback functor. So we start with a triple (Y, W, ϕ) , where $Y \in \text{Cov}(\tilde{X})$, $W \in \text{Cov}(Z)$, and $\phi : Y \times_{\tilde{X}} \tilde{Z} \xrightarrow{\cong} W \times_Z \tilde{Z}$ is an isomorphism. The idea is to construct a descent datum of Y for the morphism f , then we get the desired object in $\text{Cov}(X)$ by applying [10, Proposition 1.16, p. 33].

To construct this descent datum, we need to work on $\tilde{X} \times_X \tilde{X}$ using the equivalence

$$\text{Cov}(\tilde{X} \times_X \tilde{X}) \cong \text{Cov}(\tilde{X}) \times_{\text{Cov}(\tilde{Z})} \text{Cov}(\tilde{Z} \times_Z \tilde{Z}),$$

which follows from Lemma 2.16 because the fiber product of the two closed immersions in the current lemma is the closed immersion $\tilde{Z} \xrightarrow{\Delta_f} \tilde{X} \times_X \tilde{X}$. Under this equivalence, $p_i^*(Y)$ corresponds to the triple $(Y, q_i^*(Y|_{\tilde{Z}}), \text{can}_i)$ for $i = 1, 2$. Here $p_i : \tilde{X} \times_X \tilde{X} \rightarrow \tilde{X}$, $q_i : \tilde{Z} \times_Z \tilde{Z} \rightarrow Z$ are projections, and can_i is the obvious canonical isomorphism. The desired descent datum $\lambda : p_1^*(Y) \rightarrow p_2^*(Y)$ corresponds to $(Y, q_1^*(Y|_{\tilde{Z}}), \text{can}_1) \rightarrow (Y, q_2^*(Y|_{\tilde{Z}}), \text{can}_2)$ given by $\text{id}_Y : Y \rightarrow Y$, $\varphi : q_1^*(Y|_{\tilde{Z}}) \rightarrow q_2^*(Y|_{\tilde{Z}})$, where φ is the canonical descent datum signifying the fact that $Y|_{\tilde{Z}}$ is isomorphic to $W \times_Z \tilde{Z}$ via ϕ .

We have to show that λ is indeed a descent datum. That is, considering the triple fibred product, the fibred product and the projections:

$$\begin{array}{ccc} \tilde{X} \times_X \tilde{X} \times_X \tilde{X} & \begin{array}{c} \xrightarrow{p_{12}} \\ \xrightarrow{p_{23}} \\ \xrightarrow{p_{13}} \end{array} & \tilde{X} \times_X \tilde{X} \\ & & \begin{array}{c} \xrightarrow{p_1} \\ \xrightarrow{p_2} \end{array} \\ & & \tilde{X} \end{array}$$

we have to show the cocycle condition $p_{23}^* \lambda \circ p_{12}^* \lambda = p_{13}^* \lambda$. Since the fibred product is covered by $\Delta_f(\tilde{X})$ and $\tilde{Z} \times_Z \tilde{Z}$, the triple fibred product is covered by the triple diagonal $\Delta_f^3(\tilde{X})$ and the closed subset

$$\tilde{Z} \times_Z \tilde{Z} \times_Z \tilde{Z}$$

Thanks to [15, OBTJ] it is enough to check the equality $p_{23}^* \lambda \circ p_{12}^* \lambda = p_{13}^* \lambda$ on $\Delta_f^3(\tilde{X})$ and $\tilde{Z} \times_Z \tilde{Z} \times_Z \tilde{Z}$ separately. On $\Delta_f^3(\tilde{X})$ all the pullbacks of λ are identities, so there is nothing to check. On $\tilde{Z} \times_Z \tilde{Z} \times_Z \tilde{Z}$ the equality holds because $\lambda|_{\tilde{Z} \times_Z \tilde{Z}} = \varphi$ is a descent datum for $\tilde{Z} \rightarrow Z$. \square

2.4. Schemes with connected singularity.

Conventions 2.18. Let X be a Nagata quasi-compact scheme, and let η_1, \dots, η_n be the generic points of X . Then we denote $f : \tilde{X} \rightarrow X$ the relative normalization of the map

$$\coprod_{1 \leq i \leq n} \text{Spec}(\kappa(\eta_i)) \rightarrow X$$

in the sense of [15, OBAK]. According to [15, OAVK], f is finite (and surjective). Moreover, according to [15, O35I], the map $f : \tilde{X} \rightarrow X$ factors through the reduced induced structure $X_{\text{red}} \subseteq X$, and the map $\tilde{X} \rightarrow X_{\text{red}}$ is the normalization of X_{red} relative to the disjoint union of the spectrums of the generic points. See [15, O359] for examples of Nagata schemes.

Assume further that X is a J-2 scheme ([15, O7R2]; see [15, O7R5] for examples of J-2 schemes). Then the singular locus $Z \subseteq X$ is closed. We give Z the reduced induced structure and put $\tilde{Z} := Z \times_X \tilde{X}$. Now it is easy to see that all the hypotheses of Lemma 2.17 are satisfied.

By [15, O35A], [15, O7R4] and 2.13, we may directly assume that X is reduced in the proof of the following theorems.

Theorem 2.19. *Let X be a connected Nagata Noetherian J-2 scheme. Let X_1, \dots, X_n be its irreducible components. Let $Z \subseteq X$ be the singular locus of X_{red} . Let \tilde{X}_i be the connected component of \tilde{X} corresponding to X_i , and put $\tilde{Z} := \tilde{X} \times_X Z$. For each $1 \leq i \leq n$, suppose $\tilde{X}_i \cap \tilde{Z} = Z_{i1} \coprod \dots \coprod Z_{in_i}$ is a decomposition. Then the pullback functor*

$$\text{Cov}(X) \xrightarrow{\cong} \prod_{\text{Cov}(Z)}^{1 \leq i \leq n} \left(\text{Cov}(\tilde{X}_i) \times_{\text{Cov}(Z_{i1}) \prod \dots \prod \text{Cov}(Z_{in_i})} \text{Cov}(Z) \right)$$

is an equivalence.

Proof. By 2.17 the pullback map

$$\text{Cov}(X) \xrightarrow{\cong} \left(\text{Cov} \left(\prod_{1 \leq i \leq n} \tilde{X}_i \right) \times_{\text{Cov}(\prod_{i,j} Z_{ij})} \text{Cov}(Z) \right) = \left[\left(\prod_{1 \leq i \leq n} \text{Cov}(\tilde{X}_i) \right) \times_{\prod_{i,j} \text{Cov}(Z_{ij})} \text{Cov}(Z) \right]$$

is an equivalence. Then one checks easily that the later is the same as the right hand side in the statement of the theorem. \square

Now we come back to fundamental groups. Assumptions and notations are as above, and we assume in addition that Z is connected. Choose geometric points $x \in Z$, $x_{ij} \in Z_{ij}$ and isomorphisms of fiber functors $F_x \simeq F_{f(x_{ij})}$, hence isomorphisms $\pi_1^{\text{proét}}(Z, x) \simeq \pi_1^{\text{proét}}(Z, f(x_{ij}))$. We denote ϕ_{ij} the map

$$\pi_1^{\text{proét}}(Z_{ij}, x_{ij}) \longrightarrow \pi_1^{\text{proét}}(Z, f(x_{ij})) \xrightarrow{\simeq} \pi_1^{\text{proét}}(Z, x)$$

induced by $Z \subseteq X$, and ψ_{ij} the map

$$\pi_1^{\text{proét}}(Z_{ij}, x_{ij}) \rightarrow \pi_1^{\text{proét}}(\tilde{X}_i, x_{ij}) \xrightarrow{\simeq} \pi_1^{\text{ét}}(\tilde{X}_i, x_{i1})$$

obtained by choosing isomorphisms of fiber functors $F_{x_{i1}} \simeq \cdots \simeq F_{x_{in_i}}$, therefore isomorphisms $\pi_1^{\text{proét}}(\tilde{X}_i, x_{i1}) \simeq \cdots \simeq \pi_1^{\text{proét}}(\tilde{X}_i, x_{in_i})$. Thanks to the equivalence between the category of tame infinite Galois categories with fiber functors and the category Noohi groups, we get

Theorem 2.20. *There is an isomorphism of topological groups*

$$\pi_1^{\text{proét}}(X, x) \simeq \coprod_{1 \leq i \leq n}^{\pi_1^{\text{proét}}(Z, x)} \mathbf{VK}(\pi_1^{\text{ét}}(\tilde{X}_i, x_{i1}), \pi_1^{\text{proét}}(Z, x); \pi_1^{\text{proét}}(Z_{i1}, x_{i1}), \dots, \pi_1^{\text{proét}}(Z_{in_i}, x_{in_i}))$$

Moreover, the canonical maps $\pi_1^{\text{proét}}(Z, x) \rightarrow \pi_1^{\text{proét}}(X, x)$ and $\pi_1^{\text{ét}}(\tilde{X}_i, x_{i1}) \rightarrow \pi_1^{\text{proét}}(X, f(x_{i1})) \simeq \pi_1^{\text{proét}}(X, x)$ correspond, via this isomorphism, to the projections in the coproduct.

Proof. This is immediate from 2.10 and 2.19. \square

2.5. Disconnected singularities: a dévissage. Next, we deal with the case when the singular locus is not connected.

Setting 2.21. Let X be a connected Nagata Noetherian J-2 scheme. Let X_1, \dots, X_n be its irreducible components. Let $Z_j \subseteq X$ ($1 \leq j \leq m$) be the connected components of the singular locus Z of X_{red} . For $j = 1, \dots, m$, set

$$T_j := \left(\bigcup_{X_i \cap Z_j \neq \emptyset} X_i \right) \setminus \left(\bigcup_{t \neq j} Z_t \right).$$

We notice that if $m = 0$, then X is regular and $n = 1$. From now on we assume $m \geq 1$.

Lemma 2.22. *With the above setting,*

- (i) Each T_j is connected and open in X .
- (ii) $X = T_1 \cup \cdots \cup T_m$.
- (iii) For each $1 \leq j \leq m$, $Z_j \subset T_j$ and Z_j is disjoint from T_t for $t \neq j$.
- (iv) There exists k such that $\bigcup_{t \neq k} T_t$ is connected, $1 \leq k \leq m$.

Proof. (i) Assume $X_i \cap Z_j \neq \emptyset$ and $X_s \cap Z_j = \emptyset$. Then $X_i \cap X_s$, being a subset of the singular locus, lies in $\bigcup_{t \neq j} Z_t$. Therefore, X_s is disjoint from T_j . Thus

$$T_j = \left[\left(\bigcup_{X_i \cap Z_j \neq \emptyset} X_i \right) \setminus \left(\bigcup_{X_s \cap Z_j = \emptyset} X_s \right) \right] \setminus \left(\bigcup_{t \neq j} Z_t \right) = \left[X \setminus \left(\bigcup_{X_i \notin S_j} X_i \right) \right] \setminus \left(\bigcup_{t \neq j} Z_t \right)$$

is open in X . Since T_j is the union of the connected sets Z_j and $X_i \setminus (\bigcup_{t \neq j} Z_t)$ (over those i such that $X_i \cap Z_j \neq \emptyset$), and each of the latter intersects with Z_j , it is clear that T_j is connected.

(ii) Let $x \in X$. Assume $x \in Z$, say $x \in Z_j$. Then it is clear that $x \in T_j$. Assume $x \notin Z$, say $x \in X_i$. Take any j such that $X_i \cap Z_j \neq \emptyset$ (such j exists; otherwise X_i is a regular component and we have $n = 1$ and $m = 0$), then it is clear $x \in T_j$.

(iii) is obvious.

(iv) We claim that we can find a permutation (j_1, \dots, j_m) of $\{1, \dots, m\}$ such that $T_{j_1} \cup \cdots \cup T_{j_r}$ is connected for all $1 \leq r \leq m$. Then the desired result follows by taking $k = j_m$. To prove the claim, we pick j_1 arbitrarily. Suppose that we have picked j_1, \dots, j_r with $r < m$, then it suffices to take j_{r+1} such that $(T_{j_1} \cup \cdots \cup T_{j_r}) \cap T_{j_{r+1}} \neq \emptyset$. Indeed if such j_{r+1} could not be found, we

would deduce that $T_{j_1} \cup \cdots \cup T_{j_r}$ is open and closed in the connected scheme X by (ii). This proves the claim. \square

Theorem 2.23. *With the above setting, assume moreover that $T'_1 := T_2 \cup \cdots \cup T_m$ is connected. Each connected component of $T_1 \cap T'_1$ is a connected component of the regular scheme $X \setminus Z$. Let D_1, \dots, D_s be the connected components of $T_1 \cap T'_1$. Then the pullback functor*

$$\mathrm{Cov}(X) \xrightarrow{\simeq} \mathrm{Cov}(T_1) \prod_{\prod_{1 \leq t \leq s} \mathrm{Cov}(D_t)} \mathrm{Cov}(T'_1)$$

is an equivalence.

Proof. By Lemma 2.22 (iii), $T_1 \cap T'_1 = (T_1 \setminus Z) \cap (T'_1 \setminus Z)$. Since $T_1 \setminus Z$ and $T'_1 \setminus Z$ are both unions of some of the irreducible components of the regular scheme $X \setminus Z$, the first statement is clear. The second statement follows readily from 2.15. \square

Again, we come back to fundamental groups. With the above setting, we may and do assume that $T'_1 := T_2 \cup \cdots \cup T_m$ is connected by Lemma 2.22 (iv). Let $d_t \in D_t$. Choose isomorphisms of fiber functors $F_{d_1} \simeq \cdots \simeq F_{d_s}$ on T_1 , and those $F_{d_1} \simeq \cdots \simeq F_{d_s}$ on T'_1 , so we get

$$\begin{aligned} \pi_1^{\mathrm{pro\acute{e}t}}(T_1, d_1) &\simeq \cdots \simeq \pi_1^{\mathrm{pro\acute{e}t}}(T_1, d_s) \\ \pi_1^{\mathrm{pro\acute{e}t}}(T'_1, d_1) &\simeq \cdots \simeq \pi_1^{\mathrm{pro\acute{e}t}}(T'_1, d_s) \end{aligned}$$

Theorem 2.24. *There is an isomorphism*

$$\pi_1^{\mathrm{pro\acute{e}t}}(X, d_1) \simeq \mathbf{VK}(\pi_1^{\mathrm{pro\acute{e}t}}(T_1, d_1), \pi_1^{\mathrm{pro\acute{e}t}}(T'_1, d_1); \pi_1^{\acute{e}t}(D_1, d_1), \dots, \pi_1^{\acute{e}t}(D_s, d_s)),$$

where the relevant maps are $\phi_t : \pi_1^{\acute{e}t}(D_t, d_t) \rightarrow \pi_1^{\mathrm{pro\acute{e}t}}(T_1, d_t) \simeq \pi_1^{\mathrm{pro\acute{e}t}}(T_1, d_1)$ and $\psi_t : \pi_1^{\acute{e}t}(D_t, d_t) \rightarrow \pi_1^{\mathrm{pro\acute{e}t}}(T'_1, d_t) \simeq \pi_1^{\mathrm{pro\acute{e}t}}(T'_1, d_1)$. Moreover, the canonical maps $\pi_1^{\mathrm{pro\acute{e}t}}(T_1, d_1) \rightarrow \pi_1^{\mathrm{pro\acute{e}t}}(X, d_1)$ and $\pi_1^{\mathrm{pro\acute{e}t}}(T'_1, d_1) \rightarrow \pi_1^{\mathrm{pro\acute{e}t}}(X, d_1)$ correspond, via this isomorphism, to the projections of the coproduct.

Proof. This follows immediately from 2.23 and 2.10. \square

Remark 2.25. Observe that the singular locus of T_1 is Z_1 and that of T'_1 is $Z_2 \cup \cdots \cup Z_m$. Therefore, the above theorem provides an inductive way to compute pro-étale fundamental groups in terms of pro-étale fundamental groups of connected schemes with connected singular locus. The latter is already handled by 2.20. By further performing an induction on the dimension of the singular locus, we can describe the pro-étale fundamental group of an arbitrary (Nagata, J-2, and Noetherian) connected scheme in terms of étale fundamental groups of normal schemes.

Corollary 2.26. *Let \mathcal{C} be the smallest class of Noohi groups such that*

- \mathcal{C} contains the étale fundamental groups of connected normal schemes.
- \mathcal{C} contains discrete free groups of finite rank.
- \mathcal{C} is closed under fiber coproducts and quotients.

Then the pro-étale fundamental group of a Nagata, J-2, Noetherian connected scheme lies in \mathcal{C} .

2.6. An example. As an example, we will compute the pro-étale fundamental group of an arbitrary connected curve over a separably closed field, generalizing [10, Theorem 1.17], who treated the stable case. In fact, we can handle more than curves. Let's resume 2.18 and 2.21. The key assumption that we now make is the following:

The pro-étale fundamental group of every connected component of Z or \tilde{Z} is trivial.

This condition is satisfied if Z is a disjoint union of spectrums of separably closed fields, e.g. when X is a curve over a separably closed field. Let's choose $x \in X$, $x_i \in \tilde{X}_i$, and paths $F_x \simeq F_{f(x_1)} \simeq \cdots \simeq F_{f(x_n)}$. Then we have

Theorem 2.27. *With the above setting, $\pi_1^{\mathrm{pro\acute{e}t}}(X, x)$ is the Noohi coproduct of $\pi_1^{\acute{e}t}(\tilde{X}_i, x_i)$, $i = 1, \dots, n$, and a discrete free group of rank $\tilde{m} - m - n + 1$, where m denotes the number of connected components of Z and \tilde{m} denotes that of \tilde{Z} .*

Proof. It is a nice exercise to prove the theorem by following the route provided by 2.24 and proceed by induction on the number m . However, it is easier to use the following argument. We will apply Theorem 2.19, and argue as in the proof of Proposition 2.10. Write the connected components of Z as Z_1, \dots, Z_m , and those of \tilde{Z} as $\tilde{Z}_1, \dots, \tilde{Z}_{\tilde{m}}$. An object of $\text{Cov}(\tilde{X}) \times_{\text{Cov}\tilde{Z}} \text{Cov}(Z)$ is $(\{A_i\}_{i=1}^n, \{B_j\}_{j=1}^m, \{\lambda_k\}_{k=1}^{\tilde{m}})$, where A_i is a $\pi_1(\tilde{X}_i, x_i)$ -set, B_j is a set, and λ_k is a bijection from $A_{i(k)}$ to $B_{j(k)}$ if $\tilde{Z}_k \subset \tilde{X}_{i(k)}$ and $f(\tilde{Z}_k) = Z_{j(k)}$.

Using that X is connected, one can show that we can choose indices k_1, \dots, k_{n+m-1} so that $\lambda_{k_1}, \dots, \lambda_{k_{n+m-1}}$ identify $A_1, \dots, A_n, B_1, \dots, B_m$ with each other, and hence we will use A_1 to represent all of them. The set A_1 now carries the action of $\pi_1(\tilde{X}_i, x_i)$, $i = 1, \dots, n$, together with the action of the remaining λ_k 's, now regarded as bijections from A_1 to itself. There are $\tilde{m} - n - m + 1$ of them. This description of $\text{Cov}(\tilde{X}) \times_{\text{Cov}\tilde{Z}} \text{Cov}(Z)$ easily implies the theorem. \square

2.7. Discrete representations. Let $\lambda: \pi \rightarrow G$ be a continuous homomorphism of topological groups. We say that λ is *discrete* if it factors as $\pi \rightarrow G' \rightarrow G$ with G' discrete, or equivalently, $\ker(\lambda)$ is open in π . When π is a fundamental group, it is often interesting to know the discrete homomorphisms.

Lemma 2.28. *Let π_1 and π_2 be Noohi groups and let $\pi_1 \coprod \pi_2$ be their Noohi coproduct. Let $\lambda: \pi_1 \coprod \pi_2 \rightarrow G$ be a continuous homomorphism corresponding to $\lambda_1: \pi_1 \rightarrow G$, $\lambda_2: \pi_2 \rightarrow G$. Then λ is discrete if and only if λ_1 and λ_2 are both discrete.*

Proof. The “only if” part is obvious. Assume that both λ_1 and λ_2 are discrete. Say λ_i factors as $\pi_i \rightarrow G'_i \rightarrow G$. Then clearly λ factors as $\pi_1 \coprod \pi_2 \rightarrow G'_1 \coprod G'_2 \rightarrow G$, where $G'_1 \coprod G'_2$ is the coproduct of G'_1 and G'_2 as topological groups. Since $G'_1 \coprod G'_2$ is discrete, λ is discrete. \square

Lemma 2.29. *Let π be a Noohi group and let $\pi' := (\pi/\langle H \rangle)^{\text{Noohi}}$ be a Noohi quotient of π . Consider a commutative diagram of continuous homomorphisms with G Hausdorff:*

$$\begin{array}{ccc} \pi & & \\ \downarrow & \searrow \lambda & \\ \pi' & \xrightarrow{\lambda'} & G \end{array}$$

Then λ is discrete if and only if λ' is discrete.

Proof. Again the “only if” part is obvious. Assume that λ is discrete and factorizes as $\pi \rightarrow G' \xrightarrow{\phi} G$. We may assume that $G' \rightarrow G$ is injective. Then $\ker(\pi \rightarrow G') = \ker(\pi \rightarrow G) \supset H$. Therefore, $\pi \rightarrow G'$ factorizes as $\pi \rightarrow \pi/\langle H \rangle^c \rightarrow G'$ in the category of topological groups. Applying the functor $(-)^{\text{Noohi}}$, we get $\pi \rightarrow \pi' \xrightarrow{\psi} G'$. It remains to show $\phi \circ \psi = \lambda'$.

By construction $\phi \circ \psi$ and λ' agree on the image M of π , which is a dense subset of π' . The map $(\phi \circ \psi, \lambda'): \pi' \rightarrow G \times G$ sends M to the diagonal of $G \times G$, which is closed as G is Hausdorff. We conclude that the whole image of $(\phi \circ \psi, \lambda')$ lies in the diagonal. This shows: $\phi \circ \psi = \lambda'$. \square

Lemma 2.30. *Suppose $g: Y' \rightarrow Y$ is a finite surjective map of normal integral schemes and $y' \in Y'$ is a geometric point. Set $y := g(y')$. Consider a commutative diagram of continuous homomorphisms:*

$$\begin{array}{ccc} \pi_1^{\text{proét}}(Y', y') & & \\ \downarrow & \searrow \lambda' & \\ \pi_1^{\text{proét}}(Y, y) & \xrightarrow{\lambda} & G \end{array}$$

Then λ is discrete if and only if λ' is discrete.

Proof. Indeed, the following commutative diagram:

$$\begin{array}{ccc} \mathrm{Gal}(K(Y')) & \hookrightarrow & \mathrm{Gal}(K(Y)) \\ \downarrow & & \downarrow \\ \pi_1^{\acute{e}t}(Y', y') & \longrightarrow & \pi_1^{\acute{e}t}(Y, y) \end{array}$$

implies that $\pi_1^{\mathrm{pro}\acute{e}t}(Y', y') \simeq \pi_1^{\acute{e}t}(Y', y') \rightarrow \pi_1^{\acute{e}t}(Y, y) \simeq \pi_1^{\mathrm{pro}\acute{e}t}(Y, y)$ is open. Thus $\mathrm{Ker}(\lambda)$ is open if and only if $\mathrm{Ker}(\lambda')$ is open. \square

Proposition 2.31. *Let X be a connected Nagata Noetherian J -2 scheme, and let $x \in X$ be a geometric point. Let $\{\eta_i\}_{1 \leq i \leq n}$ be the set of all generic points of X , and let $A := \prod_{1 \leq i \leq n} K_i$, where each K_i is a finite extension of $\kappa(\eta_i)$. Let \tilde{X}_A denote the normalization of X inside $\mathrm{Spec}(A)$. Let*

$$\lambda: \pi_1^{\mathrm{pro}\acute{e}t}(X, x) \longrightarrow G$$

be a continuous map of Hausdorff topological groups. Then λ is discrete if and only if the induced map $\pi_1^{\mathrm{pro}\acute{e}t}(U, u) \rightarrow G$ is discrete for each connected component U of \tilde{X}_A and each geometric point $u \in U$.

Proof. We just have to show the “if” direction. By 2.30, we may assume that $K_i = \kappa(\eta_i)$ for all i , i.e. $\tilde{X}_A = \tilde{X}$. By 2.24, 2.28 and 2.29, we may do an induction to reduce the proposition to the case where the singular locus of $Z \subseteq X_{\mathrm{red}}$ is connected. So we are in the situation of Lemma 2.20.

We want to show that the induced map $\pi_1^{\mathrm{pro}\acute{e}t}(Z, x) \rightarrow G$ is discrete. This would finish the proof by applying 2.28 and Lemma 2.29 again.

There is a ring $C := \prod_{j \in J} L_j$, where each L_j is a finite extension of the residue field of a generic point of Z , such that the normalization \tilde{Z}_C of Z in C lifts the inclusion $Z \hookrightarrow X$ to a map $\tilde{Z}_C \rightarrow \tilde{X}$. Thanks to the lifting, all the maps from the pro-étale fundamental groups of the connected components of \tilde{Z}_C to G factor through a discrete quotient. Since the dimension of Z is smaller than that of X , we complete our proof by induction. \square

Remark 2.32. Proposition 2.31 can also be proved using [9, Remark 3.20]. Indeed, after we have reduced to the case $\tilde{X}_A = \tilde{X}$, we can directly apply 2.28 and 2.29 together with *loc. cit.* to finish the proof, because in *loc. cit.* $\pi_1^{\mathrm{pro}\acute{e}t}(X, x)$ is written as a Noohi quotient of the Noohi coproduct of the fundamental groups of the \tilde{X}_i s and a free group.

ACKNOWLEDGEMENTS

The first named author’s work was a part of the project “Kapibara” supported by the funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 802787).

The third named author would like to thank the Direct Grant (No. 4053343) of the College of Science of The Chinese University of Hong Kong for support.

REFERENCES

- [1] Piotr Achinger, Marcin Lara, and Alex Youcis. *Specialization for the pro-étale fundamental group*. 2021. arXiv: [2107.06761](https://arxiv.org/abs/2107.06761) [math.AG].
- [2] Michael Artin, Alexander Grothendieck, and Jean-Louis Verdier. *Theorie de Topos et Cohomologie Etale des Schemas I, II, III*. Vol. 269, 270, 305. Lecture Notes in Mathematics. Springer, 1971.
- [3] Bhargav Bhatt and Peter Scholze. “The pro-étale topology for schemes”. In: *Astérisque* 369 (2015), pp. 99–201. ISSN: 0303-1179.
- [4] Ronald Brown. *Topology and groupoids*. Third edition of it Elements of modern topology [McGraw-Hill, New York, 1968; MR0227979], With 1 CD-ROM (Windows, Macintosh and UNIX). BookSurge, LLC, Charleston, SC, 2006, pp. xxvi+512. ISBN: 1-4196-2722-8.
- [5] David Gieseker. “Stable vector bundles and the Frobenius morphism”. In: *Ann. Sci. École Norm. Sup. (4)* 6 (1973), pp. 95–101. ISSN: 0012-9593. URL: http://www.numdam.org/item?id=ASENS_1973_4_6_1_0.

- [6] A. Grothendieck and J. A. Dieudonné. *Eléments de géométrie algébrique. I*. Vol. 166. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 1971, pp. ix+466. ISBN: 3-540-05113-9; 0-387-05113-9.
- [7] Alexander Grothendieck. *Revêtements étales et groupe fondamental (SGA 1)*. Vol. 224. Lecture notes in mathematics. Springer-Verlag, 1971.
- [8] Robin Hartshorne. *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977, pp. xvi+496. ISBN: 0-387-90244-9.
- [9] Marcin Lara. “Homotopy Exact Sequence for the Pro-Étale Fundamental Group”. PhD thesis. 2019. URL: <http://dx.doi.org/10.17169/refubium-2773>.
- [10] Elena Lavanda. “Specialization map between stratified bundles and pro-étale fundamental group”. In: *Adv. Math.* 335 (2018), pp. 27–59. ISSN: 0001-8708. DOI: [10.1016/j.aim.2018.06.013](https://doi.org/10.1016/j.aim.2018.06.013). URL: <https://doi.org/10.1016/j.aim.2018.06.013>.
- [11] Hideyuki Matsumura. *Commutative algebra*. Second. Vol. 56. Mathematics Lecture Note Series. Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980, pp. xv+313. ISBN: 0-8053-7026-9.
- [12] Arthur Ogus. “F-isocrystals and de Rham cohomology. II: Convergent isocrystals.” English. In: *Duke Math. J.* 51 (1984), pp. 765–850. ISSN: 0012-7094; 1547-7398/e. DOI: [10.1215/S0012-7094-84-05136-6](https://doi.org/10.1215/S0012-7094-84-05136-6).
- [13] Valentina Di Proietto, Fabio Tonini, and Lei Zhang. *A crystalline incarnation of Berthelot’s conjecture and Künneth formula for isocrystals*. To appear in *Journal of Algebraic Geometry*. 2021.
- [14] David Rydh. “Submersions and effective descent of étale morphisms”. en. In: *Bulletin de la Société Mathématique de France* 138.2 (2010), pp. 181–230. DOI: [10.24033/bsmf.2588](https://doi.org/10.24033/bsmf.2588). URL: <http://www.numdam.org/articles/10.24033/bsmf.2588/>.
- [15] The Stacks project authors. *The Stacks project*. <https://stacks.math.columbia.edu>. 2019.
- [16] Jakob Stix. “A general Seifert–Van Kampen theorem for algebraic fundamental groups”. In: *Publ. Res. Inst. Math. Sci* 42.3 (2006), pp. 763–786.
- [17] Fabio Tonini and Lei Zhang. “Algebraic and Nori fundamental gerbes”. In: *J. Inst. Math. Jussieu* 18.4 (2019), pp. 855–897. ISSN: 1474-7480. DOI: [10.1017/s147474801700024x](https://doi.org/10.1017/s147474801700024x). URL: <https://doi.org/10.1017/s147474801700024x>.
- [18] Lei Zhang. “Nori’s fundamental group over a non algebraically closed field”. In: *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 18.4 (2018), pp. 1349–1394. ISSN: 0391-173X.

MARCIN LARA, INSTYTUT MATEMATYCZNY PAN, ŚNIADECKICH 8, WARSAW, POLAND
 Email address: marcin.lara@impan.pl

JIU-KANG YU, THE CHINESE UNIVERSITY OF HONG KONG, DEPARTMENT OF MATHEMATICS, SHATIN, NEW TERRITORIES, HONG KONG
 Email address: jkyu@ims.cuhk.edu.hk

LEI ZHANG, THE CHINESE UNIVERSITY OF HONG KONG, DEPARTMENT OF MATHEMATICS, SHATIN, NEW TERRITORIES, HONG KONG
 Email address: lzhang@math.cuhk.edu.hk