

March, 2021

On the D_n Spin Vertex Models for Odd n

DORON GEPNER

*Department of Particle Physics and Astrophysics, Weizmann Institute,
Rehovot 76100, Israel*

ABSTRACT

Solvable vertex models in two dimensions are of importance in conformal field theory, phase transitions and integrable models. We consider here the D_n spin vertex models, for n which is odd. The models involve also the anti-spinor representation. We describe here the Boltzmann weights for these representations using crossing symmetry from the previously known spinor representations. For calculation reasons we limit ourself to the $n = 3$ and $n = 5$ cases, which are described explicitly.

Solvable vertex models in two dimensions are of interest to test such notions as universality, conformal field theory and phase transitions. For reviews see [1, 2, 3, 4].

Recently, the Boltzmann weights for the $SO(k)$ spin vertex models were described [5, 6]. This note is a continuation of these works. Specifically, in the case of D_n spin models, only the direct Boltzmann weights were determined. Namely, the $SS \rightarrow SS$ Boltzmann weights, where S denotes the spin representation. When n is odd, there are also the anti-spinor representations, \bar{S} . Our purpose here is to describe also the Boltzmann weights involving the spinor and anti-spinor representations. For calculation reasons we do this, explicitly, only for $n = 3$ and $n = 5$.

Our solution for the Boltzmann weights is trigonometric. Note that rational solutions were known before by a different method [7].

We denote the simple roots of D_n by $\alpha_r = \epsilon_r - \epsilon_{r+1}$, where $r = 1, 2, \dots, n-1$ and $\alpha_n = \epsilon_{n-1} + \epsilon_n$, and ϵ_r are orthogonal unit vectors. The spin representation has the weights $\sum_{i=1}^n p_i \epsilon_i / 2$ where $p_i = \pm 1$ and $\prod_{i=1}^n p_i = \pm 1$ for the spinor representation (S) and the anti-spinor representation (\bar{S}), respectively. We shift these weights by $1/2$ to describe the weights by a vector m , where $m_i = 0$ or 1 . We denote $V = S \oplus \bar{S}$.

A solution for the spin vertex models, which commutes with the co-product in the quantum group $U_{q^2}(D_n)$ was described by Wenzl [8]. It is given by the element C of $\text{End}(V \otimes V)$,

$$C_{m,s}^{b,c} = \sum_{j=1}^n \delta_{m_j, 1-s_j} (-q^2)^{\{m-s\}_j} \delta_{b, \bar{m}_j} \delta_{c, \bar{s}_j}, \quad (1)$$

where

$$\{m\}_j = \sum_{r=1}^j m_r, \quad (2)$$

and \bar{s}_j is equal to s_j except for the coordinate j where it is $1-s_j$. The eigenvalues

of C are

$$\lambda_j = \pm s(n - j), \quad \text{for } j = 0, 1, \dots, n, \quad (3)$$

where

$$s(x) = \frac{q^{2x} - q^{-2x}}{q^2 - q^{-2}}. \quad (4)$$

The solution C maps $S \otimes S$ to $\bar{S} \otimes \bar{S}$. Thus, we square it, where C^2 maps the $S \otimes S$ to itself. We define the projection operators,

$$(P^a)^{b,c}_{m,s} = \prod_{p \neq a} \left[\frac{C^2 - \lambda_p^2 I}{\lambda_a^2 - \lambda_p^2} \right], \quad (5)$$

where I is the identity map and the product is in $\text{End}(V \times V)$.

To make contact with conformal field theory we define

$$q^2 = \exp \left[\frac{\pi i}{r + 2n - 2} \right], \quad (6)$$

where r is the level of the D_n WZW model [3]. The solution to the Yang–Baxter equation for $S \otimes S$ was described in ref. [6]. The Casimir of the representation V_j , which is the anti-symmetric product of j vector representations, is given by

$$C(V_j) = C_j = j(2n - j). \quad (7)$$

We assume that n is odd. We define $h_i = n - 2i$, for $i = 0, 1, \dots, (n - 1)/2$. We define the ‘crossing’ parameters,

$$\zeta_j = (C_{h_{j+1}} - C_{h_j})/2, \quad (8)$$

for $j = 0, 1, \dots, (n - 1)/2$. We define $p(x) = q^x - q^{-x}$. The solution to the Yang

Baxter equation for $S \otimes S \rightarrow S \otimes S$, or $\bar{S} \otimes \bar{S} \rightarrow \bar{S} \otimes \bar{S}$, is then given by [6],

$$R_{m,s}^{a,b}(u) = \sum_{j=0}^{(n-1)/2} f_j(u) (P^{h_j})_{m,s}^{a,b}, \quad (9)$$

where

$$f_a(u) = \frac{\left[\prod_{j=1}^a p(\zeta_{j-1} - u) \right] \left[\prod_{j=a+1}^{(n-1)/2} p(\zeta_{j-1} + u) \right]}{\left[\prod_{j=1}^{(n-1)/2} p(\zeta_{j-1}) \right]}, \quad (10)$$

where $a = 0, 1, \dots, (n-1)/2$.

The solution eq. (9) obeys the Yang–Baxter equation (YBE),

$$\sum_{\alpha,\beta,\gamma} R_{j,k}^{\beta,\alpha}(u) R_{i,\beta}^{l,\gamma}(u+v) R_{\gamma,\alpha}^{m,s}(v) = \sum_{\alpha,\beta,\gamma} R_{i,j}^{\alpha,\beta}(v) R_{\beta,k}^{\gamma,s}(u+v) R_{\alpha,\gamma}^{l,m}(u), \quad (11)$$

provided that i, j, k, l, m, s are all taken from the spinor representation or that they are all taken from the anti–spinor representation.

We wish to describe now the general solution to the YBE, as applicable to the other representations in $\text{End}(V \otimes V)$. First we equate to zero,

$$R_{m,s}^{a,b}(u) = 0, \quad (12)$$

where m, a are in the spinor representation, and s, b are in the anti–spinor representation, or m, a are in the anti–spinor representation and s, b are in the spinor representation.

To get the other cases we use the crossing symmetry,

$$R_{j,i}^{k,l}(u) = R_{\bar{k},\bar{j}}^{l,\bar{i}}(\lambda - u) \left[\frac{r(i)r(l)}{r(j)r(k)} \right], \quad (13)$$

where the crossing parameter is

$$\lambda = C_2/2, \quad (14)$$

and $r(i)$ are the crossing factors, which are as yet unknown. Here \bar{i} is the representation with the weight $-\lambda_i$ where λ_i is the weight of the representation i . We

also use the symmetry relation

$$R_{j,i}^{k,l}(u) = R_{k,l}^{j,i}(u). \quad (15)$$

The crossing symmetry gives us the R matrix $R_{m,s}^{a,b}(u)$ where m, b are in the spinor representation and s, a are in the anti-spinor representation, or vice versa. This completes all the values of $R(u)$ in $\text{End}(V \otimes V)$.

We still need to determine the crossing factors. This we do on a case by case basis. The idea is to substitute substitute some unknown crossing factors, eq. (13) and then to use the YBE with various weights and to solve it for the crossing factors. We also know the relation,

$$r(\bar{i}) = 1/r(i). \quad (16)$$

In the case of D_3 we have four weights in the representation S which are

$$\lambda_1 = \{1, 0, 0\}, \quad \lambda_2 = \{0, 1, 0\}, \quad \lambda_3 = \{0, 0, 1\}, \quad \lambda_4 = \{1, 1, 1\}. \quad (17)$$

The weight of \bar{i} is $\lambda_{\bar{i}} = \{1, 1, 1\} - \lambda_i$. By solving the YBE we get some values for the crossing factors. There are several solutions. One of them is the following,

$$\{r[1], r[2], r[3], r[4]\} = \{-1, q, -q^2, 1/q\}. \quad (18)$$

Substituting these values, we can see that the YBE indeed holds for all the values of the weights in $\text{End}(V \otimes V)$.

Let us turn now to D_5 . We find that the conservation law

$$R_{i,j}^{k,l}(u) = 0 \text{ if } \lambda_i + \lambda_j \neq \lambda_k + \lambda_l, \quad (19)$$

implies that the Boltzmann weights do not depend on the crossing factors, eq. (13), since the only solution for eq. (19) for $S\bar{S} \rightarrow \bar{S}S$ is when i and j are equal

to k and l , or vice versa, and then the crossing factors cancel. Thus, we can directly substitute the YBE and we find that it is, indeed, obeyed for various weights.

For D_n with $n \geq 7$ our limitation of the calculation power, implies that it is not practical to do the computation. So, we have to limit ourself to the explicit examples of D_3 and D_5 .

Acknowledgements: I thank Alexander Zamolodchikov, Alexander Belavin and Paul Wiegmann for helpful discussions.

REFERENCES

1. R.J. Baxter, “Exactly solved models in statistical mechanics”, Dover Publications (1982).
2. M. Wadati, T. Deguchi and Y. Akutsu, Physics Reports 180 (4) (1989) 247.
3. P. Francesco, P. Mathieu and D. Senechal, “Conformal field theory”, Springer–Verlag, Graduate texts in contemporary physics (1997).
4. P. Ginsparg, “Applied conformal field theory”, Les Houches lectures (1988).
5. D. Gepner, Nucl.Phys. B 958 (2020) 115116.
6. V. Belavin, D. Gepner and H. Wenzl, Nucl.Phys. B 959 (2020) 115160.
7. E. Ogievetsky and P. Wiegmann, Phys.Lett. B 168 (1986) 360.
8. H. Wenzl, “Dualities for spin representation”, arXiv: 2005.11299 (2020).