

# BOUNDS ON COMPLEXITY OF MATRIX MULTIPLICATION AWAY FROM CW TENSORS

ROSER HOMS, JOACHIM JELISIEJEW, MATEUSZ MICHAŁEK, AND TIM SEYNNAEVE

ABSTRACT. We present three families of minimal border rank tensors: they come from highest weight vectors, smoothable algebras, or monomial algebras. We analyse them using Strassen’s laser method and obtain an upper bound 2.431 on  $\omega$ . We also explain how in certain monomial cases using the laser method directly is less profitable than first degenerating. Our results form possible paths in the search for valuable tensors for the laser method away from Coppersmith-Winograd tensors.

## 1. INTRODUCTION

Determining the complexity of matrix multiplication is a central problem in computer science. Its algebraic counterpart translates to estimating the rank or border rank of the matrix multiplication tensor  $M_{\langle n,n,n \rangle} \in \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2} \otimes \mathbb{C}^{n^2}$  [5, 10, 11], [14, Chapter 9.3]. The complexity of matrix multiplication is measured by the constant  $\omega$ , defined as the smallest number such that for any  $\epsilon > 0$  the multiplication of  $n \times n$  matrices can be performed in time  $O(n^{\omega+\epsilon})$ . Equivalently,  $\omega$  is the smallest number such that for any  $\epsilon > 0$  the rank (or border rank) of  $M_{\langle n,n,n \rangle}$  is  $O(n^{\omega+\epsilon})$ .

The best known upper bounds on  $\omega$  are all obtained using the so-called *laser method*, which is based on the work of Strassen [19]. The idea behind the laser method is to, instead of studying the matrix multiplication tensor directly, consider a different tensor which can be proven to have low border rank, and at the same time is “close” to being a matrix multiplication tensor. Strassen obtained a bound  $\omega < 2.48$  using the laser method. Shortly thereafter, Coppersmith and Winograd introduced a new tensor, and applied (in a highly nontrivial way) the laser method to it to obtain  $\omega < 2.3755$  [9]. Since then, the improvements on the bound of  $\omega$  were made by Stothers, Williams, and Le Gall [18, 20, 13], arriving at the current state of the art  $\omega < 2.373$ . These improvements were all obtained by applying the laser method to the Coppersmith-Winograd tensor and its powers. However, several recent results [1, 2, 8] proved the existence of barriers: as a very particular case, using the Coppersmith-Winograd tensors alone in the laser method, one cannot obtain an upper bound for  $\omega$  close to 2.

Below we present other tensors appearing naturally and apply the laser method to find the bound on  $\omega$  they give. We explore two approaches of constructing such tensors. One approach is based on the highest weight vectors in  $S^3(\mathfrak{sl}_n)$ . As we show, a few of them are the Coppersmith-Winograd tensors! We use the other highest weight vectors to prove bounds on  $\omega$ . For one of those tensors we obtain a bound  $\omega < 2.45$ , which is not as good as the Coppersmith-Winograd bound, but better than Strassen’s bound.

Another approach builds upon the works [12] and [3]. Under certain genericity assumptions, a tensor is of minimal border rank if and only if it is the multiplication tensor of a smoothable finite-dimensional algebra. The Coppersmith-Winograd tensor arises in this way: it is the multiplication tensor of an algebra with Hilbert function  $(1, n, 1)$ , and it follows from a result of Cartwright et al. [6] that such an algebra is always smoothable. We give an

---

2020 *Mathematics Subject Classification.* primary: 15A69, 14Q20.

JJ was partially supported by Polish National Science Center, project 2017/26/D/ST1/00755 and by the START fellowship of the Foundation for Polish Science.

example of an algebra with Hilbert function  $(1, n, 2)$  (hence smoothable by [6]) whose multiplication tensor is suitable for the laser method. The obtained bound  $\omega < 2.431$  is better than the bound above, but still not as good as the bound obtained from the Coppersmith-Winograd tensor. We hope that our constructions will be inspiring for other mathematicians to look for “valuable” tensors different from the Coppersmith-Winograd tensor.

Finally, we look at a very simple algebra  $\mathbb{C}[x]/(x^2)$ . The associated tensor may be regarded as a very degenerate CW tensor (in  $\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$ ). By taking the third Kronecker power of that tensor, corresponding to the third tensor product of algebras, we obtain a new tensor  $T$ , for which we apply the laser method obtaining  $\omega < 2.56$ . Here however our main observation is that  $T$  degenerates to the CW tensor, which is used to obtain some of the world record bounds on  $\omega$ . This means that directly applying the laser method to  $T$  is very far from optimal. In particular, it suggests that a new method, suited to analyze tensors like  $T$ , is needed – and if found will provide new bounds on  $\omega$ .

## 2. PRELIMINARIES

We start by presenting preliminaries about tensors and fast matrix multiplication. We work in the space  $\mathcal{V} = U \otimes V \otimes W$ , where  $U, V, W$  are finite-dimensional vector spaces over  $\mathbb{C}$ . We will assume familiarity with the notions of rank and border rank of tensors (denoted by  $\text{rk}(T)$  and  $\underline{\text{rk}}(T)$ , respectively) and (border) Waring rank of symmetric tensors, see [10] for an in-depth treatment.

Let  $G := GL(U) \times GL(V) \times GL(W)$ . There is a natural action of  $G$  on  $\mathcal{V}$ , which extends to an action of the algebra  $A := \text{End}(U) \times \text{End}(V) \times \text{End}(W) \supseteq G$ .

**Definition 2.1.** Let  $T, T' \in \mathcal{V}$ .

- We say  $T'$  is a *restriction* of  $T$ , denoted  $T' \leq T$ , if  $T' \in A \cdot T$ .
- We say  $T'$  is a *degeneration* of  $T$ , denoted  $T' \leq T$  if  $T' \in \overline{G \cdot T}$ .

**Definition 2.2.** The *Kronecker product*  $T \boxtimes T'$  of two tensors  $T \in U \otimes V \otimes W$  and  $T' \in U' \otimes V' \otimes W'$  is by definition their tensor product  $T \otimes T'$ , viewed as a 3-way tensor in  $(U \otimes U') \otimes (V \otimes V') \otimes (W \otimes W')$ . The bracketing is important: the rank one tensors in this space are all tensors of the form  $u'' \otimes v'' \otimes w''$ , with  $u'' \in U \otimes U'$ ,  $v'' \in V \otimes V'$ ,  $w'' \in W \otimes W'$ .

A tensor  $T \in \mathbb{C}^a \otimes \mathbb{C}^b \otimes \mathbb{C}^c$  may be identified with a linear map  $\mathbb{C}^a \rightarrow \mathbb{C}^b \otimes \mathbb{C}^c$ . The image of that map, which may be represented as a linear space of  $b \times c$  matrices, determines  $T$  up to isomorphism. Hence, we will often represent  $T$  as a space  $L_T$  of matrices, as in (3.1).  $T$  is called *concise* if the corresponding linear maps  $\mathbb{C}^a \rightarrow \mathbb{C}^b \otimes \mathbb{C}^c$ ,  $\mathbb{C}^b \rightarrow \mathbb{C}^a \otimes \mathbb{C}^c$ ,  $\mathbb{C}^c \rightarrow \mathbb{C}^a \otimes \mathbb{C}^b$  are injective. Every tensor can be made concise by replacing  $\mathbb{C}^a, \mathbb{C}^b, \mathbb{C}^c$  with suitable subspaces. The border rank of a concise tensor is at least  $\max\{a, b, c\}$ ; in case of equality we say that  $T$  has *minimal border rank*.

We will denote the  $a \times b \times c$  *matrix multiplication tensor* by  $M_{(a,b,c)} \in \mathbb{C}^{ab} \otimes \mathbb{C}^{bc} \otimes \mathbb{C}^{ca}$ , see [10, Section 2.5.2] and [14, Chapter 9.3] for details. To prove bounds on  $\omega$ , instead of analyzing  $M_{(a,b,c)}$  directly, we will consider different tensors that are both of low (typically minimal) border rank, and degenerate to a direct sum of many large matrix multiplication tensors. This second requirement is made quantitative using the notion of degeneracy value.

**Definition 2.3** (See [11, Definition 3.4.7.1]). Let  $T \in U \otimes V \otimes W$  be a tensor. For  $N \in \mathbb{N}$ , we define

$$V_{\omega, N}(T) = \sup \left\{ \sum_{i=1}^q (a_i b_i c_i)^{\omega/3} \mid T^{\boxtimes N} \supseteq \bigoplus_{i=1}^q M_{(a_i, b_i, c_i)} \right\},$$

where we take the supremum over all possible ways of degenerating  $T^{\boxtimes N}$  to a direct sum of matrix multiplication tensors. The *degeneracy value* (or simply *value*)  $V_{\omega}(T)$  of  $T$  is defined as the supremum  $\sup_N V_{\omega, N}(T)^{\frac{1}{N}}$ .

*Remark 2.4.* Here are some easy but important properties of the degeneracy value. The first two are trivial, for the third one, see [11, Section 3.4.7].

- If  $T \succeq T'$ , then  $V_\omega(T) \geq V_\omega(T')$ .
- Supermultiplicativity:  $V_\omega(T \boxtimes T') \geq V_\omega(T)V_\omega(T')$ .
- Superadditivity:  $V_\omega(T \oplus T') \geq V_\omega(T) + V_\omega(T')$ .

The following theorem is a restatement of Schönhage's asymptotic sum inequality ([15], see also [5, (15.11)]).

**Theorem 2.5.** *For any tensor  $T$ , we have  $V_\omega(T) \leq \underline{\text{rk}}(T)$ .*

Therefore, any given tensor  $T$  with an upper bound on its border rank and a lower bound on its value yields a bound on  $\omega$ , as illustrated in Example 2.11. In the rest of this section we briefly recall Strassen's *laser method*, which can be used to estimate the value of a tensor if the tensor can be decomposed into smaller tensors for which the value is known. The version we present here was first introduced in [17].

**Definition 2.6.** Let  $T \in U \otimes V \otimes W$  be a tensor.

- A *blocking*  $D$  of  $T$  is given by decompositions  $U = \bigoplus_{i \in I} U_i$ ,  $V = \bigoplus_{j \in J} V_j$ ,  $W = \bigoplus_{k \in K} W_k$ . These induce a decomposition

$$T = \sum_{(i,j,k) \in I \times J \times K} T_{(i,j,k)}.$$

- The *support* of a blocking  $D$ , denoted  $\text{supp}_D T$ , consists of all triples  $(i, j, k) \in I \times J \times K$  for which  $T_{(i,j,k)} \neq 0$ .
- We say  $\text{supp}_D T$  is *tight*, if there are injections  $\alpha : I \rightarrow \mathbb{Z}^r$ ,  $\beta : J \rightarrow \mathbb{Z}^r$ ,  $\gamma : K \rightarrow \mathbb{Z}^r$  s.t.  $\alpha(i) + \beta(j) + \gamma(k) = 0$  for all  $(i, j, k) \in \text{supp}_D T$ .
- The *symmetrization* of  $T$ , denoted by  $\tilde{T}$ , is the Kronecker product  $T \boxtimes T' \boxtimes T'' \in (U \otimes V \otimes W)^{\otimes 3}$ , where  $T' \in V \otimes W \otimes U$  and  $T'' \in W \otimes U \otimes V$  are obtained from  $T$  by cyclically permuting the indices.

It is not hard to see that  $\underline{\text{rk}}(\tilde{T}) \leq \underline{\text{rk}}(T)^3$  and  $V_\omega(\tilde{T}) \geq V_\omega(T)^3$ . As opposed to [13, Theorem 4.1], instead of assuming lower bounds on the values of the blocks, we will only assume known bounds on the values of their symmetrizations.

*Remark 2.7.* We will apply the laser method to tensors  $T \in \mathbb{C}^m \otimes \mathbb{C}^m \otimes \mathbb{C}^m$  of minimal border rank, where the blocking  $D$  is given by partitioning the basis elements of each  $\mathbb{C}^m$  into three groups labeled by  $\{0, 1, 2\}$ . We assume that the tensor has nonzero entries only if the labels belong to the groups with indices summing up to 2, i.e. if  $(i, j, k) \in \text{supp}_D T$  then  $i + j + k = 2$ . For instance, under our assumptions  $T_{(1,1,0)}$  is typically nonzero, while  $T_{(1,0,0)} = 0$ . Such a blocking is automatically tight: we can take  $r = 1$  and define  $\alpha(i) = i$ ,  $\beta(j) = j$ ,  $\gamma(k) = 2 - k$ .

Let  $D$  be a blocking of a tensor  $T \in U \otimes V \otimes W$ , with indexing sets  $I, J, K$ . We will consider probability distributions  $P : \text{supp}_D T \rightarrow [0, 1]$  on  $\text{supp}_D T$  (this gives each block a "weight"). We will write  $P_I : I \rightarrow [0, 1]$  for the marginal distribution on  $I$ , and similar for  $J, K$ . The entropy  $H(P_I) := -\sum_{i \in I} P(i) \log P(i)$  (and similar for  $J, K$ ) will play an important role in the laser method. We will make a technical (but easy to verify) assumption on our blocking which removes the need for the extra term  $-\Gamma_S(P)$  appearing in [13, Theorem 4.1]:

**Definition 2.8.** We say that a subset  $\Phi \subseteq I \times J \times K$  is *reconstructible*, if every probability distribution on  $\Phi$  is uniquely determined by its 3 marginal distributions.

*Example 2.9.* Let  $I = J = K = \{0, 1, 2\}$ , and take

$$\Phi = \{(2, 0, 0), (0, 2, 0), (0, 0, 2), (1, 1, 0), (1, 0, 1), (0, 1, 1)\}.$$

Then  $\Phi$  is tight and reconstructible: if  $P$  is a probability distribution on  $\Phi$ , then the equalities  $P(2, 0, 0) = P_I(2)$ ,  $P(0, 1, 1) = P_I(0) - P_J(2) - P_K(2), \dots$  allow us to reconstruct  $P$  from the marginal distributions  $P_I, P_J, P_K$ . However, if we consider

$$\Phi' = \{(2, 1, 0), (1, 2, 0), (2, 0, 1), (1, 0, 2), (0, 2, 1), (0, 1, 2)\},$$

then  $\Phi'$  is again tight, but not reconstructible. For instance, the uniform distribution  $P(x, y, z) = \frac{1}{6}$  and the distribution  $P(0, 1, 2) = P(2, 0, 1) = P(1, 2, 0) = \frac{1}{3}$  have the same marginals.

**Theorem 2.10** (Laser method). *Let  $T \in U \otimes V \otimes W$ , and let  $D$  be a blocking of  $T$ , indexed by  $I \times J \times K$ . Assume that  $\text{supp}_D T$  is tight and reconstructible. Let  $P$  be any probability distribution on  $\text{supp}_D T$ . Then*

$$(2.1) \quad \log V_\omega(\tilde{T}) \geq H(P_I) + H(P_J) + H(P_K) + \sum_{\text{supp}_D T} P(i, j, k) \log V_\omega(\widetilde{T_{(i,j,k)}}).$$

*Proof.* See Appendix A. ■

*Example 2.11* (Strassen's tensor). Consider the following tensor:

$$T = T_{STR,n} := \sum_{i=1}^n u_0 \otimes v_i \otimes w_i + \sum_{i=1}^n u_i \otimes v_0 \otimes w_i \in U \otimes V \otimes W,$$

where  $U, V$  and  $W$  have respective bases  $\{u_0, u_1, \dots, u_n\}$ ,  $\{v_0, v_1, \dots, v_n\}$ , and  $\{w_1, \dots, w_n\}$ . We consider the block decomposition  $D$  with  $I = J = \{0, 1\}$ ,  $K = \{1\}$ ,  $U_0 = \langle u_0 \rangle$ ,  $U_1 = \langle u_1, \dots, u_n \rangle$ ,  $V_0 = \langle v_0 \rangle$ ,  $V_1 = \langle v_1, \dots, v_n \rangle$ ,  $W_1 = \langle w_1, \dots, w_n \rangle$ . Then  $\text{supp}_D T = \{(0, 1, 1), (1, 0, 1)\}$  which is clearly tight and reconstructible.

Since  $T_{(0,1,1)} = \sum_i u_0 \otimes v_i \otimes w_i = M_{(1,1,n)}$ , we find that  $\widetilde{T_{(0,1,1)}} = M_{(n,n,n)}$ . Similarly  $\widetilde{T_{(1,0,1)}} = M_{(n,n,n)}$ . Let  $P$  be the uniform distribution on  $\{(0, 1, 1), (1, 0, 1)\}$ , then  $H(P_1) = H(P_2) = \log(2)$  and  $H(P_3) = 0$ . Theorem 2.10 now gives

$$\log V_\omega(\tilde{T}) \geq 2 \log(2) + \log(n^\omega),$$

so that  $V_\omega(\tilde{T}) \geq 4n^\omega$ . The border rank of  $T$  is equal\* to  $n + 1$ , as can be seen from writing it as a limit

$$\lim_{t \rightarrow 0} \frac{1}{t} \left( \sum_{j=1}^n (u_0 + tu_j) \otimes (v_0 + tv_j) \otimes w_j - u_0 \otimes v_0 \otimes (w_1 + \dots + w_n) \right).$$

Hence the border rank of  $\tilde{T}$  is at most  $(n + 1)^3$ , so Theorem 2.5 yields

$$4n^\omega \leq (n + 1)^3.$$

For  $n = 5$ , this gives Strassen's bound  $\omega < 2.48$  [19].

### 3. TENSORS FROM HIGHEST WEIGHT VECTORS

Given elements  $x, y, z$  in a vector space  $V$ ,  $xyz := \frac{1}{6}(x \otimes y \otimes z + x \otimes z \otimes y + y \otimes x \otimes z + y \otimes z \otimes x + z \otimes x \otimes y + z \otimes y \otimes x)$  is a 3-way tensor in  $V \otimes V \otimes V$ . Similarly,  $xy := \frac{1}{2}(x \otimes y + y \otimes x)$  is a matrix in  $V \otimes V$ .

We recall the small and big Coppersmith-Winograd tensors  $T_{cw,m}$  and  $T_{CW,m}$  from [9]. They are symmetric tensors, precisely elements of  $\mathbb{C}[x_0, \dots, x_{m+1}]_3$ , given by

$$T_{cw,m} = \sum_{i=1}^m 3x_0 x_i^2 \quad \text{and} \quad T_{CW,m} = 3x_0^2 x_{m+1} + 3 \sum_{i=1}^m x_0 x_i^2$$

with border Waring rank equal to  $m + 2$ .

In [16], the fourth author studied the highest weight vectors of the  $\mathfrak{sl}_n$ -representation  $S^3(\mathfrak{gl}_n)$ . A basis of  $\mathfrak{gl}_n$  is given by  $\{E_{i,j}\}_{1 \leq i, j \leq n}$ , where  $E_{i,j}$  is the matrix with a 1 at position  $(i, j)$  and 0 elsewhere. One of the highest weight vectors in  $S^3(\mathfrak{gl}_n)$  is the symmetrized matrix multiplication tensor  $\sum_{i,j,k} E_{i,j} E_{j,k} E_{k,i}$  [7]. The other highest weight vectors are listed in [16, Table 2]. An interesting observation is that many of these highest weight vectors are, up to a change of variables, equal to  $T_{cw,m}$  or  $T_{CW,m}$  for some value of  $m$ .

---

\*The border rank cannot be lower than  $n + 1$ , as  $T$  is a concise tensor.

**Proposition 3.1.** *The following equalities hold, up to a change of variables:*

$$\begin{aligned}
 IE_{1,n}E_{2,n-1} - IE_{1,n-1}E_{2,n} &= T_{cw,4} \\
 E_{1,n}E_{1,n-1}E_{2,n} - E_{1,n}E_{1,n}E_{2,n-1} &= T_{CW,2} \\
 \sum_i IE_{1,i}E_{i,n} &= T_{cw,2n-2} \\
 \sum_i E_{1,n}E_{1,i}E_{i,n} &= T_{CW,2n-4} \\
 \sum_{i,j} E_{1,n}E_{i,j}E_{j,i} &= T_{CW,n^2-2},
 \end{aligned}$$

*i.e. five of the highest weight vectors are Coppersmith-Winograd tensors.*

*Proof.* We will only prove the fourth equality; the other ones are similar and left to the reader. Note that

$$\sum_{j=1}^n E_{1,n}E_{1,j}E_{j,n} = E_{1,n}^2(E_{1,1} + E_{n,n}) + \sum_{j=2}^{n-1} E_{1,n}E_{1,j}E_{j,n}.$$

If we substitute  $E_{1,n} = 3x_0$ ,  $E_{1,1} + E_{n,n} = \frac{1}{3}x_{2n-3}$ ,  $E_{1,j} = x_{2j-3} + ix_{2j-2}$  and  $E_{j,n} = x_{2j-3} - ix_{2j-2}$  for  $2 \leq j \leq n-1$  (where  $i$  is the imaginary unit, so that  $E_{1,j}E_{j,n} = x_{2j-3}^2 + x_{2j-2}^2$ ), we obtain the Coppersmith-Winograd tensor  $T_{CW,2n-4}$ .  $\blacksquare$

We will now focus on one of the other highest weight vectors from [16, Table 2], that is not a Coppersmith-Winograd tensor. Let  $n \geq 3$  and write:

$$T_{HW,n} := \sum_{i=1}^n (E_{1,n}E_{2,i}E_{i,n} - E_{2,n}E_{1,i}E_{i,n}) \in S^3(\mathbb{C}^{n^2}).$$

This is not a concise tensor, but we can make it concise by changing the ambient space: first, rewrite  $T_{HW,n}$  as

$$E_{1,n}^2E_{2,1} - E_{2,n}^2E_{1,2} + E_{1,n}E_{2,n}(E_{2,2} - E_{1,1}) + \sum_{i=3}^{n-1} (E_{1,n}E_{2,i}E_{i,n} - E_{2,n}E_{1,i}E_{i,n}).$$

For every  $i \in \{3 \dots n-1\}$ , we put  $E_{i,n} = x_{i-3}$ ,  $E_{2,i} = y_{i-3}$ , and  $E_{1,i} = z_{i-3}$ . Moreover, we put  $E_{1,1} - E_{2,2} = b_0$ ,  $E_{1,n} = a_1$ ,  $E_{2,n} = -a_2$ ,  $E_{2,1} = b_1$ ,  $E_{1,2} = -b_2$ . Then, after suitably rescaling, our tensor becomes

$$T_{HW,m} = 6a_1a_2b_0 + 3a_1^2b_1 + 3a_2^2b_2 + 6 \sum_{j=1}^m (a_1x_jy_j + a_2x_jz_j) \in S^3(V) \subseteq V \otimes V \otimes V,$$

where  $m = n - 3$  and  $V$  is the  $(3m + 5)$ -dimensional vector space with basis

$$\{a_1, a_2, x_1, \dots, x_m, y_1, \dots, y_m, z_1, \dots, z_m, b_0, b_1, b_2\}.$$

The space  $L_{T_{HW,m}}$ , made of all possible contractions of the tensor  $T_{HW,m}$ , consists of all matrices of the form

$$(3.1) \quad \left( \begin{array}{cc|ccc|ccc|ccc|ccc} b_1 & b_0 & y_1 & \dots & y_m & x_1 & \dots & x_m & 0 & \dots & 0 & a_2 & a_1 & 0 \\ b_0 & b_2 & z_1 & \dots & z_m & 0 & \dots & 0 & x_1 & \dots & x_m & a_1 & 0 & a_2 \\ \hline y_1 & z_1 & & & & a_1 & & & a_2 & & & & & \\ \vdots & \vdots & & & & & \ddots & & & & & & & \\ y_m & z_m & & & & & & a_1 & & & a_2 & & & \\ \hline x_1 & 0 & a_1 & & & & & & & & & & & \\ \vdots & \vdots & & \ddots & & & & & & & & & & \\ x_m & 0 & & & a_1 & & & & & & & & & \\ \hline 0 & x_1 & a_2 & & & & & & & & & & & \\ \vdots & \vdots & & \ddots & & & & & & & & & & \\ 0 & x_m & & & a_2 & & & & & & & & & \\ \hline a_2 & a_1 & & & & & & & & & & & & \\ a_1 & 0 & & & & & & & & & & & & \\ 0 & a_2 & & & & & & & & & & & & \end{array} \right).$$

It is easy to see that  $T_{HW,m}$  is concise, i.e. the induced map  $V^* \rightarrow V \otimes V$  is injective. From this, it follows that  $T_{HW,m}$  has border rank at least  $\dim V = 3m + 5$ . The reason why this highest weight vector in particular is of interest, is that it has minimal border rank.

**Theorem 3.2.** *The border rank of  $T_{HW,m}$  is equal to  $3m + 5$ .*

*Proof.* The case  $m = 0$  follows from Proposition 3.3. For  $m \geq 1$ , consider the following collection of rank 1 symmetric matrices in  $V \otimes V$ :

- $A_{i,t} := (t^{-1}a_1 + tx_i + t^2y_i)^2 = t^{-2}a_1^2 + 2a_1x_i + 2ta_1y_i + t^2x_i^2 + 2t^3x_iy_i + t^4y_i^2$ ,
- $B_{i,t} := (t^{-1}a_2 + tx_i + t^2z_i)^2 = t^{-2}a_2^2 + 2a_2x_i + 2ta_2z_i + t^2x_i^2 + 2t^3x_iz_i + t^4z_i^2$ ,
- $C_{i,t} := (\sqrt{2}t^{-1}a_1/2 + \sqrt{2}t^{-1}a_2/2 + \sqrt{2}tx_i)^2$ ,
- $D := (a_1 - a_2)^2 = a_1^2 + a_2^2 - 2a_1a_2$ ,
- $E_{1,t} := (t^{-2}a_1 + t^2 \sum_{j=1}^m x_j + t^3 \sum_{j=1}^m y_j + t^5b_0 - t^5b_1)^2$ ,
- $E_{2,t} := (t^{-2}a_2 + t^2 \sum_{j=1}^m x_j + t^3 \sum_{j=1}^m z_j + t^5b_0 - t^5b_2)^2$ ,
- $F_t := (\sqrt{2}t^{-2}a_1/2 + \sqrt{2}t^{-2}a_2/2 + \sqrt{2}t^2 \sum_{j=1}^m x_j + 2\sqrt{2}t^5b_0)^2$ ,
- $G_t := (t^{-4} - mt^{-2} - 1/2 - l)a_1^2 + (mt^{-2} - t^{-4} - 1/2 - l)a_2^2 + (2l - 1)a_1a_2$ ,

with  $l := (mt^2 - 1)^2/(2t^8)$ .

We want to prove that  $\lim_{t \rightarrow 0} \langle A_{i,t}, B_{i,t}, C_{i,t}, D, E_{1,t}, E_{2,t}, F_t, G_t \rangle = L_{T_{HW,m}}$ .

$L_{T_{HW,m}}|_{x_i=1}$  denotes the specialization of a matrix in  $L_{T_{HW,m}}$  to  $x_i = 1$  and the rest of variables set to 0. We use similar notation for other variables. We can build  $3m + 3$  generators of the  $(3m + 5)$ -dimensional space of matrices  $L_{T_{HW,m}}$  as follows:

- $\lim_{t \rightarrow 0} A_{i,t} - t^2 E_{1,t} = 2a_1x_i = L_T|_{y_i=1}$ ,
- $\lim_{t \rightarrow 0} B_{i,t} - t^2 E_{2,t} = 2a_2x_i = L_T|_{z_i=1}$ ,
- $\lim_{t \rightarrow 0} t^{-1} (A_{i,t} + B_{i,t} - C_{i,t} - (1/2t^2)D) = 2a_1y_i + 2a_2z_i = L_T|_{x_i=1}$ .
- $\lim_{t \rightarrow 0} t^2 A_{i,t} = a_1^2 = L_T|_{b_1=1}$ ,
- $\lim_{t \rightarrow 0} t^2 B_{i,t} = a_2^2 = L_T|_{b_2=1}$ ,
- $\lim_{t \rightarrow 0} t^2 A_{i,t} + t^2 B_{i,t} - D = 2a_1a_2 = L_T|_{b_0=1}$ .

The remaining two can be obtained as follows:

- $\lim_{t \rightarrow 0} \frac{\sum_{j=1}^m A_{j,t} + \sum_{j=1}^m B_{j,t} - \sum_{j=1}^m C_{j,t} - E_{1,t} - E_{2,t} + (\frac{1}{2t^4} - \frac{m}{2t^2})D + F_t}{t^3} = 2a_1b_0 + 2a_1b_1 + 2a_2b_0 + 2a_2b_2 + 2 \sum_{j=1}^m x_j y_j + 2 \sum_{j=1}^m x_j z_j = L_T|_{a_1=a_2=1}$ ,

$$\bullet \lim_{t \rightarrow 0} \frac{\sum_{j=1}^m A_{j,t} - \sum_{j=1}^m B_{j,t} - E_{1,t} + E_{2,t} + t^4 F_t + lD + G_t}{t^3} = -2a_1 b_0 + 2a_1 b_1 + 2a_2 b_0 - 2a_2 b_2 + 2 \sum_{j=1}^m x_j y_j - 2 \sum_{j=1}^m x_j z_j = LT|_{a_1=-a_2=1}.$$

Ever stronger, we conjecture that the border Waring rank of  $T_{HW,m}$  is equal to  $3m + 5$ . For  $m \leq 1$  we have the following symmetric border decompositions.

**Proposition 3.3.** *For  $m = 0$  and  $m = 1$ , the tensor  $T_{HW,m}$  has border Waring rank  $3m + 5$ .*

*Proof.* We provide explicit Waring rank approximations in both cases. For the case  $m = 0$ , let

$$T_{0,t} = 3(a_1 + tb_1)^3 + 6(a_2 + tb_2)^3 + (a_1 - 2a_2)^3 - 3(a_1 - a_2 + tb_0)^3 - (a_1 + a_2 - 3tb_0)^3.$$

Then

$$\lim_{t \rightarrow 0} \frac{T_{0,t}}{t} = 36a_1 a_2 b_0 + 9a_1^2 b_1 + 18a_2^2 b_2,$$

which is equal to our tensor  $T_{HW,1}$  up to rescaling the variables. For the case  $m = 1$ , let

$$\begin{aligned} T_{1,t} = & (-a_1 - a_2 + t^3 b_0)^3 + \frac{1}{3}(-a_1 + a_2)^3 + (a_1 - t^2 y_1 + t^3 b_1)^3 + \\ & \frac{1}{3}(a_1 - a_2 + 3tx_1 - 3t^3 b_0)^3 + \frac{1}{4}(2a_2 - 2tx_1 - t^2 z_1 + t^3 b_2)^3 + \\ & (-a_1 - 2tx_1 + t^2 y_1)^3 + \frac{1}{4}(-2a_2 + t^2 z_1)^3 + (a_1 + a_2 + tx_1)^3. \end{aligned}$$

Then

$$\lim_{t \rightarrow 0} \frac{T_{1,t}}{t^3} = 12a_1 a_2 b_0 + 3a_1^2 b_1 + 3a_2^2 b_2 + 12a_1 x_1 y_1 + 6a_2 x_1 z_1,$$

which is equal to  $T_{HW,1}$  up to rescaling the variables.  $\blacksquare$

*Remark 3.4.* One approach to generalizing Proposition 3.3 to all  $m$  would be using border apolarity [4]. It would be enough to find a homogeneous ideal

$$I \subset \mathbb{C}[a_1, a_2, x_1, \dots, x_m, y_1, \dots, y_m, z_1, \dots, z_m, b_0, b_1, b_2]$$

such that  $I \subset T_{HW,m}^\perp$  and prove that  $I$  is a limit of saturated ideals of points. However, the ideal  $I$  is not uniquely determined by  $T_{HW,m}$ . One possible choice of  $I$  would be  $I_0$  given by all quadrics annihilating  $T_{HW,m}$  and additionally by the element  $a_1^4 a_2 - a_1 a_2^4$  and by

$$(a_1^3 - a_2^3)x_i \text{ for } i = 1, \dots, m.$$

This ideal has the correct Hilbert function. While the ideal  $I_0$  is indeed a limit of saturated ideals for small values of  $m$ , we do not know whether it is so for all  $m$ .

We now apply the laser method to the tensor  $T_m = T_{HW,m}$ . The blocking  $D$  is given by  $V = V_0 \oplus V_1 \oplus V_2$ , with

$$V_0 = \langle a_1, a_2 \rangle, V_1 = \langle x_1, \dots, x_m, y_1, \dots, y_m, z_1, \dots, z_m \rangle, V_2 = \langle b_0, b_1, b_2 \rangle$$

The support of  $T_m$  is tight:  $\{(0, 0, 2), (0, 2, 0), (2, 0, 0), (1, 1, 0), (1, 0, 1), (0, 1, 1)\}$ . The block

$$(3.2) \quad T_{m,(1,1,0)} = \sum_{j=1}^m (x_j \otimes y_j \otimes a_1 + x_j \otimes z_j \otimes a_2 + y_j \otimes x_j \otimes a_1 + z_j \otimes x_j \otimes a_2),$$

can be identified with the Kronecker product

$$\left( \sum_{j=0}^{m-1} e_j \otimes e_j \otimes 1 \right) \boxtimes (e_0 \otimes e_1 \otimes e_1 + e_0 \otimes e_2 \otimes e_2 + e_1 \otimes e_0 \otimes e_1 + e_2 \otimes e_0 \otimes e_2).$$

The first factor is the matrix multiplication tensor  $M_{(1,m,1)}$ , whose symmetrization (with respect to  $\mathbb{Z}/3\mathbb{Z} \subset S_3$ ) has value  $V_\omega(M_{(m,m,m)}) = m^\omega$ . The second factor is Strassen's tensor  $T_{STR,2}$  from Example 2.11. We have  $V_\omega(\widetilde{T_{STR,2}}) \geq 4 \cdot 2^\omega$ . Hence

$$V_\omega(\widetilde{T_{m,(1,1,0)}}) \geq 4(2m)^\omega.$$

The same holds for  $T_{m,(1,0,1)}$  and  $T_{m,(0,1,1)}$ . To compute the value of the remaining blocks, i.e.  $T_{m,(0,0,2)}$ ,  $T_{m,(2,0,0)}$  and  $T_{m,(0,2,0)}$ , we apply the laser method to the symmetrization of the tensor

$$T_{m,(0,0,2)} = \left( \begin{array}{c|c} b_1 & b_0 \\ \hline b_0 & b_2 \end{array} \right),$$

and the blocking as above, where additionally the letters are in three separate groups. Equivalently  $T_{m,(0,0,2)} = a_1 \otimes a_1 \otimes b_1 + a_1 \otimes a_2 \otimes b_0 + a_2 \otimes a_1 \otimes b_0 + a_2 \otimes a_2 \otimes b_2$ , and the blocks are simply the summands in this expression. The value of (the symmetrization of) each block is equal to 1. Hence by assigning the blocks  $a_1 \otimes a_1 \otimes b_1$  and  $a_2 \otimes a_2 \otimes b_2$  a probability  $\frac{1}{3}$ , and the remaining two blocks a probability  $\frac{1}{6}$ , the inequality (2.1) becomes

$$\log V_\omega(\widetilde{T_{m,(0,0,2)}}) \geq \log(2) + \log(2) + \log(3) + 0,$$

so we obtain that this symmetrization  $\widetilde{T_{m,(0,0,2)}}$  has value at least 12.

With the above value estimates, Theorem 2.10 yields

$$\begin{aligned} 3 \log(3m+5) &= 3 \log \underline{\text{rk}}(T_m) \geq \log V_\omega(\widetilde{T_m}) \geq \\ &H(P_1) + H(P_2) + H(P_3) + (P(2,0,0) + P(0,2,0) + P(0,0,2)) \log(12) \\ &\quad + (P(1,1,0) + P(1,0,1) + P(0,1,1)) \log(4(2m)^\omega). \end{aligned}$$

We obtain for every  $m$  a bound on  $\omega$ , by optimizing over all probability distributions on  $\text{supp}_D T_m$ . For  $m = 7$  we obtain  $\omega < 2.45$ . We point out that our analysis may be improved.

#### 4. TENSORS FROM SMOOTHABLE ALGEBRAS

An algebra  $A$  is called *smoothable* if  $\text{Spec}(A)$  is a smoothable scheme. Its multiplication map  $V \otimes V \rightarrow V$  (where  $V$  is the underlying vector space of  $A$ ) can be seen as a tensor  $T_A \in V^* \otimes V^* \otimes V$ .

**Proposition 4.1.** [3, Corollary 3.6] *If  $A$  is a smoothable algebra, then the multiplication map of  $A$  is a tensor of minimal border rank.*

**Theorem 4.2** ([6, Propositions 4.12 and 4.13]). *If  $A$  is a local algebra with Hilbert function  $(1, n, 1)$  or  $(1, n, 2)$ , then  $A$  is smoothable.*

*Example 4.3.* The Coppersmith-Winograd tensor arises as the multiplication tensor of a smoothable algebra. Let  $V$  be an  $(n+2)$ -dimensional vector space with basis  $\{a_0, \dots, a_{n+1}\}$ , and let  $A = (V, *)$  be the algebra with unit  $a_0$  defined by  $a_i * a_i = a_{n+1}$  for  $i = 1, \dots, n$ ,  $a_i * a_j = 0$  for  $1 \leq i < j \leq n+1$ , and  $a_{n+1} * a_{n+1} = 0$ .

*Remark 4.4.* We note that we would get an isomorphic algebra if we defined  $a_0$  to be the unit,  $a_i * a_j = a_{n+1}$  for any  $i + j = n + 1$  and 0 otherwise. In particular, we would get isomorphic tensors.

Equivalently  $A = \mathbb{C}[a_1, \dots, a_{n+1}]/I$ , where  $I$  is the ideal generated by the relations above. The multiplication tensor of  $A$  is equal to

$$\alpha_0 \otimes \alpha_0 \otimes a_0 + \sum_{i=1}^{n+1} (\alpha_0 \otimes \alpha_i \otimes a_i + \alpha_i \otimes \alpha_0 \otimes a_i) + \sum_{i=1}^n (\alpha_i \otimes \alpha_i \otimes a_{n+1}),$$

where  $\{\alpha_0, \dots, \alpha_{n+1}\}$  is the dual basis of  $\{a_0, \dots, a_{n+1}\}$ . This is the Coppersmith-Winograd tensor  $T_{CW,n}$ . Now,  $A$  is a local graded algebra, with  $\mathfrak{m} = (x_1, \dots, x_{n+1})$  and grading given by  $\deg(x_i) = 1$  for  $i = 1, \dots, n$  and  $\deg(x_{n+1}) = 2$ . So the Hilbert function of  $A$  is equal



to  $(1, n, 1)$ . Hence by Theorem 4.2  $\text{Spec}(A)$  is a smoothable scheme, and by Proposition 4.1  $T_{CW,m}$  is of minimal border rank.

*Example 4.5.* Consider the  $3m + 3$ -dimensional algebra  $A = \mathbb{C}[a_1, \dots, a_{3m+2}]/I$ , where the ideal  $I$  is generated by

- $a_{3m+1} - a_i a_{m+i}$  for  $1 \leq i \leq m$ ,
- $a_{3m+2} - a_i a_{2m+i}$  for  $1 \leq i \leq m$ ,
- and all products  $a_i a_j$ ,  $1 \leq i \leq j \leq 3m+2$  not occurring in one of the above generators.

Its Hilbert function is given by  $(1, 3m, 2)$ . As in the previous example, we conclude that  $\text{Spec}(A)$  is a smoothable scheme, and hence the multiplication tensor  $T_A$  is a tensor of minimal border rank  $3m + 3$ .

Explicitly,  $T_A$  is equal to the following tensor in  $A^* \otimes A^* \otimes A$  (with  $A \cong \mathbb{C}^{3m+3}$ ):

$$\begin{aligned} T_A &= \alpha_0 \otimes \alpha_0 \otimes a_0 + \sum_{i=1}^{3m} (\alpha_0 \otimes \alpha_i \otimes a_i + \alpha_i \otimes \alpha_0 \otimes a_i) \\ &+ \sum_{i=1}^m (\alpha_i \otimes \alpha_{m+i} \otimes a_{3m+1} + \alpha_i \otimes \alpha_{2m+i} \otimes a_{3m+2} + \alpha_{m+i} \otimes \alpha_i \otimes a_{3m+1} + \alpha_{2m+i} \otimes \alpha_i \otimes a_{3m+2}) \\ &+ \alpha_0 \otimes \alpha_{3m+1} \otimes a_{3m+1} + \alpha_0 \otimes \alpha_{3m+2} \otimes a_{3m+2} + \alpha_{3m+1} \otimes \alpha_0 \otimes a_{3m+1} + \alpha_{3m+2} \otimes \alpha_0 \otimes a_{3m+2}. \end{aligned}$$

The space  $L_{T_A}$  consists of all matrices of the form

$$\begin{pmatrix} \lambda_0 & \lambda_1 & \dots & \lambda_m & \lambda_{m+1} & \dots & \lambda_{2m} & \lambda_{2m+1} & \dots & \lambda_{3m} & \lambda_{3m+1} & \lambda_{3m+2} \\ \lambda_1 & & & & \lambda_{3m+1} & & & \lambda_{3m+2} & & & & \\ \vdots & & & & & \ddots & & & \ddots & & & \\ \lambda_m & & & & & & \lambda_{3m+1} & & & & & \\ \lambda_{m+1} & \lambda_{3m+1} & & & & & & & & & & \\ \vdots & & \ddots & & & & & & & & & \\ \lambda_{2m} & & & \lambda_{3m+1} & & & & & & & & \\ \lambda_{2m+1} & \lambda_{3m+2} & & & & & & & & & & \\ \vdots & & \ddots & & & & & & & & & \\ \lambda_{3m} & & & \lambda_{3m+2} & & & & & & & & \\ \lambda_{3m+1} & & & & & & & & & & & \\ \lambda_{3m+2} & & & & & & & & & & & \end{pmatrix}.$$

We point out that this looks exactly like the ‘‘multiplication table’’ of  $A$ . Also, note that if we take the ideal generated by the  $2 \times 2$  minors of this matrix and substitute  $\lambda_0 = 1$  and  $\lambda_i = a_i$  for  $i > 0$ , we recover  $I$ .

We now apply the laser method to  $T_A$ . We take the blocking  $D$  given by

$$A_0^* = \langle \alpha_0 \rangle, A_1^* = \langle \alpha_1, \dots, \alpha_{3m} \rangle, A_2^* = \langle \alpha_{3m+1}, \alpha_{3m+2} \rangle$$

for the first 2 tensor factors, and

$$A_0 = \langle a_{3m+1}, a_{3m+2} \rangle, A_1 = \langle a_1, \dots, a_{3m} \rangle, A_2 = \langle a_0 \rangle$$

for the third tensor factor. The support  $\text{supp}_D T_A$  is again equal to

$$\{(0, 0, 2), (0, 2, 0), (2, 0, 0), (1, 1, 0), (1, 0, 1), (0, 1, 1)\},$$

which is tight. The blocks  $T_{A,(2,0,0)} = M_{\langle 2,1,1 \rangle}$ ,  $T_{A,(0,2,0)} = M_{\langle 1,2,1 \rangle}$ ,  $T_{A,(0,0,2)} = M_{\langle 1,1,1 \rangle}$ ,  $T_{A,(0,1,1)} = M_{\langle 1,3m,1 \rangle}$  and  $T_{A,(1,0,1)} = M_{\langle 1,1,3m \rangle}$  are all matrix multiplication tensors, and hence their values are known. The final block

$$\begin{aligned} T_{A,(1,1,0)} &= \sum_{i=1}^m (\alpha_i \otimes \alpha_{m+i} \otimes a_{3m+1} + \alpha_i \otimes \alpha_{2m+i} \otimes a_{3m+2} \\ &\quad + \alpha_{m+i} \otimes \alpha_i \otimes a_{3m+1} + \alpha_{2m+i} \otimes \alpha_i \otimes a_{3m+2}) \end{aligned}$$

is precisely the same as the block (3.2) from previous section. Hence its value is at least  $4(2m)^\omega$ . We now apply the laser method using this value estimate:

$$\begin{aligned} 3 \log \mathbf{rk}(T_A) &\geq \log V_\omega(\widetilde{T}_A) \geq H(P_1) + H(P_2) + H(P_3) + P(1, 1, 0) \log(4(2m)^\omega) \\ &\quad + (P(2, 0, 0) + P(0, 2, 0)) \log(2^\omega) + (P(1, 0, 1) + P(0, 1, 1)) \log((3m)^\omega). \end{aligned}$$

For fixed  $m$  we can obtain a bound on  $\omega$ , by optimizing over all probability distributions on  $\text{supp}_D T_A$ . The best bound is obtained by putting  $m = 4$ : we obtain  $\omega < 2.431$ .

## 5. TENSOR PRODUCTS OF MONOMIAL ALGEBRAS

To a polynomial  $f \in \mathbb{C}[x_1, \dots, x_n]$  we associate the apolar ideal  $f^\perp \subset \mathbb{C}[\partial_{x_1}, \dots, \partial_{x_n}]$  of differentials that annihilate  $f$  and the algebra  $\mathbb{C}[\partial_{x_1}, \dots, \partial_{x_n}]/f^\perp$ . The last family of tensors we would like to discuss are simply powers of the algebra  $\mathbb{C}[x]/(x^2)$  which is obtained as an apolar to  $f = x$ . Taking Kronecker powers of the tensor corresponds to the tensor product of algebras.

**Lemma 5.1.** *If two algebras are apolar respectively to  $f_1(x_1, \dots, x_n)$  and  $f_2(y_1, \dots, y_m)$  then their tensor product is the algebra apolar to the product  $f_1(x_1, \dots, x_n)f_2(y_1, \dots, y_m)$ .*

*Proof.* We write  $f := f_1(x_1, \dots, x_n)f_2(y_1, \dots, y_m)$  and let  $S_1 := \mathbb{C}[\partial_{x_1}, \dots, \partial_{x_n}]$ ,  $S_2 := \mathbb{C}[\partial_{y_1}, \dots, \partial_{y_m}]$  and  $S = S_1 \otimes_{\mathbb{C}} S_2 = \mathbb{C}[\partial_{x_1}, \dots, \partial_{x_n}, \partial_{y_1}, \dots, \partial_{y_m}]$ . We have  $S_1(f_2^\perp) + S_2(f_1^\perp) \subset f^\perp$  so that we get a surjection  $\pi: (S_1/f_1^\perp) \otimes_{\mathbb{C}} (S_2/f_2^\perp) \rightarrow S/f^\perp$ . But  $S/f^\perp \simeq Sf$  as vector spaces and we see directly that  $\dim(Sf) = \dim(S_1f_1) \cdot \dim(S_2f_2)$ , so  $\pi$  is an isomorphism.  $\blacksquare$

Let  $A_1 = \mathbb{C}[x]/(x^2)$ . Then  $A_2 := A \otimes A$  is apolar to the quadric  $xy$ . In particular, it coincides with the Coppersmith-Winograd tensor  $T_{CW,2}$ . The third tensor power is  $A_3 := A_2 \otimes A_1$ , which is apolar to the cubic  $xyz$ . In matrix representation we have

$$T_{CW,2}^{\otimes 3} = \left( \begin{array}{cc|cc||cc|cc} a_{0,0} & a_{0,1} & a_{1,0} & a_{1,1} & b_{0,0} & b_{0,1} & b_{1,0} & b_{1,1} \\ a_{0,1} & 0 & a_{1,1} & 0 & b_{0,1} & 0 & b_{1,1} & 0 \\ \hline a_{1,0} & a_{1,1} & 0 & 0 & b_{1,0} & b_{1,1} & 0 & 0 \\ a_{1,1} & 0 & 0 & 0 & b_{1,1} & 0 & 0 & 0 \\ \hline b_{0,0} & b_{0,1} & b_{1,0} & b_{1,1} & 0 & 0 & 0 & 0 \\ b_{0,1} & 0 & b_{1,1} & 0 & 0 & 0 & 0 & 0 \\ \hline b_{1,0} & b_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 \\ b_{1,1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right).$$

Switching rows and columns 3 and 4, we obtain:

$$\left( \begin{array}{c|ccc|ccc|c} a & b & c & d & e & f & g & h \\ \hline b & 0 & e & f & 0 & 0 & h & 0 \\ c & e & 0 & g & 0 & h & 0 & 0 \\ d & f & g & 0 & h & 0 & 0 & 0 \\ \hline e & 0 & 0 & h & 0 & 0 & 0 & 0 \\ f & 0 & h & 0 & 0 & 0 & 0 & 0 \\ g & h & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline h & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right).$$

The blocking above is not reconstructible, but we can analyze it using [13, Theorem 4.1]. The only nontrivial value is that of the block

$$\left( \begin{array}{c|cc} 0 & e & f \\ \hline e & 0 & g \\ f & g & 0 \end{array} \right),$$

for which we rerun the laser method using the blocking as above. In the end, the obtained bound is  $\omega < 2.56$ .

However, what is more interesting is that  $A_3 \neq T_{CW,6}$ , but  $A_3$  degenerates to  $T_{CW,6}$ . In the matrix representation above we can multiply rows and columns 2, 3, 4 and 8 by  $t$ ,

and the letters  $b, c, d$  and  $h$  by  $t^{-1}$  (this amounts to acting with diagonal matrices on the corresponding tensor factors). Letting  $t \rightarrow 0$  we obtain

$$\left( \begin{array}{c|ccc|ccc|c} a & b & c & d & e & f & g & h \\ \hline b & 0 & 0 & 0 & 0 & 0 & h & 0 \\ c & 0 & 0 & 0 & 0 & h & 0 & 0 \\ d & 0 & 0 & 0 & h & 0 & 0 & 0 \\ \hline e & 0 & 0 & h & 0 & 0 & 0 & 0 \\ f & 0 & h & 0 & 0 & 0 & 0 & 0 \\ g & h & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline h & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right),$$

which is the matrix representation of  $T_{CW,6}$ . Note that the Hilbert function of  $A_3$  is  $(1, 3, 3, 1)$  whereas the Hilbert function of the local graded algebra corresponding to the CW tensor  $T_{CW,6}$  is  $(1, 6, 1)$ , see Example 4.3.

In particular, this proves that high powers of the  $A_3$  tensor (and hence also  $A_1$ ) degenerate to matrix multiplications that are much larger than suggested by the laser method. In particular, the laser method is very far from optimal when analyzing those tensors. We believe this calls for a new method. One of the reasons why laser method applied to the  $A_3$  tensor is far from the optimal result is the fact that there are many blocks in  $A_3$  that form bigger matrix multiplications (one would say that their phases are adjusted, although this is not required from the input of the algorithm).

#### APPENDIX A. PROOF OF THEOREM 2.10

We present here a proof of our version of the laser method. It is based on the proof of [5, Theorem 15.41] and very similar to the proof of [13, Theorem 4.1]. The version (Theorem A.6) presented in this appendix is even a bit more general than Theorem 2.10: we will consider symmetrization with respect to an arbitrary subgroup of  $S_3$ , instead of only  $\mathbb{Z}/3\mathbb{Z}$ . We will need some preliminary definitions and results.

**Definition A.1** ([5, (15.29)]). Let  $I, J, K$  be finite sets. If  $\Psi \subseteq \Phi \subseteq I \times J \times K$ , we call  $\Psi$  a *combinatorial degeneration* of  $\Phi$ , written  $\Psi \trianglelefteq \Phi$ , if there exist functions  $\alpha : I \rightarrow \mathbb{Z}$ ,  $\beta : J \rightarrow \mathbb{Z}$ ,  $\gamma : K \rightarrow \mathbb{Z}$ , such that  $\alpha(i) + \beta(j) + \gamma(k) = 0$  whenever  $(i, j, k) \in \Psi$ , and  $\alpha(i) + \beta(j) + \gamma(k) > 0$  whenever  $(i, j, k) \in \Phi \setminus \Psi$ .

**Proposition A.2** ([5, (15.30)]). Let  $D$  be a blocking of  $T$ , with components  $T_{(i,j,k)}$ ,  $(i, j, k) \in \text{supp}_D T \subseteq I \times J \times K$ . Let  $\Psi$  be a combinatorial degeneration of  $\text{supp}_D T$ . Then

$$\sum_{(i,j,k) \in \Psi} T_{(i,j,k)} \trianglelefteq T,$$

where  $\trianglelefteq$  denotes the usual tensor degeneration.

**Definition A.3.** Let  $I, J, K$  be finite sets and  $\Delta \subseteq I \times J \times K$ .

- We say  $\Delta$  is a *diagonal*, if the three projections  $\Delta \rightarrow I$ ,  $\Delta \rightarrow J$ ,  $\Delta \rightarrow K$  are injective.
- Recall that  $\Delta$  is *tight*, if there are injections  $\alpha : I \rightarrow \mathbb{Z}^r$ ,  $\beta : J \rightarrow \mathbb{Z}^r$ ,  $\gamma : K \rightarrow \mathbb{Z}^r$  s.t.  $\alpha(i) + \beta(j) + \gamma(k) = 0$  for all  $(i, j, k) \in \Delta$ . If moreover  $\alpha, \beta, \gamma$  can be chosen such that their images are contained in  $\{-b, -b+1, \dots, b-1, b\}^r$ , we say  $\Delta$  is *b-tight*.
- We say  $\Delta$  is *balanced*, if the projection  $p_I : \Delta \rightarrow I$  is surjective, with all fibers of equal cardinality  $|\Delta|/I$ , and similar for the other projections  $p_J, p_K$ .

The following theorem explains the relevance of tight sets: they are sets which are not quite diagonal, but contain a large diagonal which is a combinatorial degeneration. In particular, if a tensor  $T$  has a tight blocking, it degenerates to a large direct sum of a subset of its blocks. Assuming lower bounds on the values of the blocks, this gives a lower bound on the value of  $T$ . The balancedness assumption below is not essential: there is also a version of the theorem without it, but the statement is more complicated.

**Theorem A.4** ([5, (15.39), attributed to Strassen]). *There exists a constant  $C_b$ , only depending on  $b$ , such that every  $b$ -tight balanced subset  $\Phi \subseteq I \times J \times K$  contains a diagonal of size at least  $C_b \cdot \min\{|I|, |J|, |K|\}$ , which is a combinatorial degeneration.*

The entropy will enter in the proof of Theorem A.6 through the following lemma, which is an easy consequence of Stirling's formula.

**Lemma A.5** (See [5, (15.40)]). *Fix a finite set  $S$ . There exists a sequence  $\rho_N$ , with  $\lim_{N \rightarrow \infty} \rho_N = 0$ , such that for every rational probability distribution  $P$  on  $S$  that can be written as  $P(i) = Q(i)/N$  for some  $Q : S \rightarrow \mathbb{N}$ , it holds that*

$$\left| \frac{1}{N} \log \binom{N}{Q} - H(P) \right| \leq \rho_N.$$

Here,  $\binom{N}{Q}$  stands for the appropriate multinomial coefficient, and  $\log$  is the logarithm in base  $e$ . Explicitly: if  $S = \{1, 2, \dots, n\}$ , then  $\binom{N}{Q} = \binom{N}{Q(1), \dots, Q(n)}$ .

The symmetric group  $S_3$  acts on  $U \otimes V \otimes W$  by permuting the factors. Note that for  $\sigma \in S_3$ , the tensor  $\sigma T \in \sigma U \otimes \sigma V \otimes \sigma W$  has a blocking  $\sigma D$ , with components  $(\sigma T)_{(\sigma(i), \sigma(j), \sigma(k))} = \sigma(T_{(i,j,k)})$ . We fix a subgroup  $G \subseteq S_3$ . For any tensor  $T$ , we will denote its *symmetrization*  $\boxtimes_{\sigma \in G} \sigma T$  by  $\tilde{T}$ . Note that  $V_\omega(\tilde{T}) \geq (V_\omega(T))^{|G|}$ , by supermultiplicativity.

**Theorem A.6.** *Let  $T \in U \otimes V \otimes W$ , and let  $D$  be a blocking of  $T$ , indexed by  $I \times J \times K$ . Assume that  $\text{supp}_D T$  is tight and reconstructible. Let  $P$  be any probability distribution on  $\text{supp}_D T$  and let  $G \subseteq S_3$  be a subgroup. Then*

$$(A.1) \quad \log V_\omega(\tilde{T}) \geq \min_{L \in \{I, J, K\}} \sum_{\sigma \in G} H(P_{\sigma L}) + \sum_{\text{supp}_D T} P(i, j, k) \log V_\omega(\widetilde{T_{(i,j,k)}}),$$

where  $\tilde{T}$  denotes the symmetrization with respect to  $G$ .

Theorem 2.10 is the special case of the above theorem with  $G = \mathbb{Z}/3\mathbb{Z}$ .

*Proof.* Assume that  $D$  is  $b$ -tight. We will assume that  $P$  is a rational probability distribution, i.e.  $P(i, j, k) \in \mathbb{Q}$  for all  $(i, j, k)$ . Since a general probability distribution can be approximated by rational ones, this suffices (see also the proof of [5, (15.41)]). There is a map  $Q : \text{supp}_D T \rightarrow \mathbb{N}$  and an  $N \in \mathbb{N}$  such that  $P(i, j, k) = Q(i, j, k)/N$  for all  $(i, j, k)$ . Let  $I_Q \subseteq I^N$  consist of all sequences in which the element  $i$  appears exactly  $Q_I(i) := N \cdot P_I(i) = \sum_{j,k} Q(i, j, k)$  times for all  $i$ . Note that  $|I_Q| = \binom{N}{Q_I}$ . We define  $J_Q \subseteq J^N$  and  $K_Q \subseteq K^N$  analogously.

Write  $\tilde{T} := \boxtimes_{\sigma \in G} (\sigma T)$ , and consider the tensor

$$\tilde{T}^{\boxtimes N} = \boxtimes_{\sigma \in G} (\sigma T)^{\boxtimes N} \in \left( \boxtimes_{\sigma \in G} (\sigma U)^{\boxtimes N} \right) \otimes \left( \boxtimes_{\sigma \in G} (\sigma V)^{\boxtimes N} \right) \otimes \left( \boxtimes_{\sigma \in G} (\sigma W)^{\boxtimes N} \right).$$

Now  $\tilde{T}$  has a blocking  $\tilde{D} := \boxtimes_{\sigma \in G} (\sigma D)^{\boxtimes N}$ , with support

$$\text{supp}_{\tilde{D}} \tilde{T}^{\boxtimes N} = \prod_{\sigma \in G} (\text{supp}_{\sigma D} \sigma T)^N \subseteq \left( \prod_{\sigma \in G} (\sigma I)^N \right) \times \left( \prod_{\sigma \in G} (\sigma J)^N \right) \times \left( \prod_{\sigma \in G} (\sigma K)^N \right),$$

which is again  $b$ -tight. We define

$$\Phi := \left( \left( \prod_{\sigma \in G} \sigma I_Q \right) \times \left( \prod_{\sigma \in G} \sigma J_Q \right) \times \left( \prod_{\sigma \in G} \sigma K_Q \right) \right) \cap \text{supp}_{\tilde{D}} \tilde{T}^{\boxtimes N}.$$

It trivially holds that  $\Phi \trianglelefteq \text{supp}_{\tilde{D}} \tilde{T}^{\boxtimes N}$ .

Let  $(x, y, z) \in \left( \prod_{\sigma \in G} (\sigma I)^N \right) \times \left( \prod_{\sigma \in G} (\sigma J)^N \right) \times \left( \prod_{\sigma \in G} (\sigma K)^N \right)$ , and write

$$(x, y, z) = \left( (i_{\sigma, \ell})_{\sigma \in G, 1 \leq \ell \leq N}, (j_{\sigma, \ell})_{\sigma \in G, 1 \leq \ell \leq N}, (k_{\sigma, \ell})_{\sigma \in G, 1 \leq \ell \leq N} \right).$$

From our reconstructibility assumption, it follows that  $(x, y, z) \in \Phi$  if and only if for every  $\sigma \in G$  and  $(\sigma(i), \sigma(j), \sigma(k)) \in \text{supp}_{\sigma D} \sigma T$ , there are exactly  $Q(i, j, k)$  indices  $\ell$  for which  $(i_{\sigma, \ell}, j_{\sigma, \ell}, k_{\sigma, \ell}) = (\sigma(i), \sigma(j), \sigma(k))$ . We find

$$\begin{aligned} \tilde{T}_{(x,y,z)}^{\boxtimes N} &= \bigotimes_{\sigma, \ell} (\sigma T)_{(i_{\sigma, \ell}, j_{\sigma, \ell}, k_{\sigma, \ell})} = \bigotimes_{\substack{(i,j,k) \in \text{supp}_D T \\ \sigma \in G}} \sigma(T_{(i,j,k)})^{Q(i,j,k)} \\ &= \bigotimes_{(i,j,k) \in \text{supp}_D T} (\widetilde{T_{(i,j,k)}})^{Q(i,j,k)}, \end{aligned}$$

hence by supermultiplicativity:

$$V_{\omega}(\tilde{T}_{(x,y,z)}^{\boxtimes N}) \geq \prod_{(i,j,k) \in \text{supp}_D T} V_{\omega}(\widetilde{T_{(i,j,k)}})^{Q(i,j,k)}.$$

We now apply Theorem A.4 to the balanced  $b$ -tight subset  $\Phi \subseteq (\prod_{\sigma} \sigma I_Q) \times (\prod_{\sigma} \sigma J_Q) \times (\prod_{\sigma} \sigma K_Q)$ , and we find a diagonal  $\Delta \trianglelefteq \Phi$ , such that  $|\Delta| \geq C_b \cdot \min_{L \in \{I, J, K\}} \prod_{\sigma} |\sigma L_Q|$ . Since  $\Phi \trianglelefteq \text{supp}_{\bar{D}} \tilde{T}^{\boxtimes N}$ , we find  $\Delta \trianglelefteq \text{supp}_{\bar{D}} \tilde{T}^{\boxtimes N}$ . So by applying Proposition A.2, we find

$$\bigoplus_{(x,y,z) \in \Delta} \tilde{T}_{(x,y,z)}^{\boxtimes N} \trianglelefteq \tilde{T}^{\boxtimes N}.$$

So we find that

$$V_{\omega}(\tilde{T}) \geq \left( \sum_{(x,y,z) \in \Delta} V_{\omega}(\tilde{T}_{(x,y,z)}^{\boxtimes N}) \right)^{\frac{1}{N}} \geq \left( |\Delta| \prod_{(i,j,k) \in \text{supp}_D T} V_{\omega}(\widetilde{T_{(i,j,k)}})^{Q(i,j,k)} \right)^{\frac{1}{N}}.$$

By taking logarithms, we obtain

$$\log V_{\omega}(\tilde{T}) \geq \frac{1}{N} \min_{L \in \{I, J, K\}} \log \left( C_b \cdot \prod_{\sigma} \binom{N}{Q_{\sigma L}} \right) + \sum_{(i,j,k) \in \text{supp}_D T} P(i, j, k) \log \left( V_{\omega}(\widetilde{T_{(i,j,k)}}) \right).$$

Now the theorem follows by taking  $N \rightarrow \infty$  and applying Lemma A.5.  $\blacksquare$

## REFERENCES

- [1] Josh Alman and Virginia Vassilevska Williams. Limits on all known (and some unknown) approaches to matrix multiplication. In *59th Annual IEEE Symposium on Foundations of Computer Science—FOCS 2018*, pages 580–591. IEEE Computer Soc., Los Alamitos, CA, 2018.
- [2] Andris Ambainis, Yuval Filmus, and François Le Gall. Fast matrix multiplication: Limitations of the coppersmith-winograd method. In *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing, STOC '15*, page 585–593, New York, NY, USA, 2015. Association for Computing Machinery.
- [3] Markus Bläser and Vladimir Lysikov. On degeneration of tensors and algebras. In *41st International Symposium on Mathematical Foundations of Computer Science*, volume 58 of *LIPICs. Leibniz Int. Proc. Inform.*, pages Art. No. 19, 11. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2016.
- [4] Weronika Buczyńska and Jarosław Buczyński. Apolarity, border rank and multigraded Hilbert scheme. *To appear in Duke Math. Journal. arXiv:1910.01944*, 2019.
- [5] Peter Bürgisser, Michael Clausen, and Mohammad A Shokrollahi. *Algebraic complexity theory*, volume 315. Springer Science & Business Media, 2013.
- [6] Dustin A. Cartwright, Daniel Erman, Mauricio Velasco, and Bianca Viray. Hilbert schemes of 8 points. *Algebra Number Theory*, 3(7):763–795, 2009.
- [7] Luca Chiantini, Jonathan D. Hauenstein, Christian Ikenmeyer, Joseph M. Landsberg, and Giorgio Ottaviani. Polynomials and the exponent of matrix multiplication. *Bull. Lond. Math. Soc.*, 50(3):369–389, 2018.
- [8] Matthias Christandl, Péter Vrana, and Jeroen Zuiddam. Barriers for fast matrix multiplication from irreversibility. In *34th Computational Complexity Conference*, volume 137 of *LIPICs. Leibniz Int. Proc. Inform.*, pages Art. No. 26, 17. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2019.
- [9] Don Coppersmith and Shmuel Winograd. Matrix multiplication via arithmetic progressions. *J. Symbolic Comput.*, 9(3):251–280, 1990.
- [10] J. M. Landsberg. *Tensors: geometry and applications*, volume 128 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012.
- [11] J. M. Landsberg. *Geometry and Complexity Theory*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2017.

- [12] Joseph M Landsberg and Mateusz Michałek. Abelian tensors. *Journal de Mathématiques Pures et Appliquées*, 108(3):333–371, 2017.
- [13] François Le Gall. Powers of tensors and fast matrix multiplication. In *Proceedings of the 39th international symposium on symbolic and algebraic computation*, pages 296–303, 2014.
- [14] Mateusz Michałek and Bernd Sturmfels. *Invitation to nonlinear algebra*, volume 211, Graduate Studies in Mathematics. American Mathematical Society, 2019.
- [15] A. Schönhage. Partial and total matrix multiplication. *SIAM J. Comput.*, 10(3):434–455, 1981.
- [16] Tim Seynnaeve. Plethysm and fast matrix multiplication. *C. R. Math. Acad. Sci. Paris*, 356(1):52–55, 2018.
- [17] Tim Seynnaeve. *Algebraic geometry for tensor networks, matrix multiplication, and flag matroids*. PhD thesis, Leipzig University, 2020. Available at <https://nbn-resolving.org/urn:nbn:de:bsz:15-qucosa2-732863>.
- [18] Andrew James Stothers. On the complexity of matrix multiplication. 2010. PhD thesis.
- [19] V. Strassen. Relative bilinear complexity and matrix multiplication. *J. Reine Angew. Math.*, 375/376:406–443, 1987.
- [20] Virginia Vassilevska Williams. Multiplying matrices faster than coppersmith-winograd. In *Proceedings of the forty-fourth annual ACM symposium on Theory of computing*, pages 887–898, 2012.

TECHNICAL UNIVERSITY OF MUNICH, DEPARTMENT OF MATHEMATICS, PARKRING 13, 85748 GARCHING BEI MÜNCHEN, GERMANY

*Email address:* `roser.homs@tum.de`

FACULTY OF MATHEMATICS, INFORMATICS AND MECHANICS, UNIVERSITY OF WARSAW, BANACHA 2, 02-097 WARSAW

*Email address:* `jjelisiejew@mimuw.edu.pl`

UNIVERSITY OF KONSTANZ, GERMANY, FACHBEREICH MATHEMATIK UND STATISTIK, FACH D 197 D-78457 KONSTANZ, GERMANY

*Email address:* `mateusz.michalek@uni-konstanz.de`

UNIVERSITY OF BERN, MATHEMATICAL INSTITUTE, SIDLERSTRASSE 5, 3012 BERN, SWITZERLAND

*Email address:* `tim.seynnaeve@math.unibe.ch`