

Bulk reconstruction of metrics inside black holes by complexity

Koji Hashimoto, Ryota Watanabe

*Department of Physics, Osaka University,
1-1 Machikaneyama, Toyonaka, Osaka 560-0043, Japan*

E-mail: koji@phys.sci.osaka-u.ac.jp,
watanabe@het.phys.sci.osaka-u.ac.jp

ABSTRACT: We provide a formula to reconstruct bulk spacetime metrics inside black holes by the time dependence of complexity in the dual quantum field theory, based on the complexity=volume (CV) conjecture in the holographic duality.

Contents

1	Introduction	1
2	Reconstructing metrics inside horizons by complexity	3
2.1	Review: calculation of the volume	3
2.2	Reconstruction formula	5
3	General case	7
3.1	Preparations	7
3.1.1	Volume of extremal surfaces in generic spacetimes	7
3.1.2	The Hartman-Maldacena entanglement entropy	9
3.2	Reconstruction formula	11
4	Example: BTZ black hole	12
5	Summary and discussions	14
A	Treatment of the special case $g = g(f)$	16

1 Introduction

It is expected that the holographic principle and the AdS/CFT correspondence [1–3] will serve as a solution to the information loss problem of black holes. For that program to be complete, it is indispensable to find out the way how the information inside black holes is encoded in the dual quantum field theories. Can we construct the spacetime metric inside a black hole horizon by a given data of the dual quantum field theory?

Bulk reconstruction is in general an inverse problem in holography, thus is difficult. Nevertheless, various methods have been developed to reconstruct bulk spacetime metrics by the data of dual quantum field theories (QFTs)¹. Successful methods include the holographic renormalization [5], the reconstruction using bulk geodesics and light cones [6–12], the reconstruction [13–26] using holographic entanglement entropy [27, 28]², the inversion formula [35] of the holographic Wilson loops [36–39], and the machine learning holography [40–45]³. However, all of these methods do not reconstruct the black hole interior, because the dual thermal QFT data is associated with the asymptotic boundary of the bulk spacetime, and the bulk classical gravity does not extract information inside the horizon.

¹Here we focus on only the bulk reconstruction of spacetime metrics. Reconstruction of operators in the bulk AdS spacetimes [4] have wider references.

²Related methods include the one [29–32] using tensor networks [33, 34] through the entanglement properties.

³The holographic bulk spacetime is identified with neural networks [40–49], and the spacetimes are emergent. See [50] for a review of data science approach to string theory, and also see [51] for applications of machine learning to material sciences.

The QFT quantity to probe the inside of the bulk black holes in holography is the quantum computational complexity, proposed by Susskind [52–54]. The conjecture originates from the idea using two-sided extended black hole geometry interpreted as a doubled QFT with thermo-field double states [55] and the use of the Ryu-Takayanagi (RT) surface connecting the left and right boundaries of the geometry [56]. There are two versions of the complexity conjecture, one is the “complexity = volume” (CV) conjecture [57, 58], and the other is the “complexity = action” (CA) conjecture [59, 60]. Both conjectures can probe the interior of the horizons of the maximally extended black hole spacetimes. Therefore, the question of how to reconstruct even the interior geometry of the black hole can be cast into a problem of unifying the bulk reconstruction techniques and the complexity conjecture.

In this paper, based on the CV conjecture, we provide a formula to reconstruct the spacetime metric of the black hole interior by the time dependence of the complexity $\mathcal{C}(t)$ of the dual QFT. The CV conjecture [57, 58] states that the quantum computational complexity of a thermo-field double state at a time in the dual QFT is equal to the volume of an extremal surface in the bulk penetrating the horizon and anchored to the boundary at the QFT time. Among CV and CA conjectures, we employ the CV conjecture, because in the bulk side it is about extremal surfaces, and therefore it allows a similarity to the RT surface and the holographic Wilson loops. For the latter quantities there exist the particular bulk reconstruction techniques for the spacetime geometry [14, 15, 35] to invert the area of extremal surfaces back to the metric. By generalizing the technique⁴ to the case of the extremal surfaces penetrating the black hole horizons, we will be able to reconstruct the metric of the black hole interior by the complexity of the dual QFT.

We focus on generic black hole spacetimes which are static, spherically symmetric and asymptotically AdS. They have the metric

$$ds^2 = -f(r)dt^2 + g(r)dr^2 + r^2d\Sigma^2. \quad (1.1)$$

Let us make technical remarks on our reconstruction. First, because the metric includes two functional degrees of freedom $f(r)$ and $g(r)$, the complexity $\mathcal{C}(t)$ can determine only one functional degree of freedom in the metric. Thus, when $g(r)$ and $f(r)$ are related, the complexity $\mathcal{C}(t)$ can reconstruct the metric. On the other hand, when they are independent, we need another QFT data, and we employ the Hartman-Maldacena entanglement entropy $S_{\text{HM}}(t)$ [56] together with the complexity $\mathcal{C}(t)$. Second, in this paper we are interested in only the interior of the black hole, so we assume that the metric outside the horizon is known by some methods listed above.

This paper is organized as follows. In Sec. 2, we provide our formula for reconstructing bulk metrics inside the black hole horizons by the time dependence of the complexity $\mathcal{C}(t)$ in the dual QFT, when $g(r) = 1/f(r)$. In Sec. 3, we provide our formula for the general case of unknown $f(r)$ and $g(r)$, by the complexity $\mathcal{C}(t)$ and the Hartman-Maldacena entanglement entropy $S_{\text{HM}}(t)$. In Sec. 4, to illustrate the validity of our formulas, we verify

⁴Note that the interesting interplay found for the RT surface and the holographic Wilson loop in thermalization and confinement [61–64] is about the outside of the black holes.

the reconstruction of the interior of the BTZ black hole. Sec. 5 is for our summary and discussions. App. A treats a special case when $g(r)$ is known as a function of $f(r)$ implicitly, $g = g(f)$.

2 Reconstructing metrics inside horizons by complexity

In this section, we provide the formula to reconstruct the interior of a black hole by the complexity in the dual QFT, based on the complexity=volume (CV) conjecture [53, 57]

$$\mathcal{C}(t) = \frac{V(t)}{G_N L}. \quad (2.1)$$

Here, $V(t)$ is the volume of the extremal surface defined in the two-sided eternal AdS black holes. The surface is anchored at time t at the two boundaries of the spacetime. G_N is the Newton's constant and L is the AdS radius.

2.1 Review: calculation of the volume

We here first summarize the known results on the calculation of the volume $V(t)$ in the gravity side, for a given metric. Following [65], in this section we consider the metric of the form⁵

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\Sigma_{k,d-1}^2. \quad (2.2)$$

The dimension of the bulk spacetime is $d+1$, and $k = \{+1, 0, -1\}$ indicates the curvature of the $(d-1)$ -dimensional line element $d\Sigma_{k,d-1}^2$. The geometry is asymptotically AdS, $f(r) \rightarrow \frac{r^2}{L^2}$ as $r \rightarrow \infty$.

The function $f(r)$ may be determined as a solution of Einstein equations. However, for our purpose of the reconstruction, we keep $f(r)$ to be a generic function. The metric (2.2) corresponds to having $g(r) = 1/f(r)$ in (1.1). For a black holes with a nonzero value of the temperature, in the vicinity of the horizon the function $f(r)$ is linear, and at the horizon $r = r_h$ we have $f(r = r_h) = 0$.

Of our interest is the codimension-one extremal surface which penetrates the horizon and connects the left and right boundaries in the maximally extended geometry. Particularly, we are interested in the case where the boundary times t_L and t_R are the same, $t_L = t_R$, so that the surface is left-right symmetric⁶. As in [65], we can express the volume V of the extremal surface and the boundary time t_R in terms of the metric function $f(r)$ and the minimal radius $r_{\min} < r_h$ of the surface⁷,

$$V = 2\Omega_{k,d-1} \int_{r_{\min}}^{r_{\max}} dr \frac{r^{2(d-1)}}{\sqrt{F(r) - F(r_{\min})}}, \quad (2.3)$$

⁵Our notations follow those in [65].

⁶We define t_L and t_R in such a way that both of them run from the past to the future.

⁷In Sec. 3.1, we will explicitly derive them for a more general metric. See (3.10) and (3.13).

$$t_{\text{R}} = \int_{r_{\text{min}}}^{r_{\text{max}}} dr \frac{r^{2(d-1)}}{F(r)} \frac{-\sqrt{-F(r_{\text{min}})}}{\sqrt{F(r) - F(r_{\text{min}})}}, \quad (2.4)$$

where $F(r) \equiv f(r)r^{2(d-1)}$, and r_{max} is a UV cutoff⁸. There are two important comments on these equations: First, since r_{min} is at the interior of the black hole, $F(r_{\text{min}}) = f(r_{\text{min}})r_{\text{min}}^{2(d-1)}$ is negative. Second, although the integrand of (2.4) blows up near the horizon $r \sim r_{\text{h}}$, we can define the integral in terms of the Cauchy principal value.

In the time evolution, the surface gets stretched and the volume grows. The growth rate dV/dt_{R} can be found as

$$\frac{dV}{dt_{\text{R}}} = 2\Omega_{k,d-1} \sqrt{-F(r_{\text{min}})}. \quad (2.5)$$

At late times, this asymptotically approaches the constant value

$$\lim_{t_{\text{R}} \rightarrow \infty} \frac{dV}{dt_{\text{R}}} = 2\Omega_{k,d-1} \sqrt{-F(r_{\text{f}})}. \quad (2.6)$$

Here, $r = r_{\text{f}}$ is the radius at which $\sqrt{-F(r)}$ is maximal⁹. As $t_{\text{R}} \rightarrow \infty$, the extremal surface asymptotically approaches this $r = r_{\text{f}}$ surface (called a “nice” slice), which in turn means that r_{min} is always greater than r_{f} .

If we knew the functional form of the metric $F(r)$, then eliminating r_{min} from (2.4) and (2.5) would determine the complexity growth $\dot{\mathcal{C}}(t_{\text{R}}) = \frac{dV}{dt_{\text{R}}}(t_{\text{R}})/G_{\text{NL}}$. Our interest in this paper is the opposite, *i.e.* the case when the metric $F(r)$ is unknown: can we reconstruct the function $F(r)$ by a given function $\frac{dV}{dt_{\text{R}}}(t_{\text{R}})$? The answer is yes, and in the next subsection we will provide the formula for the reconstruction.

⁸The integral (2.3) is divergent for $r_{\text{max}} \rightarrow \infty$ while that in (2.4) is convergent.

⁹When there are several extremal radii of $\sqrt{-F(r)}$, this $r = r_{\text{f}}$ is the one closest to the horizon. In other words, we do not consider a sudden jump in the complexity in its time evolution.

2.2 Reconstruction formula

Let us provide the reconstruction formula of the metric inside horizons by complexity.

Assume that

- the time derivative of the complexity $\dot{\mathcal{C}}(t)$ in the dual QFT is given.
- $\dot{\mathcal{C}}(t)$ is monotonic in time.
- the metric $f(r)$ of (2.2) outside the horizon ($r \geq r_h$) is known.

Then, solve

$$\frac{d\mathcal{C}}{dt_{\text{R}}}(t_{\text{R}}) = \frac{2\Omega_{k,d-1}}{G_{\text{N}}L} \sqrt{-F_{\text{min}}} \quad (2.7)$$

to get t_{R} as a function of F_{min} . In addition, outside of the horizon, invert $F = F(r) \equiv f(r)r^{2(d-1)}$ to find $r = r(F)$. Substitute them into

$$Q(F_{\text{min}}) \equiv \frac{t_{\text{R}}(F_{\text{min}})}{\sqrt{-F_{\text{min}}}} + \int_{\epsilon}^{\infty} dF \frac{dr}{dF} \frac{r(F)^{2(d-1)}}{F} \frac{1}{\sqrt{F - F_{\text{min}}}}, \quad (2.8)$$

with a positive infinitesimal parameter ϵ . For $F \leq -\epsilon$, calculate

$$y(F) \equiv \frac{1}{\pi} \frac{d}{dF} \int_F^{-\epsilon} dU \frac{Q(U)}{\sqrt{U - F}}, \quad (2.9)$$

and integrate the differential equation

$$\frac{dr}{dF} \frac{r(F)^{2(d-1)}}{F} = y(F) \quad (2.10)$$

to find $r = r(F)$. Finally, invert $r(F)$ to find $F(r)$, and the metric inside the black hole is given as $f(r) = F(r)r^{-2(d-1)}$.

Proof. First, (2.7) is nothing but (2.1) and (2.5) with $F_{\text{min}} \equiv F(r_{\text{min}}) < 0$. Next, we consider rewriting (2.4). Assuming that $F(r)$ outside the horizon is monotonic in r , $F = F(r)$ can be inverted to a single-valued function $r = r(F)$. Then we can transform the integration variable of (2.4) from r to F :

$$t_{\text{R}}(F_{\text{min}}) = \int_{F_{\text{min}}}^{\infty} dF \frac{dr}{dF} \frac{r(F)^{2(d-1)}}{F} \frac{-\sqrt{-F_{\text{min}}}}{\sqrt{F - F_{\text{min}}}}. \quad (2.11)$$

To make the expression simpler, we define the function

$$y(F) \equiv \frac{dr}{dF} \frac{r(F)^{2(d-1)}}{F}. \quad (2.12)$$

As pointed out after (2.4), we interpret (2.11) as the Cauchy principal value. Introducing an infinitesimally small parameter $\epsilon > 0$, we rewrite (2.11) as

$$-\frac{t_{\text{R}}(F_{\text{min}})}{\sqrt{-F_{\text{min}}}} = \left(\int_{F_{\text{min}}}^{-\epsilon} + \int_{-\epsilon}^{\epsilon} + \int_{\epsilon}^{\infty} \right) dF \frac{y(F)}{\sqrt{F - F_{\text{min}}}}. \quad (2.13)$$




Figure 1. The typical r dependence of $F(r) = f(r)r^{2(d-1)}$ for $d > 2$.

If we take ϵ small enough, the second integral $\int_{-\epsilon}^{\epsilon}$ cancels by itself, and we obtain

$$-\frac{t_{\text{R}}(F_{\text{min}})}{\sqrt{-F_{\text{min}}}} = \left(\int_{F_{\text{min}}}^{-\epsilon} + \int_{\epsilon}^{\infty} \right) dF \frac{y(F)}{\sqrt{F - F_{\text{min}}}}. \quad (2.14)$$

We rewrite this as

$$\int_{F_{\text{min}}}^{-\epsilon} dF \frac{y(F)}{\sqrt{F - F_{\text{min}}}} = -\frac{t_{\text{R}}(F_{\text{min}})}{\sqrt{-F_{\text{min}}}} - \int_{\epsilon}^{\infty} dF \frac{y(F)}{\sqrt{F - F_{\text{min}}}} \equiv Q(F_{\text{min}}). \quad (2.15)$$

This is nothing but (2.8). Since we do not know the metric inside the black hole, the left-hand side is unknown. On the other hand, the right-hand side can be calculated if we know the complexity and the metric outside the horizon. As can be easily checked, the solution of this integral equation is

$$y(F) = \frac{1}{\pi} \frac{d}{dF} \int_F^{-\epsilon} dU \frac{Q(U)}{\sqrt{U - F}}. \quad (2.16)$$

This provides the metric inside the horizon. □

Several comments on the formula are in order. In the proof above, first, we need to know the metric outside the horizon. In Sec. 5 we discuss it is able to be reconstructed by the methods listed in Sec. 1 using some QFT data. Second, in the proof, we have assumed¹⁰ that the function $F(r)$ outside the horizon is monotonic in r . The reconstructed $F(r)$ inside the horizon is also monotonic, which follows from the construction method. For example, in the AdS Schwarzschild geometry, $F(r)$ behaves like Fig. 1, and in the region $r_{\text{h}} > r \geq r_{\text{f}}$ it is monotonic. Our reconstruction builds the spacetime of the region $r_{\text{h}} > r \geq r_{\text{f}}$, not the whole interior of the black hole. In particular, in general the reconstruction of the black hole singularity is not possible, which is physically sensible.

¹⁰It is worthwhile to mention that the formula can be generalized such that the monotonicity of $F(r)$ outside the horizon is not necessary. The second term in (2.8) can be written as an integral over r rather than the integral over F . So even with $F(r)$ which is not monotonic outside the horizon, the evaluation of Q in (2.8) is possible. The boundary of the integration region $F = \epsilon$ needs to be rephrased in terms of r by the known functional form of $F(r)$.

3 General case

In the previous section, we have demonstrated the reconstruction of the metric by the complexity assuming that the metric takes the particular form (2.2). In general the static and spherically symmetric black hole metric is of the form

$$ds^2 = -f(r)dt^2 + g(r)dr^2 + r^2 d\Sigma_{k,d-1}^2, \quad (3.1)$$

where $f(r)$ and $g(r)$ are independent. In this section, we provide a reconstruction formula to determine both $f(r)$ and $g(r)$.¹¹

As is described in the introduction, the reconstruction in the previous section with (2.2) is possible since the time dependence of the complexity provided just enough degree of freedom to reconstruct a single unknown metric function $f(r)$. Therefore, for the reconstruction of both $f(r)$ and $g(r)$, we need another functional information in addition to the complexity. As such a quantity, we use the Hartman-Maldacena entanglement entropy [56].

3.1 Preparations

To derive our reconstruction formula, in this subsection we first formulate the CV conjecture and the Hartman-Maldacena entanglement entropy in generic black hole spacetimes (3.1).

3.1.1 Volume of extremal surfaces in generic spacetimes

We express the volume of the extremal codimension-one surface using $f(r)$ and $g(r)$. This can be worked out by slightly generalizing the calculations in [65]. Again, let $r = r_h$ be the position of the horizon. For black holes with a nonzero value of the temperature, $f(r)$ and $1/g(r)$ are linear near the horizon and vanish at the horizon. The spacetime is asymptotically AdS with radius L , $f(r) \rightarrow \frac{r^2}{L^2}$, $g(r) \rightarrow \frac{L^2}{r^2}$ as $r \rightarrow \infty$. We introduce the tortoise coordinate as

$$dr^* \equiv \frac{\sqrt{f(r)g(r)}}{f(r)} dr, \quad (3.2)$$

and then define the incoming Eddington-Finkelstein coordinate $v \equiv t + r^*$. In terms of them, the metric (3.1) is written as

$$ds^2 = -f(r)dv^2 + 2\sqrt{f(r)g(r)}dvdr + r^2 d\Sigma_{k,d-1}^2. \quad (3.3)$$

Now, let us consider the codimension-one surface which penetrates the horizon and connects the left and right boundaries in the maximally extended geometry. We focus on the

¹¹See App. A for the reconstruction formula for the case when the relation between f and g is known, $g = g(f)$.

symmetric configuration, $t_L = t_R$. The profile of the surface is determined by $v(\lambda)$ and $r(\lambda)$, where λ is a coordinate intrinsic to the surface. Then, the induced metric becomes

$$ds^2 = \left(-f(r)\dot{v}^2 + 2\sqrt{f(r)g(r)}\dot{v}\dot{r} \right) d\lambda^2 + r^2 d\Sigma_{k,d-1}^2, \quad (3.4)$$

where the dots represent $d/d\lambda$. We assume that, when the surface volume is extremized, $r(\lambda)$ is a strictly monotonically increasing function, $\dot{r} > 0$. We set $\lambda = 0$ at the left-right symmetric point on the surface, where the radius of the surface $r(\lambda)$ takes the minimum value, $r(\lambda = 0) = r_{\min} < r_h$. Smoothness requires that $\dot{r}(\lambda = 0) = 0$. Note that r_{\min} varies depending on the boundary time t_R . The volume of the surface is

$$\begin{aligned} V &= 2 \int d\lambda d\Sigma_{k,d-1} \sqrt{\left(-f(r)\dot{v}^2 + 2\sqrt{f(r)g(r)}\dot{v}\dot{r} \right)} r^{2(d-1)} \\ &= 2\Omega_{k,d-1} \int_0^{\lambda_{\max}} d\lambda \sqrt{-F(r)\dot{v}^2 + 2\sqrt{F(r)G(r)}\dot{v}\dot{r}}, \end{aligned} \quad (3.5)$$

where $F(r) \equiv f(r)r^{2(d-1)}$, $G(r) \equiv g(r)r^{2(d-1)}$ and λ_{\max} corresponds to the cutoff $r = r_{\max}$ necessary for the volume to be well-defined.

Since we want to apply the CV conjecture, we need to extremize the volume while the boundary of the surface is fixed. To work it out, instead of solving the equations of motion for $v(\lambda)$ and $r(\lambda)$, we make use of a conserved quantity and the reparametrization invariance. Since the integrand of (3.5) is written only with \dot{v} and does not contain v , when the volume is extremal,

$$E \equiv \frac{F(r)\dot{v} - \sqrt{F(r)G(r)}\dot{r}}{\sqrt{-F(r)\dot{v}^2 + 2\sqrt{F(r)G(r)}\dot{v}\dot{r}}} \quad (3.6)$$

does not depend on λ , *i.e.*, is ‘‘conserved’’. Note that E does depend on the boundary time t_R . Also, since (3.5) is invariant under the reparametrization of λ , we choose a gauge

$$\sqrt{-F(r)\dot{v}^2 + 2\sqrt{F(r)G(r)}\dot{v}\dot{r}} = 1. \quad (3.7)$$

From (3.6) and (3.7), we obtain

$$\dot{r} = \sqrt{\frac{F(r) + E^2}{F(r)G(r)}}, \quad \frac{\dot{v}}{\dot{r}} = \frac{\sqrt{F(r)G(r)}}{F(r)} \left(\frac{E}{\sqrt{F(r) + E^2}} + 1 \right). \quad (3.8)$$

Recalling $r(0) = r_{\min} < r_h$ and $\dot{r}(0) = 0$, the first equation of (3.8) with $\lambda = 0$ yields

$$E^2 = -F(r_{\min}) > 0. \quad (3.9)$$

As we will see shortly, E must be negative. Substituting (3.7) and (3.8) into (3.5), the extremal volume is written as

$$\begin{aligned} V &= 2\Omega_{k,d-1} \int_{r_{\min}}^{r_{\max}} \frac{dr}{\dot{r}} \\ &= 2\Omega_{k,d-1} \int_{r_{\min}}^{r_{\max}} dr \sqrt{\frac{F(r)G(r)}{F(r) + E^2}}. \end{aligned} \quad (3.10)$$

On the other hand, integrating $dt + dr^* = dv$ over $0 \leq \lambda \leq \lambda_{\max}$ and using (3.2) and (3.8), we obtain

$$\begin{aligned} t_{\text{R}} - t(0) + \int_{r_{\min}}^{r_{\max}} dr \frac{\sqrt{F(r)G(r)}}{F(r)} &= \int_{r_{\min}}^{r_{\max}} dr \frac{\dot{v}}{\dot{r}} \\ &= \int_{r_{\min}}^{r_{\max}} dr \frac{\sqrt{F(r)G(r)}}{F(r)} \left(\frac{E}{\sqrt{F(r) + E^2}} + 1 \right). \end{aligned} \quad (3.11)$$

Due to the symmetry of the spacetime, we find $t(0) = 0$. Near the horizon, the integrand of the integral on the left-hand side behaves like

$$\frac{\sqrt{F(r)G(r)}}{F(r)} \propto \frac{1}{r - r_{\text{h}}}. \quad (3.12)$$

Hereafter, we regard the integral as the Cauchy principal value, which is finite. Then, (3.11) simplifies to

$$t_{\text{R}} = \int_{r_{\min}}^{r_{\max}} dr \frac{\sqrt{F(r)G(r)}}{F(r)} \frac{E}{\sqrt{F(r) + E^2}}. \quad (3.13)$$

It is natural to assume that r_{\min} decreases as t_{R} increases since the wormhole should be more stretched in the time evolution. With (3.13), this implies that E is negative, and (3.9) yields

$$E = -\sqrt{-F(r_{\min})}. \quad (3.14)$$

Multiplying (3.13) by E , we get

$$Et_{\text{R}} = \int_{r_{\min}}^{r_{\max}} dr \frac{\sqrt{F(r)G(r)}}{F(r)} \sqrt{F(r) + E^2} - \frac{V}{2\Omega_{k,d-1}}. \quad (3.15)$$

Taking the derivative with respect to t_{R} , we obtain

$$\frac{dV}{dt_{\text{R}}} = -2\Omega_{k,d-1}E = 2\Omega_{k,d-1}\sqrt{F(r_{\min})}. \quad (3.16)$$

At late times, this asymptotically approaches the constant value

$$\lim_{t_{\text{R}} \rightarrow \infty} \frac{dV}{dt_{\text{R}}} = 2\Omega_{k,d-1}\sqrt{-F(r_{\text{f}})}. \quad (3.17)$$

Here, $r = r_{\text{f}}$ is the radius where $\sqrt{-F(r)}$ is maximal. As $t_{\text{R}} \rightarrow \infty$, the extremal surface asymptotically approaches this $r = r_{\text{f}}$ surface.

3.1.2 The Hartman-Maldacena entanglement entropy

Presently, f and g are treated as being independent with each other. Therefore, reconstructing both of them requires as the inputs two functional degrees of freedom, one of which is given by the complexity. In the present paper, as an additional quantity as supplement, we use the Hartman-Maldacena entanglement entropy S_{HM} [56].

Holographically, S_{HM} is given by the volume A of the Ryu-Takayanagi surface [27] as

$$S_{\text{HM}} = \frac{A}{4G_{\text{N}}}. \quad (3.18)$$

Repeating the calculations in Sec. 3.1, we obtain

$$A = 2\Omega_{k,d-2} \int_{\tilde{r}_{\text{min}}}^{r_{\text{max}}} dr \sqrt{\frac{\tilde{F}(r)\tilde{G}(r)}{\tilde{F}(r) - \tilde{F}_{\text{min}}}}, \quad (3.19)$$

$$t_{\text{R}} = \int_{\tilde{r}_{\text{min}}}^{r_{\text{max}}} dr \frac{\sqrt{\tilde{F}(r)\tilde{G}(r)}}{\tilde{F}(r)} \frac{-\sqrt{-\tilde{F}(\tilde{r}_{\text{min}})}}{\sqrt{\tilde{F}(r) - \tilde{F}(\tilde{r}_{\text{min}})}}, \quad (3.20)$$

where $\tilde{F}(r) \equiv f(r)r^{2(d-2)}$, $\tilde{G}(r) \equiv g(r)r^{2(d-2)}$. Note that the only difference from the equations for the complexity in Sec. 3.1.1 is the power of r in the definitions of $\tilde{F}(r)$ and $\tilde{G}(r)$. Resultantly, \tilde{r}_{min} may be different from r_{min} of the codimension-one extremal surface.

Also, we can show

$$\frac{dA}{dt_{\text{R}}} = 2\Omega_{k,d-2} \sqrt{-\tilde{F}(\tilde{r}_{\text{min}})}. \quad (3.21)$$

At late times, this asymptotically approaches the constant value

$$\lim_{t_{\text{R}} \rightarrow \infty} \frac{dA}{dt_{\text{R}}} = 2\Omega_{k,d-2} \sqrt{-\tilde{F}(\tilde{r}_{\text{f}})}. \quad (3.22)$$

Here, $r = \tilde{r}_{\text{f}}$ is the radius at which $\sqrt{-\tilde{F}(r)}$ is maximal. As $t_{\text{R}} \rightarrow \infty$, the extremal surface asymptotically approaches this $r = \tilde{r}_{\text{f}}$ surface.

We here have seen that all the equations for the Hartman-Maldacena entanglement entropy are of the same form as those for the complexity of the CV conjecture. The only difference is in the power of r in the integrands. For the reconstruction, we can use this slight difference to extract the information of $f(r)$ and $g(r)$ independently.

3.2 Reconstruction formula

Assume that

- the time derivative of the complexity $\dot{\mathcal{C}}(t)$ and the time derivative of the Hartman-Maldacena entanglement entropy $\dot{S}_{\text{HM}}(t)$ in the dual QFT are given.
- $\dot{\mathcal{C}}(t)$ and $\dot{S}_{\text{HM}}(t)$ are monotonic in time.
- the metric (3.1) outside the horizon ($r \geq r_{\text{h}}$) is known.

Then, solve

$$\frac{d\mathcal{C}}{dt_{\text{R}}}(t_{\text{R}}) = \frac{2\Omega_{k,d-1}}{G_{\text{NL}}} \sqrt{-F_{\text{min}}}, \quad \frac{dS_{\text{HM}}}{d\tilde{t}_{\text{R}}}(\tilde{t}_{\text{R}}) = \frac{\Omega_{k,d-2}}{2G_{\text{N}}} \sqrt{-\tilde{F}_{\text{min}}} \quad (3.23)$$

to get $t_{\text{R}} = t_{\text{R}}(F_{\text{min}})$ and $\tilde{t}_{\text{R}} = \tilde{t}_{\text{R}}(\tilde{F}_{\text{min}})$. In addition, outside the horizon, invert $F = F(r) \equiv f(r)r^{2(d-1)}$ and $\tilde{F} = \tilde{F}(r) \equiv f(r)r^{2(d-2)}$ to find $r = r(F)$ and $r = \tilde{r}(\tilde{F})$. Substitute them into

$$Q(F_{\text{min}}) \equiv \frac{t_{\text{R}}(F_{\text{min}})}{\sqrt{-F_{\text{min}}}} + \int_{\epsilon}^{\infty} dF \frac{dr}{dF} \frac{\sqrt{FG(r(F))}}{F} \frac{1}{\sqrt{F - F_{\text{min}}}}, \quad (3.24)$$

$$\tilde{Q}(\tilde{F}_{\text{min}}) \equiv \frac{\tilde{t}_{\text{R}}(\tilde{F}_{\text{min}})}{\sqrt{-\tilde{F}_{\text{min}}}} + \int_{\delta}^{\infty} d\tilde{F} \frac{d\tilde{r}}{d\tilde{F}} \frac{\sqrt{\tilde{F}\tilde{G}(\tilde{r}(\tilde{F}))}}{\tilde{F}} \frac{1}{\sqrt{\tilde{F} - \tilde{F}_{\text{min}}}}, \quad (3.25)$$

with positive infinitesimal ϵ and δ , and $G(r) \equiv g(r)r^{2(d-1)}$ and $\tilde{G}(r) \equiv g(r)r^{2(d-2)}$. For $F \leq -\epsilon$ and $\tilde{F} \leq -\delta$, calculate

$$y(F) \equiv \frac{1}{\pi} \frac{d}{dF} \int_F^{-\epsilon} dU \frac{Q(U)}{\sqrt{U - F}}, \quad (3.26)$$

$$\tilde{y}(\tilde{F}) \equiv \frac{1}{\pi} \frac{d}{d\tilde{F}} \int_{\tilde{F}}^{-\delta} d\tilde{U} \frac{\tilde{Q}(\tilde{U})}{\sqrt{\tilde{U} - \tilde{F}}}. \quad (3.27)$$

Using these, solve

$$\frac{d\tilde{F}}{dF} = \frac{\tilde{y}(\tilde{F})}{y(F)} \quad (3.28)$$

to find \tilde{F} as a function of F . Substituting it into $r = F/\tilde{F}$, we obtain $r = r(F)$, from which $f(r)$ is obtained. Finally, substitute $r = r(F)$ into the equation

$$\frac{dr}{dF} \frac{\sqrt{FG(r(F))}}{F} = y(F), \quad (3.29)$$

and solve it for $G(r(F))$. From that, $g(r)$ is obtained. This determines the metric inside the horizon, $f(r)$ and $g(r)$ for $r < r_{\text{h}}$.

Proof. For the logic until (3.27), read the proof of Sec. 2. Then the following differential

equations are derived which the metric functions should satisfy,

$$\frac{dr}{dF} \frac{\sqrt{FG(r(F))}}{F} = y(F), \quad \frac{d\tilde{r}}{d\tilde{F}} \frac{\sqrt{\tilde{F}\tilde{G}(\tilde{r}(\tilde{F}))}}{\tilde{F}} = \tilde{y}(\tilde{F}). \quad (3.30)$$

The first equation is the same as (3.29). Note that F and \tilde{F} are not independent but constrained by $r(F) = \tilde{r}(\tilde{F})$. Using $F(r) = r\tilde{F}(r)$ and $G(r) = r\tilde{G}(r)$, we find that (3.30) yields (3.28), which is the last nontrivial equation to derive $F(r)$. \square

4 Example: BTZ black hole

In this section, we provide an explicit example to show that our reconstruction formulas do work properly. The example is the static BTZ black hole,

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2d\phi^2, \quad f(r) = \frac{r^2 - r_h^2}{L^2}. \quad (4.1)$$

We employ the BTZ black hole because the analytic form of the Hartman-Maldacena entanglement entropy is known [56]. It is given by the volume of the codimension-two extremal surface linking the left and the right boundaries, which is just the geodesic length,

$$S_{\text{HM}} = \frac{A}{4G_{\text{N}}}, \quad A = 2L \left[\ln \frac{2r_{\text{max}}}{r_h} + \ln \cosh \frac{r_h t_{\text{R}}}{L^2} \right], \quad (4.2)$$

where r_{max} is the UV cutoff. Therefore, we have

$$\frac{dA}{dt_{\text{R}}} = 2 \frac{r_h}{L} \tanh \frac{r_h t_{\text{R}}}{L^2}. \quad (4.3)$$

Although we do not know the exact form of the complexity, it is obvious that the formulas we have obtained in Sec. 2 for $g(r) = 1/f(r)$ can be generalized also to the Hartman-Maldacena entanglement entropy, we can use this (4.3) as a test case to verify our reconstruction method.

Pretending that we do not know $f(r)$ for the interior of the black hole, we follow our reconstruction formula and obtain the metric of the form (2.2) inside the horizon. We assume that we know (4.3) and also $f(r)$ outside the horizon. We compare our obtained metric with the original analytic form (4.1). If the original metric (4.1) is well reconstructed, we can say that our formulas work well.

For simplicity, we set $r_h = L = 1$ in what follows. Also, since $\tilde{F} = f$ for $d = 2$, we simply write as f instead of \tilde{F} . First, we solve¹²

$$\frac{dS_{\text{HM}}}{dt_{\text{R}}} = \frac{1}{2G_{\text{N}}} \sqrt{-f_{\text{min}}}, \quad (4.4)$$

¹²For $d = 2$, $\Omega_{k,d-2}$ is unity.

to find t_R as a function of f_{\min} . The result is

$$t_R = \tilde{t}_R(f_{\min}) \equiv \text{Arctanh} \left(\sqrt{-f_{\min}} \right). \quad (4.5)$$

Next, using the metric $f(r) = r^2 - 1$ outside the horizon and $\tilde{G} = g = 1/f$, we rewrite (3.25) as

$$\begin{aligned} \tilde{Q}(f_{\min}) &= \frac{\text{Arctanh}(\sqrt{-f_{\min}})}{\sqrt{-f_{\min}}} + \int_{\delta}^{\infty} df \frac{1}{f \sqrt{(f+1)(f-f_{\min})}} \\ &= \frac{\text{Arctanh}(\sqrt{-f_{\min}})}{\sqrt{-f_{\min}}} + \int_{\xi}^{\pi/2} d\theta \frac{1}{\sin \theta \sqrt{\tan^2 \theta - f_{\min}}}, \end{aligned} \quad (4.6)$$

where we have set $f = \tan^2 \theta$ and have defined $\xi \equiv \text{Arctan} \sqrt{\delta}$ in the last line. We numerically calculate (4.6) for a various choice of the infinitesimal parameter, $\delta = 10^{-2}, 10^{-3}, 10^{-4}$. From (3.27) and (3.30), we understand that the equation we have to solve is

$$\frac{dr}{df} \frac{1}{f} = \frac{1}{\pi} \frac{d}{df} \int_f^{-\delta} du \frac{\tilde{Q}(u)}{\sqrt{u-f}}, \quad (4.7)$$

which we rewrite as

$$dr = df \frac{1}{\pi} f \frac{d}{df} \int_f^{-\delta} du \frac{\tilde{Q}(u)}{\sqrt{u-f}}. \quad (4.8)$$

Near the horizon, δ is the small cutoff for the value of f . Correspondingly, the horizon cutoff for r is found as $r_{\delta} \equiv \sqrt{1-\delta}$ since we know the analytic structure of the metric near the horizon. Using this as a upper limit of the integral region, we integrate (4.8),

$$\int_r^{\sqrt{1-\delta}} dr = \int_f^{-\delta} df' \frac{1}{\pi} f' \frac{d}{df'} \int_{f'}^{-\delta} du \frac{\tilde{Q}(u)}{\sqrt{u-f'}}. \quad (4.9)$$

A partial integration yields

$$r = \sqrt{1-\delta} + \frac{1}{\pi} \left(f \int_f^{-\delta} du \frac{\tilde{Q}(u)}{\sqrt{u-f}} + \int_f^{-\delta} df' \int_{f'}^{-\delta} du \frac{\tilde{Q}(u)}{\sqrt{u-f'}} \right). \quad (4.10)$$

Numerically calculating the right-hand side of this equation, we finally obtain $r = r(f)$ inside the horizon. See Fig.2 for the plots of our numerical results. We can see that our reconstruction formulas neatly reproduce the interior of the BTZ black hole. The δ dependence is as expected — the reconstruction works more accurately for smaller δ . This BTZ case is special in that the final RT surface at $t_R \rightarrow \infty$ hits the singularity, thus we are eventually able to reconstruct the whole interior of the BTZ black hole.




Figure 2. The numerical results of (4.10) for $\delta = 10^{-2}, 10^{-3}, 10^{-4}$ (dotted lines, from top to bottom for each value of δ respectively), and the expected analytic form $r = \sqrt{f+1}$ (solid line). The metric of the interior of the BTZ black hole is properly reconstructed. We can see the reconstruction works more accurately for smaller δ .

5 Summary and discussions

In this paper, we have obtained formulas to reconstruct a metric inside a black hole horizon by the time derivative of the complexity measured in the dual QFT, based on the “complexity = volume” (CV) conjecture in the holographic duality. We have assumed only that the metrics outside the horizon is known.

The bulk metric in the stationary spacetimes takes the form $ds^2 = -f(r)dt^2 + g(r)dr^2 + r^2d\Omega^2$. If $g(r)$ is known as a function of $f(r)$ (such as the case $g(r) = 1/f(r)$), we need only the time dependence of the complexity in the dual QFT as the data to reconstruct the metric of the black hole interior, as we have seen in Sec. 2.2 and in App. A. On the other hand, in the general case when both $f(r)$ and $g(r)$ are not known, we need, in addition to the complexity, the Hartman-Maldacena entanglement entropy of the doubled QFT [56] to reconstruct the interior metric, as we have seen in Sec. 3.2.

For our reconstruction formula, the metric outside the horizon needs to be known. We note here that other QFT quantities can determine it through the holographic duality. The entanglement entropy can determine only $g(r)$ [14, 15] because the RT surface is on a time slice. The Wilson loop [35] can determine both $f(r)$ and $g(r)$ but they are in the string frame, and they differ from the Einstein frame by a conformal factor of unknown dilaton profile. The method using the light-cone cuts [8, 9, 12] determines the metric up to a conformal factor. Therefore, the combination of the entanglement entropy and either the Wilson loops or the light-cone cuts can determine the metric outside the black hole horizon completely.

Another assumption for the reconstruction of the metric is that the time derivative

of the complexity needs to be monotonic. The charged and uncharged AdS black holes in various dimensions satisfy this criterion [65], while some field-theoretic definitions of complexity may not satisfy it [66]. There exist several different proposals on the definition of the complexity in QFTs [58, 67–73], and the time dependence can vary depending on the definitions (see [65, 66, 74–78] for the study of the time dependence). We plan to come back to the question of what our formula can say for a holographically consistent definition of the QFT complexity.

How far can we go with this reconstruction method to reveal the mechanism of the AdS/CFT correspondence? Since the gravity side of the AdS/CFT are described mainly by geometric quantities, once the inverse problems to obtain spacetimes by the probes can be solved analytically as demonstrated in this paper, enough information of the dual QFT data will be able to reconstruct completely the bulk metrics and the bulk theory. Using the complexity would be beneficial since it can go even inside of the horizons, and our spirit is close to the derivation of the Einstein equations by varying the complexity [79]. The holographic complexity is a version of surfaces in quantum gravity spacetimes, and string theory has been developed based on probes of surfaces, which started with the brane democracy [80]. Needless to mention a similarity between the complexity surface and the spacelike branes [81, 82], the various surfaces in the bulk may keep playing a central role in the study of the AdS/CFT correspondence.

Acknowledgments

We are grateful to K. Y. Kim for discussions. K. H. was supported in part by JSPS KAKENHI Grant No. JP17H06462.

A Treatment of the special case $g = g(f)$

As we have seen in Sec. 2, if we assume $g = 1/f$, we can reconstruct the metric inside the horizon by the time dependence of the complexity once the metric outside the horizon is known. More generally, if g is known as a function of f , we can reconstruct the geometry by only the complexity. Here we present our formula for that case.

Assume that

- the time derivative of the complexity $\dot{\mathcal{C}}(t)$ in the dual QFT is given.
- $\dot{\mathcal{C}}(t)$ is monotonic in time.
- the metric (3.1) outside the horizon ($r \geq r_h$) is known.

Then, solve

$$\frac{d\mathcal{C}}{dt_R}(t_R) = \frac{2\Omega_{k,d-1}}{G_{\text{NL}}} \sqrt{-F_{\text{min}}} \quad (\text{A.1})$$

to get t_R as a function of F_{min} . In addition, outside the horizon, invert $F = F(r) \equiv f(r)r^{2(d-1)}$ to find $r = r(F)$. Substitute them into

$$Q(F_{\text{min}}) \equiv \frac{t_R(F_{\text{min}})}{\sqrt{-F_{\text{min}}}} + \int_{\epsilon}^{\infty} dF \frac{dr}{dF} \frac{\sqrt{FG(F, r(F))}}{F} \frac{1}{\sqrt{F - F_{\text{min}}}}, \quad (\text{A.2})$$

with a positive infinitesimal parameter ϵ . $G(F, r(F))$ is defined as

$$G(F, r(F)) \equiv g(Fr(F)^{-2(d-1)})r(F)^{2(d-1)}. \quad (\text{A.3})$$

For $F \leq -\epsilon$, calculate

$$y(F) \equiv \frac{1}{\pi} \frac{d}{dF} \int_F^{-\epsilon} dU \frac{Q(U)}{\sqrt{U - F}}, \quad (\text{A.4})$$

and integrate the differential equation

$$\frac{dr}{dF} \frac{\sqrt{FG(F, r(F))}}{F} = y(F) \quad (\text{A.5})$$

to find $r = r(F)$. Finally, invert it to find $F(r)$, and the metric inside the black hole is given by $f(r) = F(r)r^{-2(d-1)}$.

The proof is almost identical to that of Sec. 2.2 since (3.13) and (3.16) have the same form as (2.4) and (2.5), respectively.

References

- [1] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” *Int. J. Theor. Phys.* **38** (1999), 1113-1133 doi:10.1023/A:1026654312961 [arXiv:hep-th/9711200 [hep-th]].
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from noncritical string theory,” *Phys. Lett. B* **428** (1998), 105-114 doi:10.1016/S0370-2693(98)00377-3 [arXiv:hep-th/9802109 [hep-th]].
- [3] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2** (1998), 253-291 doi:10.4310/ATMP.1998.v2.n2.a2 [arXiv:hep-th/9802150 [hep-th]].
- [4] X. Dong, D. Harlow and A. C. Wall, “Reconstruction of Bulk Operators within the Entanglement Wedge in Gauge-Gravity Duality,” *Phys. Rev. Lett.* **117**, no.2, 021601 (2016) doi:10.1103/PhysRevLett.117.021601 [arXiv:1601.05416 [hep-th]].
- [5] S. de Haro, S. N. Solodukhin and K. Skenderis, “Holographic reconstruction of space-time and renormalization in the AdS / CFT correspondence,” *Commun. Math. Phys.* **217** (2001), 595-622 doi:10.1007/s002200100381 [arXiv:hep-th/0002230 [hep-th]].
- [6] J. Hammersley, “Extracting the bulk metric from boundary information in asymptotically AdS spacetimes,” *JHEP* **12** (2006), 047 doi:10.1088/1126-6708/2006/12/047 [arXiv:hep-th/0609202 [hep-th]].
- [7] V. E. Hubeny, H. Liu and M. Rangamani, “Bulk-cone singularities & signatures of horizon formation in AdS/CFT,” *JHEP* **01** (2007), 009 doi:10.1088/1126-6708/2007/01/009 [arXiv:hep-th/0610041 [hep-th]].
- [8] N. Engelhardt and G. T. Horowitz, “Towards a Reconstruction of General Bulk Metrics,” *Class. Quant. Grav.* **34** (2017) no.1, 015004 doi:10.1088/1361-6382/34/1/015004 [arXiv:1605.01070 [hep-th]].
- [9] N. Engelhardt and G. T. Horowitz, “Recovering the spacetime metric from a holographic dual,” *Adv. Theor. Math. Phys.* **21** (2017), 1635-1653 doi:10.4310/ATMP.2017.v21.n7.a2 [arXiv:1612.00391 [hep-th]].
- [10] S. R. Roy and D. Sarkar, “Bulk metric reconstruction from boundary entanglement,” *Phys. Rev. D* **98** (2018) no.6, 066017 doi:10.1103/PhysRevD.98.066017 [arXiv:1801.07280 [hep-th]].
- [11] P. Burda, R. Gregory and A. Jain, “Holographic reconstruction of bubble spacetimes,” *Phys. Rev. D* **99** (2019) no.2, 026003 doi:10.1103/PhysRevD.99.026003 [arXiv:1804.05202 [hep-th]].
- [12] S. Hernández-Cuenca and G. T. Horowitz, “Bulk reconstruction of metrics with a compact space asymptotically,” [arXiv:2003.08409 [hep-th]].
- [13] J. Hammersley, “Numerical metric extraction in AdS/CFT,” *Gen. Rel. Grav.* **40** (2008), 1619-1652 doi:10.1007/s10714-007-0564-6 [arXiv:0705.0159 [hep-th]].
- [14] S. Bilson, “Extracting spacetimes using the AdS/CFT conjecture,” *JHEP* **08** (2008), 073 doi:10.1088/1126-6708/2008/08/073 [arXiv:0807.3695 [hep-th]].
- [15] S. Bilson, “Extracting Spacetimes using the AdS/CFT Conjecture: Part II,” *JHEP* **02** (2011), 050 doi:10.1007/JHEP02(2011)050 [arXiv:1012.1812 [hep-th]].
- [16] V. E. Hubeny, “Extremal surfaces as bulk probes in AdS/CFT,” *JHEP* **07** (2012), 093 doi:10.1007/JHEP07(2012)093 [arXiv:1203.1044 [hep-th]].

- [17] V. Balasubramanian, B. Czech, B. D. Chowdhury and J. de Boer, “The entropy of a hole in spacetime,” *JHEP* **10** (2013), 220 doi:10.1007/JHEP10(2013)220 [arXiv:1305.0856 [hep-th]].
- [18] V. Balasubramanian, B. D. Chowdhury, B. Czech, J. de Boer and M. P. Heller, “Bulk curves from boundary data in holography,” *Phys. Rev. D* **89** (2014) no.8, 086004 doi:10.1103/PhysRevD.89.086004 [arXiv:1310.4204 [hep-th]].
- [19] R. C. Myers, J. Rao and S. Sugishita, “Holographic Holes in Higher Dimensions,” *JHEP* **06** (2014), 044 doi:10.1007/JHEP06(2014)044 [arXiv:1403.3416 [hep-th]].
- [20] B. Czech, X. Dong and J. Sully, “Holographic Reconstruction of General Bulk Surfaces,” *JHEP* **11** (2014), 015 doi:10.1007/JHEP11(2014)015 [arXiv:1406.4889 [hep-th]].
- [21] B. Czech and L. Lamprou, “Holographic definition of points and distances,” *Phys. Rev. D* **90** (2014), 106005 doi:10.1103/PhysRevD.90.106005 [arXiv:1409.4473 [hep-th]].
- [22] S. A. Gentle and C. Keeler, “On the reconstruction of Lifshitz spacetimes,” *JHEP* **03** (2016), 195 doi:10.1007/JHEP03(2016)195 [arXiv:1512.04538 [hep-th]].
- [23] A. Saha, S. Karar and S. Gangopadhyay, “Bulk geometry from entanglement entropy of CFT,” *Eur. Phys. J. Plus* **135** (2020) no.2, 132 doi:10.1140/epjp/s13360-020-00110-7 [arXiv:1807.04646 [hep-th]].
- [24] N. Bao, C. Cao, S. Fischetti and C. Keeler, “Towards Bulk Metric Reconstruction from Extremal Area Variations,” *Class. Quant. Grav.* **36** (2019) no.18, 185002 doi:10.1088/1361-6382/ab377f [arXiv:1904.04834 [hep-th]].
- [25] C. Cao, X. L. Qi, B. Swingle and E. Tang, “Building Bulk Geometry from the Tensor Radon Transform,” [arXiv:2007.00004 [hep-th]].
- [26] N. Jokela and A. Pönni, “Towards precision holography,” [arXiv:2007.00010 [hep-th]].
- [27] S. Ryu and T. Takayanagi, “Holographic derivation of entanglement entropy from AdS/CFT,” *Phys. Rev. Lett.* **96** (2006), 181602 doi:10.1103/PhysRevLett.96.181602 [arXiv:hep-th/0603001 [hep-th]].
- [28] S. Ryu and T. Takayanagi, “Aspects of Holographic Entanglement Entropy,” *JHEP* **08** (2006), 045 doi:10.1088/1126-6708/2006/08/045 [arXiv:hep-th/0605073 [hep-th]].
- [29] B. Swingle, “Constructing holographic spacetimes using entanglement renormalization,” [arXiv:1209.3304 [hep-th]].
- [30] N. Bao, G. Penington, J. Sorce and A. C. Wall, “Beyond Toy Models: Distilling Tensor Networks in Full AdS/CFT,” *JHEP* **19** (2020), 069 doi:10.1007/JHEP11(2019)069 [arXiv:1812.01171 [hep-th]].
- [31] A. Milsted and G. Vidal, “Geometric interpretation of the multi-scale entanglement renormalization ansatz,” [arXiv:1812.00529 [hep-th]].
- [32] N. Bao, G. Penington, J. Sorce and A. C. Wall, “Holographic Tensor Networks in Full AdS/CFT,” [arXiv:1902.10157 [hep-th]].
- [33] B. Swingle, “Entanglement Renormalization and Holography,” *Phys. Rev. D* **86** (2012), 065007 doi:10.1103/PhysRevD.86.065007 [arXiv:0905.1317 [cond-mat.str-el]].
- [34] F. Pastawski, B. Yoshida, D. Harlow and J. Preskill, “Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence,” *JHEP* **06** (2015), 149 doi:10.1007/JHEP06(2015)149 [arXiv:1503.06237 [hep-th]].

- [35] K. Hashimoto, “Building bulk from Wilson loops,” *PTEP* **2021** (2021), 023B04 doi:10.1093/ptep/ptaa183 [arXiv:2008.10883 [hep-th]].
- [36] J. M. Maldacena, “Wilson loops in large N field theories,” *Phys. Rev. Lett.* **80** (1998), 4859-4862 doi:10.1103/PhysRevLett.80.4859 [arXiv:hep-th/9803002 [hep-th]].
- [37] S. J. Rey and J. T. Yee, “Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity,” *Eur. Phys. J. C* **22** (2001), 379-394 doi:10.1007/s100520100799 [arXiv:hep-th/9803001 [hep-th]].
- [38] A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, “Wilson loops in the large N limit at finite temperature,” *Phys. Lett. B* **434** (1998), 36-40 doi:10.1016/S0370-2693(98)00730-8 [arXiv:hep-th/9803137 [hep-th]].
- [39] S. J. Rey, S. Theisen and J. T. Yee, “Wilson-Polyakov loop at finite temperature in large N gauge theory and anti-de Sitter supergravity,” *Nucl. Phys. B* **527** (1998), 171-186 doi:10.1016/S0550-3213(98)00471-4 [arXiv:hep-th/9803135 [hep-th]].
- [40] K. Hashimoto, S. Sugishita, A. Tanaka and A. Tomiya, “Deep learning and the AdS/CFT correspondence,” *Phys. Rev. D* **98** (2018) no.4, 046019 doi:10.1103/PhysRevD.98.046019 [arXiv:1802.08313 [hep-th]].
- [41] K. Hashimoto, S. Sugishita, A. Tanaka and A. Tomiya, “Deep Learning and Holographic QCD,” *Phys. Rev. D* **98** (2018) no.10, 106014 doi:10.1103/PhysRevD.98.106014 [arXiv:1809.10536 [hep-th]].
- [42] J. Tan and C. B. Chen, “Deep learning the holographic black hole with charge,” *Int. J. Mod. Phys. D* **28** (2019) no.12, 1950153 doi:10.1142/S0218271819501530 [arXiv:1908.01470 [hep-th]].
- [43] Y. K. Yan, S. F. Wu, X. H. Ge and Y. Tian, “Deep learning black hole metrics from shear viscosity,” *Phys. Rev. D* **102**, no.10, 101902 doi:10.1103/PhysRevD.102.101902 [arXiv:2004.12112 [hep-th]].
- [44] T. Akutagawa, K. Hashimoto and T. Sumimoto, “Deep Learning and AdS/QCD,” *Phys. Rev. D* **102** (2020) no.2, 026020 doi:10.1103/PhysRevD.102.026020 [arXiv:2005.02636 [hep-th]].
- [45] K. Hashimoto, H. Y. Hu and Y. Z. You, “Neural ODE and Holographic QCD,” to be published in *Machine Learning: Science and Technology* [arXiv:2006.00712 [hep-th]].
- [46] Y. Z. You, Z. Yang and X. L. Qi, “Machine Learning Spatial Geometry from Entanglement Features,” *Phys. Rev. B* **97** (2018) no.4, 045153 doi:10.1103/PhysRevB.97.045153 [arXiv:1709.01223 [cond-mat.dis-nn]].
- [47] H. Y. Hu, S. H. Li, L. Wang and Y. Z. You, “Machine Learning Holographic Mapping by Neural Network Renormalization Group,” *Phys. Rev. Res.* **2** (2020) no.2, 023369 doi:10.1103/PhysRevResearch.2.023369 [arXiv:1903.00804 [cond-mat.dis-nn]].
- [48] K. Hashimoto, “AdS/CFT correspondence as a deep Boltzmann machine,” *Phys. Rev. D* **99** (2019) no.10, 106017 doi:10.1103/PhysRevD.99.106017 [arXiv:1903.04951 [hep-th]].
- [49] M. Song, M. S. H. Oh, Y. Ahn and K. Y. Kim, “AdS/Deep-Learning made easy: simple examples,” [arXiv:2011.13726 [physics.class-ph]].
- [50] F. Ruehle, “Data science applications to string theory,” *Phys. Rept.* **839** (2020), 1-117 doi:10.1016/j.physrep.2019.09.005

- [51] G. Carleo, I. Cirac, K. Cranmer, L. Daudet, M. Schuld, N. Tishby, L. Vogt-Maranto and L. Zdeborová, “Machine learning and the physical sciences,” *Rev. Mod. Phys.* **91** (2019) no.4, 045002 doi:10.1103/RevModPhys.91.045002 [arXiv:1903.10563 [physics.comp-ph]].
- [52] L. Susskind, “Computational Complexity and Black Hole Horizons,” *Fortsch. Phys.* **64**, 24 (2016) doi:10.1002/prop.201500092 [arXiv:1403.5695 [hep-th], arXiv:1402.5674 [hep-th]].
- [53] L. Susskind, “Entanglement is not enough,” *Fortsch. Phys.* **64**, 49 (2016) doi:10.1002/prop.201500095 [arXiv:1411.0690 [hep-th]].
- [54] L. Susskind, “Addendum to computational complexity and black hole horizons,” *Fortsch. Phys.* **64**, 44 (2016). doi:10.1002/prop.201500093
- [55] J. M. Maldacena, “Eternal black holes in anti-de Sitter,” *JHEP* **0304**, 021 (2003) doi:10.1088/1126-6708/2003/04/021 [hep-th/0106112].
- [56] T. Hartman and J. Maldacena, “Time Evolution of Entanglement Entropy from Black Hole Interiors,” *JHEP* **1305**, 014 (2013) doi:10.1007/JHEP05(2013)014 [arXiv:1303.1080 [hep-th]].
- [57] D. Stanford and L. Susskind, “Complexity and Shock Wave Geometries,” *Phys. Rev. D* **90**, no. 12, 126007 (2014) doi:10.1103/PhysRevD.90.126007 [arXiv:1406.2678 [hep-th]].
- [58] L. Susskind and Y. Zhao, “Switchbacks and the Bridge to Nowhere,” arXiv:1408.2823 [hep-th].
- [59] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle and Y. Zhao, “Holographic Complexity Equals Bulk Action?,” *Phys. Rev. Lett.* **116**, no. 19, 191301 (2016) doi:10.1103/PhysRevLett.116.191301 [arXiv:1509.07876 [hep-th]].
- [60] A. R. Brown, D. A. Roberts, L. Susskind, B. Swingle and Y. Zhao, “Complexity, action, and black holes,” *Phys. Rev. D* **93**, no. 8, 086006 (2016) doi:10.1103/PhysRevD.93.086006 [arXiv:1512.04993 [hep-th]].
- [61] V. Balasubramanian, A. Bernamonti, J. de Boer, N. Copland, B. Craps, E. Keski-Vakkuri, B. Muller, A. Schafer, M. Shigemori and W. Staessens, “Thermalization of Strongly Coupled Field Theories,” *Phys. Rev. Lett.* **106** (2011), 191601 doi:10.1103/PhysRevLett.106.191601 [arXiv:1012.4753 [hep-th]].
- [62] V. Balasubramanian, A. Bernamonti, J. de Boer, N. Copland, B. Craps, E. Keski-Vakkuri, B. Muller, A. Schafer, M. Shigemori and W. Staessens, “Holographic Thermalization,” *Phys. Rev. D* **84** (2011), 026010 doi:10.1103/PhysRevD.84.026010 [arXiv:1103.2683 [hep-th]].
- [63] U. Kol, C. Nunez, D. Schofield, J. Sonnenschein and M. Warschawski, “Confinement, Phase Transitions and non-Locality in the Entanglement Entropy,” *JHEP* **06** (2014), 005 doi:10.1007/JHEP06(2014)005 [arXiv:1403.2721 [hep-th]].
- [64] N. Bao, A. Chatwin-Davies, B. E. Niehoff and M. Usatyuk, “Bulk Reconstruction Beyond the Entanglement Wedge,” *Phys. Rev. D* **101** (2020) no.6, 066011 doi:10.1103/PhysRevD.101.066011 [arXiv:1911.00519 [hep-th]].
- [65] D. Carmi, S. Chapman, H. Marrochio, R. C. Myers and S. Sugishita, “On the Time Dependence of Holographic Complexity,” *JHEP* **11**, 188 (2017) doi:10.1007/JHEP11(2017)188 [arXiv:1709.10184 [hep-th]].
- [66] R. Q. Yang, C. Niu, C. Y. Zhang and K. Y. Kim, “Comparison of holographic and field theoretic complexities for time dependent thermofield double states,” *JHEP* **02**, 082 (2018) doi:10.1007/JHEP02(2018)082 [arXiv:1710.00600 [hep-th]].

- [67] S. Chapman, M. P. Heller, H. Marrochio and F. Pastawski, “Toward a Definition of Complexity for Quantum Field Theory States,” *Phys. Rev. Lett.* **120**, no.12, 121602 (2018) doi:10.1103/PhysRevLett.120.121602 [arXiv:1707.08582 [hep-th]].
- [68] P. Caputa, N. Kundu, M. Miyaji, T. Takayanagi and K. Watanabe, “Anti-de Sitter Space from Optimization of Path Integrals in Conformal Field Theories,” *Phys. Rev. Lett.* **119**, no.7, 071602 (2017) doi:10.1103/PhysRevLett.119.071602 [arXiv:1703.00456 [hep-th]].
- [69] K. Hashimoto, N. Iizuka and S. Sugishita, “Time evolution of complexity in Abelian gauge theories,” *Phys. Rev. D* **96**, no.12, 126001 (2017) doi:10.1103/PhysRevD.96.126001 [arXiv:1707.03840 [hep-th]].
- [70] R. Q. Yang, “Complexity for quantum field theory states and applications to thermofield double states,” *Phys. Rev. D* **97**, no.6, 066004 (2018) doi:10.1103/PhysRevD.97.066004 [arXiv:1709.00921 [hep-th]].
- [71] R. Q. Yang, Y. S. An, C. Niu, C. Y. Zhang and K. Y. Kim, “Principles and symmetries of complexity in quantum field theory,” *Eur. Phys. J. C* **79**, no.2, 109 (2019) doi:10.1140/epjc/s10052-019-6600-3 [arXiv:1803.01797 [hep-th]].
- [72] H. A. Camargo, M. P. Heller, R. Jefferson and J. Knaute, “Path integral optimization as circuit complexity,” *Phys. Rev. Lett.* **123**, no.1, 011601 (2019) doi:10.1103/PhysRevLett.123.011601 [arXiv:1904.02713 [hep-th]].
- [73] R. Q. Yang, Y. S. An, C. Niu, C. Y. Zhang and K. Y. Kim, “To be unitary-invariant or not?: a simple but non-trivial proposal for the complexity between states in quantum mechanics/field theory,” [arXiv:1906.02063 [hep-th]].
- [74] R. Q. Yang, *Phys. Rev. D* **95**, no.8, 086017 (2017) doi:10.1103/PhysRevD.95.086017 [arXiv:1610.05090 [gr-qc]].
- [75] A. R. Brown and L. Susskind, *Phys. Rev. D* **97**, no.8, 086015 (2018) doi:10.1103/PhysRevD.97.086015 [arXiv:1701.01107 [hep-th]].
- [76] W. Cottrell and M. Montero, *JHEP* **02**, 039 (2018) doi:10.1007/JHEP02(2018)039 [arXiv:1710.01175 [hep-th]].
- [77] K. Hashimoto, N. Iizuka and S. Sugishita, “Thoughts on Holographic Complexity and its Basis-dependence,” *Phys. Rev. D* **98**, no.4, 046002 (2018) doi:10.1103/PhysRevD.98.046002 [arXiv:1805.04226 [hep-th]].
- [78] T. Ali, A. Bhattacharyya, S. Shajidul Haque, E. H. Kim and N. Moynihan, “Time Evolution of Complexity: A Critique of Three Methods,” *JHEP* **04**, 087 (2019) doi:10.1007/JHEP04(2019)087 [arXiv:1810.02734 [hep-th]].
- [79] B. Czech, “Einstein Equations from Varying Complexity,” *Phys. Rev. Lett.* **120**, no.3, 031601 (2018) doi:10.1103/PhysRevLett.120.031601 [arXiv:1706.00965 [hep-th]].
- [80] P. K. Townsend, “P-brane democracy,” [arXiv:hep-th/9507048 [hep-th]].
- [81] M. Gutperle and A. Strominger, “Space - like branes,” *JHEP* **04**, 018 (2002) doi:10.1088/1126-6708/2002/04/018 [arXiv:hep-th/0202210 [hep-th]].
- [82] K. Hashimoto, P. M. Ho and J. E. Wang, “S-brane actions,” *Phys. Rev. Lett.* **90**, 141601 (2003) doi:10.1103/PhysRevLett.90.141601 [arXiv:hep-th/0211090 [hep-th]].