

Meta-Analysis of Bayes Factors ^{*}

Stavros Nikolakopoulos^{†1,2} and Ioannis Ntzoufras¹

¹Department of Statistics, Athens University of Economics and Business

²Department of Biostatistics, University Medical Center Utrecht

Abstract

Bayes Factors, the Bayesian tool for hypothesis testing, are receiving increasing attention in the literature. Compared to their frequentist rivals (p -values or test statistics), Bayes Factors have the conceptual advantage of providing evidence both for and against a null hypothesis and they can be calibrated so that they do not depend so heavily on the sample size. However, research on the synthesis of Bayes Factors arising from individual studies has received very limited attention. In this work we review and propose methods for combining Bayes Factors from multiple studies, depending on the level of information available. In the process, we provide insights with respect to the interplay between frequentist and Bayesian evidence. We also clarify why some intuitive suggestions in the literature can be misleading. We assess the performance of the methods discussed via a simulation study and apply the methods in an example from the field of psychology.

^{*}*Acknowledgements:* This work was supported by the DRASI-2 funding scheme, Athens University of Economics and Business

[†]corresponding author; e:sknikolak@aueb.gr

Meta-Analysis of Bayes Factors

1 Introduction

In evidence-based empirical sciences, research results arise from a variety of sources and accumulate in an all-increasing rate. In the context of synthesizing aggregate data, the field of meta-analysis has received considerable attention from the statistical community, mostly within the classical (frequentist) framework of statistical inference.

In classical statistics, hypothesis-testing-related inference is typically based on p -values, and on point or interval estimates, when the focus is parameter estimation. If K (independent) estimates $\hat{\theta} = \hat{\theta}_1, \dots, \hat{\theta}_K$ are available for the same parameter θ , then the meta-analytic estimate of θ is of the general form $\hat{\theta}_{MA} = f(\hat{\theta})$. If instead, interest lies in testing a null hypothesis for θ by synthesizing the evidence from the K datasets, then one should synthesize the test statistics $\mathbf{T} = \mathcal{T}_1, \dots, \mathcal{T}_K$ or the corresponding p -values $\mathbf{p} = \{p_1, \dots, p_K\}$. Hence, hypothesis-testing based meta-analysis can provide an overall p -value $\mathcal{P}_{MA} = f_p(\mathbf{p})$ or meta-analytic test statistic $\mathcal{T}_{MA} = f_t(\mathbf{T})$, taking into account the evidence against H_0 arising from each individual study.

The published research focusing on frequentist meta-analysis is vast and an exhaustive review is beyond the scope of this paper. Methods for combining p -values date back to Pearson (1933) and Fisher (1934). For a recent overview and methodological developments see Heard and Rubin-Delanchy (2018). For a more general overview of meta-analysis methods, see Hedges and Olkin (2014).

The shortcomings of p -values as a measure of evidence have long been the subject of a heated discussion among statisticians. Their inability to quantify evidence in favour of a null hypothesis as well as their overstatement of evidence against it have been overly stressed in the statistical literature (Berger and Sellke 1987; Sellke et al. 2001; Held and Ott 2018; Benjamin et al. 2018). Therefore, combining p -values for quantifying overall evidence inevitably suffers from similar shortcomings.

Within the Bayesian framework, each hypothesis test is viewed as a pairwise model comparison problem. The most natural tool for comparing two hypotheses or models in Bayesian theory are the posterior model odds (PO) or the corresponding Bayes Factors (BF) (Jeffreys 1961; Kass and Raftery 1995). From a scientific point of view, PO s and BF s have the significant conceptual advantage of providing evidence both for and against a null hypothesis. This feature can be of paramount importance since null hypotheses often correspond to theoretically useful statements of invariance and constraint (Kass 1993).

Synthesis of BF s arising from individual datasets is, therefore, of direct concern in the meta-analysis of published research. Even though the appearance of BF s as the sole form of evidence quantification is still rather rare, transformation of frequentist evidence to a corresponding Bayesian measure is a challenging task which is gaining increasing attention. Examples include the Bayesian assessment for the strength of evidence suggested by the FDA for approved antidepressants (Monden et al. 2016), as well as re-examining an unexpected claim for increased mortality with paclitaxel-eluting balloons and stents to treat femoropopliteal arterial disease (Bittl et al. 2019).

Nevertheless, research on the methods for synthesizing BF s has been very limited. Maybe this can be partly attributed to the fallacious intuition that the BF resulting from a series of BF s comparing the same two models is merely the product of those BF s, by sequential updating of the BF s (Goodman 2016). This intuition that the meta-analytic BF is just the product of individual BF s is fundamentally false (Rouder and Morey 2011).

In this paper we present and evaluate methods for calculating the overall BF from a set of available BF s that compare the same two models. We focus on the simple linear regression model with either a continuous or a binary covariate, the latter representing the very common case in meta analysis of synthesizing the effect estimates from randomized studies with a continuous outcome (t -tests). We make use of results that provide an analytic expression connecting BF with the frequentist test statistics in combination with meta-analytic methodology. Since a meta-analysis can include varying levels of detail concerning the effect of variables within each study, we present appropriate methods for several scenarios. Thus, the aims of the methodology we review and propose are the following:

- The translation of classical (frequentist) evidence to Bayesian (BF)

- The synthesis of (frequentist or Bayesian) evidence in terms of BF s
- Meta-analytic BF s for varying levels of information availability

We demonstrate the methods through an example from the field of public health. The paper concludes with a discussion.

2 Preliminaries

2.1 The Bayes Factor

For a given set of response values \mathbf{y} and two competing models $M_0 = \{f(\cdot|\theta_0, M_0), \theta_0 \in \Theta_0\}$ and $M_1 = \{f(\cdot|\theta_1, M_1), \theta_1 \in \Theta_1\}$ the Bayes Factor B_{10} is the ratio of the marginal densities of \mathbf{y} under M_1 and M_0 given by

$$BF_{10} = \frac{f(\mathbf{y}|M_1)}{f(\mathbf{y}|M_0)} = \frac{\int_{\Theta_1} f(\mathbf{y}|\theta_1, M_1)\pi(\theta_1|M_1)d\theta_1}{\int_{\Theta_0} f(\mathbf{y}|\theta_0, M_0)\pi(\theta_0|M_0)d\theta_0}$$

where $\pi(\theta_\ell|M_\ell)$ represents the prior distribution of θ_ℓ under model M_ℓ , for $\ell \in \{0, 1\}$. Note also that

$$PO_{10} = \frac{Pr(M_1|\mathbf{y})}{Pr(M_0|\mathbf{y})} = BF_{10} \times \frac{Pr(M_1)}{Pr(M_0)}$$

so the PO_{10} quantifies the increase in the odds of M_1 relative to M_0 after the data is observed.

The Bayes factor is a special case of posterior model odds when considering equal prior probabilities for each hypothesis or model under consideration. In this article we focus on the synthesis of BF s rather than PO s for several reasons. First of all, the choice of equal probabilities for the two competing models/hypothesis is considered as a reasonable non-informative choice. Moreover, when synthesizing Bayesian evidence, it is sensible to leave outside of the meta-analysis procedure any study specific prior preferences. Thus, we estimate a combined BF across all studies and then incorporate any overall prior model preferences at the end of the meta analytic procedure.

Since the BF does not depend on the prior probabilities of the models, it is usually interpreted as the odds provided by the data for M_1 to M_0 . However, BF_{10} obviously depends on the prior distributions of the model parameters. Large values of $BF_{k\ell}$ indicate strong posterior support of model M_k against model M_ℓ ; for details see, e.g. Kass and Raftery (1995).

2.2 Bayes Factors with conjugate priors

We consider the standard general hypothesis testing setting in linear regression with normal independent errors:

$$\begin{aligned} H_0 : \boldsymbol{\beta} = 0 &\Rightarrow M_0 : \mathbf{y}|\boldsymbol{\alpha}, \tau \sim N_n(\mathbf{X}_0\boldsymbol{\alpha}, \tau^{-1}\mathbf{I}_n) \\ H_1 : \boldsymbol{\beta} \neq 0 &\Rightarrow M_1 : \mathbf{y}|\boldsymbol{\alpha}, \boldsymbol{\beta}, \tau \sim N_n(\mathbf{X}_0\boldsymbol{\alpha} + \mathbf{X}_1\boldsymbol{\beta}, \tau^{-1}\mathbf{I}_n) \end{aligned} \tag{1}$$

where N_n stands for the multivariate normal distribution of dimension n , \mathbf{I}_n is an $n \times n$ identity matrix, \mathbf{X}_0 is a full-rank $n \times q$ matrix representing possible nuisance covariates (including a column of ones which corresponds to the constant parameter), \mathbf{X}_1 is an $n \times p$ design matrix including the covariates of interest additionally to the ones included in M_0 , $\boldsymbol{\alpha}$ is a q -dimensional vector of parameters of the null model, $\boldsymbol{\beta}$ is a p -dimensional vector of regression coefficients related to the covariates of \mathbf{X}_1 additionally to the ones included in the null model, and τ^{-1} is the error variance.

The null model (M_0) is nested within M_1 and represents the model with no effect of the covariates involved in \mathbf{X}_1 . Note that this is a generalization of the commonly considered as M_0 , the constant model (without any covariates) which corresponds to setting $\mathbf{X}_0 = \mathbf{1}_{n \times 1}$ in (1).

For the parameterization (1), the commonly employed conjugate normal-inverse-gamma prior (NIG , O'Hagan and Forster (2004)) is formulated as

$$(\boldsymbol{\alpha}, \boldsymbol{\beta})|\tau \sim N_{q+p}(\mathbf{0}, \mathbf{V}\tau^{-1}).$$

Assuming prior independence between α and β , then:

$$\begin{aligned}\alpha|\tau &\sim N_q(\mathbf{0}, \mathbf{V}_\alpha \tau^{-1}) \\ \beta|\tau &\sim N_p(\mathbf{0}, \mathbf{V}_\beta \tau^{-1}) \\ \tau &\sim \text{Gamma}(k_1/2, k_2/2)\end{aligned}$$

where \mathbf{V}_α and \mathbf{V}_β are some positive definite matrices, and the gamma distribution is in the shape-rate parameterization. A limiting case of the *NIG* prior is Zellner's *g*-prior (Zellner 1986), as modified and implemented by Liang et al. (2008), which results to

$$\begin{aligned}\mathbf{X}_0 &= \mathbf{1}, \quad \alpha = \alpha \\ p(\alpha, \tau) &\propto \tau^{-1} \\ \beta|\tau &\sim N_n(\mathbf{0}, g(\mathbf{X}_1^t \mathbf{X}_1)^{-1} \tau^{-1})\end{aligned}$$

by imposing $\mathbf{V}_\alpha \rightarrow 0$ and $k_1, k_2 \rightarrow 0$; where α is now a scalar. Liang et al. (2008) have shown that for the *g*-prior setup:

$$2 \log BF_{10} = (n - p - 1) \log(1 + g) - (n - 1) \log [1 + g(1 - R_1^2)] \quad (2)$$

where R_1^2 is the coefficient of determination for model M_1 . For the model (1) with q nuisance covariates, the resulting BF_{01} is given by

$$2 \log BF_{01} = -(n - p - 1) \log(1 + g) + (n - q - 1) \log \left[1 + g \frac{1 - R_1^2}{1 - R_0^2} \right] \quad (3)$$

where R_0^2 the coefficient of determination for model M_0 . Zhou and Guan (2018) further show that for the latter formulation and $p(\alpha, \tau) \propto \tau^{(q-2)/2}$, then

$$\begin{aligned}BF_{10} &= |\mathbf{V}_\beta|^{-1/2} |\mathbf{X}^t \mathbf{X} + \mathbf{V}_\beta^{-1}|^{-1/2} \\ &\times \left\{ 1 - \frac{\mathbf{y}^t \mathbf{X} (\mathbf{X}^t \mathbf{X} + \mathbf{V}_\beta^{-1})^{-1} \mathbf{X}^t \mathbf{y}}{\mathbf{y}^t \mathbf{y} - \mathbf{y}^t \mathbf{X}_0 (\mathbf{X}_0^t \mathbf{X}_0)^{-1} \mathbf{X}_0^t \mathbf{y}} \right\}^{-n/2}\end{aligned} \quad (4)$$

with

$$\mathbf{X} = \left(\mathbf{I}_n - \mathbf{X}_0 (\mathbf{X}_0^t \mathbf{X}_0)^{-1} \mathbf{X}_0^t \right) \mathbf{X}_1$$

the residuals of \mathbf{X}_1 after regressing out \mathbf{X}_0 and $|\cdot|$ denotes the determinant. It can easily be shown that for the *g*-prior setting and no nuisance covariates, (4) results to (2), with a small difference in the exponents (n instead of $n - 1$ for the *BF* scale), caused by the slightly different priors on τ . This difference is negligible for large n .

2.3 Bayes Factors with *g*-priors via test statistics

Let us assume a model M_1 with parameters $\theta = (\theta_0, \theta_1)$ and a sub-model M_0 with $\theta_1 = 0$. Then, the likelihood ratio statistic for a null hypothesis $H_0 : \theta_1 = 0, \theta_0 \in \Theta_0$ vs. the alternative $H_1 : \theta \in \Theta = \Theta_0 \times \Theta_1$ (which correspond to M_0 and M_1 as defined above) is defined as

$$\Lambda_{10} = 2 \log \left[\frac{\sup_{\theta \in \Theta_1} f(\mathbf{y}|\theta_0, \theta_1, M_1)}{\sup_{\theta_0 \in \Theta_0} f(\mathbf{y}|\theta_0, \theta_1 = 0, M_0)} \right].$$

Commonly, Λ_{10} is represented as a function of $\hat{\theta}$, the MLE of θ , and hence it can be expressed as

$$\Lambda_{10} = -2 \left[\ell_0(\hat{\theta}_0) - \ell_1(\hat{\theta}) \right]$$

with $\ell_k(\cdot) = \log f(\mathbf{y}|\cdot, M_k)$ denoting the log-likelihood function. This is a functional form which stresses the difference with *BFs*; in order to calculate the *BF* one has to integrate the likelihood function over the respective prior distribution for each model, while Λ calls for conditional evaluation of the likelihood function given the MLE. Note also that Λ is defined for nested hypotheses (or models) and thus $\Theta_0 \subseteq \Theta$ (or equivalently $M_0 \subseteq M_1$). For a comprehensive overview concerning the relationship as well as approximations between *BFs* and Λ see Kass and Raftery (1995).

Equations (2) & (4) make clear that in the case of conjugate analysis for the linear model described in (1), the *BF* comparing the two models is also merely a statistic, i.e. a function of the data only. Considering that $\Lambda_\ell = -n \log(1 - R_\ell^2)$ for a model M_ℓ (Wang et al. 2017), equation (2) shows that for linear regression and *g*-priors, the *BF* is a simple transformation of Λ . Furthermore, for the case of simple linear regression (and thus $p=1$), hypothesis in (1) can also be tested via the T_ν statistic with $\nu = n - 2$ degrees of freedom for which we have (Engle 1984):

$$T_\nu^2 = \nu \left[\exp\left(\frac{\Lambda}{n}\right) - 1 \right].$$

It is straightforward to show that for $p = 1$:

$$2 \log BF_{10} = (n - 2) \log(1 + g) - (n - 1) \log \left[1 + g \left(\frac{T_{n-2}^2}{n - 2} + 1 \right)^{-1} \right]. \quad (5)$$

Thus for simple linear regression and the *g*-priors setting, the resulting *BF* can be expressed analytically as a function of the T_ν statistic. Therefore, by combining individual T statistics, we can eventually also obtain the overall Bayesian evidence in the form of a *BF*.

3 Bayesian Evidence Synthesis

3.1 General Setting and notation

Suppose we have K datasets each resulting from the same data generating model. The sample size of each dataset is denoted by n_k and the size of the combined dataset is given by $N = \sum_{k=1}^K n_k$. Here we focus on the "simple" regression case where $p = 1$ and X_1 can be either a continuous scalar covariate or a binary factor. The latter represents the case of a simple *t*-test comparing two groups, probably the most common problem encountered in the meta-analytic context. We assume that each dataset (or study) provides us with a statistic \mathcal{T}_k arising from dataset k with response data \mathbf{y}_k and it evaluates the (same) null hypothesis H_0 . This can be a Student *t*-test statistic T_k (i.e. the *t* statistic from study k with $\nu_k = n_k - 2$ degrees of freedom), a likelihood ratio statistic Λ_k , or, quite unlikely currently, a Bayes factor $BF_{10k} = BF_k$. We will refer to a T_k statistic as a *directional* statistic, as it retains information about the sign of the effect, while Λ and *BF* are *undirectional* (with respect to the null and alternative defined in (1)).

Furthermore, we will denote by $\widehat{\beta}_k$ the absolute estimated effect from study k . Note that for each of the $T_k = \widehat{\beta}_k / SE(\widehat{\beta}_k)$, it holds that $T_k \sim t_{n_k-2}(\sqrt{ss_k} \beta_k^{sd})$, where $t_\nu(c)$ a *t* distribution with ν degrees of freedom and non-centrality parameter c , $ss_k = (n_k - 1) s_k^2$ with s_k^2 denoting the sample variance of X_1 in dataset k and $\beta^{sd} = \beta \tau^{-\frac{1}{2}}$ is the corresponding standardized, common across studies, effect size parameter; reminder: τ^{-1} is the error variance in the regression model. For the case where X_1 is a binary factor (i.e. we have a two-sample *t*-test), then $ss_k = \frac{n_{k(1)} n_{k(2)}}{n_{k(1)} + n_{k(2)}}$, where $n_{k(j)}$ is the sample size of group j at study k .

Each \mathcal{T}_k is comparing the same models M_0 and M_1 , with the latter describing the addition of one covariate to M_0 . In what follows, we will stick to the case where M_0 is the null model which assumes a constant expected response across all observations and M_1 describes either a difference in means between two groups for a dichotomous X_1 , or the linear effect of a continuous X_1 on the response Y . The proposed methodology can be easily employed to accommodate model comparisons within the multiple linear regression setup as discussed in Section 2.2 (see Equation 4).

We denote by *BF*, Λ , T the true, unknown, values of the statistics that we would obtain if all raw

data were available, BF_k, Λ_k, T_k are the corresponding statistics calculated from the k -th dataset and $\widetilde{BF}, \widetilde{\Lambda}, \widetilde{T}$ denote the synthesized statistics from the respective sample specific test statistics $\{BF_k\}_{k=1}^K, \{\Lambda_k\}_{k=1}^K$ or $\{T_k\}_{k=1}^K$. Our interest lies in calculating $BF_{10} = BF$, i.e. the evidence taking into account all K datasets. Thus, we are interested in methods that result in $\widetilde{BF} \approx BF$.

3.2 The product of Bayesian evidence

It has been suggested that $\widetilde{BF} = \prod_{k=1}^K BF_k$ (Goodman 2016). Nevertheless, the multiplication of the BF s from the individual studies or sub-samples does not lead to the same value as the full combined BF . This approach suggests that the marginal likelihood of the full dataset can be obtained as the product of the marginal likelihoods of each sub-sample and thus $\widehat{m}^{(K)} = \prod_{k=1}^K m_k$ where $\widehat{m}^{(K)}$ denotes the estimate of the marginal likelihood of the whole sample based on the m_k marginal likelihoods of the K sub-samples. This claim is not true since (ignoring parameters and denoting data by \mathbf{y})

$$\begin{aligned} m^{(K)} &= f(\mathbf{y}^{(K)}) = f(\mathbf{y}_1, \mathbf{y}_1, \dots, \mathbf{y}_K) \\ &= f(\mathbf{y}_1) \times f(\mathbf{y}_2|\mathbf{y}_1) \times \dots \times f(\mathbf{y}_{K-1}|\mathbf{y}_{K-2}, \dots, \mathbf{y}_1) \times f(\mathbf{y}_K|\mathbf{y}_{K-1}, \dots, \mathbf{y}_1) \\ &= \prod_{k=1}^K f(\mathbf{y}_k|\mathbf{h}_k) \end{aligned}$$

where \mathbf{h}_k is the history of sub-sample k , i.e. the sub-samples incorporated to the analysis previously to \mathbf{y}_k , given by $\mathbf{h}_k = (\mathbf{y}_1, \dots, \mathbf{y}_{k-1})$ and $\mathbf{h}_1 = \emptyset$. Here, we assume exchangeability between studies, which means that the ordering of the samples does not matter for these calculations. Thus, the time ordering will not influence our model and the association between Y and X_1 . Therefore the combined marginal likelihood $m^{(K)}$ is the product of the marginal likelihoods, which are the *conditional* marginal likelihoods of \mathbf{y}_k given the previous history \mathbf{h}_k , given by

$$f(\mathbf{y}_k|\mathbf{h}_k) = \int f(\mathbf{y}_k|\boldsymbol{\theta})p(\boldsymbol{\theta}|\mathbf{h}_k)d\boldsymbol{\theta}$$

where $p(\boldsymbol{\theta}|\mathbf{h}^{(k)})$ the posterior distribution of the relevant parameters for the data $\mathbf{y}^{(1)}, \dots, \mathbf{y}^{(k-1)}$. This expression is obviously different from the original intuition about the simple product of the marginal likelihoods and its computation requires the full set of raw data.

Another way to realize why simple multiplication of BF s is not an appropriate evidence synthesis method is to consider that BF is a function of Λ and under the null hypothesis, $\Lambda|H_0 \sim \chi_1^2$. From a frequentist point of view, multiplication of the likelihood functions would further imply that $\widetilde{\Lambda} \approx \sum_k \Lambda_k$ which is not true. In order for this to become more clear one has to recognize that $\sum_k \Lambda_k|H_0 \sim \chi_k^2$ while $\Lambda \sim \chi_1^2$. A demonstrative example of why the product of BF_k s resulting from partitioning a dataset is not the same as the BF computed from the pooled data is provided in Rouder and Morey (2011).

3.3 Bayesian Evidence Synthesis using Cauchy priors

Bayesian t -tests employing BF s have received attention mainly in the psychometric literature; see for example Gönen et al. (2005), Rouder et al. (2009) and Gronau et al. (2019).

An alternative to the conjugate prior setup and the g -prior specification for testing (1), is to use the a standard Cauchy prior on β^{sd} as originally suggested by Zellner and Siow (1980). The Zellner-Siow prior is frequently combined with a Jeffreys prior for τ which sometimes is referred to as the Jeffreys-Zellner-Siow prior (JZS) (Rouder et al. 2009). This prior can be considered as a mixture of g -priors (Liang et al. 2008) since it can be written in the following hierarchical manner:

$$\beta|\sigma_\beta^2 \sim N(0, \sigma_\beta^2) \text{ and } \sigma_\beta^2 \sim \text{Inv-}\chi_1^2.$$

The Cauchy prior is a less popular choice in variable selection than the g -priors or their recent extension, the hyper- g priors (Liang et al. 2008). The main reason was their computational intractability

which was partially solved by the recent work of Liang et al. (2008), where the mixture of g -prior representation lead to one-dimensional integrals which can be easily solved numerically.

Rouder et al. (2009) adopt a slightly different parameterization for the two sample t -test (originally suggested in (Gönen et al. 2005)) so that inference is based directly on the β^{sd} parameter as described here. This parameterization essentially translates to the X_1 variable being centered in the univariable regression setup (Wetzels and Wagenmakers 2012). We may write the JZS Bayes factor as a function of the $T = T_\nu$ statistic, by the following expression (Gronau et al. 2019)

$$BF_{10} = \frac{\int_{-\infty}^{+\infty} f_\nu(T; \sqrt{ss_x}\beta^{sd})g(\beta^{sd})d\beta^{sd}}{f_\nu(T; 0)}$$

where $f_\nu(\cdot; c)$ the pdf of a t -distributed random variable with ν degrees of freedom and non-centrality parameter c and $g(\cdot) = f_1(\cdot; 0)$ the pdf of a standard Cauchy distribution. The number of degrees of freedom depends on whether one- or two- sample t -tests are employed ($n - 1$ or $n - 2$ respectively).

Rouder and Morey (2011) suggest treating the individual T_k statistics that arise from different studies as a sample of data points from respective t distributions. With this formulation the only parameter in both models is β^{sd} which is 0 under H_0 and follows a standard Cauchy distribution under H_1 . Based on that formulation, a meta-analytic BF is suggested (here generalized for the univariable regression setting):

$$\widetilde{BF}_{10} = \frac{\int_{-\infty}^{+\infty} \prod_{k=1}^K f_{\nu_k}(T_k; \sqrt{ss_k}\beta^{sd})g(\beta^{sd})d\beta^{sd}}{\prod_{k=1}^K f_{\nu_k}(T_k; 0)}.$$

This method for the synthesis of Bayesian evidence will be abbreviated in the following sections as *Meta-BF_{JZS}*.

4 Proposed methods: Bayes Factors Synthesis under the g -priors

4.1 Evidence synthesis and availability of data

In order to calculate *Meta-BF_{JZS}*, one needs to have access to T_k , n_k and ss_k for all studies. However, following a systematic review, the level of detail of the available information concerning the effect of a variable can vary across different studies (Higgins et al. 2019). For example, it is not uncommon that information concerning ss_k , i.e. the sum of squares of the covariate X_1 , is not available in all studies. Another possibility is that, in some studies, only the effect estimates are presented in the form of $\widehat{\beta}_k$ but not the observed values of the test statistics T_k (thus with unknown $SE(\widehat{\beta}_k)$).

We therefore present methods for Bayesian evidence synthesis for three scenarios of minimum data availability. We employ the terms directional/undirectional and one-sided/two-sided interchangeably, since they coincide for the case of a testing the effect of a single covariate.

- **Case 1 - Detailed Information (D):** In this scenario, one- or two- sided \mathcal{T}_k (T , Λ , T^2 , p -value, or BF_k), n_k , effect direction and ss_k are available from all K studies. This scenario reasonably appears when considering evidence synthesis for the t -test case as it simply requires the sample size per group to be reported. For the simple linear regression case, it requires to record the observed sum of squares ss_k of the covariate X_1 for each study k .
- **Case 2 - Partial Information (P):** One- or two- sided \mathcal{T}_k (T , Λ , T^2 , p -value, or BF_k), n_k and effect direction available. For binary factors X_1 this scenario translates to unknown sample sizes per group. For continuous covariates X_1 , this corresponds to unreported ss_k - which is a common scenario (Becker and Wu 2007).
- **Case 3 - Limited Information (L):** Only two-sided \mathcal{T}_k (Λ , T^2 , two-sided p -value, or BF_k) and n_k available without knowing the direction of the effect. In this, admittedly extreme, scenario, there is no information on the direction of the effect from all the K datasets and effect measures are limited to an unidirectional statistic, e.g. (Cucherat et al. 2000).

Below we present methods for calculating g -priors based \widetilde{BF} for the above varying levels of available information. We present an alternative for JZS -based \widetilde{BF} when the same level of detail is obtainable (Case 1- D), but also for the scenario with more restricted access to information regarding the effect under study (Cases 2 & 3).

As with $Meta-BF_{JZS}$, we approach the problem through the prism of synthesizing the frequentist test statistics (mainly T_k) and then transform the combined evidence into \widetilde{BF} . This serves a pragmatic purpose, since most published studies employ the classical statistical approach, but it also demonstrates how the methods can be used for translating frequentist evidence into Bayesian. We, therefore, aim at arriving at a \widetilde{BF}_g (as in: g -prior based meta analytic BF), by synthesizing the K available evidence measures. As all the statistics discussed here (Λ_k , T_k^2 , two-sided p -value, BF_k) can be transformed to a T^2 statistic, we discuss below methods for synthesizing those T_k^2 statistics into a \widetilde{BF}_g .

Note that, the general problem presented in (1) is a two-sided hypothesis test. Evidence synthesis on the basis of unidirectional statistics (like the ones mentioned in Case 3) comes with additional complications. Specifically, two-sided statistics are not one-to-one transformations of the underlying one-sided statistics, and thus evidence synthesis methods might fall short (Held and Ott 2018). As an illustration, consider a standard t -test and a set of data that arises from H_0 , that is, the true $\beta = 0$. A random split of the data in $K=2$ partitions could result in $\hat{\beta}_1 > 0$ and $\hat{\beta}_2 < 0$, or vice-versa. Depending on the magnitude of the sub-samples' deviations from H_0 , this in turn could translate to a pair of small two-sided p -values or, equivalently, large test statistics (i.e. T^2 , Λ or BF). Synthesizing any of these two-sided statistics without taking into account the sign of β_k would result to a misleading combined estimate. Therefore, for Cases 1 & 2 evidence synthesis will be on the basis of the T_k statistics, while for Case 3, we will employ two-sided p -value combination methods.

4.2 Methods for Detailed information (D)

In this section we deal with the full information scenario where the study specific original test statistics T_k are directly available or can be obtained from other reported test statistics \mathcal{T}_k and the signs of $\hat{\beta}_k$.

By employing a technique that involves a normal approximation for the T_k statistics, the overall \widetilde{T} can be estimated by

$$\widetilde{T} = \sum_{k=1}^K w_k \left[\left(\mathbb{1}_{\{\hat{\beta}_k > 0\}} - \mathbb{1}_{\{\hat{\beta}_k < 0\}} \right) \times \sqrt{T_k^2} \right] \quad (6)$$

where $\mathbb{1}_{\{\cdot\}}$ the indicator function. Note that we present results on the basis of $\sqrt{T_k^2} = |T_k|$ in order to make the connection with BF_k obvious. This meta-analytic test statistic is based on the inverse normal method, commonly employed in the field of adaptive trial design where test statistics from different stages are combined (Lehmacher and Wassmer 1999). Note that for known and equal variance across studies, and weights that equal $w_k = \sqrt{n_k/N}$, then - by abusing notation and referring by T to the corresponding Z statistic - $\widetilde{T} = T$.

By combining (5) & (6) one can obtain \widetilde{BF} by utilizing either T_k^2 or BF_k . Weights w_k should be chosen appropriately and satisfy $\sum_{k=1}^K w_k^2 = 1$ (Lehmacher and Wassmer 1999). By incorporating inverse variance weighting and taking into account the relative variances of the T_k statistics, the weights are given by $w_k = \sqrt{\frac{v_k^{-1}}{\sum_{k=1}^K v_k^{-1}}}$ where:

$$v_k = H(\nu_k/2)^{-2} \frac{\nu_k}{ss_k(\nu_k - 2)} + \left[\left(\frac{\nu_k}{\nu_k - 2} \right) H(\nu_k/2)^{-2} - 1 \right] d_k^2 \quad (7)$$

$$\text{with } d_k = \frac{T_k}{H(\nu_k/2)\sqrt{ss_k}} \text{ and } H(z) = \sqrt{z} \left[\frac{\Gamma(z - 1/2)}{\Gamma(z)} \right].$$

These weights are the result of combining Equation 23 from Malzahn et al. (2000), and the requirement for $\sum_{k=1}^K w_k^2 = 1$. They essentially apply inverse-variance weighting for the individual T_k statistics and treats their sum as a sum of independent $N(\sqrt{ss_k}\beta_k^{sd}, 1)$ variables. In the following, we will refer to this meta-analytic estimator as $Meta-BF_g^{(D)}$.

4.3 Methods for Partial Information (P)

In this section we deal with the more realistic scenario where partial information is available. Hence, if the variance component ss_k is not available for every study then we cannot implement neither the Meta- $BF_g^{(D)}$ approach proposed in Section 4.2 nor the JZS based \widetilde{BF} of Rouder et al. (2009) presented in Section 3.3. In our proposed Meta- $BF_g^{(D)}$, the problem is mainly that we cannot obtain the weight w_k of each study k since (6) requires the values of the corresponding sum of squares $ss_k = (n_k - 1)s_k^2$.

In such cases, a simple but reasonable solution might be given by setting $w_k = \omega_k = \sqrt{n_k/N}$ in equation 7 and then employ a normal approximation-based inverse variance weighing for all the T_k statistics. This is equivalent to adopting the assumption that all variances are known and equal across the K studies. This approach will be abbreviated as $Meta-BF_g^{(P)}$.

4.4 Methods for Limited Information (L)

Finally, we consider the case where the available information is restricted to p -values or test statistics where the direction of the effect is not evident. Hence, if we only have access to p -values from an unidirectional test then we can think of using any p -value combination method for the synthesis of frequentist evidence, and then transform the result to a Meta-BF (\widetilde{BF}). In order to synthesize a vector of p -values $\mathbf{p} = \{p_1, \dots, p_K\}$ testing the same H_0 vs. H_1 coming from K independent studies the result that

$$p_i | H_0 \sim Uniform(0, 1) \quad \forall i \in \{1 \dots K\} \quad (8)$$

is of fundamental value. By using (8), several combination methods exist in order to obtain an overall p -value $\tilde{p} = f(\mathbf{p})$ which will act as a global measure of departure from H_0 . Two of the most common statistics for combining p -values, for which the distribution under the null hypothesis is available are given by:

(a) the Fisher's method (Fisher 1934), and (b) the Stouffer's method (Stouffer et al. 1949). The p -value based statistics for these two methods are given by

$$\text{Fisher's method: } S_F = -2 \sum_{k=1}^K \log(p_k) \sim \chi_{2K}^2, \quad (9)$$

$$\text{Stouffer's method: } S_S = \sum_{k=1}^K \Phi^{-1}(1 - p_k) \sim N(0, K). \quad (10)$$

The following strategy could be adopted in order to obtain the Meta-BF, \widetilde{BF} , from a (two-sided only) \mathcal{T}_k :

1. For each \mathcal{T}_k , calculate the associated (two-sided) p -value;
2. Synthesize the resulting vector $\mathbf{p} = \{p_1, \dots, p_K\}$ via (9) or (10) and obtain \tilde{p} ;
3. Transform \tilde{p} to an appropriate $\tilde{\mathcal{T}}_k$.
4. Obtain \widetilde{BF} from $\tilde{\mathcal{T}}_k$ using (5).

However, as discussed above, such an approach would suffer from the loss of directionality in each p_k .

Owen et al. (2009) addresses this issue by exploring a long-ignored method for p -value combination dating back to Pearson (1934). It can be briefly described as running a Fisher or Stouffer style test for left sided alternatives and one for right-sided alternatives and then considering the most extreme of the two. In other words, for a set of study specific two-sided only p -values $\mathbf{p} = \{p_1, \dots, p_K\}$, and by implementing the Stouffer's method, we then consider:

$$S_S^L = \sum_{k=1}^K \Phi^{-1}(1 - p_k/2), \quad S_S^R = \sum_{k=1}^K \Phi^{-1}(p_k/2), \quad \text{and} \quad S_S^C = \max(S_S^R, S_S^L). \quad (11)$$

Note that $S_S^R = -S_S^L$ and $S_S^C = |S_S^L| = |S_S^R|$. This p -value based statistic can be adopted as a combination method if no information on the direction of the β_k is available from the K studies. A modified weighted version of (11) could be employed with $S_S^C = \sum_{k=1}^K \omega_k |T_k|$ as an approximation to \widetilde{T} which can be transformed to a \widetilde{BF} via (5).

We will refer to this method as $Meta-BF_g^{(L)}$. The methods described alongside the required ingredients for their implementation and abbreviations used in what follows are summarized in Table 1.

5 Simulation Study

In this section we compare the performance of the discussed methods through two simulation studies. In the first one the total sample size N is retained fixed. The main goal is to assess whether the meta-analytic \widetilde{BF} s are providing the same level of evidence as the BF obtained from the full data.

The second simulation follows the standard simulation approach typically employed for the comparison of meta-analytic methods, where the total and within-study sample size is random (Pateras et al. 2018). The main goal of this exercise is to compare the operational characteristics of the approaches under study for increasing K .

5.1 Fixed sample size

5.1.1 Simulation Plan

The data generating model was $Y \sim N(\beta X_1, 1)$ and X_1 was either a group indicator, representing an independent samples t -test, or a fixed realization of a $N(0, \frac{1}{4})$ variate, materializing a simple linear regression scenario with variance approximately equal to the variance used in the t -test simulation scenario. The total sample size was set to $N = 1000$ and we explored $K = \{2, 5, 10\}$ partitions of the full data set. Group sizes were balanced in the case of the t -test. We employ two over-arching scenarios for the partitions, one where $n_k = N/K \Rightarrow \omega_k^2 = 1/K$ and thus all sub-samples were balanced and one with unequal partition where:

- For $K = 2$, $\omega_1^2 = 0.7$ and $\omega_2^2 = 0.3$
- For $K = 5$, $\{\omega_k^2\}_{k=1}^5 = \{0.05, 0.1, 0.15, 0.3, 0.4\}$
- For $K = 10$, $\omega_1^2 = \dots = \omega_6^2 = 0.05$, $\omega_7^2 = 0.1$ and $\omega_8^2 = \omega_9^2 = \omega_{10}^2 = 0.2$.

We will refer to the former as the EQ scenario, and the latter as the UNEQ. Concerning the effect size, we assess the null hypothesis/model and levels of moderate departure from the null, i.e. $\beta \in \{0, 0.1, 0.2, 0.3\}$. We conducted $R = 1000$ replications per scenario.

For each of the K sub-samples per scenario, BF_k and T_k were calculated as well as the BF from the full dataset. For the g -prior setup we consider $g_k = n_k$ and $g = N$. As a basic assessment of the

	Minimum data requirements per study		
Two sided only evidence (Λ , T^2 , p -value, or BF_k)	✓	✓	✓
Sample size per study (n_k)	✓	✓	✓
Effect direction ($sign(\hat{\beta}_k)$)	✓	✓	
Sum of squares of covariate (ss_k)	✓		
Method (Section)	$Meta-BF_{JZS}$ (3.3)	$Meta-BF_g^{(P)}$ (4.3)	$Meta-BF_g^{(L)}$ (4.4)
	$Meta-BF_g^{(D)}$ (4.2)		

Table 1: Summary of methods for Bayesian evidence synthesis with respect to the available data from included studies in a meta analysis

performance of each method, we focus on the quantification of evidence in a discrete sense. According to Kass and Raftery (1995), the evidence provided by a BF_{10} can be categorized in four levels of evidence presented in in Table 2. Per simulation scenario and for each method, a contingency table is constructed presenting the cross-classification of the levels of evidence for the (full dataset) BF and each of the meta-analytic \widetilde{BF} proposed here. Agreement in terms of correct classification in the categories of strength of evidence is assessed via the weighted version of Cohen’s κ statistic, using quadratic weights to assign larger importance to larger discordances (Fleiss et al. 1969). The categories used to evaluate agreement are shown in Table 3, alongside the colour scale used for visualization of the level of agreement.

$2 \log BF_{10}$	BF_{10}	Evidence against H_0
0 - 2	1 - 3	Not worth more than a bare mention
2 - 6	3 - 20	Positive
6 - 10	20 - 150	Strong
>10	>150	Very strong

Table 2: Level of evidence provided by BF

Cohen’s κ	Level of Agreement
< 0	No agreement
0 – 0.2	Slight agreement
0.2 – 0.4	Fair agreement
0.4 – 0.6	Moderate agreement
0.6 – 0.8	Substantial agreement
0.8 – 1.0	Almost Perfect agreement

Table 3: Level of agreement suggested by (weighted) Cohen’s κ

5.1.2 Results

The results of the first simulation exercise are shown in Figure 1. It presents the association scatterplots between $2 \log BF$ and $2 \log \widetilde{BF}$, for the case of the two-sample comparison of means (t -test) and the EQ scenario. Results for linear regression as well as for the UNEQ scenario were very similar and can be found in the online supplementary material.

The performance of all $Meta-BF_g$ methods except for the limited information $Meta-BF_g^{(L)}$ is very satisfactory for the settings studied - see Fig. 1. This suggests that the known variance approximation of the T_k statistics is very adequate and knowledge of the individual ss_k is not of cardinal importance for the successful synthesis of a small to moderate number of BF s. From Fig. 1 we can observe a relative discordance for large values of BF s corresponding to larger effect sizes (i.e. when $\beta = 0.3$, on the top right corner of the graphs). Nevertheless, this does not seem to be of major importance in terms of evidence classification.

For the limited information scenario, the effect of the availability of only two-sided evidence is obvious for $Meta-BF_g^{(L)}$; see Fig. 1. When $\beta = 0$, the combined evidence systematically understate evidence for H_0 as a result of the shortcomings of synthesizing undirected p -values (or test statistics) as discussed in Section 4.4. For $K = 2$ the method performs surprisingly well for $\beta > 0$. But as K increases, it overstates the evidence for H_1 an effect partly mitigated for increasing β , as then, the number of T_k s with concordant signs naturally increases.

5.2 Random sample size

5.2.1 Simulation Plan

For this simulation scenario, the total sample size was drawn from a discrete $Uni(800, 10000)$ and was split in K parts of random size, where $K \in \{2, \dots, 50\}$. Model parameters were the same as in the fixed sample size case. We compared all *Meta-BF* methods according to their empirical bias, Root Mean Squared Error (RMSE) as well as the weighted Cohen's κ statistic. For the calculation of the empirical bias and the RMSE, the BF was treated as a parameter while the respective \widetilde{BF} as the estimator. Therefore, for each simulation scenario:

$$Bias = \frac{\sum_{j=1}^R (\widetilde{BF} - BF)}{R}$$

$$RMSE = \sqrt{\frac{\sum_{j=1}^R (\widetilde{BF} - BF)^2}{R}}$$

where R the number of simulations. The limited information estimator $Meta-BF_g^{(L)}$ was not included in this comparison as it utilizes substantially less information from each study and, as it is shown in Section 5.1.2 it does not perform satisfactorily for a large number of studies.

Comparisons are made on the $2 \log BF$ scale. \widetilde{BF} s based on the g -prior and JZS prior are compared to the g -prior-based and JZS prior-based BF respectively.

5.2.2 Results

Figure 2 presents the results for the bias, RMSE and weighted Cohen's κ statistic for increasing K , when the total sample size as well as ω_l for subsample $k \in \{1, \dots, K\}$ is random. For the two-sample comparison of means (t -test), the effect of the lack of appropriate weighting imposed by $Meta-BF_g^{(P)}$ becomes more apparent as K grows. It exhibits higher bias and RMSE. Nevertheless, when it comes to the evidence quantification in a discrete manner, its performance is more than adequate, as indicated by the large (> 0.9) values of the weighted Cohen's κ statistic. The methods that utilize information about ss_k , the coefficient's sum of squares in each subsample ($Meta-BF_g^{(D)}$ and $Meta-BF_{JZS}$), are not substantially influenced by increasing K , with a notable difference on the bias: $Meta-BF_{JZS}$ exhibits negative bias, while $Meta-BF_g^{(D)}$ yields on average larger values than BF , with the bias being of slightly smaller absolute value compared to $Meta-BF_{JZS}$. $Meta-BF_g^{(D)}$ performs marginally better in terms of RMSE, for increasing β .

Results for the regression case are slightly different. One notable difference is that $ss_k = (n_k - 1)\hat{\sigma}_{X_1}^2$ and thus exhibiting more variability than in the t -test case where $ss_k = \frac{n_k}{2}$. This increased heterogeneity embodied in the ss_k , is handled in a better way by the $Meta-BF_g^{(D)}$ approach compared to $Meta-BF_{JZS}$, in terms of bias and RMSE. Such a behavior does not come as a surprise given that the meta-analytic modeling is only partially implemented on the (variable) non-centrality parameter of the individual t statistics in the $Meta-BF_{JZS}$ formulation. For the t -test, a comparable situation would be the case where the allocation ratios among the studies were different and not all equal to one as we have considered in this simulation. Note that the $Meta-BF_{JZS}$ approach is performing better under H_0 in specific metrics (concerning the bias in the t -test and the Cohen's κ in regression). This may be attributed to the smaller variability in the (observed) non-centrality parameters imposed by $\beta = 0$.

6 Illustrative Example

We apply the methods discussed in a meta-analysis from the field of psychology. In Bolier et al. (2013), the authors present a systematic review and meta-analysis of randomized controlled trials assessing the effects of positive psychology interventions. The use of positive psychological interventions may be considered as a complementary strategy in mental health promotion and treatment. The efficacy of the interventions was assessed in several outcomes. Here, we focus on the outcome of psychological




Figure 1: Scatter plots and weighted Cohen's κ statistics for the agreement between BF and \widetilde{BF} for binary X (t -test) and EQ scenario. $2 \log BF_{10}$ are shown, except for $\beta = 0$ where $2 \log BF_{01}$ is shown. Grid lines indicate the categories of level of evidence as described in Table 2. Values <-1 and >12 are shown as equal.




Figure 2: Bias, RMSE and weighted Cohen’s κ versus K (number of subsamples) for random sample sizes. The scale is $2 \log BF$. Columns indicate different models (regression and t -test) while sub-rows within each metric correspond to different values of β . The value of 0 for the Bias is highlighted by a dashed line.

well being which exhibited low levels of heterogeneity and thus a fixed-effects model (like the one treated in the present work) is deemed appropriate. Psychological well being was measured by means of questionnaires and it was considered a continuous outcome.

There were 20 studies included in the meta analysis of the psychological well being outcome measured at the end of the intervention. Since all measurements were continuous but instruments (i.e. questionnaires) varied across studies, the parameter of interest was the standardized mean difference. Table 4 shows the studies' results in terms of T_k statistics, as well as the sample sizes per arm. We have added the calculated BF_k s using the g -prior ($BF_k^{(g)}$) as well as the ones using the JZS prior ($BF_k^{(JZS)}$). The last two columns show the resulting (squared) weights of the studies, for the $Meta-BF_g^{(L)}/Meta-BF_g^{(P)}$ as well as for the $Meta-BF_g^{(D)}$ approach.

Table 4 presents Bayes Factors in the form BF_{10} . The relationship between the $BF^{(g)}$ s and $BF^{(JZS)}$ can be sketched. In general, $BF^{(JZS)} > BF^{(g)}$, except for when the sample size is relatively small, in which case the $BF^{(JZS)}$ is more conservative (e.g. Study 5). $BF^{(JZS)}$ gives less support for H_0 even when observed differences are very close to 0 (Studies 6 & 19).

For the \widetilde{BF}_g s, differences in the weighting schemes are also of interest. When the weights take into account the allocation ratio within the studies as well as the observed effects ($w_k^{(v)}$), both factors that contribute to the variance and mean of the t -statistics, studies with largely unequal allocation ratio are downweighted compared to the weights based solely on the total sample sizes of the studies ($w_k^{(\omega)}$). Take for example studies 12 & 17: They are comparable with respect to their relative total sample sizes (364 vs 411) but only 70/411 from study 17 were allocated to group 2. This has considerable impact on its relative weight as $w_{17}^{(v)}=0.097$ while $w_{12}^{(v)}=0.15$.

The results for the meta-analytic \widetilde{BF} s are presented in Table 5. \widetilde{BF}_{JZS} results to a considerably larger value, but belonging in the same evidence level as \widetilde{BF}_g with respect to Table 2, indicating very strong evidence for H_1 . The two methods which employ less data information result to slightly smaller BF s ($2 \log BF \approx 9$). We see that $Meta-BF_g^{(L)}$ performs very well in this case. This is due to the fact that only two of the studies demonstrated a T statistic in the opposite direction. Based on the simulation results one would expect $Meta-BF_g^{(P)}$ to exhibit a larger value in comparison to $Meta-BF_g^{(D)}$, but the effect of the discrepant weights due to combinations of unequal allocations and observed effect sizes results to the opposite.

7 Discussion

In this work we explored methods for synthesizing evidence in the form of Bayes Factors. We do so for the standard meta analysis paradigm, i.e. when the available data from the separate studies are on the aggregate level. We present methods for varying levels of detail in the available information and compare their performance. By utilizing the common g -prior setup, we show how the problem of BF synthesis can be translated to the synthesis of T statistics for simple linear regression. An analytic relationship between BF 's and standard test statistics can be found in several hypothesis testing contexts, outside the g -prior setup (Held and Ott 2018; Johnson 2005, 2008). This is also the case for the JZS prior though it involves numerical integration.

The JZS -prior can be seen as a mixture of g priors. Alongside popularity, computational efficiency and understandable interpretation of g -priors, resulting BF s have been shown to be monotone functions of the F statistic for testing (1) and therefore gives the uniformly most powerful invariant test (Shively and Walker 2018). This holds also for the mixture of g -priors, and it is evident via the results presented here for the univariate regression example, considering that $F = T^2$.

Meta-analysis is mostly involved with parameter estimates. The synthesis of BF s involves the synthesis of evidence concerning hypothesis testing. In frequentist inference, the analogous would be p -value combination methods. However, p -value combination methods are meant to result to a statistic that has a known distributional form under the intersection of all null hypotheses being tested $H_0 = \cap_{i=1}^K \{H_{0i}\}$. Consequently their purpose is to retain frequentist testing operational characteristics (type I error control) and not to synthesize test statistics. They thus fall short for the BF synthesis context. We mitigate this shortcoming by employing the weighted test statistic from Stouffer's method,

Study	T_k	$n_{k(1)}$	$n_{k(2)}$	n_k	$2 \log BF^{(g)}$	$2 \log BF^{(JZS)}$	$BF^{(g)}$	$BF^{(JZS)}$	$w_k^{(\omega)}$	$w_k^{(v)}$
1	-0.22	26	27	53	-3.9	-3.1	0.1	0.2	0.015	0.022
2	0.50	17	15	32	-3.2	-2.5	0.2	0.3	0.009	0.013
3	2.58	33	32	65	2.1	2.5	2.9	3.5	0.019	0.026
4	0.71	10	10	20	-2.5	-1.9	0.3	0.4	0.006	0.008
5	3.6	8	8	16	6.2	5.4	22.2	14.8	0.005	0.004
6	0.00	37	29	66	-4.2	-3.3	0.1	0.2	0.019	0.027
7	0.82	79	86	165	-4.4	-3.6	0.1	0.2	0.047	0.069
8	0.77	559	63	622	-5.8	-3.9	0.1	0.1	0.178	0.095
9	-0.08	35	38	73	-4.3	-3.4	0.1	0.2	0.021	0.030
10	1.72	25	25	50	-1.1	-0.5	0.6	0.8	0.014	0.020
11	0.86	80	37	117	-4.0	-3.1	0.1	0.2	0.033	0.042
12	2.19	187	177	364	-1.1	-0.3	0.6	0.9	0.104	0.150
13	2.68	153	89	242	1.6	2.4	2.2	3.3	0.069	0.093
14	1.44	48	54	102	-2.6	-1.8	0.3	0.4	0.029	0.042
15	1.21	804	138	942	-5.4	-3.8	0.1	0.1	0.269	0.198
16	1.35	13	10	23	-1.4	-1.0	0.5	0.6	0.007	0.009
17	0.11	341	70	411	-6.0	-4.5	0.0	0.1	0.117	0.097
18	1.84	11	9	20	0.1	0.2	1.0	1.1	0.006	0.007
19	0.00	36	42	78	-4.4	-3.5	0.1	0.2	0.022	0.032
20	0.51	20	17	37	-3.4	-2.6	0.2	0.3	0.011	0.015

Table 4: Study result from Bolier et al. (2013), alongside the resulting BF s and weights based on sample fraction as well as on v_k

Method	$2 \log \widetilde{BF}$	\widetilde{BF}
$Meta-BF_g^{(P)}$	9.00	90.1
$Meta-BF_g^{(L)}$	9.64	124.3
$Meta-BF_g^{(D)}$	11.83	372.1
$Meta-BF_{JZS}$	12.75	588.9

Table 5: Meta-analytic Bayes Factors for the illustrative example of Section 6

which, for proper weights (that sum to 1) and original T_k 's that are (approximately) normal, result to a correct \widetilde{T} and \widetilde{BF} for one-sided statistics (it is equivalent to the inverse normal method).

Our demonstration is limited to the case of a single covariate. Such a task could be very useful in simple but very common settings, as is the synthesis of evidence from a series of experiments, where testing is usually concerned with a difference between groups (e.g. randomized controlled trials). For more general problems where BF s are commonly employed, like variable/model selection, the extension is not straightforward. A similar strategy could be sought where regression slopes arising from independent datasets would be synthesized, though such a task does not come without complexity (Becker

and Wu 2007). Such complexities include not reported covariance matrices as well as systematically missing covariates.

We present meta-analytic Bayes Factors for varying levels of detail in the reported data from available studies. It is highly likely that in realistic scenarios one can encounter a combination of such levels, i.e. some studies reporting only sample sizes and unidirectional statistics, while others all the required ingredients for the computation of $Meta-BF_g^{(D)}$ or $Meta-BF_{JZS}$. The reasonable choice in such cases would be to synthesize the evidence with a method that balances between using the most of the available information and is supported by some theoretical justification. For example, if information on the ss_k is missing for some k it is not straightforward how the weights should be differentiated between studies with and without available ss_k , and perhaps the preferred solution would be to proceed with weighting all studies by the sample sizes.

A limitation of our work is the treatment of only of fixed-effects models, i.e. absence of heterogeneity. Hypothesis testing in the presence of heterogeneity involves several considerations which are further perplexed by the employment of Bayes Factors (Higgins et al. 2009). The study of Meta-Analytic Bayes Factors under the random-effects model requires considerable attention beyond the scope of this work.

The difficulties arising in the computation of BF s makes meta-analytic methods also relevant in data partitioning for Big Data when computations become CPU-intensive as well as data-intensive and cannot be handled by a single machine (Zhao et al. 2013). P -value combination and other meta-analytic methods have been proposed in this field in order to tackle parallel processing of data partition for frequentist analyses (Tsamardinos et al. 2019).

Utilizing frequentist techniques in order to acquire Bayesian evidence may leave a bitter taste to Bayesians. A fully Bayesian approach would require placing a likelihood model on the BF s which is not straightforward. One could take advantage of recent results concerning the sampling distribution of BF s (a shifted non-central chi-square for g priors, see Zhou and Guan 2018) and pursue inference based on a full posterior or predictive distribution. However, such an exercise would strip BF s of the attractive and easily communicable feature of a single piece of evidence for or against a model.

Clearly, a reality where BF s are the only level of evidence available in published studies (even more BF s employing the same priors) is far from close. However, the translation and interplay between frequentist and Bayesian evidence is of interest and recent meta-analyses are pointed towards re-assessing frequentist evidence through a Bayesian prism (Bittl et al. 2019; Monden et al. 2016). The methods developed and assessed in this work are fit towards this direction.

References

- Becker, B. J. and Wu, M.-J. (2007). “The synthesis of regression slopes in meta-analysis”. *Statistical Science* 22 (3), pp. 414–429.
- Benjamin, D. J., Berger, J. O., Johannesson, M., Nosek, B. A., Wagenmakers, E.-J., Berk, R., Bollen, K. A., Brembs, B., Brown, L., Camerer, C., et al. (2018). “Redefine statistical significance”. *Nature Human Behaviour* 2 (1), pp. 6–10.
- Berger, J. O. and Sellke, T. (1987). “Testing a point null hypothesis: The irreconcilability of p values and evidence”. *Journal of the American Statistical Association* 82 (397), pp. 112–122.
- Bittl, J. A., He, Y., Baber, U., Feldman, R. L., Mering, G. O. von, and Kaul, S. (2019). “Bayes factor meta-analysis of the mortality claim for peripheral paclitaxel-eluting devices”. *JACC: Cardiovascular Interventions* 12 (24), pp. 2528–2537.
- Bolier, L., Haverman, M., Westerhof, G. J., Riper, H., Smit, F., and Bohlmeijer, E. (2013). “Positive psychology interventions: a meta-analysis of randomized controlled studies”. *BMC public health* 13 (1), p. 119.
- Cucherat, M, Haugh, M., Gooch, M, and Boissel, J.-P. (2000). “Evidence of clinical efficacy of homeopathy”. *European journal of clinical pharmacology* 56 (1), pp. 27–33.
- Engle, R. F. (1984). “Wald, likelihood ratio, and Lagrange multiplier tests in econometrics”. *Handbook of Econometrics* 2, pp. 775–826.
- Fisher, R. (1934). *Statistical methods for research workers*. Edinburgh: Oliver and Boyd.

- Fleiss, J. L., Cohen, J., and Everitt, B. S. (1969). “Large sample standard errors of kappa and weighted kappa.” *Psychological bulletin* 72 (5), p. 323.
- Gönen, M., Johnson, W. O., Lu, Y., and Westfall, P. H. (2005). “The Bayesian two-sample t test”. *The American Statistician* 59 (3), pp. 252–257.
- Goodman, S. N. (2016). “Aligning statistical and scientific reasoning”. *Science* 352 (6290), pp. 1180–1181.
- Gronau, Q. F., Ly, A., and Wagenmakers, E.-J. (2019). “Informed Bayesian t-tests”. *The American Statistician*, pp. 1–14.
- Heard, N. A. and Rubin-Delanchy, P. (2018). “Choosing between methods of combining-values”. *Biometrika* 105 (1), pp. 239–246.
- Hedges, L. V. and Olkin, I. (2014). *Statistical methods for meta-analysis*. Academic press.
- Held, L. and Ott, M. (2018). “On p-values and Bayes factors”. *Annual Review of Statistics and Its Application* 5, pp. 6.1–6.27.
- Higgins, J. P., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., and Welch, V. A. (2019). *Cochrane handbook for systematic reviews of interventions*. John Wiley & Sons.
- Higgins, J. P., Thompson, S. G., and Spiegelhalter, D. J. (2009). “A re-evaluation of random-effects meta-analysis”. *Journal of the Royal Statistical Society: Series A (Statistics in Society)* 172 (1), pp. 137–159.
- Jeffreys, H. (1961). *Theory of Probability (3rd edition)*. Oxford University Press.
- Johnson, V. E. (2005). “Bayes factors based on test statistics”. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 67 (5), pp. 689–701.
- (2008). “Properties of Bayes factors based on test statistics”. *Scandinavian Journal of statistics* 35 (2), pp. 354–368.
- Kass, R. E. (1993). “Bayes factors in practice”. *Journal of the Royal Statistical Society: Series D (The Statistician)* 42 (5), pp. 551–560.
- Kass, R. E. and Raftery, A. E. (1995). “Bayes factors”. *Journal of the American statistical association* 90 (430), pp. 773–795.
- Lehmacher, W. and Wassmer, G. (1999). “Adaptive sample size calculations in group sequential trials”. *Biometrics* 55 (4), pp. 1286–1290.
- Liang, F., Paulo, R., Molina, G., Clyde, M. A., and Berger, J. O. (2008). “Mixtures of g priors for Bayesian variable selection”. *Journal of the American Statistical Association* 103 (481), pp. 410–423.
- Malzahn, U., Böhning, D., and Holling, H. (2000). “Nonparametric estimation of heterogeneity variance for the standardised difference used in meta-analysis”. *Biometrika* 87 (3), pp. 619–632.
- Monden, R., Vos, S. de, Morey, R., Wagenmakers, E.-J., Jonge, P. de, and Roest, A. M. (2016). “Toward evidence-based medical statistics: a Bayesian analysis of double-blind placebo-controlled antidepressant trials in the treatment of anxiety disorders”. *International Journal of Methods in Psychiatric Research* 25 (4), pp. 299–308.
- O’Hagan, A. and Forster, J. J. (2004). *Kendall’s advanced theory of statistics, volume 2B: Bayesian inference*. Vol. 2. Arnold.
- Owen, A. B. et al. (2009). “Karl Pearson’s meta-analysis revisited”. *The Annals of Statistics* 37 (6B), pp. 3867–3892.
- Pateras, K., Nikolakopoulos, S., and Roes, K. (2018). “Data-generating models of dichotomous outcomes: heterogeneity in simulation studies for a random-effects meta-analysis”. *Statistics in medicine* 37 (7), pp. 1115–1124.
- Pearson, K. (1933). “On a method of determining whether a sample of size n supposed to have been drawn from a parent population having a known probability integral has probably been drawn at random”. *Biometrika* 25 (3/4), pp. 379–410.
- (1934). “On a New Method of Determining” Goodness of Fit”. *Biometrika* 26 (4), pp. 425–442.
- Rouder, J. N. and Morey, R. D. (2011). “A Bayes factor meta-analysis of Bem’s ESP claim”. *Psychonomic Bulletin & Review* 18 (4), pp. 682–689.
- Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., and Iverson, G. (2009). “Bayesian t tests for accepting and rejecting the null hypothesis”. *Psychonomic bulletin & review* 16 (2), pp. 225–237.

- Sellke, T., Bayarri, M., and Berger, J. O. (2001). “Calibration of p values for testing precise null hypotheses”. *The American Statistician* 55 (1), pp. 62–71.
- Shively, T. S. and Walker, S. G. (2018). “On Bayes factors for the linear model”. *Biometrika* 105 (3), pp. 739–744.
- Stouffer, S. A., Suchman, E. A., DeVinney, L. C., Star, S. A., and Williams Jr, R. M. (1949). “The american soldier: Adjustment during army life.”
- Tsamardinos, I., Borboudakis, G., Katsogridakis, P., Pratikakis, P., and Christophides, V. (2019). “A greedy feature selection algorithm for Big Data of high dimensionality”. *Machine learning* 108 (2), pp. 149–202.
- Wang, X., Jiang, B., and Liu, J. S. (2017). “Generalized R-squared for detecting dependence”. *Biometrika* 104 (1), pp. 129–139.
- Wetzels, R. and Wagenmakers, E.-J. (2012). “A default Bayesian hypothesis test for correlations and partial correlations”. *Psychonomic bulletin & review* 19 (6), pp. 1057–1064.
- Zellner, A. (1986). “On assessing prior distributions and Bayesian regression analysis with g-prior distributions”. *Bayesian inference and decision techniques*.
- Zellner, A. and Siow, A. (1980). “Posterior odds ratios for selected regression hypotheses”. *Trabajos de estadística y de investigación operativa* 31 (1), pp. 585–603.
- Zhao, Z., Zhang, R., Cox, J., Duling, D., and Sarle, W. (2013). “Massively parallel feature selection: an approach based on variance preservation”. *Machine learning* 92 (1), pp. 195–220.
- Zhou, Q. and Guan, Y. (2018). “On the null distribution of Bayes factors in linear regression”. *Journal of the American Statistical Association* 113 (523), pp. 1362–1371.