

Variable Selection Using Nearest Neighbor Gaussian Processes

Konstantin Posch^{a,*}, Maximilian Arbeiter^a, Martin Pleschberger^b, Juergen Pilz^a

^a*Department of Statistics, University Klagenfurt, Austria*

^b*KAI-Kompetenzzentrum Automobil- und Industrieelektronik GmbH, Villach, Austria*

Abstract

A novel Bayesian approach to the problem of variable selection using Gaussian process regression is proposed. The selection of the most relevant variables for a problem at hand often results in an increased interpretability and in many cases is an essential step in terms of model regularization. In detail, the proposed method relies on so-called nearest neighbor Gaussian processes, that can be considered as highly scalable approximations of classical Gaussian processes. To perform a variable selection the mean and the covariance function of the process are conditioned on a random set \mathcal{A} . This set holds the indices of variables that contribute to the model. While the specification of a priori beliefs regarding \mathcal{A} allows to control the number of selected variables, so-called reference priors are assigned to the remaining model parameters. The application of the reference priors ensures that the process covariance matrix is (numerically) robust. For the model inference a Metropolis within Gibbs algorithm is proposed. Based on simulated data, an approximation problem from computer experiments and two real-world datasets, the performance of the new approach is evaluated.

Keywords: Variable Selection, Hierarchical Bayes, Nearest Neighbor Gaussian Process, Model Uncertainty, Metropolis-Hastings Algorithm

1. Introduction

Due to the omnipresent and ongoing digitalization process more and more data is collected by companies, governments and organizations of all kinds. Consequently, the urge of analyzing dependencies between observed variables is also growing. Regression techniques are often the methods of choice for this task. The application of regression models is straight forward if there is a large number n of observations of a comparatively small number d of observed variables. However, often this does not hold true and things get challenging. Models start to over-fit, since the model complexity cannot be supported by the available sample size (Buehlmann et al., 2016). Moreover, it becomes difficult to identify the most influential variables. For this reason, approaches that automatically detect the most important variables gain increasing attention in many fields including genetics (Lee et al., 2003), astronomy (Zheng and Zhang, 2007), and economics (Foster and Stine, 2004).

In case that the dependencies of interest are purely linear simple linear regression models are sufficient for the analysis. There are many well-established approaches to perform variable selection in linear regression models (Tibshirani, 1996; Fan and Li, 2001; Zou and Hastie, 2005; Zou, 2006; Park and Casella, 2008; Ročková and George, 2014; Alhamzawi and Taha Mohammad Ali, 2018; Posch et al., 2020). However, in many practical applications linearity does not hold and more sophisticated methods are required. Recently, Gaussian process regression models (GPRMs), originally proposed by O'Hagan (1978), gained a lot of attention. GPRMs can be considered as an alternative to neural networks since a large class of neural network based regression models converges to a Gaussian Process (GP) in the limit of an infinite network (Neal, 1996). Alternatively, GPs can also be viewed at from the perspective of non-parametric Bayesian

*Corresponding author

Email address: Konstantin.Posch@aau.at (Konstantin Posch)

regression (Rasmussen and Williams, 2005). Here, Gaussian priors are directly assigned to the space of regression functions. There are many empirical studies which confirm that GPRMs perform very well compared to other non-linear models (Chen et al., 2007; Yuan et al., 2008). Moreover, GP regression has become a very popular approach for the approximation of time-expensive computer models (Vollert et al., 2019).

A large class of GPRMs automatically provide some information regarding the importance of the individual variables the model is fitted with. This property is known as automatic relevance determination (Rasmussen and Williams, 2005). However, in case of limited training data such approaches perform poorly, resulting in a high generalization error. One possibility to overcome this problem is to add a penalty term, that encourages model fits, where only some of the variables are considered as important. Respective approaches were proposed by Yi et al. (2011) and Wu et al. (2017). A related Bayesian approach was proposed by Linkletter et al. (2006), who used a spike and slab prior (mixture prior between a uniform distribution and a point mass) for the variable selection task. Chen and Wang (2010) proposed a Bayesian approach that uses a random indicator vector to specify which variables contribute to the model.

A possible weakness of GPRMs is that the estimation process sometimes becomes numerically unstable. To address this issue Gu et al. (2018) proposed the concept of robust estimation. In a Bayesian manner, robust estimates can be achieved by assigning so-called reference priors to the GP parameters. These priors are recommended by many authors (Berger et al., 2001; Paulo, 2005; Kazianka and Pilz, 2012; Vollert et al., 2019). Another disadvantage of GPRMs is that they are computationally expensive. Recently, (Datta et al., 2016) introduced so-called nearest neighbor Gaussian processes (NNGPs). They can be considered as highly scalable approximations derived from classical GPs.

In this work we propose a variable selection approach combining NNGPs and reference priors. To the best of our knowledge such a combination has not been proposed before. The variable selection is based on a random set \mathcal{A} used to specify variables that contribute to the model. For the model inference we propose a Metropolis within Gibbs algorithm to sample from the posterior distribution. In particular, the proposed MCMC algorithm makes use of Hamiltonian Monte Carlo (HMC) (Duane et al., 1987). HMC is a family of MCMC methods based on the concept of dynamic systems in physics. Virtual dynamical systems are created by augmenting hyperparameters with momentum variables and in the end a sampling from the distribution of the combined system is performed. In a Bayesian context HMC uses the derivatives of the posterior density to generate efficient proposals. In many applications HMC converges significantly faster than conventional approaches (Duane et al., 1987; Neal, 1997; Rasmussen and Williams, 2005).

2. Methodology

2.1. Nearest Neighbor Gaussian Process

Formally, a stochastic process $\{y(\mathbf{x}); \mathbf{x} \in X \subseteq \mathbb{R}^d\}$ is called a Gaussian Process iff for every finite set of indices $\{\mathbf{x}_1, \dots, \mathbf{x}_k\} \subseteq X$ the random vector $y_{\mathbf{x}_1, \dots, \mathbf{x}_k} = (y(\mathbf{x}_1), \dots, y(\mathbf{x}_k))^T$ is multivariate normally distributed. A GP is fully determined by its mean function $\mu : X \rightarrow \mathbb{R}$, $\mathbf{x} \mapsto \mathbb{E}(y(\mathbf{x}))$ and its covariance function $C : X \times X \rightarrow \mathbb{R}$, $(\mathbf{x}_1, \mathbf{x}_2) \mapsto \text{Cov}(y(\mathbf{x}_1), y(\mathbf{x}_2))$. In the context of nonparametric Bayesian regression GPs are often used as prior distribution for the unknown regression function. Performing inference in such a model reduces to the estimation of some unknown parameters used to define the mean function $\mu(\cdot)$ and the covariance function $C(\cdot, \cdot)$. In the regression context the indices \mathbf{x}_i are also called covariates or predictors. Let $D = \{(\mathbf{x}_i, y(\mathbf{x}_i)) | i = 1, \dots, n\}$ be a finite realization of a GP, i.e., some training data. Further, let $S = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ denote the set of covariate constellations for which observations are available. Then, $\mathbf{C}_S = [C(\mathbf{x}_i, \mathbf{x}_j)]_{\mathbf{x}_i, \mathbf{x}_j \in S}$ denotes the corresponding covariance matrix. Due to the form of the Gaussian density, the model inference is based on the quantities \mathbf{C}_S^{-1} and $|\mathbf{C}_S|$. Since the computation of inverse matrices and also the calculation of determinants is computationally expensive, the application of GP regression models becomes challenging in case that n is large. Recently, Datta et al. (2016) introduced so-called nearest neighbor Gaussian processes as highly scalable approximations derived from classical GP models. Starting point is a zero mean GP, i.e., $\mu(\cdot) \equiv 0$. Then the main idea behind NNGPs is to write

the joint density of $\mathbf{y}_S = (y(\mathbf{x}_1), \dots, y(\mathbf{x}_n))^T$ as a product of conditional densities, using the multiplication theorem, and subsequently replacing the conditioning sets by smaller ones of size at most $m < n$. This is

$$p(\mathbf{y}_S) = p(y(\mathbf{x}_1)) \prod_{i=2}^n p(y(\mathbf{x}_i) | y(\mathbf{x}_{i-1}), \dots, y(\mathbf{x}_1)) \quad (2.1)$$

$$\tilde{p}(\mathbf{y}_S) = \prod_{i=1}^n p(y(\mathbf{x}_i) | y_{N(\mathbf{x}_i)}) \quad (2.2)$$

where $y_{N(\mathbf{x}_i)}$ denotes the vector formed by stacking all the $y(\mathbf{x})$ for which \mathbf{x} lies in the $N(\mathbf{x}_i) \subset S \setminus \{\mathbf{x}_i\}$, $|N(\mathbf{x}_i)| \leq m$, the so-called neighbor set of \mathbf{x}_i . Defining the neighbor set $N(\mathbf{x}_i)$ to be any subset of $\{\mathbf{x}_1, \dots, \mathbf{x}_{i-1}\}$ ensures that Equation 2.2 specifies a valid probability density (Datta et al., 2016). In this work we focus on the specification of Vecchia (1988), who specified $N(\mathbf{x}_i)$ to be the m nearest neighbors of \mathbf{x}_i among $\{\mathbf{x}_1, \dots, \mathbf{x}_{i-1}\}$ with respect to the Euclidean distance. Note that the density $\tilde{p}(\mathbf{y}_S)$ is given by

$$\tilde{p}(\mathbf{y}_S) = \prod_{i=1}^n \mathcal{N}(y(\mathbf{x}_i) | \mathbf{B}_{\mathbf{x}_i} y_{N(\mathbf{x}_i)}, F_{\mathbf{x}_i}) \quad (2.3)$$

where $\mathbf{B}_{\mathbf{x}_i} = \mathbf{C}_{\mathbf{x}_i, N(\mathbf{x}_i)} \mathbf{C}_{N(\mathbf{x}_i)}^{-1}$ and $F_{\mathbf{x}_i} = \mathbf{C}(\mathbf{x}_i, \mathbf{x}_i) - \mathbf{C}_{\mathbf{x}_i, N(\mathbf{x}_i)} \mathbf{C}_{N(\mathbf{x}_i)}^{-1} \mathbf{C}_{N(\mathbf{x}_i), \mathbf{x}_i}$. Equation 2.3 holds due to Equation 2.2 and the fact that for

$$\mathbf{x} = \begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{pmatrix} \sim \mathcal{N} \left(\begin{pmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{pmatrix}, \begin{pmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{pmatrix} \right)$$

$(\mathbf{x}_1 | \mathbf{x}_2 = \mathbf{a}) \sim \mathcal{N}(\bar{\boldsymbol{\mu}}, \bar{\boldsymbol{\Sigma}})$ with $\bar{\boldsymbol{\mu}} = \boldsymbol{\mu}_1 + \boldsymbol{\Sigma}_{12} \boldsymbol{\Sigma}_{22}^{-1} (\mathbf{a} - \boldsymbol{\mu}_2)$ and $\bar{\boldsymbol{\Sigma}} = \boldsymbol{\Sigma}_{11} - \boldsymbol{\Sigma}_{12} \boldsymbol{\Sigma}_{22}^{-1} \boldsymbol{\Sigma}_{21}$.

Following Datta et al. (2016) the density $\tilde{p}(\mathbf{y}_S)$ can be extended to a legitimate stochastic process conform to Komogorov's consistency criteria. Let $\mathbf{u} \in X \setminus S$. Further, let $N(\mathbf{u})$ be the set of m -nearest neighbors of \mathbf{u} in S . For any finite set $U = \{\mathbf{u}_1, \dots, \mathbf{u}_r\}$ with $S \cap U = \emptyset$ we define the nearest neighbor density of \mathbf{y}_U conditional on \mathbf{y}_S as

$$\tilde{p}(\mathbf{y}_U | \mathbf{y}_S) = \prod_{i=1}^r p(y(\mathbf{u}_i) | \mathbf{y}_{N(\mathbf{u}_i)}) \quad (2.4)$$

This conditional density is similar to 2.2 except that all the neighbor sets are subsets of S . This ensures a proper conditional density. Indeed Equation 2.2 and Equation 2.4 are sufficient to describe the joint (nearest neighbor) density of any finite set $V = \{\mathbf{v}_1, \dots, \mathbf{v}_n\} \subseteq X$

$$\tilde{p}(\mathbf{y}_V) = \int \tilde{p}(\mathbf{y}_U | \mathbf{y}_S) \tilde{p}(\mathbf{y}_S) \prod_{\{\mathbf{s}_i \in S \setminus V\}} dy(\mathbf{s}_i) \quad (2.5)$$

where $U = V \setminus S$. If $U = \emptyset$, then Equation 2.4 implies that $\tilde{p}(\mathbf{y}_U | \mathbf{y}_S) = 1$ in Equation 2.5. If $S \setminus V = \emptyset$ the integration in Equation 2.5 is not needed. Moreover, Datta et al. (2016) could show that $\tilde{p}(\mathbf{y}_V)$ is the density of a zero mean GP with covariance function

$$\tilde{\mathbf{C}}(\mathbf{v}_1, \mathbf{v}_2) = \begin{cases} \tilde{\mathbf{C}}_{\mathbf{x}_i, \mathbf{x}_j} & \text{if } \mathbf{v}_1 = \mathbf{x}_i, \mathbf{v}_2 = \mathbf{x}_j \text{ are both in } S \\ \mathbf{B}_{\mathbf{v}_1} \tilde{\mathbf{C}}_{N(\mathbf{v}_1), \mathbf{x}_j} & \text{if } \mathbf{v}_1 \notin S, \mathbf{v}_2 = \mathbf{x}_j \in S \\ \mathbf{B}_{\mathbf{v}_1} \tilde{\mathbf{C}}_{N(\mathbf{v}_1), N(\mathbf{v}_2)} \mathbf{B}_{\mathbf{v}_2}^T + \delta_{(\mathbf{v}_1 = \mathbf{v}_2)} F_{\mathbf{v}_1} & \text{if } \mathbf{v}_1, \mathbf{v}_2 \notin S \end{cases} \quad (2.6)$$

where

$$\tilde{\mathbf{C}}_S = \mathbf{B}_S^{-1} \mathbf{F}_S (\mathbf{B}_S^T)^{-1}, \quad (2.7)$$

with

$$\mathbf{B}_S = \begin{pmatrix} 1 & & & & \\ & 1 & & \mathcal{O} & \\ & & \ddots & & \\ & B_{\mathbf{x}_i, j} & & 1 & \\ & & & & 1 \end{pmatrix}, \quad B_{\mathbf{x}_i, j} = \begin{cases} 1 & i = j \\ -\mathbf{B}_{\mathbf{x}_i}[l] & \mathbf{x}_j = \mathbf{x}_i^l \text{ for some } l, \\ 0 & \text{else} \end{cases} \quad (2.8)$$

$$\mathbf{B}_{\mathbf{v}_i} = \mathbf{C}_{\mathbf{v}_i, N(\mathbf{v}_i)} \mathbf{C}_{N(\mathbf{v}_i)}^{-1}, \quad (2.9)$$

$$\mathbf{F}_S = \text{diag}(F_{\mathbf{x}_1}, \dots, F_{\mathbf{x}_n}), \quad (2.10)$$

$$F_{\mathbf{v}_i} = C_{\mathbf{v}_i} - \mathbf{B}_{\mathbf{v}_i} \mathbf{C}_{N(\mathbf{v}_i), \mathbf{v}_i}, \quad (2.11)$$

$\mathbf{x}_j = \mathbf{x}_i^l$ means that \mathbf{x}_j is the l -th neighbor of \mathbf{x}_i ($N(\mathbf{x}_i) = \{\mathbf{x}_i^1, \mathbf{x}_i^2, \dots, \mathbf{x}_i^l, \dots, \mathbf{x}_i^{N(\mathbf{x}_i)}\}$), $\mathbf{v}_1, \mathbf{v}_2 \in X$, $\tilde{\mathbf{C}}_{A,B}$ denotes submatrices of $\tilde{\mathbf{C}}$ indexed by the elements of the sets A and B , and $\delta_{(\mathbf{v}_1=\mathbf{v}_2)}$ is the Kronecker delta.

2.2. Reference Prior

In terms of robust estimation of the learnable parameters of a GP the application of so-called reference priors has become popular (Gu et al., 2018; Vollert et al., 2019). Paulo (2005) has computed the general form of the reference prior for a broad class of Gaussian processes. Assume that $y(\cdot)$ is a GP on $X \subseteq \mathbb{R}^d$ with mean and covariance given by

$$\mu(\mathbf{x}) = \mathbf{x}^T \boldsymbol{\beta}, \quad C(\mathbf{x}_1, \mathbf{x}_2) = \sigma^2 K(\mathbf{x}_1, \mathbf{x}_2 | \boldsymbol{\xi}), \quad (2.12)$$

where $\boldsymbol{\beta} \in \mathbb{R}^d$, $\boldsymbol{\xi} \in \mathbb{R}_+^r$, and $\sigma^2 > 0$ is the process variance. In order to define the likelihood function corresponding to the observed data D , consider the following additional notations:

$$\mathbf{X} = (\mathbf{x}_1 \cdots \mathbf{x}_n)^T, \quad (2.13)$$

$$\boldsymbol{\eta} = (\sigma^2, \boldsymbol{\beta}^T, \boldsymbol{\xi}^T)^T, \quad (2.14)$$

$$\mathbf{K}_S = [K(\mathbf{x}_i, \mathbf{x}_j | \boldsymbol{\xi})]_{\mathbf{x}_i, \mathbf{x}_j \in S}. \quad (2.15)$$

Then, $\mathbf{y}_S | \boldsymbol{\eta} \sim \mathcal{N}(\mathbf{X}\boldsymbol{\beta}, \sigma^2 \mathbf{K}_S)$, which implies that the likelihood is given by:

$$L(\boldsymbol{\eta} | \mathbf{y}_S) \propto (\sigma^2)^{-\frac{n}{2}} |\mathbf{K}_S|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2\sigma^2} (\mathbf{y}_S - \mathbf{X}\boldsymbol{\beta})^T \mathbf{K}_S^{-1} (\mathbf{y}_S - \mathbf{X}\boldsymbol{\beta}) \right\} \quad (2.16)$$

In order to define the reference prior $(\sigma^2, \boldsymbol{\xi}^T)^T$ is considered as parameter of interest and $\boldsymbol{\beta}$ is considered as nuisance parameter. Moreover, the reference prior distribution $\pi^R(\boldsymbol{\eta})$ is factored as $\pi^R(\boldsymbol{\eta}) = \pi^R(\boldsymbol{\beta} | \sigma^2, \boldsymbol{\xi}) \pi^R(\sigma^2, \boldsymbol{\xi})$. The conditional prior of $\boldsymbol{\beta}$ is specified proportional to 1, $\pi^R(\boldsymbol{\beta} | \sigma^2, \boldsymbol{\xi}) \propto 1$, because this is the Jeffreys-rule prior for the model at hand when σ^2 and $\boldsymbol{\xi}$ are considered to be known. Next $\pi^R(\sigma^2, \boldsymbol{\xi})$ is calculated as the Jeffreys-rule prior for the marginal experiment defined via the integrated likelihood with respect to $\pi^R(\boldsymbol{\beta} | \sigma^2, \boldsymbol{\xi})$. The integrated likelihood is given by:

$$L^I(\sigma^2, \boldsymbol{\xi} | \mathbf{y}_S) = \int_{\mathbb{R}^d} L(\boldsymbol{\eta} | \mathbf{y}_S) \pi^R(\boldsymbol{\beta} | \sigma^2, \boldsymbol{\xi}) d\boldsymbol{\beta} \quad (2.17)$$

$$= \int_{\mathbb{R}^d} L(\boldsymbol{\eta} | \mathbf{y}_S) d\boldsymbol{\beta} \quad (2.18)$$

$$\propto (\sigma^2)^{-\frac{n-d}{2}} |\mathbf{K}_S|^{-\frac{1}{2}} \|\mathbf{X}^T \mathbf{K}_S^{-1} \mathbf{X}\|^{-\frac{1}{2}} \exp \left\{ -\frac{S_{\boldsymbol{\xi}}^2}{2\sigma^2} \right\} \quad (2.19)$$

where $S_{\xi}^2 = \mathbf{y}_S^T \mathbf{Q} \mathbf{y}_S$, $\mathbf{Q} = \mathbf{K}_S^{-1} \mathbf{P}$, and $\mathbf{P} = \mathbf{I} - \mathbf{X}(\mathbf{X}^T \mathbf{K}_S^{-1} \mathbf{X})^{-1} \mathbf{X}^T \mathbf{K}_S^{-1}$. Finally, the reference prior $\pi^R(\boldsymbol{\eta})$ is of the form $\pi^R(\boldsymbol{\eta}) \propto \frac{\pi^R(\boldsymbol{\xi})}{\sigma^2}$ with $\pi^R(\boldsymbol{\xi}) \propto |\tilde{I}_R(\boldsymbol{\xi})|^{\frac{1}{2}}$ where

$$\mathbf{W}_k = \frac{\partial \mathbf{K}_S}{\partial \xi_k} \mathbf{Q} \quad k = 1, \dots, r, \quad (2.20)$$

$$\tilde{I}_R(\boldsymbol{\xi}) = \begin{pmatrix} n-d & \text{tr}(\mathbf{W}_1) & \text{tr}(\mathbf{W}_2) & \cdots & \text{tr}(\mathbf{W}_r) \\ \text{tr}(\mathbf{W}_1) & \text{tr}(\mathbf{W}_1^2) & \text{tr}(\mathbf{W}_1 \mathbf{W}_2) & \cdots & \text{tr}(\mathbf{W}_1 \mathbf{W}_r) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \text{tr}(\mathbf{W}_r) & \text{tr}(\mathbf{W}_1 \mathbf{W}_r) & \text{tr}(\mathbf{W}_2 \mathbf{W}_r) & \cdots & \text{tr}(\mathbf{W}_r^2) \end{pmatrix}, \quad (2.21)$$

see (Paulo, 2005).

2.3. Proposed Variable Selection Model

In this section, the proposed variable selection model is described. To this aim at first the likelihood function is defined. Afterwards, in a Bayesian manner, diverse prior distributions are assigned to the model parameters.

2.3.1. Likelihood Function

Let $y \in \mathbb{R}$ denote a target variable of interest. Further, let $\mathbf{x} \in \mathbb{R}^d$ denote predictors. It is assumed that the relationship

$$y = \mathbf{x}^T \boldsymbol{\beta} + f(\mathbf{x}) + \varepsilon \quad (2.22)$$

holds, where f denotes an unknown function and ε additive noise. Moreover, let \mathcal{A} denote a (random) subset of the index set $\{1, \dots, d\}$, with $|\mathcal{A}| = k > 0$. This set is used to specify the covariates that contribute to the regression model. For given \mathcal{A} we assign a GP prior with zero mean and covariance function \tilde{C} to the sum $z := f + \varepsilon$:

$$z \sim GP(0, \tilde{C}(\cdot, \cdot | \sigma^2, \gamma, \rho, \mathcal{A})), \quad (2.23)$$

with $\sigma^2 > 0$, $\rho > 0$ and $\gamma \in (0, 1)$. In particular, $GP(0, \tilde{C}(\cdot, \cdot | \sigma^2, \gamma, \rho, \mathcal{A}))$ is defined as the NNGP derived from the parent $GP(0, C(\cdot, \cdot | \sigma^2, \gamma, \rho, \mathcal{A}))$, with

$$C(\mathbf{x}, \mathbf{x}' | \sigma^2, \gamma, \rho, \mathcal{A}) = \sigma^2 \underbrace{[\delta_{(\mathbf{x}=\mathbf{x}')} (1-\gamma) + \gamma K(\mathbf{x}, \mathbf{x}' | \rho, \mathcal{A})]}_{=\mathcal{K}(\mathbf{x}, \mathbf{x}' | \gamma, \rho, \mathcal{A})}, \quad (2.24)$$

where

$$\mathcal{K}(\mathbf{x}, \mathbf{x}' | \rho, \mathcal{A}) = \left(1 + \frac{\sqrt{5} d_{\mathcal{A}}}{\rho} + \frac{5 d_{\mathcal{A}}^2}{3 \rho^2} \right) \exp \left(-\frac{\sqrt{5} d_{\mathcal{A}}}{\rho} \right), \quad (2.25)$$

with

$$d_{\mathcal{A}} = \sqrt{\sum_{i \in \mathcal{A}} (x_i - x'_i)^2}. \quad (2.26)$$

Therefore, the parent GP has a Matern- $\frac{5}{2}$ covariance function. Moreover, the Euclidean distance function $d_{\mathcal{A}}$ is based on \mathcal{A} and considers only those predictors with indices in \mathcal{A} . Please note that $d_{\mathcal{A}}$ is also used to determine the neighbor sets required for the computation of the NNGP.

We want to point out that the parent GP results from assigning a zero mean GP with covariance function $\sigma_f^2 K(\mathbf{x}, \mathbf{x}' | \rho, \mathcal{A})$ to f and, moreover, assuming the noise ε to be *i.i.d.* $\mathcal{N}(0, \sigma_\varepsilon^2)$. The reparameterization $\sigma^2 = \sigma_f^2 + \sigma_\varepsilon^2, \gamma = 1 - \frac{\sigma_\varepsilon^2}{\sigma^2}$ in (2.24) leads to:

$$C(\mathbf{x}, \mathbf{x}' | \sigma_f^2, \sigma_\varepsilon^2, \rho, \mathcal{A}) = \sigma_\varepsilon^2 \delta_{(\mathbf{x}=\mathbf{x}')} + \sigma_f^2 K(\mathbf{x}, \mathbf{x}' | \rho, \mathcal{A}) \quad (2.27)$$

$$= \sigma^2 \left[\frac{\sigma_\varepsilon^2}{\sigma^2} \delta_{(\mathbf{x}=\mathbf{x}')} + \frac{\sigma^2 - \sigma_\varepsilon^2}{\sigma^2} K(\mathbf{x}, \mathbf{x}' | \rho, \mathcal{A}) \right] \quad (2.28)$$

$$= \sigma^2 [(1 - \gamma) \delta_{(\mathbf{x}=\mathbf{x}')} + \gamma K(\mathbf{x}, \mathbf{x}' | \rho, \mathcal{A})]. \quad (2.29)$$

Above specifications directly imply that y follows a Gaussian process with mean function $\mu(\mathbf{x}) = \mathbf{x}^T \boldsymbol{\beta}$ and covariance function $\tilde{C}(\mathbf{x}, \mathbf{x}' | \sigma^2, \gamma, \rho, \mathcal{A})$. Let $D = \{(\mathbf{x}_i, y_i) | i = 1, \dots, n\}$ denote some training data and, further, let $S = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ denote the corresponding set of covariate constellations. Consider the following additional notations:

$$\tilde{\mathbf{C}}_S = [\tilde{C}(\mathbf{x}_i, \mathbf{x}_j | \sigma^2, \gamma, \rho, \mathcal{A})]_{\mathbf{x}_i, \mathbf{x}_j \in S}. \quad (2.30)$$

According to Section 2.1 $\tilde{\mathbf{C}}_S$ computes as:

$$\tilde{\mathbf{C}}_S = \mathbf{B}_S^{-1} \mathbf{F}_S (\mathbf{B}_S^T)^{-1} \quad (2.31)$$

with

$$\mathbf{B}_S = \begin{pmatrix} 1 & & & & \\ & 1 & & \mathcal{O} & \\ & & \ddots & & \\ & B_{\mathbf{x}_i, j} & & 1 & \\ & & & & 1 \end{pmatrix}, \quad B_{\mathbf{x}_i, j} = \begin{cases} 1 & i = j \\ -\mathbf{B}_{\mathbf{x}_i}[l] & \mathbf{x}_j = \mathbf{x}_i^l \text{ for some } l \\ 0 & \text{else} \end{cases} \quad (2.32)$$

$$\mathbf{B}_{\mathbf{x}_i} = \mathbf{C}_{\mathbf{x}_i, N(\mathbf{x}_i)} \mathbf{C}_{N(\mathbf{x}_i)}^{-1} = \sigma^2 \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)} (\sigma^2)^{-1} \mathcal{K}_{N(\mathbf{x}_i)}^{-1} \quad (2.33)$$

$$= \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)} \mathcal{K}_{N(\mathbf{x}_i)}^{-1} \quad (\text{independent of } \sigma^2), \quad (2.34)$$

$$\mathbf{F}_S = \text{diag}(F_{\mathbf{x}_1}, \dots, F_{\mathbf{x}_n}), \quad (2.35)$$

$$F_{\mathbf{x}_i} = C_{\mathbf{x}_i} - \mathbf{B}_{\mathbf{x}_i} \mathbf{C}_{N(\mathbf{x}_i), \mathbf{x}_i} \quad (2.36)$$

$$= \sigma^2 \mathcal{K}_{\mathbf{x}_i} - \mathbf{B}_{\mathbf{x}_i} \sigma^2 \mathcal{K}_{N(\mathbf{x}_i), \mathbf{x}_i} \quad (2.37)$$

$$= \sigma^2 \underbrace{[\mathcal{K}_{\mathbf{x}_i} - \mathbf{B}_{\mathbf{x}_i} \mathcal{K}_{N(\mathbf{x}_i), \mathbf{x}_i}]}_{= \mathcal{F}_{\mathbf{x}_i} \quad (\text{independent of } \sigma^2)}. \quad (2.38)$$

Overall one obtains:

$$\tilde{\mathbf{C}}_S = \sigma^2 \underbrace{\mathbf{B}_S^{-1} \text{diag}(\mathcal{F}_{\mathbf{x}_1}, \dots, \mathcal{F}_{\mathbf{x}_n}) (\mathbf{B}_S^T)^{-1}}_{=: \tilde{\mathcal{K}}_S \quad (\text{independent of } \sigma^2, \text{ dependent of } \rho, \gamma, \mathcal{A})} \quad (2.39)$$

$$= \sigma^2 \tilde{\mathcal{K}}_S. \quad (2.40)$$

Thus, $\mathbf{y}_S | \sigma^2, \gamma, \rho, \mathcal{A} \sim \mathcal{N}(\mathbf{X}\boldsymbol{\beta}, \sigma^2 \tilde{\mathcal{K}}_S)$, which implies that the likelihood function is given by:

$$L(\boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} | \mathbf{y}_S) \propto (\sigma^2)^{-\frac{n}{2}} |\tilde{\mathcal{K}}_S|^{-\frac{1}{2}} \exp \left\{ -\frac{1}{2\sigma^2} (\mathbf{y}_S - \mathbf{X}\boldsymbol{\beta})^T \tilde{\mathcal{K}}_S^{-1} (\mathbf{y}_S - \mathbf{X}\boldsymbol{\beta}) \right\}. \quad (2.41)$$

2.3.2. Reference Prior

For given \mathcal{A} we denote the components of $\boldsymbol{\beta}$ with indices in \mathcal{A} as $\boldsymbol{\beta}_{\mathcal{A}}$ and the remaining ones as $\boldsymbol{\beta}_{\mathcal{A}^c}$. We assume that $\boldsymbol{\beta}_{\mathcal{A}^c}$ is equal to zero and, moreover, a uniform prior is assigned to $\boldsymbol{\beta}_{\mathcal{A}}$. The joint prior density with respect to the measure $(\lambda + \delta_0)^d$ is given by

$$p(\boldsymbol{\beta} | \mathcal{A}) = p(\boldsymbol{\beta}_{\mathcal{A}} | \mathcal{A}) p(\boldsymbol{\beta}_{\mathcal{A}^c} | \mathcal{A}) \propto I_{\{\boldsymbol{\beta}_{\mathcal{A}^c} = \mathbf{0}\}}(\boldsymbol{\beta}_{\mathcal{A}^c}) I_{\{\boldsymbol{\beta}_i \neq 0 | \forall i \in \mathcal{A}\}}(\boldsymbol{\beta}_{\mathcal{A}}), \quad (2.42)$$

where λ denotes the Lebesgue measure and δ_0 denotes the Dirac measure concentrated at zero. Thus, the integrated likelihood (compare to Section 2.2) is given by:

$$L^I(\sigma^2, \rho, \gamma, \mathcal{A} | \mathbf{y}_S) = \int_{\mathbb{R}^d} L(\sigma^2, \rho, \gamma, \mathcal{A}, \boldsymbol{\beta} | \mathbf{y}_S) p(\boldsymbol{\beta} | \mathcal{A}) d\lambda(\boldsymbol{\beta}_{\mathcal{A}}) d(\delta_0(\boldsymbol{\beta}_{\mathcal{A}}^C)) \quad (2.43)$$

$$\propto (\sigma^2)^{-\frac{n-|\mathcal{A}|}{2}} |\tilde{\mathcal{K}}_S|^{-\frac{1}{2}} \|\mathbf{X}_{\mathcal{A}}^T \tilde{\mathcal{K}}_S^{-1} \mathbf{X}_{\mathcal{A}}\|^{-\frac{1}{2}} \exp\left\{-\frac{S_{\lambda}^2}{2\sigma^2}\right\} \quad (2.44)$$

where $\boldsymbol{\lambda} = (\rho, \gamma, \mathcal{A})^T$, $S_{\lambda}^2 = \mathbf{y}_S^T \mathbf{Q} \mathbf{y}_S$, $\mathbf{Q} = \tilde{\mathcal{K}}_S^{-1} \mathbf{P}$, $\mathbf{P} = \mathbf{I} - \mathbf{X}_{\mathcal{A}} (\mathbf{X}_{\mathcal{A}}^T \tilde{\mathcal{K}}_S^{-1} \mathbf{X}_{\mathcal{A}})^{-1} \mathbf{X}_{\mathcal{A}}^T \tilde{\mathcal{K}}_S^{-1}$, and $\mathbf{X}_{\mathcal{A}}$ denotes the corresponding to the set \mathcal{A} reduced design matrix. In accordance with Section 2.2 we use the prior

$$p(\sigma^2, \rho, \gamma) \propto \frac{|\tilde{I}_R(\rho, \gamma)|^{\frac{1}{2}}}{\sigma^2} \quad (2.45)$$

where

$$\mathbf{W}_{\rho} = \frac{\partial \tilde{\mathcal{K}}_S}{\partial \xi_k} \mathbf{Q} \quad (2.46)$$

$$\mathbf{W}_{\gamma} = \frac{\partial \tilde{\mathcal{K}}_S}{\partial \gamma} \mathbf{Q} \quad (2.47)$$

$$\tilde{I}_R(\boldsymbol{\lambda}) = \begin{pmatrix} n - |\mathcal{A}| & \text{tr}(\mathbf{W}_{\rho}) & \text{tr}(\mathbf{W}_{\gamma}) \\ \text{tr}(\mathbf{W}_{\rho}) & \text{tr}(\mathbf{W}_{\rho}^2) & \text{tr}(\mathbf{W}_{\rho} \mathbf{W}_{\gamma}) \\ \text{tr}(\mathbf{W}_{\gamma}) & \text{tr}(\mathbf{W}_{\rho} \mathbf{W}_{\gamma}) & \text{tr}(\mathbf{W}_{\gamma}^2) \end{pmatrix}. \quad (2.48)$$

The computation of the derivatives presented in Equation 2.46 and Equation 2.47 is given in Appendix A.

2.3.3. Prior for Random Set

Inspired by Posch et al. (2020) we propose the following prior for \mathcal{A} :

$$p(\mathcal{A} = \{\alpha_1, \dots, \alpha_k\}) \propto (p_{\alpha_1} + \dots + p_{\alpha_k}) \frac{1}{k} \tilde{p}(k), \quad (2.49)$$

where

- (i) $\{p_{\alpha_1}, \dots, p_{\alpha_k}\} \subseteq \{p_1, \dots, p_d\}$ with $\sum_{i=1}^d p_i = 1$ and $p_i \geq 0$ for $i = 1, \dots, d$ and
- (ii) $\tilde{p}: \{1, \dots, d\} \rightarrow \mathbb{R}_0^+$.

The mapping \tilde{p} can be arbitrarily chosen and should represent the a priori belief in the number of predictors contributing to the model. On the other hand the parameters p_1, \dots, p_d are used to represent the a priori belief in the importance of the predictors x_1, \dots, x_d to the model.

2.4. Model Inference

Clearly, the joint posterior $p(\boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} | \mathbf{y}_S, \mathbf{X})$ is analytically intractable. Therefore, we propose a Metropolis within Gibbs algorithm to simulate from this distribution. The algorithm produces a Markov Chain with the true posterior as limiting distribution. In particular the algorithm includes 3 main steps:

1. Sample from the conditional posterior $p(\boldsymbol{\beta}_{t+1}, \mathcal{A}_{t+1} | \mathbf{y}_S, \mathbf{X}, \sigma_t^2, \gamma_t, \rho_t)$ using a classical Metropolis Hastings (MH) algorithm.
2. Sample from the conditional posterior $p(\sigma_{t+1}^2 | \mathbf{y}_S, \mathbf{X}, \gamma_t, \rho_t, \boldsymbol{\beta}_{t+1}, \mathcal{A}_{t+1})$.
3. Sample from the conditional posterior $p(\gamma_{t+1}, \rho_{t+1} | \mathbf{y}_S, \mathbf{X}, \boldsymbol{\beta}_{t+1}, \mathcal{A}_{t+1}, \sigma_{t+1}^2)$ using Hamiltonian MCMC (HMC).

In order to run the procedure one has to define an initial sample $\boldsymbol{\beta}_0, \sigma_0^2, \gamma_0, \rho_0, \mathcal{A}_0$ that lies in the support of $p(\boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} | \mathbf{y}_S, \mathbf{X})$. Additionally, some tuning parameters for the proposal distribution of \mathcal{A} are required and also the step size ε and the number of steps L for the leap frog method, that is used within the HMC step. In the following sections details regarding the individual steps are presented.

2.4.1. Details Step 1:

Following Posch et al. (2020) to specify the proposal for the random set \mathcal{A} at first a Bernoulli distributed random variable c_h is introduced

$$c_h \sim \text{Bernoulli}(p_h), \quad (2.50)$$

where $p_h \in [0, 1]$ can be considered as a tuning parameter. The event $c_h = 1$ means that the model size (number of predictors used) changes, i.e. $k_{t+1} \neq k_t$. Moreover, a random variable α with support on the index set $\{1, \dots, d\}$ is introduced in order to describe the model transition probabilities in case of changing model size. A realization of the conditional random variable $\alpha|\mathcal{A}_t$ corresponds to the index of a predictor which is going to be added to or removed from the model. Transitions between models, which differ by two or more parameters, are not allowed. The probability mass function of $\alpha|\mathcal{A}_t$ is defined as

$$q(\alpha|\mathcal{A}_t) = \mathbb{I}_{\{1, \dots, d\}}(\alpha) \begin{cases} \tilde{p}_\alpha & \text{if } k_t > 1, \alpha \notin \mathcal{A}_t \\ \frac{\sum_{i \in \mathcal{A}_t} \tilde{p}_i}{\tilde{p}_\alpha \sum_{i \in \mathcal{A}_t} 1/\tilde{p}_i} & \text{if } k_t > 1, \alpha \in \mathcal{A}_t \\ 0 & \text{if } k_t = 1, \alpha \in \mathcal{A}_t \\ \frac{\tilde{p}_\alpha}{\sum_{i \in \{1, \dots, p\} \setminus \mathcal{A}_t} \tilde{p}_i} & \text{else} \end{cases}, \quad (2.51)$$

where the parameters $\tilde{p}_1, \dots, \tilde{p}_d$ are greater or equal to zero, sum up to one, and can but must not be identical to the parameters p_1, \dots, p_d already used in the prior $p(\mathcal{A})$. Using Equation 2.50 and Equation 2.51 the proposal distribution $q(\mathcal{A}_{t+1}|\mathcal{A}_t)$ is finally defined by:

$$q(\mathcal{A}_{t+1}|\mathcal{A}_t, c_h = 1) = q([\mathcal{A}_{t+1} \setminus \mathcal{A}_t] \cup [\mathcal{A}_t \setminus \mathcal{A}_{t+1}]|\mathcal{A}_t) \mathbb{I}_{\{1\}}(|k_{t+1} - k_t|), \quad (2.52)$$

$$q(\mathcal{A}_{t+1}|\mathcal{A}_t, c_h = 0) = \begin{cases} 1 & \text{if } \mathcal{A}_{t+1} = \mathcal{A}_t \\ 0 & \text{else} \end{cases}, \quad (2.53)$$

$$q(\mathcal{A}_{t+1}|\mathcal{A}_t) = \sum_{c_h \in \{0,1\}} q(\mathcal{A}_{t+1}, c_h|\mathcal{A}_t) \quad (2.54)$$

$$= \sum_{c_h \in \{0,1\}} q(\mathcal{A}_{t+1}|c_h, \mathcal{A}_t) q(c_h). \quad (2.55)$$

Simple algebra shows that:

$$\boldsymbol{\beta}_{t+1, \mathcal{A}_{t+1}} | \mathbf{y}_S, \mathbf{X}, \sigma_t^2, \gamma_t, \rho_t, \mathcal{A}_{t+1} \sim \mathcal{N}(\boldsymbol{\mu} = \boldsymbol{\Sigma} \mathbf{X}_{\mathcal{A}_{t+1}}^T \tilde{\mathbf{C}}_S^{-1} \mathbf{y}_S, \boldsymbol{\Sigma} = (\mathbf{X}_{\mathcal{A}_{t+1}}^T \tilde{\mathbf{C}}_S^{-1} \mathbf{X}_{\mathcal{A}_{t+1}})^{-1}), \quad (2.56)$$

$$\boldsymbol{\beta}_{t+1, \mathcal{A}_{t+1}^c} | \mathbf{y}_S, \mathbf{X}, \sigma_t^2, \gamma_t, \rho_t, \mathcal{A}_{t+1} \sim \delta_0. \quad (2.57)$$

We use the conditional posterior 2.56-2.57 as proposal for $\boldsymbol{\beta}_{t+1}$. The Hastings ratio is then given by:

$$r = \frac{L(\boldsymbol{\beta}_{t+1}, \sigma_t^2, \gamma_t, \rho_t, \mathcal{A}_{t+1} | \mathbf{y}_S) p(\boldsymbol{\beta}_{t+1} | \mathcal{A}_{t+1}) p(\mathcal{A}_{t+1}) q(\boldsymbol{\beta}_t | \sigma_t^2, \gamma_t, \rho_t, \mathcal{A}_t) q(\mathcal{A}_t | \mathcal{A}_{t+1})}{L(\boldsymbol{\beta}_t, \sigma_t^2, \gamma_t, \rho_t, \mathcal{A}_t | \mathbf{y}_S) p(\boldsymbol{\beta}_t | \mathcal{A}_t) p(\mathcal{A}_t) q(\boldsymbol{\beta}_{t+1} | \sigma_t^2, \gamma_t, \rho_t, \mathcal{A}_{t+1}) q(\mathcal{A}_{t+1} | \mathcal{A}_t)}. \quad (2.58)$$

2.4.2. Details Step 2:

The conditional posterior $p(\sigma_{t+1}^2 | \mathbf{y}_S, \mathbf{X}, \gamma_t, \rho_t, \boldsymbol{\beta}_{t+1}, \mathcal{A}_{t+1})$ can be computed in closed analytical form. Simple algebra shows that it is given by:

$$\sigma_{t+1}^2 | \mathbf{y}_S, \mathbf{X}, \gamma_t, \rho_t, \boldsymbol{\beta}_{t+1}, \mathcal{A}_{t+1} \sim \text{IG}(\alpha = \frac{n}{2}, \beta = \frac{1}{2} (\mathbf{y}_S - \mathbf{X}\boldsymbol{\beta})^T \tilde{\boldsymbol{\mathcal{K}}}_S^{-1} (\mathbf{y}_S - \mathbf{X}\boldsymbol{\beta})). \quad (2.59)$$

2.4.3. Details Step 3:

For the HMC algorithm we reparameterize the posterior $p(\gamma_{t+1}, \rho_{t+1} | \mathbf{y}_S, \mathbf{X}, \boldsymbol{\beta}_{t+1}, \mathcal{A}_{t+1}, \sigma_{t+1}^2)$, such that the support of the reparameterized distribution is unbounded. This can be easily done using Jacobi transformation:

$$\gamma_{t+1} = \frac{1}{1 + \exp(-\tilde{\gamma}_{t+1})} \text{ with } \tilde{\gamma}_{t+1} \in \mathbb{R}, \quad (2.60)$$

$$\rho_{t+1} = \log(1 + \exp(\tilde{\rho}_{t+1})) \text{ with } \tilde{\rho}_{t+1} \in \mathbb{R}, \quad (2.61)$$

$$p(\tilde{\gamma}_{t+1}, \tilde{\rho}_{t+1} | \mathbf{y}_S, \mathbf{X}, \boldsymbol{\beta}_{t+1}, \mathcal{A}_{t+1}, \sigma_{t+1}^2) \propto L(\boldsymbol{\beta}_{t+1}, \sigma_{t+1}^2, \gamma_{t+1}, \rho_{t+1}, \mathcal{A}_{t+1} | \mathbf{y}_S) p(\rho_{t+1}, \gamma_{t+1}) \frac{\exp(\tilde{\rho}_{t+1} - \tilde{\gamma}_{t+1})}{(1 + \exp(-\tilde{\gamma}_{t+1})^2)(1 + \exp(\tilde{\rho}_{t+1}))} \quad (2.62)$$

Next we specify the so-called potential energy function:

$$E(\tilde{\rho}_{t+1}, \tilde{\gamma}_{t+1}) = -\log(p(\tilde{\gamma}_{t+1}, \tilde{\rho}_{t+1} | \mathbf{y}_S, \mathbf{X}, \boldsymbol{\beta}_{t+1}, \mathcal{A}_{t+1}, \sigma_{t+1}^2)) \quad (2.63)$$

Due to Equation 2.62 $E(\cdot|\cdot)$ is known up to an additive constant, that vanishes if derivatives are computed. Moreover, the kinetic energy is defined as the negative logarithm of the density of a zero mean normal distribution with a diagonal covariance matrix:

$$K(\mathbf{v} = (v_{\tilde{\rho}}, v_{\tilde{\gamma}})^T) = c + \frac{1}{2} \left(\frac{v_{\tilde{\rho}}^2}{m_{\tilde{\rho}}} + \frac{v_{\tilde{\gamma}}^2}{m_{\tilde{\gamma}}} \right). \quad (2.64)$$

The parameters $m_{\tilde{\rho}}$ and $m_{\tilde{\gamma}}$ are tuning parameters of the algorithm and, further, c denotes an additive constant. Consequently, the partial derivatives of the kinetic energy function are given by:

$$\frac{\partial K(\mathbf{v})}{\partial v_{\tilde{\rho}}} = \frac{v_{\tilde{\rho}}}{m_{\tilde{\rho}}}, \quad \frac{\partial K(\mathbf{v})}{\partial v_{\tilde{\gamma}}} = \frac{v_{\tilde{\gamma}}}{m_{\tilde{\gamma}}}. \quad (2.65)$$

With above considerations the HMC algorithm proposes a sample from the conditional posterior by applying the following procedure:

1. Draw a sample $\mathbf{v} = (v_{\tilde{\rho}}, v_{\tilde{\gamma}})^T$ from $\mathcal{N}(\mathbf{0}, \text{diag}(m_{\tilde{\rho}}, m_{\tilde{\gamma}}))$.
2. Use the leap-frog method to simulate $\tilde{\rho}^*, \tilde{\gamma}^*, v_{\tilde{\rho}}^*, v_{\tilde{\gamma}}^*$.
 - (i) Set $\mathbf{v}^* = \mathbf{v}, \tilde{\rho}^* = \tilde{\rho}_t, \tilde{\gamma}^* = \tilde{\gamma}_t$.
 - (ii) For $j = 1, \dots, L$:
 - (a) $v_{\tilde{\rho}}^* = v_{\tilde{\rho}}^* - \frac{\varepsilon}{2} \frac{\partial E}{\partial \tilde{\rho}^*}(\tilde{\rho}^*, \tilde{\gamma}^*), v_{\tilde{\gamma}}^* = v_{\tilde{\gamma}}^* - \frac{\varepsilon}{2} \frac{\partial E}{\partial \tilde{\gamma}^*}(\tilde{\rho}^*, \tilde{\gamma}^*)$
 - (b) $\tilde{\rho}^* = \tilde{\rho}^* + \varepsilon \frac{v_{\tilde{\rho}}^*}{m_{\tilde{\rho}}}, \tilde{\gamma}^* = \tilde{\gamma}^* + \varepsilon \frac{v_{\tilde{\gamma}}^*}{m_{\tilde{\gamma}}}$
 - (c) $v_{\tilde{\rho}}^* = v_{\tilde{\rho}}^* - \frac{\varepsilon}{2} \frac{\partial E}{\partial \tilde{\rho}^*}(\tilde{\rho}^*, \tilde{\gamma}^*), v_{\tilde{\gamma}}^* = v_{\tilde{\gamma}}^* - \frac{\varepsilon}{2} \frac{\partial E}{\partial \tilde{\gamma}^*}(\tilde{\rho}^*, \tilde{\gamma}^*)$
3. Calculate the Hastings ratio $r = \exp[K(\mathbf{v}) + E(\tilde{\rho}_t, \tilde{\gamma}_t) - K(\mathbf{v}^*) - E(\tilde{\rho}^*, \tilde{\gamma}^*)]$.
4. With probability $\min(r, 1)$ set $\tilde{\rho}_{t+1} = \tilde{\rho}^*, \tilde{\gamma}_{t+1} = \tilde{\gamma}^*$. Otherwise set $\tilde{\rho}_{t+1} = \tilde{\rho}_t, \tilde{\gamma}_{t+1} = \tilde{\gamma}_t$.

2.4.4. Tuning Parameters

This section provides some insights regarding the tuning parameter specifications used for the evaluation of our approach (Section 3). The tuning parameter p_h is set to the value 0.6. Thus, the proposal for \mathcal{A} slightly prefers to modify \mathcal{A} over keeping it unchanged. Moreover, the parameters $\tilde{p}_1, \dots, \tilde{p}_d$ are all specified as $1/d$. Consequently, the proposal does not favor special predictors. The \tilde{p}_i are specified like this, since there is no a priori knowledge available and a specification using empirical investigations regarding the

importance of the individual predictors is not necessary. The MCMC chains converge without informative values for the \hat{p}_i after a reasonable amount of MCMC iterations (a few thousand). For simplicity, nearly all tuning parameters of the HMC step are fixed and set to the values $m_{\hat{p}} = m_{\hat{\gamma}} = 1, L = 2$. The only exception is the step size ε of the leap-frog method. This parameter turns out to be the most influential one and values between 0.15 and 0.5 are chosen. The actual specification is selected in such a way (per trial and error) that the MCMC chains show a good mixing.

2.5. Prediction

Let $\mathbf{X}^* = (\mathbf{x}_1^* \cdots \mathbf{x}_v^*)^T$ be the design matrix corresponding to some predictor specifications for which predictions $\hat{\mathbf{y}}^*$ are required. Note that \mathbf{y}_S and \mathbf{y}^* are joint normally distributed:

$$(\mathbf{y}^*, \mathbf{y}_S)^T | \mathbf{X}^*, \mathbf{X}, \boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} \sim \mathcal{N} \left(\begin{pmatrix} \mathbf{X}^* \boldsymbol{\beta} \\ \mathbf{X} \boldsymbol{\beta} \end{pmatrix}, \begin{pmatrix} \tilde{\mathbf{C}}_{\mathbf{X}^*} & \tilde{\mathbf{C}}_{\mathbf{X}^*, \mathbf{S}} \\ \tilde{\mathbf{C}}_{\mathbf{X}^*, \mathbf{S}} & \tilde{\mathbf{C}}_{\mathbf{S}} \end{pmatrix} \right). \quad (2.66)$$

This directly implies:

$$\mathbf{y}^* | \mathbf{y}_S, \mathbf{X}^*, \mathbf{X}, \boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} \sim \mathcal{N}(\mathbf{X}^* \boldsymbol{\beta} + \tilde{\mathbf{C}}_{\mathbf{X}^*, \mathbf{S}} \tilde{\mathbf{C}}_{\mathbf{S}}^{-1} (\mathbf{y}_S - \mathbf{X} \boldsymbol{\beta}), \tilde{\mathbf{C}}_{\mathbf{X}^*} - \tilde{\mathbf{C}}_{\mathbf{X}^*, \mathbf{S}} \tilde{\mathbf{C}}_{\mathbf{S}}^{-1} \tilde{\mathbf{C}}_{\mathbf{X}^*, \mathbf{S}}) \quad (2.67)$$

We use the expected value of the posterior predictive distribution as prediction. Using Monte Carlo Integration this quantity can easily be approximated:

$$\hat{\mathbf{y}}^* = \int \mathbf{y}^* p(\mathbf{y}^* | \mathbf{y}_S, \mathbf{X}^*, \mathbf{X}) d\mathbf{y}^* \quad (2.68)$$

$$= \int \mathbf{y}^* \left\{ \int \int \int \int \int p(\mathbf{y}^* | \mathbf{y}_S, \mathbf{X}^*, \mathbf{X}, \boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A}) p(\boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} | \mathbf{y}_S, \mathbf{X}) d\sigma^2 d\gamma d\rho \#(d\mathcal{A}) (\lambda + \delta_0)^d(d\boldsymbol{\beta}) \right\} d\mathbf{y}^* \quad (2.69)$$

$$= \int \int \int \int \int \left\{ \int \mathbf{y}^* p(\mathbf{y}^* | \mathbf{y}_S, \mathbf{X}^*, \mathbf{X}, \boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A}) d\mathbf{y}^* \right\} p(\boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} | \mathbf{y}_S, \mathbf{X}) d\sigma^2 d\gamma d\rho \#(d\mathcal{A}) (\lambda + \delta_0)^d(d\boldsymbol{\beta}) \quad (2.70)$$

$$\stackrel{2.67}{=} \int \int \int \int \int \left\{ \mathbf{X}^* \boldsymbol{\beta} + \tilde{\mathbf{C}}_{\mathbf{X}^*, \mathbf{S}} \tilde{\mathbf{C}}_{\mathbf{S}}^{-1} (\mathbf{y}_S - \mathbf{X} \boldsymbol{\beta}) \right\} p(\boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} | \mathbf{y}_S, \mathbf{X}) d\sigma^2 d\gamma d\rho \#(d\mathcal{A}) (\lambda + \delta_0)^d(d\boldsymbol{\beta}) \quad (2.71)$$

$$\approx \frac{1}{N} \sum_{i=1}^N \mathbf{X}^* \boldsymbol{\beta}^{(i)} + \tilde{\mathbf{C}}_{\mathbf{X}^*, \mathbf{S}}^{(i)} \tilde{\mathbf{C}}_{\mathbf{S}}^{(i)-1} (\mathbf{y}_S - \mathbf{X} \boldsymbol{\beta}^{(i)}), \quad (2.72)$$

where $\#$ denotes the counting measure. The (i) in Equation 2.72 is used to outline that each of the individual summands is computed for an *iid* sample from the posterior $p(\boldsymbol{\beta}, \sigma^2, \gamma, \rho, \mathcal{A} | \mathbf{y}_S, \mathbf{X})$.

3. Evaluation

We have implemented the proposed model using the package `Rcpp` (Eddelbuettel and François, 2011), which allows for a seamless R and C++ integration. Additionally, we have used the package `RcppArmadillo` (Eddelbuettel and Sanderson, 2014) in order to accelerate R with high-performance linear algebra. In this section, we evaluate and compare the predictive performance of our approach (VRNNGP) with the accuracy of some other well-established feature selection methods. The methods under comparison are:

- **Lasso.** The least absolute shrinkage and selection operator (Lasso) (Tibshirani, 1996) penalizes the absolute values of the regression coefficients in the classical linear model in order to obtain a sparse solution. The R package `glmnet` (Friedman et al., 2010) implements the lasso. Cross-validation is used to determine the penalization strength.

- **Adaptive lasso (ALasso)**. The adaptive Lasso (Zou, 2006) is an extension of the classical lasso that allows for individual penalization strengths regarding the regression coefficients. In this comparison study the individual penalization factors are defined by multiplying a global factor with the inverse numbers of the absolute values of the optimized regression coefficients from a ridge-penalized model. Again, the package `glmnet` is used for this task.
- **Bayes linear selection (BLS)**. Recently, Posch et al. (2020) proposed a Bayesian variable selection approach for linear regression models. Since our approach shares some common features (e.g. the prior for the random set \mathcal{A}) we also include their method in our evaluation study.
- **Random forest (RF)**. A combination of tree predictors proposed by Breiman (2001). The method is implemented in the R package `randomForest` (Liaw and Wiener, 2002).
- **Robust Gaussian process (RGP)**. The R package `RobustGaSP` (Gu et al., 2020) implements robust Gaussian process regression models using reference priors and separable (anisotropic) covariance functions. We use this package with the default specifications (i.e. a Matern 5/2 kernel etc.), except that we force the method to estimate a nugget term. Note, that GP models with anisotropic kernels allow for an automatic relevance determination (Rasmussen and Williams, 2005). To this aim, the predictors have to be scaled prior to model inference.
- **Maximum likelihood Gaussian process (MLGP)**. The R package `mlegp` (Dancik and Dorman, 2008) implements Gaussian Process regression using maximum likelihood estimates and an anisotropic covariance function. We use this package with the default settings for the model inference, except that we specify the mean function to be non-constant.

The accuracy measures used for the performance comparison are the mean squared error (MSE) and the mean absolute deviation (MAD). Accuracies are always computed based on testing datasets that do not share common observations with the training data. Finally, it should be mentioned that the number of nearest neighbors is specified as $m = 10$ in all subsequent experiments. It turned out that 10 neighbors are sufficient to obtain promising results.

3.1. Pepelyshev Function

In this section, the Pepelyshev function (Dette and Pepelyshev, 2010), a commonly used test function in computer experiments, is considered. In this field it has become common to use a GP in order to model the output of a complex system (Vollert et al., 2019). The highly curved Pepelyshev function is given by

$$f(\mathbf{x}) = 4(x_1 - 2 + 8x_2 - 8x_2^2)^2 + (3 - 4x_2)^2 + 16\sqrt{x_3 + 1}(2x_3 - 1)^2, \quad (3.1)$$

for $x_1, x_2, x_3 \in [0, 1]$. To make the approximation problem more challenging and also in order to obtain a true variable selection task, we extend Function 3.1 by 17 inert variables $x_4, \dots, x_{20} \in [0, 1]$.

The evaluation of our approach is based on a training data set with sample size 31 and a testing dataset of size 100. Both datasets are constructed by generating a maxmin latin hypercube design (LHD), respectively. Maxmin LHDs are standard designs for computer experiments (Santner et al., 2003). They can be easily constructed by using the R package `lhs` (Carnell, 2020). Prior to the analysis the training and the testing dataset are scaled. In Table 1 the accuracy of the methods considered in this study is shown. It can be seen that VRNNGP outperforms the other ones. Reason for the superior performance is that the other methods fail to identify the actual variables x_1, x_2, x_3 , i.e. they also use the inert variables. However, it should be mentioned that also VRNNGP is unable to identify x_1, x_2, x_3 . In this approach x_2, x_3 have a posterior probability of 1 to contribute to the models, while all other predictors have a probability near zero.

Finally, we want to illustrate that the proposed Metropolis within Gibbs algorithm (Section 2.4) converges fast. To this aim the Figures 1-4 show the Markov chains of some representative model parameters. It can be seen that the algorithm converges after about 1000 iterations.

Table 1

Accuracy of individual approaches for the Pepelyshev function with inert variables. The proposed approach performs best.

Method	MSE	MAD
VRNNGP	0.1949179	0.3148653
MLGP	1.999522	1.176881
RGP	1.483502	1.01951
RF	0.7351364	0.7156615
Lasso	1.003959	0.8617405
ALasso	1.63588	1.065925
BLS	1.30754	0.972762

3.2. Sine Function

In this simulation study data are generated from the model

$$y = \sin(x_3) + \sin(5x_4) + \varepsilon, \quad (3.2)$$

$$\varepsilon \sim \mathcal{N}(0, 0.0025), \quad (3.3)$$

$$(x_1, \dots, x_{20})^T \sim \mathcal{N}(\mathbf{0}, \mathbf{\Sigma}), \quad (3.4)$$

$$\text{diag}(\mathbf{\Sigma}) = \mathbf{1}, \quad (3.5)$$

$$\Sigma_{ij} = \begin{cases} 0.4 & \text{if } (i, j) \in \{(3, 13), (13, 3)\}, \\ 0.3 & \text{if } (i, j) \in \{(3, 12), (12, 3)\}, \\ 0.4 & \text{if } (i, j) \in \{(4, 14), (14, 4)\}, \\ 0.3 & \text{if } (i, j) \in \{(4, 15), (15, 4)\}, \\ 0 & \text{else,} \end{cases} \quad (3.6)$$

where $x_1, x_2, x_5, \dots, x_{20}$ are inert variables. In particular, 100 samples are simulated from model 3.2-3.6 to obtain a dataset, that is subsequently zero centered and scaled. The performance evaluation is based on a 5-fold cross validation. In Table 2 the mean MSEs and the mean MADs of the comparison methods is given. VRNNGP performs significantly better than the other methods. Indeed, this method identifies the active variables x_3, x_4 in four of the five considered train/tests splits, while the other methods are unable to do so.

Table 2

Mean accuracy of individual approaches for the simulation model 3.2-3.6. The proposed approach performs best.

Method	MSE	MAD
VRNNGP	0.3580486	0.4316482
MLGP	1.219693	0.9032923
RGP	0.7396772	0.7100051
RF	0.7706562	0.7031102
Lasso	0.8173408	0.7418092
ALasso	0.8286106	0.7483805
BLS	0.8528274	0.7460358

3.3. APC data

In this section, real-world data originating from the so-called semiconductor frontend production is considered (Pleschberger, 2021). In this production chain, a wafer, i.e. a slice based on semiconductor material, goes through hundreds of process steps resulting in thousands of semiconductor devices on the wafer. During each process step, several sensors mounted on the process equipment record physical parameters like gas flow, voltage, temperature, etc. resulting in so-called time-dependent trace data. From this trace data,




Fig. 1. Markov chain of the cardinality of the random set \mathcal{A} .




Fig. 2. Markov chain of the covariance parameter γ .




Fig. 3. Markov chain of the covariance parameter ρ .




Fig. 4. Markov chain of the covariance parameter σ^2 .

so-called key numbers are extracted, which are simple statistical quantities like mean, standard deviation or maximum used to monitor the process stability. After frontend production, each device on the wafer is fastidiously tested in order to meet the customer’s requirements resulting in the so-called wafer test data. These are analog test procedures measuring quantities like current, voltage or resistance. Now, if certain wafer test parameters are conspicuous and lead to potentially high failures, the root cause must be found in frontend production. The following analysis is intended to identify important influencing variables in order to identify processes in frontend production causing the variation at the end. The selected and prepared dataset includes 228 observations from 77 possible predictors (key numbers) extracted from trace data of two crucial process steps. The target variable is an analog measurement from the wafer test data describing the wafer quality. The higher the measured value the worse the quality. Prior to the analysis we have scaled the predictors and zero centered the target variable. Moreover, the complete dataset was randomly partitioned into a training and a testing dataset. Both datasets are equal sized.

In Table 3 the accuracy of the methods considered in this study is shown. VRNNGP returns the best results followed by RF. MLGP fails completely, while the robust version RGP performs slightly worse than the linear approaches. In the Figures 5-7 the importance of the individual predictors obtained from the methods VRNNGP, RF, and BLS is visualized. For VRNNGP and BLS the importance of a predictor is the relative frequency how often its index lies in the posterior samples from \mathcal{A} . This can be interpreted as the posterior probability of the predictor being included in the model. For RF the visualized importance is the mean decrease in accuracy if a particular variable is removed. This quantity can be easily computed using the `importance()` function included in the R package `randomForest`. It can be seen that the importance values differ from method to method. However, the two most important variables in NNGP (variable 34, variable 41) are also important in RF. In BLS at least the variable 41 is considered as important. We assume that the influence of the variable 34 is non-linear, such that a linear model fails to detect this influence. Since the variables 34 and 41 appear to be important across models we train the two models RGP and RF again, but with a reduced dataset including only these two variables. In Table 4 the respective accuracies are presented. Both models perform comparable, but significantly better than before. In particular, the reduction to the variables 34 and 41 led to the best results overall. The Figures show the functions learned by RGP and RF on basis of the reduced dataset. Both models agree on parameter specifications with a poor production quality. Finally, in Figure 10 the true production quality of the testing dataset (y_{Test}) is plotted vs the predicted quality from the RGP model with the reduced dataset. It can be observed, that all of the really poor production results ($y_{\text{Test}} > 0.5$) can be identified. This shows, that the variables selected by NNGP are useful for this application.

Table 3
Accuracy of individual approaches on the APC data. The proposed approach performs best.

Method	MSE	MAD
VRNNGP	0.2389607	0.4197644
MLGP	32.49251	1.44142
RGP	0.3108199	0.4695633
RF	0.2687667	0.4441394
Lasso	0.2843513	0.4568827
ALasso	0.2903017	0.4529153
BLS	0.2846291	0.4471566

Table 4
Accuracy of RGP and RF on reduced dataset. Only the variables 34 and 41 are considered.

Method	MSE	MAD
RGP	0.2053222	0.3692166
RF	0.2022327	0.3539577




Fig. 5. Variable Importance obtained from VRNNGP.




Fig. 6. Variable Importance obtained from RF.




Fig. 7. Variable Importance obtained from BLS.




Fig. 8. Model fit obtained from RGP.




Fig. 9. Model fit obtained from RF.




Fig. 10. True production quality of the testing dataset (y_{Test}) vs predicted quality. Used model: RGP on the reduced dataset.

3.4. Body Fat Data

In this section, the popular body fat dataset (Johnson, 1996) available in the R package `mplot` (Tarr et al., 2018) is considered. The dataset includes 128 observations on 14 variables. The target variable is the body fat percentage of a man and the predictors are age, weight, height, and ten body circumference measurements. A 5-fold cross validation is performed to evaluate and compare the performance of VRNNGP with other methods. Prior to the analysis that the whole dataset was zero centered and scaled. In Table 5 the mean MSEs and the mean MADs of the comparison methods is given. VRNNGP achieves the best results. In the Figures 11-13 the importance of the individual predictors (compare to Section 3.3) obtained from the methods VRNNGP, RF, and BLS is visualized. While all of these methods agree that the sixth predictor (abdomen circumference) is the most influential, there are some disagreements regarding the order of importance of the other predictors.

Table 5

Mean accuracy of individual approaches for the body fat dataset. The proposed approach performs best.

Method	MSE	MAD
VRNNGP	0.2762897	0.4211418
MLGP	0.3657517	0.4865936
RGP	0.308341	0.4434407
RF	0.3035638	0.4458874
Lasso	0.3113187	0.4484568
ALasso	0.314493	0.4521927
BLS	0.3014364	0.4356817




Fig. 11. Variable Importance obtained from VRNNGP.




Fig. 12. Variable Importance obtained from RF.




Fig. 13. Variable Importance obtained from BLS.

3.5. Prior Specifications

In the Sections 3.1-3.4 the proposed model is evaluated based on different datasets. The good performance is not least attributable to the fact that valuable prior information regarding the random set \mathcal{A} can be formulated (see Section 2.3.3). Now we provide the individual prior specifications used in the preceding evaluations. All evaluations share the definition of the a priori importance $p_1 = \dots = p_d = 1/d$ of the predictors. No predictor is preferred over another. However, the specification of the mapping $\tilde{p}(\cdot)$, that represents the a priori belief regarding the number of predictors contributing to the model, differs among the evaluation studies. In Section 3.1, in Section 3.2, and in Section 3.4 the mapping is defined as $\tilde{p}(k) = 1/k \forall k \in \{1, \dots, d\}$. Thus, smaller models are quite intuitively preferred. In Section 3.3 the number of potential predictors is larger than in the other evaluation studies. Therefore, $\tilde{p}(\cdot)$ is specified using the recommendations from Posch et al. (2020), who proposed the prior for \mathcal{A} . In particular, $\tilde{p}(\cdot)$ is defined to be proportional to the third power of the probability mass function of a zero truncated binomial distribution with size parameter d and second parameter $1/d$.

4. Conclusion

In this work, a novel Bayesian approach to the problem of variable selection was proposed. To this aim nearest neighbor Gaussian processes, reference priors, and model selection strategies based on random sets were combined. To the best of our knowledge, such a combination has not been investigated in previous works. While the application of the nearest neighbor processes reduces the computational cost compared to classical processes, the usage of reference priors guarantees a (numerically) robust model inference. The random set approach allows for a selection of the most predictive variables, whilst incorporating appropriate a priori knowledge. For the model inference a Metropolis within Gibbs algorithm was proposed. The algorithm allows for transitions between different sets of active predictors (predictors that truly contribute to the model) and, further, uses Hamiltonian dynamics for the inference of the Gaussian process correlation parameters. The performance of the proposed method was extensively evaluated. In evaluations including simulated data and an approximation problem corresponding to computer experiments, the new approach could identify truly significant numbers. Purely linear models, anisotropic Gaussian processes (automatic relevance determination), and random forest failed at these variable selection tasks. Finally, a real-world application related to the semiconductor manufacturing industry showed that the proposed method is of real value for companies and organizations of many kinds.

Acknowledgments

The publication is one of the results of the project iDev40 (www.idev40.eu). This project has received funding from the ECSEL Joint Undertaking (JU) under grant agreement No 783163. The JU receives support from the European Unions Horizon 2020 research and innovation programme and Austria (BMVIT-IKT der Zukunft, FFG project no. 865348), Germany, Belgium, Italy, Spain, Romania. The information and results set out in this publication are those of the authors and do not necessarily reflect the opinion of the ECSEL Joint Undertaking.



Appendix A. Derivatives for Reference Prior

Calculation rules for matrices:

$$\frac{\partial \mathbf{AB}}{\partial t} = \frac{\partial \mathbf{A}}{\partial t} \mathbf{B} + \mathbf{A} \frac{\partial \mathbf{B}}{\partial t} \quad (\text{A.1})$$

$$\frac{\partial \mathbf{A}^{-1}}{\partial t} = -\mathbf{A}^{-1} \frac{\partial \mathbf{A}}{\partial t} \mathbf{A}^{-1} \quad (\text{A.2})$$

$$\frac{\partial \tilde{\mathbf{K}}_S}{\partial \rho} = \frac{\partial}{\partial \rho} \mathbf{B}_S^{-1} \overbrace{\text{diag}(\mathcal{F}_{\mathbf{x}_1}, \dots, \mathcal{F}_{\mathbf{x}_n})}^{=: \mathcal{F}_S} (\mathbf{B}_S^T)^{-1} \quad (\text{A.3})$$

$$= \frac{\partial \mathbf{B}_S^{-1} \mathcal{F}_S}{\partial \rho} (\mathbf{B}_S^T)^{-1} + \mathbf{B}_S^{-1} \mathcal{F}_S \frac{\partial (\mathbf{B}_S^T)^{-1}}{\partial \rho} \quad (\text{A.4})$$

$$= \left(\frac{\partial \mathbf{B}_S^{-1}}{\partial \rho} \mathcal{F}_S + \mathbf{B}_S^{-1} \frac{\partial \mathcal{F}_S}{\partial \rho} \right) (\mathbf{B}_S^T)^{-1} + \mathbf{B}_S^{-1} \mathcal{F}_S \frac{\partial (\mathbf{B}_S^T)^{-1}}{\partial \rho} \quad (\text{A.5})$$

$$= \left(-\mathbf{B}_S^{-1} \frac{\partial \mathbf{B}_S}{\partial \rho} \mathbf{B}_S^{-1} \mathcal{F}_S + \mathbf{B}_S^{-1} \frac{\partial \mathcal{F}_S}{\partial \rho} \right) (\mathbf{B}_S^T)^{-1} - \mathbf{B}_S^{-1} \mathcal{F}_S (\mathbf{B}_S^T)^{-1} \frac{\partial (\mathbf{B}_S^T)}{\partial \rho} (\mathbf{B}_S^T)^{-1} \quad (\text{A.6})$$

$$= -\mathbf{B}_S^{-1} \left\{ \underbrace{\frac{\partial \mathbf{B}_S}{\partial \rho} \mathbf{B}_S^{-1} \mathcal{F}_S}_{=: \mathbf{A}} - \frac{\partial \mathcal{F}_S}{\partial \rho} + \underbrace{\mathcal{F}_S (\mathbf{B}_S^T)^{-1} \frac{\partial (\mathbf{B}_S^T)}{\partial \rho}}_{\mathbf{A}^T} \right\} (\mathbf{B}_S^T)^{-1} \quad (\text{A.7})$$

$$= -\mathbf{B}_S^{-1} \left\{ \mathbf{A} + \mathbf{A}^T - \frac{\partial \mathcal{F}_S}{\partial \rho} \right\} (\mathbf{B}_S^T)^{-1} \quad (\text{A.8})$$

$$\frac{\partial \mathbf{B}_S}{\partial \rho} = \frac{\partial [B_{\mathbf{x}_i, j}]_{i, j=1, \dots, n}}{\partial \rho} \quad B_{\mathbf{x}_i, j} = \begin{cases} 1 & i = j \\ -\mathbf{B}_{\mathbf{x}_i}[l] & \mathbf{x}_j = \mathbf{x}_i^l \text{ for some } l \\ 0 & \text{else} \end{cases} \quad (\text{A.9})$$

$$(\text{A.10})$$

$$\frac{\partial \mathbf{B}_{\mathbf{x}_i}}{\partial \rho} = \frac{\partial \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)} \mathcal{K}_{N(\mathbf{x}_i)}^{-1}}{\partial \rho} \quad (\text{A.11})$$

$$= \frac{\partial \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)}}{\partial \rho} \mathcal{K}_{N(\mathbf{x}_i)}^{-1} + \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)} \frac{\partial \mathcal{K}_{N(\mathbf{x}_i)}^{-1}}{\partial \rho} \quad (\text{A.12})$$

$$= \frac{\partial \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)}}{\partial \rho} \mathcal{K}_{N(\mathbf{x}_i)}^{-1} - \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)} \mathcal{K}_{N(\mathbf{x}_i)}^{-1} \frac{\partial \mathcal{K}_{N(\mathbf{x}_i)}}{\partial \rho} \mathcal{K}_{N(\mathbf{x}_i)}^{-1} \quad (\text{A.13})$$

$$= \left(\frac{\partial \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)}}{\partial \rho} - \mathbf{B}_{\mathbf{x}_i} \frac{\partial \mathcal{K}_{N(\mathbf{x}_i)}}{\partial \rho} \right) \mathcal{K}_{N(\mathbf{x}_i)}^{-1} \quad (\text{A.14})$$

Thus:

$$\frac{\partial \mathbf{B}_S}{\partial \rho} = \begin{cases} 0 & i \geq j \\ \left\{ \left(\mathbf{B}_{\mathbf{x}_i} \frac{\partial \mathcal{K}_{N(\mathbf{x}_i)}}{\partial \rho} - \frac{\partial \mathcal{K}_{\mathbf{x}_i, N(\mathbf{x}_i)}}{\partial \rho} \right) \mathcal{K}_{N(\mathbf{x}_i)}^{-1} \right\} [l] & i < j \text{ and } \mathbf{x}_j = \mathbf{x}_i^l \text{ for some } l \\ 0 & \text{else} \end{cases} \quad (\text{A.15})$$

$$\frac{\partial \mathcal{F}_S}{\partial \rho} = \text{diag} \left(\frac{\partial \mathcal{F}_{\mathbf{x}_1}}{\rho}, \dots, \frac{\partial \mathcal{F}_{\mathbf{x}_n}}{\rho} \right) \quad (\text{A.16})$$

$$\frac{\partial \mathcal{F}_{\mathbf{x}_i}}{\rho} = \frac{\partial \mathcal{K}_{\mathbf{x}_i} - \mathbf{B}_{\mathbf{x}_i} \mathcal{K}_{N(\mathbf{x}_i), \mathbf{x}_i}}{\partial \rho} \quad (\text{A.17})$$

$$= \frac{\partial \mathcal{K}_{\mathbf{x}_i}}{\partial \rho} - \left(\frac{\partial \mathbf{B}_{\mathbf{x}_i}}{\partial \rho} \mathcal{K}_{N(\mathbf{x}_i), \mathbf{x}_i} + \mathbf{B}_{\mathbf{x}_i} \frac{\partial \mathcal{K}_{N(\mathbf{x}_i), \mathbf{x}_i}}{\partial \rho} \right) \quad (\text{A.18})$$

All the calculations above hold also for \mathbf{W}_γ when replacing ρ with γ .

$$\frac{\partial \mathcal{K}_{\mathbf{x}_i, \mathbf{x}_j}}{\partial \rho} = \gamma \frac{\partial K_{\mathbf{x}_i, \mathbf{x}_j}}{\partial \rho} \quad (\text{A.19})$$

$$\frac{\partial \mathcal{K}_{\mathbf{x}_i, \mathbf{x}_j}}{\partial \gamma} = -\delta_{(\mathbf{x}_i = \mathbf{x}_j)} + K_{\mathbf{x}_i, \mathbf{x}_j} \quad (\text{A.20})$$

$$\frac{\partial K_{\mathbf{x}_i, \mathbf{x}_j}}{\partial \rho} = \left[-\frac{\sqrt{5}d_A}{\rho^2} - \frac{10d_A^2}{3\rho^3} \right] \exp \left(-\frac{\sqrt{5}d_A}{\rho} \right) + K_{\mathbf{x}_i, \mathbf{x}_j} \frac{\sqrt{5}d_A}{\rho^2} \quad (\text{A.21})$$

References

- Alhamzawi, R., Taha Mohammad Ali, H., 2018. The bayesian adaptive lasso regression. *Mathematical Biosciences* 303, 75 – 82.
- Berger, J.O., de Oliveira, V., Sanso, B., 2001. Objective bayesian analysis of spatially correlated data. *Journal of the American Statistical Association* 96, 1361–1374. URL: <http://www.jstor.org/stable/3085905>.
- Breiman, L., 2001. Machine learning, volume 45, number 1 - springerlink. *Machine Learning* 45, 5–32. doi:10.1023/A:1010933404324.
- Buehlmann, P., Drineas, P., Kane, M., van der Laan, M., 2016. *Handbook of Big Data*. Chapman & Hall/CRC.
- Carnell, R., 2020. lhs: Latin Hypercube Samples. URL: <https://CRAN.R-project.org/package=lhs>. r package version 1.1.1.
- Chen, T., Morris, J., Martin, E., 2007. Gaussian process regression for multivariate spectroscopic calibration. *Chemometrics and Intelligent Laboratory Systems* 87, 59 – 71. doi:10.1016/j.chemolab.2006.09.004.
- Chen, T., Wang, B., 2010. Bayesian variable selection for gaussian process regression: Application to chemometric calibration of spectrometers. *Neurocomputing* 73, 2718 – 2726. doi:10.1016/j.neucom.2010.04.014. pattern Recognition in Bioinformatics Advances in Neural Control.

- Dancik, G.M., Dorman, K.S., 2008. mlegp: statistical analysis for computer models of biological systems using r. *Bioinformatics* 24, 1966–1967.
- Datta, A., Banerjee, S., Finley, A.O., Gelfand, A.E., 2016. Hierarchical nearest-neighbor gaussian process models for large geostatistical datasets. *Journal of the American Statistical Association* 111, 800–812.
- Dette, H., Pepelyshev, A., 2010. Generalized latin hypercube design for computer experiments. *Technometrics* 52, 421–429. doi:10.1198/TECH.2010.09157.
- Duane, S., Kennedy, A., Pendleton, B.J., Roweth, D., 1987. Hybrid monte carlo. *Physics Letters B* 195, 216 – 222. doi:10.1016/0370-2693(87)91197-X.
- Eddelbuettel, D., François, R., 2011. Rcpp: Seamless R and C++ integration. *Journal of Statistical Software* 40, 1–18. doi:10.18637/jss.v040.i08.
- Eddelbuettel, D., Sanderson, C., 2014. Rcpparmadillo: Accelerating r with high-performance c++ linear algebra. *Computational Statistics and Data Analysis* 71, 1054–1063. URL: <http://dx.doi.org/10.1016/j.csda.2013.02.005>.
- Fan, J., Li, R., 2001. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American Statistical Association* 96, 1348–1360.
- Foster, D.P., Stine, R.A., 2004. Variable selection in data mining: Building a predictive model for bankruptcy. *Journal of the American Statistical Association* 99, 303–313.
- Friedman, J., Hastie, T., Tibshirani, R., 2010. Regularization paths for generalized linear models via coordinate descent. *Journal of Statistical Software* 33, 1–22. URL: <http://www.jstatsoft.org/v33/i01/>.
- Gu, M., Palomo, J., Berger, J., 2020. RobustGaSP: Robust Gaussian Stochastic Process Emulation. URL: <https://CRAN.R-project.org/package=RobustGaSP>. r package version 0.6.0.
- Gu, M., Wang, X., Berger, J.O., 2018. Robust gaussian stochastic process emulation. *Ann. Statist.* 46, 3038–3066. URL: <https://doi.org/10.1214/17-AOS1648>, doi:10.1214/17-AOS1648.
- Johnson, R.W., 1996. Fitting percentage of body fat to simple body measurements. *Journal of Statistics Education* 4. doi:10.1080/10691898.1996.11910505.
- Kazianka, H., Pilz, J., 2012. Objective bayesian analysis of spatial data with uncertain nugget and range parameters. *The Canadian Journal of Statistics / La Revue Canadienne de Statistique* 40, 304–327. URL: <http://www.jstor.org/stable/41724530>.
- Lee, K.E., Sha, N., Dougherty, E.R., Vannucci, M., Mallick, B.K., 2003. Gene selection: a bayesian variable selection approach. *Bioinformatics* 19 1, 90–7.
- Liaw, A., Wiener, M., 2002. Classification and regression by randomforest. *R News* 2, 18–22. URL: <https://CRAN.R-project.org/doc/Rnews/>.
- Linkletter, C., Bingham, D., Hengartner, N., Higdon, D., Ye, K.Q., 2006. Variable selection for gaussian process models in computer experiments. *Technometrics* 48, 478–490. doi:10.1198/004017006000000228.
- Neal, R., 1997. Monte carlo implementation of gaussian process models for bayesian regression and classification. arXiv: Data Analysis, Statistics and Probability .
- Neal, R.M., 1996. *Bayesian Learning for Neural Networks*. Springer-Verlag, Berlin, Heidelberg.
- O’Hagan, A., 1978. Curve fitting and optimal design for prediction. *Journal of the Royal Statistical Society. Series B (Methodological)* 40, 1–42. URL: <http://www.jstor.org/stable/2984861>.
- Park, T., Casella, G., 2008. The bayesian lasso. *Journal of the American Statistical Association* 103, 681–686.
- Paulo, R., 2005. Default priors for gaussian processes. *The Annals of Statistics* 33, 556–582.
- Pleschberger, M., 2021. Data set of extracted summary statistics from equipment sensor data. doi:10.5281/zenodo.4462777. dataset on Zenodo, <https://doi.org/10.5281/zenodo.4462777>.
- Posch, K., Arbeiter, M., Pilz, J., 2020. A novel bayesian approach for variable selection in linear regression models. *Computational Statistics & Data Analysis* 144, 106881. doi:10.1016/j.csda.2019.106881.
- Rasmussen, C.E., Williams, C.K.I., 2005. *Gaussian Processes for Machine Learning*. MIT Press.
- Ročková, V., George, E.I., 2014. Emvs: The em approach to bayesian variable selection. *Journal of the American Statistical Association* 109, 828–846.
- Santner, T.J., Williams, B.J., Notz, W.I., 2003. *The Design and Analysis of Computer Experiments*. Springer.
- Tarr, G., Müller, S., Welsh, A.H., 2018. mplot: An R package for graphical model stability and variable selection procedures. *Journal of Statistical Software* 83, 1–28. doi:10.18637/jss.v083.i09.
- Tibshirani, R., 1996. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society. Series B (Methodological)* 58, 267–288.
- Vecchia, A.V., 1988. Estimation and model identification for continuous spatial processes. *Journal of the Royal Statistical Society. Series B (Methodological)* 50, 297–312.
- Vollert, N., Ortner, M., Pilz, J., 2019. Robust additive gaussian process models using reference priors and cut-off-designs. *Applied Mathematical Modelling* 65, 586–596. doi:10.1016/j.apm.2018.07.050.
- Wu, X., Chan, L.L.T., Chen, J., Xie, L., 2017. Application of gaussian processes with variable shrinkage method and just-in-time modeling in the semiconductor industry, in: 2017 6th International Symposium on Advanced Control of Industrial Processes (AdCONIP), pp. 78–83. doi:10.1109/ADCONIP.2017.7983759.
- Yi, G., Shi, J.Q., Choi, T., 2011. Penalized gaussian process regression and classification for high-dimensional nonlinear data. *Biometrics* 67, 1285–1294. URL: <http://www.jstor.org/stable/41434433>.
- Yuan, J., Wang, K., yu, z., Fang, M., 2008. Reliable multi-objective optimization of high-speed wedm process based on gaussian process regression. *International Journal of Machine Tools and Manufacture* 48, 47–60. doi:10.1016/j.ijmactools.2007.07.011.
- Zheng, H., Zhang, Y., 2007. Feature selection for high dimensional data in astronomy. *Advances in Space Research* 41, 1960 –

- 1964.
- Zou, H., 2006. The adaptive lasso and its oracle properties. *Journal of the American Statistical Association* 101, 1418–1429.
- Zou, H., Hastie, T., 2005. Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 67, 301–320.