

A quantum bound on the compactness

Roberto Casadio^{ab*}

^a*Dipartimento di Fisica e Astronomia, Università di Bologna
via Irnerio 46, 40126 Bologna, Italy*

^b*I.N.F.N., Sezione di Bologna, I.S. FLAG
viale B. Pichat 6/2, 40127 Bologna, Italy*

October 11, 2021

Abstract

We present a simple quantum description of the gravitational collapse of a ball of dust which excludes those states whose width is arbitrarily smaller than the gravitational radius of the matter source and supports the conclusion that black holes are macroscopic extended objects. We also comment briefly on the relevance of this result for the ultraviolet self-completion of gravity and the connection with the corpuscular picture of black holes.

1 Introduction and motivation

It has been often argued that quantum gravity should remove the singularity predicted by General Relativity at the endpoint of the gravitational collapse [1]. We study this issue by considering a simple quantum description of the gravitational collapse of a ball of dust [2]. Since gravity is the only interaction acting on dust, the areal radius R of the ball classically follows a radial geodesic in the (outer) Schwarzschild spacetime ¹

$$ds^2 = - \left(1 - \frac{2 G_N M}{r} \right) dt^2 + \left(1 - \frac{2 G_N M}{r} \right)^{-1} dr^2 + r^2 d\Omega^2, \quad (1.1)$$

where M is the Arnowitt-Deser-Misner (ADM) [3] mass of the dust. In analogy with the quantum mechanics of the hydrogen atom, in which one quantises the position of the electron with respect to the centre-of-mass of the system, only the radius R of the ball will be quantised here. ² The radial geodesic equation will then become a time-independent Schrödinger equation for a particle in the Newtonian potential, with the important feature that quantum states with width significantly smaller than the gravitational radius

$$R_H = 2 G_N M \quad (1.2)$$

*E-mail: casadio@bo.infn.it

¹We use units with $c = 1$, the reduced Planck constant $\hbar = \ell_p m_p$ and the Newton constant $G_N = \ell_p/m_p$, where ℓ_p is the Planck length and m_p the Planck mass.

²For more refined approaches in which the singularity is also removed, see *e.g.* Refs. [4].

are not physically allowed. This is similar to the quantum removal of the classical UV catastrophe in the hydrogen atom, except that the mass M naturally introduces a lower bound on the energy spectrum which would be absent in the Newtonian approximation. One could then infer that the quantum nature of black holes as extended objects follows from the non-linearity of the gravitational interaction as it is described by General Relativity.

The main results will be obtained in the next Section, with more speculative considerations summarised in Section 3.

2 Minisuperspace model for the gravitational collapse

We consider a self-gravitating ball of dust of radius R and ADM mass M [2,5]. In General Relativity, the surface of the ball follows a radial geodesic in the Schwarzschild spacetime (1.1) and its areal radius $R = R(\tau)$ must therefore satisfy the equation

$$\left(\frac{dR}{d\tau}\right)^2 + 1 - \frac{2G_N M}{R} = \frac{E^2}{M^2}, \quad (2.1)$$

where τ is the proper time and $E < M$ is the conserved energy of a bound trajectory. The above can be rewritten as

$$H \equiv \frac{P^2}{2M} - \frac{G_N M^2}{R} = \frac{M}{2} \left(\frac{E^2}{M^2} - 1 \right) \equiv \mathcal{E}, \quad (2.2)$$

where we introduced the momentum $P = M(dR/d\tau)$. One can immediately notice that Eq. (2.2) is formally the same as the Newtonian equation for energy conservation.³

The study of spherically symmetric gravitational collapse in the quantum theory has a long history (for a very limited selection of papers, see Refs. [4,6]). We here proceed with a simple quantisation prescription for the above equation by introducing the momentum operator $\hat{P} = -i\hbar\partial/\partial R$. This is physically equivalent to assuming that the radius of the ball R satisfies an uncertainty relation stemming from the usual canonical commutator, that is

$$[\hat{R}, \hat{P}] = i\hbar \quad \Rightarrow \quad \Delta R \Delta P \gtrsim \hbar = \ell_p m_p, \quad (2.3)$$

where $\Delta O \equiv \langle \hat{O}^2 \rangle - \langle \hat{O} \rangle^2$ for $\hat{O} = \hat{R}$ or \hat{P} . Moreover, all of the expectation values are taken on wavefunctions $\Psi = \Psi(R)$ satisfying

$$\hat{H} \Psi = \mathcal{E} \Psi, \quad (2.4)$$

which is just the time-independent Schrödinger equation for a gravitational (Newtonian) atom. In particular, the energy spectrum contains the eigenstates

$$\Psi_n = \sqrt{\frac{M^9}{\pi n^5 \ell_p^3 m_p^9}} e^{-\frac{M^3 r}{n m_p^3 \ell_p}} L_{n-1}^1 \left(\frac{2 M^3 r}{n m_p^3 \ell_p} \right). \quad (2.5)$$

³In fact, Eq. (2.2) is the mass-shell condition for the surface of the ball and the Hamiltonian constraint of General Relativity for dust.

where the integer number $n \geq 1$ and L_{n-1}^1 are the generalised Laguerre polynomials (for zero angular momentum). The corresponding eigenvalues are given by ⁴

$$\frac{\mathcal{E}_n}{M} \simeq -\frac{G_{\text{N}}^2 M^4}{2 \hbar^2 n^2} = -\frac{1}{2 n^2} \left(\frac{M}{m_{\text{p}}} \right)^4 = \frac{1}{2} \left(\frac{E_n^2}{M^2} - 1 \right), \quad (2.6)$$

and one also has

$$R_n \equiv \langle \Psi_n | R | \Psi_n \rangle \simeq \frac{\hbar^2 n^2}{G_{\text{N}} M^3} = n^2 \ell_{\text{p}} \left(\frac{m_{\text{p}}}{M} \right)^3. \quad (2.7)$$

At first sight, it thus appears that the spectrum contains states Ψ_n of infinitesimally small width, since $R_1 \sim \ell_{\text{p}} (m_{\text{p}}/M)^3 \ll \ell_{\text{p}}$ for any macroscopic objects, like stars, whose mass $M \gg m_{\text{p}}$. This state would have an energy density of the order of $M/R_1^3 \sim (M^{10}/m_{\text{p}}^9) \ell_{\text{p}}^{-3}$, which can hardly be considered a satisfying alternative to the classical singularity of infinite energy density.

However, Eq. (2.6) yields

$$0 \leq \frac{E_n^2}{M^2} \simeq 1 - \frac{1}{n^2} \left(\frac{M}{m_{\text{p}}} \right)^4, \quad (2.8)$$

tantamount to $\mathcal{E} \geq -M/2$ in Eq. (2.2), and we thus find that acceptable states Ψ_n must satisfy

$$n \geq N_M \simeq \left(\frac{M}{m_{\text{p}}} \right)^2. \quad (2.9)$$

Correspondingly, we obtain

$$R_n \gtrsim R_{N_M} \sim R_{\text{H}}, \quad (2.10)$$

which results in the quantum upper bound

$$\frac{G_{\text{N}} M}{R_n} \lesssim 1. \quad (2.11)$$

Moreover, we observe that the Hamiltonian eigenvalues are also bounded from below as

$$\mathcal{E}_n \geq \mathcal{E}_{N_M} \simeq -\frac{M}{2}, \quad (2.12)$$

which corresponds to

$$E_n^2 \geq E_{N_M}^2 \simeq 0. \quad (2.13)$$

Since any semiclassical state representing the collapsing ball must be described in terms of superpositions of the Ψ_n 's with $n \geq N_M$, the above quantum bound (2.11) implies that any ball of dust must have compactness $G_{\text{N}} M/R \lesssim 1$. This result holds up to a factor of order one and better estimates should be obtained only by considering more realistic and complete models.

⁴Numerical coefficients of order one will often be omitted or approximated for the sake of clarity, given that the simplicity of the model diminishes their physical relevance.




Figure 1: Probability density $\mathcal{P} = \mathcal{P}(R)$ for: a) ground state with $n = N_M = M^2/m_p^2$ [solid (dashed) line represents the region inside (outside) the gravitational radius $R_H = 2 G_N M$]; b) first excited state with $n = N_M + 1$ (dotted line).

The present minisuperspace description only contains the observable R . From the wavefunction $\Psi = \Psi(R)$, we can therefore only determine such information as the expectation value of R and the probability that the ball be inside the gravitational radius,

$$P(R \leq R_H) \equiv \int_0^{R_H} \mathcal{P}(R) dR = 4\pi \int_0^{R_H} |\Psi(R)|^2 R^2 dR, \quad (2.14)$$

which can be viewed as the probability that the dust ball is a black hole (when the mass M is treated as a fixed parameter⁵). For the ground state, whose wavefunction is given by

$$\Psi = \Psi_{N_M} \simeq \sqrt{\frac{m_p}{\pi \ell_p^3 M}} e^{-\frac{M r}{m_p \ell_p}} L_{\frac{M^2}{m_p^2} - 1}^1 \left(\frac{2 M r}{m_p \ell_p} \right), \quad (2.15)$$

the probability density $\mathcal{P} = \mathcal{P}_{N_M}(R)$ is plotted in Fig. 1 for a few cases with very small M . For values of $M \gg m_p$, the probability density narrowly peaks around a value of R slightly below R_H and one thus finds

$$P_{N_M}(R \leq R_H) \simeq 1, \quad (2.16)$$

with uncertainty

$$\Delta R_{N_M} \sim \frac{R_{N_M}}{N_M} \sim \ell_p \frac{m_p}{M}. \quad (2.17)$$

⁵Alternative viewpoints are considered in Refs. [7], where the mass M is quantised, and in Ref. [8], where M is allowed a statistical uncertainty.

As expected, $\Delta R_{N_M} \ll R_{N_M}$, hence the radius of a very massive matter source behaves like a classical variable.

Fig. 1 also displays the first excited state, which asymptotically approaches the corresponding ground state for increasing M . This implies that $P_n(R \leq R_H) \simeq 1$ for a range of states $n \gtrsim N_M$, which suggests that large astrophysical black holes of a given mass M might not necessarily be in their ground state with $n = N_M$. In fact, for $n \gtrsim N_M$, the quantum of the Hamiltonian H in Eq. (2.2) is given by

$$\delta H \equiv |\mathcal{E}_{n+1} - \mathcal{E}_n| \simeq m_p \frac{m_p}{M}, \quad (2.18)$$

so that $\delta H \sim m_p (\Delta R_{N_M} / \ell_p) \ll m_p$ for a macroscopic object of mass $M \gg m_p$. Furthermore, since

$$\delta E \equiv |E_{n+1} - E_n| \simeq m_p, \quad (2.19)$$

it appears that the proper source “energy” is naturally quantised in units of the fundamental Planck mass m_p , but this quantum is redshifted down to the much smaller δH measured by outer observers.

3 Concluding remarks and outlook

We have shown how the General Relativistic description of gravity leads to an upper bound for the compactness of a ball of dust in the quantum theory. It is interesting to notice that the lower bounds on the eigenvalue $\mathcal{E}_n \geq \mathcal{E}_{N_M}$ and radius $R_n \gtrsim G_N M$ could not be obtained in the Newtonian theory, since $\mathcal{E} = E$ in that approximation, and the entire spectrum Ψ_n with $n \geq 1$ would be physically acceptable therein. The bound (2.11) on the compactness therefore follows from the nonlinearity of General Relativity and agrees with previous results [9] obtained by adding a gravitational self-interaction term to the Newtonian theory [10]. It also agrees with those results following from the quantum description of the gravitational radius and black hole horizon [7]. In particular, the latter approach leads to very similar conclusions to the ones shown in the previous Section, like the black hole probability (2.16) and radius uncertainty (2.17), when the self-gravitating object is described by an extended many-body system with a very large occupation number of order $N_M \sim M^2/m_p^2$ in its ground state [11]. The first excited modes could then be populated thermally [11] and reproduce the Hawking radiation [12].

It is indeed very intriguing that Eq. (2.9) and the bound (2.11) allow for recovering the fundamental scaling relations employed in the corpuscular description of black holes [13]. In fact, the allowed state of minimum energy (2.15) has principal quantum number

$$N_M \simeq N_G, \quad (3.1)$$

where N_G is the number of soft gravitons in the coherent state representing the gravitational potential generated by a source of mass M [14, 15]. We recall that the corpuscular picture naturally reproduces Bekenstein’s area spectrum $G_N^2 M^2 \sim N_G \ell_p^2$ [16], so that the black hole ADM mass is quantised in units of

$$\delta M \simeq \delta H, \quad (3.2)$$

where δH is precisely the quantum of our Hamiltonian given in Eq. (2.18). This is also the typical energy of particles emitted by the Hawking evaporation process, which the corpuscular picture describes as the depletion of the quantum state of gravity.

In this perspective, the wavefunction $\Psi_{N_M} = \Psi_{N_G}$ appears as the “non-perturbative ground state” for self-gravitating macroscopic objects of mass M and should thus be the closest possible to a classical black hole (as we have already commented at the end of the previous Section). This result is consistent with the fact that a static gravitational field must be fully determined by the state of the source, as it does not represent independent degrees of freedom in a static system [7]. It can further be interpreted as the fact that the quantum state of a macroscopic self-gravitating system of mass M is very far from the vacuum Ψ_0 of quantum gravity, for which $N_G = 0$. The number $N_M \sim N_G \sim M^2$ hence provides a quantitative measure for this “distance” from the vacuum in the Hilbert space of quantum states. The appearance of the ground state $\Psi_{N_M} = \Psi_{N_G}$ when the ADM energy is given by Eq. (2.9), in turn, can be interpreted as a form of *classicalization* ensuring the ultraviolet self-completeness of gravity [17], because states with spatial momenta much larger than $\hbar/R_{N_M} \sim \hbar/R_H$ have $n < N_M$ and cannot be populated at energy scales of the order of M .

We conclude by noting that, although states corresponding to the classical singularity appear to be removed from the spectrum, it would be phenomenologically very important to determine a more precise maximum value of the compactness in order to assess the size of quantum deviations from the exterior Schwarzschild geometry. Moreover, in order to make contact with actual observations, one needs an explicit description of the exterior spacetime and of the interaction among the collapsing object and signals that can reach out to our detectors. For example, an effectively finite size of the self-gravitating system is expected to induce the kind of quantum deviations obtained in Ref. [15] from a ultraviolet cut-off for the momenta generating the outer mean-field geometry. A unified description of the quantum states of the collapsing ball presented here and the quantum gravitational potential of Ref. [15] is left for future investigations. That could serve as the starting point for reconstructing the exterior spacetime geometry of a quantum black hole, and allow us to search for phenomenological signatures of quantum physics in the geodesic motion of test bodies.

Acknowledgments

R.C. is partially supported by the INFN grant FLAG and his work has also been carried out in the framework of activities of the National Group of Mathematical Physics (GNFM, INdAM) and COST action *Cantata*.

References

- [1] S. W. Hawking and G. F. R. Ellis, “The Large Scale Structure of Space-Time” (Cambridge University Press, Cambridge, 1973)
- [2] J. R. Oppenheimer and H. Snyder, Phys. Rev. **56** (1939) 455.
- [3] R.L. Arnowitt, S. Deser and C.W. Misner, Phys. Rev. **116** (1959) 1322.
- [4] V. P. Frolov and G. A. Vilkovisky, Phys. Lett. B **106** (1981) 307; D. Malafarina, Universe **3** (2017) 48 [arXiv:1703.04138 [gr-qc]]; R. Casadio, Int. J. Mod. Phys. D **9** (2000) 511 [arXiv:gr-qc/9810073 [gr-qc]]; H. M. Haggard and C. Rovelli, Phys. Rev. D **92** (2015) 104020 [arXiv:1407.0989 [gr-qc]]; W. Piechocki and T. Schmitz, Phys. Rev. D **102** (2020) 046004 [arXiv:2004.02939 [gr-qc]]; T. Schmitz, Phys. Rev. D **103** (2021) 064074 [arXiv:2012.04383 [gr-qc]].

- [5] H. Stephani, “Relativity” (Cambridge University Press, Cambridge, 2004)
- [6] P. Hajicek, B. S. Kay and K. V. Kuchar, *Phys. Rev. D* **46** (1992) 5439; P. Hajicek and C. Kiefer, *Int. J. Mod. Phys. D* **10** (2001) 775 [arXiv:gr-qc/0107102 [gr-qc]]; R. Casadio and G. Venturi, *Class. Quant. Grav.* **13** (1996) 2715 [arXiv:gr-qc/9512032 [gr-qc]].
- [7] R. Casadio, A. Giugno, O. Micu and A. Orlandi, *Phys. Rev. D* **90** (2014) 084040 [arXiv:1405.4192 [hep-th]]; R. Casadio, “Localised particles and fuzzy horizons: A tool for probing Quantum Black Holes,” [arXiv:1305.3195 [gr-qc]]; R. Casadio and F. Scardigli, *Eur. Phys. J. C* **74** (2014) 2685 [arXiv:1306.5298 [gr-qc]]; R. Casadio, A. Giugno and A. Giusti, *Gen. Rel. Grav.* **49** (2017) 32 [arXiv:1605.06617 [gr-qc]].
- [8] R. Casadio, A. Giusti and A. Mentrelli, *Phys. Rev. D* **100** (2019) 024036 [arXiv:1901.06206 [gr-qc]].
- [9] R. Casadio, M. Lenzi and A. Ciarfella, *Phys. Rev. D* **101** (2020)124032 [arXiv:2002.00221 [gr-qc]].
- [10] R. Casadio, M. Lenzi and O. Micu, *Phys. Rev. D* **98** (2018) 104016 [arXiv:1806.07639 [gr-qc]]; *Eur. Phys. J. C* **79** (2019) 894 [arXiv:1904.06752 [gr-qc]]; R. Casadio and I. Kuntz, *Eur. Phys. J. C* **80** (2020) 581 [arXiv:2003.03579 [gr-qc]]; R. Casadio, A. Giusti, I. Kuntz and G. Neri, *Phys. Rev. D* **103** (2021) 064001 [arXiv:2101.12471 [gr-qc]].
- [11] R. Casadio, A. Giugno and A. Orlandi, *Phys. Rev. D* **91** (2015) 124069 [arXiv:1504.05356 [gr-qc]].
- [12] S. W. Hawking, *Commun. Math. Phys.* **43** (1975) 199 [erratum: *Commun. Math. Phys.* **46** (1976) 206].
- [13] G. Dvali and C. Gomez, *Fortsch. Phys.* **61** (2013) 742 [arXiv:1112.3359 [hep-th]]; G. Dvali, C. Gomez and S. Mukhanov, “Black Hole Masses are Quantized,” arXiv:1106.5894 [hep-ph]. G. Dvali and C. Gomez, *Phys. Lett. B* **719** (2013) 419 [arXiv:1203.6575 [hep-th]]; *Phys. Lett. B* **716** (2012) 240 [arXiv:1203.3372 [hep-th]]; *Eur. Phys. J. C* **74** (2014) 2752 [arXiv:1207.4059 [hep-th]]; S. Hofmann and T. Rug, *Nucl. Phys. B* **902** (2016) 302 [arXiv:1403.3224 [hep-th]]; A. Giusti, *Int. J. Geom. Meth. Mod. Phys.* **16** (2019) 1930001.
- [14] R. Casadio, A. Giugno and A. Giusti, *Phys. Lett. B* **763** (2016) 337 [arXiv:1606.04744 [gr-qc]]; W. Mück, *Can. J. Phys.* **92** (2014) 973 [arXiv:1306.6245 [hep-th]]; R. Casadio, A. Giugno, A. Giusti and M. Lenzi, *Phys. Rev. D* **96** 044010 (2017) [arXiv:1702.05918 [gr-qc]].
- [15] R. Casadio, “Quantum black holes and resolution of the singularity,” [arXiv:2103.00183 [gr-qc]].
- [16] J. D. Bekenstein, *Phys. Rev. D* **7** (1973) 2333.
- [17] G. Dvali, G. F. Giudice, C. Gomez and A. Kehagias, *JHEP* **1108** (2011) 108 [arXiv:1010.1415 [hep-ph]]; G. Dvali and D. Pirtskhalava, *Phys. Lett. B* **699** (2011) 78 [arXiv:1011.0114 [hep-ph]]; G. Dvali, C. Gomez and A. Kehagias, *JHEP* **1111** (2011) 070 [arXiv:1103.5963 [hep-th]]; R. Percacci and L. Rachwal, *Phys. Lett. B* **711** (2012) 184 [arXiv:1202.1101 [hep-th]].