

Testing For a Parametric Baseline-Intensity in Dynamic Interaction Networks

Alexander Kreiss
ORSTAT, KU Leuven
and
Enno Mammen
Institute for Applied Mathematics, Heidelberg University
and
Wolfgang Polonik *
Department of Statistics, University of California at Davis

October 11, 2021

Abstract

In statistical network analysis it is common to observe so called interaction data. Such data is characterized by the actors who form the vertices of a network. These are able to interact with each other along the edges of the network. One usually assumes that the edges in the network are randomly formed and dissolved over the observation horizon. In addition covariates are observed and the interest is to model the impact of the covariates on the interactions. In this paper we develop a framework to test if a non-parametric form of the baseline intensity allows for more flexibility than a baseline which is parametrically dependent on system-wide covariates (i.e. covariates which take the same value for all individuals, e.g. time). This allows to test if certain seasonality effects can be explained by simple covariates like the time. The procedure is applied to modeling the baseline intensity in a bike-sharing network by using weather and time information.

1 Introduction

One branch in statistical network analysis is concerned with the analysis of so called interaction data. Some references for this type of data are for example [Butts \(2008\)](#); [Perry and Wolfe \(2013\)](#); [Kreiss et al. \(2019\)](#). The data typically comprises a sequence of random networks where the vertices are some sort of actors who can interact with each other along the edges of the network. Typically, it is assumed that the vertex set, i.e. the actors, are fixed

*W. Polonik acknowledges partial support by the National Science Foundation under Grant No. DMS-1713108.

but that the edge set, i.e. their relations, are random. In this paper we will also consider dynamically changing networks, in which it is allowed that the edge set randomly changes over the observation horizon. A classical example would be social contact networks: Here the vertices of the network under consideration are people. An interaction, or an event, between two people could be an instance of close contact like having a conversation. Over the course of the day the potential interaction partners change because people tend to be in different locations during the day (e.g. at work or at home). In this example we would include an edge between two people if they are near each other (e.g. in a 50m radius) and there is an interaction if two people have a conversation. In addition to the network and the interactions, one typically also observes a set of covariates on the ‘pair-level’. Such covariates describe the relation of each pair in the network and can be composed of covariates on actor-level, pair specific covariates or external covariates. The interest lies in modeling the relation between the interactions and the covariates. We emphasize already here that the edge set does not necessarily need to describe physical (or otherwise established) relations like in the previous example. It is also possible that one imposes a network on the data, e.g. by specifying which pairs are relevant for the question of interest. In this case, the edges of the network can be understood as model inclusion dummies, indicating at which point in time which pair is relevant for the study. Such scenarios might be relevant when the interest lies on interactions that are only of interest if other conditions are met, such as the use of social media while using the smart-phone.

One possibility to model interaction data on networks is to formulate a counting process model in which the covariates are used to specify a model for the intensity function. This has the advantage that the covariates may depend on their past in an arbitrary way. Thus, in particular, we may consider covariates that depend on the past of the pair of consideration itself or on other pairs of actors. Such models have been studied in both parametric and non-parametric settings, see, for instance, [Perry and Wolfe \(2013\)](#) and [Kreiss et al. \(2019\)](#), respectively. A question of interest is therefore whether there is a need to use a non-parametric model or whether a parametric model is sufficient. In [Kreiss \(2020+\)](#) it has been shown how such a test can be constructed based on a test-statistic which compares the parametric and the non-parametric estimator by means of L^2 distances similar to [Härdle and Mammen \(1993\)](#). A significant short-coming of the test suggested in [Kreiss \(2020+\)](#) is that all the covariates are required to be specified on edge-level. Thus in particular system wide covariates like time, weather, or seasons may not be used. The reason for this is that a constant effect of time cannot be separated from a time varying intercept. For a single covariate there might be work-arounds like combining the intercept and the covariate in one variable, this seems to be very ad-hoc and a rigorous treatment would be preferable. On the other hand, if one has a specific hypothesis about what causes the seasonality, our testing framework provides a methodology to test for this hypothesis.

Being unable to properly include system wide covariates in the testing is restrictive because many data-sets show a clear seasonality and therefore the superiority of a non-parametric model might simply stem from the situation that the parametric model under consideration

does not accurately account for this seasonality. The aim of this paper is to explore whether extending a parametric model by a non-parametric baseline (thereby allowing for a non-parametric seasonality) is producing a meaningful extension of the model. To be more specific we suppose that the edge-specific covariates, stochastic processes $X_{n,ij}$ on $[0, T]$, have a parametric (constant over time) impact on the interactions. We then test whether a parametric specification of the baseline through system-wide covariates, deterministic processes Z (which are the same for all pairs of actors), is sufficient for the data or if a non-parametric specification of the baseline increases the model fit.

The complex dependence structure in network settings makes the mathematics behind such types of analysis significantly more challenging. While in standard situations individuals are typically considered to behave independently, such an assumption is rarely plausible in a networks set-up. Most of the time it is rather the case that neighboring individuals influence each other and should therefore not be treated as independent. On the other hand it is intuitively clear that this dependence diminishes exponentially as the distance of actors grows in the network: If you are equally influenced by your k friends and they, in turn, are equally influenced by their k friends, the impact of the friends of your friends (i.e. of actors of distance 2) on you, is of the order k^{-2} . Moreover, we need to assume that there is no single actor who influences a considerable fraction of the actors (so called hubs). Intuitively it is also clear that in a situation where everybody is copying one person, there is no new information from observing new people. This intuition has been made precise in [Kreiss \(2020+\)](#) and we make use of these ideas.

The organization of this paper is as follows: In Section 2 we introduce the exact model and present the hypothesis and our suggested test statistic. Afterwards in Section 3 we present the theory for our test statistic. The test is applied to real-world data in Section 4. Section 5 concludes. Proofs and other technical details are collected in the Appendix Section 6 and the supplementary Section 7.

2 Model Specification and Test Strategy

In the following we introduce the exact data generating process considered throughout the paper, and formulate the testing problem of interest. Suppose that we observe a sequence of networks $G_{n,t} = (V_n, E_{n,t})$ for $n \in \mathbb{N}$ over a time interval $t \in [0, T]$, where the deterministic population V_n is of growing size $|V_n| = n$. The edge set $E_{n,t} \subseteq V_n \times V_n$ is random and time varying. For $i, j \in V_n$, the functions $C_{n,ij}(t) = \mathbb{1}((i, j) \in E_{n,t})$ indicate whether i and j are connected at time $t \in [0, T]$. $C_{n,ij}$ can also be understood as the random, time varying adjacency matrix. Within the population, individuals can interact with each other when they are connected by an edge. For $i, j \in V_n$ we denote by $N_{n,ij}(t)$ the number of interactions up to and including time $t \in [0, T]$. The processes $N_{n,ij} : [0, T] \rightarrow \mathbb{N}$ are thus counting processes. While networks can be directed or undirected, we only consider undirected networks for simplicity. Thus, we assume throughout $C_{n,ij} = C_{n,ji}$ as well as $N_{n,ij} = N_{n,ji}$ for all $i, j \in V_n$. Moreover, also for simplicity, we exclude self-interactions,

meaning that we set $C_{n,ii} \equiv 0$ and $N_{n,ii} \equiv 0$ for all $i \in V_n$. We suppose that the network process (via $C_{n,ij}$) and the interactions (via $N_{n,ij}$) are observed. In addition to these we also observe random covariates $X_{n,ij} : [0, T] \rightarrow \mathbb{R}^p$ which are specific for the pair $i, j \in V_n$. Throughout we will assume that the array $(N_{n,ij})_{i,j \in V_n}$ forms a multivariate counting process with respect to a filtration $(\mathcal{F}_t^n)_{t \in [0, T]}$. If not further specified all counting processes and martingales are understood to be defined with respect to this filtration. A particular consequence of this assumption is that no two counting processes jump at the same time (with probability one). We suppose that the covariates and the interactions are connected through the intensity functions $\lambda_{n,ij} : [0, T] \rightarrow [0, \infty)$ for which we assume the proportional hazards model (cf. Andersen et al. (1993); Martinussen and Scheike (2006); Cox (1972); Andersen and Gill (1982)), that is, we suppose that the intensity function with respect to the filtration $(\mathcal{F}_t^n)_{t \in [0, T]}$ is given by

$$\lambda_{n,ij}(t) = C_{n,ij}(t)\alpha_0(t)\Psi(X_{n,ij}(t); \beta_0), \quad (2.1)$$

where $X_{n,ij}(t) \in \mathbb{R}^p$ are random covariates depending on time, $\beta_0 \in \mathbb{R}^q$ is an unknown parameter and $\alpha_0 : [0, T] \rightarrow [0, \infty)$ is an unknown, deterministic baseline intensity. The link function Ψ is supposed to be known to the researcher, e.g. in a Cox-type model $p = q$ and $\Psi(X_{n,ij}(t), \beta_0) = \exp(\beta_0' X_{n,ij}(t))$ (cf. Scheike and Martinussen (2004)). In order to assure identifiability, Ψ (or $X_{n,ij}$) may not include an intercept, and we have to impose, e.g. $\Psi(0; \beta_0) = 1$. The intercept and the deviation from the reference intensity is specified by α_0 . Our interest then lies in testing if the baseline α_0 can be adequately modeled by deterministic system wide covariates, i.e., covariates that are the same for all individuals, such as time, weather or economic development. We denote these covariates by $Z : [0, T] \rightarrow \mathbb{R}^d$. By ‘deterministic’ we here mean ‘measurable with respect to \mathcal{F}_0^n ’. The covariates Z may also contain basis expansions of single covariates. Since the covariates Z are supposed to be deterministic and to be the same for the entire network (regardless of its size), we assume also that they do not change with n . Consider the following example: If we consider the weather in a city as a covariate, then it is plausible that the weather is not depending on which network in the city we study. Therefore we may condition on the weather and keep it fixed for our analysis. Our aim is testing the hypothesis

$$H_0 : \alpha_0(t) = \alpha(Z(t); \theta_0; t) \text{ for some } \theta_0 \in \Theta,$$

where $\Theta \subseteq \mathbb{R}^d$ is a suitable parameter space and $\alpha : \mathbb{R}^d \times \Theta \times [0, T]$ is a known link function. To simplify notation, we let $\alpha(\theta, t) := \alpha(Z(t); \theta; t)$. The test statistic we use to test for this hypothesis is along the lines of Hardle and Mammen (1993), that is, we compare a parametric and a non-parametric estimator. In order to define those estimators, we need the following definitions

$$\begin{aligned} N_n(t) &:= \sum_{i,j \in V_n} N_{n,ij}(t), & X_n(t; \beta) &:= \sum_{i,j \in V_n} C_{n,ij}(t)\Psi(X_{n,ij}(t); \beta) \\ C_n(t) &:= \mathbb{1}(\exists i, j \in V_n : C_{n,ji}(t) = 1), & \lambda_n(t, \beta) &:= \alpha_0(t)X_n(t, \beta), \end{aligned}$$

where the above sums over $(i, j) \in V_n$ are understood as *Sum over all undirected pairs* $(i, j) \in V_n \times V_n$ with $i \neq j$. We will use this notation throughout the paper. The process N_n is again a counting process with respect to $(\mathcal{F}_t^n)_{t \in [0, T]}$ because by assumption no two individual processes jump at the same time. The intensity function of N_n is given by $\lambda_n(t, \beta_0)$. The function C_n is an indicator which equals 1 at time t if there is at least one edge present in the network. We denote by $M_{n,ij}(t) := N_{n,ij}(t) - \int_0^t \lambda_{n,ij}(s) ds$ the martingale associated with the counting process $N_{n,ij}$. Then, $M_n := \sum_{i,j \in V_n} M_{n,ij}$ is the martingale associated with N_n (all with respect to $(\mathcal{F}_t^n)_{t \in [0, T]}$).

The estimators used for our test statistic are as follows. Our parametric estimator is the maximum likelihood estimator

$$\left(\hat{\theta}_n, \hat{\beta}_n\right) := \operatorname{argmax}_{(\theta, \beta)} \sum_{i,j \in V_n} \int_0^T \log(\alpha(\theta, t) \Psi(X_{n,ij}(t); \beta)) dN_{n,ij}(t) - \int_0^T \alpha(\theta, t) X_n(t; \beta) dt.$$

For the non-parametric estimator, we first use an initial estimator for β_0 not depending on any specific form of α_0 . For this we take the partial-maximum likelihood estimator (cf. [Cox \(1975\)](#); [Perry and Wolfe \(2013\)](#)).

$$\tilde{\beta}_n := \operatorname{argmax}_{\beta} \sum_{i,j \in V_n} \int_0^T \log \Psi(X_{n,ij}(t); \beta) - \log(X_n(t; \beta)) dN_{n,ij}(t). \quad (2.2)$$

With this, our non-parametric estimator is the smoothed Nelson-Aalen estimator (cf. [Andersen et al. \(1993\)](#); [Aalen \(1978\)](#); [Nelson \(1969\)](#); [Ramlau-Hansen \(1983\)](#))

$$\hat{\alpha}_n(t, \tilde{\beta}_n) := \int_0^T K_{h,t}(s) \frac{C_n(s)}{X_n(s; \tilde{\beta}_n)} dN_n(s), \quad (2.3)$$

where $K : \mathbb{R} \rightarrow [0, \infty)$ is a kernel function and $K_{h,t}(s) := \frac{1}{h} K\left(\frac{s-t}{h}\right)$. Above we use the convention $0/0 := 0$. Let further

$$\alpha_{0,s}(t) := \int_0^T K_{h,t}(s) \alpha_0(s) ds \quad \text{and} \quad \alpha_s(\theta, t) := \int_0^T K_{h,t}(s) \alpha(\theta, s) ds \quad \text{for all } \theta \in \Theta,$$

be smoothed versions of the true baseline and the parametric estimator, respectively. The smoothing is necessary because the non-parametric estimator is inherently biased due to the smoothing. Hence, in order to compare the non-parametric and the parametric estimators, we need to add this bias also to the parametric estimator (cf. [Härdle and Mammen \(1993\)](#)). In order to test the hypothesis H_0 we therefore study the following test statistic

$$T_n := \int_0^T \left(\hat{\alpha}_n(t; \tilde{\beta}_n) - \alpha_s(\hat{\theta}_n, t)\right)^2 w(t) dt, \quad (2.4)$$

where $w : [0, T] \rightarrow [0, \infty)$ is a weight function with $\mathbb{T} := \operatorname{supp} w \subset (0, T)$, thus w cuts off the boundary and therefore we may ignore possible boundary issues of the kernel type Nelson-Aalen estimator.

3 Main Results

In this section we present the main theoretical result of the paper. We begin by stating our main result on the asymptotic behavior of T_n under the null-hypothesis and under local alternatives (Section 3.1). The assumptions needed for this result to hold are presented and discussed in detail in Section 3.2.

3.1 Main Result

In order to state our main result, we introduce the following notation (by Assumption (ID) below, all these quantities are well defined):

$$\bar{X}_n(t; \beta) := \mathbb{E}(\Psi(X_{n,ij}(t); \beta) | C_{n,ij}(t) = 1), \quad (3.1)$$

$$p_n(t) := \mathbb{P}(C_{n,ij}(t) = 1), \quad (3.2)$$

$$a_n := \left(\int_0^T \int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} w(t) ds dt \right)^{-\frac{1}{2}}. \quad (3.3)$$

where $r_n = n(n-1)/2$ denotes the total number of undirected pairs.

Theorem 3.1. *Let all conditions from Section 3.2 hold. Suppose that the data is generated by model (2.1) with $\alpha_0(t) = \alpha(\theta_0; t) + c_n \Delta_n(t)$, where $c_n = a_n^{-1} h^{-1/4}$ and Δ_n is uniformly bounded and continuously differentiable with uniformly bounded derivative (that is, in both t and n). When $\Delta_n \equiv 0$ we are on the null hypothesis H_0 . Then,*

$$a_n^2 \sqrt{h} T_n - A_n / \sqrt{h} - \int_0^T \left(\int_0^T K_{h,t}(s) \Delta_n(s) ds \right)^2 w(t) dt \xrightarrow{d} \mathcal{N}(0, B),$$

where

$$A_n := a_n^2 \sum_{i,j \in V_n} \int_0^T f_n(r, r) \left(\frac{1}{r_n p_n(r)} \bar{X}_n(r, \beta_0)^{-1} \right)^2 dN_{n,ij}(r),$$

$$B := 4K^{(2)} \int_0^T \bar{X}_n(s; \beta_0)^{-2} w(s)^2 \alpha(\theta_0, s)^2 \gamma(s) ds,$$

$$K^{(2)} := \int_0^2 \left(\int_{-1}^1 K(u+v) K(u) du \right)^2 dv,$$

$$f_n(r, s) := \int_0^T h K_{h,t}(s) K_{h,t}(r) w(t) dt.$$

In other situations it has been pointed out that the convergence to a normal limit for L^2 type non-parametric tests is very slow. As a remedy, resampling techniques have been proposed for finding critical values. We are not aware of a resampling scheme that is suitable for our model of dependent counting processes and let this question open for future research.

3.2 Assumptions

In this section we collect the assumptions that we use to prove our main result. We begin with fairly standard assumptions on the data.

Assumption (ID): (Identical Distribution)

All quantities which are indexed by $(i, j) \in V_n \times V_n$ are identically distributed.

This assumption is implied by the stronger assumption that relabeling the vertices of the network is not changing the distribution. Since in observational studies names are normally uninformative, we regard this assumption as not very restrictive. A particular consequence of (ID) is that $\bar{X}_n(t; \beta)$ and $p_n(t)$ as defined in (3.1) and (3.2), respectively, are well defined. We continue with standard assumptions on the kernel and the bandwidth:

Assumption (K): (Kernel)

The kernel K and the weight function w are bounded. Moreover, K is supported on $[-1, 1]$ and $\mathbb{T} := \text{supp } w \subset (0, T)$. The bandwidth h fulfills $h \rightarrow 0$, $hr_n p_n \rightarrow \infty$, $\sqrt{h} \log r_n \rightarrow 0$ and $\log r_n / hr_n p_n \rightarrow 0$.

Recall that in our set-up, the observations are the connected pairs in the network, and their expected number is $r_n p_n$. While we do not discuss optimal bandwidth choice, a reasonable choice for h appears to be $h = O((r_n p_n)^{-r})$ for a small power $r > 0$ with the standard choice $r = 1/5$. In this case the above requirements on the bandwidth are true. The assumptions on the kernel hold for example for a triangular kernel.

Assumption (SC) (Sparsity Condition)

There is a bounded function $\gamma(t)$, which is the uniform limit of $\frac{a_n^4}{r_n^2 p_n(t)^2}$, where a_n and $p_n(t)$ are as in (3.1) and (3.3), respectively.

As a motivation of (SC) denote $p_n := \inf_{t \in [0, T]} p_n(t)$ (this definition will be important in other places too). Then, $p_n(t) = p_n \cdot p_n^*(t)$ for some functions $p_n^* : [0, T] \rightarrow [1, \infty)$ (mind the range). By definition of a_n ,

$$\begin{aligned} \frac{a_n^4}{r_n^2 p_n(t)^2} &= \left(\int_0^T \int_0^T K_{h,t}(s) \frac{p_n(t)}{p_n(s)} w(t) ds dt \right)^{-2} \\ &= \left(\int_0^T \int_0^T K_{h,t}(s) w(t) ds dt + \int_0^T \int_0^T K_{h,t}(s) \frac{p_n^*(t) - p_n^*(s)}{p_n^*(s)} w(t) ds dt \right)^{-2}. \end{aligned}$$

Under (K), the first integral above equals $\int_0^T w(t) dt$ for $h > 0$ small enough. The second integral converges to zero under, e.g., an equi-continuity assumption because $p_n^*(s) \geq 1$ for all $n \in \mathbb{N}$ and $s \in [0, T]$. Note also that we explicitly allow that $p_n \rightarrow 0$ so that our set-up includes sparse networks in which the expected number of edges $r_n p_n = O(n)$. In the following, let $B_n(c) := \left\{ \beta : \|\beta - \beta_0\| \leq \frac{c}{\sqrt{r_n p_n}} \right\}$ for an arbitrary constant $c > 0$.

Assumption (C): (Continuity and Boundedness of the Model)

The link functions Ψ and α are bounded, i.e. $\|\Psi(\cdot; \beta_0)\|_\infty < \infty$ and $\|\alpha(\theta_0; \cdot)\|_\infty < \infty$, and

fulfill the following continuity property

$$|\Psi(x, \beta_1) - \Psi(x; \beta_2)| \leq L_\Psi(x)\omega_\Psi(\|\beta_1 - \beta_2\|) \quad \text{and} \quad |\alpha(\theta_1, t) - \alpha(\theta_2, t)| \leq L_\alpha(t)\omega_\alpha(\|\theta_1 - \theta_2\|),$$

where ω_Ψ and ω_α are increasing functions satisfying, for all $c^* > 0$,

$$a_n^2\omega_\Psi(c^*/\sqrt{r_n p_n})^2 = O(1), a_n^2\omega_\alpha(c^*/\sqrt{r_n p_n})^2 = O(1) \quad \text{and} \quad hr_n p_n \omega_\Psi\left(\sqrt{\log r_n/r_n p_n}\right)^2 = o(1).$$

The functions L_Ψ and L_α are non-negative and satisfy, for any $c > 0$,

$$\int_0^T \left(\frac{\mathbb{E}(L_\Psi(X_{n,ij}(s)) | C_{n,ij}(s=1))}{\inf_{\beta \in B_n(c)} \bar{X}_n(s; \beta)^2} \right)^2 ds = O(1) \quad \text{and} \quad \int_0^T L_\alpha(t)^2 dt < \infty.$$

Moreover, $\alpha(\theta_0; \cdot)$, w and $\bar{X}_n(\cdot; \beta_0)$ are continuously differentiable with bounded first derivative. Finally,

$$\sup_{s \in [0, T]} \sup_{r \in [s, s+2h]} \frac{p_n(s)}{p_n(r)} = O(1).$$

Assumption (C) collects continuity and boundedness statements on expectations and probabilities. Intuitively speaking they all mean that the model is not changing too rapidly over time, which is a mild assumption. In view of (SC), we see that the convergence assumptions on the moduli of continuity ω_α and ω_Ψ hold if the link functions are Lipschitz continuous, e.g., if ω_Ψ and ω_α are the identity function. Such an assumption is for example plausible if we assume a continuously differentiable link function (for the true parameters) and bounded covariates. The assumption on $p_n(t)$ can be translated to a continuity assumption on $p^*(t)$ if we assume that $p_n(t) = p_n p^*(t)$. What we really exclude here is that the network density is changing its order, i.e., we cannot have that $p_n(t_0) \approx 1/n$ and $p_n(t_1) \approx 1/\sqrt{n}$.

Assumption (Co): (Covariates)

The conditional expectation of the covariates is bounded from above and below, that is, for any $c > 0$, $\sup_{\beta \in B_n(c)} \|\bar{X}_n(\cdot; \beta)\|_\infty = O(1)$ and $\sup_{\beta \in B_n(c)} \|\bar{X}_n(\cdot; \beta)^{-1}\|_\infty = O(1)$.

The upper bound holds for example if Ψ is bounded and the lower bound ($\|\bar{X}_n(\cdot; \beta_0)^{-1}\|_\infty < \infty$) essentially only means that at no point in time the intensity converges to zero (provided that there is a link). Since we condition on the existence of a link in the definition of \bar{X}_n it might even be plausible to assume that \bar{X}_n does not even depend on n as was argued in [Kreiss et al. \(2019\)](#).

Assumption (P) (Parametric Estimation)

The estimators $\hat{\theta}_n$ and $\tilde{\beta}_n$ are independent of the observations and fulfill $\sqrt{r_n p_n} \|\hat{\theta}_n - \theta_0\| = O_P(1)$ and $\sqrt{r_n p_n} \|\tilde{\beta}_n - \beta_0\| = O_P(1)$ and let for some $c^* > 0$

$$r_n p_n \mathbb{P} \left(\left\| \tilde{\beta}_n - \beta_0 \right\| \geq c^* \sqrt{\frac{\log n}{r_n p_n}} \right) = O(1).$$

The independence assumption is made for simplicity. In simple regression settings such an assumption can be avoided if one has expansions of the estimators around true parameters available which are of approximating order $o_P(1/\sqrt{r_n p_n})$. We expect that with such expansions we can prove our results without the independence assumptions. But this would increase the (already high) amount of technicalities in the proof. A simple case where (P) holds trivially is in the case of testing simple hypotheses in which β_0 and θ_0 are known. There, we can simply put $\tilde{\beta}_n = \beta_0$ and $\hat{\theta}_n = \theta_0$. The assumption also holds if we use data splitting in time. More precisely, let us observe data on an interval $[0, T]$ which has a strong periodicity, like the rental bike data which we will consider in Section 4. Then it is often plausible to assume that the data on two certain time intervals $[a, b] \cap [c, d] = \emptyset$ (both forming one period) is independent due to short memory effects, if they lie far enough apart. We can then use the first interval to do the parametric estimation and the second interval for the non-parametric estimation. Note also that the intervals do not need to have the same length, the interval for the parametric estimator can be chosen to be shorter.

In order to understand the rate assumption, recall that $r_n p_n$ is the effective number of observations. Therefore, we would expect that any reasonable parametric estimator has the rate $1/\sqrt{r_n p_n}$ which is exactly what we assume here.

We turn now to assumptions that are more difficult to interpret. For any $c_1, c_2 > 0$, define the following events (recall that $B_n(c) := \{\beta : \|\beta - \beta_0\| \leq c/\sqrt{r_n p_n}\}$):

$$\mathcal{A}_n(c_1, c_2) := \left\{ \sup_{t \in [0, T], \beta \in B_n(c_2)} \sqrt{p_n(t)} \left| \left(\frac{1}{r_n p_n(t)} X_n(t; \beta) \right)^{-1} - \bar{X}_n(t; \beta)^{-1} \right| \leq c_1 \sqrt{\frac{\log r_n}{r_n}} \right\}$$

$$\tilde{\mathcal{A}}_n(c_1, c_2) := \left\{ \sup_{t \in [0, T], \beta \in B_n(c_2)} \sqrt{p_n(t)} \left| \frac{1}{r_n p_n(t)} X_n(t; \beta) - \bar{X}_n(t; \beta) \right| \leq c_1 \sqrt{\frac{\log r_n}{r_n}} \right\}.$$

where $\bar{X}_n(t; \beta)$ was defined as $\bar{X}_n(t; \beta) := \mathbb{E}(\Psi(X_{n,ij}(t), \beta) | C_{n,ij}(t) = 1)$.

Assumption (CC): (Concentration of Covariates)

For any $c_2 > 0$ and any $\delta > 0$ there is $c_1 > 0$ such that for all $n \in \mathbb{N}$

$$\mathbb{P}(\mathcal{A}_n(c_1, c_2)) \geq 1 - \delta \quad \text{and} \quad \mathbb{P}(\tilde{\mathcal{A}}_n(c_1, c_2)) \geq 1 - \delta$$

and for any $c > 0$

$$\sup_{s \in [0, T], \beta \in B_n(c)} \mathbb{E} \left(\left(\frac{X_n(s, \beta)}{r_n p_n(s)} \right)^2 \right) = O(1).$$

Assumption (CC) essentially requires that the hazards $\Psi(X_{n,ij}(t), \beta)$ concentrate around their mean. In Section 6.2 we will discuss how this assumption can be proven by assuming certain mixing assumptions as introduced in Kreiss (2020+). Intuitively speaking, it can be

shown that (CC) holds whenever we are comfortable to assume that there is an exponential decay in the dependence if the distance in the network grows. This is for example the case if one considers exponentially growing friendship circles (friends, friends of friends, friends of friends of friends, ...).

In order to motivate why the last set of assumptions is necessary, we consider the following. Let $\varepsilon_1, \dots, \varepsilon_n$ be iid random variables with finite variance and zero expectation and let $X_{n,i}$ be further random variables. Our aim is to show that $1/\sqrt{n} \sum_{i=1}^n X_{n,i} \varepsilon_i = o_P(1)$. This clearly holds if $X_{n,i}$ is deterministically bounded by a sequence $\delta_n \rightarrow 0$ and if $(X_{n,i} \varepsilon_i)_{i=1, \dots, n}$ is an iid sequence with $\mathbb{E}(\varepsilon_i | X_{n,i}) = 0$. But if we drop the latter, i.e. if we allow that $X_{n,i}$ might be correlated with all $\varepsilon_1, \dots, \varepsilon_n$ at the same time, we have a harder time to handle the sum:

$$\mathbb{E} \left(\left(\frac{1}{\sqrt{n}} \sum_{i=1}^n \varepsilon_i X_{n,i} \right)^2 \right) = \left| \frac{1}{n} \sum_{i,j=1}^n \mathbb{E}(X_{n,i} X_{n,j} \varepsilon_i \varepsilon_j) \right| \leq \delta_n^2 n \mathbb{E}(\varepsilon_1^2).$$

So even $\delta_n = 1/\sqrt{n}$ is not sufficient to let the sum go to zero. However, we have a chance if $X_{n,i}$ is only weakly correlated with ε_j for $j \neq i$ and if there is $\tilde{X}_{n,i}$ which is even uncorrelated with ε_j for $j \neq i$ and such that $X_{n,i} - \tilde{X}_{n,i}$ is very small. Such ideas have been transferred to counting processes by [Mammen and Nielsen \(2007\)](#) and to counting processes on networks in [Kreiss \(2020+\)](#). Suppose to this end that we have a random distance function $d_t^n : V_n^2 \times V_n^2 \rightarrow [0, \infty)$ indexed by time t and the number of actors n , assigning a distance to all pairs. For $(i, j) \in V_n \times V_n$ and $I \subseteq V_n \times V_n$, we define $d_t^n(ij, I) := \inf_{(k,l) \in I} d_t^n(ij, kl)$ and by definition $d_t^n(ij, \emptyset) := \infty$. In order to be able to define the correct analogue to uncorrelatedness as mentioned above we introduce the following augmented σ -field for any sets $I, J \subseteq V_n \times V_n$ and any $m > 0$, where we denote by $\mathcal{F}_1 \wedge \mathcal{F}_2$ the σ -field generated by the union of the σ -fields \mathcal{F}_1 and \mathcal{F}_2 :

$$\begin{aligned} \tilde{\mathcal{F}}_{I,t}^{n,J,m} &:= \sigma(X_{n,ij}(\tau) : (i, j) \in I, \tau \leq t) \\ &\wedge \sigma(\mathcal{N}_{n,ij}(r) \mathbb{1}(d_s^n((i, j), J) \geq m) : s \leq \max(0, t - 4h), r \leq s + 6h, (i, j) \in V_n \times V_n), \\ \mathcal{N}_{n,ij}(r) &:= (N_{n,ij}(r), C_{n,ij}(r), C_{n,ij}(r)). \end{aligned}$$

Intuitively speaking the above filtration contains all information of the covariates belonging to pairs in I and it contains extra (future) information about pairs that are far from J . We construct now functions using exactly this information. Recall that $f_n(r, s) = \int_0^T h K_{h,t}(s) K_{h,t}(r) w(t) dt$. Note that $f_n(r, s)$ is symmetric. Moreover, $f_n(r, s)$ functions as a kernel, and if K is supported on $[-1, 1]$ it is different from zero only if $|r - s| \leq 2h$. For $s_2 \leq s_1$, $(i, j), (k, l) \in V_n \times V_n$ and $I \subseteq V_n \times V_n$ we consider the following function:

$$\begin{aligned} &\varphi_{n,ij,kl}^I(s_1, s_2) \\ &:= \frac{4r_n a_n^4}{h} \int_{s_1}^{s_2+2h} f_n(r, s_1) f_n(r, s_2) \frac{\bar{X}_n(r; \beta_0)^{-2}}{r_n p_n(r)} \alpha_0(r) \frac{\bar{X}_n(s_1; \beta_0)^{-1}}{r_n p_n(s_1)} \frac{\bar{X}_n(s_2; \beta_0)^{-1}}{r_n p_n(s_2)} \end{aligned} \quad (3.4)$$

$$\times \frac{1}{r_n p_n(r)} \sum_{\substack{(i',j') \in V_n \times V_n \\ (i',j') \neq (i,j),(k,l)}} \Psi(X_{n,i'j'}(r); \beta_0) C_{n,i'j'}(r) \mathbb{1}(d_{s_1-4h}^n(i'j', I) \geq m) dr$$

We will write $\varphi_{n,ij}^I(r)$ instead of $\varphi_{n,ij,ij}^I(r, r)$ and let $\varphi_{n,ij,kl}(s_1, s_2) := \varphi_{n,ij,kl}^\emptyset(s_1, s_2)$. The functions $\varphi_{n,ij,kl}(s_1, s_2)$ will be our $X_{n,i}$ in the initial example and $\varphi_{n,ij,ij}^I(s, r)$ will be the approximation. We see that $t \mapsto \int_{t-2h}^t \varphi_{n,ij,kl}^I(t, s_2) ds_2$ is predictable with respect to $\tilde{\mathcal{F}}_{ij,kl,t}^{n,I,m}$. In order for the approximating functions to be uncorrelated with the other variables, we make the following assumption.

Assumption (mDep): (Momentary- m -Dependence)

There is a number $m > 0$ such that

$$\begin{aligned} \forall n \in \mathbb{N}, \forall t_0 \in [0, T], \forall J \subseteq V_n \times V_n : \text{ Given } \mathcal{F}_{t_0}^n, \\ (\mathcal{N}_{n,ij}(t))_{(i,j) \in J, t \in [t_0, t_0+6h]} \text{ is cond. independent of} \\ \sigma(\mathcal{N}_{n,ij}(r) \mathbb{1}(d_s^n(ij, J) \geq m) : s \leq t_0, r \leq s + 6h, (i, j) \in V_n \times V_n). \end{aligned}$$

Intuitively speaking, the above assumption means that, conditional on the past, the immediate short future of far apart processes can be treated as if they were independent. This is plausible if information needs some time to *travel* through the network. For a more detailed discussion and an example under which this assumption holds we refer to [Kreiss \(2020+\)](#). To finish the analogy, we have to discuss why the approximation is good enough. To this end we need to introduce the concept of a hub in a network. For a subset $A \subseteq V_n \times V_n$ we let

$$K_m^A := \sup_{t \in [0, T]} \sup_{(k,l) \in A} \sum_{i,j \in V_n} \mathbb{1}(d_{t-4h}^n(ij, kl) < m) \sup_{r \in [t-4h, t+2h]} C_{n,ij}(r)$$

be the maximal number of close neighbor pairs of any of the pairs in A . For a given $F > 0$, we call a pair (i, j) a hub if $K_m^{ij} \geq F$, and any random variable H_{UB}^A with

$$H_{UB}^A \geq \mathbb{1}(K_m^A \geq F),$$

the hub-ability of the set A . With these definitions, the following Lemma is shown in the Appendix (cf. Lemma 6.2):

Lemma

Under the assumptions of Theorem 3.1 there is a constant $C > 0$ and a random function $\Phi_n : [0, T] \rightarrow [0, \infty)$ such that for $I \subseteq J \subseteq V_n \times V_n$

$$\begin{aligned} |\varphi_{n,ij,kl}^I(t, r) - \varphi_{n,ij,kl}^J(t, r)| &\leq |J \setminus I| \cdot \frac{C}{r_n p_n(t) p_n(r)} B_n^{J \setminus I}, \\ |\varphi_{n,ij,kl}^I(t, r)| &\leq \frac{1}{p_n(r)} \Phi_n(t), \end{aligned}$$

where $B_n^A := (F + H_{UB}^A K_m^{V_n \times V_n})$ for $A \subseteq V_n \times V_n$.

The number B_n^A describes a trade-off between the size of a hub $K_m^{V_n \times V_n}$ and the likelihood of its appearance H_{UB}^A . Note moreover that the function Φ_n behaves like $X_n(\cdot; \beta)$ and thus in view of Assumption (CC) we may expect that it concentrates around its expectation. These considerations show that the following two assumptions are requirements on the trade-off between hub-size and frequency.

Assumption (HSR1): (Hub Size Restriction I)

Let

$$\begin{aligned} \mathbb{E} \left(\left(\sum_{ij} \int_0^T \frac{B_n^{ij}}{r_n p_n(t)} \right)^2 \sup_{t \in [0, T]} \Phi_n(t) \right) &= o(r_n p_n), \\ \mathbb{E} \left(\left(\sum_{ij} \int_0^T \frac{(B_n^{ij})^2}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \right) &= o(r_n^2 p_n^2), \\ \mathbb{E} \left(\sum_{ij} \int_0^T \frac{C_{n,ij}(t)}{r_n p_n(t)} dt \sup_{t \in [0, T]} \Phi_n(t)^2 \right) &= o(r_n p_n), \end{aligned}$$

Suppose moreover that the approximations $\varphi_{n,ij,kl}^I$ fulfill the conditions conditions (2.4)-(2.8) in [Kreiss \(2020+\)](#) with $\delta_n = h$.

The assumptions (2.4)-(2.8) are essentially requirements on the quality of approximation of the functions $\varphi_{n,ij,kl}^I$. The main problem is that they involve complicated sums of martingale integrals that are difficult to handle. However, their domain of integration has length of order $h \rightarrow 0$ and their integrands may be bounded by using [Lemma 6.2](#) (as above). Therefore, assuming $K_m^{V_n \times V_n}$ to be deterministic and slowly growing relative to $\mathbb{P}(H_{UB}^A = 1)$ together with integrability assumptions similar to the ones in (HSR1), will allow to prove (2.4)-(2.8) in [Kreiss \(2020+\)](#).

Assumption (HSR2): (Hub Size Restriction II)

Let

$$\begin{aligned} \int_0^T \mathbb{E} \left(\left(\sup_{r \in [s, s+2h]} \frac{1}{r_n p_n(r)} \sum_{k,l \in V_n} C_{n,kl}(r) \right)^2 \middle| C_{n,ij}(s) = 1 \right) ds &= O(1), \\ \mathbb{E} \left(\sup_{r \in [0, T]} \sum_{k,l \in V_n} \frac{C_{n,kl}(r)}{r_n p_n(r)} \left(\sum_{ij \in V_n} \int_0^T \frac{B_n^{ij}}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \right) &= O(1), \\ \mathbb{E} \left(\left(\sum_{ij \in V_n} \int_0^T \frac{(B_n^{ij})^2}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \right) &= O(r_n p_n). \end{aligned}$$




Figure 1: Kernel density estimate for bike a rides over a period of two weeks in 2018 (May 13 - May 26).

4 Empirical Application

¹ Here we apply the methodology from Section 2 to bike sharing data. More precisely, the data is based on 527 bike stations in Washington D.C. and its surrounding. We consider these bike stations to be the vertices of a network. An interaction from bike station i to bike station j happens if someone rents a bike at i and returns it at j . We consider such data from May 13, 2018 (Sunday) to May 26, 2018 (Saturday). The data is publicly available and can be downloaded from <https://www.capitalbikeshare.com/system-data>. Since many of the connections are rarely (or never) used, we attempt to only model the frequently used connections. To this end we construct a network of active pairs as follows: There is a link from bike station i to station j if there were at least ten bike rides in April 2018 from i to j . So we only consider pairs that were active at least twice a week on average over a period of one month. This convention is somewhat arbitrary and a full analysis would require a sensitivity analysis with respect to this choice. Also note that this is a directed network. This is no problem because the result in Theorem 3.1 holds analogously for directed networks.

Figure 1 shows a kernel density estimate of the times at which bike rides happened over the period from May 13 till May 26. The scale of the x-axis are days, with day 1 and 8

¹The code for this example is available at <https://github.com/akreiss/Baseline-Estimation.git>.

being Sundays and days 7 and 14 being Saturdays. One clearly sees a different pattern for weekdays and weekends. Moreover it appears that in the second week there were more bike rides than in the first week. One might suspect that this is due to the weather situation: Figure 2 shows that there was more rain in the first week than in the second week. The weather information is publicly available from <https://www.wunderground.com/> and was collected at the Ronald Reagan Airport in Washington D.C. Even though the Washington D.C. area is large, it is plausible that this local weather information is a valid indicator for the weather in the entire region. Therefore, the temperature and the precipitation are system-wide covariates that are identical for all pairs. Let $T(t)$ denote the log of the temperature at time t and let $P(t)$ denote the precipitation. As a parametric model for α_0 we consider the following ($\theta = (\theta_1, \dots, \theta_{17})$)

$$\begin{aligned} \alpha(t, \theta) = & \exp \left(\theta_1 + (T(t) \quad T(t)^2)' \begin{pmatrix} \theta_2 \\ \theta_3 \end{pmatrix} + (P(t) \quad P(t)^2)' \begin{pmatrix} \theta_4 \\ \theta_5 \end{pmatrix} \right. \\ & + \begin{pmatrix} \sin \left(t \frac{2\pi}{24} \right) \\ \sin \left(2t \frac{2\pi}{24} \right) \\ \sin \left(3t \frac{2\pi}{24} \right) \end{pmatrix}' \begin{pmatrix} \theta_6 \\ \theta_7 \\ \theta_8 \end{pmatrix} + \begin{pmatrix} \cos \left(t \frac{2\pi}{24} \right) \\ \cos \left(2t \frac{2\pi}{24} \right) \\ \cos \left(3t \frac{2\pi}{24} \right) \end{pmatrix}' \begin{pmatrix} \theta_9 \\ \theta_{10} \\ \theta_{11} \end{pmatrix} \\ & \left. + W(t) \begin{pmatrix} \sin \left(t \frac{2\pi}{24} \right) \\ \sin \left(2t \frac{2\pi}{24} \right) \\ \sin \left(3t \frac{2\pi}{24} \right) \end{pmatrix}' \begin{pmatrix} \theta_{12} \\ \theta_{13} \\ \theta_{14} \end{pmatrix} + W(t) \begin{pmatrix} \cos \left(t \frac{2\pi}{24} \right) \\ \cos \left(2t \frac{2\pi}{24} \right) \\ \cos \left(3t \frac{2\pi}{24} \right) \end{pmatrix}' \begin{pmatrix} \theta_{15} \\ \theta_{16} \\ \theta_{17} \end{pmatrix} \right), \end{aligned}$$

where $W(t)$ is an indicator function which equals 1 if the t lies on a weekend. Since the weather data is only available for every hour, we consider the functions $P(t)$ and $T(t)$ as piece-wise constant. Note that the essential assumption of continuity of the link function is not violated because we only require continuity in the parameter. The constant $L_\alpha(t)$ is bounded because temperature and precipitation as well as the other functions are bounded. The differentiability required in Assumption (C) is more of an issue. However, in the limit this is not a problem because the differentiability is used for kernel approximations, thus these approximations are only problematic at the discontinuity. Since we integrate over the entire observation period, these approximation errors on short intervals are not of big importance.

We consider covariates on pair-wise level based on distances. Let $d_{i,j}$ denote the logarithm of the biking time in minutes from station i to station j as returned by Google maps. Then, we consider a simple bi-variate covariate vector

$$X_{n,ij} = (d_{i,j} \quad d_{i,j}^2)'$$

As a link function we consider $\Psi(x; \beta) = \exp(x'\beta)$ and hence we have a Cox proportional hazards model. We compute the estimates $\hat{\beta}_n$ and $(\hat{\theta}_n, \hat{\beta}_n)$ as it was described in Section 2 based on the first week, i.e, from May 13 to May 19. The estimates $\hat{\beta}_n$ and $\hat{\theta}_n$ are almost




Figure 2: Temperature and Precipitation in Washington D.C.




Figure 3: Parametric (P) and Non-parametric (NP) estimates of the baseline intensity for the second week.

identical and have roughly the values $(0.219, -0.147)'$. The negative sign of the parameter for $d_{i,j}^2$ shows that the intensity (as a function of the distance) has a maximum. This is plausible because people will likely not use the bikes for very short or very large tours. Now we compute the non-parametric Nelson-Aalen estimator of the baseline intensity based on the second week and compare the estimate with the parametric estimator. Here we have chosen the bandwidth to be 30 min. This has been obtained by *eye-balling*. The assumption of the independence of the two weeks is plausible because we expect that people do not base their biking decisions on the past week other than possibly indirectly through the weather which we control for.

Both estimators are shown in Figure 3. We can see that in particular the drop in bike rides on Tuesday (Day 3) (possibly due to the rain) is well captured by the parametric model. Also the difference between weekends and weekdays is well visible in both estimators. When computing our test statistic, we obtain a value $T_n \approx 0.04868$ which appears small but it should be noted that the scale of the estimators is very low too. We estimate A_n and B by plugging in the sample quantities (and estimates for α and β) and obtain $a_n \approx 7.41666$, $A_n \approx 0.00933$ and $B \approx 0.00003$. Thus, if the hypothesis holds, the test static is very large. Thus, we would reject the hypothesis of the parametric model on all conventional significance levels.

5 Conclusion

In this paper we have applied weak dependence assumptions on networks in order to be able to prove the asymptotic behavior of a test static which can be used to test a parametric model of baseline specification. We have shown an asymptotic distribution for the test statistic which can be used to obtain critical values. Moreover, from a mathematical point of view, we have illustrated how weak dependence assumptions on networks can be used in order to prove asymptotic results for counting processes on networks. These assumptions allow for a certain dependence between neighbors in a network but require that the dependence is weak for vertices which are far apart in the network. Such assumptions allow a rigorous study of counting processes on networks under realistic assumptions.

As an illustration we have applied the test to bike share data. We saw that even though the weather and time information used by the model, appear to explain the data well, there is still some structure that is not unraveled by the particular model under study. This motivates further study in the direction of modeling interactions on networks. As a direct consequence, an interesting further research question would be to test if the fully non-parametric model from [Kreiss et al. \(2019\)](#) offers needed flexibility or if a hybrid model like it was studied here offers enough flexibility. In addition, a thorough treatment of the bandwidth would be desirable.

6 Appendix

6.1 Proof of Theorem 3.1

In this Section, we assume throughout that all conditions of Theorem 3.1 hold. However, in order to make it more clear for the reader which of the assumptions are important for which step, we formulate each lemma and each proposition with its own set of assumptions. While doing this, we focused only on the assumptions not universally necessary. So we do not mention for every lemma that we assume the model from Theorem 3.1 as well as Assumption (ID) and the fact that the kernel is non-negative and integrates to 1. All assumptions made in the lemmas and propositions of this subsection are contained in the set of conditions made for Theorem 3.1.

Before starting with the proof of Theorem 3.1 we make the following simple observation on the continuity of \bar{X}_n which will be useful in several places.

Lemma 6.1. *Suppose that $|\Psi(x; \beta_1) - \Psi(x; \beta_2)| \leq L_\Psi(x)\omega_\Psi(\|\beta_1 - \beta_2\|)$. Then,*

$$|\bar{X}_n(t; \beta_1) - \bar{X}_n(t; \beta_2)| \leq \mathcal{L}_n(t)\omega_\Psi(\|\beta_1 - \beta_2\|),$$

where $\mathcal{L}_n(t) := \mathbb{E}(L_\Psi(X_{n,ij}(t)) | C_{n,ij}(1) = 1)$.

Proof of Lemma 6.1. We have

$$\begin{aligned} |\bar{X}_n(t; \beta_1) - \bar{X}_n(t; \beta_2)| &\leq \mathbb{E}(|\Psi(X_{n,ij}(t), \beta_1) - \Psi(X_{n,ij}(t), \beta_2)| | C_{n,ij}(1) = 1) \\ &\leq \mathbb{E}(L_\Psi(X_{n,ij}(t)) | C_{n,ij}(1) = 1) \omega_\Psi(\|\beta_1 - \beta_2\|) \end{aligned}$$

□

The general outline of the proof is simple. However there are many technical details that are very tedious to handle. We provide a comprehensive treatment of all the details in the supplementary Section 7 and show here the main steps only. Note that $C_n(t) = 1$ if one of the processes $N_{n,ij}$ jumps at time t . We begin by rewriting the integrand in the test statistic in the following way

$$\begin{aligned} &\hat{\alpha}_n(t; \tilde{\beta}_n) - \alpha_s(\hat{\theta}_n, t) \\ &= \int_0^T K_{h,t}(s) \frac{1}{X_n(s; \beta_0)} dN_n(s) - \int_0^T K_{h,t}(s) \alpha(\hat{\theta}_n, s) ds \\ &\quad + \int_0^T K_{h,t}(s) \left(\frac{1}{X_n(s; \tilde{\beta}_n)} - \frac{1}{X_n(s; \beta_0)} \right) dN_n(s) \\ &= \int_0^T K_{h,t}(s) \frac{1}{X_n(s; \beta_0)} dM_n(s) + \int_0^T K_{h,t}(s) \left(\alpha(\theta_0, s) - \alpha(\hat{\theta}_n, s) + c_n \Delta_n(s) \right) ds \\ &\quad + \int_0^T K_{h,t}(s) \left(\frac{1}{X_n(s; \tilde{\beta}_n)} - \frac{1}{X_n(s; \beta_0)} \right) dN_n(s) \end{aligned}$$

$$=I_1(t) + I_2(t) + I_3(t) + I_4(t),$$

where

$$I_1(t) := \int_0^T K_{h,t}(s) \frac{1}{\bar{X}_n(s; \beta_0)} dM_n(s), \quad (6.1)$$

$$I_2(t) := c_n \int_0^T K_{h,t}(s) \Delta_n(s) ds. \quad (6.2)$$

$$I_3(t) := \int_0^T K_{h,t}(s) (\alpha(\theta_0, s) - \alpha(\hat{\theta}_n, s)) ds, \quad (6.3)$$

$$I_4(t) = \int_0^T K_{h,t}(s) \frac{X_n(s; \beta_0) - X_n(s; \tilde{\beta}_n)}{X_n(s; \tilde{\beta}_n) X_n(s; \beta_0)} dN_n(s). \quad (6.4)$$

With this notation, we obtain that

$$\begin{aligned} a_n^2 \sqrt{h} T_n &= a_n^2 \sqrt{h} \int_0^T \left(\hat{\alpha}_n(t; \tilde{\beta}_n) - \alpha_s(\hat{\theta}_n, t) \right)^2 w(t) dt \\ &= a_n^2 \sqrt{h} \int_0^T I_1(t)^2 w(t) dt + a_n^2 \sqrt{h} \int_0^T I_2(t)^2 w(t) dt + \sum_{i=3}^4 a_n^2 \sqrt{h} \int_0^T I_i(t)^2 w(t) dt \\ &\quad + a_n^2 \sqrt{h} \sum_{\substack{i,j=1 \\ i \neq j}}^4 \int_0^T I_i(t) I_j(t) w(t) dt. \end{aligned}$$

Now, Lemmas 7.2-7.7 in the supplementary Section 7 together with the Cauchy-Schwarz Inequality show that the entire second row as well as the two terms in the sum in the first line converge to zero. Thus, the asymptotic behavior of T_n is entirely determined by the terms I_1 and I_2 . The convergence of the term involving $I_1(t)$ is studied in the following proposition, the proof of which is deferred to the end of this section.

Proposition 6.1. *Assume (K), (Co), (SC), (CC), (HSR1), (mDep), that*

$$\|K\|_\infty + \|\Psi(\cdot; \beta_0)\|_\infty + \|\alpha_0(\cdot; \beta_0)\|_\infty + \sup_{n \in \mathbb{N}} \|\Delta_n\|_\infty + \|w\|_\infty < \infty$$

and that $hr_n p_n \rightarrow \infty$, $h \rightarrow 0$. Then,

$$a_n^2 \sqrt{h} \int_0^T I_1(t)^2 w(t) dt - h^{-\frac{1}{2}} A_n \xrightarrow{d} \mathcal{N}(0, B), \quad (6.5)$$

where

$$A_n := a_n^2 \sum_{i,j \in V_n} \int_0^T f_n(r, r) \left(\frac{1}{r_n p_n(r)} \bar{X}_n(r, \beta_0)^{-1} \right)^2 dN_{n,ij}(r),$$

$$B := 4K^{(2)} \int_0^T \bar{X}_n(s; \beta_0)^{-2} w(s)^2 \alpha(\theta_0, s)^2 \gamma(s) ds,$$

$$K^{(2)} := \int_0^2 \left(\int_{-1}^1 K(u+v) K(u) du \right)^2 dv.$$

Thus we conclude that

$$a_n^2 \sqrt{h} T_n - h^{\frac{1}{2}} A_n - a_n^2 \sqrt{h} \int_0^T I_2(t)^2 w(t) dt \xrightarrow{d} \mathcal{N}(0, B)$$

and the proof of Theorem 3.1 is completed by computing the term involving I_2 (mind the definition of c_n)

$$a_n^2 \sqrt{h} \int_0^T I_2(t)^2 w(t) dt = \int_0^T \left(\int_0^T K_{h,t}(s) \Delta_n(s) ds \right)^2 w(t) dt.$$

Let us now turn to the proof of Proposition 6.1. The proof involves the momentary- m -dependence assumption. As it was motivated in Section 3.2 we need that the functions φ (defined in (3.4)) are useful approximations to each other. This is the content of the next lemma.

Lemma 6.2. *Let $\|\Psi(\cdot; \beta_0)\|_\infty + \|\alpha_0\|_\infty + \|\bar{X}_n(\cdot; \beta_0)^{-1}\|_\infty < \infty$. There is a constant $C > 0$ and a random function $\Phi_n : [0, T] \rightarrow [0, \infty)$ such that for $I \subseteq J \subseteq V_n \times V_n$*

$$|\varphi_{n,ij,kl}^I(t, r) - \varphi_{n,ij,kl}^J(t, r)| \leq |J \setminus I| \cdot \frac{C}{r_n p_n(t) p_n(r)} B_n^{J \setminus I}, \quad (6.6)$$

$$|\varphi_{n,ij,kl}^I(t, r)| \leq \frac{1}{p_n(r)} \Phi_n(t) \quad (6.7)$$

for some function Φ_n and $B_n^A := (F + H_{UB}^A K_m^{V_n \times V_n})$ for $A \subseteq V_n \times V_n$. If, in addition, the assumption on \mathcal{A}_n in (CC) holds, then there is $C > 0$ such that $\mathbb{P}(\sup_{t \in [0, T]} \Phi_n(t) > C) \rightarrow 0$.

Proof of Lemma 6.2. Recall that $K_m^{V_n \times V_n}$ is the maximal size of a hub and that H_{UB}^A is the indicator function whether any edge from $A \subseteq V_n \times V_n$ has the potential of being a hub (this is independent of time). Since, $I \subseteq J$, we get for any $(k, l) \in V_n \times V_n$

$$\begin{aligned} & |\mathbb{1}(d_{t-4h}^n(kl, I) \geq m) - \mathbb{1}(d_{t-4h}^n(kl, J) \geq m)| \\ & \leq \mathbb{1}(\exists (i, j) \in J \setminus I : d_{t-4h}^n(kl, ij) \leq m) \leq \sum_{(i,j) \in J \setminus I} \mathbb{1}(d_{t-4h}^n(kl, ij) \leq m). \end{aligned}$$

Then, we obtain

$$\sum_{\substack{(i', j') \in V_n \times V_n \\ (i', j') \neq (i, j), (k, l)}} \Psi(X_{n, i' j'}(r), \beta_0) C_{n, i' j'}(r) |\mathbb{1}(d_{t-4h}^n(i' j', I) \geq m) - \mathbb{1}(d_{t-4h}^n(i' j', J) \geq m)|$$

$$\begin{aligned}
&\leq \|\Psi(\cdot; \beta_0)\|_\infty |J \setminus I| \sup_{(a,b) \in J \setminus I} \left(\mathbb{1} \left(H_{UB}^{ab} = 0 \right) + \mathbb{1} \left(H_{UB}^{ab} = 1 \right) \right) \\
&\quad \times \sum_{\substack{(i',j') \in V_n \times V_n \\ i',j' \neq (i,j),(k,l)}} C_{n,ab}(r) \mathbb{1}(d_{t-4h}^n(i'j', ab) \leq m) \\
&\leq \|\Psi(\cdot; \beta_0)\|_\infty \cdot |J \setminus I| \cdot \left(F + H_{UB}^{J \setminus I} K_m^{G_n} \right).
\end{aligned}$$

Replacing the definition of $B_n^{J \setminus I}$ yields (6.6). In order to show (6.7) we bound the $\varphi_{n,ij,kl}$ by the direct bounds and obtain this way an expression for Φ_n . When looking at the definition (3.4), we see that $\sup_{t \in [0, T]} \Phi_n(t)$ is essentially a constant times $\sup_{t \in [0, T]} \frac{1}{r_n p_n(t)} X_n(t; \beta_0)$ and hence (CC) gives the boundedness property of Φ_n . \square

We also need a result on the asymptotic behavior of the φ 's.

Lemma 6.3. *Suppose (CC) holds and that $\bar{X}_n(\cdot; \beta_0)$, w , $\alpha(\theta_0; \cdot)$ and all Δ_n are continuously differentiable with uniformly bounded first derivative. In addition, we require that p_n is smooth enough such that*

$$\sup_{s \in [0, T]} \sup_{r \in [s, s+2h]} \frac{p_n(s)}{p_n(r)} = O(1).$$

Then we have that

$$\sup_{s \in [0, T]} \sup_{(i,j) \in V_n \times V_n} \left| \frac{r_n^2 p_n^3(s)}{a_n^4} \varphi_{n,ij}(s) - 4K^{(2)} \omega(s)^2 \bar{X}_n(s)^{-3} \alpha_0(s) \right| = o_P(1),$$

where

$$K^{(2)} := \int_0^2 \left(\int_{-1}^1 K(u+v) K(u) du \right)^2 dv.$$

Proof. Let $\delta > 0$ be arbitrary and let $c > 0$ be so large such that $\mathbb{P}(\tilde{\mathcal{A}}_n(c, 0)) \geq 1 - \delta$. Now, on $\tilde{\mathcal{A}}_n$ we have for any $r \in [0, T]$ and any $(i, j) \in V_n \times V_n$

$$\begin{aligned}
&\left| \frac{1}{r_n p_n(r)} \sum_{(i',j') \neq (i,j)} \Psi(X_{n,i'j'}(r); \beta_0) C_{n,i'j'}(r) - \bar{X}_n(r; \beta_0) \right| \\
&\leq \left| \frac{1}{r_n p_n(r)} X_n(r; \beta_0) - \bar{X}_n(r; \beta_0) \right| + \frac{C_{n,ij}(r)}{r_n p_n(r)} \Psi(X_{n,ij}(r); \beta_0) \\
&\leq c \sqrt{\frac{\log r_n}{r_n p_n(r)}} + \frac{1}{r_n p_n(r)} \|\Psi(\cdot; \beta_0)\|_\infty \leq c^* \sqrt{\frac{\log r_n}{r_n p_n}},
\end{aligned}$$

where $c^* > 0$ is a suitable constant only depending on c . Thus, for any $\gamma > 0$

$$\mathbb{P} \left(\sup_{s \in [0, T]} \sup_{(i,j) \in V_n \times V_n} \left| \frac{r_n^2 p_n^3(s)}{a_n^4} \varphi_{n,ij}(s) - 4K^{(2)} \omega(s)^2 \bar{X}_n(s)^{-3} \alpha_0(s) \right| > \gamma \right)$$

$$\begin{aligned}
&\leq \mathbb{P} \left(\sup_{s \in [0, T]} \sup_{(i, j) \in V_n \times V_n} \frac{4p_n(s)}{h} \int_s^{s+2h} f_n(r, s)^2 \frac{\bar{X}_n(r; \beta_0)^{-2} \bar{X}_n(s; \beta_0)^{-2} \alpha_0(r)}{p_n(r)} \right. \\
&\quad \times \left| \frac{1}{r_n p_n(r)} \sum_{(i', j') \neq (i, j)} \Psi(X_{n, i' j'}(r); \beta_0) C_{n, i' j'}(r) - \bar{X}_n(r; \beta_0) \right| \\
&\quad + \sup_{s \in [0, T]} \left| \frac{4p_n(s)}{h} \int_s^{s+2h} f_n(r, s)^2 \frac{\bar{X}_n(r; \beta_0)^{-1} \bar{X}_n(s; \beta_0)^{-2} \alpha_0(r)}{p_n(r)} dr \right. \\
&\quad \quad \left. - 4K^{(2)} \omega(s)^2 \bar{X}_n(s)^{-3} \alpha_0(s) \right| > \gamma \Big) \\
&\leq \mathbb{P} \left(\sup_{s \in [0, T]} \frac{4p_n(s)}{h} \int_s^{s+2h} f_n(r, s)^2 \frac{1}{p_n(r)} c^* \sqrt{\frac{\log r_n}{r_n p_n}} dr \|\alpha_0\|_\infty \|\bar{X}_n(\cdot; \beta_0)^{-1}\|_\infty^4 \right. \\
&\quad + \sup_{s \in [0, T]} \left| \frac{4p_n(s)}{h} \int_s^{s+2h} f_n(r, s)^2 \frac{\bar{X}_n(r; \beta_0)^{-1} \bar{X}_n(s; \beta_0)^{-2} \alpha_0(r)}{p_n(r)} dr \right. \\
&\quad \quad \left. - 4K^{(2)} \omega(s)^2 \bar{X}_n(s)^{-3} \alpha_0(s) \right| > \gamma \Big) + \delta.
\end{aligned}$$

Both lines in the probability above are actually not random, so we need to prove convergence to zero of both lines separately. For the second line we use standard kernel approximation arguments and the fact that all involved functions are continuously differentiable with bounded first derivative (recall that $\alpha_0(s) = \alpha(\theta_0, s) + c_n \Delta_n(s)$). Moreover, the specific continuity requirement for p_n is important for this step because it allows us to estimate $p_n(s)^{-1} - p_n(r)^{-1}$ when s and r are close. This is also necessary for handling the first line. \square

With the above two lemmas at hand, we can finally prove Proposition 6.1.

Proof of Proposition 6.1. By Lemma 7.1 from the supplementary Section 7 it is sufficient to prove

$$a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \bar{X}_n(s; \beta_0)^{-1} dM_n(s) \right)^2 w(t) dt - h^{-\frac{1}{2}} A_n \xrightarrow{d} N(0, B). \quad (6.8)$$

Recall that

$$f_n(r, s) := \int_0^T h K_{h,t}(s) K_{h,t}(r) w(t) dt.$$

Then, we can rewrite (6.8) as follows

$$a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \bar{X}_n(s; \beta_0)^{-1} dM_n(s) \right)^2 w(t) dt$$

$$\begin{aligned}
&= 2a_n^2 h^{-\frac{1}{2}} \int_0^T \int_0^{r-} f_n(r, s) \frac{1}{r_n p_n(s)} \bar{X}_n(s; \beta_0)^{-1} \frac{1}{r_n p_n(r)} \bar{X}_n(r; \beta_0)^{-1} dM_n(s) dM_n(r) \quad (6.9) \\
&\quad + a_n^2 h^{-\frac{1}{2}} \int_0^T \int_{\{r\}} f_n(r, s) \frac{1}{r_n p_n(r)} \bar{X}_n(r; \beta_0)^{-1} \frac{1}{r_n p_n(s)} \bar{X}_n(s; \beta_0)^{-1} dM_n(s) dM_n(r).
\end{aligned} \tag{6.10}$$

Note that two martingales $M_{n,ij}$ and $M_{n,kl}$ do not jump simultaneously when $(i, j) \neq (k, l)$ and that the height of each jump is exactly one. We obtain

$$(6.10) = a_n^2 h^{-\frac{1}{2}} \sum_{(i,j) \in V_n \times V_n} \int_0^T \frac{f_n(r, r)}{r_n^2 p_n(r)^2} \bar{X}_n(r; \beta_0)^{-2} dN_{n,ij}(r) = h^{-\frac{1}{2}} A_n.$$

In the next step, we split the double summation in (6.9) in two steps as well,

$$\begin{aligned}
&(6.9) \\
&= 2a_n^2 \sum_{\substack{(i,j), (k,l) \in V_n \times V_n \\ (i,j) \neq (k,l)}} h^{-\frac{1}{2}} \int_0^T \int_0^{r-} \frac{f_n(r, s)}{r_n^2 p_n(s) p_n(r)} \bar{X}_n(s; \beta_0)^{-1} \bar{X}_n(r; \beta_0)^{-1} dM_{n,ij}(s) dM_{n,kl}(r)
\end{aligned} \tag{6.11}$$

$$\begin{aligned}
&\quad + 2a_n^2 \sum_{(i,j) \in V_n \times V_n} h^{-\frac{1}{2}} \int_0^T \int_0^{r-} \frac{f_n(r, s)}{r_n^2 p_n(s) p_n(r)} \bar{X}_n(s, \beta_0)^{-1} \bar{X}_n(r, \beta_0)^{-1} dM_{n,ij}(s) dM_{n,ij}(r).
\end{aligned} \tag{6.12}$$

By assumption $hK_{h,t}(s) \leq \|K\|_\infty < \infty$. Using this and the bounded support of K we can prove the useful bound $\int_0^T \int_0^T f_n(r, s)^2 \frac{1}{r_n^2 p_n(r) p_n(s)} ds dr \leq \|K\|_\infty^2 a_n^{-4}$. Note, moreover, that we achieved by the splitting that (6.12) is a martingale and thus we may compute the expectation of the square as the expectation of the quadratic variation. Therefore, we get for $n \rightarrow \infty$ with $\eta_0 := \inf_{t \in [0, T]} \bar{X}_n(t; \beta_0)$

$$\begin{aligned}
&\mathbb{E}((6.12)^2) \\
&\leq 4a_n^4 \sum_{(i,j) \in V_n \times V_n} \int_0^T \int_0^r \frac{f_n(r, s)^2 \|\Psi(\cdot; \beta_0)\|_\infty}{r_n^4 p_n(s)^2 p_n(r)^2 h \eta_0^4} \mathbb{E}(\Psi(X_{n,ij}(s); \beta_0) C_{n,ij}(s)) \alpha_0(s) ds \alpha_0(r) dr \\
&\leq 4 \frac{\|\Psi(\cdot; \beta_0)\|_\infty^2 \|\alpha_0\|_\infty^2}{h r_n p_n \eta_0^4} \|K\|_\infty^2 \rightarrow 0.
\end{aligned}$$

Lastly, we will prove the convergence of (6.11) to a normal distribution. We will do this by showing the assumptions of Rebolledo's martingale central limit theorem (cf. Theorem 6.3 in the Appendix). We note to this end that (6.11) is indeed a martingale evaluated at T . We therefore have to study the variation process as well as the jump process. Recall the definition of $\varphi_{n,ij}^I$ in (3.4). With that the variation process can be written as follows

$$\langle (6.11) \rangle$$

$$= \frac{1}{r_n} \sum_{(i,j),(i',j') \in V_n \times V_n} \int_0^T \int_{s_1-2h}^{s_1-} 2\varphi_{n,ij,i'j'}(s_1, s_2) dM_{n,i'j'}(s_2) dM_{n,ij}(s_1) \quad (6.13)$$

$$+ \frac{1}{r_n} \sum_{(i,j),(i',j') \in V_n \times V_n} \int_0^T \int_{\{s_1\}} \varphi_{n,ij,i'j'}(s_1, s_2) dM_{n,i'j'}(s_2) dM_{n,ij}(s_1). \quad (6.14)$$

We will prove that (6.13) converges to zero and (6.14) converges to a non-zero quantity which is going to be the asymptotic variance of the normal distribution. For the latter we note again that for $(i, j) \neq (i', j')$ the martingales $M_{n,ij}$ and $M_{n,i'j'}$ never jump simultaneously and that their jump height always equals one. So we get

$$(6.14) = \frac{1}{r_n} \sum_{(i,j) \in V_n \times V_n} \int_0^T \varphi_{n,ij}(s) dM_{n,ij}(s) + \frac{1}{r_n} \sum_{(i,j) \in V_n \times V_n} \int_0^T \varphi_{n,ij}(s) \Psi(X_{n,ij}(s); \beta_0) C_{n,ij}(s) \alpha_0(s) ds. \quad (6.15)$$

The first part of (6.15) converges to zero by Proposition 2.9 from Kreiss (2020+): The upper bound provided in the proposition converges to zero for $\tilde{\varphi}_{n,ij}^I(t) = \varphi_{n,ij}^{ij}(t)$ by the boundedness assumptions which allow us to use the bounds from Lemma 6.2. In the following we check the conditions of Lemma 2.9 in Kreiss (2020+). We use also the bound $B_n^{ij,kl} \leq B_n^{ij} B_n^{kl}$. Then, we have

$$\begin{aligned} & \frac{1}{r_n^2} \sum_{ij,kl} \mathbb{E} \left(\int_0^T \varphi_{n,ij}^{ij,kl}(t) dM_{n,ij}(t) \int_0^T (\varphi_{n,kl}(t) - \varphi_{n,kl}^{ij,kl}(t)) dM_{n,kl}(t) \right) \\ & \leq \frac{1}{r_n^2} \sum_{ij,kl} \mathbb{E} \left(\int_0^T \frac{1}{p_n(t)} \Phi_n(t) d|M_{n,ij}|(t) \int_0^T \frac{2CB_n^{ij,kl}}{r_n p_n(t)^2} d|M_{n,kl}|(t) \right) \\ & \leq \frac{2C}{r_n p_n} \mathbb{E} \left(\left(\sum_{ij} \int_0^T \frac{B_n^{ij}}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \sup_{t \in [0, T]} \Phi_n(t) \right). \end{aligned}$$

The expectation converges to zero by assumption and the first condition holds. The second condition holds because of Lemma 6.2 and

$$\begin{aligned} & \frac{1}{r_n^2} \sum_{ij,kl} \mathbb{E} \left(\int_0^T (\varphi_{n,ij}(t) - \varphi_{n,ij}^{ij,kl}(t)) dM_{n,ij}(t) \int_0^T (\varphi_{n,kl}(t) - \varphi_{n,kl}^{ij,kl}(t)) dM_{n,kl}(t) \right) \\ & \leq \frac{1}{r_n^2} \sum_{ij,kl} \mathbb{E} \left(\int_0^T \frac{2CB_n^{ij,kl}}{r_n p_n(t)^2} d|M_{n,ij}|(t) \int_0^T \frac{2CB_n^{ij,kl}}{r_n p_n(t)^2} d|M_{n,kl}|(t) \right) \\ & \leq \frac{4C^2}{r_n^2 p_n^2} \mathbb{E} \left(\left(\sum_{ij} \int_0^T \frac{(B_n^{ij})^2}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \right). \end{aligned}$$

We have again convergence to zero by assumption. And finally, we obtain for the third condition again by Lemma 6.2

$$\begin{aligned}
& \frac{1}{r_n^2} \sum_{ij} \int_0^T \mathbb{E} \left(\varphi_{n,ij}^{ij}(t)^2 C_{n,ij}(t) \Psi(X_{n,j}(t), \beta_0) \right) dt \\
& \leq \frac{1}{r_n} \mathbb{E} \left(\sum_{ij} \int_0^T \frac{1}{r_n p_n(t)^2} \Phi_n(t)^2 \|\Psi(\cdot; \beta_0)\|_\infty C_{n,ij}(t) dt \right) \\
& \leq \frac{1}{r_n p_n} \mathbb{E} \left(\sum_{ij} \int_0^T \frac{C_{n,ij}(t)}{r_n p_n(t)} dt \sup_{t \in [0, T]} \Phi_n(t)^2 \right) \|\Psi(\cdot; \beta_0)\|_\infty
\end{aligned}$$

which converges to zero by assumption. Hence, we can employ Proposition 2.9 in Kreiss (2020+) to have convergence of the first part of (6.15) to zero.

For the second part of (6.15) we note that, by Lemma 6.3, $\frac{r_n^2 p_n(s)^3}{a_n^4} \varphi_{n,ij}(s)$ converges uniformly in s and (i, j) to $\ell_n(s) := 4K^{(2)} w(s)^2 \bar{X}_n(s; \beta_0)^{-3} \alpha_0(s)$ (note that α_0 depends on n). By boundedness of Δ_n we get uniformly $\ell_n(s) \rightarrow \ell(s) := 4K^{(2)} w(s)^2 \bar{X}_n(s; \beta_0)^{-3} \alpha(s; \theta_0)$. Thus, we obtain by also using Assumption (SC)

$$\begin{aligned}
& \frac{1}{r_n} \sum_{(i,j) \in V_n \times V_n} \int_0^T \varphi_{n,ij}(s) \Psi(X_{n,ij}(s); \beta_0) C_{n,ij}(s) \alpha_0(s) ds \\
& = \sum_{(i,j) \in V_n \times V_n} \int_0^T \frac{a_n^4}{r_n^3 p_n(s)^3} \left(\frac{r_n^2 p_n(s)^3}{a_n^4} \varphi_{n,ij}(s) - \ell_n(s) \right) \Psi(X_{n,ij}(s); \beta_0) C_{n,ij}(s) \alpha_0(s) ds \\
& \quad + \int_0^T \frac{a_n^4}{r_n^2 p_n(s)^2} \left(\frac{1}{r_n p_n(s)} \sum_{(i,j) \in V_n \times V_n} \Psi(X_{n,ij}(s); \beta_0) C_{n,ij}(s) - \bar{X}_n(s; \beta_0) \right) \ell(s) \alpha_0(s) ds \\
& \quad + \int_0^T \frac{a_n^4}{r_n^2 p_n(s)^2} \bar{X}_n(s; \beta_0) \ell_n(s) \alpha_0(s) ds \\
& \rightarrow 4K^{(2)} \int_0^T \bar{X}_n(s; \beta_0)^{-2} w(s)^2 \alpha(\theta_0, s)^2 \gamma(s) ds = B.
\end{aligned}$$

Lastly, we split (6.13) in two parts and obtain

$$(6.13) = \frac{1}{r_n} \sum_{\substack{(i,j), (i',j') \in V_n \times V_n \\ (i,j) \neq (i',j')}} \int_0^T \int_{s_1-2h}^{s_1-} 2\varphi_{n,ij,i'j'}(s_1, s_2) dM_{n,i'j'}(s_2) dM_{n,ij}(s_1) \quad (6.16)$$

$$+ \frac{1}{r_n} \sum_{(i,j) \in V_n \times V_n} \int_0^T \int_{s_1-2h}^{s_1-} 2\varphi_{n,ij,ij}(s_1, s_2) dM_{n,ij}(s_2) dM_{n,ij}(s_1). \quad (6.17)$$

We have that (6.16) converges to zero by Theorem 2.11 in Kreiss (2020+). In order to see that (6.17) converges to zero we apply the bounds from Lemma 6.2. More precisely, let $\gamma, \delta > 0$ be arbitrary and choose $C > 0$ such that $\mathbb{P}(\sup_{t \in [0, T]} \Phi_n(t) > C) \leq \delta$. Then, we firstly make crude bounds in the probability such that we can apply Markov's Inequality and use martingale properties:

$$\begin{aligned}
\mathbb{P}((6.17) > \gamma) &\leq \mathbb{P}\left(\frac{1}{r_n} \sum_{(i,j)} \int_0^T \int_{s_1-2h}^{s_1-} \frac{2}{p_n(s_2)} \Phi_n(s_1) d|M_{n,ij}|(s_2) d|M_{n,ij}|(s_1) > \gamma\right) \\
&\leq \mathbb{P}\left(\frac{2C}{r_n} \sum_{(i,j)} \int_0^T \int_{s_1-2h}^{s_1-} \frac{1}{p_n(s_2)} d|M_{n,ij}|(s_2) d|M_{n,ij}|(s_1) > \gamma\right) + \delta \\
&\leq 2\mathbb{E}\left(\int_0^T \left(\int_{s_1-2h}^{s_1-} \frac{1}{p_n(s_2)} dM_{n,ij}(s_2) + \int_{s_1-2h}^{s_1} \frac{C_{n,ij}(s_2)}{p_n(s_2)} \|\Psi(\cdot; \beta_0)\|_\infty ds_2\right) \|\Psi(\cdot; \beta_0)\|_\infty ds_1\right) + \delta \\
&\leq 2hT \|\Psi(\cdot; \beta_0)\|_\infty^2 + \delta.
\end{aligned}$$

The above converges to δ since $h \rightarrow 0$. Since δ is arbitrary we conclude that (6.17) = $o_P(1)$ and finish this part of the proof.

Finally, we are concerned with the jump process associated with (6.11). Let therefore $\eta > 0$ be given. Then, the variation process of the jump process is given by

$$\begin{aligned}
&\frac{4a_n^4}{h} \sum_{i,j \in V_n} \int_0^T \mathbb{1}\left(2a_n^2 h^{-\frac{1}{2}} \left| \sum_{\substack{(k,l) \in V_n \\ (k,l) \neq (i,j)}} \int_0^{r-} f_n(r,s) \frac{\bar{X}_n(s; \beta_0)^{-1} \bar{X}_n(r; \beta_0)^{-1}}{r_n p_n(s) \cdot r_n p_n(r)} dM_{n,kl}(s) \right| > \eta\right) \\
&\quad \times \left(\sum_{\substack{k,l \in V_n \\ (k,l) \neq (i,j)}} \int_0^{r-} f_n(r,s) \frac{\bar{X}_n(s; \beta_0)^{-1} \bar{X}_n(r; \beta_0)^{-1}}{r_n p_n(s) \cdot r_n p_n(r)} dM_{n,kl}(s) \right)^2 X_{n,ij}(r) C_{n,ij}(r) \alpha_0(r) dr.
\end{aligned}$$

Note that up to the indicator function the above expression is identical to the variation process of (6.11) which we just proved to converge. Hence, we have convergence of the above to zero if we can prove that (mind the supremum in the indicator)

$$\mathbb{1}\left(\sup_{\substack{i,j \in V_n \\ r \in [0, T]}} 2a_n^2 h^{-\frac{1}{2}} \left| \sum_{\substack{k,l \in V_n \\ (k,l) \neq (i,j)}} \int_0^{r-} f_n(r,s) \frac{\bar{X}_n(s; \beta_0)^{-1} \bar{X}_n(r; \beta_0)^{-1}}{r_n p_n(s) \cdot r_n p_n(r)} dM_{n,kl}(s) \right| > \eta\right) = o_P(1).$$

Proving this is significantly easier than handling the previous expressions because we have no double integral with respect to two martingales. We can make the following estimate:

(recall the notation $\eta_0 := \inf_{t \in [0, T]} \bar{X}_n(t; \beta_0)$)

$$\begin{aligned} & \sup_{\substack{r \in [0, T] \\ i, j \in V_n}} 2a_n^2 h^{-\frac{1}{2}} \left| \sum_{\substack{k, l \in V_n \\ (k, l) \neq (i, j)}} \int_0^{r^-} f_n(r, s) \frac{\bar{X}_n(s; \beta_0)^{-1} \bar{X}_n(r; \beta_0)^{-1}}{r_n p_n(s) \cdot r_n p_n(r)} dM_{n, kl}(s) \right| \\ & \leq \sup_{r \in [0, T]} 2a_n^2 h^{-\frac{1}{2}} \sum_{k, l \in V_n} \int_0^r f_n(r, s) \frac{\bar{X}_n(s; \beta_0)^{-1} \bar{X}_n(r; \beta_0)^{-1}}{r_n p_n(s) \cdot r_n p_n(r)} dM_{n, kl}(s) \end{aligned} \quad (6.18)$$

$$\begin{aligned} & + \sup_{r \in [0, T]} \frac{4\|\alpha_0\|_\infty}{\eta_0} \cdot \frac{a_n^2}{r_n p_n} h^{-\frac{1}{2}} \int_0^r f_n(r, s) \bar{X}_n(s; \beta_0)^{-1} \\ & \quad \times \frac{1}{r_n p_n(s)} \sum_{k, l \in V_n} \Psi(X_{n, kl}(s), \beta_0) C_{n, kl}(s) ds. \end{aligned} \quad (6.19)$$

We note firstly that (6.19) is $o_P(1)$ because the sum converges by assumption, since $\frac{a_n^2}{r_n p_n}$ is bounded and because $f_n(r, s)$ acts like a kernel and causes the integral to be $O(h)$ (uniformly over r). For (6.18) we note that we take the supremum of a martingale. Hence, we can apply Doob's Martingale inequality which applies to all right continuous martingales (cf. Theorem 5.1.3 in [Cohen and Elliott \(2015\)](#)) to obtain:

$$\begin{aligned} & \mathbb{E} \left(\left(\sup_{r \in [0, T]} 2a_n^2 h^{-\frac{1}{2}} \left| \sum_{k, l \in V_n} \int_0^r f_n(r, s) \frac{\bar{X}_n(s; \beta_0)^{-1} \bar{X}_n(r; \beta_0)^{-1}}{r_n p_n(s) \cdot r_n p_n(r)} dM_{n, kl}(s) \right| \right)^2 \right) \\ & \leq 4\mathbb{E} \left(\left(2a_n^2 h^{-\frac{1}{2}} \left| \sum_{k, l \in V_n} \int_0^T f_n(r, s) \frac{\bar{X}_n(s; \beta_0)^{-1} \bar{X}_n(r; \beta_0)^{-1}}{r_n p_n(s) \cdot r_n p_n(r)} dM_{n, kl}(s) \right| \right)^2 \right) \\ & \leq \frac{16\|\Psi(\cdot; \beta_0)\|_\infty \|\alpha_0\|_\infty}{\eta_0^4} \cdot \frac{a_n^4}{r_n^2 p_n^2} \cdot \frac{1}{r_n p_n} \cdot \frac{1}{h} \int_0^T f_n(r, s)^2 ds, \end{aligned}$$

which converges since $a_n^4/r_n^2 p_n^2 = O(1)$, $r_n p_n \rightarrow \infty$ and by definition of f_n . \square

6.2 Mixing

Assumption (CC) in the main text, which involves the concentration behavior of the covariates, can be shown to hold under certain mixing assumptions. Mixing on networks has been discussed in [Kreiss \(2020+\)](#) and we give here just a quick overview. In order to formulate the mixing assumptions we need the following notation. Let $\delta_n = a \log n$ for some fixed $a > 0$ and consider a random, time-varying partition $(G^t(k, m, \delta_n))_{k, m} \subseteq E_{n, t}$ of the edge set, where $k = 1, \dots, \mathcal{K}$ and $m = 1, \dots, r_n$ (so some of the sets are empty). We call $G^t(k, m, \delta_n)$ the m -th group of type k . Note that \mathcal{K} is fixed. This partition shall have the following properties

1. $(k, m) \neq (k', m') \Rightarrow G^t(k, m, \delta_n) \cap G^t(k', m', \delta_n) = \emptyset$,
2. $\bigcup_{k,m} G^t(k, m, \delta_n) = E_{n,t}$,
3. For $m \neq m'$: $(i, j) \in G^t(k, m, \delta_n), (k, l) \in G^t(k, m', \delta_n) \Rightarrow d_i^n(ij, kl) \geq \delta_n$.

Let $I_{n,ij}^{k,m,t} := \mathbb{1}((i, j) \in G^t(k, m, \delta_n))$ be the indicator which states if the pair (i, j) is contained in the set $G^t(k, m, \delta_n)$. The idea is now to rewrite the sum of interest in \mathcal{A}_n and $\tilde{\mathcal{A}}_n$ as follows

$$\sum_{i,j \in V_n} C_{n,ij}(t) \Psi(X_{n,ij}(t); \beta) = \sum_{k,m} U_{k,m}^t, \text{ where } U_{k,m}^t := \sum_{i,j \in V_n} C_{n,ij}(t) \Psi(X_{n,ij}(t); \beta) I_{n,ij}^{k,m,t}.$$

The β -mixing coefficient of the network is then defined as follows

$$\beta_t(n) := \max_{\substack{M=1, \dots, r_n \\ k=1, \dots, \mathcal{K}}} \beta \left([U_{k,m}^t]_{m \leq M-1}, U_{k,M}^{n,t} \right).$$

In order to make use of these concepts, we have to define the maximal size of a group of type k , $S_k(t) := \max_{m=1, \dots, r_n} |G^t(k, m, \delta_n)|$ and the indicator

$$\Gamma_n^t(c) := \left(\max_{k=1, \dots, \mathcal{K}} S_k^2 \leq c^2 \frac{r_n p_n(t)}{\log r_n p_n(t)} \right)$$

for $c > 0$ which indicates if no single group is too large. Moreover, we have to define the restriction of the random variables of interest to the event that the partition fulfills $\Gamma_n^t(c) = 1$,

$$Y_{n,ij}(t) := (C_{n,ij}(t) \Psi(X_{n,ij}(t); \beta) - \mathbb{E}(C_{n,ij}(t) \Psi(X_{n,ij}(t); \beta) | \Gamma_n^t = 1)) \Gamma_n^t.$$

Let $E_{k,m}^{n,t} = \mathbb{E}(|G^t(k, m, \delta_n)|)$ be the expected size of a group. We need to make the following assumptions.

Assumption (Mixing)

Suppose that for any $b > 0$ we can choose $a, c_1 > 0$ such that for $\delta_n = a \log n$, $\beta_t(n) \leq n^{-b}$ and $\mathbb{P}(\Gamma_n^t(c_1) = 0) \leq n^{-b}$ for all $t \in [0, T]$. Suppose furthermore that for each such a , there are constants $c_2, c_3 > 0$ such that for all $t \in [0, T]$

$$\frac{1}{r_n p_n(t)} \sum_{m=1}^{r_n} E_{k,m}^{n,t} \geq c_2,$$

and that for pairwise different vertices $v_1, \dots, v_4 \in V_n$ and all $t \in [0, T]$

$$\begin{aligned} \text{Var} \left(Y_{n,v_1 v_2}(t) I_{n,v_1 v_2}^{k,m,t} \right) &\leq c_3 \|\Psi(\cdot; \beta_0)\|_\infty^2 E_{k,m}^{n,t} r_n^{-1}, \\ \text{Cov} \left(Y_{n,v_1 v_2}(t) I_{n,v_1 v_2}^{k,m,t}, Y_{n,v_2 v_3}(t) I_{n,v_2 v_3}^{k,m,t} \right) &\leq c_3 \|\Psi(\cdot; \beta)\|_\infty^2 E_{k,m}^{n,t} r_n^{-\frac{3}{2}}, \end{aligned}$$

$$\text{Cov}\left(Y_{n,v_1v_2}(t)I_{n,v_1v_2}^{k,m,t}, Y_{n,v_3v_4}(t)I_{n,v_3v_4}^{k,m,t}\right) \leq c_3 \|\Psi(\cdot; \beta)\|_\infty^2 E_{k,m}^{n,t} r_n^{-2}.$$

The existence of such a partition is in particular plausible in networks which are related to geographic locations (like the bike stations). The exponential decay of the β -mixing coefficient is motivated through an exponential growth of the δ_n -neighborhoods, i.e., the number of pairs in a distance smaller than δ_n . For an example we refer to [Kreiss \(2020+\)](#). For the assumptions on the correlations note that $\mathbb{E}(Y_{n,ij}(t)) \approx \|\Psi(\cdot; \beta)\|_\infty p_n(t)$ and that, at least for reasonable partitions, $\mathbb{E}(I_{n,ij}^{k,m,t}) \approx E_{k,m}^{n,t}/r_n p_n(t)$.

We can prove that under these conditions the Assumption (CC) can be proven. We do this separately for the two statements in (CC).

Proposition 6.2. *Let $c_2 > 0$ be arbitrary but fixed. Suppose that the (Mixing) Assumption holds. Suppose furthermore that $p_n(t)$ is Hölder continuous with exponent α and constant $H_{n,p}$ such that for $m > 0$ large enough $\frac{H_{n,p}}{p_n^2} n^{-m\alpha} \rightarrow 0$. Suppose that $\sup_{\beta \in B_n(c_2)} \|\Psi(\cdot; \beta)\|_\infty = O(1)$, and that*

$$|\Psi(x, \beta_1) - \Psi(x, \beta_2)| \leq L_\Psi \omega_\Psi(\|\beta_1 - \beta_2\|), \quad |\Psi(x_1, \beta) - \Psi(x_2, \beta)| \leq L_\Psi(\beta) \tilde{\omega}_\Psi(\|x_1 - x_2\|),$$

where $\sup_{\beta \in B_n(c_2)} L_\Psi(\beta) = O(1)$ and $\omega_\Psi(n^{-k}) \sqrt{r_n/p_n \log r_n} \rightarrow 0$ for some $k > 0$. For the same k , let $\#\text{jumps}(C_n)[n^{-k}]$ denote the maximal number of jumps of C_n in any interval of length n^{-k} , and assume that

$$\frac{\#\text{jumps}(C_n)[n^{-k}]}{r_n p_n} = O_P\left(\sqrt{\frac{\log r_n}{r_n p_n}}\right),$$

$$\sup_{t_1, t_2: |t_1 - t_2| \leq n^{-k}} \sum_{(i,j) \in V_n \times V_n} \frac{\tilde{\omega}_\Psi(\|X_{n,ij}(t_1) - X_{n,ij}(t_2)\|)}{r_n p_n} = O_P\left(\sqrt{\frac{\log r_n}{r_n p_n}}\right).$$

Finally, let $\eta_0 = \inf_{\beta \in B_n(c_2)} \|\bar{X}_n(\cdot; \beta)\|_\infty$ and assume that $\eta_0^{-1} = O(1)$. Then, for any $\delta > 0$, there is $c_1 > 0$ such that

$$\mathbb{P}(\mathcal{A}_n(c_1, c_2)) \geq 1 - \delta \quad \text{and} \quad \mathbb{P}(\tilde{\mathcal{A}}_n(c_1, c_2)) \geq 1 - \delta.$$

Lemma 6.4. *Let $c_2 > 0$ be fixed and suppose that $\sup_{\beta \in B_n(c_2)} \|\Psi(\cdot; \beta)\|_\infty < \infty$ and (Mixing) hold. Then,*

$$\sup_{s \in [0, T], \beta \in B_n(c_2)} \mathbb{E}\left(\left(\frac{X_n(s, \beta)}{r_n p_n(s)}\right)^2\right) = O(1).$$

The proofs of both results are presented in Section 7.2 in the Supplement.

6.3 Rebolledos Martingale Central Limit Theorem

For easier reference we state here the exact form of Rebolledo's Martingale Central Limit Theorem. We state here a version based on Theorem II.5.1 in [Andersen et al. \(1993\)](#), the original work is [Rebolledo \(1980\)](#).

Let $M^n = (M_1^n, \dots, M_k^n)$ be a vector of sequences of locally square integrable martingales on an interval \mathcal{T} . For $\varepsilon > 0$ we denote by M_ε^n a vector of locally square integrable martingales that contain all jumps of components of M^n which are larger in absolute value than ε , i.e., $M_i^n - M_{\varepsilon,i}^n$ is a local square integrable martingale for all $i = 1, \dots, k$ and $|\Delta M_i^n - \Delta M_{\varepsilon,i}^n| \leq \varepsilon$. Furthermore, we denote by $\langle M^n \rangle := \left(\langle M_i^n, M_j^n \rangle \right)_{i,j=1,\dots,k}$ the $k \times k$ matrix of quadratic covariations.

Moreover, we denote by M a multivariate, continuous Gaussian martingale with $\langle M \rangle_t = V_t$, where $V : \mathcal{T} \rightarrow \mathbb{R}^{k \times k}$ is a continuous deterministic $k \times k$ positive semi-definite matrix valued function on \mathcal{T} such that its increments $V_t - V_s$ are also positive semi-definite for $s \leq t$, then $M_t - M_s \sim \mathcal{N}(0, V_t - V_s)$ is independent of $(M_r : r \leq s)$. Given such a function V , such a Gaussian process M always exists. We can now formulate the central limit theorem for martingales.

Theorem 6.3. *Let $\mathcal{T}_0 \subseteq \mathcal{T}$. Assume that for all $t \in \mathcal{T}_0$ as $n \rightarrow \infty$ we have $\langle M^n \rangle_t \xrightarrow{\mathbb{P}} V_t$ and $\langle M_\varepsilon^n \rangle \xrightarrow{\mathbb{P}} 0$. Then, $M_t^n \xrightarrow{d} M_t$ as $n \rightarrow \infty$ for all $t \in \mathcal{T}_0$.*

7 Technical Proofs

7.1 Supporting Lemmas

Similarly as in Section 6.1, we assume throughout that the model from Theorem 3.1 as well as Assumption (ID) hold and do not mention this explicitly every time. Moreover, all assumptions made in the lemmas below are contained in the set of conditions made in Section 3.2.

Lemma 7.1. *Let I_1 be defined as in (6.1) and assume all notation from Theorem 3.1. Suppose that (CC) holds and let $\|\alpha(\theta_0, \cdot)\|_\infty + \sup_{n \in \mathbb{N}} \|\Delta\|_\infty + \|K\|_\infty + \|w\|_\infty < \infty$ as well as $\|\bar{X}_n(\cdot; \beta_0)\|_\infty + \|\bar{X}_n(\cdot; \beta_0)^{-1}\|_\infty < \infty$. Suppose moreover that $\log r_n / hr_n p_n \rightarrow 0$, $\sqrt{h} \log r_n \rightarrow 0$ and $a_n^2 / r_n p_n = O(1)$. Assume finally that $\int_{-\infty}^{\infty} K(u)^2 < \infty$. Then,*

$$a_n^2 \sqrt{h} \int_0^T I_1(t)^2 w(t) dt - a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \bar{X}_n(s; \beta_0)^{-1} dM_n(s) \right)^2 w(t) dt = o_P(1).$$

Lemma 7.2. *Let $I_2(t)$ be defined as in (6.2) and let otherwise all notation be as in Theorem 3.1. Suppose that $\sup_{n \in \mathbb{N}} \|\Delta_n\|_\infty + \|w\|_\infty < +\infty$. Then,*

$$a_n^2 \sqrt{h} \int_0^T I_2(t)^2 w(t) dt = O(1). \tag{7.1}$$

Lemma 7.3. Let I_3 be defined as in (6.3) and let otherwise all notation be the same as in Theorem 3.1. Suppose that the conditions on α in (C) and the conditions on $\hat{\alpha}_n$ in (P) (without the independence) hold and let $\|w\|_\infty < \infty$. Then,

$$a_n^2 \sqrt{h} \int_0^T I_3(t)^2 w(t) dt = o_P(1). \quad (7.2)$$

Lemma 7.4. Let I_4 be defined as in (6.4) and let otherwise all notation be the same as in Theorem 3.1. Suppose (CC), the assumptions on $\tilde{\beta}_n$ in (P) (without the independence) and the assumptions on Ψ in (C) hold and that $\|\alpha(\theta_0; \cdot)\|_\infty + \|K\|_\infty + \|w\|_\infty + \sup_{n \in \mathbb{N}} \|\Delta_n\|_\infty < +\infty$. For the bandwidth we require that $\sqrt{h} \log r_n \rightarrow 0$ and $hr_n p_n \rightarrow \infty$, moreover, $a_n^2 / r_n p_n = O(1)$. Then,

$$a_n^2 \sqrt{h} \int_0^T I_4(t)^2 w(t) dt = o_P(1). \quad (7.3)$$

Lemma 7.5. Let I_1, I_2 be defined as in (6.1), (6.2). Suppose that all assumptions of the Lemmas 7.1 and 7.2 hold. Then,

$$a_n^2 \sqrt{h} \int_0^T I_1(t) I_2(t) w(t) dt = o_P(1).$$

Lemma 7.6. Let I_1, I_3 be defined as in (6.1), (6.3). Suppose that, in addition to all assumptions of the Lemmas 7.1 and 7.3, the estimator $\hat{\theta}_n$ is obtained independently of the main data. Then,

$$a_n^2 \sqrt{h} \int_0^T I_1(t) I_3(t) w(t) dt = o_P(1).$$

Lemma 7.7. Let I_1, I_4 be defined as in (6.1), (6.4). Suppose that, in addition to all assumptions of the Lemmas 7.8, 7.1 and 7.4, the estimator $\tilde{\beta}_n$ is obtained independently of the main data. Then,

$$a_n^2 \sqrt{h} \int_0^T I_1(t) I_4(t) w(t) dt = o_P(1).$$

Proof of Lemma 7.1. We rewrite the quantity of interest as follows

$$\begin{aligned} & a_n^2 \sqrt{h} \int_0^T I_1(t)^2 w(t) dt - a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \bar{X}_n(s; \beta_0)^{-1} dM_n(s) \right)^2 w(t) dt \\ &= a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\left(\frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}(s; \beta_0)^{-1} \right) dM_n(s) \right)^2 w(t) dt \\ & \quad + 2a_n^2 \sqrt{h} \int_0^T \int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \bar{X}_n(s; \beta_0)^{-1} dM_n(s) \end{aligned} \quad (7.4)$$

$$\times \int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\left(\frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}_n(s; \beta_0)^{-1} \right) dM_n(s) w(t) dt. \quad (7.5)$$

Part (7.4) converges to zero in probability as we show next. We estimate

$$(7.4) \leq 2a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{w(t)}{r_n p_n(s)} \left(\left(\frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}_n(s; \beta_0)^{-1} \right) dN_n(s) \right)^2 dt \\ + 2a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\left(\frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}_n(s; \beta_0)^{-1} \right) \right. \\ \left. \times X_n(s; \beta_0) \alpha_0(s) ds \right)^2 w(t) dt.$$

By assumption we can find for any $\delta > 0$ a $c > 0$ such that $\mathbb{P}(\mathcal{A}_n(c, 0)) \geq 1 - \delta$ for all $n \in \mathbb{N}$. This can be used as follows for any $\gamma > 0$

$$\mathbb{P}((7.4) > \gamma) \leq \mathbb{P}((7.4) > \gamma, \mathcal{A}_n(c, 0)) + \mathbb{P}(\mathcal{A}_n(c, 0)^c) \\ \leq \mathbb{P} \left(2a_n^2 \sqrt{h} \int_0^T \left[\left(\int_0^T K_{h,t}(s) \frac{c \sqrt{\log r_n}}{(r_n p_n(s))^{\frac{3}{2}}} dN_n(s) \right)^2 \right. \right. \\ \left. \left. + \left(\int_0^T K_{h,t}(s) \frac{c \sqrt{\log r_n}}{(r_n p_n(s))^{\frac{3}{2}}} X_n(s; \beta_0) \alpha_0(s) ds \right)^2 \right] w(t) dt > \gamma \right) + \delta \\ \leq \mathbb{P} \left(\int_0^T \left[4a_n^2 \sqrt{h} \left(\int_0^T K_{h,t}(s) \frac{\sqrt{\log r_n}}{(r_n p_n(s))^{\frac{3}{2}}} dM_n(s) \right)^2 \right. \right. \\ \left. \left. + 6a_n^2 \sqrt{h} \left(\int_0^T K_{h,t}(s) \frac{\sqrt{\log r_n}}{(r_n p_n(s))^{\frac{3}{2}}} X_n(s; \beta_0) \alpha_0(s) ds \right)^2 \right] w(t) dt > \frac{\gamma}{c^2} \right) + \delta.$$

The above expression can be bounded completely analogously by a version of (7.13) as in the proof of Lemma 7.4: The only difference is that the two 2's have to be replaced by 4 and a 6. But this does not change the convergence behavior and hence we get that the above expression converges to δ and we conclude that (7.4) = $o_P(1)$.

In order to show that (7.5) = $o_P(1)$, we make the following abbreviations

$$f_n(t, s) := K_{h,t}(s) \frac{1}{r_n p_n(s)} \bar{X}_n(s; \beta_0)^{-1}, \\ g_n(t, s) := K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\left(\frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}_n(s; \beta_0)^{-1} \right).$$

Then, we get

$$\begin{aligned}
(7.5) &= 2a_n^2 \sqrt{h} \int_0^T \int_0^T \int_0^T f_n(t, s) g_n(t, r) w(t) dt dM_n(s) dM_n(r) \\
&= 2a_n^2 \sqrt{h} \int_0^T \int_0^{r-} \int_0^T f_n(t, s) g_n(t, r) w(t) dt dM_n(s) dM_n(r) \tag{7.6}
\end{aligned}$$

$$+ 2a_n^2 \sqrt{h} \int_0^T \int_{r+}^T \int_0^T f_n(t, s) g_n(t, r) w(t) dt dM_n(s) dM_n(r) \tag{7.7}$$

$$+ 2a_n^2 \sqrt{h} \int_0^T \int_{\{r\}} \int_0^T f_n(t, s) g_n(t, r) w(t) dt dM_n(s) dM_n(r). \tag{7.8}$$

We begin by treating (7.6). Note that the two inner integrals are a predictable function of r and therefore, we may treat the stochastic integral like a martingale. From Lenglart's Inequality (see equation (2.15.18) in Andersen et al. (1993)) we see that we may alternatively prove that the quadratic variation converges to zero in probability. Hence, we prove in the following that the quadratic variation of (7.6) converges to zero in probability. Let to this end $\delta > 0$ be arbitrary and choose $c > 0$ so large such that $\mathbb{P}(\mathcal{A}_n(c, 0) \cap \tilde{\mathcal{A}}_n(c, 0)) \geq 1 - \delta$ for all $n \in \mathbb{N}$. For simplicity of notation we let $\xi_n(s) := c\sqrt{\log r_n/r_n p_n(s)}$ and note that on $\mathcal{A}_n(c, 0) \cap \tilde{\mathcal{A}}_n(c, 0)$ we have

$$\left| \left(\frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}_n(s; \beta_0)^{-1} \right| \leq \xi_n(s), \quad \left| \frac{1}{r_n p_n(s)} X_n(s; \beta_0) - \bar{X}_n(s; \beta_0) \right| \leq \xi_n(s)$$

We use these considerations to obtain for every $\gamma > 0$

$$\begin{aligned}
&\mathbb{P} \left(4a_n^4 h \int_0^T \left(\int_0^{r-} \int_0^T f_n(t, s) g_n(t, r) w(t) dt dM_n(s) \right)^2 \alpha_0(r) X_n(r, \beta_0) dr > \gamma \right) \\
&\leq \mathbb{P} \left(4a_n^4 h \int_0^T \left(\int_0^{r-} \int_0^T f_n(t, s) g_n(t, r) w(t) dt dM_n(s) \right)^2 \alpha_0(r) X_n(r, \beta_0) dr > \gamma, \right. \\
&\quad \left. \mathcal{A}_n(c, 0) \cap \tilde{\mathcal{A}}_n(c, 0) \right) + \delta \\
&\leq \mathbb{P} \left(8a_n^4 h \int_0^T \alpha_0(r) X_n(r, \beta_0) \xi_n(r)^2 \left[\left(\int_0^{r-} \int_0^T \frac{K_{h,t}(s) K_{h,t}(r)}{r_n^2 p_n^2 \bar{X}_n(s; \beta_0)} w(t) dt dN_n(s) \right)^2 \right. \right. \\
&\quad \left. \left. + \left(\int_0^{r-} \int_0^T \frac{K_{h,t}(s) K_{h,t}(r)}{r_n^2 p_n^2 \bar{X}_n(s; \beta_0)} w(t) \alpha_0(s) X_n(s; \beta_0) dt ds \right)^2 \right] dr > \gamma \right) + \delta \\
&\leq \mathbb{P} \left(8a_n^4 h \int_0^T \alpha_0(r) (\bar{X}_n(r, \beta_0) + \xi_n(r)) \xi_n(r)^2 \left[2 \left(\int_0^{r-} \int_0^T \frac{K_{h,t}(s) K_{h,t}(r)}{r_n^2 p_n^2 \bar{X}_n(s; \beta_0)} w(t) dt dM_n(s) \right)^2 \right. \right. \\
&\quad \left. \left. + 3 \left(\int_0^{r-} \int_0^T \frac{K_{h,t}(s) K_{h,t}(r)}{r_n^2 p_n^2 \bar{X}_n(s; \beta_0)} w(t) \alpha_0(s) X_n(s; \beta_0) dt ds \right)^2 \right] dr > \gamma \right) + \delta
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{\gamma} \left(8a_n^4 h \int_0^T \alpha_0(r) (\bar{X}_n(r, \beta_0) + \xi_n(r)) \xi_n(r)^2 \right. \\
&\quad \times \left[2\mathbb{E} \left(\left(\int_0^{r-} \int_0^T \frac{K_{h,t}(s)K_{h,t}(r)}{r_n^2 p_n^2 \bar{X}_n(s; \beta_0)} w(t) dt dM_n(s) \right)^2 \right) \right. \\
&\quad \left. \left. + \mathbb{E} \left(3 \left(\int_0^{r-} \int_0^T \frac{K_{h,t}(s)K_{h,t}(r)}{r_n^2 p_n^2 \bar{X}_n(s; \beta_0)} w(t) \alpha_0(s) X_n(s; \beta_0) dt ds \right)^2 \right) \right] dr \right) + \delta \\
&\leq \frac{1}{\gamma} \left(8a_n^4 h \int_0^T \alpha_0(r) (\bar{X}_n(r, \beta_0) + \xi_n(r)) \xi_n(r)^2 \right. \\
&\quad \times \left[2 \left(\int_0^r \left(\int_0^T \frac{K_{h,t}(s)K_{h,t}(r)}{r_n^2 p_n^2 \bar{X}_n(s; \beta_0)} w(t) dt \right)^2 \alpha_0(s) \mathbb{E}(X_n(s; \beta_0)) ds \right) \right. \\
&\quad \left. + 3 \int_0^r \int_0^T K_{h,t}(s)K_{h,t}(r) \mathbb{E} \left(\left(\frac{w(t)\alpha_0(s)X_n(s; \beta_0)}{r_n^2 p_n^2 \bar{X}_n(s; \beta_0)} \right)^2 \right) dt ds \right] dr \right) + \delta \\
&\leq \frac{c^*}{\gamma} \left(\frac{c^2 h \log r_n}{r_n p_n} \int_0^T \left[2 \left(\int_0^r \left(\int_0^T K_{h,t}(s)K_{h,t}(r) dt \right)^2 \frac{1}{r_n p_n} ds \right) \right. \right. \\
&\quad \left. \left. + \int_0^r \int_0^T K_{h,t}(s)K_{h,t}(r) dt ds \right] dr \right) + \delta,
\end{aligned}$$

where the last inequality follows from boundedness of w , α_0 , $\bar{X}_n(\cdot; \beta_0)$ and $\bar{X}_n(\cdot; \beta_0)^{-1}$ as well as by assumption on the expectation and $a_n^2/r_n p_n = O(1)$ for a suitable c^* . The above converges to δ because

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} K_{h,t}(s)K_{h,t}(r) dt ds = O(1) \text{ and } \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} K_{h,t}(s)K_{h,t}(r) dt \right)^2 ds = O\left(\frac{1}{h}\right).$$

Therefore, we have proven that (7.6) = $o_p(1)$. For (7.7) we note that interchanging the order of integration yields

$$(7.7) = 2a_n^2 \sqrt{h} \int_0^T \int_0^{s-} \int_0^T f_n(t, s) g_n(t, r) w(t) dt dM_n(r) dM_n(s).$$

Hence, we have an expression which is almost identical to (7.6) and may therefore be treated by the same arguments. So we skip the proof of (7.7) = $o_p(1)$. Instead, we detail the arguments for handling (7.8). Note firstly that in the inner integral with respect to $M_n(s)$ the integral over the intensity can be ignored and only the counting process integral remains. Similarly for the outer integral only the counting process integral remains. In consideration of this we get (let as usually $\gamma, \delta > 0$ be arbitrary and $c > 0$ such that $\mathbb{P}(\mathcal{A}_n(c, 0)) \geq 1 - \delta$)

$$\mathbb{P}((7.8) > \gamma) = \mathbb{P} \left(2a_n^2 \sqrt{h} \int_0^T \int_0^T f_n(t, r) g_n(t, r) w(t) dt dN_n(r) > \gamma \right)$$

$$\begin{aligned}
&= \mathbb{P} \left(2a_n^2 \sqrt{h} \int_0^T \int_0^T f_n(t, r) g_n(t, r) w(t) dt dN_n(r) > \gamma, \mathcal{A}_n(c, 0) \right) + \delta \\
&\leq \mathbb{P} \left(\frac{2a_n^2}{r_n^2 p_n^2} \sqrt{h} \int_0^T \int_0^T K_{h,t}(r)^2 \bar{X}_n(r; \beta_0)^{-1} \xi_n(r) w(t) dt dN_n(r) > \gamma \right) + \delta \\
&\leq \frac{2a_n^2}{\gamma r_n^2 p_n^2} \sqrt{h} \int_0^T \int_0^T K_{h,t}(r)^2 \bar{X}_n(r; \beta_0)^{-1} \xi_n(r) w(t) \alpha_0(r) \mathbb{E}(X_n(r; \beta_0)) dt dr + \delta \quad (7.9)
\end{aligned}$$

We bound the above expression by assumption which implies that $\mathbb{E}(X_n(r; \beta_0)/r_n p_n)$ is bounded as well as boundedness of $a_n^2/r_n p_n$, α_0 , $\bar{X}_n(\cdot, \beta_0)^{-1}$ and w and the definition of $\xi_n(r)$. So there is a constant c^* such that

$$(7.9) \leq \frac{c^*}{\gamma} \sqrt{\frac{h \log r_n}{r_n p_n}} \int_0^T \int_0^T K_{h,t}(r)^2 dt dr + \delta \leq \frac{T c^*}{\gamma} \sqrt{\frac{\log r_n}{h r_n p_n}} \int_{-\infty}^{\infty} K(u)^2 du + \delta.$$

Since, $\log r_n / h r_n p_n \rightarrow 0$ the proof of the Lemma is complete. \square

Proof of Lemma 7.2. Since we're only interested in boundedness and convergence to zero we may ignore the function w because it is bounded. For (7.1) we note that simply by bounding Δ_n

$$\begin{aligned}
a_n^2 \sqrt{h} \int_0^T I_2(t)^2 dt &= a_n^2 \sqrt{h} \int_0^T \left(c_n \int_0^T K_{h,t}(s) \Delta_n(s) ds \right)^2 dt \\
&\leq a_n^2 c_n^2 \sqrt{h} T \left(\sup_{n \in \mathbb{N}} \|\Delta_n\|_{\infty} \right)^2 < +\infty
\end{aligned}$$

by assumption and (7.1) is shown. \square

Proof of Lemma 7.3. Since we're only interested in boundedness and convergence to zero we may ignore the function w because it is bounded. By the Cauchy-Schwarz Inequality and the continuity assumption on α

$$\begin{aligned}
a_n^2 \sqrt{h} \int_0^T I_3(t)^2 dt &= a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \left(\alpha(\theta_0, s) - \alpha(\hat{\theta}_n, s) \right) ds \right)^2 dt \\
&\leq a_n^2 \sqrt{h} \int_0^T \int_0^T K_{h,t}(s) \left(\alpha(\theta_0, s) - \alpha(\hat{\theta}_n, s) \right)^2 ds dt \\
&\leq a_n^2 \sqrt{h} \omega_{\alpha} \left(\|\theta_0 - \hat{\theta}_n\| \right)^2 \int_0^T L_{\alpha}(t)^2 dt.
\end{aligned}$$

Thus, we conclude by the assumptions that for any $\varepsilon, \delta > 0$ and a suitably chosen $c^* > 0$ we find for n large enough

$$\mathbb{P} \left(a_n^2 \sqrt{h} \int_0^T I_3(t)^2 dt > \varepsilon \right)$$

$$\leq \mathbb{P} \left(\left\| \hat{\theta}_n - \theta_0 \right\| > \frac{c^*}{\sqrt{r_n p_n}} \right) + \mathbb{P} \left(a_n^2 \sqrt{h} \omega_\alpha \left(\frac{c^*}{\sqrt{r_n p_n}} \right)^2 \int_0^T L_\alpha(t)^2 dt > \varepsilon \right) \leq \delta$$

□

Proof of Lemma 7.4. Since we're only interested in boundedness and convergence to zero we may ignore the function w because it is bounded. Finally, we show (7.2). We firstly insert the limit of X_n as follows

$$\begin{aligned} & \frac{1}{3} a_n^2 \sqrt{h} \int_0^T I_4(t)^2 dt \\ &= \frac{1}{3} a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\frac{1}{\frac{1}{r_n p_n(s)} X_n(s; \tilde{\beta}_n)} - \frac{1}{\frac{1}{r_n p_n(s)} X_n(s; \beta_0)} \right) dN_n(s) \right)^2 dt \\ &\leq a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\frac{1}{\frac{1}{r_n p_n(s)} X_n(s; \tilde{\beta}_n)} - \frac{1}{\bar{X}_n(s; \tilde{\beta}_n)} \right) dN_n(s) \right)^2 dt \quad (7.10) \end{aligned}$$

$$+ a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\frac{1}{\bar{X}_n(s; \tilde{\beta}_n)} - \frac{1}{\bar{X}_n(s; \beta_0)} \right) dN_n(s) \right)^2 dt \quad (7.11)$$

$$+ a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\frac{1}{\frac{1}{r_n p_n(s)} X_n(s; \beta_0)} - \frac{1}{\bar{X}_n(s; \beta_0)} \right) dN_n(s) \right)^2 dt. \quad (7.12)$$

The arguments for (7.10) and (7.12) are identical. Therefore, we only detail (7.10). Define to this end for a constant $c_1 > 0$ the event

$$\mathcal{B}(c_2) := \left\{ \left\| \tilde{\beta}_n - \beta \right\| \leq \frac{c_2}{\sqrt{r_n p_n}} \right\}.$$

By the assumptions on $\tilde{\beta}_n$ and $\mathcal{A}(c_1, c_2)$ we see that for any given $\delta > 0$ we can choose c_1, c_2 so large such that $\mathbb{P}(\mathcal{A}_n(c_1, c_2) \cap \mathcal{B}_n(c_2)) \geq 1 - \delta$ for all $n \in \mathbb{N}$. Let now δ, c_1, c_2 be fixed. Then, we obtain for any $\gamma > 0$ (use the Cauchy-Schwarz Inequality in the last line)

$$\begin{aligned} & \mathbb{P}((7.10) > \gamma) \leq \mathbb{P}((7.10) > \gamma, \mathcal{A}_n(c_1, c_2) \cap \mathcal{B}_n(c_2)) + \mathbb{P}((\mathcal{A}_n(c_1, c_2) \cap \mathcal{B}_n(c_2))^c) \\ &\leq \mathbb{P} \left(a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \frac{c_1 \sqrt{\log r_n}}{(r_n p_n(s))^{\frac{3}{2}}} dN_n(s) \right)^2 > \gamma \right) + \delta \\ &\leq \mathbb{P} \left(\int_0^T \left[2a_n^2 \sqrt{h} \left(\int_0^T K_{h,t}(s) \frac{\sqrt{\log r_n}}{(r_n p_n(s))^{\frac{3}{2}}} dM_n(s) \right)^2 \right. \right. \\ &\quad \left. \left. + 2a_n^2 \sqrt{h} \left(\int_0^T K_{h,t}(s) \frac{\sqrt{\log r_n}}{(r_n p_n(s))^{\frac{3}{2}}} X_n(s; \beta_0) \alpha_0(s) ds \right)^2 \right] dt > \frac{\gamma}{c_1^2} \right) + \delta \end{aligned}$$

$$\begin{aligned}
&\leq \frac{c_1^2}{\gamma} \int_0^T 2a_n^2 \sqrt{h} \int_0^T K_{h,t}(s) \frac{\log r_n}{r_n p_n(s)} \alpha_0(s) \\
&\quad \times \left[\frac{K_{h,t}(s)}{r_n p_n(s)} \mathbb{E} \left(\frac{X_n(s; \beta_0)}{r_n p_n(s)} \right) + \mathbb{E} \left(\left(\frac{X_n(s; \beta_0)}{r_n p_n(s)} \right)^2 \right) \alpha_0(s) \right] ds dt + \delta. \tag{7.13}
\end{aligned}$$

By assumption the expectations are uniformly bounded. Moreover, by assumption $\|\alpha_0\|_\infty < +\infty$ and the kernel K is bounded too. We find therefore a constant $c^* > 0$ such that we can continue

$$(7.13) \leq c^* \frac{T c_1^2}{\gamma} \sqrt{h} \log r_n \frac{a_n^2}{r_n p_n} \left[\frac{1}{hr_n p_n} + 1 \right] + \delta \rightarrow \delta$$

since $\sqrt{h} \log r_n \rightarrow 0$, $hr_n p_n \rightarrow 0$ and $a_n^2/r_n p_n = o(1)$ by assumption. Thus, we have shown that (7.10), (7.12) = $o_P(1)$. Thus, we have left to prove that (7.11) = $o(1)$. By Lemma 6.1 we get for any $\gamma > 0$ that (let c_2 and δ be as above)

$$\begin{aligned}
&\mathbb{P}((7.11) > \gamma) \leq \mathbb{P}((7.11) > \gamma, \mathcal{B}_n(c_2)) + \delta \\
&\leq \mathbb{P} \left(a_n^2 \sqrt{h} \int_0^T \left(\int_0^T K_{h,t}(s) \cdot \frac{\mathcal{L}_n(s)}{r_n p_n(s) \cdot \inf_{\beta \in B(c_2)} \bar{X}_n(s; \beta)^2} \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) dN_n(s) \right)^2 dt > \gamma \right) \\
&\quad + \delta \\
&\leq \mathbb{P} \left(2a_n^2 \sqrt{h} \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right)^2 \int_0^T \left[\left(\int_0^T K_{h,t}(s) \frac{\mathcal{L}_n(s)}{r_n p_n(s) \cdot \inf_{\beta \in B(c_2)} \bar{X}_n(s; \beta)^2} dM_n(s) \right)^2 \right. \right. \\
&\quad \left. \left. + \left(\int_0^T K_{h,t}(s) \frac{\mathcal{L}_n(s)}{r_n p_n(s) \cdot \inf_{\beta \in B(c_2)} \bar{X}_n(s; \beta)^2} X_n(s; \beta_0) \alpha_0(s) ds \right)^2 \right] dt > \gamma \right) + \delta \\
&\leq \frac{1}{\gamma} \int_0^T 2a_n^2 \sqrt{h} \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right)^2 \int_0^T K_{h,t}(s) \alpha_0(s) \left(\frac{\mathcal{L}_n(s)}{\inf_{\beta \in B(c_2)} \bar{X}_n(s; \beta)^2} \right)^2 \\
&\quad \times \left[\frac{K_{h,t}(s)}{r_n p_n(s)} \mathbb{E} \left(\frac{X_n(s; \beta_0)}{r_n p_n(s)} \right) + \mathbb{E} \left(\left(\frac{X_n(s; \beta_0)}{r_n p_n(s)} \right)^2 \alpha_0(s) \right) \right] ds dt + \delta. \tag{7.14}
\end{aligned}$$

Similarly as before, we obtain by assumption that the expectations are bounded and we find that for a certain constant $c^* > 0$ (which might be different from before)

$$\begin{aligned}
(7.14) &\leq \frac{2c^*}{\gamma} a_n^2 \sqrt{h} \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right)^2 \int_0^T \left(\frac{\mathcal{L}_n(s)}{\inf_{\beta \in B(c_2)} \bar{X}_n(s; \beta)^2} \right)^2 \left[\frac{1}{hr_n p_n(s)} + 1 \right] ds + \delta \\
&\rightarrow \delta
\end{aligned}$$

since $a_n^2 \sqrt{h} \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right)^2 \rightarrow 0$ and since

$$\int_0^T \left(\frac{\mathcal{L}_n(s)}{\inf_{\beta \in B(c_2)} \bar{X}_n(s; \beta)^2} \right)^2 = O(1)$$

by assumption. Thus, the proof is complete. \square

Proof of Lemma 7.5. We have

$$a_n^2 \sqrt{h} \int_0^T I_1(t) I_2(t) w(t) dt = a_n^2 \sqrt{h} c_n \int_0^T \int_0^T \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{\Delta_n(r) w(t)}{X_n(s; \beta_0)} dt dr dM_n(s)$$

Since the integrand is predictable, the stochastic integral is a martingale. Therefore, we may use Lenglart's Inequality (cf. (2.5.18) in [Andersen et al. \(1993\)](#)) to argue that it is enough to prove that the quadratic variation of the process converges to zero. Suppose to this end that $\delta > 0$ is arbitrary and that $c > 0$ is chosen such that $\mathbb{P}(\mathcal{A}_n(c, 0)) \geq 1 - \delta$. Let $\xi_n = \sqrt{\log r_n / r_n p_n}$. On the set $\mathcal{A}_n(c, 0)$ it holds for all $r \in [0, T]$ that

$$\left| \left(\frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}_n(s; \beta_0)^{-1} \right| \leq \xi_n.$$

Then we obtain for the quadratic variation and for any $\gamma > 0$ (replace also the definition of c_n and use the Cauchy-Schwarz Inequality together with $\int_0^T \int_0^T K_{h,t}(s) K_{h,t}(r) dt dr \leq 1$)

$$\begin{aligned} & \mathbb{P} \left(a_n^4 h c_n^2 \int_0^T \left(\int_0^T \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{\Delta_n(r) w(t)}{X_n(s; \beta_0)} dt dr \right)^2 \alpha_0(s) X_n(s; \beta_0) ds > \gamma \right) \\ & \leq \mathbb{P} \left(a_n^2 \sqrt{h} \int_0^T \left(\int_0^T \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{\Delta_n(r) w(t)}{X_n(s; \beta_0)} dt dr \right)^2 \alpha_0(s) X_n(s; \beta_0) ds > \gamma \right) \\ & \leq \mathbb{P} \left(a_n^2 \sqrt{h} \int_0^T \int_0^T \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{\Delta_n(r)^2 w(t)^2 \alpha_0(s)}{X_n(s; \beta_0)} dt dr ds > \gamma, \mathcal{A}_n(c, 0) \right) + \delta \\ & \leq \mathbb{P} \left(a_n^2 \sqrt{h} \frac{T \|\Delta_n\|_\infty^2 \|w\|_\infty^2 \|\alpha_0\|_\infty}{r_n p_n} (\|\bar{X}_n(\cdot; \beta_0)^{-1}\|_\infty + \xi_n) > \gamma \right) + \delta. \end{aligned}$$

Now we see that the expression in the probability is not random and hence the probability equals zero if the inequality inside is violated. This is the case for large n since $a_n^2 / r_n p_n = O(1)$, $h \rightarrow 0$ and the boundedness assumptions on $\|\Delta_n\|_\infty$, $\|\alpha_0\|$ and $\|\bar{X}_n(\cdot; \beta_0)^{-1}\|_\infty$. Thus, we have that the quadratic variation of $a_n^2 \sqrt{h} \int_0^T I_1(t) I_2(t) w(t) dt$ converges to zero in probability and thus also the martingale itself. \square

Proof of Lemma 7.6. Let $\delta > 0$ be arbitrary and let c_1, c_2 be such that

$$\mathbb{P} \left(\left\| \hat{\theta}_n - \theta_0 \right\| \leq c_2 \frac{1}{\sqrt{r_n p_n}}, \mathcal{A}_n(c_1, c_2) \right) \geq 1 - \delta.$$

By the independence of $\hat{\theta}_n$ from the remaining random quantities, we have that

$$a_n^2 \sqrt{h} \int_0^T I_1(t) I_3(t) w(t) dt$$

$$= a_n^2 \sqrt{h} \int_0^T \int_0^T \int_0^T K_{h,t}(r) K_{h,t}(s) \left(\alpha_0(\theta_0, r) - \alpha(\hat{\theta}_n, r) \right) \frac{w(t)}{X_n(s; \beta_0)} dr dt dM_n(s)$$

is a martingale. As a consequence we can prove the convergence to zero by applying Lenglart's Inequality (cf. (2.5.18) in [Andersen et al. \(1993\)](#)) and prove that the quadratic variation converges to zero in probability. We do this in the following. Let $\gamma > 0$ be arbitrary

$$\begin{aligned} & \mathbb{P} \left(a_n^4 h \int_0^T \left(\int_0^T \int_0^T K_{h,t}(r) K_{h,t}(s) \left(\alpha_0(\theta_0, r) - \alpha(\hat{\theta}_n, r) \right) \frac{w(t)}{X_n(s; \beta_0)} dr dt \right)^2 \right. \\ & \quad \left. \times \alpha_0(s) X_n(s; \beta_0) ds > \gamma \right) \\ & \mathbb{P} \left(a_n^4 h \int_0^T \int_0^T \int_0^T K_{h,t}(r) K_{h,t}(s) \left(\alpha_0(\theta_0, r) - \alpha(\hat{\theta}_n, r) \right)^2 w(t)^2 dr dt \frac{\alpha_0(s)}{X_n(s; \beta_0)} ds > \gamma \right) \\ & \mathbb{P} \left(a_n^4 h \int_0^T \int_0^T \int_0^T K_{h,t}(r) K_{h,t}(s) L_\alpha(r)^2 \omega_\alpha \left(\frac{c_2}{\sqrt{r_n p_n}} \right)^2 w(t)^2 dr dt \frac{\alpha_0(s)}{X_n(s; \beta_0)} ds > \gamma \right) + \delta. \end{aligned}$$

At this point we can proceed in exactly the same way as at the end of the proof of [Lemma 7.5](#) by using the properties of L_α and ω_α . \square

Proof of Lemma 7.7. We firstly write

$$\begin{aligned} & a_n^2 \sqrt{h} \int_0^T I_1(t) I_4(t) w(t) dt \\ & = a_n^2 \sqrt{h} \int_0^T \int_0^T \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \beta_0) X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} w(t) dN_n(r) dt dM_n(s) \\ & = a_n^2 \sqrt{h} \int_0^T \int_0^T \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \beta_0) X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} w(t) dM_n(r) dt dM_n(s) \end{aligned} \tag{7.15}$$

$$+ a_n^2 \sqrt{h} \int_0^T \int_0^T \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} \alpha_0(r) w(t) dr dt dM_n(s). \tag{7.16}$$

We begin with (7.16). We write

$$\begin{aligned} & (7.16) \\ & = a_n^2 \sqrt{h} \int_0^T \int_0^s \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} \alpha_0(r) w(t) dt dr dM_n(s) \end{aligned} \tag{7.17}$$

$$+ a_n^2 \sqrt{h} \int_0^T \int_0^r \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} \alpha_0(r) w(t) dt dM_n(s) dr. \tag{7.18}$$

We see that (7.17) is a martingale because the integrand is predictable. Thus, by Lenglar's Inequality it is sufficient to prove that the quadratic variation converges to zero. We do this in the following. Let as usual $\delta, \gamma > 0$ and choose $c_1, c_2 > 0$ such that the probability of the event $\|\tilde{\beta}_n - \beta_0\| \leq \frac{c_2}{\sqrt{r_n p_n}} \cap \mathcal{A}_n(c_1, c_2) \cap \tilde{\mathcal{A}}_n(c_1, c_2)$ is at least $1 - \delta$. On this event, we have by Lemma 6.1

$$\begin{aligned} \left| \left(\frac{1}{r_n p_n(r)} X_n(r; \beta_0) \right)^{-1} - \bar{X}_n(r, \beta_0)^{-1} \right| &\leq c_1 \sqrt{\frac{\log r_n}{r_n p_n(r)}}, \\ \left| \frac{1}{r_n p_n(r)} X_n(r; \beta_0) - \bar{X}_n(r, \beta_0) \right| &\leq c_1 \sqrt{\frac{\log r_n}{r_n p_n(r)}}, \\ \left| \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n) X_n(r; \beta_0)} \right| &\leq \frac{1}{r_n p_n} \left(2c_1 \sqrt{\frac{\log r_n}{r_n p_n}} + \mathcal{L}_n(r) \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) \right). \end{aligned}$$

Then we obtain for the quadratic variation of (7.17)

$$\begin{aligned} &\mathbb{P} \left(a_n^4 h \int_0^T \left(\int_0^s \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} \alpha_0(r) w(t) dt dr \right)^2 \right. \\ &\quad \left. \alpha_0(s) X_n(s; \beta_0) ds > \gamma \right) \\ &\leq \mathbb{P} \left(a_n^4 h \int_0^T \int_0^s \int_0^T K_{h,t}(s) K_{h,t}(r) \left(\frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n)} \right)^2 \frac{\alpha_0(r)^2 w(t)^2 \alpha_0(s)}{X_n(s; \beta_0)} dt dr ds > \gamma \right) \\ &\leq \mathbb{P} \left(a_n^4 h \|\alpha_0\|_\infty^3 \|w\|_\infty^2 \int_0^T \frac{1}{r_n p_n} \left(2c_1 \sqrt{\frac{\log r_n}{r_n p_n}} + \mathcal{L}_n(r) \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) \right)^2 dr \right. \\ &\quad \left. \times \left(\|\bar{X}_n(\cdot; \beta_0)^{-1}\|_\infty + c_1 \sqrt{\frac{\log r_n}{r_n p_n}} \right) \left(\|\bar{X}_n(\cdot; \beta_0)\|_\infty + c_1 \sqrt{\frac{\log r_n}{r_n p_n}} \right)^2 > \gamma \right) + \delta \end{aligned}$$

The probability above converges to zero since $a_n^2 / r_n p_n = O(1)$, $\int_0^T \mathcal{L}_n(r)^2 dr < \infty$ and $a_n^2 \sqrt{h} \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right)^2 \rightarrow 0$ and $h \log r_n \rightarrow 0$. So we have (7.17) = $o_P(1)$. The proof of (7.18) = $o_P(1)$ requires momentary- m -dependence and is given in Lemma 7.8.

Let us now turn to (7.15). Note to this end that (7.15) and (7.5) have the same structure. Therefore, we may employ very similar ideas. In fact, when letting

$$f_n(t, s) := K_{h,t}(s) X_n(s; \beta_0)^{-1} \text{ and } g_n(t, r) := K_{h,t}(r) \left(X_n(r; \tilde{\beta}_n)^{-1} - X_n(r; \beta_0)^{-1} \right)$$

we see that we can repeat the arguments which lead to (7.6)-(7.8). The independence of $\tilde{\beta}_n$ from the main observations ensured that the correspondences to (7.6), (7.7) and (7.8)

have the same measurability structure and therefore the same martingales arguments can be applied. In order to be able to finish the proof in an analogue fashion it is required that f_n and g_n can be bounded on a set with large probability. This is the case because for any $\delta > 0$ there are $c_1, c_2 > 0$ such that

$$\mathbb{P} \left(\|\tilde{\beta}_n - \beta_0\| \leq \frac{c_2}{\sqrt{r_n p_n}}, \mathcal{A}_n(c_1, c_2) \right) \geq 1 - \delta.$$

Then, on this event,

$$f_n(t, s) \leq K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(\bar{X}_n(s; \beta_0)^{-1} + c_1 \sqrt{\frac{\log r_n}{r_n p_n(s)}} \right)$$

and by Lemma 6.1

$$g_n(t, s) \leq K_{h,t}(s) \frac{1}{r_n p_n(s)} \left(2c_1 \sqrt{\frac{\log r_n}{r_n p_n(s)}} + \mathcal{L}_n(s) \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) \right).$$

The proof, can then be carried out in the same way. \square

The following result uses momentary- m -dependence in a similar way as in 6.1.

Lemma 7.8. *Suppose (mDep), (CC), (Co), (HSR2), the assumptions on Ψ in (C) and the assumptions on $\tilde{\beta}_n$ in (P) hold. Suppose moreover that $a_n/\sqrt{r_n p_n} = O(1)$, $\sqrt{h} \log r_n \rightarrow 0$. Then,*

$$a_n^2 \sqrt{h} \int_0^T \int_0^r \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} \alpha_0(r) w(t) dt dM_n(s) dr = o_P(1).$$

Proof. Recall that $f_n(r, s) := \int_0^T h K_{h,t}(s) K_{h,t}(r) w(t) dt$. Then, we may rewrite

$$\begin{aligned} & a_n^2 \sqrt{h} \int_0^T \int_0^r \int_0^T K_{h,t}(s) K_{h,t}(r) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} \alpha_0(r) w(t) dt dM_n(s) dr \\ &= a_n^2 h^{-\frac{1}{2}} \int_0^T \int_0^r f_n(r, s) \frac{X_n(r; \beta_0) - X_n(r; \tilde{\beta}_n)}{X_n(r; \tilde{\beta}_n) X_n(s; \beta_0)} \alpha_0(r) dM_n(s) dr \\ &= a_n^2 h^{-\frac{1}{2}} \int_0^T \int_0^r f_n(r, s) \frac{1}{r_n p_n(s)} \left(\frac{1}{r_n p_n(r)} X_n(r; \beta_0) - \frac{1}{r_n p_n(r)} X_n(r; \tilde{\beta}_n) \right) \\ & \quad \times \left(\left(\frac{1}{r_n p_n(r)} X_n(r; \tilde{\beta}_n) \frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}_n(r, \tilde{\beta}_n)^{-1} \bar{X}_n(s; \beta_0)^{-1} \right) \alpha_0(r) dM_n(s) dr \end{aligned} \tag{7.19}$$

$$+ a_n^2 h^{-\frac{1}{2}} \int_0^T \int_0^r f_n(r, s) \frac{1}{r_n p_n(s)} \frac{\frac{1}{r_n p_n(r)} X_n(r; \beta_0) - \frac{1}{r_n p_n(r)} X_n(r; \tilde{\beta}_n)}{\bar{X}_n(r, \tilde{\beta}_n) \bar{X}_n(s; \beta_0)} \alpha_0(r) dM_n(s) dr. \tag{7.20}$$

Let now $\eta_0 := \inf_{t \in [0, T]} \bar{X}_n(t; \beta_0)$ and choose n so large such that $\sqrt{\log r_n / r_n p_n} \leq \min(1, 1/\eta_0)$. For the remainder of the proof, n will always be so large such that this relation holds. Suppose that we are on the event $\|\tilde{\beta}_n - \beta\| \leq c_2 / \sqrt{r_n p_n}$ intersected with $\mathcal{A}_n(c_1, c_2) \cap \tilde{\mathcal{A}}_n(c_1, c_2)$ for $c_1 \geq 1$ and $c_2 > 0$ arbitrary. Then,

$$\begin{aligned} \sup_{r \in [0, T]} \left| \frac{1}{r_n p_n(r)} X_n(r; \beta_0) - \frac{1}{r_n p_n(r)} X_n(r; \tilde{\beta}_n) \right| &\leq c_1 \sqrt{\frac{\log r_n}{r_n p_n}} + L_\Psi \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right), \\ \sup_{r \in [0, T]} \sup_{\beta \in B_n(c_2)} \left(\frac{1}{r_n p_n(r)} X_n(r; \beta) \right)^{-1} &\leq c_1 \sqrt{\frac{\log r_n}{r_n p_n}} + \sup_{r \in [0, T]} \bar{X}_n(r; \beta_0)^{-1} \leq \frac{2}{\eta_0}. \end{aligned}$$

From the last two we obtain in particular that there is a constant (independent of c_1, c_2) such that

$$\begin{aligned} &\left| \left(\frac{1}{r_n p_n(r)} X_n(r; \tilde{\beta}_n) \frac{1}{r_n p_n(s)} X_n(s; \beta_0) \right)^{-1} - \bar{X}_n(r, \tilde{\beta}_n)^{-1} \bar{X}_n(s; \beta_0)^{-1} \right| \\ &\leq \frac{c_1 c^*}{\eta_0} \sqrt{\frac{\log r_n}{r_n p_n}}. \end{aligned}$$

Let now $\delta > 0$ be arbitrary and choose $c_1 \geq 1$ and $c_2 > 0$ such that the event $\|\tilde{\beta}_n - \beta\| \leq c_2 / \sqrt{r_n p_n}$ intersected with $\mathcal{A}_n(c_1, c_2) \cap \tilde{\mathcal{A}}_n(c_1, c_2)$ has probability at least $1 - \delta$ for all $n \in \mathbb{N}$. Then we obtain for every $\gamma > 0$ (note that f_n acts like a kernel in the sense that $f_n(r, s) \neq 0$ only if $|r - s| \leq 2h$)

$$\begin{aligned} &\mathbb{P}(|(7.19)| > \gamma) \\ &\leq \mathbb{P} \left(a_n^2 h^{-\frac{1}{2}} \int_0^T \int_0^r f_n(r, s) \frac{\|\alpha_0\|_\infty}{r_n p_n(s)} \left(c_1 \sqrt{\frac{\log r_n}{r_n p_n}} + L_\Psi \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) \right) \right. \\ &\quad \times \left. \frac{c_1 c^*}{\eta_0} \sqrt{\frac{\log r_n}{r_n p_n}} d|M_n|(s) dr > \gamma \right) \\ &\leq \sum_{i, j \in V_n} \frac{2}{\gamma} \mathbb{E} \left(a_n^2 h^{-\frac{1}{2}} \int_0^T \int_0^r f_n(r, s) \frac{\|\alpha_0\|_\infty}{r_n p_n(s)} \left(c_1 \sqrt{\frac{\log r_n}{r_n p_n}} + L_\Psi \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) \right) \right. \\ &\quad \times \left. \frac{c_1 c^*}{\eta_0} \sqrt{\frac{\log r_n}{r_n p_n}} \alpha_0(s) \Psi(X_{n, ij}(s); \beta_0) C_{n, ij}(s) ds dr \right) \\ &\leq \frac{4c_1 c^*}{\gamma \eta_0} T \|K\|_\infty \|\alpha_0\|_\infty^2 \|\Psi(\cdot; \beta_0)\|_\infty \sqrt{h} a_n^2 \left(c_1 \sqrt{\frac{\log r_n}{r_n p_n}} + L_\Psi \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) \right) \sqrt{\frac{\log r_n}{r_n p_n}} \end{aligned}$$

which converges to zero since $\eta_0^{-1} = O(1)$, $a_n / \sqrt{r_n p_n} = O(1)$ and $a_n \omega_\Psi(c_2 / \sqrt{r_n p_n}) = O(1)$ and $\sqrt{h} \log r_n \rightarrow 0$. So we have shown that (7.19) converges to zero. In order to handle

(7.20) we have to use momentary- m -dependence ideas and define to this end for any set $I \subseteq V_n \times V_n$

$$\begin{aligned} F_n(r, s) &:= \frac{\frac{1}{r_n p_n(r)} X_n(r; \beta_0) - \frac{1}{r_n p_n(r)} X_n(r; \tilde{\beta} + n)}{\bar{X}_n(r; \tilde{\beta}_n) \bar{X}_n(s; \beta_0)}, \\ F_n^I(r, s) &:= \frac{\frac{1}{r_n p_n(r)} X_n^I(r, s; \beta_0) - \frac{1}{r_n p_n(r)} X_n^I(r, s; \tilde{\beta} + n)}{\bar{X}_n(r; \tilde{\beta}_n) \bar{X}_n(s; \beta_0)}, \\ X_n^I(r, s; \beta) &:= \sum_{k, l \in V_n} \Psi(X_{n, kl}(r); \beta) C_{n, kl}(r) \mathbb{1}(d_{s-4h}^n(kl, I) \geq m). \end{aligned}$$

Note that in the definition of X_n^I , the distance is always indexed by $s - 4h$ regardless which value we plug in for r . In particular $X_n(r; \beta) = X_n^\emptyset(r, s; \beta)$ for all $s \in [0, T]$. These definitions ensure that

$$\tilde{\varphi}_{n, ij}^I(s) := \frac{a_n^2}{r_n p_n(s) \sqrt{h}} \int_s^{s+2h} f_n(r, s) \alpha_0(r) F_n^I(r, s) dr$$

is predictable with respect to $\tilde{F}_{ij, t}^{n, I, m}$. Note that the index ij is not necessary, but we keep it here to reproduce the notation from Proposition 2.9 in Kreiss (2020+). We write moreover $\tilde{\varphi}_{n, ij} := \tilde{\varphi}_{n, ij}^\emptyset$. In the following we show two bounds similar to Lemma 6.2. On the event $\|\tilde{\beta}_n - \beta_0\| \leq c_2/\sqrt{r_n p_n}$ we have that

$$\begin{aligned} &|\tilde{\varphi}_{n, ij}^I(s)| \\ &\leq \frac{a_n^2 \|\alpha_0\|_\infty \|K\|_\infty}{r_n p_n(s) \sqrt{h} \eta_0^2} \int_s^{s+2h} \frac{1}{r_n p_n(r)} \sum_{k, l \in V_n} \left| \Psi(X_{n, kl}(r); \beta_0) - \Psi(X_{n, kl}(r); \tilde{\beta}_n) \right| C_{n, kl}(r) dr \\ &\leq \frac{2a_n^2 \|\alpha_0\|_\infty \|K\|_\infty \sqrt{h}}{r_n p_n(s) \eta_0^2} L_\Psi \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) \sup_{r \in [s, s+2h]} \frac{1}{r_n p_n(r)} \sum_{k, l \in V_n} C_{n, kl}(r). \end{aligned} \quad (7.21)$$

Similarly, on the same event

$$\begin{aligned} &|\tilde{\varphi}_{n, ij}(s) - \tilde{\varphi}_{n, ij}^I(s)| \\ &\leq \frac{a_n^2 \|\alpha_0\|_\infty \|K\|_\infty}{r_n p_n(s) \sqrt{h} \eta_0^2} \int_s^{s+2h} \frac{1}{r_n p_n(r)} \sum_{k, l \in V_n} \left| \Psi(X_{n, kl}(r); \beta_0) - \Psi(X_{n, kl}(r); \tilde{\beta}_n) \right| \\ &\quad \times |1 - \mathbb{1}(d_{s-4h}^n(kl, I) \geq m)| \cdot C_{n, kl}(r) dr \\ &\leq \frac{2a_n^2 \|\alpha_0\|_\infty \|K\|_\infty \sqrt{h}}{r_n^2 p_n(s) p_n \eta_0^2} L_\Psi \omega_\Psi \left(\frac{c_2}{\sqrt{r_n p_n}} \right) |I| B_n^I. \end{aligned} \quad (7.22)$$

Note that in (7.21) and (7.22) we may drop the assumption that we are on the event $\|\tilde{\beta}_n - \beta_0\| \leq c_2/\sqrt{r_n p_n}$ when we replace $L_\Psi \omega_\Psi(c_1 \sqrt{\log r_n / r_n p_n})$ by $2\|\Psi\|_\infty$. With these inequalities and since

$$(7.20) = \sum_{i, j \in V_n} \int_0^T \tilde{\varphi}_{n, ij}(s) dM_{n, ij}(s)$$

we may apply Lemma 2.9 from Kreiss (2020+) to show that (7.20) = $o_P(1)$. Thus, we need to show the three condition from this proposition. We show them all by the same scheme: Add the indicator that $\|\tilde{\beta}_n - \beta_0\| \leq c_2/\sqrt{r_n p_n}$ or not and apply the corresponding bound. Then we can use that $\tilde{\beta}_n$ is constructed independently and split the expectation. We begin with the first condition: We apply (7.21) with $c_2 = c^* \sqrt{\log r_n}$

$$\begin{aligned}
& \sum_{i,j \in V_n} \int_0^T \mathbb{E} \left(\tilde{\varphi}_{n,ij}^{ij}(s)^2 C_{n,ij}(s) \alpha_0(s) \Psi(X_{n,ij}(s); \beta_0) \right) ds \\
& \leq \frac{4a_n^4 \|\alpha_0\|_\infty^3 \|K\|_\infty^2 h}{r_n p_n \eta_0^4} L_{\Psi}^2 \omega_{\Psi} \left(c^* \sqrt{\frac{\log r_n}{r_n p_n}} \right)^2 \|\Psi(\cdot; \beta_0)\|_\infty \\
& \quad \times \int_0^T \mathbb{E} \left(\left(\sup_{r \in [s, s+2h]} \frac{1}{r_n p_n(r)} \sum_{k,l \in V_n} C_{n,kl}(r) \right)^2 \middle| C_{n,ij}(s) = 1 \right) ds \\
& \quad + \frac{4a_n^4 \|\alpha_0\|_\infty^3 \|K\|_\infty^2 h}{r_n p_n \eta_0^4} \|\Psi\|_\infty^2 \|\Psi(\cdot; \beta_0)\|_\infty \cdot \mathbb{P} \left(\|\tilde{\beta}_n - \beta_0\| > c^* \sqrt{\frac{\log r_n}{r_n p_n}} \right) \\
& \quad \times \int_0^T \mathbb{E} \left(\left(\sup_{r \in [s, s+2h]} \frac{1}{r_n p_n(r)} \sum_{k,l \in V_n} C_{n,kl}(r) \right)^2 \middle| C_{n,ij}(s) = 1 \right) ds.
\end{aligned}$$

The above converges to zero because $a_n^2/r_n p_n = O(1)$, $hr_n p_n \omega_{\Psi} \left(\sqrt{\log r_n / r_n p_n} \right)^2 = o(1)$ and the boundedness. We continue with the second condition by using (7.21) and (7.22) in the same way for $c_2 = c^* \sqrt{\log r_n}$ (recall also that $B_n^{ij,kl} \leq B_n^{ij} B_n^{kl}$)

$$\begin{aligned}
& \sum_{ij,kl \in V_n} \mathbb{E} \left(\int_0^T \tilde{\varphi}_{n,ij}^{ij,kl}(t) dM_{n,ij}(t) \int_0^T (\varphi_{n,kl}(s) - \tilde{\varphi}_{n,kl}^{ij,kl}(s)) dM_{n,kl}(s) \right) \\
& \leq \frac{8a_n^4 \|\alpha_0\|_\infty^2 \|K\|_\infty^2 h}{r_n p_n \eta_0^4} L_{\Psi}^2 \omega_{\Psi} \left(c^* \sqrt{\frac{\log r_n}{r_n p_n}} \right)^2 \\
& \quad \times \mathbb{E} \left(\sup_{r \in [0, T]} \sum_{k,l \in V_n} \frac{C_{n,kl}(r)}{r_n p_n(r)} \left(\sum_{ij \in V_n} \int_0^T \frac{B_n^{ij}}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \right) \\
& \quad + \frac{8a_n^4 \|\alpha_0\|_\infty^2 \|K\|_\infty^2 h}{r_n p_n \eta_0^4} \|\Psi\|_\infty^2 \mathbb{P} \left(\|\tilde{\beta}_n - \beta_0\| > c^* \sqrt{\frac{\log r_n}{r_n p_n}} \right) \\
& \quad \times \mathbb{E} \left(\sup_{r \in [0, T]} \sum_{k,l \in V_n} \frac{C_{n,kl}(r)}{r_n p_n(r)} \left(\sum_{ij \in V_n} \int_0^T \frac{B_n^{ij}}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \right)
\end{aligned}$$

which converges to zero because $a_n^2/r_n p_n = O(1)$, $hr_n p_n \omega_{\Psi} \left(\sqrt{\log r_n / r_n p_n} \right)^2 = o(1)$ and

the boundedness of the expectation. And finally for the third assumption we use (7.22) again with $c_2 = c^* \sqrt{\log r_n}$

$$\begin{aligned}
& \sum_{ij,kl \in V_n} \mathbb{E} \left(\int_0^T \left(\tilde{\varphi}_{n,ij}(t) - \tilde{\varphi}_{n,ij}^{ij,kl}(t) \right) dM_{n,ij}(t) \int_0^T \left(\varphi_{n,kl}(s) - \tilde{\varphi}_{n,kl}^{ij,kl}(s) \right) dM_{n,kl}(s) \right) \\
& \leq \frac{16a_n^4 \|\alpha_0\|_\infty^2 \|K\|_\infty^2 h}{r_n^2 p_n^2 \eta_0^4} L_{\Psi}^2 \omega_{\Psi} \left(c^* \sqrt{\frac{\log r_n}{r_n p_n}} \right)^2 \mathbb{E} \left(\left(\sum_{ij \in V_n} \int_0^T \frac{(B_n^{ij})^2}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \right) \\
& \quad + \frac{16a_n^4 \|\alpha_0\|_\infty^2 \|K\|_\infty^2 h}{r_n^2 p_n^2 \eta_0^4} \|\Psi\|_\infty^2 \mathbb{P} \left(\|\tilde{\beta}_n - \beta_0\| > c^* \sqrt{\frac{\log r_n}{r_n p_n}} \right) \\
& \quad \quad \quad \times \mathbb{E} \left(\left(\sum_{ij \in V_n} \int_0^T \frac{(B_n^{ij})^2}{r_n p_n(t)} d|M_{n,ij}|(t) \right)^2 \right)
\end{aligned}$$

which converges to zero by the assumptions and the proof is complete. \square

7.2 Mixing

Proof of Proposition 6.2. The proof uses a chaining type argument: Let

$$T_{n,k} := \left\{ \left(\frac{a_j}{n^k} \right)_{j=0,\dots,q} : a_j \in \mathbb{Z} \cdot \frac{2}{\sqrt{1+q}}, j = 0, \dots, q \right\} \cap [0, T] \times B_n(c_2)$$

be a finite grid of $[0, T] \times B_n(c_2)$ of mesh $2n^{-k}/\sqrt{1+q}$. Then $\#T_{n,k} = O(n^{(1+q)k}/(r_n p_n)^{q/2})$ and denote by $\pi_{n,k} : [0, T] \times B_n(c_2) \rightarrow T_{n,k}$ the projection, i.e., for any $(t, \beta) \in [0, T] \times B_n(c_2)$ it holds that $\|(t, \beta) - \pi_{n,k}(t, \beta)\| \leq n^{-k}$. In order to avoid too complicated notation we write $\pi_{n,k}(t, \beta) = (\pi_{n,k}(t), \pi_{n,k}(\beta))$ such that $\pi_{n,k}(t) \in [0, T]$ is the time grid point which lies closest to t and $\pi_{n,k}(\beta) \in B_n(c_2)$ is the parameter grid point lying closest to β .

We begin by studying $\mathcal{A}_n(c_1, c_2)$ and rewrite the supremum in its definition in the following way (the following holds for all $c_1, c_2 > 0$)

$$\begin{aligned}
& \sup_{t \in [0, T], \beta \in B_n(c_2)} \sqrt{p_n(t)} \left| \frac{1}{r_n p_n(t)} X_n(t; \beta) - \bar{X}_n(t; \beta) \right| \\
& \leq \sup_{t \in [0, T], \beta \in B_n(c_2)} \left| \frac{1}{r_n \sqrt{p_n(t)}} X_n(t; \beta) - \frac{1}{r_n \sqrt{p_n(\pi_{n,k}(t))}} X_n(\pi_{n,k}(t), \beta) \right| \tag{7.23}
\end{aligned}$$

$$+ \sup_{t \in [0, T], \beta \in B_n(c_2)} \left| \sqrt{p_n(\pi_{n,k}(t))} \bar{X}_n(\pi_{n,k}(t), \beta) - \sqrt{p_n(t)} \bar{X}_n(t; \beta) \right| \tag{7.24}$$

$$+ \sup_{t \in [0, T], \beta \in B_n(c_2)} \left| \frac{1}{r_n \sqrt{p_n(\pi_{n,k}(t))}} X_n(\pi_{n,k}(t), \beta) - \sqrt{p_n(\pi_{n,k}(t))} \bar{X}_n(\pi_{n,k}(t), \beta) \right|. \tag{7.25}$$

Let now $c_2 > 0$ arbitrary but fixed. We note firstly that, by Holder continuity of p_n and \bar{X}_n , we can choose c_1 and k suitably such that (7.24) converges to zero with the prescribed rate. For (7.23), we note that

$$\begin{aligned} & \left| \frac{1}{r_n \sqrt{p_n(t)}} X_n(t; \beta) - \frac{1}{r_n \sqrt{p_n(\pi_{n,k}(t))}} X_n(\pi_{n,k}(t), \beta) \right| \\ &= \frac{1}{r_n} |X_n(t; \beta)| \cdot \left| \frac{1}{\sqrt{p_n(t)}} - \frac{1}{\sqrt{p_n(\pi_{n,k}(t))}} \right| + \frac{1}{r_n \sqrt{p_n(\pi_{n,k}(t))}} |X_n(t; \beta) - X_n(\pi_{n,k}(t), \beta)|. \end{aligned}$$

The first part converges to zero when k is chosen large enough by the assumption of Hoelder continuity of p_n and since

$$\frac{1}{r_n} X_n \leq \sup_{\beta \in B_n(c_2)} \|\Psi(\cdot; \beta)\|_\infty$$

is bounded by assumption. For the second part we note that for general $t_1, t_2 \in [0, T]$ and $\beta_1, \beta_2 \in B_n(c_2)$ (below $\#\text{jumps}(C_n)[t_1, t_2]$ denotes the number of jumps of the process C_n in the interval $[t_1, t_2]$)

$$\begin{aligned} & |X_n(t_1; \beta_1) - X_n(t_2; \beta_2)| \\ & \leq \sum_{(i,j) \in V_n \times V_n} \left[|C_{n,ij}(t_1) - C_{n,ij}(t_2)| \Psi(X_{n,ij}(t_1), \beta_1) \right. \\ & \quad \left. + C_{n,ij}(t_2) |\Psi(X_{n,ij}(t_1), \beta_1) - \Psi(X_{n,ij}(t_1), \beta_2) + \Psi(X_{n,ij}(t_1), \beta_2) - \Psi(X_{n,ij}(t_2), \beta_2))| \right] \\ & \leq \|\Psi(\cdot; \beta_1)\|_\infty \cdot \#\text{jumps}(C_n)[t, s] + r_n L_\Psi \omega_\Psi (\|\beta_1 - \beta_2\|) \\ & \quad + \sum_{(i,j) \in V_n \times V_n} L_\Psi(\beta_2) \tilde{\omega}_\Psi (\|X_{n,ij}(t_1) - X_{n,ij}(t_2)\|). \end{aligned}$$

Hence we obtain

$$\begin{aligned} & \sup_{t \in [0, T], \beta \in B(c_2)} \frac{1}{r_n \sqrt{p_n(\pi_{n,k}(t))}} |X_n(t; \beta) - X_n(\pi_{n,k}(t), \beta)| \\ & \leq \sup_{\beta \in B_n(c_2)} \frac{\|\Psi(\cdot; \beta_1)\|_\infty \cdot \#\text{jumps}(C_n)[n^{-k}]}{r_n \sqrt{p_n}} + \frac{L_\Psi \omega(n^{-k})}{\sqrt{p_n}} \\ & \quad + \sup_{\beta \in B_n(c_2)} L_\Psi(\beta) \cdot \sup_{t_1, t_2: |t_1 - t_2| \leq n^{-k}} \sum_{(i,j) \in V_n \times V_n} \frac{\tilde{\omega}_\Psi (\|X_{n,ij}(t_1) - X_{n,ij}(t_2)\|)}{r_n \sqrt{p_n}}. \end{aligned}$$

By assumption, we may choose k and c_1 large enough to obtain that the above exceeds $c_1 \sqrt{\log r_n / r_n}$ with probability at most δ . Let now k and c_1 be so large that all previous arguments hold and note that we will be fine as long as do not decrease them in the remainder of the proof. For this choice of k , the supremum (7.25) can be handled by the

exponential inequality in Lemma 2.16 in [Kreiss \(2020+\)](#) (we can apply this Lemma by Assumption (Mixing)). We have that for c_1 (after possibly increasing it further) chosen large enough (note that $\log r_n p_n(t) \leq \log r_n$)

$$\begin{aligned} & \mathbb{P} \left(\sup_{t \in [0, T], \beta \in B_n(c_2)} \sqrt{p_n(\pi_{n,k}(t))} \left| \frac{1}{r_n p_n(\pi_{n,k}(t, \beta))} X_n(\pi_{n,k}(t)) - \bar{X}_n(\pi_{n,k}(t, \beta)) \right| > c_1 \sqrt{\frac{\log r_n}{r_n}} \right) \\ & \leq \#T_{n,k} \sup_{t \in [0, T], \beta \in B_n(c_2)} \mathbb{P} \left(\left| \frac{1}{r_n p_n(t)} X_n(t, \beta) - \bar{X}_n(t; \beta) \right| > c_1 \sqrt{\frac{\log r_n}{r_n p_n(t)}} \right) \rightarrow 0. \end{aligned}$$

So we have proved the convergence rate of the process X_n itself. For the convergence of the reciprocals denote

$$\eta_0 := \inf_{\beta \in B_n(c_2)} \|\bar{X}_n(\cdot; \beta)\|.$$

We make the observation that on $\tilde{\mathcal{A}}_n(c_1, c_2)$

$$\begin{aligned} & \sqrt{p_n(s)} \left| \left(\frac{1}{r_n p_n(s)} X_n(s; \beta) \right)^{-1} - \bar{X}_n(s; \beta)^{-1} \right| \\ & = \left(\frac{1}{r_n p_n(s)} X_n(s; \beta) \right)^{-1} \bar{X}_n(s; \beta)^{-1} \sqrt{p_n(s)} \left| \frac{1}{r_n p_n(s)} X_n(s; \beta) - \bar{X}_n(s; \beta) \right| \\ & \leq \left(\eta_0 - c_1 \sqrt{\frac{\log r_n}{r_n p_n(s)}} \right)^{-1} \eta_0^{-1} \sqrt{p_n(s)} \left| \frac{1}{r_n p_n(s)} X_n(s; \beta) - \bar{X}_n(s; \beta) \right| \\ & \leq \left(\eta_0 - c_1 \sqrt{\frac{\log r_n}{r_n p_n(s)}} \right)^{-1} \eta_0^{-1} c_1 \sqrt{\frac{\log r_n}{r_n}} \end{aligned}$$

which implies the statement because $\eta_0^{-1} = O(1)$. \square

Proof of Lemma 6.4. We estimate

$$\mathbb{E} \left(\left(\frac{X_n(s, \beta)}{r_n p_n(s)} \right)^2 \right) \leq \sum_{k=0}^{\infty} (k+1) \mathbb{P} \left(\left(\frac{X_n(s, \beta)}{r_n p_n(s)} \right)^2 \geq k \right).$$

By using the exponential Inequality 2.16 in [Kreiss \(2020+\)](#), the probability to the right can be bounded by an exponentially decreasing function such that the sum converges. \square

References

- O. Aalen. Nonparametric inference for a family of counting processes. *Ann. Statist.*, 6(4): 701–726, 07 1978.
- P. K. Andersen and R. D. Gill. Cox’s regression model for counting processes: A large sample study. *Ann. Statist.*, 10(4):1100–1120, 12 1982.

- P. K. Andersen, O. Borgan, R. D. Gill, and N. Keiding. *Statistical Models Based on Counting Processes*. Springer, 1993.
- C. T. Butts. A relational event framework for social action. *Sociological Methodology*, 38(1):155–200, 2008.
- S. N. Cohen and R. J. Elliott. *Stochastic Calculus and Applications*. Birkhäuser, 2015.
- D. R. Cox. Regression models and life-tables. *Journal of the Royal Statistical Society. Series B (Methodological)*, 34(2):187–220, 1972.
- D. R. Cox. Partial likelihood. *Biometrika*, 62(2):269–276, 1975.
- W. Härdle and E. Mammen. Comparing nonparametric versus parametric regression fits. *Ann. Statist.*, 21(4):1926–1947, 12 1993.
- A. Kreiss. Correlation bounds, mixing and m-dependence under random time-varying network distances with an application to cox-processes. *Bernoulli (to appear)*, 2020+.
- A. Kreiss, E. Mammen, and W. Polonik. Nonparametric inference for continuous-time event counting and link-based dynamic network models. *Electron. J. Statist.*, 13(2):2764–2829, 2019.
- E. Mammen and J. P. Nielsen. A general approach to the predictability issue in survival analysis with applications. *Biometrika*, 94(4):873–892, 2007.
- T. Martinussen and T. H. Scheike. *Dynamic Regression Models for Survival Data*. Statistics for Biology and Health. Springer, 2006.
- W. Nelson. Hazard plotting for incomplete failure data. *Journal of Quality Technology*, 1(1):27–52, 1969.
- P. O. Perry and P. J. Wolfe. Point process modelling for directed interaction networks. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 75(5):821–849, 2013.
- H. Ramlau-Hansen. Smoothing counting process intensities by means of kernel functions. *Ann. Statist.*, 11(2):453–466, 06 1983.
- R. Rebolledo. Central limit theorems for local martingales. *Z. Wahrsch. verw. Geb.*, 51: 269–286, 1980.
- T. H. Scheike and T. Martinussen. On estimation and tests of time-varying effects in the proportional hazards model. *Scandinavian Journal of Statistics*, 31(1):51–62, 2004.