

On the local behavior of weighted orbital integrals and the affine Springer fibers

Zongbin CHEN

November 30, 2021

Abstract

We show the rationality of a generating series from the affine Springer fibers for the group GL_d . The main ingredient is the homogeneity of the Arthur-Shalika germ expansion for the weighted orbital integrals.

Contents

1	Introduction	1
2	The weighted orbital integrals	7
3	Homogeneity of the Arthur-Shalika germ expansion	9
3.1	The Shalika germ expansion	9
3.2	The Arthur-Shalika expansion	11
3.3	The parabolic descent	13
4	Counting points on the fundamental domain	18
5	Rationality from homogeneity	24
5.1	Generating series of weighted orbital integrals	24
5.2	Generating series of the affine Springer fibers	28

1 Introduction

Let $k = \mathbb{F}_q$ be a finite field, $F = k((\epsilon))$ the field of Laurent series, $\mathcal{O} = k[[\epsilon]]$ the ring of integers of F . Let $\text{val} : F^\times \rightarrow \mathbf{Z}$ be the standard valuation on F and let $|\cdot|$ be the associated norm. Let $\mathbf{G} = GL_{d+1}$ with $d < p = \text{char}(k)$, let \mathbf{A} be the maximal torus of \mathbf{G} of the diagonal matrices, let \mathbf{B} be the Borel subgroup of \mathbf{G} consisting of the upper triangular matrices. For a closed algebraic subgroup \mathbf{H} of \mathbf{G} over k , we denote its Lie algebra by the corresponding gothic letter and we denote $H = \mathbf{H}(F)$.

Let $\gamma \in \mathfrak{g}$ be a regular semisimple element, recall that the *affine Springer fiber* at γ is the closed sub ind- k -scheme of the *affine grassmannian* $\mathcal{X} = \mathbf{G}((\epsilon))/\mathbf{G}[[\epsilon]]$ defined by

$$\mathcal{X}_\gamma = \{g \in \mathcal{X} \mid \text{Ad}(g^{-1})\gamma \in \mathfrak{g}[[\epsilon]]\}.$$

It is a finite dimensional k -scheme with a large group of symmetry. Let T be the centralizer of γ in G , then T acts on \mathcal{X}_γ by left translation. It is known that T acts transitively on a dense open subscheme $\mathcal{X}_\gamma^{\text{reg}}$ of \mathcal{X}_γ . In particular, all the irreducible components of \mathcal{X}_γ are isomorphic.

Conjecture 1.1 (Goresky-Kottwitz-MacPherson [GKM1]). *The cohomology of \mathcal{X}_γ is pure in the sense of Deligne.*

In an attempt to attack the conjecture, we [C1] are led to construct a replacement of the irreducible components of the affine Springer fiber, called the *fundamental domain* of the affine Springer fiber, denoted F_γ . We are able to construct an associated *Arthur-Kottwitz reduction*, which is a decomposition of the complement $\mathcal{X}_\gamma \setminus F_\gamma$ into union of locally closed subschemes which admits iterated affine fibrations onto the fundamental domains F_γ^M for the Levi subgroups M of G containing T . In particular, the cohomological purity of the affine Springer fibers are reduced to that of their fundamental domains.

In the split case, according to Chaudouard and Laumon [CL1], under the purity hypothesis, the cohomology of F_γ depends only on the moment graph of the T -action on it. In other words, it depends only on the *root valuation datum* of γ . (It can be shown that there exists a set of simple roots $\{\alpha_i\}_{i=1}^d$ in the root system $\Phi(G, T)$ such that for any root $\alpha = \sum_{l=i}^j \alpha_l$, we have $\text{val}(\alpha(\gamma)) = \min\{\alpha_l(\gamma)\}_{l=i}^j$. The ordered d -tuple $(\text{val}(\alpha_l(\gamma)))_{l=1}^d$ is called the *root valuation datum* of γ .) In [C1], we look carefully into the geometry of F_γ for the group GL_2 and GL_3 . Our investigation, together with the calculations of [CL1], leads us to the conjecture:

Conjecture 1.2 ([C1]). *Let $\gamma \in \mathfrak{a}$ be a regular element with root valuation datum $\mathbf{n} \in \mathbf{Z}_{\geq 0}^d$. Then the Poincaré polynomial of F_γ depends only on \mathbf{n} , we denote it by $P_{\mathbf{n}}(q)$. Moreover, the generating series*

$$P(q; \vec{t}) := \sum_{n_1=0}^{+\infty} \cdots \sum_{n_d=0}^{+\infty} P_{(n_1, \dots, n_d)}(q) t_1^{n_1} \cdots t_d^{n_d} \in \mathbf{Z}[[q; t_1, \dots, t_d]]$$

is a rational fraction, i.e. it is an element of $\mathbf{Z}(q; t_1, \dots, t_d)$.

This is a very strong finiteness statement about the affine Springer fibers. It implies that to understand all the affine Springer fibers, it is enough to understand a finite number of them. The conjecture seems to be out of reach by direct attack, for the lack of effective ways to calculate the Poincaré polynomials of F_γ . Instead, we turn to a closely related problem—counting the number of points on F_γ . Our main result states:

Theorem 1.1. *Let $\gamma \in \mathfrak{a}$ be a regular element with root valuation datum $\mathbf{n} \in \mathbf{Z}_{\geq 0}^d$. Then the number of points $|F_\gamma(\mathbb{F}_q)|$ is a polynomial in q depending only on \mathbf{n} , denoted $F_{\mathbf{n}}(q)$. Moreover, the generating series*

$$F(q; \vec{t}) := \sum_{n_1=0}^{+\infty} \cdots \sum_{n_d=0}^{+\infty} F_{(n_1, \dots, n_d)}(q) t_1^{n_1} \cdots t_d^{n_d} \in \mathbf{Z}[[q; t_1, \dots, t_d]]$$

is a rational fraction, i.e. it is an element of $\mathbf{Z}(q; t_1, \dots, t_d)$.

Assuming the purity hypothesis, the theorem is actually equivalent to the conjecture. Indeed, as $|F_\gamma(\mathbb{F}_q)|$ is polynomial in q , we have $|F_\gamma(\mathbb{F}_q)| = P_{\mathbf{n}}(q)$ by the purity hypothesis, whence the equivalence of the two assertions. It will be interesting to see whether the rationality behavior of the counting points implies the purity hypothesis.

The proof of theorem 1.1 involves the study of the local behavior of the weighted orbital integrals. In [C2], we have explained a transition between counting points on the fundamental domain and the weighted orbital integrals. But unfortunately the recursion that we discovered in *loc. cit.* is hard to solve. Here we proceed differently and we are able to write down an explicit transition between $|F_\gamma^M(\mathbb{F}_q)|$ and the weighted orbital integral $J_A^M(\gamma, \mathbb{1}_{\mathfrak{g}(\mathcal{O})})$, with M running through the Levi subgroups of G containing A , see theorem 4.5 and 4.7. The question is then reduced to a similar rationality assertion for the weighted orbital integrals, which is also essentially a finiteness statement as we have explained for the generating series $P(q; \vec{t})$. Recall that the weighted orbital integrals do satisfy a finiteness property—the *Arthur-Shalika germ expansion*. In the group-theoretic setting, Arthur [A3] defined the weighted orbital integrals for the unipotent elements, and showed that the weighted orbital integrals can be expanded in terms of the unipotent ones. In [W], Waldspurger adapted Arthur’s work to the Lie-algebraic setting. But unfortunately, both authors made no efforts to determine the domain of validity of the expansion. We fill in this gap. Let M be a Levi subgroup of G containing γ , let $\mathcal{L}(M)$ be the set of Levi subgroups of G containing M . For $L \in \mathcal{L}(M)$, let \mathcal{N}_L be the nilpotent cone of \mathbf{L} , and let $[\mathcal{N}_L(F)]$ be a set of representatives for the $\mathbf{L}(F)$ -conjugacy classes in $\mathcal{N}_L(F)$. An element $\gamma \in \mathfrak{g}$ is said to be *integral* if its characteristic polynomial has coefficients in \mathcal{O} .

Theorem 1.2. *For each element $u \in [\mathcal{N}_L(F)]$, $L \in \mathcal{L}(M)$, there exists unique function $g_M^L(\cdot, u)$ on $\mathfrak{m} \cap \mathfrak{g}^{\text{rs}}$ such that for any $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{g}(\mathcal{O}))$ and any integral element $\gamma \in \mathfrak{m} \cap \mathfrak{g}^{\text{rs}}$, we have*

$$J_M(\gamma, f) = \sum_{L \in \mathcal{L}(M)} \sum_{u \in [\mathcal{N}_L(F)]} g_M^L(\gamma, u) J_L(u, f).$$

This expansion will be called *Arthur-Shalika germ expansion*, and the functions g_M^L will be called *Arthur-Shalika germs*. Consider the scaling action on \mathfrak{g} , which preserves the nilpotent cone and in fact all the nilpotent orbits. By the uniqueness of the Arthur-Shalika germ expansion, we can deduce a homogeneous property of the Arthur-Shalika germs with respect to the scaling action, see theorem 3.7. In general, we can consider the germ expansion around an element $s \in \mathfrak{a}_M$. This is achieved by a careful analysis of the parabolic descent of the weighted orbital integrals. We refer the reader to theorem 3.11 for more details.

Coming back to the proof of theorem 1.1. As we have explained before, it is equivalent to a rationality statement for a similar generating series of the weighted orbital integrals $J_A(\gamma, \mathbb{1}_{\mathfrak{g}(\mathcal{O})})$. With the Arthur-Shalika germ expansion, this is further reduced to the rationality of a similar generating series for the Arthur-Shalika germs.

Theorem 1.3. *For $M \in \mathcal{L}(A)$, $u \in \mathcal{N}_M(F)$, $\gamma \in \mathfrak{a}$ regular, the Arthur-Shalika germ $g_A^M(\gamma, u)$ depends only on the root valuation datum \mathbf{n} of γ , and we denote it by $g_A^M(\mathbf{n}, u)$. Moreover, the generating series*

$$g_A^M(u; \vec{t}) := \sum_{n_1=0}^{+\infty} \cdots \sum_{n_d=0}^{+\infty} g_A^M((n_1, \dots, n_d), u) t_1^{n_1} \cdots t_d^{n_d} \in \mathbf{C}[[t_1, \dots, t_d]]$$

is a rational fraction, i.e. it is an element of $\mathbf{C}(t_1, \dots, t_d)$.

We prove it by induction on the root valuation datum of γ and on the semisimple rank of \mathbf{G} , using the homogeneity and the parabolic descent of the Arthur-Shalika germs, see theorem 5.4.

Notations and conventions

The structures from the group The algebraic subgroups \mathbf{H} of \mathbf{G} are always assumed to be defined over k , and we use the convention that $H = \mathbf{H}(F)$. To simplify the notations, we will not refer to the group \mathbf{H} if the context is clear. This applies particularly to the Levi subgroups and the parabolic subgroups.

Let M be a Levi subgroup of G . We denote by $\mathcal{P}(M)$ the set of parabolic subgroups of G whose Levi factor is M , by $\mathcal{L}(M)$ the set of Levi subgroups of G containing M , and by $\mathcal{F}(M)$ the set of parabolic subgroups of G containing M . For $P \in \mathcal{P}(M)$, we denote by $P^- \in \mathcal{P}(M)$ the opposite of P with respect to M . Let

$$X^*(M) = \text{Hom}(M, \mathbb{G}_m), \quad X_*(M) = \text{Hom}(X^*(M), \mathbf{Z}).$$

Let $\mathfrak{a}_M^* = X^*(M) \otimes \mathbf{R}$ and $\mathfrak{a}_M = X_*(M) \otimes \mathbf{R}$. Let A_M be the central connected component of the center of M , then the restriction $X^*(M) \rightarrow X^*(A_M)$ induces an isomorphism $\mathfrak{a}_M^* \cong X^*(A_M) \otimes \mathbf{R}$. For a Levi subgroup $L \in \mathcal{L}(M)$, the restriction $X^*(L) \rightarrow X^*(M)$ induces an injection $\mathfrak{a}_L^* \hookrightarrow \mathfrak{a}_M^*$, hence a dual surjection $\mathfrak{a}_M \rightarrow \mathfrak{a}_L$. On the other hand, the inclusion $A_L \subset A_M$ induces an inclusion $X_*(A_L) \otimes \mathbf{R} \hookrightarrow X_*(A_M) \otimes \mathbf{R}$ and a dual surjection $X^*(A_M) \otimes \mathbf{R} \rightarrow X^*(A_L) \otimes \mathbf{R}$. Taken together, we get splittings

$$\mathfrak{a}_M = \mathfrak{a}_M^L \oplus \mathfrak{a}_L, \quad (\mathfrak{a}_M)^* = (\mathfrak{a}_M^L)^* \oplus (\mathfrak{a}_L)^*.$$

Let π^L, π_L be the projections to the two factors for these two splittings.

To define Arthur's weight factor $v_M(g)$, we need to choose a Lebesgue measure on \mathfrak{a}_M^G . We fix a W -invariant positive definite inner product $\langle \cdot, \cdot \rangle$ on the vector space \mathfrak{a}_A^G , where W is the Weyl group of G with respect to A . Notice that \mathfrak{a}_A^M and \mathfrak{a}_M are orthogonal to each other with respect to the inner product for any $M \in \mathcal{L}(A)$. We fix a Lebesgue measure on \mathfrak{a}_M^G normalised by the condition that the lattice generated by the orthonormal bases in \mathfrak{a}_M^G has covolume 1.

For $P \in \mathcal{F}(A)$, let M_P be its Levi factor containing A and let N_P be its unipotent radical. Let Φ_P be the set of roots of (P, A_P) for $A_P := A_{M_P}$, let $\overline{\Phi}_P$ be the reduced root system and let Δ_P be the set of simple roots in $\overline{\Phi}_P$, they can be seen as elements in $(\mathfrak{a}_M^G)^*$. Let Δ_P^\vee be the associated simple coroots.

(\mathbf{G}, M)-family We need Arthur's notion of (G, M) -family. The reader can find a beautiful exposition in [A4], §17. It is a family of smooth functions $(r_P(\lambda))_{P \in \mathcal{P}(M)}$ on $i\mathfrak{a}_M^* \subset \mathfrak{a}_M^* \otimes \mathbf{C}$ which satisfy for any adjacent parabolic subgroups (P, P') the property that $r_P(\lambda) = r_{P'}(\lambda)$ for any λ on the hyperplane defined by the unique coroot $\beta_{P, P'}^\vee$ in $\Delta_P^\vee \cap (-\Delta_{P'}^\vee)$. An important source of example comes from the (G, M) -orthogonal sets. A collection $r = (r_P)_{P \in \mathcal{P}(M)}$ with $r_P \in \mathfrak{a}_M$ is said to be a (G, M) -orthogonal set if for any adjacent parabolic subgroups $P, P' \in \mathcal{P}(M)$,

$$r_P - r_{P'} = c_{P, P'} \cdot \beta_{P, P'}^\vee,$$

for some $c_{P, P'} \in \mathbf{R}$. It is said to be *positive* if all the $c_{P, P'}$ are positive. Notice that the family of function $r = (r_P(\lambda))_{P \in \mathcal{P}(M)}$ with

$$r_P(\lambda) := e^{\langle \lambda, r_P \rangle}, \quad \lambda \in i\mathfrak{a}_M^*,$$

forms a (G, M) -family, called the (G, M) -family associated to the (G, M) -orthogonal set.

For $Q = LN_Q \in \mathcal{F}(M)$, let

$$r_R^Q(\lambda) = r_{RN_Q}(\lambda), \quad \forall R \in \mathcal{P}^L(M).$$

It is easy to see that $(r_R^Q(\lambda))_{R \in \mathcal{P}^L(M)}$ form a (L, M) -family. We call it the Q -facet of r , denoted by r^Q . Notice that it can be viewed as a (G, M) -family, denoted $\iota(r^Q)$, by $\iota(r^Q)_P(\lambda) = r_{P \cap L}^Q(\lambda)$ for $P \in \mathcal{P}(M)$. On the other hand, let $P \in \mathcal{P}(M)$ be a parabolic subgroup contained in Q , it is easy to see that the restriction of the function $r_P(\lambda)$ to \mathfrak{a}_L^* doesn't depend on P , we denote it by $r_Q(\lambda)$. It is easy to see that the family $(r_Q(\lambda))_{Q \in \mathcal{P}(L)}$ forms a (G, L) -family, we denote it by $\pi_L(r)$. Similar construction works for the (G, M) -orthogonal sets, we preserve the same notations for them. Notice that there is no ambiguity about the notation $\pi_L(r)$ because it is also the image of r under the projection $\pi_L : \mathfrak{a}_M \rightarrow \mathfrak{a}_L$.

In case that the (G, M) -orthogonal set r is positive, we denote by $\mathbf{E}(r)$ the convex hull of the points r_P , it is a convex polytope in \mathfrak{a}_M . Its volume with respect to the Lebesgue measure on \mathfrak{a}_M can be calculated to be

$$r_M = \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} d_P(\lambda)^{-1} r_P(\lambda),$$

with $d_P(\lambda) = \text{vol}(\mathfrak{a}_M^G / \mathbf{Z}(\Delta_P^\vee))^{-1} \prod_{\alpha \in \Delta_P} \lambda(\alpha^\vee)$. For a general (G, M) -family $(r_P(\lambda))_{P \in \mathcal{P}(M)}$, we define

$$r_M(\lambda) = \sum_{P \in \mathcal{P}(M)} d_P(\lambda)^{-1} r_P(\lambda),$$

for generic $\lambda \in i\mathfrak{a}_M^*$. It can be shown that the function extends smoothly over all $i\mathfrak{a}_M^*$. Let

$$r_M = \lim_{\lambda \rightarrow 0} r_M(\lambda) = \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} d_P(\lambda)^{-1} r_P(\lambda), \quad (1.1)$$

and we call it the *volume* of the (G, M) -family $(r_P(\lambda))_{P \in \mathcal{P}(M)}$.

Let c and d be two (G, M) -orthogonal sets, we define their sum $c + d$ to be

$$(c + d)_P = c_P + d_P, \quad \forall P \in \mathcal{P}(M).$$

Obviously the sum remains a (G, M) -orthogonal set. Similarly, we can define the product of two (G, M) -families. If both c and d are positive, and that $0 \in \text{Ec}(d)$, we have inclusion of polytopes $\mathbf{E}(c) \subset \mathbf{E}(c + d)$, and the complement $\mathbf{E}(c + d) \setminus \mathbf{E}(c)$ can be partitioned as: For $Q = LN_Q \in \mathcal{F}(M)$, let

$$\mathfrak{a}_Q^+ = \{a \in \mathfrak{a}_M \mid \alpha(a) > 0, \forall \alpha \in \Delta_Q; \beta(a) = 0, \forall \beta \in \Phi(L, A_M)\}.$$

Then we have disjoint partition

$$\mathfrak{a}_M = \bigsqcup_{Q \in \mathcal{F}(M)} \mathfrak{a}_Q^+.$$

Let $\mathbf{E}^Q(c)$ be the Q -facet of the polytope $\mathbf{E}(c)$, let $R_Q(c) = \mathbf{E}^Q(c) + \mathfrak{a}_Q^+$, then

$$\mathfrak{a}_M = \bigsqcup_{Q \in \mathcal{F}(M)} R_Q(c).$$

Let $R_Q(c, d) = R_Q(c) \cap \mathbf{E}(c + d)$, we get a decomposition

$$\mathbf{E}(c + d) = \bigsqcup_{Q \in \mathcal{F}(M)} R_Q(c, d),$$

with $R_G(c, d) = \mathbf{E}(c)$. The decomposition enables us to express the volume of $\mathbf{E}(c + d)$ in terms of that of c and d .

The construction generalizes to the (G, M) -families. Let r, s be two (G, M) -families. Then the collection $\{r_P(\lambda)s_P(\lambda)\}_{P \in \mathcal{P}(M)}$ forms a (G, M) -family. We denote it by $r \cdot s$ and call it the product of r and s . From the (G, M) -family $\{s_P(\lambda)\}_{P \in \mathcal{P}(M)}$, Arthur has defined a smooth function $s'_Q(\lambda)$ on \mathfrak{a}_Q^* for $Q \in \mathcal{F}(M)$, which is too complicated to recall here and we simply refer the reader to [A4], equation (17.7). Let $s'_Q = s'_Q(0)$. Then Arthur's formula for the volume of the product of two (G, M) -families states

$$(r \cdot s)_M = \sum_{Q \in \mathcal{F}(M)} r_M^Q \cdot s'_Q. \quad (1.2)$$

Setting $r = 1$, we get

$$s_M = \sum_{P \in \mathcal{P}(M)} s'_P. \quad (1.3)$$

In case that the (G, M) -family r satisfies the extra condition

$$r_M^L = r_M^Q, \quad \forall L \in \mathcal{L}(M), Q \in \mathcal{P}(L),$$

the above product formula further simplifies to

$$(r \cdot s)_M = \sum_{L \in \mathcal{L}(M)} r_M^L \cdot s_L. \quad (1.4)$$

For $L \in \mathcal{L}(M)$, we will need a descent formula for the volume r_L of the (G, L) -family $(r_R(\lambda))_{R \in \mathcal{P}(L)}$. For $L' \in \mathcal{L}(M)$, we define $\theta_M^G(L, L')$ to be 0 unless the natural map

$$\mathfrak{a}_M^L \oplus \mathfrak{a}_M^{L'} \rightarrow \mathfrak{a}_M^G$$

is an isomorphism, in which case $\theta_M^G(L, L')$ is the factor by which the product Haar measure on $\mathfrak{a}_M^L \oplus \mathfrak{a}_M^{L'}$ must be multiplied in order to be equal to the Haar measure on \mathfrak{a}_M^G . Let ξ be a fixed generic vector in \mathfrak{a}_M^L , then for any $L' \in \mathcal{L}(M)$ satisfying $\theta_M^G(L, L') \neq 0$, the affine space $\xi + \mathfrak{a}_M^G$ intersects non-trivially with $\mathfrak{a}_{L'}^G \subset \mathfrak{a}_M^G$ at one point, which belongs to a chamber \mathfrak{a}_Q^+ in $\mathfrak{a}_{L'}^G$, for a unique $Q \in \mathcal{P}(L')$. We will denote it by $Q_{L'}^\xi$. Then

$$r_L = \sum_{L' \in \mathcal{L}(M)} \theta_M^G(L, L') \cdot r_M^{Q_{L'}^\xi}. \quad (1.5)$$

Measures on the groups To define the weighted orbital integrals in a uniform way, we need to make a uniform choice of Haar measure on G and its closed subgroups. Let dx be the Haar measure on F with $dx(\mathcal{O}) = 1$. Let \mathbf{H} be a connected closed algebraic subgroup of \mathbf{G} over F . Following Weil [We], let $\omega_{\mathfrak{h}} \in \wedge^{\text{top}} \mathfrak{h}$ be a volume form on \mathfrak{h} , then it defines a Haar measure $|\omega_{\mathfrak{h}}|$ on \mathfrak{h} . By left translation, $\omega_{\mathfrak{h}}$ defines a left-invariant volume form $\omega_{\mathbf{H}}$ on \mathbf{H} . This determines a left-invariant measure $|\omega_{\mathbf{H}}|$ on H . We normalize $|\omega_{\mathfrak{h}}|$ and $|\omega_{\mathbf{H}}|$ by taking a volume form $\omega_{\mathfrak{h}}$ such that $\text{vol}_{|\omega_{\mathbf{H}}|}(H \cap \mathbf{G}(\mathcal{O})) = 1$. We denote the resulting measure by dh if the context is clear. Given an element $\gamma \in \mathfrak{g}$, it is known that its stabilizer G_γ for the adjoint action of G is unimodular, hence G_γ carries a normalized Haar measure dh and we can define an invariant measure $\frac{dq}{dh}$ on the G -orbit of γ .

Moy-Prasad subgroups Let $\mathcal{B}(G)$ be the Bruhat-Tits building of G . For any point $x \in \mathcal{B}(G)$, Moy and Prasad [MP] defines a decreasing exhaustive filtration $\{\mathfrak{g}_{x,r}\}_{r \in \mathbf{R}}$ of \mathfrak{g} . For $r \geq 0$, $\mathfrak{g}_{x,r}$ is an ideal of \mathfrak{g} , let $G_{x,r}$ be the associated Lie group, it is a normal subgroup of G . Let $\mathfrak{g}_{x,r+} = \bigcup_{t > r} \mathfrak{g}_{x,t}$ and let $\mathfrak{g}_x(r) = \mathfrak{g}_{x,r}/\mathfrak{g}_{x,r+}$. For $x = 0$, we simplify $\mathfrak{k} = \mathfrak{g}_{0,0}$ and $\mathfrak{k}_r = \mathfrak{g}_{0,r}$, and accordingly $K = G_{0,0}$, $K_r = G_{0,r}$ for $r \geq 0$. For $r = 0$, we abbreviate $\mathfrak{g}_x = \mathfrak{g}_{x,0}$ and accordingly $G_x = G_{x,0}$. Let

$$\mathfrak{g}_r = \bigcup_{x \in \mathcal{B}(G)} \mathfrak{g}_{x,r}, \quad r \in \mathbf{R}.$$

Let $\mathcal{C}_c^\infty(\mathfrak{g})$ be the space of compactly supported locally constant functions on \mathfrak{g} . We define its subspaces

$$\mathcal{H}_r = \sum_{x \in \mathcal{B}(G)} \mathcal{C}_c(\mathfrak{g}/\mathfrak{g}_{x,r}).$$

Similar definitions for \mathfrak{g}_{r+} and \mathcal{H}_{r+} .

Connected components of the affine grassmannian We identify $X_*(A)$ with $A(F)/A(\mathcal{O})$ by sending χ to $\chi(\epsilon)$. With this identification, the canonical surjection $A(F) \rightarrow A(F)/A(\mathcal{O})$ can be viewed as

$$A(F) \rightarrow X_*(A). \quad (1.6)$$

We use Λ_G to denote the quotient of $X_*(A)$ by the coroot lattice of G (the subgroup of $X_*(A)$ generated by the coroots of A in G). According to Kottwitz [K1], we have a canonical homomorphism

$$\nu_G : G \rightarrow \Lambda_G, \quad (1.7)$$

which is characterized by the following properties: it is trivial on the image of G_{sc} in G (\mathbf{G}_{sc} is the simply connected cover of \mathbf{G}^{der}), and its restriction to A coincides with the composition of (1.6) with the projection of $X_*(A)$ to Λ_G . Since the morphism (1.7) is trivial on K , it descends to a map

$$\nu_G : \mathcal{X} \rightarrow \Lambda_G,$$

whose fibers are the connected components of \mathcal{X} . For $\mu \in \Lambda_G$, we denote the connected component $\nu_G^{-1}(\mu)$ by \mathcal{X}^μ .

2 The weighted orbital integrals

We make a brief recall on the weighted orbital integrals for the Lie algebra, following Waldspurger [W]. For the weighted orbital integrals, we can work in a broader context: Let F be a complete discrete valued field, let \mathbf{G} be a connected reductive algebraic group over F . Assume that $\text{char}(F)$ is either 0 or large enough with respect to $\text{rk}(\mathbf{G})$. We fix a minimal Levi subgroup $\mathbf{M}_{\text{min}} \subset \mathbf{G}$, and a special vertex in the apartment associated to $A_{\mathbf{M}_{\text{min}}}$ in the Bruhat-Tits building $\mathcal{B}(G)$ of G , let K be the stabilizer in G of this special vertex. We work exclusively with the Levi subgroups $M \in \mathcal{L}(M_{\text{min}})$.

For $P = MN \in \mathcal{P}(M)$, the Iwasawa decomposition $G = NMK$ defines a parabolic reduction on the affine grassmannian:

$$f_P : \mathcal{X} \rightarrow \mathcal{X}^M, \quad [g] = [nmk] \mapsto [m],$$

Let $H_M : M \rightarrow \mathfrak{a}_M$ be the unique group homomorphism¹ satisfying

$$\chi(H_M(m)) = \text{val}(\chi(m)), \quad \forall \chi \in X^*(M), m \in M.$$

Notice that it is invariant under the right K -action, so it induces a map from \mathcal{X}^M to \mathfrak{a}_M , still denoted by H_M . Let $H_P : \mathcal{X} \rightarrow \mathfrak{a}_M$ be the composition

$$H_P : \mathcal{X} \xrightarrow{f_P} \mathcal{X}^M \xrightarrow{H_M} \mathfrak{a}_M.$$

It has been shown by Arthur [A1] that for any point $x \in \mathcal{X}$, the reductions

$$\text{Ec}_M(x) := (H_P(x))_{P \in \mathcal{P}(M)}$$

forms a positive (G, M) -orthogonal set. Let

$$v_P(\lambda, x) = e^{\langle \lambda, H_P(x) \rangle}, \quad \lambda \in i\mathfrak{a}_M^*,$$

then $(v_P(\lambda, x))_{P \in \mathcal{P}(M)}$ forms a (G, M) -family. Let $v_M(x)$ be its volume as defined by the equation (1.1), we call it *Arthur's weight factor*.

Let $\gamma \in \mathfrak{m}$ be an element satisfying $G_\gamma = M_\gamma$. For $\varphi \in C_c^\infty(\mathfrak{g})$, the *weighted orbital integral* $J_M(\gamma, \varphi)$ for γ is defined to be

$$J_M(\gamma, \varphi) = |D(\gamma)|^{1/2} \int_{G_\gamma \backslash G} \varphi(\text{Ad}(g^{-1})\gamma) v_M(g) \frac{dg}{dt}, \quad (2.1)$$

which is well-defined because v_M is left M -invariant and $G_\gamma = M_\gamma$ is contained in M . Here $D(\gamma)$ is the *generalized Weyl discriminant*

$$D(\gamma) := \det(\text{ad}(\gamma_s) | \mathfrak{g}/\mathfrak{g}_{\gamma_s}),$$

where γ_s is the semisimple part of γ .

In general, without the assumption that $G_\gamma = M_\gamma$, we can perturb γ by an element $a \in \mathfrak{a}_M^{\text{reg}}$ sufficiently close to 0 such that $G_{\gamma+a} = M_{\gamma+a}$, and define $J_M(\gamma, \varphi)$ by a limit process. More precisely, let $\gamma = \gamma_s + \gamma_u$ be the Jordan decomposition, define

$$J_M(\gamma, \varphi) = \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \sum_{L \in \mathcal{L}(M)} r_M^L(\exp(\gamma_u), \exp(a)) \cdot J_L(\gamma + a, \varphi), \quad (2.2)$$

where r_M^L is the volume of a (G, M) -family constructed by Arthur [A3], §5. The existence of the above limit is one of the main results of Arthur [A3], and can be adapted to the Lie algebraic setting without difficulty. We simplify the notation $r_M^L(\exp(\gamma_u), \exp(a))$ to $r_M^L(\gamma_u, a)$ if there is no risk of confusion.

We want to emphasize the approximation of γ by $\gamma + a$ with $a \in \mathfrak{a}_M^{\text{reg}}$ in the definition (2.2). This is to ensure that the M -orbit of $\gamma + a$ approaches the M -orbit of γ . If we used the same approximation to calculate $J_L(\gamma, \varphi)$ for $L \in \mathcal{L}(M)$, we will arrive at $J_L(\gamma_M^L, \varphi)$ with γ_M^L the Lusztig-Spaltenstein induction of γ , because the L -orbit of $\gamma + a$ will approach the L -orbit of γ_M^L rather than that of γ . Recall that the *Lusztig-Spaltenstein induction* γ_M^G of an element $\gamma \in \mathfrak{m}$ is the union of the G -orbits $\{\gamma_i\}_G$ which intersects the coset $\gamma + \mathfrak{n}$ in a dense open subset, where \mathfrak{n} is the Lie algebra of the unipotent radical of a parabolic subgroup $P \in \mathcal{P}(M)$. There are only

¹Our definition differs from the conventional one by a minus sign.

finitely many such orbits and they belong to the same geometric orbit. They can be characterized analytically by the formula

$$J_G(\gamma_M^G, \varphi) := \sum_i J_G(\gamma_i, \varphi) = \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} J_G(\gamma + a, \varphi), \quad \forall \varphi \in \mathcal{C}_c^\infty(\mathfrak{g}),$$

which is a reflection of the geometric fact that the G -orbit of $\gamma + a$ approaches the G -orbits in γ_M^G when $a \in \mathfrak{a}_M^{\text{reg}} \mapsto 0$. In general, let $\gamma_M^L = \bigsqcup_i \{\gamma'_i\}_L$ be the decomposition into L -orbits, we define

$$J_L(\gamma_M^L, \varphi) = \sum_i J_L(\gamma'_i, \varphi).$$

This is well-defined because $J_L(\gamma'_i, \varphi)$ depends only on the L -orbit of γ'_i . Then following the same argument as in proof of [A3], cor. 6.3, we can show

$$\lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \sum_{L' \in \mathcal{L}(L)} r_{L'}^L(\gamma_u, a) \cdot J_{L'}(\gamma + a, \varphi) = J_L(\gamma_M^L, \varphi). \quad (2.3)$$

3 Homogeneity of the Arthur-Shalika germ expansion

The great advantage of definition (2.2) is that $J_M(\gamma, \varphi)$ has the same behavior under conjugation, whether $\gamma \in \mathfrak{m}$ satisfies the condition $M_\gamma = G_\gamma$ or not. For $\varphi \in \mathcal{C}_c^\infty(\mathfrak{g})$, $y \in G$, let

$$\varphi^y(X) = \varphi(\text{Ad}(y)X), \quad \forall X \in \mathfrak{g}.$$

Following the same lines as [A3], Lem. 8.1, one can show

$$J_M(\gamma, \varphi^y) - J_M(\gamma, \varphi) = \sum_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} J_M^{MQ}(\gamma, \varphi_{Q,y}), \quad (3.1)$$

where

$$\varphi_{Q,y}(Y) = \int_{\mathfrak{n}_Q} \int_K \varphi(\text{Ad}(k^{-1})(Y + n)) v'_Q(ky) dk dn, \quad \forall Y \in \mathfrak{m}_Q.$$

Despite the complexity of the expression, the function $\varphi_{Q,y}$ has the nice property:

$$\text{If } \varphi \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_r), \text{ then } \varphi_{Q,y} \in \mathcal{C}_c(\mathfrak{m}_Q/\mathfrak{m}_Q \cap \mathfrak{k}_r), \quad \forall r \in \mathbf{R}, \quad (3.2)$$

because \mathfrak{k}_r is stable under the adjoint action of K . As a consequence of this nice behavior, we can deduce the Arthur-Shalika germ expansion.

3.1 The Shalika germ expansion

We make a brief recall on the Shalika germ expansion following Kottwitz [K2]. By definition, a *distribution* on \mathfrak{g} is a linear functional on $\mathcal{C}_c^\infty(\mathfrak{g})$. Let $\mathcal{D}(\mathfrak{g})$ be the space of distributions on \mathfrak{g} , it is endowed with the topology as explained in [K2], §26.3. Let $\mathcal{D}(\mathfrak{g})^G$ be the subspace of G -invariant distributions on \mathfrak{g} . For any $X \in \mathfrak{g}$, it is known that the orbital integral

$$J_G(X, f) = |D(X)|^{\frac{1}{2}} \int_{G_X \backslash G} f(\text{Ad}(g^{-1})X) d\bar{g}$$

is absolutely convergent for any $f \in \mathcal{C}_c^\infty(\mathfrak{g})$, hence it defines a G -invariant distribution \mathcal{O}_X . Let $\mathcal{D}(\mathfrak{g})_{\text{orb}}$ be the subspace of $\mathcal{D}(\mathfrak{g})^G$ spanned by the orbital integral $\mathcal{O}_X, X \in \mathfrak{g}$. It is well known that the subspace $\mathcal{D}(\mathfrak{g})_{\text{orb}}$ has dense image in $\mathcal{D}(\mathfrak{g})^G$.

Theorem 3.1 (Shalika). *Let \mathbf{G} be a connected semisimple algebraic group over F . There exists functions $\Gamma_G(\cdot, u)$ on \mathfrak{g}^{rs} , $u \in [\mathcal{N}_G(F)]$, satisfying the property: For any $f \in \mathcal{C}_c^\infty(\mathfrak{g})$, there exists an open neighbourhood U_f of 0 in \mathfrak{g} such that*

$$\mathcal{O}_X(f) = \sum_{u \in [\mathcal{N}_G(F)]} \Gamma_G(X, u) \mathcal{O}_u(f), \quad \forall X \in U_f \cap \mathfrak{g}^{\text{rs}}.$$

The germs around 0 of the functions $\Gamma_G(\cdot, u)$ are uniquely determined, called Shalika germs.

The nilpotent orbits in \mathfrak{g} have a nice property: they are preserved under scaling. For $u \in \mathcal{N}_G(F)$, let \mathcal{O}_u be its $G(F)$ -orbit. It is known that if $t \in F^\times$, then u and tu are conjugate over \overline{F} ; and if $t \in (F^\times)^2$, they will be $G(F)$ -conjugate. Moreover,

$$d(tx) = |t|^{\dim(\mathcal{O}_u)/2} dx \quad (3.3)$$

for the canonical measure dx on \mathcal{O}_u . By the uniqueness of the Shalika germ expansion, we get the homogeneous property of the Shalika germs

$$\Gamma_G(tX, u) = |t|^{\frac{1}{2}[\dim(G_u) - \text{rk}(G)]} \Gamma_G(X, u).$$

This reduces the study of the orbital integrals to elements with small root valuations, which is much easier. The problem arises then to determine the exact domain of validity of the Shalika germ expansion.

For a compact subset ω of \mathfrak{g} , we denote

$${}^G\omega := \{\text{Ad}(g)X \mid g \in G, X \in \omega\}.$$

For a G -invariant subset $\Omega \subset \mathfrak{g}$, we denote by $J(\Omega)$ the invariant distributions on \mathfrak{g} with support in Ω . The following theorem determines the exact domain of validity of the Shalika germ expansion.

Theorem 3.2 (DeBacker [D2]). *For any $r \in \mathbf{R}$, we have equality of distributions:*

$$\text{Res}_{\mathcal{H}_r}(J(\mathfrak{g}_r)) = \text{Res}_{\mathcal{H}_r}(J(\mathcal{N}_G)).$$

In general, we can consider the germ expansion around any element $s \in \mathfrak{a}_{M_{\min}}$.

Theorem 3.3. *Let \mathbf{G} be a connected reductive algebraic group over F , let $s \in \mathfrak{a}_{M_{\min}}$. There exists functions $\Gamma_G(s + \cdot, s + u)$ on $\mathfrak{g}_s^{\text{der,rs}}$, $u \in [\mathcal{N}_{G_s}(F)]$, satisfying the property: For any $f \in \mathcal{C}_c^\infty(\mathfrak{g})$, there exists an open neighbourhood V_f of 0 in $\mathfrak{g}_s^{\text{der}}$ such that*

$$J_G(s + Y, f) = \sum_{u \in [\mathcal{N}_{G_s}(F)]} \Gamma_{G_s}(s + Y, s + u) J_{G_s}(s + u, f), \quad \forall Y \in V_f \cap \mathfrak{g}_s^{\text{der,rs}}.$$

Proof. Take a parabolic subgroup $Q \in \mathcal{P}(G_s)$, recall that for $Y \in \mathfrak{g}_s^{\text{der}}$, $f \in \mathcal{C}_c^\infty(\mathfrak{g})$, we have the parabolic descent

$$J_G(s + Y, f) = J_{G_s}^{G_s}(s + Y, f^Q), \quad (3.4)$$

where

$$f^Q(Y') = \int_K \int_{\mathfrak{n}_Q} f(\text{Ad}(k)(Y' + \nu)) d\nu dk, \quad \forall Y' \in \mathfrak{g}_s. \quad (3.5)$$

Applying the Shalika germ expansion for G_s , there exists an open neighbourhood V_f of 0 in $\mathfrak{g}_s^{\text{der}}$ such that

$$\begin{aligned} J_G(s+Y, f) &= J_{G_s}^{G_s}(s+Y, f^Q) = \sum_{u \in [\mathcal{N}_{G_s}(F)]} \Gamma_{G_s}(s+Y, s+u) J_{G_s}^{G_s}(s+u, f^Q) \\ &= \sum_{u \in [\mathcal{N}_{G_s}(F)]} \Gamma_{G_s}(s+Y, s+u) J_G(s+u, f). \end{aligned} \quad (3.6)$$

□

Theorem 3.4. *Let $s \in \mathfrak{a}_{M_{\min}}$, we have the equality of distributions*

$$\text{Res}_{\mathcal{H}_r} J(G(s + (\mathfrak{g}_s^{\text{der}})_r)) = \text{Res}_{\mathcal{H}_r} J(G(s + \mathcal{N}_{G_s})), \quad \forall r \in \mathbf{R}.$$

Proof. According to Harish-Chandra, every invariant distribution on \mathfrak{g} lies in the closure of the linear span of the subset $\{\mathcal{O}_X \mid X \in \mathfrak{g}^{\text{rs}}\}$. (We refer the reader to [K2], §27, for a beautiful proof of this result.) Consequently, $J(G(s + (\mathfrak{g}_s^{\text{der}})_r))$ lies in the closure of the linear span of the subset

$$\left\{ \mathcal{O}_X \mid X \in \mathfrak{g}^{\text{rs}} \cap G(s + (\mathfrak{g}_s^{\text{der}})_r) \right\}.$$

For $Y \in (\mathfrak{g}_s^{\text{der}})_r$, we have parabolic descent as before:

$$J_G(s+Y, f) = J_{G_s}^{G_s}(s+Y, f^Q).$$

Notice that if $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_r)$, then $f^Q \in \mathcal{C}_c(\mathfrak{g}_s/\mathfrak{g}_s \cap \mathfrak{k}_r)$ as \mathfrak{k}_r is stable under the adjoint action of K . Hence the expansion (3.6) holds for $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_r)$ if we apply theorem 3.2 to the group \mathbf{G}_s . The expansion, together with the above density result, implies the inclusion of distributions

$$\text{Res}_{\mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_r)} J(G(s + (\mathfrak{g}_s^{\text{der}})_r)) \subset \text{Res}_{\mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_r)} J(G(s + \mathcal{N}_{G_s})).$$

The inclusion in the other direction is clear, hence the equality of distributions as claimed.

□

3.2 The Arthur-Shalika expansion

For this section, we work with the group $\mathbf{G} = \text{GL}_{d+1}$. Let \mathbf{A} be the maximal torus of \mathbf{G} of the diagonal matrices, let \mathbf{B}_0 be the Borel subgroup of \mathbf{G} of the upper triangular matrices. Let $K = \mathbf{G}(\mathcal{O})$, let I be the standard Iwahori subgroup, it is the pre-image of $\mathbf{B}_0(k)$ under the reduction map $\mathbf{G}(\mathcal{O}) \rightarrow \mathbf{G}(k)$. In terms of Moy-Prasad filtration, we have $K = G_{0,0}$ and $I = G_{x_0,0}$ with $x_0 = (1, \frac{d}{d+1}, \dots, \frac{1}{d+1})$. For $n \in \mathbf{Z}$, let $\mathfrak{k}_n = \mathfrak{g}_{0,n}$ and $\mathfrak{i}_n = \mathfrak{g}_{x_0,n}$, and similarly for \mathfrak{k}_{n+} and \mathfrak{i}_{n+} .

To begin with, we can reinforce DeBacker's homogeneity result. Let \mathcal{L}_{n+}^G be the vector space of G -invariant linear functionals $L : \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_{n+}) \rightarrow \mathbf{C}$ such that $L(f) = 0$ for $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_{n+})$ with $\text{supp}(f) \cap (\mathfrak{k}_{n+} + \mathcal{N}_G) = \emptyset$.

Theorem 3.5. *For any $n \in \mathbf{Z}$, we have*

$$\mathcal{L}_{n+}^G = \text{Res}_{\mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_{n+})} (J(\mathcal{N}_G)).$$

Proof. After scaling by ϵ^n on \mathfrak{g} , the assertion is reduced to the $n = 0$ case, which is essentially the main result of DeBacker [D1]. Indeed, the key lemma 4 of *loc. cit.* holds more generally for the linear functionals in \mathcal{L}_{0+}^G . We don't need the much stronger assumption that D is a *distribution*, because the test functions involved in the proof lie in $\mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_{0+})$.

□

Theorem 3.6. *Let $M \in \mathcal{L}(A)$. For each element $u \in [\mathcal{N}_L(F)]$, $L \in \mathcal{L}(M)$, there exists a unique function $g_M^L(\cdot, u)$ on \mathfrak{g} , such that for any $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_{n+})$ and any element $\gamma \in G_{i_{n+}}$, $n \in \mathbf{Z}$, we have*

$$J_M(\gamma, f) = \sum_{L \in \mathcal{L}(M)} \sum_{u \in [\mathcal{N}_L(F)]} g_M^L(\gamma, u) J_L(u, f).$$

Proof. Up to scaling, it is enough to prove the assertion for $n = 0$. We will prove it in this case by induction on $\text{rk}(G)$. For $G = M$, the theorem is exactly the Shalika germ expansion, combined with the theorem of DeBacker.

Now suppose that the theorem holds for the proper Levi subgroups of G containing M , then

$$\begin{aligned} J_M(\gamma, f^y) - J_M(\gamma, f) &= \sum_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} J_M^{MQ}(\gamma, f_{Q,y}) \\ &= \sum_{\substack{Q \in \mathcal{F}(M) \\ Q \neq G}} \sum_{L \in \mathcal{L}^{MQ}(M)} \sum_{u \in [\mathcal{N}_L(F)]} g_M^L(\gamma, u) \cdot J_L^{MQ}(u, f_{Q,y}) \\ &= \sum_{\substack{L \in \mathcal{L}(M) \\ L \neq G}} \sum_{u \in [\mathcal{N}_L(F)]} g_M^L(\gamma, u) \sum_{\substack{Q \in \mathcal{F}(L) \\ Q \neq G}} J_L^{MQ}(u, f_{Q,y}) \\ &= \sum_{\substack{L \in \mathcal{L}(M) \\ L \neq G}} \sum_{u \in [\mathcal{N}_L(F)]} g_M^L(\gamma, u) [J_L(u, f^y) - J_L(u, f)], \end{aligned}$$

where the first and the last equation is an application of the equation (3.1) to the element γ and u respectively, and the second equation is just the induction hypothesis applied to the function $f_{Q,y}$, which belongs to $\mathcal{C}_c(\mathfrak{m}_Q/\mathfrak{m}_Q \cap \mathfrak{k}_{0+})$ as we have observed in (3.2). The above equation implies that the linear functional on $\mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_{0+})$ sending $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_{0+})$ to

$$J'_M(\gamma, f) := J_M(\gamma, f) - \sum_{\substack{L \in \mathcal{L}(M) \\ L \neq G}} \sum_{u \in [\mathcal{N}_L(F)]} g_M^L(\gamma, u) J_L(u, f) \quad (3.7)$$

is G -conjugation invariant. Moreover, as $G_{i_{0+}} \subset \mathfrak{k}_{0+} + \mathcal{N}_G$, we have $J_M(\gamma, f) = 0$ for $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_{0+})$ with $\text{supp}(f) \cap (\mathfrak{k}_{0+} + \mathcal{N}_G) = \emptyset$, and similarly for $J_M(u, f)$, $u \in [\mathcal{N}_G(F)]$, by taking limits. Hence $J'_M(\gamma, \cdot) \in \mathcal{L}_{0+}^G$, and so it is a linear combination of the invariant distributions \mathcal{O}_u , $u \in [\mathcal{N}_G(F)]$ by theorem 3.5, this defines the functions $g_M^G(\gamma, u)$. Their uniqueness follows from the linear independence of the invariant distributions \mathcal{O}_u 's. \square

The functions g_M^G are called the *Arthur-Shalika germs*. In [A3], Arthur has shown that the *germs* of these functions satisfies a homogeneous property and a semisimple descent formula. Now that we have removed all the ambiguities on the Arthur-Shalika germ expansion in *loc. cit.* and determine its domain of validity, we can reinforce these results to the functions themselves. For $u \in \mathcal{N}_M(F)$, Arthur has defined a (G, M) -family

$$c_P(\lambda; \exp(u), t), \quad P \in \mathcal{P}(M).$$

By abuse of notation, we will denote it by $c_P(\lambda; u, t)$. For $v \in [\mathcal{N}_L(F)]$, $L \in \mathcal{L}(M)$, let

$$[u_M^L : v] = \begin{cases} 1, & \text{if } v \in u_M^L, \\ 0, & \text{otherwise,} \end{cases}$$

and $d^M(u) = \frac{1}{2}(\dim(M_u) - \text{rk}(M))$. Obviously $d^M(u) = d^L(v)$ if $v \in u_M^L$.

Theorem 3.7. For $u \in [\mathcal{N}_G(F)]$ and $\gamma \in \mathfrak{m} \cap \mathfrak{g}^{\text{rs}}$, we have

$$g_M^G(t\gamma, tu) = |t|^{d^G(u)} \sum_{L \in \mathcal{L}(M)} \sum_{v \in [\mathcal{N}_L(F)]} g_M^L(\gamma, v) c_L(v, t) [v_L^G : u], \quad \forall t \in F^\times.$$

Proof. The proof proceeds along the same lines as in [A3], §10, taking into account the fact that the scaling on \mathfrak{g} sends

$$G_i \xrightarrow{\cdot t} G_{i+\text{val}(t)_+}, \quad C_c(\mathfrak{g}/\mathfrak{k}_{n+}) \xrightarrow{\cdot t} C_c(\mathfrak{g}/\mathfrak{k}_{n+\text{val}(t)_+}), \quad \forall n \in \mathbf{Z},$$

which preserves the validity of theorem 3.6. □

3.3 The parabolic descent

Let $\mathbf{G} = \text{GL}_{d+1}$ as before. Let $\chi : \mathfrak{g} \rightarrow \mathfrak{g} // G$ be the characteristic morphism. The above results describe the local behavior of $J_M(\gamma, f)$ as $\chi(\gamma) \rightarrow 0$. We can describe its local behavior as $\chi(\gamma)$ approaches a wall. For $s \in \mathfrak{a}$, let $M \in \mathcal{L}(A)$ be a Levi subgroup satisfying $A_{M_s} = A_M$. Let $\mu \in \mathfrak{m} \cap \mathfrak{g}_s^{\text{der}}$, then $G_{s+\mu} \subset G_s$. Following Arthur [A3], we can express $J_M(s + \mu, f)$ in terms of weighted orbital integrals of μ for the group G_s . To begin with, we write

$$\begin{aligned} J_M(s + \mu, f) &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \sum_{L \in \mathcal{L}(M)} r_M^L(s + \mu, a) J_L(a + s + \mu, f) \\ &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} |D(a + s + \mu)|^{\frac{1}{2}} \int_{G_s \backslash G} dg_1 \int_{G_{a+s+\mu} \backslash G_s} dg_2 \cdot f(\text{Ad}(g_2 g_1)^{-1}(a + s + \mu)) \\ &\quad \cdot \left[\sum_{L \in \mathcal{L}(M)} r_M^L(s + \mu, a) v_L(g_2 g_1) \right]. \end{aligned} \tag{3.8}$$

Lemma 3.8 (Arthur [A3]). For $L \in \mathcal{L}(M)$, we have

$$r_M^L(s + \mu, a) = \begin{cases} r_{M_s}^L(\mu, a), & \text{if } A_L = A_{L_s}, \\ 0, & \text{otherwise.} \end{cases}$$

For the weight factor $v_L(g_2 g_1)$, it can be written as the volume of the product of two (G, L) -families. For $Q = LN_Q \in \mathcal{P}(L)$, we can write $g_2 = lnk_Q(g_2)$ with $l \in L(F)$, $n \in N_Q(F)$, $k_Q(g_2) \in K$, then

$$\begin{aligned} v_Q(\lambda; g_2 g_1) &= e^{\langle \lambda, H_Q(g_2 g_1) \rangle} = e^{\langle \lambda, H_Q(g_2) \rangle} e^{\langle \lambda, H_Q(k_Q(g_2) g_1) \rangle} \\ &= v_Q(\lambda; g_2) u_Q(\lambda; g_2, g_1), \end{aligned}$$

with $u_Q(\lambda; g_2, g_1) = e^{\langle \lambda, H_Q(k_Q(g_2) g_1) \rangle}$. Notice that $(u_Q(\lambda; g_2, g_1))_{Q \in \mathcal{P}(L)}$ forms a (G, L) -family, hence by the product formula (1.2), we have

$$v_L(g_2 g_1) = \sum_{Q' \in \mathcal{F}(L)} v_{L'}^{Q'}(g_2) u_{Q'}'(g_2, g_1). \tag{3.9}$$

By construction, it is clear that for any $k \in K$, we have

$$u_Q(\lambda; g_2, k g_1) = u_Q(\lambda; g_2 k, g_1),$$

hence

$$u'_{Q'}(g_2, kg_1) = u'_{Q'}(g_2k, g_1), \quad \forall Q' \in \mathcal{F}(L). \quad (3.10)$$

We can further simplify the formula with the condition $g_2 \in G_s$. By the above formula, it is enough to consider the Levi subgroups $L \in \mathcal{L}(M)$ such that $A_{L_s} = A_L$. We identify all the structures on \mathfrak{a}_{L_s} and \mathfrak{a}_L .

Lemma 3.9 (Arthur [A3]). *We have*

$$v_L(g_2g_1) = \sum_{R \in \mathcal{F}^{G_s}(L_s)} v_{L_s}^R(g_2) u'_R(g_2, g_1),$$

where

$$u'_R(g_2, g_1) = \sum_{\{Q' \in \mathcal{F}(M) | Q'_s = R, \mathfrak{a}_{Q'} = \mathfrak{a}_R\}} u'_{Q'}(g_2, g_1).$$

The mapping $L \rightarrow L_s$ sets up a bijection from the set of elements in $\mathcal{L}(M)$ satisfying $A_L = A_{L_s}$ with the set $\mathcal{L}^{G_s}(M_s)$. Hence combining the above two lemmas, we get

$$\sum_{L \in \mathcal{L}(M)} r_M^L(s + \mu, a) v_L(g_2g_1) = \sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{J \in \mathcal{L}^{M_R}(M_s)} r_{M_s}^J(\mu, a) v_J^R(g_2) u'_R(g_2, g_1).$$

Plug it into the equation (3.8), we get

$$\begin{aligned} J_M(s + \mu, f) &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}}(F) \\ a \rightarrow 0}} |D(a + s + \mu)|^{\frac{1}{2}} \int_{G_s \backslash G} dg_1 \int_{G_{a+s+\mu} \backslash G_s} dg_2 \cdot f(\text{Ad}(g_2g_1)^{-1}(a + s + \mu)) \\ &\quad \cdot \left[\sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{J \in \mathcal{L}^{M_R}(M_s)} r_{M_s}^J(\mu, a) v_J^R(g_2) u'_R(g_2, g_1) \right] \\ &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}}(F) \\ a \rightarrow 0}} |D(a + s + \mu)|^{\frac{1}{2}} \int_{G_s \backslash G} dg_1 \sum_{R \in \mathcal{F}^{G_s}(M_s)} \int_{M_{s+\mu} \backslash M_R} dm \int_{N_R} dn \int_{K^{G_s}} dk \\ &\quad \cdot f(\text{Ad}(g_1)^{-1}[s + \text{Ad}(mnk)^{-1}(a + \mu)]) \left[\sum_{J \in \mathcal{L}^{M_R}(M_s)} r_{M_s}^J(\mu, a) v_J^R(m) u'_R(k, g_1) \right] \\ &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}}(F) \\ a \rightarrow 0}} |D(a + s + \mu)|^{\frac{1}{2}} \int_{G_s \backslash G} dg_1 \sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{J \in \mathcal{L}^{M_R}(M_s)} r_{M_s}^J(\mu, a) \int_{M_{s+\mu} \backslash M_R} \\ &\quad \cdot \left[\int_{N_R} \int_{K^{G_s}} f(\text{Ad}(g_1)^{-1}[s + \text{Ad}(mnk)^{-1}(a + \mu)]) u'_R(k, g_1) dk dn \right] v_J^R(m) dm. \quad (3.11) \end{aligned}$$

For $Y \in \mathfrak{m}_R$, let

$$\tilde{f}_{R, g_1}(Y) = \int_{N_R} \int_{K^{G_s}} f(\text{Ad}(g_1)^{-1}[s + \text{Ad}(nk)^{-1}Y]) u'_R(k, g_1) dk dn. \quad (3.12)$$

The above equation can be simplified to

$$\begin{aligned}
J_M(s + \mu, f) &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \left| \frac{D(a + s + \mu)}{D^{G_s}(a + \mu)} \right|^{\frac{1}{2}} \int_{G_s \backslash G} dg_1 \sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{J \in \mathcal{L}^{MR}(M_s)} r_{M_s}^J(\mu, a) J_J^{MR}(a + \mu, \tilde{f}_{R, g_1}) \\
&= \left| \frac{D(s + \mu)}{D^{G_s}(\mu)} \right|^{\frac{1}{2}} \int_{G_s \backslash G} dg_1 \sum_{R \in \mathcal{F}^{G_s}(M_s)} \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \sum_{J \in \mathcal{L}^{MR}(M_s)} r_{M_s}^J(\mu, a) J_J^{MR}(a + \mu, \tilde{f}_{R, g_1}) \\
&= \left| \frac{D(s + \mu)}{D^{G_s}(\mu)} \right|^{\frac{1}{2}} \int_{G_s \backslash G} dg_1 \sum_{R \in \mathcal{F}^{G_s}(M_s)} J_{M_s}^{MR}(\mu, \tilde{f}_{R, g_1}), \tag{3.13}
\end{aligned}$$

where in the second line we are able to interchange the order of taking limit and integration because the integration is essentially finite as \tilde{f}_{R, g_1} vanishes identically for g_1 outside of a compact region in $G_s \backslash G$. We summarize the above calculations as

Theorem 3.10. For $s \in \mathfrak{a}$, let $M \in \mathcal{L}(A)$ be such that $A_{M_s} = A_M$, let $\mu \in \mathfrak{m} \cap \mathfrak{g}_s^{\text{der}}$, then

$$J_M(s + \mu, f) = \left| \frac{D(s + \mu)}{D^{G_s}(\mu)} \right|^{\frac{1}{2}} \int_{G_s \backslash G} \sum_{R \in \mathcal{F}^{G_s}(M_s)} J_{M_s}^R(\mu, \tilde{f}_{R, g}) dg.$$

Let $U_s^+ \subset \mathfrak{g}_s^{\text{der}}$ be the invariant neighborhood of 0 consisting of elements u such that all the eigenvalues of $\text{ad}(u)\text{ad}(s)^{-1}$ on $\mathfrak{g}/\mathfrak{g}_s$ have valuations ≥ 0 .

Theorem 3.11. For $s \in \mathfrak{a}$, $M \in \mathcal{L}(A)$ be such that $A_{M_s} = A_M$, let r be the valuation of the eigenvalues of $\text{ad}(s)$ on $\mathfrak{g}/\mathfrak{g}_s$. Let $\eta \in \mathfrak{m} \cap U_s^+$ be such that $s + \eta \in \mathfrak{g}^{\text{rs}}$, then for any $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}_r)$, we have expansion

$$J_M(s + \eta, f) = \sum_{L_s \in \mathcal{L}^{G_s}(M_s)} \sum_{u \in [\mathcal{N}_{L_s}(F)]} g_{M_s}^{L_s}(\eta, u) J_{L_s}(s + u, f).$$

Proof. By the homogeneity of the Arthur-Shalika germ and by induction on $\dim(\mathfrak{a}_{G_s})$, it is enough to treat the case that $\dim(\mathfrak{a}_{G_s}) = \dim(\mathfrak{a}_G) + 1$ and that the eigenvalues of the action of $\text{ad}(s)$ on $\mathfrak{g}/\mathfrak{g}_s$ has valuation 0. We fix a parabolic subgroup $Q \in \mathcal{P}(G_s)$.

According to the construction in the proof of theorem 3.6, the functions g_M^G are defined inductively as coefficients of the expansion

$$J'_M(s + \eta, f) = \sum_{u \in [\mathcal{N}_G(F)]} g_M^G(s + \eta, u) J_G(u, f), \quad \text{for } f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k}).$$

To determine them, it is then enough to examine the local behavior of $J_M(s + \eta, f)$ for a set of test functions $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k})$ which forms a dual basis of $J_G(u, \cdot)$, $u \in [\mathcal{N}_G(F)]$. Since we are working with $\mathbf{G} = \text{GL}_{d+1}$, the functions f can be taken to be the characteristic functions $[\mathbf{G}^{(k)}(\epsilon^{-1}u) + \mathfrak{k}]$, $u \in \mathcal{N}_G(k)$. For such f and $\eta \in \mathfrak{m} \cap U_s^+$ such that $s + \eta \in \mathfrak{g}^{\text{rs}}$, by theorem 3.10, we have

$$\begin{aligned}
J_M(s + \eta, f) &= \left| \frac{D(s + \eta)}{D^{G_s}(\eta)} \right|^{\frac{1}{2}} \int_{G_s \backslash G} \sum_{R \in \mathcal{F}^{G_s}(M_s)} J_{M_s}^R(\eta, \tilde{f}_{R, g}) dg \\
&= \delta_Q(s) \int_{N_Q} \sum_{R \in \mathcal{F}^{G_s}(M_s)} J_{M_s}^R(\eta, \tilde{f}_{R, g}) dg, \tag{3.14}
\end{aligned}$$

where in the second line we have used the Iwasawa decomposition and the fact that f is invariant under K -conjugation, and we have set $\delta_Q(s) = |\det(\text{ad}(s); \mathfrak{n}_Q)|$.

Look at the integral $\int_{N_Q} J_{M_s}^R(\eta, \tilde{f}_{R,g}) dg$. With the expression (3.12) and the fact that f is K -conjugation invariant, we can manipulate

$$\begin{aligned}
& \int_{N_Q} dg \int_{M_{s+\eta} \backslash M_R} \left[\int_{N_R} \int_{K^{G_s}} f(\text{Ad}(g)^{-1}[s + \text{Ad}(mnk)^{-1}\eta]) u'_R(k, g) dk dn \right] v_{M_s}^R(m) dm \\
&= \int_{N_Q} dg \int_{M_{s+\eta} \backslash M_R} \left[\int_{N_R} \int_{K^{G_s}} f(\text{Ad}(kgk^{-1})^{-1}[s + \text{Ad}(mn)^{-1}\eta]) u'_R(k, g) dk dn \right] v_{M_s}^R(m) dm \\
&= \int_{N_Q} dg' \int_{M_{s+\eta} \backslash M_R} \left[\int_{N_R} \int_{K^{G_s}} f(\text{Ad}(g')^{-1}[s + \text{Ad}(mn)^{-1}\eta]) u'_R(k, k^{-1}g'k) dk dn \right] v_{M_s}^R(m) dm \\
&= \int_{N_Q} dg' \int_{M_{s+\eta} \backslash M_R} \left[\int_{N_R} f(\text{Ad}(g')^{-1}[s + \text{Ad}(mn)^{-1}\eta]) u'_R(1, g') dn \right] v_{M_s}^R(m) dm, \quad (3.15)
\end{aligned}$$

where the last equality is due to equation (3.10) and the right- K -invariance of Arthur's weight factor. The factor $u'_R(1, g')$ can be further simplified. By construction,

$$u'_R(1, g') = \sum_{\{Q' \in \mathcal{F}(M) \mid Q'_s = R, \mathfrak{a}_{Q'} = \mathfrak{a}_R\}} u'_{Q'}(1, g') = \sum_{\{Q' \in \mathcal{F}(M) \mid Q'_s = R, \mathfrak{a}_{Q'} = \mathfrak{a}_R\}} v'_{Q'}(g').$$

In particular, it is left- R -invariant and right- K -invariant in the variable g' .

For the integral to be non-zero (and hence positive), we need to solve the equation

$$\text{Ad}(g')^{-1}[s + \text{Ad}(mn)^{-1}\eta] \in \mathbf{G}^{(k)}(\epsilon^{-1}u) + \mathfrak{k}, \quad (3.16)$$

with $g' \in N_Q, m \in M_R, n \in N_R$. Let $g' = \exp(Z)$ with $Z \in \mathfrak{n}_Q$, then the equation can be rewritten

$$(s + \text{Ad}(mn)^{-1}\eta) + [s + \text{Ad}(mn)^{-1}\eta, Z] \in \mathbf{G}^{(k)}(\epsilon^{-1}u) + \mathfrak{k}.$$

Notice that the left hand side is an element of \mathfrak{q} . To solve the equation, we need to understand the intersection $\mathbf{G}^{(k)}u \cap \mathfrak{q}(k)$. It is clear that it is stable under $\mathbf{G}_s(k)$ -conjugation, hence there exists finitely many orbits $\mathbf{G}_s^{(k)}u_i, i \in I$, in $\mathcal{N}_{G_s}(k)$ such that

$$\mathbf{G}^{(k)}u \cap \mathfrak{q}(k) = \bigsqcup_{i \in I} \mathbf{G}^{(k)}u \cap (\mathbf{G}_s^{(k)}u_i + \mathfrak{n}_Q).$$

Hence the set of solutions Ω to the equation (3.16) is the disjoint union of the set of solutions Ω_i to the equations

$$\text{Ad}(mn)^{-1}\eta \in \mathbf{G}_s^{(k)}(\epsilon^{-1}u_i) + \mathfrak{k}^{G_s}, \quad (3.17)$$

$$[s + \text{Ad}(mn)^{-1}\eta, Z] \in \epsilon^{-1}(\mathfrak{n}_Q \cap \mathfrak{k}), \quad (3.18)$$

such that their sum satisfies

$$\text{Ad}(mn)^{-1}\eta + [s + \text{Ad}(mn)^{-1}\eta, Z] \in \mathbf{G}^{(k)}(\epsilon^{-1}u) + \mathfrak{k}. \quad (3.19)$$

The equation (3.17) can be further refined. Notice that its left hand side is an element of \mathfrak{t} . As before, we can consider the $R(k)$ -conjugation and make the decomposition

$$\mathbf{G}_s^{(k)}u_i \cap \mathfrak{t}(k) = \bigsqcup_{j \in J_i} R^{(k)}u_{ij}, \quad \text{with } u_{ij} \in \mathfrak{t}(k).$$

Let Ω_{ij} be the set of solutions to the equation

$$\text{Ad}(mn)^{-1}\eta \in {}^{R(k)}(\epsilon^{-1}u_{ij}) + (\mathfrak{r} \cap \mathfrak{k}), \quad (3.20)$$

and the equations (3.18) and (3.19), then $\Omega_i = \bigsqcup_{j \in J_i} \Omega_{ij}$.

Notice that if we conjugate the equation (3.20) on both sides by an element $h \in R(k)$, we get a solution to the equations (3.18) and (3.19) if we substitute Z by $\text{Ad}(h)Z$. Hence we can normalize the equation (3.20) as $\text{Ad}(mn)^{-1}\eta = \epsilon^{-1}u_{ij} + \eta'$ for some $\eta' \in \mathfrak{r} \cap \mathfrak{k}$, then the equation (3.18) simplifies to

$$[s + \epsilon^{-1}u_{ij} + \eta', Z] \in \epsilon^{-1}(\mathfrak{n}_Q \cap \mathfrak{k}). \quad (3.21)$$

By assumption, $\eta \in U_s^+$, hence the eigenvalues of $\text{ad}(\eta')$ on $\mathfrak{g}/\mathfrak{g}_s$ is ≥ 0 . This implies that there exists $k' \in R \cap K$ such that $Z' = \text{Ad}(k')Z$ satisfies the equation

$$[s + \epsilon^{-1}u_{ij}, Z'] \in \epsilon^{-1}(\mathfrak{n}_Q \cap \mathfrak{k}), \quad (3.22)$$

and that this sets up a bijection between the set of solution of equation (3.21) and (3.22).

Looking back at the expression (3.15) for $\int_{N_Q} J_{M_s}^R(\eta, \tilde{f}_{R,g})$. The above discussion implies that, for each set of solution Ω_{ij} , the element $g' = \exp(Z)$ depends only on $s + \epsilon^{-1}u_{ij}$ up to conjugation by an element of $R \cap K$. Observe that $u'_R(1, g')$ is invariant under this conjugation, as it is left- R -invariant and right- K -invariant in the variable g' . Hence $u'_R(1, g')$ takes the same value on Ω_{ij} , depending only on s and u_{ij} , we denote it by $\phi_R(s, u_{ij})$.

Let $\mathbb{1}_{\Omega_{ij}}$ be the characteristic function of Ω_{ij} . Let Ω_{ij}^R be the set of solutions to equation (3.20) and $\mathbb{1}_{\Omega_{ij}^R}$ its characteristic function. The expression (3.14) for $J_M(s + \eta, f)$ can be simplified as

$$\begin{aligned} J_M(s + \eta, f) &= \delta_Q(s) \int_{N_Q} \sum_{R \in \mathcal{F}^{G_s}(M_s)} J_{M_s}^R(\eta, \tilde{f}_{R,g}) dg \\ &= \delta_Q(s) \sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{i \in I, j \in J_i} \int_{N_Q} \int_{M_s + \eta \backslash M_R} \left[\int_{N_R} \mathbb{1}_{\Omega_{ij}} \cdot f(\text{Ad}(g)^{-1}[s + \text{Ad}(mn)^{-1}\eta]) \phi_R(s, u_{ij}) dn \right] v_{M_s}^R(m) dmdg \\ &= \delta_Q(s) \sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{i \in I, j \in J_i} \phi_R(s, u_{ij}) \int_{M_s + \eta \backslash M_R} \int_{N_R} \left[\int_{N_Q} \mathbb{1}_{\Omega_{ij}} \cdot f(\text{Ad}(g)^{-1}[s + \text{Ad}(mn)^{-1}\eta]) dg \right] v_{M_s}^R(m) dndm \\ &= \delta_Q(s) \sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{i \in I, j \in J_i} \phi_R(s, u_{ij}) \int_{M_s + \eta \backslash M_R} \int_{N_R} \mathbb{1}_{\Omega_{ij}^R} \cdot f^Q(s + \text{Ad}(mn)^{-1}\eta) v_{M_s}^R(m) dndm \\ &= \delta_Q(s) \sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{i \in I, j \in J_i} \phi_R(s, u_{ij}) J_{M_s}^R(s + \eta, \mathbb{1}_{\Omega_{ij}^R} \cdot f^Q), \end{aligned} \quad (3.23)$$

where in the integrals $\mathbb{1}_{\Omega_{ij}}$ actually means $\mathbb{1}_{\Omega_{ij}}(g; m, n, k)$ and similarly for $\mathbb{1}_{\Omega_{ij}^R}$. For general elements $\mu \in \mathfrak{m} \cap U_s^+$, similar calculations apply to the weighted orbital integrals $J_J^{MR}(a + \mu, \tilde{f}_{R,g_1})$ in the expression (3.13) for $J_M(s + \mu, f)$. Taking limits, we get

$$J_M(s + \mu, f) = \delta_Q(s) \sum_{R \in \mathcal{F}^{G_s}(M_s)} \sum_{i \in I, j \in J_i} \phi_R(s, u_{ij}) J_{M_s}^{MR}(s + \mu, \mathbb{1}_{\Omega_{ij}^R} \cdot f^Q). \quad (3.24)$$

Now that $f^Q \in \mathcal{C}_c(\mathfrak{g}_s/\mathfrak{k}^{G_s})$ and that $s, \eta \in (\mathfrak{g}_s)_0$, we can apply the Arthur-Shalika germ expansion

to $J_{M_s}^{MR}(s + \eta, \mathbb{1}_{\Omega_{ij}^R} \cdot f^Q)$ and obtain the searched-for expansion

$$\begin{aligned}
J_M(s + \eta, f) &= \delta_Q(s) \sum_{R \in \mathcal{F}^{Gs}(M_s)} \sum_{i \in I, j \in J_i} \phi_R(s, u_{ij}) J_{M_s}^{MR}(s + \eta, \mathbb{1}_{\Omega_{ij}^R} \cdot f^Q) \\
&= \delta_Q(s) \sum_{R \in \mathcal{F}^{Gs}(M_s)} \sum_{i \in I, j \in J_i} \phi_R(s, u_{ij}) \sum_{L_s \in \mathcal{L}^{MR}(M_s)} \sum_{u \in [\mathcal{N}_{L_s}(F)]} g_{M_s}^{L_s}(s + \eta, s + u) J_{L_s}^{MR}(s + u, \mathbb{1}_{\Omega_{ij}^R} \cdot f^Q) \\
&= \delta_Q(s) \sum_{L_s \in \mathcal{L}^{Gs}(M_s)} \sum_{u \in [\mathcal{N}_{L_s}(F)]} g_{M_s}^{L_s}(\eta, u) \sum_{R \in \mathcal{F}^{Gs}(L_s)} \sum_{i \in I, j \in J_i} \phi_R(s, u_{ij}) J_{L_s}^{MR}(s + u, \mathbb{1}_{\Omega_{ij}^R} \cdot f^Q) \\
&= \sum_{L_s \in \mathcal{L}^{Gs}(M_s)} \sum_{u \in [\mathcal{N}_{L_s}(F)]} g_{M_s}^{L_s}(\eta, u) J_{L_s}(s + u, f),
\end{aligned}$$

where the first and last line are due to equation (3.23) and (3.24) respectively. \square

4 Counting points on the fundamental domain

For the affine Springer fibers, we need to take $F = \mathbb{F}_q((\epsilon))$. For the group \mathbf{G} , we assume that it is a connected reductive algebraic group such that \mathbf{G}^{der} is simply connected. In [C2], we have explained a transition between counting points on the fundamental domain of the affine Springer fibers and the calculation of Arthur's weighted orbital integrals. The strategy is to count the number of points on the truncated affine Springer fibers in two ways: by the Arthur-Kottwitz reduction and by the Harder-Narasimhan reduction. Here we explain another way to count the number of points on the fundamental domains.

Let T be a maximal torus of G , let $\gamma \in \mathfrak{t}$ be a regular integral element. Recall that the affine Springer fiber is the sub ind- k -scheme of the affine grassmannian $\mathcal{X} = \mathbf{G}((\epsilon))/\mathbf{G}[\epsilon]$ defined by the equation

$$\mathcal{X}_\gamma = \{g \in \mathcal{X} \mid \text{Ad}(g)^{-1}\gamma \in \mathfrak{g}[\epsilon]\}.$$

The group T acts on it by left translation. Let $S \subset T$ be the maximal F -split subtorus of T , let $\Lambda = \{\epsilon^\nu \mid \nu \in X_*(S)\}$. Then Λ acts freely on \mathcal{X}_γ and the quotient $\Lambda \backslash \mathcal{X}_\gamma$ is a projective k -scheme [KL]. Moreover, there is an open dense subscheme $\mathcal{X}_\gamma^{\text{reg}}$ consisting of the points $[g]$ such that $\text{Ad}(g)^{-1}\gamma \bmod \epsilon$ is regular in \mathfrak{g} , and the group T acts transitively on $\mathcal{X}_\gamma^{\text{reg}}$ [Bez]. Let $M_0 = Z_{\mathbf{G}_F}(S)$ be the centralizer of S in \mathbf{G}_F , then M_0 is a Levi subgroup of G and T is elliptic in M_0 . Fix a point $x_0 \in \mathcal{X}_\gamma^{\text{reg}}$, the fundamental domain of \mathcal{X}_γ is defined as:

$$F_\gamma = \{x \in \mathcal{X}_\gamma \mid \text{Ec}_{M_0}(x) \subset \text{Ec}_{M_0}(x_0), x \text{ and } x_0 \text{ lie on the same connected component}\}.$$

Then F_γ is a projective k -scheme, and for different choices of x_0 , the fundamental domains are isomorphic. As explained in [C1], the study of the geometry of \mathcal{X}_γ can be reduced to that of F_γ by the Arthur-Kottwitz reduction.

From now on we assume that the splitting field of T is totally ramified over F . Let $T^1 = T \cap \ker(H_{M_0})$. By [C2], lemma 2.6, T^1 is of finite volume and we have an exact sequence

$$1 \rightarrow T^1 \rightarrow T \xrightarrow{H_{M_0}} X_*(M_0) \rightarrow 1. \quad (4.1)$$

Let Λ_{M_0} be the quotient of $X_*(A)$ by the coroot lattice of M_0 , recall that the connected components of $\mathcal{X}_\gamma^{M_0}$ are parametrized by Λ_{M_0} . Now that M_0^{der} is also simply connected, we have $X_*(M_0) = \Lambda_{M_0}$ by [CL2], lemma 11.6.1. For any point $x \in \mathcal{X}_\gamma$, we define a weight factor

$$v_\gamma(x) = |\{\lambda \in \Lambda_{M_0} \mid \lambda + \text{Ec}_{M_0}(x) \subset \text{Ec}_{M_0}(x_0)\}|.$$

Lemma 4.1. *We have*

$$|F_\gamma(\mathbb{F}_q)| = \text{vol}(T^1) \int_{T \backslash G} \mathbb{1}_{\mathfrak{t}}(\text{Ad}(g^{-1})\gamma)v_\gamma(g)d\bar{g}.$$

Proof. Take a parabolic subgroup $P = M_0N \in \mathcal{P}(M_0)$, consider the parabolic reduction

$$f_P : \mathcal{X}_\gamma \rightarrow \mathcal{X}_\gamma^{M_0}.$$

It is known that its restriction to the pre-image of each connected component $\mathcal{X}_\gamma^{M_0, \nu}$, $\nu \in \Lambda_{M_0}$, is an iterated fibration in affine spaces, and that the translation with T identifies them [KL]. In particular, every point in F_γ can be moved to $f_P^{-1}(\mathcal{X}_\gamma^{M_0, 0})$ by translation with T . Combined with the fact that $H_P = H_{M_0} \circ f_P$, and that T fits into the exact sequence (4.1), we get

$$\begin{aligned} |F_\gamma(\mathbb{F}_q)| &= \sum_{x \in f_P^{-1}(\mathcal{X}_\gamma^{M_0, 0})(\mathbb{F}_q)} |\{\lambda \in \Lambda_{M_0} \mid \lambda + \text{Ec}(x) \subset \text{Ec}(x_0)\}| \\ &= \text{vol}(T^1) \int_{T \backslash G} \mathbb{1}_{\mathfrak{t}}(\text{Ad}(g^{-1})\gamma)v_\gamma(g)d\bar{g}. \end{aligned}$$

□

For $x \in \mathcal{X}_\gamma$, let $\lambda(x, x_0)$ be the positive (G, M_0) -orthogonal set

$$\lambda_P(x, x_0) = H_P(x_0) - H_P(x), \quad \forall P \in \mathcal{P}(M_0).$$

In other words, it is the difference of the (G, M_0) -orthogonal sets $\text{Ec}_{M_0}(x_0)$ and $\text{Ec}_{M_0}(x)$. Here the positivity of $\lambda(x, x_0)$ follows from the characterization of the regular points on \mathcal{X}_γ by Goresky, Kottwitz and MacPherson [GKM2].

Lemma 4.2. *For any point $x \in \mathcal{X}_\gamma$, we have*

$$v_\gamma(x) = |\mathbf{E}(\lambda(x, x_0)) \cap \Lambda_{M_0}|.$$

Proof. By definition, the inclusion of polytopes $\lambda + \text{Ec}(x) \subset \text{Ec}(x_0)$ is equivalent to the condition

$$\lambda + H_P(x) \prec_P H_P(x_0), \quad \forall P \in \mathcal{P}(M_0),$$

where $a \prec_{P_0} b$ means that $b - a$ is a positive linear combination of elements in $\Delta_{P_0}^\vee$. Obviously, this is equivalent to the condition that $\lambda \in \mathbf{E}(\lambda(x, x_0))$.

□

To count the number of lattice points in the polytope generated by a positive (G, M) -orthogonal set, the question has been addressed in [C2], §2.2.3, following the work of Arthur [A2] and Chaudouard-Laumon [CL2]. Let $h = (h_P)_{P \in \mathcal{P}(M)}$ be an *integral* positive (G, M) -orthogonal set. By the word *integral*, we mean that $h_P \in \Lambda_M$ for all $P \in \mathcal{P}(M)$. Let $w_M(h)$ be the number of lattice points in Λ_M contained in the polytope $\mathbf{E}(h)$, let $v_M(h)$ be the volume of the polytope $\mathbf{E}(h)$.

Lemma 4.3. *Let $\bar{c}_P(\lambda) = \prod_{\alpha \in \Delta_P} (1 - e^{-\langle \lambda, \alpha^\vee \rangle})$, then*

$$w_M(h) = \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} \bar{c}_P(\lambda)^{-1} e^{\langle \lambda, h_P \rangle}.$$

Proof. The proof follows the same line as the deduction of the equations (2.8) in [C2], we indicate only the modifications. Let φ_P^λ be the characteristic function of the cone

$$\{a \in \mathfrak{a}_M \mid \varpi_\alpha(a) > 0, \forall \alpha \in \Delta_P^\lambda; \varpi_\alpha(a) \leq 0, \forall \alpha \in \Delta_P \setminus \Delta_P^\lambda\},$$

where λ is a generic point in \mathfrak{a}_M^* , and

$$\Delta_P^\lambda = \{\alpha \in \Delta_P \mid \langle \lambda, \alpha \rangle < 0\}.$$

Then the characteristic function of the *closed* convex polytope $\mathbf{E}(h)$ is equal to the function

$$a \in \mathfrak{a}_M \mapsto \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} \varphi_P^\lambda(-h_P + a),$$

and the function $w_M(h)$ can be expressed as

$$w_M(h) = \sum_{\chi \in X_*(M)} \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} \varphi_P^\lambda(-h_P + \chi).$$

Again we introduce an extra exponential factor to treat the infinite sum in $X_*(M)$, let

$$\overline{S}_P(\lambda) = \sum_{\chi \in X_*(M)} \varphi_P^\lambda(-h_P + \chi) e^{\langle \lambda, \chi \rangle}.$$

The series converges absolutely for generic λ , and

$$w_M(h) = \lim_{\lambda \rightarrow 0} \sum_{P \in \mathcal{P}(M)} (-1)^{|\Delta_P^\lambda|} \overline{S}_P(\lambda),$$

where the limit is taken for generic $\lambda \in \mathfrak{a}_M^*$. The geometric series $\overline{S}_P(\lambda)$ can be calculated to be

$$\overline{S}_P(\lambda) = (-1)^{|\Delta_P^\lambda|} e^{\langle \lambda, h_P \rangle} \prod_{\alpha \in \Delta_P} \frac{1}{1 - e^{-\langle \lambda, \alpha^\vee \rangle}},$$

whence the result. □

Let $e_P(\lambda) = d_P(\lambda) \overline{c}_P(\lambda)^{-1}$, they form a (G, M) -family, still denoted by e . Lemma 4.3 expresses the lattice counting $w_M(h)$ as the volume of the product $(h \cdot e)$ of (G, M) -families, where h is the (G, M) -family associated to h . By the product formula (1.2) of Arthur, we get

$$w_M(h) = \sum_{Q \in \mathcal{F}(M)} v_M^Q(h) e'_Q. \quad (4.2)$$

Proposition 4.4. *We have*

$$v_\gamma(x) = \sum_{Q=LN_Q \in \mathcal{F}(M_0)} e'_Q \sum_{R=MN_R \in \mathcal{F}^L(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^M)} \cdot v_{M_0}^M(\mathrm{Ec}^{RN_Q}(x)) \cdot [\mathrm{Ec}^Q(x_0)]'_R.$$

Proof. By lemma 4.2, $v_\gamma(x)$ counts the number of lattice points in the polytope $\mathbf{E}(\lambda(x, x_0))$. Hence by the above formula,

$$v_\gamma(x) = w_{M_0}(\lambda(x, x_0)) = \sum_{Q \in \mathcal{F}(M_0)} v_{M_0}^Q(\lambda(x, x_0)) e'_Q.$$

Now that $\lambda(x, x_0) = \text{Ec}(x_0) - \text{Ec}(x)$, for $Q = LN_Q \in \mathcal{F}(M_0)$, we have

$$\begin{aligned} v_{M_0}^Q(\lambda(x, x_0)) &= v_{M_0}^L(\text{Ec}^Q(x_0) - \text{Ec}^Q(x)) \\ &= \sum_{R \in \mathcal{F}^L(M_0)} v_{M_0}^R(-\text{Ec}^Q(x)) \cdot [\text{Ec}^Q(x_0)]'_R \\ &= \sum_{R=MN_R \in \mathcal{F}^L(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^M)} \cdot v_{M_0}^M(\text{Ec}^{RN_Q}(x)) \cdot [\text{Ec}^Q(x_0)]'_R, \end{aligned}$$

where the last equality follows from a simple change of variable $\lambda \mapsto -\lambda$ in the definition (1.1) of the volume of a (G, M) -family. Putting everything together, we get the assertion. \square

Theorem 4.5. *We have*

$$|F_\gamma(\mathbb{F}_q)| = \text{vol}(T^1) |D(\gamma)|^{-\frac{1}{2}} \sum_{\substack{M, L \in \mathcal{L}(M_0) \\ M \subset L}} (-1)^{\dim(\mathfrak{a}_{M_0}^M)} J_{M_0}^M(\gamma, \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}) v_M^L(\text{Ec}(x_0)) e_L.$$

Proof. By lemma 4.1 and proposition 4.4, we have

$$\begin{aligned} |F_\gamma(\mathbb{F}_q)| &= \text{vol}(T^1) \int_{T \setminus G} \mathbb{1}_{\mathfrak{k}}(\text{Ad}(g^{-1})\gamma) v_\gamma(g) d\bar{g} \\ &= \text{vol}(T^1) \sum_{Q=LN_Q \in \mathcal{F}(M_0)} e'_Q \sum_{R=MN_R \in \mathcal{F}^L(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^M)} \cdot [\text{Ec}^Q(x_0)]'_R \\ &\quad \int_{T \setminus G} \mathbb{1}_{\mathfrak{k}}(\text{Ad}(g^{-1})\gamma) v_{M_0}^M(\text{Ec}^{RN_Q}(g)) d\bar{g}. \end{aligned}$$

With the Iwasawa decomposition $g = mnk$, $m \in M$, $n \in N_R N_Q$, $k \in K$, the factor $v_{M_0}^M(\text{Ec}^{RN_Q}(g))$ simplifies to $v_{M_0}^M(m)$, and

$$\begin{aligned} &\int_{T \setminus G} \mathbb{1}_{\mathfrak{k}}(\text{Ad}(g^{-1})\gamma) v_{M_0}^M(\text{Ec}^{RN_Q}(g)) d\bar{g} \\ &= \int_{T \setminus M} \int_{N_R N_Q} \int_K \mathbb{1}_{\mathfrak{k}}(\text{Ad}(k^{-1}n^{-1}m^{-1})\gamma) v_{M_0}^M(m) dk dnd\bar{m} \\ &= |\det(\text{ad}(\gamma); \mathfrak{g}/\mathfrak{m})|^{-\frac{1}{2}} \int_{T \setminus M} \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}(\text{Ad}(m^{-1})\gamma) v_{M_0}^M(m) d\bar{m} \\ &= |D(\gamma)|^{-\frac{1}{2}} J_{M_0}^M(\gamma, \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}). \end{aligned}$$

Plug in the above equation, $|F_\gamma(\mathbb{F}_q)|$ equals

$$\begin{aligned} &\text{vol}(T^1) |D(\gamma)|^{-\frac{1}{2}} \sum_{Q=LN_Q \in \mathcal{F}(M_0)} e'_Q \sum_{R=MN_R \in \mathcal{F}^L(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^M)} \cdot [\text{Ec}^Q(x_0)]'_R \cdot J_{M_0}^M(\gamma, \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}) \\ &= \text{vol}(T^1) |D(\gamma)|^{-\frac{1}{2}} \sum_{L \in \mathcal{L}(M_0)} e_L \sum_{M \in \mathcal{L}^L(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^M)} \cdot \text{Ec}_M^L(x_0) \cdot J_{M_0}^M(\gamma, \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}), \end{aligned}$$

where the second equality follows the fact that $\text{Ec}^Q(x_0) = \text{Ec}^L(x_0)$ and that

$$\sum_{R \in \mathcal{P}(M)} [\text{Ec}^L(x_0)]'_R = \text{Ec}_M^L(x_0), \quad \sum_{Q \in \mathcal{P}(L)} e'_Q = e_L,$$

as implied by the equation (1.3). □

We can reverse the above process to get an expression of $J_{M_0}(\gamma, \mathbb{1}_\mathfrak{k})$ in terms of $|F_\gamma^M(\mathbb{F}_q)|$, $M \in \mathcal{L}(M_0)$. Lemma 4.3 expresses $w_M(h)$ as the volume of the product $(h \cdot e)$. Since $h = (h \cdot e) \cdot e^{-1}$, we get inversely

$$v_{M_0}(h) = \sum_{Q \in \mathcal{F}(M_0)} w_{M_0}^Q(h)(e^{-1})'_Q. \quad (4.3)$$

Proposition 4.6. *For any $x \in \mathcal{X}_\gamma$, we have*

$$v_{M_0}(x) = \sum_{Q=LN_Q \in \mathcal{F}(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^L)} \text{Ec}(x_0)'_Q \sum_{R=MN_R \in \mathcal{F}^L(M_0)} v_\gamma^M(f_{RN_Q}(x)) [(e^{-1})^L]'_R.$$

Proof. By construction, $\lambda(x, x_0) = \text{Ec}(x_0) - \text{Ec}(x)$, hence

$$\begin{aligned} v_{M_0}(\text{Ec}(x)) &= v_{M_0}(\text{Ec}(x_0) - \lambda(x, x_0)) = \sum_{Q=LN_Q \in \mathcal{F}(M_0)} v_{M_0}^Q(-\lambda(x, x_0)) \cdot \text{Ec}(x_0)'_Q \\ &= \sum_{Q=LN_Q \in \mathcal{F}(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^L)} v_{M_0}^L(\lambda^Q(x, x_0)) \cdot \text{Ec}(x_0)'_Q \\ &= \sum_{Q=LN_Q \in \mathcal{F}(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^L)} \text{Ec}(x_0)'_Q \sum_{R=MN_R \in \mathcal{F}^L(M_0)} w_{M_0}^R(\lambda^Q(x, x_0)) [(e^{-1})^L]'_R. \end{aligned}$$

By definition, $w_{M_0}^R(\lambda^Q(x, x_0))$ counts the number of lattice points in the R -facet of $\lambda^Q(x, x_0)$, hence it equals $v_\gamma^M(f_{RN_Q}(x))$, whence the assertion. □

Theorem 4.7. *We have*

$$J_{M_0}(\gamma, \mathbb{1}_\mathfrak{k}) = \text{vol}(T^1)^{-1} \sum_{\substack{M, L \in \mathcal{L}(M_0) \\ M \subset L}} (-1)^{\dim(\mathfrak{a}_{M_0}^L)} |D^M(\gamma)|^{\frac{1}{2}} \cdot |F_\gamma^M(\mathbb{F}_q)| \cdot (e^{-1})_M^L \cdot \text{Ec}(x_0)_L.$$

Proof. By the above result, $J_{M_0}(\gamma, \mathbb{1}_\mathfrak{k})$ equals

$$\begin{aligned} |D(\gamma)|^{\frac{1}{2}} \int_{T \setminus G} \mathbb{1}_\mathfrak{k}(\text{Ad}(g^{-1})\gamma) \sum_{Q=LN_Q \in \mathcal{F}(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^L)} \text{Ec}(x_0)'_Q \cdot \\ \sum_{R=MN_R \in \mathcal{F}^L(M_0)} v_\gamma^M(f_{RN_Q}(g)) [(e^{-1})^L]'_R d\bar{g}. \end{aligned}$$

With the Iwasawa decomposition, we calculate

$$\begin{aligned}
& |D(\gamma)|^{\frac{1}{2}} \int_{T \backslash G} \mathbb{1}_{\mathfrak{k}}(\text{Ad}(g^{-1})\gamma) v_{\gamma}^M(f_{RN_Q}(g)) d\bar{g} \\
&= |D(\gamma)|^{\frac{1}{2}} \int_{T \backslash M} \int_{N_R N_Q} \int_K \mathbb{1}_{\mathfrak{k}}(\text{Ad}(k^{-1}n^{-1}m^{-1})\gamma) v_{\gamma}^M(m) dk dnd\bar{m} \\
&= |D^M(\gamma)|^{\frac{1}{2}} \int_{T \backslash M} \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}(\text{Ad}(m^{-1})\gamma) v_{\gamma}^M(m) d\bar{m} \\
&= |D^M(\gamma)|^{\frac{1}{2}} \cdot \text{vol}(T^1)^{-1} \cdot |F_{\gamma}^M(\mathbb{F}_q)|,
\end{aligned}$$

Plug in the above equation, $J_{M_0}(\gamma, \mathbb{1}_K)$ equals

$$\begin{aligned}
& \text{vol}(T^1)^{-1} \sum_{Q=LN_Q \in \mathcal{F}(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^L)} \text{Ec}(x_0)'_Q \cdot \sum_{R=MN_R \in \mathcal{F}^L(M_0)} |D^M(\gamma)|^{\frac{1}{2}} \cdot |F_{\gamma}^M(\mathbb{F}_q)| \cdot [(e^{-1})^L]'_R \\
&= \text{vol}(T^1)^{-1} \sum_{L \in \mathcal{L}(M_0)} (-1)^{\dim(\mathfrak{a}_{M_0}^L)} \text{Ec}(x_0)_L \cdot \sum_{M \in \mathcal{L}^L(M_0)} |D^M(\gamma)|^{\frac{1}{2}} \cdot |F_{\gamma}^M(\mathbb{F}_q)| \cdot (e^{-1})_M^L,
\end{aligned}$$

where in the second equation we have used the equation (1.3) to get

$$\sum_{Q \in \mathcal{P}(L)} \text{Ec}(x_0)'_Q = \text{Ec}(x_0)_L, \quad \sum_{R \in \mathcal{P}^L(M)} [(e^{-1})^L]'_R = (e^{-1})_M^L.$$

Rearrange the terms, we get the assertion. □

In [C2], the transition between weighted orbital integrals and counting points on the fundamental domains was extended to general Levi subgroup $M \in \mathcal{L}(M_0)$. This can be achieved also with our current method. Indeed, we have a reduction formula for $J_M(\gamma, \varphi)$.

Proposition 4.8. *Let $\gamma \in \mathfrak{g}$ be an element satisfying $M_{\gamma} = G_{\gamma}$, let $\varphi \in \mathcal{C}_c^{\infty}(\mathfrak{g})$, then*

$$J_M(\gamma, \varphi) = \sum_{L \in \mathcal{L}(M_0)} \theta_{M_0}^G(M, L) \cdot J_{M_0}^L(\gamma, \varphi^{Q_L^{\xi}}), \quad M \in \mathcal{L}(M_0),$$

where $Q_L^{\xi} \in \mathcal{P}(L)$ is uniquely determined by a generic element $\xi \in \mathfrak{a}_M^L$ as in equation (1.5).

Proof. By equation (1.5), we have

$$\begin{aligned}
J_M(\gamma, \varphi) &= |D(\gamma)|^{\frac{1}{2}} \int_{T \backslash G} \varphi(\text{Ad}(g^{-1})\gamma) \sum_{L \in \mathcal{L}(M_0)} \theta_{M_0}^G(M, L) v_{M_0}^{Q_L^{\xi}}(g) d\bar{g} \\
&= |D(\gamma)|^{\frac{1}{2}} \sum_{L \in \mathcal{L}(M_0)} \theta_{M_0}^G(M, L) \int_{T \backslash L} \int_{N_{Q_L^{\xi}}} \int_K \varphi(\text{Ad}(k^{-1}n^{-1}l^{-1})\gamma) v_{M_0}^L(l) d\bar{l} \\
&= \sum_{L \in \mathcal{L}(M_0)} \theta_{M_0}^G(M, L) |D^L(\gamma)|^{\frac{1}{2}} \int_{T \backslash L} \varphi^{Q_L^{\xi}}(\text{Ad}(l^{-1})\gamma) v_{M_0}^L(l) d\bar{l}.
\end{aligned}$$

□

Remark 4.1. The reduction formula is reminiscent of the corollary 3.10 in [C2], which reduces the counting points of the intermediate fundamental domain to that of the fundamental domains.

5 Rationality from homogeneity

This section is devoted to the proof of theorem 1.1. We work with the group $\mathbf{G} = \mathrm{GL}_{d+1}$, let \mathbf{A} be its maximal torus of the diagonal matrices, let $\gamma \in \mathfrak{a}$ be a regular integral element. Following Goresky, Kottwitz and MacPherson [GKM3], we define the *root valuation datum* of γ to be the function

$$R_\gamma : \Phi = \Phi(\mathbf{G}, \mathbf{A}) \rightarrow \mathbf{Z}, \quad \alpha \mapsto \mathrm{val}(\alpha(\gamma)).$$

Lemma 5.1. *There exists a set $\Delta = \{\alpha'_i\}_{i=1}^d$ of simple roots of Φ such that for any root $\alpha = \sum_{i=1}^d a_i \alpha'_i \in \Phi$, we have*

$$R_\gamma(\alpha) = \min \{R_\gamma(\alpha'_i) \mid a_i \neq 0\}.$$

In this case, we say that γ is in minimal form with respect to Δ .

Proof. The question involves only the root system, we proceed by induction on the dimension of the root system. The assertion holds obviously in dimension 1. Suppose that it holds for dimension less than d .

For $m \in \mathbf{Z}$, let $\Phi_{\gamma, m} = \{\alpha \in \Phi \mid R_\gamma(\alpha) \geq m\}$, then we get a decreasing filtration

$$\Phi = \Phi_{\gamma, 0} \supset \Phi_{\gamma, 1} \supset \Phi_{\gamma, 2} \supset \cdots. \quad (5.1)$$

According to Goresky, Kottwitz and MacPherson [GKM3], lemma 4.8.2, $\Phi_{\gamma, m}$ is \mathbf{Q} -closed, i.e. any \mathbf{Q} -linear combination of roots in $\Phi_{\gamma, m}$, if it is a root, must be an element of $\Phi_{\gamma, m}$. In particular, $\Phi_{\gamma, m}$ is itself a root system. Let m_0 be the smallest integer such that $\Phi_{\gamma, 0} \supsetneq \Phi_{\gamma, m_0}$. By induction hypothesis, there exists a set Δ' of simple roots in Φ_{γ, m_0} such that the restriction of the root valuation datum to Φ_{γ, m_0} satisfies the conclusion of the assertion. By [B], chap. 6, §7, prop. 24, the set Δ' can be extended to a set of simple roots Δ of Φ . It is easy to verify that Δ satisfies the property in the assertion with the strong triangular inequality of the valuation. \square

Since the affine Springer fiber \mathcal{X}_γ depends only on the conjugacy class of γ and that all the sets of simple roots are conjugate under the Weyl group W , by conjugating γ with a suitable element in W , we can assume that γ is in minimal form with respect to the set of simple roots $\Delta_0 = \{\alpha_i\}_{i=1}^d$ associated to the standard Borel subgroup \mathbf{B}_0 of the upper triangular matrices, and we use also the name *root valuation datum* of γ for the d -tuple $\mathbf{n} = (R_\gamma(\alpha_i))_{i=1}^d$. Obviously, for any $\mathbf{n} \in \mathbf{Z}_{\geq 0}^d$, we can find γ in minimal form with root valuation datum \mathbf{n} .

5.1 Generating series of weighted orbital integrals

To begin with, we investigate a similar generating series of the weighted orbital integrals. Suppose that γ is in minimal form with respect to Δ_0 and let \mathbf{n} be its root valuation. Let $m_1 < \cdots < m_l$ be the breaking points of the filtration (5.1), i.e. the filtration reduces to

$$\Phi_{\gamma, m_1} \supsetneq \Phi_{\gamma, m_2} \supsetneq \cdots \supsetneq \Phi_{\gamma, m_l}. \quad (5.2)$$

Then for any root $\alpha \in \Phi_{\gamma, m_i} \setminus \Phi_{\gamma, m_{i+1}}$, we have $R_\gamma(\alpha) = m_i$. Let $M_i \in \mathcal{L}(A)$ be the unique Levi subgroup with root system Φ_{γ, m_i} .

We have the following basic reduction: Let $\gamma' = \epsilon^{-m_1} \gamma$, then γ' remains integral with root valuation datum dropped by m_1 . In particular, for the roots in $\mathfrak{g}/\mathfrak{m}_2$, the root valuations are all 0, and we can write

$$\gamma' = s + \eta,$$

with $s \in \mathfrak{a}_{M_2}^{\text{reg}}$ and $\eta \in \mathfrak{m}_2 \cap U_s^+$ both integral and that the eigenvalues of $\text{ad}(s)$ on $\mathfrak{g}/\mathfrak{m}_2$ all have valuation 0. With the Arthur-Shalika expansion, i.e. theorem 3.6, for $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k})$, we can reduce the study of the weighted orbital integral $J_A(\gamma, f)$ to that of the Arthur-Shalika germs $g_A^L(\gamma, u)$ and the unipotent weighted orbital integrals $J_L(u, f)$, for $L \in \mathcal{L}(A)$ and $u \in \mathcal{N}_L(F)$. By theorem 3.7, $g_A^L(\gamma, u)$ can be expressed in terms of $g_A^M(\gamma', v) = g_A^M(s + \eta, v)$ with $M \in \mathcal{L}^L(A)$ and $v \in \mathcal{N}_M(F)$. We can make a further parabolic reduction for $g_A^M(s + \eta, v)$. Combining theorem 3.6 and theorem 3.11, we get two expansions of $J_A(s + \eta, f)$, which give rise to the linear equation

$$\sum_{M \in \mathcal{L}(A)} \sum_{v \in [\mathcal{N}_M(F)]} g_A^M(s + \eta, v) J_M(v, f) = \sum_{M' \in \mathcal{L}^{M_2}(A)} \sum_{u' \in [\mathcal{N}_{M'}(F)]} g_A^{M'}(\eta, u') J_{M'}(s + u', f).$$

Recall that g_A^M is constructed by induction on M , hence it is enough to work out g_A^G . Let f run through a dual basis of the distributions $J_G(u, \cdot)$, $u \in [\mathcal{N}_G(F)]$, we get a linear system. Solving them inductively in M and $v \in [\mathcal{N}_M(F)]$, we can express $g_A^M(s + \eta, v)$ in terms of $g_A^{M'}(\eta, u')$ with coefficients in $J_M(v, f)$ and $J_{M'}(s + u', f)$. Hence the problem is further reduced to the study of $g_A^{M'}$, $M' \in \mathcal{L}^{M_2}(A)$, i.e. to the study of the local behavior of $J_A^{M_2}(\eta, \cdot)$ for the group M_2 . Briefly, the study of $J_A(\gamma, \cdot)$ can be reduced to that of

$$\begin{cases} J_M(u, \cdot), & \text{for } M \in \mathcal{L}(A), u \in [\mathcal{N}_M(F)], \\ J_{M'}(s + u', \cdot), & \text{for } M' \in \mathcal{L}^{M_2}(A), u' \in [\mathcal{N}_{M'}(F)], \\ J_A^{M_2}(\eta, \cdot). \end{cases}$$

This process can be carried on for $J_A^{M_2}(\eta, \cdot)$ and so on. At the end, the study of $J_A(\gamma, \cdot)$ can be reduced to

$$\begin{cases} J_M^{M_i}(u, \cdot), & \text{for } M \in \mathcal{L}^{M_i}(A), u \in [\mathcal{N}_M(F)], \\ J_{M'}^{M_i}(s_i + u', \cdot), & \text{for } M' \in \mathcal{L}^{M_{i+1}}(A), u' \in [\mathcal{N}_{M'}(F)], \end{cases} \quad (5.3)$$

where $s_i \in \mathfrak{a}_{M_{i+1}}^{\text{reg}}$ are integral such that the eigenvalues of $\text{ad}(s_i)$ on $\mathfrak{m}_i/\mathfrak{m}_{i+1}$ all have valuation 0, $i = 1, \dots, l$. For $L \in \mathcal{L}(A)$, let $\mathfrak{a}_L^{\text{reg},0}$ be the open subset of $\mathfrak{a}_L^{\text{reg}}$ consisting of the integral elements s such that the eigenvalues of $\text{ad}(s)$ on $\mathfrak{g}/\mathfrak{l}$ all have valuation 0.

Lemma 5.2. *Let $L \in \mathcal{L}(A)$ be such that $\dim(A_L) = \dim(A_G) + 1$, let $s \in \mathfrak{a}_L^{\text{reg},0}$. Then for any K -invariant $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k})$, $M \in \mathcal{L}^L(A)$ and $u \in [\mathcal{N}_M(F)]$, as a function of $s \in \mathfrak{a}_L^{\text{reg},0}$, the weighted orbital integral $J_M(s + u, f)$ is constant.*

Proof. By definition,

$$J_M(s + u, f) = \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \sum_{M' \in \mathcal{L}(M)} r_M^{M'}(u, a) J_{M'}(s + u + a, f).$$

Take $Q \in \mathcal{P}(L)$, then Q is a maximal parabolic subgroup of G . With the Iwasawa decomposition and the K -invariance of f , we have

$$J_{M'}(s + u + a, f) = \int_{M_u \backslash L} \int_{N_Q} f(\text{Ad}(n^{-1}l^{-1})(s + u + a)) \nu_{M'}(ln) \, dn d\bar{l} \quad (5.4)$$

Without loss of generality, we can assume that f is the characteristic function $[\omega + \mathfrak{k}]$ for a K -invariant subset $\omega \subset \mathfrak{g}$. For the integrand to be non-zero, we need to solve the equation

$$\text{Ad}(n^{-1}l^{-1})(s + u + a) = \text{Ad}(n^{-1})(s + \text{Ad}(l^{-1})(a + u)) \in \omega + \mathfrak{k}. \quad (5.5)$$

Let $\gamma = \text{Ad}(l^{-1})(a + u)$ and $n = \exp(\nu), \nu \in \mathfrak{n}_Q$, the equation can be rewritten as

$$(s + \gamma) + \text{ad}(s + \gamma)\nu \in \omega + \mathfrak{k},$$

which further simplifies to

$$\gamma + \text{ad}(s + \gamma)\nu \in \omega + \mathfrak{k}. \quad (5.6)$$

Notice that the left hand side belongs to \mathfrak{q} , hence it actually lies in $\mathfrak{q} \cap (\omega + \mathfrak{k})$. Consider the conjugation action of L on $\mathfrak{q} \cap (\omega + \mathfrak{k})$, there exist finitely L -orbits $\omega_i \subset \mathfrak{l}, i \in I$, such that

$$\mathfrak{q} \cap (\omega + \mathfrak{k}) = \bigsqcup_{i \in I} (\omega_i + (\mathfrak{l} \cap \mathfrak{k}) + \mathfrak{n}_Q) \cap (\omega + \mathfrak{k}).$$

Hence the set of solutions Ω to the equation (5.6) is the disjoint union of the set of solutions Ω_i to the equation (5.6) and the equation

$$\gamma = \text{Ad}(l^{-1})(a + u) \in \omega_i + (\mathfrak{l} \cap \mathfrak{k}), \quad (5.7)$$

which is independent of s .

Given another $s' \in \mathfrak{a}_L^{\text{reg},0}$, we can find $k_\gamma \in L \cap K$ such that

$$\text{ad}(s' + \gamma) = \text{ad}(s + \gamma) \circ \text{Ad}(k_\gamma) \quad \text{on } \mathfrak{n}_Q.$$

Then the equation (5.6) for s' can be rewritten as

$$\gamma + [s + \gamma, \text{Ad}(k_\gamma)\nu] \in \omega + \mathfrak{k}.$$

Hence the set of solutions of the equation (5.6) for s and s' are related by

$$\nu' = \text{Ad}(k_\gamma^{-1})\nu, \quad l' = l. \quad (5.8)$$

We make this change of variable to relate $J_{M'}(s + u + a)$ and $J_{M'}(s' + u + a)$. Notice that as $a \rightarrow 0$ in $\mathfrak{a}_M^{\text{reg}}$, the action of $\text{Ad}(k_\gamma)$ approaches to that of $\text{Ad}(k_0)$ for an element $k_0 \in A_L \cap K$, hence

$$\lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \nu_{M'}(l \cdot k_\gamma^{-1} n k_\gamma) = \nu_{M'}(l \cdot k_0^{-1} n k_0) = \nu_{M'}(k_0^{-1} l \cdot n k_0) = \nu_{M'}(ln), \quad (5.9)$$

as $\nu_{M'}$ is left- M' -invariant and right- K -invariant. Moreover, as $\nu_{M'}$ is discrete, the above equality holds for a sufficiently close to 0, but depending on l . Since f is compactly supported on \mathfrak{g} , we have

$$\begin{aligned} J_M(s' + u, f) &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \sum_{M' \in \mathcal{L}(M)} r_M^{M'}(u, a) J_{M'}(s' + u + a, f) \\ &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \int_{M_u \setminus L} \int_{N_Q} f(\text{Ad}(n^{-1}l^{-1})(s' + u + a)) \sum_{M' \in \mathcal{L}(M)} r_M^{M'}(u, a) \nu_{M'}(ln) \, dnd\bar{l} \\ &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \int_{M_u \setminus L} \int_{N_Q} f(\text{Ad}(n^{-1}l^{-1})(s + u + a)) \sum_{M' \in \mathcal{L}(M)} r_M^{M'}(u, a) \nu_{M'}(l \cdot k_\gamma^{-1} n k_\gamma) \, dnd\bar{l} \\ &= \lim_{\substack{a \in \mathfrak{a}_M^{\text{reg}} \\ a \rightarrow 0}} \int_{M_u \setminus L} \int_{N_Q} f(\text{Ad}(n^{-1}l^{-1})(s + u + a)) \sum_{M' \in \mathcal{L}(M)} r_M^{M'}(u, a) \nu_{M'}(ln) \, dnd\bar{l} \\ &= J_M(s + u, f), \end{aligned}$$

where the next to the last equality is due to the fact that for l in a bounded domain of L the limit (5.9) holds for a sufficiently close to 0. □

Theorem 5.3. Let $\mathbf{G} = \mathrm{GL}_{d+1}$, let $\gamma \in \mathfrak{a}$ be a regular integral element, then for any $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k})$, the weighted orbital integral $J_A(\gamma, f)$ depends only on the root valuation datum \mathbf{n} of γ , and we denote it by $J_A(\mathbf{n}, f)$. Moreover, the generating series

$$J_f(\vec{t}) := \sum_{n_1=0}^{+\infty} \cdots \sum_{n_d=0}^{+\infty} J_A((n_1, \dots, n_d), f) t_1^{n_1} \cdots t_d^{n_d} \in \mathbf{C}[[t_1, \dots, t_d]]$$

is a rational fraction, i.e. it is an element of $\mathbf{C}(t_1, \dots, t_d)$.

The theorem is an immediate consequence of a similar assertion about the Arthur-Shalika germs by theorem 3.6, we state it separately because it has independent interest.

Theorem 5.4. Let $\mathbf{G} = \mathrm{GL}_{d+1}$, let $\gamma \in \mathfrak{a}$ be a regular integral element. For $M \in \mathcal{L}(A)$, $u \in \mathcal{N}_M(F)$, the Arthur-Shalika germ $g_A^M(\gamma, u)$ depends only on the root valuation datum \mathbf{n} of γ , and we denote it by $g_A^M(\mathbf{n}, u)$. Moreover, the generating series

$$g_A^M(u; \vec{t}) := \sum_{n_1=0}^{+\infty} \cdots \sum_{n_d=0}^{+\infty} g_A^M((n_1, \dots, n_d), u) t_1^{n_1} \cdots t_d^{n_d} \in \mathbf{C}[[t_1, \dots, t_d]]$$

is a rational fraction, i.e. it is an element of $\mathbf{C}(t_1, \dots, t_d)$.

Proof. We proceed by induction on $\mathrm{rk}(\mathbf{G})$. The theorem holds trivially if $\mathrm{rk}(\mathbf{G}) = 0$. In general, the Arthur-Shalika germs satisfy the homogeneity and parabolic descent by theorem 3.7 and 3.11. Applying the basic reduction at the beginning of the section, the first assertion can be reduced to the trivial case of $\mathrm{rk}(\mathbf{G}) = 0$. Indeed, the homogeneity reduces us to the case that $\gamma' = s + \eta$ with $s \in \mathfrak{a}_{M_2}^{\mathrm{reg},0}$ and $\eta \in \mathfrak{m}_2 \cap U_s^+$ both integral. By theorem 3.7, this process depends only on the root valuation datum of γ . Then we apply the parabolic descent of the Arthur-Shalika germ, i.e. theorem 3.11. By the transitivity of the parabolic descent, we can assume that $\dim(A_{M_2}) = \dim(A_G) + 1$. By lemma 5.2, for any K -invariant test function $f \in \mathcal{C}_c(\mathfrak{g}/\mathfrak{k})$, $M' \in \mathcal{L}^{M_2}(A)$, $u' \in \mathcal{N}_{M'}(F)$, the weighted orbital integral $J_{M'}(s + u', f)$ is constant as a function of $s \in \mathfrak{a}_{M_2}^{\mathrm{reg},0}$. Solving the linear system from two expansions of $J_A(s + \eta, f)$:

$$\sum_{M \in \mathcal{L}(A)} \sum_{u \in [\mathcal{N}_M(F)]} g_A^M(s + \eta, u) J_M(u, f) = \sum_{M' \in \mathcal{L}^{M_2}(A)} \sum_{u' \in [\mathcal{N}_{M'}(F)]} g_A^{M'}(\eta, u') J_{M'}(s + u', f),$$

with f running through the characteristic functions $[\mathbf{G}^{(k)}(\epsilon^{-1}u) + \mathfrak{k}]$, $u \in [\mathcal{N}_G(k)]$, we get a linear transformation between $\{g_A^M(s + \eta, u)\}$ and $\{g_A^{M'}(\eta, u')\}$, which is constant as a function of $s \in \mathfrak{a}_{M_2}^{\mathrm{reg},0}$. Repeating these process, the question is reduced to the case of $\mathrm{rk}(\mathbf{G}) = 0$.

Let $\Phi_\bullet(\mathbf{n})$ be the reduced strict filtration associated to (5.2) with $\Phi_i(\mathbf{n}) = \Phi_{\gamma, m_i}$, $i = 1, \dots, l$. We regroup the summation in the expression of $g_A^M(u; \vec{t})$ according to the filtration of roots $\Phi_\bullet(\mathbf{n})$. Let Φ_\bullet be a filtration

$$\Phi_1 = \Phi \supsetneq \Phi_2 \supsetneq \cdots \supsetneq \Phi_l$$

of Φ by \mathbf{Q} -closed sub root systems. We denote loosely $\Phi_\bullet(\mathbf{n}) = \Phi_\bullet$ if the filtrations have the same length and that $\Phi_{m_i}(\mathbf{n}) = \Phi_i$ for all the breaking points m_i of the filtration $\Phi_\bullet(\mathbf{n})$. Let

$$S_{\Phi_\bullet}^{G,M}(u; \vec{t}) := \sum_{\mathbf{n} \in \mathbf{Z}_{>0}^d, \Phi_\bullet(\mathbf{n}) = \Phi_\bullet} g_A^M(\mathbf{n}, u) t_1^{n_1} \cdots t_d^{n_d}.$$

Then the assertion is reduced to showing that $S_{\Phi_\bullet}^{G,M}(u; \vec{t})$ is rational, as there are only finitely many filtrations Φ_\bullet . We will prove it by induction on the semisimple rank of the groups. As a starting point, the assertion is clearly true for $\text{rk}(\mathbf{G}) = 0$.

We break the summation in $S_{\Phi_\bullet}^{G,M}(u; \vec{t})$ into two terms, $S_1(u; \vec{t})$ and $S_2(u; \vec{t})$, according to whether $m_1 = 0$ or not. For the term $S_1(u; \vec{t})$, as we have explained before with parabolic descent of Arthur-Shalika germs, there is a linear transformation between $\{g_A^M(\mathbf{n}, u)\}$ and $\{g_A^{M'}(\bar{\mathbf{n}}, u')\}$, where $\bar{\mathbf{n}}$ is obtained from \mathbf{n} by removing all the 0's. Then the power series $S_1(u; \vec{t})$ can be written as a linear combination of $S_{\Phi'_\bullet}^{M_2, M'}(u'; \vec{t})$ for $M' \in \mathcal{L}^{M_2}(A)$ and $u' \in [\mathcal{N}_{M'}(F)]$. By induction, it must be rational.

Now that the rationality of $S_1(u; \vec{t})$ is established, with the homogeneity of the Arthur-Shalika germs, we can make the equation

$$S_{\Phi_\bullet}^{G,M}(u; \vec{t}) = S_1(u; \vec{t}) + S_2(u; \vec{t})$$

into a recurrence equation. Let $\mathbf{1} = (1, \dots, 1) \in \mathbf{Z}^d$. By theorem 3.7, we have

$$\begin{aligned} S_{\Phi_\bullet}^{G,M}(u; \vec{t}) &= S_1(u; \vec{t}) + \sum_{\mathbf{n} \in \mathbf{N}^d, \Phi_\bullet(\mathbf{n}) = \Phi_\bullet} g_A^M(\mathbf{n}, u) t_1^{n_1} \cdots t_d^{n_d} \\ &= S_1(u; \vec{t}) + \sum_{\mathbf{n} \in \mathbf{Z}_{\geq 0}^d, \Phi_\bullet(\mathbf{n}) = \Phi_\bullet} g_A^M(\mathbf{n} + \mathbf{1}, u) t_1^{n_1+1} \cdots t_d^{n_d+1} \\ &= S_1(u; \vec{t}) + \sum_{\mathbf{n} \in \mathbf{Z}_{\geq 0}^d, \Phi_\bullet(\mathbf{n}) = \Phi_\bullet} q^{-d^M(u)} \sum_{M' \in \mathcal{L}^M(A)} \sum_{v \in [\mathcal{N}_{M'}(F)]} g_A^{M'}(\mathbf{n}, v) c_{M'}^M(v, \epsilon) \\ &\quad \cdot [v_{M'}^M : u/\epsilon] t_1^{n_1+1} \cdots t_d^{n_d+1} \\ &= S_1(u; \vec{t}) + q^{-d^M(u)} \left[c_M^M(u/\epsilon, \epsilon) S_{\Phi_\bullet}^{G,M}(u/\epsilon; \vec{t}) t_1 \cdots t_d + \right. \\ &\quad \left. \sum_{\substack{M' \in \mathcal{L}^M(A) \\ M' \neq M}} \sum_{v \in [\mathcal{N}_{M'}(F)]} [v_{M'}^M : u/\epsilon] c_{M'}^M(v, \epsilon) S_{\Phi_\bullet}^{G,M'}(v; \vec{t}) t_1 \cdots t_d \right]. \end{aligned}$$

Since u and u/ϵ belongs to the same nilpotent orbit in $\mathcal{N}_M(F)$, we have

$$S_{\Phi_\bullet}^{G,M}(u; \vec{t}) = S_{\Phi_\bullet}^{G,M}(u/\epsilon; \vec{t}),$$

and the above equation defines a linear system of recurrence relation among the $S_{\Phi_\bullet}^{G,M}(u; \vec{t})$'s. For $M = A$, the system reduces to

$$S_{\Phi_\bullet}^{G,A}(0; \vec{t}) = S_1(0; \vec{t}) + c_A^A(0, \epsilon) S_{\Phi_\bullet}^{G,A}(0; \vec{t}) t_1 \cdots t_d.$$

Now that $S_1(0; \vec{t})$ is rational, we get the rationality of $S_{\Phi_\bullet}^{G,A}(0; \vec{t})$. The rationality of $S_{\Phi_\bullet}^{G,M}(u; \vec{t})$ then follows from the above recurrence relation by induction on the Levi subgroups M and the nilpotent orbits. \square

5.2 Generating series of the affine Springer fibers

We resume the discussions in §4. By theorem 4.5, we have

$$|F_\gamma(\mathbb{F}_q)| = |D(\gamma)|^{-\frac{1}{2}} \sum_{\substack{M, L \in \mathcal{L}(A) \\ M \subset L}} (-1)^{\dim(\mathfrak{a}_A^M)} J_A^M(\gamma, \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}) v_M^L(\text{Ec}(x_0)) e_L. \quad (5.10)$$

Theorem 5.5. Let $\mathbf{G} = \mathrm{GL}_{d+1}$. For regular element $\gamma \in \mathfrak{a}$, the number of points $|F_\gamma(\mathbb{F}_q)|$ is a polynomial in q depending only on the root valuation datum \mathbf{n} of γ , denoted $F_{\mathbf{n}}(q)$. Moreover, the generating series

$$F(q; \vec{t}) := \sum_{n_1=0}^{+\infty} \cdots \sum_{n_d=0}^{+\infty} F_{(n_1, \dots, n_d)}(q) t_1^{n_1} \cdots t_d^{n_d} \in \mathbf{Z}[[q; t_1, \dots, t_d]]$$

is a rational fraction, i.e. it is an element of $\mathbf{Z}(q; t_1, \dots, t_d)$.

Proof. The assertion that $|F_\gamma(\mathbb{F}_q)|$ is polynomial in q results from the fact that the group A acts on F_γ with finitely many fixed points. Its dependence on the root valuation datum results from the similar property for $J_A^M(\gamma, \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}})$ by equation (5.10). For the second assertion, it is enough to show that the generating series for $J_A^M(\gamma, \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}) v_M^L(\mathrm{Ec}(x_0))$ is rational. By equation (1.5), we have

$$v_M^L(\mathrm{Ec}(x_0)) = \sum_{M' \in \mathcal{L}^L(A)} \theta_A^L(M, M') v_A^{M'}(\mathrm{Ec}(x_0)),$$

with $\theta_A^L(M, M') \neq 0$ if and only if the natural projection

$$\mathfrak{a}_A^M \oplus \mathfrak{a}_A^{M'} \rightarrow \mathfrak{a}_A^L$$

is an isomorphism. We call M, M' *transversal* if the condition holds for some $L \in \mathcal{L}(A)$. The question is then reduced to showing that the generating series for $J_A^M(\gamma, \mathbb{1}_{\mathfrak{m} \cap \mathfrak{k}}) v_A^{M'}(\mathrm{Ec}(x_0))$ is rational for all transversal M, M' . With the Arthur-Shalika germ expansion, the question is further reduced to the rationality of the generating series of $g_A^{M''}(\gamma, u) v_A^{M'}(\mathrm{Ec}(x_0))$ for $M'' \in \mathcal{L}^M(A)$ and $u \in [\mathcal{N}_{M''}(F)]$. Notice that M', M'' are transversal again.

Suppose that γ is in minimal form with root valuation $\mathbf{n} \in \mathbf{Z}_{\geq 0}^d$ with respect to \mathbf{B}_0 . It is then also in minimal form for the groups M', M'' with respect to $\mathbf{B}_0 \cap M'$ and $\mathbf{B}_0 \cap M''$ respectively. The recursion of $g_A^{M''}(\gamma, u)$ for $\mathbf{n} \mapsto \mathbf{n} + \mathbf{1}$ is given by the homogeneity of Shalika germs, the theorem 3.7. For $v_A^{M'}(\mathrm{Ec}(x_0))$, notice that the (G, A) -family associated to $-\mathrm{Ec}(x_0)$ is equal to $r(\lambda; 0, \gamma)$ up to multiplication by a global function on $(\mathfrak{a}_A^G)^*$. To emphasize its dependence on the root valuation datum, we denote $\mathrm{Ec}(x_0)$ by $\Pi_{\mathbf{n}}$. Then similar to [A3], lemma 10.1, with the product formula (1.4), we can show

$$v_A^{M'}(\Pi_{\mathbf{n}+\mathbf{1}}) = \sum_{J \in \mathcal{L}^{M'}(A)} v_A^J(\Pi_{\mathbf{n}}) c_J^{M'}(0, \epsilon),$$

as $\Pi_{\mathbf{n}}$ satisfies the extra conditions on the facets leading to equation (1.4). Combining their recursion relations, we get

$$v_A^{M'}(\Pi_{\mathbf{n}+\mathbf{1}}) g_A^{M''}(\mathbf{n} + \mathbf{1}, u) = q^{-d^{M''}(u)} \sum_{J' \in \mathcal{L}^{M'}(A)} \sum_{J'' \in \mathcal{L}^{M''}(A)} \sum_{v \in [\mathcal{N}_{J''}(F)]} v_A^{J'}(\Pi_{\mathbf{n}}) g_A^{J''}(\mathbf{n}, v) c_{J'}^{M'}(0, \epsilon) c_{J''}^{M''}(v, \epsilon) [v_{J''}^{M''} : u/\epsilon].$$

Notice that J' and J'' are transversal, like M' and M'' . The rationality of the generating series for $v_A^{M'}(\Pi_{\mathbf{n}}) g_A^{M''}(\mathbf{n}, u)$ then results from the recurrence relation, following the same lines of reasoning as in the proof of theorem 5.4. \square

References

- [A1] J. Arthur, *The characters of discrete series as orbital integrals*, Invent. math. **32**(1976), 205-261.
- [A2] J. Arthur, *The trace formula in invariant form*, Ann. of Math., 114 (1981), 1-74.
- [A3] J. Arthur, *The local behavior of weighted orbital integrals*, Duke. Math. J., Vol 56, No. 2, 1988.
- [A4] J. Arthur, *An introduction to the trace formula*. Harmonic analysis, the trace formula, and Shimura varieties, 1–263, Clay Math. Proc., 4, Amer. Math. Soc., Providence, RI, 2005.
- [B] N. Bourbaki, *Groupes et algèbres de Lie*. Chapitre 4-6. Masson, Paris, 1982.
- [Bez] R. Bezrukavnikov, *The dimension of the fixed point set on affine flag manifolds*, Math. Res. Lett. 3 (1996), 185–189.
- [C1] Z. Chen, *On the fundamental domain of affine Springer fibers*. Math. Z. 286 (2017), no. 3-4, 1323-1356.
- [C2] Z. Chen, *Truncated affine Springer fibers and Arthur’s weighted orbital integrals*, <https://arxiv.org/abs/1701.03202>.
- [CL1] P-H. Chaudouard, G. Laumon, *Sur l’homologie des fibres de Springer affines tronquées*, Duke Math. J. 145 (2008), no. 3, 443–535.
- [CL2] P-H. Chaudouard, G. Laumon, *Le lemme fondamental pondéré. I. Constructions géométriques*. Compos. Math. 146 (2010), no. 6, 1416–1506.
- [D1] S. DeBacker, *Homogeneity of certain invariant distributions on the Lie algebra of p -adic GL_n* , Compos. Math. 124 (1) (2000) 11-16.
- [D2] S. DeBacker, *Homogeneity results for invariant distributions of a reductive p -adic group*. Ann. Sci. École Norm. Sup. (4) 35 (2002), no. 3, 391-422.
- [D3] S. DeBacker, *Parametrizing nilpotent orbits via Bruhat-Tits theory*. Ann. of Math. (2) 156 (2002), no. 1, 295-332.
- [GKM1] M. Goresky, R. Kottwitz, R. MacPherson, *Homology of affine Springer fibers in the unramified case*, Duke Math. J. 121 (2004), no. 3, 509-561.
- [GKM2] M. Goresky, R. Kottwitz, R. MacPherson, *Regular points in affine Springer fibers*, Michigan Math. J. 53 (2005), no. 1, 97-107.
- [GKM3] M. Goresky, R. Kottwitz, R. MacPherson, *Codimensions of root valuation strata*, Pure. Appl. Math. Q. 5 (2009), no. 4, 1253-1310.
- [K1] R. E. Kottwitz, *Isocrystals with additional structure*. Compositio Math. 56 (1985), no. 2, 201–220.
- [K2] R. E. Kottwitz, *Harmonic analysis on reductive p -adic groups and Lie algebras*. Harmonic analysis, the trace formula, and Shimura varieties, 393–522, Clay Math. Proc., 4, Amer. Math. Soc., Providence, RI, 2005

- [KL] D. Kazhdan, G. Lusztig, *Fixed point varieties on affine flag manifolds*, Israel. J. Math. **62**(1988), 129-168.
- [KV] R. Kottwitz, E. Viehman, *Generalized affine Springer fibers*, J. Inst. Math. Jussieu 11 (2012), no. 3, 569–609.
- [MP] A. Moy, G. Prasad, *Unrefined minimal K -types for p -adic groups*, Invent. Math. 116 (1994), 393–408.
- [R] R. Ranga Rao, *Orbital integrals in reductive groups*, Annals of Mathematics, Second Series, Vol. 96, No. 3 (Nov., 1972), pp. 505-510.
- [W] J.-L. Waldspurger, *Une formule des traces locale pour les algèbres de Lie p -adiques*. J. Reine Angew. Math. 465 (1995), 41–99.
- [We] A. Weil, Adeles and algebraic groups, with appendices by M. Demazure and Takashi Ono. Progress in Mathematics, 23. Birkhäuser, Boston, Mass., 1982.

Zongbin CHEN

Yau Mathematical Science Center,
Jinchunyuan west building, office 255, Tsinghua University,
Haidian district, 100084,
Beijing, China
email: zbchen@tsinghua.edu.cn