

Solving Systems of Equations of Raising-to-Powers Type

Francesco Paolo Gallinaro

November 25, 2021

Abstract

We address special cases of the analogues of the exponential algebraic closedness conjecture relative to the exponential maps of semiabelian varieties and to the modular j function. In particular, we show that the graph of the exponential of an abelian variety intersects products of free rotund varieties in which the subvariety of the domain is a sufficiently generic linear subspace, and that the graph of j intersects products of free broad varieties in which the subvariety of the domain is a Möbius subvariety.

1 Introduction

Zilber’s study of the model theory of complex exponentiation has originated a lot of work around systems of exponential polynomial equation. The main conjecture, known as *exponential algebraic closedness* and first stated in [Zil05], predicts sufficient conditions for an algebraic subvariety of $\mathbb{C}^n \times (\mathbb{C}^\times)^n$ to intersect the graph of the exponential function. It can be interpreted as a dual statement to Schanuel’s conjecture, a famous conjecture from transcendental number theory.

Conjecture 1.1 (Schanuel). *Let $z_1, \dots, z_n \in \mathbb{C}^n$ be \mathbb{Q} -linearly independent complex numbers. Then*

$$\text{trdeg}(z_1, \dots, z_n, e^{z_1}, \dots, e^{z_n}) \geq n.$$

Schanuel’s conjecture predicts that algebraic varieties of low dimension defined over \mathbb{Q} can’t intersect the graph of the exponential function, unless they are forced to do so by the fact that \exp is a group homomorphism. The exponential algebraic closedness conjecture can then be informally stated as “all varieties which are allowed to intersect the graph of \exp according to Schanuel’s conjecture do intersect it.” This is interpreted precisely through the technical conditions of “freeness” and “rotundity”, which are expected to be sufficient for an algebraic subvariety of $\mathbb{C}^n \times (\mathbb{C}^\times)^n$ to intersect the graph of \exp . For example, the case $n = 1$ states that if $p \in \mathbb{C}[X, Y]$ is a polynomial depending on both variables then there are infinitely many points $z \in \mathbb{C}$ such that $p(z, e^z) = 0$;

as discussed in [Mar06], this can be proved by using standard techniques from complex analysis.

The geometric interpretation of the exponential algebraic closedness conjecture has led to the formulation of other questions, in connection with other Schanuel-type statements. In fact, Schanuel’s conjecture fits in a rather broad framework in transcendental number theory; we refer the interested reader to [Ber02]. In particular, similar statements have been considered for the exponential maps of semiabelian varieties (see [BK18]) and the modular j -function (see [AK20]). These statements take similar forms, predicting sufficient conditions for some complex algebraic varieties to intersect the graph of the analytic function under consideration. Various special cases of these conjectures have been addressed: we recall [Zil02], [Zil15], [MZ16], [BM17], [DFT18], [Kir19] for the exponential function, [Kir09] for semiabelian varieties, [Asl18], [EH20], [AK20], [AEK21] for the modular j invariant.

We are particularly interested in [Zil02] and [Zil15]. There, Zilber proved a special case of the conjecture for \exp , studying intersections between the graph of \exp and varieties of the form $L \times W$, satisfying the freeness and rotundity conditions mentioned above, where L is an \mathbb{R} -linear subspace of \mathbb{C}^n and W is an algebraic subvariety of $(\mathbb{C}^\times)^n$. In particular, [Zil02] finds intersections for this kind of varieties assuming a diophantine conjecture, while [Zil15] removes the dependence on the conjecture but requires L to be defined over some “generic” subfield of the reals.

Here, drawing inspiration from Zilber’s results in [Zil02] and [Zil15], we solve some special cases of the existential closedness conjectures for exponential maps of abelian varieties and for the j function. In particular, we consider subvarieties which split as products of two varieties, one of which is a subvariety of the domain of a particular form (suggested by the structure induced by the relative analytic function) while the other one is a general subvariety of the codomain.

In Section 2 we consider abelian varieties. Using the fact that abelian varieties are complex Lie groups, and their subgroups are thus well-understood thanks to Cartan’s closed subgroup theorem, we obtain intersections between the graph of the exponential \exp of an abelian variety A and a sufficiently generic product of the form $L \times W$ which satisfies the technical conditions of *freeness* and *rotundity*, which will be defined later. The genericity condition is precise and stated in terms of the lattice Λ such that $A \cong \mathbb{C}^g/\Lambda$; the fundamental notions concerning the correspondence between lattices and abelian varieties will also be recalled.

Theorem 2.12. *Let (L, W) be a free rotund pair in $LA \times A$, and suppose L is not contained in any hyperplanes defined over the conjugate of the dual of the lattice Λ . Then $\exp(L) \cap W$ is dense in W in the Euclidean topology.*

The additional assumption on L makes $\exp(L)$ a dense subgroup of A in the Euclidean topology; in general, it is only Zariski-dense. It seems that it should be possible to remove this assumption; that is the subject of ongoing work.

In Section 3 we obtain a similar result for j ; the assumption on subvarieties of the domain is now that they are *Möbius subvarieties* of \mathbb{H}^n (\mathbb{H} denotes the

complex upper half plane.) These are intersections with \mathbb{H}^n of complex algebraic varieties which are defined by using the action of the group $\mathrm{SL}_2(\mathbb{R})$ on \mathbb{H} by Möbius transformations: a matrix $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ acts on z by taking it to $gz = \frac{az+b}{cz+d}$, and a Möbius variety is a variety that can be defined by only using conditions of the form $z_i = c$ (constant coordinates) and $z_j = gz_i$. The geometry of Möbius transformations, and other general facts about the j function, will be reviewed at the beginning of Section 3. Again, the statement also relies on the technical conditions of *freeness* and *broadness*, which will be explained later on.

Theorem 3.31. *Let L be a Möbius variety, (L, W) be a free broad pair. Then W contains a dense subset of points of $j(L)$.*

As the j function is algebraically independent from its first two derivatives, results in this area often take into consideration equations which involve j, j' and j'' (see for example [AK20]). The methods of this paper don't seem to work if we consider all three functions together, and only allow us to work with one function at a time, yielding only some partial results. This is explained in Section 4, where we sketch the proof of a partial result that can be obtained for j' ; we also note that some technical difficulties prevent us from getting a similar result for j'' .

While the motivation for these questions comes from model theory, the methods that we use are mainly geometric, ergodic- and number-theoretic. Hence, we have tried to make this exposition as self-contained as possible, introducing all the necessary notions at the beginning of each section and providing detailed proofs.

Acknowledgements. I would like to thank my supervisor, Vincenzo Mantova, for suggesting to work on this topic, for his countless suggestions, and for the valuable feedback that he has provided. Thanks to Gareth Jones, whose suggestion to use Ratner's theorem to tackle Lemma 3.13 proved correct, and to Vahagn Aslanyan and Jonathan Kirby for many interesting conversations around these topics. This research was done as part of the author's Ph.D. project, supported by a scholarship from the School of Mathematics of the University of Leeds.

2 Abelian Varieties

2.1 Background and Notation

In this section we will deal with complex abelian varieties, i.e., connected complex projective algebraic groups.

A will be used to denote an abelian variety. The set $A(\mathbb{C})$ of complex points of A has a differentiable structure as a complex manifold, and hence it is a complex Lie group. As such, we can consider the Lie algebra LA , its tangent space at identity. We'll denote by TA the tangent bundle of A : this is a fibre

bundle over A in which the fibre over each point $\bar{x} \in A$ is the tangent space to A at \bar{x} . Tangent spaces are functorial constructions, hence we'll freely use the fact that to any regular morphism $f : A \rightarrow B$ of abelian varieties we can associate morphisms $Lf : LA \rightarrow LB$ and $Tf : TA \rightarrow TB$.

We will use $+$ and $-$ to denote the operations on the Lie algebra, while \oplus and \ominus will denote the operations on both A and the tangent bundle TA (it will be clear from context which one we'll be referring to.)

It is well-known that there is an analytic covering map $\exp : LA(\mathbb{C}) \rightarrow A(\mathbb{C})$. $LA(\mathbb{C})$ is a complex vector space of the same dimension g as the abelian variety, and as such it is isomorphic to \mathbb{C}^g ; the exponential map is used to identify A with the quotient \mathbb{C}^g/Λ , where Λ is a lattice in \mathbb{C}^g . Hence the class of abelian varieties is contained in a larger class of Lie groups, those that are known as complex tori.

Definition 2.1. A *complex torus* is a compact complex Lie group that is obtained as a quotient \mathbb{C}^g/Λ where Λ is a discrete subgroup of rank $2g$.

In fact, abelian varieties are completely characterized as the complex tori which allow an embedding into a projective space (see Section A.5 of [HS00] for details.)

We're interested in finding intersections between the graph Γ_{\exp} of the exponential function and products of varieties of the form $L \times W$, where L is a translate of a linear subspace of LA and W is an algebraic subvariety of A .

2.2 Dense Subgroups of Abelian Varieties

The first property of pairs of varieties that we're interested in is freeness.

Definition 2.2. Let (L, W) be a pair with $L \subseteq LA(\mathbb{C})$ a translate of a linear space and $W \subseteq A(\mathbb{C})$ an algebraic variety.

We say L is *free* if it is not contained in a translate of LB for any abelian subvariety B of A .

We say W is *free* if it is not contained in a translate of B for any abelian subvariety B of A .

We say the pair (L, W) is *free* if both L and W are.

In this section we describe the linear subspaces of LA whose image through the exponential function is dense in A in the Euclidean topology, showing that this property holds for a generic subspace, even if the required property is in general stronger than freeness. This will allow us to apply a similar argument to the one used in Section 6 of [Kir19] to prove an existential closedness property for blurred complex exponentiation. This characterization is not new as it appears implicitly for example in Section 3 of [UY18], but we work out the precise statements.

For the rest of this subsection we use \bar{z} to denote the complex conjugate of the complex number z , rather than the fact that z is a tuple as in the rest of the article.

We endow \mathbb{C}^g with the usual Hermitian product, defined for $z = (z_1, \dots, z_g)$ and $w = (w_1, \dots, w_g)$ as $z \cdot w = z_1 \bar{w}_1 + \dots + z_g \bar{w}_g$. Consider, for complex vectors $z, z', \langle z, z' \rangle := \Re(z \cdot z')$.

Remark 2.3. Identify $z = (z_1, \dots, z_g) = (x_1 + iy_1, \dots, x_g + iy_g)$ with the real vector $(x_1, y_1, \dots, x_g, y_g) \in \mathbb{R}^{2g}$. Then we have that $\langle z, z' \rangle$ coincides with the real scalar product in \mathbb{R}^{2g} , as

$$\begin{aligned} \Re((x_1 + iy_1, \dots, x_g + iy_g) \cdot (x'_1 + iy'_1, \dots, x'_g + iy'_g)) &= \\ &= x_1 x'_1 + y_1 y'_1 + \dots + x_g x'_g + y_g y'_g. \end{aligned}$$

We use Remark 2.3 to define the dual of a lattice.

Definition 2.4. Let Λ be a lattice of rank $2g$ in \mathbb{C}^g . The *dual lattice* Λ^* is the lattice defined by

$$\Lambda^* := \{\theta \in \mathbb{C}^g \mid \langle \theta, \lambda \rangle \in \mathbb{Z} \ \forall \lambda \in \Lambda\}.$$

We use the dual lattice to describe the real hyperplanes of \mathbb{C}^g whose image in \mathbb{C}^g/Λ is closed.

Lemma 2.5 (See Section 3 of [UY18]). *Let H be a real hyperplane in \mathbb{C}^g . Then $H + \Lambda$ is closed in \mathbb{C}^g/Λ if and only if H can be defined by an equation of the form $\Re(\sum_{i=1}^g \bar{\theta}_i z_i) = 0$ for $\theta = (\theta_1, \dots, \theta_g) \in \Lambda^*$.*

Proof. As a real torus, \mathbb{C}^g/Λ is isomorphic to $\mathbb{R}^{2g}/\mathbb{Z}^{2g}$. It is well-known that the hyperplanes H of \mathbb{R}^{2g} which have closed image in the quotient are those defined by a \mathbb{Q} -linear equation, as these have a set of generators in \mathbb{Z}^{2g} ; in other words, $H \cap \mathbb{Z}^{2g}$ is a lattice in H . Hence, in the setting of \mathbb{C}^g/Λ we must characterize the spaces H for which $H \cap \Lambda$ is a lattice in H .

So suppose $H \cap \Lambda$ is a lattice in H . Then H contains $2g - 1$ linearly independent elements of Λ , and there is an element of Λ^* which is orthogonal to each of those for the real scalar product. Then by Remark 2.3 we have the claim.

Conversely, suppose H is defined by an equation of the form above. Then H is orthogonal to a space defined by an element of the dual, and so it is generated by points in the lattice. \square

This property easily translates to complex hyperplanes.

Corollary 2.6. *Suppose L is a complex hyperplane of \mathbb{C}^g which can't be defined by an equation of the form $\sum_{i=1}^g \bar{\theta}_i z_i = 0$ for $\theta = (\theta_1, \dots, \theta_g) \in \Lambda^*$ (we say such an hyperplane is not defined over the conjugate of the dual lattice). Then $L + \Lambda$ is dense in \mathbb{C}^g .*

Proof. If a complex linear space satisfies $\Re(\sum_{i=1}^g \bar{\theta}_i z_i) = 0$ for coefficients in Λ^* , then it must satisfy $\Im(\sum_{i=1}^g \bar{\theta}_i z_i) = 0$, and hence the full equation $\sum_{i=1}^g \bar{\theta}_i z_i = 0$. Hence if a complex linear hyperplane does not have dense exponential, it is defined over the conjugate of the dual lattice. \square

Thus it is trivial to obtain the following lemma, which clears the picture for general complex linear spaces.

Lemma 2.7. *Let L be a complex linear subspace of \mathbb{C}^g . Then $L + \Lambda$ is dense in \mathbb{C}^g if and only if L is not contained in a complex hyperplane defined over the conjugate of Λ^* .*

We provide explicit computations in the case of powers of an elliptic curve $E \cong \mathbb{C}/\Lambda$, where Λ is a lattice of the form $\mathbb{Z} + \tau\mathbb{Z}$ for some complex number $\tau = a + ib$.

Identifying \mathbb{C} with \mathbb{R}^2 , the lattice Λ is generated by the points $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} a \\ b \end{pmatrix}$. Clearly then a pair of generators of Λ^* is given by $\begin{pmatrix} 0 \\ \frac{1}{b} \end{pmatrix}$ and $\begin{pmatrix} 1 \\ -\frac{a}{b} \end{pmatrix}$. These correspond to the complex numbers $-\frac{i}{b}$ and $1 + \frac{ia}{b}$. Note that, multiplying both for ib , we obtain 1 and $-a + ib = -\bar{\tau}$, which generate the conjugate of Λ . Hence, in the case of powers of an elliptic curve, the conjugate of the dual lattice is actually the original lattice up to multiplication by a scalar, and Lemma 2.7 takes the following form.

Corollary 2.8. *Let $E \cong \mathbb{C}/\Lambda$ be an elliptic curve, $L \leq \mathbb{C}^g$ a complex linear space, \exp the exponential map of the abelian variety E^g . Then $\exp(L)$ is dense in E^g if and only if L is not contained in an hyperplane defined over Λ .*

Recall that an elliptic curve E is said to have *Complex Multiplication* when its endomorphism ring is strictly larger than \mathbb{Z} , and that these are characterized as the curves whose period lattice has the form $\mathbb{Z} + \tau\mathbb{Z}$ for τ an imaginary quadratic number (see [HS00], Example A.5.1.3). Corollary 2.8 has a particularly nice interpretation in this case. In fact, if E has CM, linear spaces defined over the lattice are Lie algebras of abelian subvarieties of E^g , and the corollary can be seen as stating that all free complex hyperplanes have dense exponentials.

Remark 2.9. A similar description can be given by the well-known characterization of complex abelian varieties as complex tori which admit a *polarization*, that is, an Hermitian form on \mathbb{C}^g whose imaginary part takes integer values on $\Lambda \times \Lambda$. We chose to follow [UY18] in using the dual lattice as it provides a more concrete description.

2.3 Intersections

In this section L will always be a translate of a linear subspace of $LA(\mathbb{C})$.

Density of $\exp(L)$ is not enough to prove the existence of intersections with W ; we need to eliminate possible parallelism phenomena. This is done via the notion of rotundity. This definition is adapted from the one given in [Kir09].

Definition 2.10. Let (L, W) be a pair of varieties in $LA \times A$. We say the pair (L, W) is *rotund* if for every abelian subvariety B and quotient map $f : A \rightarrow A/B$,

$$\dim Tf(L \times W) = \dim(Lf(L)) + \dim f(W) \geq \dim A/B.$$

Note that in particular this implies that $\dim L + \dim W \geq \dim A$, by considering the trivial subgroup of A . If A is simple, that is, it has no proper non-trivial abelian subvariety, this is actually the only condition.

The fibre dimension theorem prevents the parallelism phenomena we were referring to for rotund pairs, as seen in the next proposition. The next two statements have essentially the same proofs as Lemma 6.1 and Proposition 6.2 from [Kir19], translated to the context of abelian varieties.

Proposition 2.11. *Let (L, W) be an irreducible rotund pair in $LA \times A$ such that $\dim L + \dim W = \dim A$, and B an abelian subvariety of A . Then there is a non-empty Zariski-open subset V_B of $L \times W$ such that if $\gamma \in V_B$,*

$$\dim((L \times W) \cap (\gamma \oplus TB)) \leq \dim B.$$

Proof. Let $f : A \rightarrow A/B$ be a quotient map, and $Tf : TA \rightarrow T(A/B)$ the tangent map. Consider the restriction of Tf to $L \times W$, so that for every $\gamma \in L \times W$, $((\gamma \oplus TB) \cap (L \times W))$ is a fibre of Tf . Then the fibre dimension theorem implies that there is a Zariski-open subset V_B of L , such that for all $\gamma \in V_B$,

$$\begin{aligned} \dim((L \times W) \cap (\gamma \oplus TB)) &= \dim Tf^{-1}(\gamma) = \\ &= \dim L + \dim W - \dim(Tf(L \times W)) \leq \\ &\leq \dim L + \dim W - \dim A/B = \\ &= \dim A - \dim A + \dim B = \dim B \end{aligned}$$

where we have used rotundity in the inequality between the second and third line. \square

Theorem 2.12. *Let (L, W) be a free rotund pair in $LA \times A$, and suppose L is not contained in any hyperplane defined over the conjugate of the dual lattice. Then $\exp(L) \cap W$ is dense in W in the Euclidean topology.*

Proof. By translating W if necessary, we may assume without loss of generality that L is a linear space. By intersecting with a generic hyperplane if necessary, we assume that $\dim L + \dim W = \dim A$.

Consider the function $\theta : L \times W \rightarrow A$, mapping (\bar{l}, \bar{w}) to $\bar{w} \ominus \exp(\bar{l})$. Let $\bar{a} = (\bar{l}_0, \bar{w}_0)$ be a point in $L \times W$. Consider a small open neighbourhood U of \bar{a} in $L \times W$ in the Euclidean topology, and the restriction of θ to U , denoted $\theta|_U$.

The set $\theta|_U^{-1}(\theta(\bar{a}))$ is analytic, and thus we can assume it is connected by shrinking U if necessary; if it has dimension 0, then $\theta|_U$ is locally finite-to-one and therefore open, by the Remmert open mapping theorem. So, under the assumption that $\dim \theta|_U^{-1}(\theta(\bar{a})) = 0$, $\theta(U)$ is an open subset of A , and as $\exp(L)$ is dense in A , $\theta(U) \cap \exp(L) \neq \emptyset$. But then there is a point $(\bar{l}, \bar{w}) \in U$ such that $\bar{w} \ominus \exp(\bar{l}) = \exp(\bar{l}')$, and so $\bar{w} = \exp(\bar{l} + \bar{l}') \in \exp(L)$.

Therefore it remains to show that $\theta|_U^{-1}(\theta(\bar{a}))$ can be taken to have dimension 0 for almost every $\bar{a} \in L \times W$. Suppose then that for some \bar{a} , the set

$$S := \{(\bar{l}, \bar{w}) \in L \times W \mid \theta(\bar{l}, \bar{w}) = \theta(\bar{a})\}$$

has positive dimension.

For any $\bar{t} \in LA$ such that $\exp(\bar{t}) = \theta(\bar{a})$, consider the set

$$S' := \{(\bar{l} + \bar{t}, \bar{w}) \in LA \times A \mid (\bar{l}, \bar{w}) \in S\}.$$

For any point $\bar{b} \in S'$, we have

$$\theta(\bar{b}) = \bar{w} \ominus (\exp(\bar{l}) \oplus \exp(\bar{t})) = \theta(\bar{a}) \ominus \theta(\bar{a}) = 0_A$$

so S' is a positive dimensional component of the intersection

$$\bar{t} + L \times W \cap \Gamma_{\exp}$$

where Γ_{\exp} denotes the graph of \exp .

Recall that such a component is said to be *atypical* when $\dim S > \dim L + \dim W - \dim A$. In this case, since $\dim L + \dim W = \dim A$ by assumption, every positive dimensional component is atypical.

Consider the family of pairs of varieties $\mathcal{V} := \{(\bar{s} + L, W) \mid s \in LA(\mathbb{C})\}$. By Theorem 8.1 in [Kir06], there is a finite set \mathcal{B} of proper abelian subvarieties of A such that for every pair $(\bar{s} + L, W) \in \mathcal{V}$, every atypical component of the intersection $(\bar{s} + L \times W) \cap \Gamma_{\exp}$ is contained in a translate of TB for some $B \in \mathcal{B}$. Hence there are $B \in \mathcal{B}$ and $\gamma \in TA(\mathbb{C})$ such that $S' \subseteq \gamma \oplus TB$, and therefore

$$S \subseteq (\gamma \ominus (\bar{t}, 0_A)) \oplus TB.$$

Note that since $\bar{a} \in S$, this can be rewritten as

$$S \subseteq \bar{a} \oplus TB.$$

Such a B can be chosen to be minimal, and by the contrapositive of the same theorem we must have that

$$\dim S = \dim(\bar{a} \oplus TB \cap L \times W) - \dim B.$$

Since $\dim S$ is positive, $\bar{a} \in V_B$ (in the notation of Proposition 2.11). Then, to avoid that $\dim S > 0$, we just need to take $\bar{a} \in V_B$ for every abelian subvariety $B \in \mathcal{B}$.

We have then shown that there is a Zariski-open subset O (the intersection of the V_B 's) of $L \times W$ such that if $\bar{a} \in O$ then for an arbitrarily small neighbourhood U of \bar{a} in $L \times W$ there is a point $(\bar{l}, \bar{w}) \in U$ such that $\bar{w} \in \exp(L)$, as desired. \square

As pointed out in the introduction, this method relies on density of $\exp(L)$ to work, and it doesn't give a definite answer for linear spaces L such that the closure of $\exp(L)$ is a real subtorus of the abelian variety A .

Using Corollary 2.8, we can give a very explicit interpretation of this result in the case in which A is a power of an elliptic curve.

Corollary 2.13. *Let (L, W) be a free rotund pair in $LE^g \times E^g$ such that $\dim L + \dim W = \dim A$, and L is not defined over $\text{End}(E) \otimes \mathbb{R}$. Then $\exp(L) \cap W$ is dense in W in the Euclidean topology.*

3 The j Function

3.1 Background and Notation

Let \mathbb{H} denote the complex upper half plane, i.e. the set $\{z \in \mathbb{C} \mid \Im(z) > 0\}$. It is well-known that the special linear group $\mathrm{SL}_2(\mathbb{R})$ acts on \mathbb{H} by Möbius transformations, that is, the matrix $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ acts on $z \in \mathbb{H}$ by mapping it to $\frac{az+b}{cz+d}$.

Remark 3.1. This action can actually be seen as a restriction of a “bigger” action by Möbius transformations, that of $\mathrm{GL}_2(\mathbb{C})$ on the Riemann sphere $\mathbb{P}^1(\mathbb{C})$. This has two consequences that are of interest for us.

The first one is that we can consider $\mathrm{SL}_2(\mathbb{R})$ acting not just on \mathbb{H} , but also on $\mathbb{R} \cup \{\infty\}$. This will be needed to give the definition of cusp.

The second one is that, while the action can be defined for any matrix in $\mathrm{GL}_2(\mathbb{C})$, it is clearly invariant by scalar multiplication, meaning that given $\lambda \in \mathbb{C}$ and $h \in \mathrm{GL}_2(\mathbb{C})$, $hz = \lambda hz$ for all $z \in \mathbb{H}$; hence this can be seen as an action of $\mathrm{SL}_2(\mathbb{C})$. The situation is not exactly the same for real matrices: let $\mathrm{GL}_2(\mathbb{R})^+ := \{h \in \mathrm{GL}_2(\mathbb{R}) \mid \det(h) > 0\}$. The action of $\mathrm{SL}_2(\mathbb{R})$ can be seen as the restriction of the action on the same space of $\mathrm{GL}_2(\mathbb{R})^+$. We’ll be especially concerned with matrices which up to scalar multiplication have rational entries, as these define the so-called modular relations on \mathbb{H} , the only algebraic relations which are preserved by the j function. We will freely abuse notation and consider the action of $\mathrm{GL}_2(\mathbb{Q})^+$ on \mathbb{H} , claiming that a matrix $h \in \mathrm{SL}_2(\mathbb{R})$ is in $\mathrm{GL}_2(\mathbb{Q})^+$ when actually what we mean is that it is a scalar multiple of such a matrix.

We introduce the notion of cusp.

Definition 3.2. Let Γ be a discrete subgroup of $\mathrm{SL}_2(\mathbb{R})$. A *cusp* for Γ is a point $x \in \mathbb{R} \cup \{\infty\}$ such that $\gamma x = x$ for every $\gamma \in \Gamma$ of the form $\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}$ for some $t \in \mathbb{R}$ (such points are called *parabolic*.)

For the rest of this subsection, let $G = \mathrm{SL}_2(\mathbb{R})$ and $\Gamma = \mathrm{SL}_2(\mathbb{Z})$.

Recall the following definition of commensurability of subgroups and the characterization for commensurability of conjugates of Γ in G .

Definition 3.3. Two subgroups H and K of a group are *commensurable* if $H \cap K$ has finite index in both H and K .

Fact 3.4 ([Mil97], Lemmas 5.29 and 5.30). Let $g \in G$. Then $g\Gamma g^{-1}$ and Γ are commensurable if and only if $g \in \mathrm{GL}_2(\mathbb{Q})^+$.

We recall the definition of the j function. To do this, we first give the general definition of modular function as presented in [Mil97], Part I, Section 4, and then specialize to the case of j .

Definition 3.5. We say a function $f : \mathbb{H} \rightarrow \mathbb{C}$ is Γ -invariant if for any $\gamma \in \Gamma$ and $z \in \mathbb{H}$, $f(\gamma z) = f(z)$.

According to Definition 3.2, the only cusp for Γ is ∞ . As the Γ -orbit of ∞ can easily be checked to be \mathbb{Q} , studying the behaviour of a Γ -invariant function at infinity is the same as studying it at any rational point.

Definition 3.6. Let f be a Γ -invariant function on \mathbb{H} , \mathbb{D} denote the open unit disk. Then we can define $f^* : \mathbb{D} \setminus \{0\} \rightarrow \mathbb{C}$ as $f^*(z) = f(e^{2\pi iz})$. We say f is meromorphic (resp. has a pole of order n) at the cusp if f^* is meromorphic (resp. has a pole of order n) at 0.

Now we have everything that is needed to define the j -function.

Definition 3.7. The modular invariant j is the unique function $j : \mathbb{H} \rightarrow \mathbb{C}$ that is holomorphic, Γ -invariant, with a simple pole at the cusp and such that $j(i) = 1728$ and $j(e^{\frac{i\pi}{3}}) = 0$.

It is well-known that j establishes an analytic isomorphism between $\Gamma \backslash \mathbb{H}$ and \mathbb{C} . We will be studying intersections between images through j of certain algebraic subvarieties of \mathbb{H}^n and algebraic subvarieties of \mathbb{C}^n .

By algebraic subvariety of \mathbb{H}^n we mean the intersection with \mathbb{H}^n of an algebraic subvariety of \mathbb{C}^n .

The varieties we will mostly be concerned with are Möbius varieties.

Definition 3.8. Let L be an algebraic subvariety of \mathbb{H}^n . We say L is a *Möbius subvariety* if L can be defined just by using conditions of the form $z_k = gz_i$ for $g \in G, i, k \leq n$ and $z_i = \tau$ for $\tau \in \mathbb{H}, i \leq n$.

So for example the variety $\{(z_1, z_2) \in \mathbb{H}^2 | z_2 = z_1 + \sqrt{2}\}$ is a Möbius variety, defined by the matrix $\begin{pmatrix} 1 & \sqrt{2} \\ 0 & 1 \end{pmatrix}$.

Weakly special varieties are Möbius varieties of particular interest, as they are an example of bialgebraic varieties in the sense of [KUY16].

Definition 3.9. A Möbius variety L is *weakly special* if it can be defined by using just conditions specifying constant coordinates and equations of the form $z_i = hz_k$ where $h \in \text{GL}_2(\mathbb{Q})^+$.

As the name suggest, this is a generalization of the notion of *special* variety, which is a weakly special variety in which all constant coordinates are special points, i.e. imaginary quadratic elements of \mathbb{H} : these are the only algebraic numbers in \mathbb{H} for whose image through the j function is also algebraic.

Fact 3.10 ([UY11], Theorem 1.2). Let V be an algebraic subvariety of \mathbb{H}^n . Then $j(V)$ is an algebraic subvariety of \mathbb{C}^n if and only if V is weakly special.

As is well-known, the algebraic relations in the codomain are defined by the modular polynomials.

Definition 3.11. For each natural number N , the N -th *modular polynomial* is the polynomial $\Phi_N \in \mathbb{Z}[X, Y]$ such that for all $w_1, w_2 \in \mathbb{C}$, $\Phi_N(w_1, w_2) = 0$ holds if and only if there are $z \in \mathbb{H}$ and $g \in \text{GL}_2(\mathbb{Q})^+$ with coprime integer entries and determinant N such that $w_1 = j(z)$, $w_2 = j(gz)$.

Therefore if a Möbius variety L is (contained in a) weakly special variety, we understand the behaviour of $j(L)$, and in particular its intersections with algebraic subvarieties in \mathbb{C}^n . We want to study what happens when this is not the case.

3.2 Density of Images of Möbius Subvarieties

As outlined above, we're interested in Möbius varieties that are not contained in weakly special varieties, as the image through j of a weakly special variety is an algebraic variety. This suggests an analogy with the case of abelian varieties, where the varieties for which algebraicity is preserved under \exp are the Lie algebras of abelian subvarieties. Hence, the natural definition of freeness in the context of the j function is the following.

Definition 3.12. A Möbius variety $L \subseteq \mathbb{H}^n$ is *free* if it is not contained in any weakly special variety, i.e., if no coordinate is constant on L and for no $i, k \leq n$ there is $h \in \mathrm{GL}_2(\mathbb{Q})^+$ such that $z_k = h z_i$ for all $\bar{z} \in L$.

The goal of this subsection is to establish the following.

Lemma 3.13. *Let L be a free Möbius variety. Then $j(L)$ is dense in \mathbb{C}^n .*

Lemma 3.13 is an application of Ratner's theorem on unipotent flows, first proved by Ratner in [Rat91]. We'll provide the details for completeness; the reader who is interested in this topic is referred to the exposition in [Mor05].

Recall that on a Lie group we can always define a left-invariant measure μ , the *Haar measure*, which is unique up to multiplication by scalars. Among other things, this is used to define lattices.

Definition 3.14. Let Γ be a subgroup of a Lie group G . A *fundamental domain* for Γ is a measurable subset \mathcal{F} of G such that $\Gamma\mathcal{F} = G$ and that $\gamma\mathcal{F} \cap \mathcal{F}$ has measure zero for all non-identity $\gamma \in \Gamma$.

Definition 3.15. A subgroup Γ of a Lie group G is a *lattice* in G if it is discrete and it has a fundamental domain of finite measure.

There is an important link between the notions of lattice and commensurability which will be needed later.

Proposition 3.16. *Let Γ and Γ' be lattices in G , Δ be the diagonal of G^2 , i.e. the group $\{(g, g) \in G^2 | g \in G\}$. Then the following are equivalent:*

1. $\Gamma \cap \Gamma'$ is a lattice in G ;
2. $(\Gamma \times \Gamma') \cap \Delta$ is a lattice in Δ ;
3. Γ and Γ' are commensurable.

Proof. That 1 and 2 are equivalent is clear once we observe that $(\Gamma \times \Gamma') \cap \Delta$ is the set $\{(\gamma, \gamma) \in \Delta \mid \gamma \in \Gamma \cap \Gamma'\}$.

For the equivalence of 1 and 3 we have the more general property that a subgroup L' of a lattice L is a lattice if and only if $[L : L']$ is finite. To see this, observe that a fundamental domain for L' is given by the union of $[L : L']$ fundamental domains for L ; and hence it has finite measure if and only if $[L : L']$ is finite. \square

We can now state (a form of) Ratner's theorem. The following statement is the combination of Theorem 1.1.14, Remark 1.1.15 and Remark 1.1.19 from [Mor05].

Theorem 3.17 (Ratner's Orbit Closure Theorem). *Let G be a Lie group, Γ a lattice in G , H a subgroup generated by unipotent elements.*

Then for every $x \in G$, there is a closed subgroup S of G with the following properties:

- a. $H \subseteq S$;
- b. Every connected component of S contains an element of H ;
- c. $x^{-1}\Gamma x \cap S$ is a lattice in S ;
- d. The double coset $\Gamma x H$ is dense in $\Gamma x S$.

We will first apply the theorem to the case of $G = \mathrm{SL}_2(\mathbb{R})$ and $\Gamma = H = \mathrm{SL}_2(\mathbb{Z})$, and then use this to extend this to powers of $\mathrm{SL}_2(\mathbb{R})$.

Lemma 3.18 (See Exercise 1.27 in [Mor05]). *Let $g \in G$. Then the double coset $\Gamma g \Gamma$ is discrete if and only if $g \in \mathrm{GL}_2(\mathbb{Q})^+$, and it is dense otherwise.*

Proof. First note that Γ is generated by the matrices $s = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $t = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, which are unipotent. Hence we consider the action of Γ on $\Gamma \backslash G$ by right multiplication. Fix $g \in G$; by Ratner's theorem there is a subgroup S satisfying the properties listed in Theorem 3.17. Consider the connected component of the identity S° ; this is a normal subgroup of S and, because $\Gamma \subseteq S$, S° is normalized by Γ .

Since G is a simple Lie group (it doesn't have non-trivial connected normal subgroups) and Γ is Zariski-dense in G as a consequence of the Borel density theorem ([Mor05], Theorem 4.7.1), the only connected subgroups of G that are normalized by Γ are the trivial subgroup and G . Suppose $S^\circ = \{\mathbb{I}_2\}$; then, because every connected component of S contains an element of Γ , $S = \Gamma$. But then $g^{-1}\Gamma g \cap \Gamma$ is a lattice in Γ , which amounts to say that $g^{-1}\Gamma g$ and Γ are commensurable by Proposition 3.16. By Fact 3.4 this is the case if and only if $g \in \mathrm{GL}_2(\mathbb{Q})^+$, so we must have $S = G$; then $\Gamma g \Gamma$ is dense in $\Gamma g G = G$. \square

We now describe the full setting to which we want to apply the theorem. Fix $n \geq 2$ and let $G = \mathrm{SL}_2(\mathbb{R})^{n-1}$ and $\Gamma = \mathrm{SL}_2(\mathbb{Z})^{n-1}$. Consider now the subgroup $H = \Delta(\Gamma)$, where by Δ we mean the diagonal, i.e. the subgroup $\{(\gamma_2, \dots, \gamma_n) \in \Gamma \mid \gamma_2 = \dots = \gamma_n\}$. (It will hopefully become clear later why we start the enumeration at 2.)

We can see H as a subgroup generated by unipotent elements by identifying G with the subgroup of $\mathrm{SL}_{2n-2}(\mathbb{R})$ of all matrices of the form

$$\begin{pmatrix} k_2 & \mathbb{O}_2 & \cdots & \mathbb{O}_2 \\ \mathbb{O}_2 & k_3 & \cdots & \mathbb{O}_2 \\ \vdots & \ddots & \cdots & \vdots \\ \mathbb{O}_2 & \cdots & \cdots & k_n \end{pmatrix}$$

with each $k_n \in \mathrm{SL}_2(\mathbb{R})$, where each \mathbb{O}_2 denotes a 2 by 2 block of zeros, and Γ and H with the corresponding subgroups. Then, as $\mathrm{SL}_2(\mathbb{Z})$ is generated by the unipotent elements $s = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $t = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$, H is generated by the matrices

$$\begin{pmatrix} s & \mathbb{O}_2 & \cdots & \mathbb{O}_2 \\ \mathbb{O}_2 & s & \cdots & \mathbb{O}_2 \\ \vdots & \ddots & \cdots & \vdots \\ \mathbb{O}_2 & \cdots & \cdots & s \end{pmatrix} \text{ and } \begin{pmatrix} t & \mathbb{O}_2 & \cdots & \mathbb{O}_2 \\ \mathbb{O}_2 & t & \cdots & \mathbb{O}_2 \\ \vdots & \ddots & \cdots & \vdots \\ \mathbb{O}_2 & \cdots & \cdots & t \end{pmatrix}$$

and so it makes sense to apply the theorem on cosets of groups generated by unipotent elements.

While representing Γ, G and H this way fits better with Ratner's theorem, we'll stick with the previous definitions because they are equivalent and easier to work with.

For the general case we're going to need two group theoretic facts: Goursat's lemma and a proposition on normal subgroups of powers of simple Lie groups.

Fact 3.19 (Goursat's Lemma, [Lan04], Exercise 1.5). Let G_1 and G_2 be groups, T a subgroup of $G_1 \times G_2$ such that the projections $\pi_i : T \rightarrow G_i$ are surjective for $i = 1, 2$. Identify the kernel N_1 of π_1 with a normal subgroup of G_2 , and the kernel N_2 of π_2 with a normal subgroup of G_1 .

Then the image of T in $G_1/N_2 \times G_2/N_1$ is the graph of an isomorphism.

We want to study subgroups of $\mathrm{SL}_2(\mathbb{R})^{n-1}$, in order to understand which subgroups are associated to elements of the group by Ratner's theorem. First we remark that closed normal subgroups are products of trivial subgroups and copies of the group. Then we move on to study general connected subgroups, and see that graphs of inner automorphisms are the only additional possibility.

Proposition 3.20. *Let G be a simple Lie group, $N \trianglelefteq G^n$ a normal connected subgroup. Then N is a product of trivial groups and copies of G .*

Proof. Observe that if the k -th coordinate of an element is not the identity, then it can be moved to any element of the group by conjugating by an element that is the identity of all the coordinates except the k -th one. \square

This suggests that in order to move forward we need an understanding of the group of automorphisms of $\mathrm{SL}_2(\mathbb{R})$. This has a very clear description as follows.

Proposition 3.21. *The Lie group automorphisms of $\mathrm{SL}_2(\mathbb{R})$ are exactly the automorphisms of the form $g \mapsto hgh^{-1}$ for matrices $h \in \mathrm{GL}_2(\mathbb{R})$.*

Proof. It is clear that every conjugation induces an isomorphism. To see that this is actually all of them we appeal to Section XV in Chapter IV of [Che46], where it is shown that the group of automorphisms of a Lie group embeds in the group of automorphisms of its Lie algebras, and Theorem 5 in Chapter 9 of [Jac], where it is proved that all automorphisms of the Lie algebra of $\mathrm{SL}_2(\mathbb{R})$ are induced by conjugation by an element of $\mathrm{GL}_2(\mathbb{R})$. \square

Proposition 3.22. *Let S be a closed connected subgroup of $G = \mathrm{SL}_2(\mathbb{R})^{n-1}$ such that all the projections $\pi_i : S \rightarrow \mathrm{SL}_2(\mathbb{R})$ are surjective. Then if $S \neq G$ there are $i, k \leq n - 1$ and an element $h \in \mathrm{GL}_2(\mathbb{R})$ such that for all $\bar{g} \in G$, $g_k = hg_i h^{-1}$.*

Proof. We do this by induction on n , with the base case $n = 3$.

So suppose $S \leq \mathrm{SL}_2(\mathbb{R})^2$ is a proper Lie subgroup with surjective projections. Then by Goursat's lemma there are normal subgroups (the kernels of the projections) N_1 and N_2 of $\mathrm{SL}_2(\mathbb{R})$ such that the image of S in $\mathrm{SL}_2(\mathbb{R})/N_2 \times \mathrm{SL}_2(\mathbb{R})/N_1$ is the graph of an isomorphism. If the kernels N_1 and N_2 are trivial, then S is the graph of an automorphism of $\mathrm{SL}_2(\mathbb{R})$. But since S is closed, it is a Lie subgroup, and hence it must be the graph of a continuous automorphism; but by Proposition 3.21, there are given by conjugation by elements of $\mathrm{GL}_2(\mathbb{R})$. The only other non-trivial normal subgroup of $\mathrm{SL}_2(\mathbb{R})$ is the group $\{\mathbb{I}_2, -\mathbb{I}_2\}$, and in that case S would be given by the union of the graph of an automorphism φ and the set of elements of the form $(g, -\varphi(g))$. However, this would contradict the assumption that S is connected, so we are done.

Now suppose $n > 3$, and the result holds for $n - 1$. If S is a subgroup of G , see it as a subgroup of $\mathrm{SL}_2(\mathbb{R})^{n-2} \times \mathrm{SL}_2(\mathbb{R})$. If the projection of S on $\mathrm{SL}_2(\mathbb{R})^{n-2}$ is not surjective, then it is a proper subgroup and we can conclude by the inductive hypothesis, so suppose it is surjective. Note that in this case any two of the first $n - 2$ coordinates are not the graph of an automorphism.

We can now apply Goursat's lemma to S as a subgroup of $\mathrm{SL}_2(\mathbb{R})^{n-2} \times \mathrm{SL}_2(\mathbb{R})$, since both projections are surjective, and find normal subgroups N_1 and N_2 such that the image of S in $\mathrm{SL}_2(\mathbb{R})^{n-2}/N_2 \times \mathrm{SL}_2(\mathbb{R})/N_1$ is the graph of an isomorphism; again, N_1 must be trivial (for the same reasons why it had to be trivial in the base case: if it is $\mathrm{SL}_2(\mathbb{R})$ then the projection is not surjective, and if it is $\{\mathbb{I}_2, -\mathbb{I}_2\}$ then the resulting subgroup is not connected) and N_2 is a proper subgroup of $\mathrm{SL}_2(\mathbb{R})^{n-2}$. We focus on N_2 : this is a normal subgroup of $\mathrm{SL}_2(\mathbb{R})^{n-2}$ such that the quotient $\mathrm{SL}_2(\mathbb{R})^{n-2}/N_2$ is isomorphic to $\mathrm{SL}_2(\mathbb{R})$, so by Proposition 3.20, and the assumption that S is connected, it must be a product of $n - 3$ copies of $\mathrm{SL}_2(\mathbb{R})$ and a trivial group, say at coordinate k . Then $S = \{\bar{g} \in \mathrm{SL}_2(\mathbb{R})^{n-1} | g_{n-1} = hg_k h^{-1}\}$ for some $h \in \mathrm{GL}_2(\mathbb{R})$. \square

In the proof of Lemma 3.18 we used the fact that connected subgroups of $\mathrm{SL}_2(\mathbb{R})$ that are normalized by Γ are indeed normal, so they can't be non-trivial as $\mathrm{SL}_2(\mathbb{R})$ is a simple Lie group. The corresponding statement for powers of $\mathrm{SL}_2(\mathbb{R})$ and the diagonal action we are considering is then the following.

Proposition 3.23. *Let S be a closed connected subgroup of $\mathrm{SL}_2(\mathbb{R})^n$, and suppose S is normalised by H . Then either S is normal, or there are $i, k \leq n$ such that $g_i = g_k$ for every $\bar{g} \in S$.*

Proof. Let S be normalised by H , and suppose it is not normal. By restricting to a subgroup of $\mathrm{SL}_2(\mathbb{R})^m$ for some $m < n$ if necessary we can assume without loss of generality that the projections are surjective. Then by Proposition 3.22 there are $i, k \leq n$ and $h \in \mathrm{GL}_2(\mathbb{R})$ such that $g_k = hg_i h^{-1}$ for every $\bar{g} \in S$. Hence it's enough to prove this when $n = 2$, and the rest will follow inductively.

So suppose S is a closed connected subgroup of $\mathrm{SL}_2(\mathbb{R})^2$; in this case, H is the diagonal of $\mathrm{SL}_2(\mathbb{Z})^2$, and S is the graph of conjugation by some matrix $h \in \mathrm{GL}_2(\mathbb{R})$. Hence,

$$S = \{(g, hgh^{-1}) \in \mathrm{SL}_2(\mathbb{R})^2 \mid g \in \mathrm{SL}_2(\mathbb{R})\}.$$

Since S is normalized by H , for every $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ and $g \in \mathrm{SL}_2(\mathbb{R})$ the point

$$(\gamma g \gamma^{-1}, \gamma h g h^{-1} \gamma^{-1})$$

belongs to S . However, as S is the graph of conjugation by H , it contains exactly one point whose first coordinate is $\gamma g \gamma^{-1}$, i.e.

$$(\gamma g \gamma^{-1}, h \gamma g \gamma^{-1} h^{-1}).$$

As this holds for all g and all γ , we can conclude that h commutes with all $\gamma \in \Gamma$. But then, by Zariski density of $\mathrm{SL}_2(\mathbb{Z})$, it commutes with all of $\mathrm{SL}_2(\mathbb{R})$, and thus it is a scalar matrix; but then S is the diagonal of $\mathrm{SL}_2(\mathbb{R})$, as required. \square

We are now in a position to prove the following lemma.

Lemma 3.24. *Fix $(g_2, \dots, g_n) \in G$. If the Möbius subvariety $L \subseteq \mathbb{H}^n$ defined by the conditions $z_i = g_i z_1$ for each $i = 2, \dots, n$ is free, then the subgroup S associated to (g_2, \dots, g_n) by Ratner's theorem is G .*

Hence, with notations as above for Γ and H , $\Gamma(g_2, \dots, g_n)H$ is dense in G .

Proof. We do this by induction on $n \in \mathbb{N} \setminus \{0, 1\}$, so the base case is $n = 2$. In that case the variety is free if and only if g_2 is not a scalar multiple of a rational matrix, so we just need to apply Lemma 3.18.

Now assume $n > 2$, and suppose S is a proper subgroup of G . Consider the connected component of the identity, S° . This is a normal subgroup of S , and since S contains H each coordinate of S° is normalised by Γ . Hence the projections of S° are either $\mathrm{SL}_2(\mathbb{R})$ or the trivial group. If the projection of S° to the k -th component is trivial then, because every element of S contains an

element of H , $\pi_k(S) = \mathrm{SL}_2(\mathbb{Z})$. But then $g^{-1}\Gamma g \cap \Gamma$ is a lattice in Γ , and $g^{-1}\Gamma g$ and Γ are commensurable. Hence $g \in \mathrm{GL}_2(\mathbb{Q})^+$, contradicting the assumption that L was free.

Therefore we may assume that all the projections are surjective. By Proposition 3.22, then, there are two coordinates on which S is the graph of an automorphism. Therefore it suffices to prove the Lemma for $n = 3$, as any other case can be reduced to this one.

Suppose then that $n = 3$, so that $G = \mathrm{SL}_2(\mathbb{R})^2$ and $\Gamma = \mathrm{SL}_2(\mathbb{Z})^2$, and S is a proper subgroup of $G = \mathrm{SL}_2(\mathbb{R})^2$. Then S° must be the graph of the identity by Proposition 3.23; Since every connected component of S must contain an element of H , and $H \subseteq \Delta(\mathbb{R})$, $S = \Delta(\mathbb{R})$. By property *c.* in Ratner's theorem, $\bar{g}^{-1}\Gamma\bar{g} \cap \Delta(\mathbb{R})$ is a lattice in $\Delta(\mathbb{R})$: but by Proposition 3.16 this implies that $g_2^{-1}\Gamma g_2$ and $g_3^{-1}\Gamma g_3$ are commensurable. Hence, that Γ and $g_2 g_3^{-1} \Gamma g_3 g_2^{-1}$ are commensurable and therefore $g_2 g_3^{-1}$ is a scalar multiple of a rational matrix, contradicting the assumption that the Möbius variety L was free. Hence, if L is free, then $\Gamma(g_2, g_3)H$ is dense in $\Gamma(g_2, g_3)G = G$, as required. \square

We get density of $j(L)$ as a corollary of Lemma 3.24. Note that, since free Möbius varieties can be seen as products of one-dimensional Möbius varieties, we just need to take care of this case and the extension will be immediate as products of dense sets are dense.

Proof of Lemma 3.13. Suppose that

$$L = \{\bar{z} \in \mathbb{H}^n \mid z_i = g_i z_1 \text{ for } i = 2, \dots, n\}$$

and let U be an open subset in \mathbb{C}^n . Take the first projection $U_1 = \pi_1(U)$, and let $z \in \mathbb{H}$ be any point such that $j(z) \in U_1$. Because j is surjective and the action of $\mathrm{SL}_2(\mathbb{R})$ on \mathbb{H} is transitive and continuous, there is an open subset O of $\mathrm{SL}_2(\mathbb{R})^{n-1}$ such that $(j(z), j(h_2 z), \dots, j(h_n z)) \in U$ if and only if $(h_2, \dots, h_n) \in O$. O is open, so by Lemma 3.24 there are $\gamma, \gamma_2, \dots, \gamma_n$ such that $(\gamma_2 g \gamma, \dots, \gamma_n g \gamma) \in O$. Then

$$(j(z), j(\gamma_2 g \gamma z), \dots, j(\gamma_n g \gamma z)) = (j(\gamma z), j(g_2 \gamma z), \dots, j(g_n \gamma z)) \in j(L) \cap U.$$

\square

3.3 Intersections

We are now ready to prove the existence of intersections between images of free Möbius subvarieties and algebraic subvarieties of \mathbb{C}^n . The first easy, but crucial, remark concerns the set of matrices that join two points in the upper half plane via Möbius transformations.

Proposition 3.25. *Let $z_1, z_2 \in \mathbb{H}$, with $z_l = x_l + iy_l$ for $l = 1, 2$. Then for any $c, d \in \mathbb{R}$ such that $|cz_1 + d|^2 = \frac{y_1}{y_2}$, there are $a, b \in \mathbb{R}$ such that if g is the matrix*

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R}), \quad g z_1 = z_2.$$

Proof. Direct calculations show that $\begin{pmatrix} \sqrt{y} & \frac{x}{\sqrt{y}} \\ 0 & \frac{1}{\sqrt{y}} \end{pmatrix} i = x + iy$, for all x and y . Therefore, since the stabilizer of i is $\mathrm{SO}_2(\mathbb{R})$, we find that for all θ :

$$\begin{pmatrix} \sqrt{y_2} & \frac{x_2}{\sqrt{y_2}} \\ 0 & \frac{1}{\sqrt{y_2}} \end{pmatrix} \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{y_1}} & -\frac{x_1}{\sqrt{y_1}} \\ 0 & \sqrt{y_1} \end{pmatrix} z_1 = z_2$$

and that the lower entries of the product matrix are $-\frac{\cos \theta}{\sqrt{y_1 y_2}}$ and $x_1 \frac{\cos \theta}{\sqrt{y_1 y_2}} + (\sin \theta) \sqrt{\frac{y_1}{y_2}}$. Therefore,

$$cz_1 + d = \sqrt{\frac{y_1}{y_2}} (\sin \theta - i \cos \theta).$$

□

This together with density of images of Möbius varieties is enough to prove the existence of intersections in the case $\dim L = 1$.

Lemma 3.26. *Let (L, W) be pair in $\mathbb{H}^n \times \mathbb{C}^n$ such that L is a free Möbius variety of dimension 1, W is irreducible, and $\dim L + \dim W \geq n$. Then W has a dense subset of points of $j(L)$.*

We will obtain this as a corollary of a stronger result, which doesn't require W to be an algebraic variety.

Lemma 3.27. *Let $g_2, \dots, g_n \in \mathrm{SL}_2(\mathbb{R})$; V an open subset of \mathbb{C}^n , $f : V \rightarrow \mathbb{C}$ a holomorphic function. Denote by W the zero locus of f . Then W has a dense subset of points of the form $(j(z_1), j(g_2 z_1), \dots, j(g_n z_1))$.*

Proof. Let $L = \{(z_1, \dots, z_n) \mid z_i = g_i z_1 \text{ for } i = 2, \dots, n\}$.

If f is of the form $X_i - c$ for some $c \in \mathbb{C}$, without loss of generality we can assume $i = 1$, and then the result follows directly from density of $j(L)$: take z_1 to be any element such that $j(z_1) = c$, and if O is any open subset of \mathbb{C}^{n-1} , then we find $\gamma_2 g_2 \gamma, \dots, \gamma_n g_n \gamma$ such that

$$(j(\gamma_2 g_2 \gamma z_1), \dots, j(\gamma_n g_n \gamma z_1)) \in O.$$

Then the point $(j(\gamma z), j(g_2 \gamma z), \dots, j(g_n \gamma z))$ belongs to the open subset $\{c\} \times O$ of W . Hence from now on we assume W has no constant coordinates.

Let $\bar{w} \in W$ be a regular point such that no coordinate of \bar{w} is 0 or ∞ . Then find a point $\bar{z} \in \mathbb{H}$ such that $j(\bar{z}) = \bar{w}$, and find $h_2, \dots, h_n \in \mathrm{SL}_2(\mathbb{R})$ such that $z_i = h_i z_1$ for $i = 2, \dots, n$. Then consider the function $F : \mathbb{H} \rightarrow \mathbb{C}$, mapping z to $f(j(z), j(h_2 z), \dots, j(h_n z))$: by construction $F(z_1) = 0$, so consider a small compact neighbourhood U of z_1 .

By density, there is a sequence $\{\bar{g}^i\}_{i \in \mathbb{N}}$, converging to \bar{h} , such that each \bar{g}^i is a tuple of the form

$$(g_2^i, \dots, g_n^i) = (\gamma_2^i g_2 \gamma^i, \dots, \gamma_n^i g_n \gamma^i)$$

for some $\gamma^i, \gamma_2^i, \dots, \gamma_n^i \in \mathrm{SL}_2(\mathbb{Z})$, where g_2, \dots, g_n are the matrices defining L .

Then consider the sequence of functions $\{F_i\}_{i \in \mathbb{N}}$, where each F_i is defined on U as $F_i(z) = f(j(z), j(g_2^i z), \dots, j(g_n^i z))$. It is then clear that

$$\lim_{i \in \mathbb{N}} F_i(z_1) = F(z_1) = 0.$$

Now consider the derivative of F . Writing $j(\bar{h}z_1)$ for the tuple

$$(j(z_1), j(h_2 z_1), \dots, j(h_n z_1)),$$

we have that

$$\begin{aligned} & \frac{d}{dz} F(z_1) = \\ &= \frac{\partial f}{\partial Y_1}(j(\bar{h}z_1))j'(z_1) + \frac{\partial f}{\partial Y_2}(j(\bar{h}z_1))(j(h_2 z_1))' + \dots + \frac{\partial f}{\partial Y_n}(j(\bar{h}z_1))(j(h_n z_1))' \end{aligned}$$

where if $h_i = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $(j(h_i z_1))' = \frac{j'(h_i z_1)}{(cz_1 + d)^2}$.

It is clear that $\{\frac{d}{dz} F_i(z_1)\} \rightarrow_{i \in \mathbb{N}} \frac{d}{dz} F(z_1)$. Therefore if $\frac{d}{dz} F(z_1) \neq 0$, then for sufficiently large i the value of $F_i(z_1)$ is small with respect to its derivative. Furthermore, since each F_i is defined on a compact set, its second derivative is bounded on its domain. This implies that the conditions for the convergence of Newton's method are satisfied (see the result of Kantorovich, Theorem 5.3.1 in [DS96]), and therefore there is a zero of F_i close to z_1 .

Claim: Without loss of generality we can assume that $\frac{d}{dz} F(z_1) \neq 0$.

Proof of Claim: Suppose $\frac{d}{dz} F(z_1) = 0$. Because $\bar{w} = j(\bar{h}z_1)$ is a regular point in W and no coordinate of \bar{w} is 0 or 1728, the summands in $\frac{d}{dz} F(z_1)$ are not all zero. Then in particular at least two of them are not zero, and so there is $l > 1$ such that $\frac{\partial f}{\partial Y_l}(j(\bar{h}z_1)) \frac{j'(h_l z_1)}{(cz_1 + d)^2} \neq 0$. But then it's enough to change the matrix h_l , which we are free to do by Proposition 3.25: there is a matrix h'_l such that $h'_l z_1 = h_l z_1$, but $(j(h'_l z_1))' \neq (j(h_l z_1))'$. This proves the claim.

Then we are done: for sufficiently large i apply Newton's method and find z_0 close to z_1 such that $F_i(z_0) = 0$, i.e., such that

$$f(j(z_0), j(\gamma_2 g_2 \gamma z_0), \dots, j(\gamma_n g_n \gamma z_0)) = 0$$

for some $\gamma, \gamma_2, \dots, \gamma_n$. Then $(j(\gamma z_0), j(g_2 \gamma z_0), \dots, j(g_n \gamma z_0)) \in j(L) \cap W$. \square

Proof of Lemma 3.26. Apply Lemma 3.27 in the case where $V = \mathbb{C}^n$ and f is a polynomial. \square

To extend this to the general case we need the notions of broadness and freeness in the codomain, introduced by Aslayan in [Asl18].

Definition 3.28. Let (L, W) be a pair such that L is a Möbius subvariety of \mathbb{H}^n and W is an algebraic subvariety of \mathbb{C}^n .

For every $I = (i_1, \dots, i_k)$ where $1 \leq i_1 < \dots < i_k \leq n$ are natural numbers, let π_I denote the projection

$$\pi_I : \mathbb{H}^n \times \mathbb{C}^n \rightarrow \mathbb{H}^k \times \mathbb{C}^k$$

which maps $(z_1, \dots, z_n, w_1, \dots, w_n)$ to $(z_{i_1}, \dots, z_{i_k}, w_{i_1}, \dots, w_{i_k})$.

We say the pair (L, W) is *broad* if for every I ,

$$\dim \pi_I(L \times W) \geq k.$$

Definition 3.29. The algebraic variety $W \subseteq \mathbb{C}^n$ is *free* if no coordinate is constant and for no two coordinates $i, j \leq n$ and $N \in \mathbb{N}$, $\Phi_N(w_i, w_j)$ holds for all $w \in W$, where the Φ_N 's are the modular polynomials from Definition 3.11.

The pair (L, W) is *free* if both L and W are free.

The following technical lemma will be used to prove the main result.

Lemma 3.30. *Let $W \subseteq \mathbb{C}^n$ be an irreducible algebraic variety. There is a non-empty Zariski open subset W' of W such that for any Möbius variety $L \subseteq \mathbb{H}^n$ such that (L, W) is free broad, any intersection between $j(L)$ and W' is typical.*

Proof. Let I be any subset of $\{1, \dots, n\}$, and $\pi_I : \mathbb{C}^n \rightarrow \mathbb{C}^{n_I}$ denote the corresponding projection.

For any such projection there is a non-empty Zariski-open subset $W_I \subseteq W$ such that for every $\bar{w} \in W_I$, $\dim(\pi_I^{-1}(\pi_I(\bar{w}))) = \dim W - \dim \pi_I(W)$.

Moreover, by Aslanyan's weak modular Zilber-Pink theorem for parametric families (Theorem 7.8 in [Asl20]), if we consider the family

$$\{W_{\bar{w}_I} \mid \bar{w}_I \in \pi_I(W)\},$$

there are finitely many weakly special subvarieties with no constant coordinates of \mathbb{C}^n (and hence a Zariski-closed subset S_I of W) such that any maximal atypical component of one of the varieties $W_{\bar{w}_I}$ is contained in S_I . Thus, define

$$W' := W \setminus \bigcup_{I \subseteq [n]} (S_I \cup W_I).$$

Now let L be a Möbius variety such that (L, W) is free broad. Suppose U is a bounded open subset of L , that C is an irreducible component of the intersection $j(U) \cap W$, and that

$$\dim C > \dim L + \dim W - n \geq 0.$$

Now let I be the maximal subset of $\{1, \dots, n\}$ for which coordinates with the corresponding indices are constant on C ; then consider its projection π_I .

Now C is contained in a single fibre of π_I ; consider the fibres $L_{\bar{c}}$ and $W_{j(\bar{c})}$, for some $\bar{c} \in \mathbb{H}^{n_I}$, and the complement I_0 of the set I . The projection π_{I_0} clearly has zero-dimensional fibres on $L_{\bar{c}}$ and on $W_{j(\bar{c})}$, so it preserves dimensions. The component $\pi_{I_0}(C)$ of the intersection

$$j(\pi_{I_0}(U_{\bar{c}})) \cap \pi_{I_0}(W_{\bar{c}})$$

must have typical dimension: if it didn't it would give rise to an atypical component of the intersection of $L \times W$ with the graph of j . By Pila and Tsimerman's Ax-Schanuel theorem ([PT16]) it would then have to be contained in a weakly special subvariety, but it can't have constant coordinates, by maximality of I , and it does not satisfy modular relations because we assumed that C is not contained in $W \setminus W'$. Thus,

$$\begin{aligned} \dim C = \dim \pi_{I_0}(C) &= \dim L_{\bar{c}} + \dim W_{j(\bar{c})} - (n - n_I) > \dim L + \dim W - n \\ \dim W_{j(\bar{c})} &> \dim L - \dim L_{\bar{c}} + \dim W - n_I. \end{aligned}$$

However, the fibre dimension theorem and broadness of the pair imply that

$$\dim W_{j(\bar{c})} = \dim W - \dim \pi_I(W) \leq \dim W - n_I + \dim \pi_I(L).$$

Comparing these, we obtain:

$$\begin{aligned} \dim W - n_I + \dim \pi_I(L) &> \dim L - \dim L_{\bar{c}} + \dim W - n_I \\ \dim L_{\bar{c}} + \dim \pi_I(L) &> \dim L \end{aligned}$$

which can't hold as L is a Möbius variety. Therefore the component C cannot be atypical, and any atypical component must be contained in $W \setminus W'$. \square

We can now prove the main result.

Theorem 3.31. *Let L be a Möbius variety, (L, W) a free broad pair. Then W contains a dense subset of points of $j(L)$.*

Proof. We do this by induction on $d = \dim L$. The case $d = 1$ is Lemma 3.26, so suppose the theorem holds for d , L has dimension $d + 1$, and W has dimension $n - d - 1$.

Write L as a product $L_1 \times \cdots \times L_{d+1}$. There are numbers n_1, n_2 such that $n_1 + n_2 = n$, $L' := L_1 \times \cdots \times L_d$ is a d -dimensional Möbius subvariety of \mathbb{H}^{n_1} and L_{d+1} is a 1-dimensional Möbius subvariety of \mathbb{H}^{n_2} ; let $\pi_i : \mathbb{C}^n \rightarrow \mathbb{C}^{n_i}$, for $i = 1, 2$ denote the corresponding projections on the codomain. By broadness, $\pi_1(W)$ has dimension at least $n_1 - d$, and hence by the inductive hypothesis it contains a dense subset of points of $j(L')$; now there are two cases.

If $\dim(\pi_1(W)) = n_1 - d$, then by the fibre dimension theorem any point $\bar{w}_1 \in \pi_1(W) \cap j(L')$ has a fibre $W_{\bar{w}_1}$ of dimension at least $\dim W - (n_1 - d) = n_2 - 1$. Therefore by Lemma 3.27 $\pi_2(W_{\bar{w}_1})$ has a dense subset of points of $j(L_{d+1})$, and we are done.

If $\dim(\pi_1(W)) = n_1 - d + k$ for some positive k , then for a generic point $\bar{w} \in \pi_1(W)$ the fibre $W_{\bar{w}}$ has dimension $n_2 - 1 - k$, so we can't argue as in the previous case.

As the pair $(L', \pi_1(W))$ is broad, by the inductive hypothesis and intersecting $\pi_1(W)$ with generic hyperplanes as in the proof of Lemma 4.31 in [Asl18] if necessary, we may assume $L' \times \pi_1(W)$ intersects the graph of j in an analytic set of dimension $k' \geq k$. Then, denoting by Γ_j the graph of j ,

$$\pi_1^{-1}(L' \times \pi_1(W) \cap \Gamma_j) = (L' \times \mathbb{C}^{n_2}) \times W \cap \Gamma_j$$

has dimension $k' + n_2 - 1 - k \geq n_2 - 1$.

Now let U be a small open ball in L' , so that $(j(U) \times \mathbb{C}^{n_2}) \cap W$ is an analytic set in $j(U) \times \mathbb{C}^{n_2}$, and π_{res} denote the restriction of the second projection π_2 to the set $(j(U) \times \mathbb{C}^{n_2}) \cap W$.

Suppose that π_{res} is finite: then it is actually a *proper map*, i.e. the preimage of a compact set is compact. To prove this we use the following fact.

Fact 3.32 ([Chi12], p.3.1). Let X and Y be topological spaces, $G \subseteq X$ and $D \subseteq Y$ subsets with \overline{G} compact. If A is a closed subset of $G \times D$, the projection $\pi_2 : A \rightarrow D$ is proper if and only if A has no limit points in $\partial G \times D$.

In our setting, $X = \mathbb{C}^{n_1}$, $Y = D = \mathbb{C}^{n_2}$, $G = j(U)$, and $A = j(U) \times \mathbb{C}^{n_2} \cap W$; let \overline{w}_2 be a point in the image of π_{res} with finite fibre. Given that π_{res} is finite, there is a ball $B \subseteq j(U)$ such that $W_{\overline{w}_2} \cap j(U) \times \mathbb{C}^{n_2}$ does not intersect the set $\partial B \times \{\overline{w}_2\}$. As $W \cap j(U) \times \mathbb{C}^{n_2}$ is closed in $j(U) \times \mathbb{C}^{n_2}$ and π_{res} is finite and hence open, this property actually holds in a neighbourhood of \overline{w}_2 ; therefore, by Fact 3.32, by taking U sufficiently small we can make sure that the map π_{res} is proper.

So, under the assumption that π_{res} is finite, we can apply the proper mapping theorem (see p.5.8 in [Chi12]), which states that the image of π_{res} is an analytic set. Since we proved $j(U) \times \mathbb{C}^{n_2} \cap W$ has dimension at least $n_2 - 1$, the image of π_{res} is either an open subset of \mathbb{C}^{n_2} , or an analytic set in \mathbb{C}^{n_2} of codimension 1; either way, using density of $j(L_{d+1})$ in the first case and Lemma 3.27 otherwise, we can find a point $\overline{w} \in W \cap j(U) \times \mathbb{C}^{n_2}$ such that $\pi_2(\overline{w}) \in j(L_{d+1})$; therefore $\overline{w} \in j(L) \cap W$, as we wanted.

Therefore, it remains to show that if we choose the ball U appropriately, then the map π_{res} is finite. Consider a generic point $\overline{w}_2 \in \pi_2(W)$, and its fibre

$$W_{\overline{w}_2} := \{\overline{w} \in W \mid \pi_2(\overline{w}) = \overline{w}_2\}.$$

Using once again broadness and the fibre dimension theorem, we know that

$$\dim W_{\overline{w}_2} = \dim W - \dim \pi_2(W) \leq n - d - 1 - n_2 + 1 = n_1 - d.$$

Hence,

$$\dim W_{\overline{w}_2} + \dim(j(L') \times \mathbb{C}^{n_2}) \leq n_1 - d + d + n_2 = n.$$

Therefore any positive dimensional intersection between these varieties is of atypical dimension; by Lemma 3.30, it suffices to make sure that $j(U) \times \mathbb{C}^{n_2} \cap W$ is contained in the Zariski-open non-empty subset W' of W for this property to be verified. This set clearly contains points of $j(L') \times \mathbb{C}^{n_2}$, because $\pi_1(W) \cap j(L')$ is dense in $\pi_1(W)$, so we're done. \square

4 Derivatives of the j Function

4.1 Background and Notation

We conclude with some remarks on extensions of the results in the previous section to the first and second derivative of the j function.

First of all we remark that while j is a modular function, j' and j'' are not: j' is a modular form of weight 2, and the transformation law for j'' under the action of $\mathrm{SL}_2(\mathbb{Z})$ is more complicated. This means that the methods in the previous section do not apply directly if we look for intersections of the form $j'(L) \cap W$ for L a Möbius subvariety of \mathbb{H}^n .

Therefore to get more general results we need to work in jet spaces. The action of $\mathrm{SL}_2(\mathbb{R})$ on \mathbb{H} induces an action on $J_2\mathbb{H}$, $J_2\mathcal{F}$ is a fundamental domain for the action of $\mathrm{SL}_2(\mathbb{Z})$ and J_2j is invariant under the action of $\mathrm{SL}_2(\mathbb{Z})$, so this seems like a better framework to work in.

We recall some general facts about jet spaces, and provide explicit computations for the facts stated above.

Definition 4.1. Let M be a complex analytic manifold. The k -th jet space of M for a natural number k is the space of equivalence classes of holomorphic functions from a small neighbourhood of $0 \in \mathbb{C}$ into M , identifying maps that are equal up to order k .

We will only be interested in second jets, so we assume $k = 2$ in the following.

An element in $J_2\mathbb{H}$ is then a triple (z, r, s) , where $z \in \mathbb{H}$, $r, s \in \mathbb{C}$, that corresponds to the function $f : U \rightarrow \mathbb{H}$ taking w to $z + rw + s\frac{w^2}{2}$.

Jets are a functorial construction: given a map $\varphi : M \rightarrow N$, there is an induced map $J_k\varphi : J_kM \rightarrow J_kN$, that takes the equivalence class of the function $f : U \rightarrow M$ to that of $\varphi \circ f$. Therefore, if for a fixed $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R})$ we consider the map $g \cdot (-) : \mathbb{H} \rightarrow \mathbb{H}$, we can see what the action induced on $J_2\mathbb{H}$ is:

$$g \cdot (z, r, s) = \left(\frac{az + b}{cz + d}, \frac{r}{(cz + d)^2}, \frac{s}{(cz + d)^2} - \frac{2cr^2}{(cz + d)^3} \right)$$

Similarly we have to consider the second jet of the j function itself, which is obtained as:

$$J_2j(z, r_1, r_2) = (j(z), j'(z)r, j''(z)r^2 + j'(z)s)$$

so that in particular for example $J_2j(z, 1, 0) = (j(z), j'(z), j''(z))$.

Using the transformation laws for j' and j'' , i.e., for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$,

$$j'(\gamma z) = (cz + d)^2 j'(z)$$

and

$$j''(\gamma z) = (cz + d)^4 j''(z) + 2c(cz + d)^3 j'(z)$$

one can prove that $J_2j(\gamma \cdot (z, r, s)) = J_2j(z, r, s)$.

4.2 Intersections for the Derivatives

4.2.1 The First Derivative

For simplicity, we deal with the case of a pair (L, W) of subvarieties of $\mathbb{H}^2 \times \mathbb{C}^2$, as this clearly implies the general case.

The fact that j' is not $\mathrm{SL}_2(\mathbb{Z})$ -invariant leads us to consider a blurring of a Möbius subvariety. This is similar to what is done in [Kir19] for the exponential function and in [AK20] for j , but note that here the blurring is not fixed, and it depends on the variety that we are considering.

Definition 4.2. Let L be the Möbius subvariety of \mathbb{H}^2 of points of the form (z, gz) , for some $g = \begin{pmatrix} a_g & b_g \\ c_g & d_g \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R})$. Consider the tangent variety

$$L' := \left\{ (z_1, z_2, z_3, z_4) \in J_1\mathbb{H}^2 \mid z_2 = gz_1, z_3 = 1, z_4 = \frac{1}{(c_g z_1 + d_g)^2} \right\}.$$

The *blurring* of L' by $\mathrm{SL}_2(\mathbb{Z})$ is the set L^* , defined as

$$\left\{ (z_1, gz_1, z_3, z_4) \in J_1\mathbb{H}^2 \mid z_3 = \frac{1}{(c_\gamma(\gamma^{-1}z_1) + d_\gamma)^2}, z_4 = \frac{1}{(c_{g\gamma}(\gamma^{-1}z_1) + d_{g\gamma})^2} \right\}$$

where $\gamma = \begin{pmatrix} a_\gamma & b_\gamma \\ c_\gamma & d_\gamma \end{pmatrix}$ varies in $\mathrm{SL}_2(\mathbb{Z})$ and $g\gamma = \begin{pmatrix} a_{g\gamma} & b_{g\gamma} \\ c_{g\gamma} & d_{g\gamma} \end{pmatrix}$.

Definition 4.2 needs to be explained. A point in the first jet of \mathbb{H}^2 carries two bits of information: the first two coordinates give a point in \mathbb{H}^2 , while the second two coordinates refer to the direction from which the point is approached. Hence, the variety L' is the variety of points in L , where to each point we associate the natural direction of the parametrization $z \mapsto (z, gz)$. When we move to the blurred version L^* , its projection to the first two coordinates is still the original Möbius variety L ; however, each of these points is considered together with the directions associated to the parametrizations $z \mapsto (\gamma z, g\gamma z)$, for all $\gamma \in \mathrm{SL}_2(\mathbb{Z})$.

Let $T_1j : J_2\mathbb{H}^2 \rightarrow \mathbb{C}^2$ denote the composition $\pi \circ J_1j$, where $\pi : J_2\mathbb{C}^2 \cong \mathbb{C}^4 \rightarrow \mathbb{C}^2$ is the projection on the third and fourth coordinate. Hence,

$$T_1(j)(z_1, z_2, r_1, r_2) = (j'(z_1)r_1, j'(z_2)r_2).$$

This allows to prove the following proposition:

Proposition 4.3. *Let L be a free Möbius subvariety of \mathbb{H}^2 . Then $T_1j(L^*)$ is dense in \mathbb{C}^2 .*

Proof. Let (w_1, w_2) be a point in \mathbb{C}^2 . As j' is surjective, find z_1 such that $j'(z_1) = w_1$, and $h = \begin{pmatrix} a_h & b_h \\ c_h & d_h \end{pmatrix} \in \mathrm{SL}_2(\mathbb{R})$ such that $\frac{j'(hz_1)}{(c_h z_1 + d_h)^2} = w_2$. Then, as in the proof of Lemma 3.13, we can consider a sequence of matrices in $\Gamma g\Gamma$ that tends to h ; denote a specific element of this sequence by $\gamma_1 g\gamma$. Then we just need to prove that $\left(j'(z_1), \frac{j'(\gamma_1 g\gamma z_1)}{(c_{\gamma_1 g\gamma} z_1 + d_{\gamma_1 g\gamma})^2} \right)$ is in $T_1j(L^*)$. Consider the point

$$\bar{z} = \left(z_1, \gamma_1 g\gamma z_1, 1, \frac{1}{(c_{\gamma_1 g\gamma} z_1 + d_{\gamma_1 g\gamma})^2} \right).$$

It's clear that

$$J_1 j(\bar{z}) = \left(j(z_1), j(\gamma_1 g \gamma z), j'(z_1), \frac{j'(\gamma_1 g \gamma z_1)}{(c_{\gamma_1 g \gamma z_1} + d_{\gamma_1 g \gamma})^2} \right)$$

Since $J_1 j$ is invariant under the action of $\mathrm{SL}_2(\mathbb{Z})$, it is then sufficient to find an element $\bar{\gamma} \in \mathrm{SL}_2(\mathbb{Z})^2$ such that $\bar{\gamma} \cdot \bar{z} \in L^*$. Then of course we consider $\bar{\gamma} = (\gamma, \gamma_1^{-1})$.

Claim: $\bar{\gamma} \cdot \bar{z} = \left(\gamma z_1, g \gamma z_1, \frac{1}{(c_{\gamma z_1} + d_{\gamma})^2}, \frac{1}{(c_{g \gamma z_1} + d_{g \gamma})^2} \right)$.

Proof of Claim: It's immediate for the first three coordinates, the only issue is on the fourth one. Using the well-known cocycle relation for automorphy factors we obtain that

$$(c_{g \gamma z} + d_{g \gamma z})^2 = (c_{\gamma_1 g \gamma z} + d_{\gamma_1 g \gamma})^2 (c_{\gamma_1^{-1}}(\gamma_1 g \gamma z) + d_{\gamma_1^{-1}})^2.$$

But the fourth coordinate of $\bar{\gamma} \cdot \bar{z}$ is

$$\frac{1}{(c_{\gamma_1 g \gamma z} + d_{\gamma_1 g \gamma})^2 (c_{\gamma_1^{-1}}(\gamma_1 g \gamma z) + d_{\gamma_1^{-1}})^2}$$

which is then equal to

$$\frac{1}{(c_{g \gamma z} + d_{g \gamma z})^2}$$

proving the claim.

It remains to show that $\bar{\gamma} \cdot \bar{z}$ is indeed a point in L^* . The condition $z_2 = g z_1$ is clearly satisfied. The second condition in the definition of the blurring is given by $\frac{1}{(c_{\gamma(\gamma^{-1} \gamma z_1)} + d_{\gamma})^2} = \frac{1}{(c_{\gamma z_1} + d_{\gamma})^2} = z_3$. Similarly the last condition is satisfied.

This concludes the proof: the sequence $T_1 j((\bar{g}_i) z_1)$ approximates w_2 and hence $T_1 j(L^*)$ is dense in \mathbb{C}^2 . \square

Remark 4.4. The crucial piece of information is that while j' is not a modular function, it is a modular form of weight 2 for $\mathrm{SL}_2(\mathbb{Z})$, and hence it is easy to come up with the blurring that makes the density argument work. This leads to speculations about similar results for modular forms in more generality - although we are not aware of any framework that could motivate such an investigation.

We now close the argument, showing that given an open neighbourhood U inside a complex algebraic variety $W \subseteq \mathbb{C}^2$, the intersection $U \cap T_1 j(L^*)$ is non-empty. While in the previous section we used transversality and Newton's method, that approach is more complicated here as it requires explicit, hands-on computations involving j' and j'' which are hard to solve. Instead, we go down a different road and, recalling that the restrictions of j and its derivatives to fundamental domains are definable in the o-minimal structure $\mathbb{R}_{\mathrm{an}, \mathrm{exp}}$ by well-known results of Peterzil-Starchenko (see [PS04], Theorem 4.1), we use o-minimality. In particular, we'll need the following result of Johns. A similar statement also holds in the j case, but we find it instructive to show that this can be done in two different ways.

Theorem 4.5 ([Joh01], Theorem). *Let $R = (R, <, \dots)$ be an o-minimal structure, $U \subseteq R^n$ an open set, $f : U \rightarrow R^n$ a continuous, definable, injective function. Then f is open.*

We'll also need an analogue for j' of Proposition 3.25, giving that the fibre of certain maps are one-dimensional subsets of $\mathrm{SL}_2(\mathbb{R})$.

Proposition 4.6. *Let $(w_1, w_2) \in W$. Then given z such that $j'(z) = w_1$, there is a definable one-dimensional subset H of $\mathrm{SL}_2(\mathbb{R})$ such that for every $h = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in H$, $\frac{j'(hz)}{(cz+d)^2} = w_2$.*

Proof. Fix \bar{w} and z as in the statement. First we note that, as $\frac{1}{|cz+d|^2} = \frac{\Im(hz)}{\Im(z)}$, $\frac{j'(hz)}{(cz+d)^2} = w_2$ implies

$$|w_2| = \left| \frac{j'(hz)}{(cz+d)^2} \right| = \left| j'(hz) \frac{\Im(hz)}{\Im(z)} \right|.$$

Hence, for h to satisfy $(j(hz))' = \frac{j'(hz)}{(cz+d)^2} = w_2$, it is necessary that

$$|j'(hz)|\Im(hz) = |w_2|\Im(z).$$

Now let us consider the function $\varphi : w \mapsto j'(w)\Im(w)$. By using the Fourier series expansion of j ,

$$j(z) = \sum_{n=-1}^{\infty} c_n e^{2\pi n i z}$$

where all c_n 's are positive integers (see for example [Rad38]) we see that j' takes imaginary values on $i\mathbb{R}_{\geq 1}$ and that its modulus is strictly increasing; a direct computation shows that the Jacobian of φ as a function $\mathbb{R} \times \mathbb{R}_{>0} \rightarrow \mathbb{R}^2$ is nonsingular on points of $\{0\} \times \mathbb{R}_{\geq 1}$. Hence, as $j'(i) = 0$ and j' has a simple pole at infinity, for every $x \in \mathbb{R}_{\geq 0}$ there is a $z \in i\mathbb{R}_{\geq 1}$ such that $|\varphi(z)| = x$ and φ is open around z .

Then the definable set of w 's such that $|j'(w)|\Im(w) = |w_2|\Im(z)$ is non-empty and one-dimensional. For each of these w 's, by Proposition 3.25, there is a unique $h \in \mathrm{SL}_2(\mathbb{R})$ such that $hz = w$ and $j(hz)' = w_2$, so we obtain H as required. \square

Lemma 4.7. *Let $W \subseteq \mathbb{C}^2$ be a one-dimensional algebraic variety not contained in the image through $T_1 j$ of a weakly special variety, $U \subseteq W$ an open subset. Then there is an open subset $O \subseteq \mathrm{SL}_2(\mathbb{R})$ such that for every $h \in O$, there is a point in U of the form $(j'(z), j(hz)')$.*

In particular, by the results in Subsection 3.2, $O \cap \Gamma g \Gamma \neq \emptyset$ for every g that is not a multiple of a rational matrix.

Proof. Let f be the polynomial defining W , and let S be a connected component of the set

$$\{(z, h) \in \mathbb{C} \times \mathrm{SL}_2(\mathbb{R}) \mid f(j'(z), (j(hz))') = 0\}.$$

Denote by $\pi_1(S)$ the projection to \mathbb{C} and by $\pi_2(S)$ the projection to $\mathrm{SL}_2(\mathbb{R})$.

S is definable in $\mathbb{R}_{\mathrm{an}, \mathrm{exp}}$. Since Skolem functions are definable in o-minimal structures, there is a definable function $F : \pi_1(S) \rightarrow \mathrm{SL}_2(\mathbb{R})$ that satisfies $(z, F(z)) \in S$ for each $z \in \pi_1(S)$. We can assume F is injective, as W is not contained in the image of a weakly special variety, and therefore by Theorem 4.5 it is open on its image. But every point $F(z)$ in the image has a small one-dimensional neighbourhood V such that actually $(z, v) \in S$ for every $v \in V$, by Proposition 4.6. Therefore $\pi_2(S)$ has non-empty interior, and we are done. \square

Now the following analogue of Theorem 3.31 for j' can be obtained by applying the same argument.

Theorem 4.8. *Let (L, W) be a free broad pair in $\mathbb{H}^n \times \mathbb{C}^n$. Then W has a dense subset of points of $T_1(j(L^*))$.*

4.2.2 The Second Derivative

There is no real motivation for finding solutions of systems of equations which only involve the second derivative of the j function: the Ax-Schanuel theorem for j and its derivatives involves j, j' , and j'' , and it seems that new ideas are needed to tackle that problem. As the present method does not directly apply to j'' , we do not deal with that case.

The situation once we consider j'' becomes more complicated: while j' is a modular form of weight 2, j'' is a *quasimodular form of weight 4 and depth 1*. In particular, to apply the same argument as in the j' case we would have to consider the blurring of the space L'' of points with the following form:

$$\left(z, gz, \frac{1}{(c_\gamma(\gamma^{-1}z) + d_\gamma)^2}, \frac{1}{(c_{g\gamma}(\gamma^{-1}z) + d_{g\gamma})^2}, \right. \\ \left. - \frac{2c_\gamma}{(c_\gamma(\gamma^{-1}z) + d_\gamma)^3}, - \frac{2c_{g\gamma}}{(c_{g\gamma}(\gamma^{-1}z) + d_{g\gamma})^3} \right)$$

Denoting by T_2j the composition of J_2j with the projection on the fifth and sixth coordinates, we see that given a point \bar{z} as above,

$$T_2j(\bar{z}) = (j(\gamma z_1)'', j(g\gamma z_1)'')$$

for some $\gamma \in \mathrm{SL}_2(\mathbb{Z})$.

However, we note that it seems likely that a result similar to Theorems 3.31 and 4.8 holds for j'' . However, to apply the same methods it would be necessary to have a version of Propositions 3.25 and 4.6 for j'' : in short, we would require the set

$$\{(z, h) \in \mathbb{C} \times \mathrm{SL}_2(\mathbb{R}) \mid j''(z) = w_1, j''(hz) = w_2\}$$

to be a real one-dimensional subset of $\mathbb{C} \times \mathrm{SL}_2(\mathbb{R})$ for every $w_1, w_2 \in \mathbb{C}^2$.

References

- [AEK21] Vahagn Aslanyan, Sebastian Eterović, and Jonathan Kirby. “Differential existential closedness for the j -function”. In: *Proceedings of the American Mathematical Society* 149.4 (2021), pp. 1417–1429. DOI: 10.1090/proc/15333.
- [AK20] Vahagn Aslanyan and Jonathan Kirby. *Blurrings of the j -function*. 2020. arXiv: 2005.10167.
- [Asl18] Vahagn Aslanyan. *Adequate Predimension Inequalities in Differential Fields*. 2018. arXiv: 1803.04753.
- [Asl20] Vahagn Aslanyan. *Weak Modular Zilber-Pink with Derivatives*. 2020. arXiv: 1803.05895.
- [Ber02] Cristiana Bertolin. “Périodes de 1-motifs et Transcendance”. In: *Journal of Number Theory* 97.2 (2002), pp. 204–221. DOI: 10.1016/S0022-314X(02)00002-1.
- [BK18] Martin Bays and Jonathan Kirby. “Pseudo-exponential maps, variants, and quasiminimality”. In: *Algebra And Number Theory* 12.3 (June 2018), pp. 493–549. DOI: 10.2140/ant.2018.12.493.
- [BM17] Woodrow Brownawell and David Masser. “Zero estimates with moving targets”. In: *Journal of the London Mathematical Society* 95 (Apr. 2017). DOI: 10.1112/jlms.12014.
- [Che46] Claude Chevalley. *Theory of Lie Groups*. Princeton University Press, 1946.
- [Chi12] Evgeni Mikhailovich Chirka. *Complex analytic sets*. Vol. 46. Springer Science & Business Media, 2012.
- [DFT18] Paola D’Aquino, Antongiulio Fornasiero, and Giuseppina Terzo. “Generic Solutions of Equations with Iterated Exponentials”. In: *Transactions of the American Mathematical Society* 370.2 (2018), pp. 1393–1407. DOI: 10.1090/tran/7206.
- [DS96] John E. Dennis Jr and Robert B. Schnabel. *Numerical Methods for Unconstrained Optimization and Nonlinear Equations*. SIAM, 1996.
- [EH20] Sebastian Eterović and Sebastián Herrero. “Solution of Equations Involving the Modular j Function”. In: *Transactions of the American Mathematical Society (to appear)* (2020).
- [HS00] Marc Hindry and Joseph H Silverman. *Diophantine Geometry: An introduction*. Vol. 201. 2000.
- [Jac] Nathan Jacobson. *Lie Algebras*. Interscience tracts in pure and applied mathematics. Interscience Publishers.
- [Joh01] Joseph Johns. “An Open Mapping Theorem for O-Minimal Structures”. In: *J. Symbolic Logic* 66.4 (Dec. 2001), pp. 1817–1820. DOI: 10.2307/2694977.

- [Kir06] Jonathan Kirby. “The Theory of Exponential Differential Equations”. PhD thesis. Magdalen College, University of Oxford, 2006.
- [Kir09] Jonathan Kirby. “The theory of the exponential differential equations of semiabelian varieties”. English. In: *Selecta Mathematica* 15.3 (2009), pp. 445–486. DOI: 10.1007/s00029-009-0001-7.
- [Kir19] Jonathan Kirby. “Blurred Complex Exponentiation”. In: *Selecta Mathematica* 25.5 (Dec. 2019). DOI: 10.1007/s00029-019-0517-4.
- [KUY16] Bruno Klingler, Emmanuel Ullmo, and Andrei Yafaev. “Bi-algebraic geometry and the André-Oort conjecture”. In: *Algebraic geometry (Salt Lake City, 2015)* (2016), pp. 319–359.
- [Lan04] Serge Lang. “Algebra, volume 211 of”. In: *Graduate Texts in Mathematics* (2004), pp. 29–30.
- [Mar06] David Marker. “A Remark on Zilber’s Pseudoexponentiation”. In: *The Journal of Symbolic Logic* 71.3 (2006), pp. 791–798. DOI: 10.2178/jsl/1154698577.
- [Mil97] James S Milne. *Modular Functions and Modular Forms*. 1997.
- [Mor05] Dave Witte Morris. *Ratner’s theorems on unipotent flows*. University of Chicago Press, 2005.
- [MZ16] Vincenzo Mantova and Umberto Zannier. “Polynomial–exponential equations and Zilber’s conjecture”. In: *Bulletin of the London Mathematical Society* 48.2 (Feb. 2016), pp. 309–320. DOI: 10.1112/blms/bdv096.
- [PS04] Ya’acov Peterzil and Sergei Starchenko. “Uniform Definability of the Weierstrass \wp Functions and Generalized Tori of Dimension One”. In: *Selecta Math.(NS)* 10.4 (2004), pp. 525–550. DOI: 10.1007/s00029-005-0393-y.
- [PT16] Jonathan Pila and Jacob Tsimerman. “Ax–Schanuel for the j -function”. In: *Duke Mathematical Journal* 165.13 (2016), pp. 2587–2605. DOI: 10.1215/00127094-3620005.
- [Rad38] Hans Rademacher. “The Fourier Coefficients of the Modular Invariant $J(\tau)$ ”. In: *American Journal of Mathematics* 60.2 (1938), pp. 501–512.
- [Rat91] Marina Ratner. “On Raghunathan’s Measure Conjecture”. In: *Annals of Mathematics* 134.3 (1991), pp. 545–607. DOI: 10.2307/2944357.
- [UY11] Emmanuel Ullmo and Andrei Yafaev. “A Characterization of Special Subvarieties”. In: *Mathematika* 57.2 (2011), pp. 263–273. DOI: 10.1112/S0025579311001628.
- [UY18] Emmanuel Ullmo and Andrei Yafaev. “Algebraic Flows on Abelian Varieties”. In: *Journal für die reine und angewandte Mathematik* 2018.741 (2018), pp. 47–66. DOI: 10.1515/crelle-2015-0085.
- [Zil02] Boris Zilber. “Exponential Sums Equations and the Schanuel Conjecture”. In: *Journal of the London Mathematical Society* 65.1 (2002), pp. 27–44. DOI: 10.1112/S0024610701002861.

- [Zil05] Boris Zilber. “Pseudo-exponentiation on algebraically closed fields of characteristic zero”. In: *Annals of Pure and Applied Logic* 132.1 (2005), pp. 67–95. DOI: 10.1016/j.apal.2004.07.001.
- [Zil15] Boris Zilber. *The theory of exponential sums*. 2015. arXiv: 1501.03297.