

# First Identification of a CMB Lensing Signal Produced by 1.5 Million Galaxies at $z \sim 4$ : Constraints on Matter Density Fluctuations at High Redshift

Hironao Miyatake<sup>1,2,3</sup>, Yuichi Harikane<sup>4,5</sup>, Masami Ouchi<sup>6,4,3</sup>, Yoshiaki Ono<sup>4</sup>, Nanaka Yamamoto<sup>2</sup>,

Atsushi J. Nishizawa<sup>1,2</sup>, Neta Bahcall<sup>7</sup>, Satoshi Miyazaki<sup>6</sup>, Andrés A. Plazas Malagón<sup>7</sup>

<sup>1</sup>*Institute for Advanced Research, Nagoya University, Nagoya, 464-8601, Japan*

<sup>2</sup>*Division of Physics and Astrophysical Science, Graduate School of Science, Nagoya University, Furo-cho, Chikusa, Nagoya 464-8602, Japan*

<sup>3</sup>*Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU, WPI), UTIAS, The University of Tokyo, 5-1-5 Kashiwa-no-ha, Kashiwa, Chiba 277-8583, Japan*

<sup>4</sup>*Institute for Cosmic Ray Research, The University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8582, Japan*

<sup>5</sup>*Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK*

<sup>6</sup>*National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan and*

<sup>7</sup>*Department of Astrophysical Sciences, Peyton Hall, Princeton University, Princeton, NJ 08540, USA*

(Dated: January 6, 2022)

We report the first detection of the dark matter distribution around Lyman break galaxies (LBGs) at high redshift through the Cosmic Microwave Background (CMB) lensing measurements with the public *Planck* PR3  $\kappa$  map. The LBG sample consists of 1,473,106 objects with the median redshift of  $z \sim 4$  that are identified in a total area of 305 deg<sup>2</sup> observed by the Hyper Suprime-Cam (HSC) Strategic Survey Program (SSP) survey. After careful investigations of systematic uncertainties, such as contamination from foreground galaxies and Cosmic Infrared Background (CIB), we obtain the significant detection of the CMB lensing signal at  $5.1\sigma$  that is dominated by 2-halo term signals of the LBGs. Fitting a simple model consisting of the Navarro–Frenk–White (NFW) profile and the linear-bias model, we obtain the typical halo mass of  $3.1_{-2.8}^{+9.3} \times 10^{11} h^{-1} M_{\odot}$ . Combining the CMB lensing and galaxy-galaxy clustering signals on the large scales, we demonstrate the first cosmological analysis at  $z \sim 4$  that constrains  $(\Omega_{m0}, \sigma_8)$ . We find that our constraint on  $\sigma_8$  is roughly consistent with the *Planck* best-fit cosmology, while this  $\sigma_8$  constraint is lower than the *Planck* cosmology over the  $1\sigma$  level. This study opens up a new window for constraining cosmological parameters at high redshift by the combination of CMB and high- $z$  galaxies as well as studying the interplay between galaxy evolution and large-scale structure at such high redshift, by upcoming CMB and optical and near-infrared imaging surveys.

## INTRODUCTION

Understanding the interplay between galaxy evolution and large-scale structure is key to understanding cosmic evolution. Since galaxies are formed in dark matter halos through gas cooling, such an interplay can be studied by measuring the connection between dark matter halos and galaxies [see reviews by 1, 2]. Galaxy-galaxy clustering, the auto-correlation of galaxy positions, is one of the statistical probes that has been widely used to investigate the galaxy-halo connection [e.g., 3–7]. Galaxy-galaxy lensing, the cross-correlation between galaxy positions and weak lensing shear of background galaxies, is rapidly emerging as another powerful probe because it enables the direct measurement of dark matter distribution around galaxies [e.g., 8–11].

The combined measurement of galaxy-galaxy clustering and galaxy-galaxy lensing allows us to constrain cosmological parameters through growth of structure, and thus it is now the standard technique in cosmological galaxy surveys. Such studies can be divided into those using both small and large scales which enable us to investigate the galaxy-halo connection simultaneously [e.g., 12, 13], and those focusing on large scales to avoid pos-

sible systematics due to our incomplete understanding of galaxy physics [e.g., 13–16]. These measurements advanced our understanding of cosmic evolution of large-scale structure at  $z < 1$  through the constraint on the amplitude of matter power spectrum  $\sigma_8$ , which are consistent with the *Planck* cosmology so far. As an independent test, it is of great importance to add such a measurement at higher redshift. Having structure growth measurements with a wide redshift may also allow us to address the  $H_0$  tension between measurements of late time probes such as the type Ia supernova distance and early time probes such as cosmic microwave background (CMB) [e.g., 17, 18]

Recently, detailed study results of galaxy properties and galaxy-halo connection through clustering measurement of Lyman break galaxies (LBGs) at  $z \sim 4 - 7$  are reported [7]. Although one of the next steps is measuring weak lensing on background galaxies around the LBGs, it is practically impossible due to the lack of data for shear measurements of galaxies behind LBGs. Instead, we can use weak lensing on the CMB anisotropies, which is sensitive to the large-scale structure at  $z > 1$ . In this paper, using CMB lensing measurements in the *Planck* satellite data, we present the first detection of dark-matter distri-

bution around LBGs at  $z \sim 4$ . We also show cosmological constraints from clustering and lensing measurements.

## DATA

### HSC GOLDRUSH Catalog

We use the second public data release (PDR2) product taken by the Hyper Suprime-Cam (HSC) Strategic Survey Program (SSP) from 2014 March to 2018 January [19]. We use the Wide layer data whose total full color full depth area is  $305 \text{ deg}^2$ . The HSC data are reduced by the HSC SSP collaboration with `hscPipe` version 6.7 [20, 21]. The typical  $5\sigma$  limiting magnitudes in a  $1.5''$  diameter aperture are 26.4, 25.9, 25.8, 25.1, and 24.2 mag in  $g$ ,  $r$ ,  $i$ ,  $z$ , and  $y$ -bands, respectively.

To construct a  $z \sim 4$  LBG catalog of Great Optically Luminous Dropout Research Using Subaru HSC (GOLDRUSH) [22], we use the dropout selection method. Here we briefly describe the selection method; full details of the catalog construction will be presented in Harikane et al. (in prep). We use the following color criteria to select  $z \sim 4$  galaxies:

$$g - r > 1.0 \quad (1)$$

$$r - i < 1.0 \quad (2)$$

$$g - r > 1.5(r - i) + 0.8, \quad (3)$$

which are the same as the GOLDRUSH catalog construction with the first HSC data release [7, 22, 23]. In addition to these criteria, we apply a detection limit of a  $> 5\sigma$  level in the  $i$ -band. We evaluate source detections with aperture magnitudes, and measure total fluxes and colors of sources with `convolvedflux` magnitudes, which show good agreement with the `cModel` magnitude in the first public data release [24]. We mask out regions affected by bright object halos or unreliable background subtractions. Finally, we select a total of  $\sim 2,000,000$  LBGs. Ono *et al.* [22] used spectroscopically identified high- $z$  galaxies in the dropout sample to estimate the average redshift of the sample  $\langle z \rangle = 3.8$  (median redshift of  $z \sim 4$ ), which we adopt for our analysis. The surface number density and angular correlation functions of the LBGs agree well with previous studies. In our analysis, we use 1,643,863 LBGs whose total  $i$ -band magnitudes are brighter than 25.5 mag, which is reduced to 1,473,106 LBGs after applying the *Planck* mask described below. Some foreground objects such as red galaxies at intermediate redshift can satisfy our selection criteria by photometric errors. A contamination rate is estimated to be  $f_{\text{cont}} \sim 0.25$  based on HSC WIDE layer results obtained by [22] who derive a fraction of contaminating galaxies as a function of  $i$ -band magnitude by calculating the number of foreground galaxies that satisfy the color criteria.

### *Planck* $\kappa$ map

We construct the convergence map using lensing products in the third public data release from the *Planck* satellite mission [25] [26]. We use the minimum variance (MV) estimator and masks in the baseline lensing potential estimates. The MV estimators are provided as harmonic coefficients with the maximum multipole  $L_{\text{max}} = 4096$ , while the mask is in real space. To construct the convergence map, we subtract the MV mean field from the MV convergence, and transform the harmonic coefficients to a real-space map applying a Gaussian filter with  $\sigma = 2'$ . We apply the mask to remove the galactic plane, point sources, and Sunyaev-Zel'dovich (SZ) clusters, which leaves a total unmasked sky fraction  $f_{\text{sky}} = 0.671$ . This results in the healpix map with NSIDE=2048 where the area of each healpixel is  $2.95 \text{ arcsec}^2$ . Note that the HSC-SSP survey does not include the galactic plane, and that we have a full overlap between our LBG sample and the *Planck* convergence map. After removing regions of point sources and SZ clusters masked in the *Planck* data, the total overlap area is  $270 \text{ deg}^2$ .

## MEASUREMENT

To measure the stacked convergence profile, we first create galaxy-centric angular bins for each galaxy by linearly dividing  $0' < \theta < 20'$  into 14 bins, and identify healpixels that belong to each bin. We then collect the healpixels over all galaxies, and compute the average convergence in each bin. To subtract the residual mean field, we also perform the same measurement around random points, and subtract the random signal from the observed convergence profile. The number of random points is 40 times larger than that of the LBGs.

We estimate the jackknife covariance dividing the PDR2 area into 262 subregions such that each subregion has a similar number of galaxies, and measure the mean-subtracted signal for each jackknife subregion. Note that the covariance is highly correlated due to the low-pass filter applied in the process of MV convergence estimation.

To test the contamination from cosmic infrared background (CIB), we stack *Planck* SZ-nulled temperature map (`SMICA_noSZ`) around the LBGs. We do not find a gradient in the signal, and thus conclude that there is no significant CIB contamination. In addition we confirm that there is no significant B-mode signal.

## RESULTS

In Fig. 1, we show the measured convergence profile together with the mean-field profile and the convergence profile before the mean-field subtraction. Defining the




FIG. 1. *Planck* CMB lensing measurements around  $z \sim 4$  LBGs. The red circles denote the measured lensing signal after the mean-field subtraction, where the filled (open) circles denote data points used for (excluded from) our model fit. The black crosses (triangles) denote the lensing signal before the mean-field subtraction (mean-field signal measured by stacking the *Planck*  $\kappa$  map around random points). The red solid (dashed) line represents our best-fit model of the lensing signal at the scales used for (excluded from) our model fit. The red dotted line shows the lensing signal of low-redshift contaminating galaxies in the LBG sample constrained by optical lensing measurements with the HSC shapes (for details, see text). The significance of the signal is  $5.1\sigma$  ( $3.5\sigma$ ) against null (the contamination signal).

significance of the convergence profile after the mean-field subtraction as  $[S/N]_{\kappa} = \sum_{ij} \kappa_i \text{Cov}_{ij}^{-1} \kappa_j$ , we obtain the significance level of  $[S/N]_{\kappa} = 5.1$ .

The convergence profile is contaminated by foreground galaxies as described in the previous section. We thus model our signal as  $\kappa_{\text{tot}}^{\text{obs}} = (1 - f_{\text{cont}})\kappa_{\text{LBG}}^{\text{obs}} + f_{\text{cont}}\kappa_{\text{cont}}^{\text{obs}}$ . We use the halo models for the convergence profiles of real LBGs and contaminating galaxies with different halo masses and redshifts.

Let us first consider the convergence profile without the *Planck* beam. In the halo model approach, the convergence profile is composed of 1-halo and 2-halo terms;  $\kappa(\theta) = \kappa_{1\text{h}}(\theta) + \kappa_{2\text{h}}(\theta)$ , where  $\kappa_{1\text{h}}(\theta)$  and  $\kappa_{2\text{h}}(\theta)$  are the contribution from dark matter halos around galaxies and neighboring halos, respectively. We model the 1-halo term as  $\kappa_{1\text{h}}(R) = \Sigma_{\text{cr}}^{-1}(z_1, z_{\text{CMB}})\bar{\rho}_m \int (kdk/2\pi)\tilde{u}_m(k; M_h, z_1)J_0(kR)$ , where  $z_1$  ( $z_{\text{CMB}}$ ) is the redshift of lens (CMB),  $\bar{\rho}_m$  is the mean matter density at present,  $\tilde{u}_m(k; z_1)$  is the Fourier-space Navarro–Frenk–White (NFW) [27, 28] profile truncated at the radius where the enclosed mass is 200 times the mean density  $R_{200}$ ,  $J_0(x)$  is the zeroth-order Bessel function of the first kind. We adopt the analytical form of  $\tilde{u}_m(k; z_1)$  provided by Takada and Jain [29], and assume the concentration-mass relation derived by Duffy *et al.* [30]. The critical surface density is defined as  $\Sigma_{\text{cr}}(z_1, z_s) = c^2[4\pi G]^{-1}d_A(z_s)[d_A(z_1)d_A(z_1, z_s)]^{-1}(1 + z_1)^{-2}$ , where  $c$  is the speed of light,  $G$  is the gravitational constant,

$d_A(z_s)$ ,  $d_A(z_1)$ , and  $d_A(z_1, z_s)$  are the angular diameter distances for the lens-source system. To model the 2-halo term, we adopt the linear bias model, i.e.,  $\kappa_{2\text{h}}(R) = \Sigma_{\text{cr}}^{-1}(z_1, z_{\text{CMB}})b(M_h, z_1)\bar{\rho}_m \int (kdk/2\pi)P_m(k, z_1)J_0(kR)$ , where  $b(M_h, z_1)$  is the linear bias parameter, and  $P_m(k, z_1)$  is the linear matter power spectrum at  $z_1$ . We adopt the fitting function for the linear bias derived by Tinker *et al.* [31], and compute the matter power spectrum using CAMB [32]. Note that in our model the 1-halo and 2-halo terms are connected through halo mass. Finally, we convolve the *Planck* beam by applying the low pass filter  $L \leq 4096$  and the Gaussian filter with  $\sigma = 2'$  to obtain a model for the observed signal  $\kappa^{\text{obs}}(\theta)$ .

To estimate the halo mass of contaminating galaxies, we measure the weak lensing signal around the LBGs with the HSC first-year shear catalog [33, 34]. This measurement enables us to pick up the lensing signal from contaminating galaxies, since most of the HSC source galaxies reside at  $z < 4$  and thus are insensitive to the real LBGs at  $z = 3.8$ . We first measure the excess surface mass density  $\Delta\Sigma(R)$ , following the procedure described in Mandelbaum *et al.* [33]. We then fit the signal with a model based on halo occupation distribution with off-centering, which is similar to the one described in More *et al.* [13], using the publicly-available Markov Chain Monte Carlo (MCMC) sampler called *emcee* [35]. We translate the best-fit model in  $\Delta\Sigma$  into convergence  $\kappa$  to obtain the contamination signal.

We then fit the convergence model to the signal varying the halo mass of LBGs. We use the signals at  $6' < \theta < 20'$  to remove small scales that can possibly suffer from residual signals at  $L > 2048$  [25]. We show the best-fit model in Fig. 1 together with the expected contamination signal based on the halo mass derived above. Note that the contamination level is much lower than the observed signal because of the insensitivity of CMB lensing at low redshift. In fact, the significance of the observed signal against the contamination signal is  $3.5\sigma$ . The LBG halo mass is constrained as  $M_h = 3.1_{-2.8}^{+9.3} \times 10^{11} h^{-1}M_{\odot}$  (68% credible level) [36]. Note that the signal is dominated by the 2-halo term, as expected from the fact that  $R_{200}$  corresponding to this halo mass is at  $\theta \sim 0.1'$ .

## DISCUSSIONS

A measurement of the matter correlation function can constrain cosmological parameters such as the matter energy density at present  $\Omega_{m0}$  and the amplitude of the matter power spectrum  $\sigma_8$ . Since galaxies are a biased tracer of the underlying matter distributions, we cannot directly measure the matter correlation function from the spatial distributions of galaxies (galaxy clustering). The matter correlation function, however, can be recovered by combining the galaxy clustering and galaxy lensing signal [37], because the clustering and lensing signal depend on

the bias in a different manner. In the limit of large scales where the galaxy density fluctuation follows the linear bias model  $\delta_g = b\delta_m$ , the clustering signal provides the galaxy-galaxy correlation function  $\xi_{gg} = b^2\xi_{mm}$ , while the lensing signal indicates the galaxy-matter correlation function  $\xi_{gm} = b\xi_{mm}$ .

Adding a projected clustering measurement of the LBG sample (Harikane et al, in prep.) to our lensing signal measurement, we demonstrate the first cosmological analysis through the growth of large-scale structure at  $z \sim 4$ . We measure the projected clustering signal and its jackknife covariance of the LBGs in the same manner as the previous measurements [7]. To ensure large-scale signals are used for cosmological constraints, we use the clustering signal at  $1.5' < \theta < 6.5'$  and lensing signal at  $6.0' < \theta < 20'$ , which corresponds to the comoving scale ranges of  $2.4 < R/[h^{-1}\text{Mpc}] < 9.4$  and  $8.7 < R/[h^{-1}\text{Mpc}] < 30$ , respectively, at  $z = 3.8$ .

We constrain the cosmological parameters  $\Omega_{m0}$  and  $\sigma_8$  by simultaneously fitting the large scale clustering and lensing signal, while the other cosmological parameters are fixed to the *Planck* cosmology [38]. We model the lensing signal in the same manner as the modeling procedures described in the previous section with the following modifications. For the convergence profile of LBGs, we use only the 2-halo term and vary  $b$  as a fitting parameter. We also vary  $\Omega_{m0}$  and  $A_s$  on which the linear power spectrum depends. Both 1-halo and 2-halo term in the lensing signal from contaminating galaxies are fixed. We model the observed clustering signal as  $\omega_{\text{tot}}(\theta) = (1 - f_{\text{cont}})^2\omega_{\text{LBG}}(\theta)$ , where  $\omega_{\text{LBG}}(\theta)$  is the clustering signal from LBGs. Note that we ignore the contamination signal for clustering signal because it is suppressed by  $f_{\text{cont}}^2 \sim 0.06$ . The projected clustering signal of LBGs is modeled as  $\omega_{\text{LBG}}(\theta) = \int dz N^2(z) (dr/dz)^{-1} \int dk (k/2\pi) b^2 P_{\text{mm}}(k, z; \Omega_{m0}, A_s) J_0(r(z)\theta k)$ , where  $N(z)$  is the normalized redshift distribution of galaxies taken from [22].

We perform an MCMC analysis with flat priors,  $\Omega_{m0} \in [0.01, 0.95]$ ,  $\ln 10^{10} A_s \in [0.1, 5.0]$ , and  $b \in [0.1, 30]$ . Fig. 2 shows constraints on these parameters for the combined lensing and clustering results, together with constraints based on the lensing or clustering results alone. Since  $\sigma_8$  and  $b$  degenerate in a different manner for lensing and clustering at small  $b$ , the combination of lensing and clustering probes improves the constraint on  $\sigma_8$ , yielding  $\sigma_8 = 0.48_{-0.22}^{+0.24}$  and  $b = 6.6_{-4.4}^{+4.9}$  (68% credible level). Although it is known that there is a strong degeneracy between  $\Omega_{m0}$  and  $\sigma_8$  in a combined analysis of lensing and clustering [e.g., 13, 15, 16], the degeneracy appears moderate in our results, due to the low signal-to-noise ratios of our measurements.

Fig. 3 compares  $\sigma_8$  constraints from our high- $z$  study with those from low- $z$  studies in the literature. Our constraint on  $f\sigma_8(z)$  is derived by converting our constraints on  $(\Omega_{m0}, \sigma_8)$  into  $f\sigma_8(z)$ , where  $f = -d \ln D(z)/d(1+z)$

is the logarithmic growth rate and  $\sigma_8(z) = \sigma_8 D(z)/D(0)$  is the linear matter fluctuation at redshift  $z$ . Our constraints on both  $\sigma_8$  and  $f\sigma_8(z)$  are roughly consistent with *Planck* results [38], while our constraints are lower than the *Planck* cosmology slightly over the  $1\sigma$  levels. Motivated by the potential deviations from the *Planck* constraints on  $\Lambda$ CDM cosmology we repeat the same analysis with the assumption that the time dependent dark energy is characterized by the equation-of-state parameter  $w(a) = w_0 + (1-a)w_a$ . We find that again our constraint is consistent with the *Planck* cosmology.

The HSC SSP survey will be completed by the end of 2021, which will cover  $\sim 1,400 \text{ deg}^2$  of the sky and provide the GOLDRUSH sample three times as large as the one used in this paper. The Advanced Atacama Cosmology Telescope Polarimeter (AdvACT) recently released the temperature and polarization maps much deeper than *Planck*, and will increase sensitivity in coming years [39, 40]. Combining these data sets, we can significantly improve the signal-to-noise ratios of the lensing and clustering signals. Moreover, in the near future, optical and near-infrared (O-NIR) programs of the Rubin Observatory Legacy Survey of Space and Time (LSST) [41] and the Nancy Grace Roman Space Telescope [42] will provide much larger and cleaner LBG samples by their deeper and redder imaging. Spectroscopic observations planned with Roman will enable us to evaluate completeness and purity of the LBG samples. On the CMB side, Simons Observatory [43] and CMB-S4 [44] are scheduled to be deployed at around the same time as the O-NIR surveys start. The combination of these data sets will allow us to extend precision cosmology studies via the structure growth from high ( $z \sim 4-6$ ) to low redshift ( $z < 1$ ) [45]. The structure growth measurement at such a high redshift may enable us to explore a non-standard cosmological model that can resolve the  $H_0$  tension such as an early dark energy model [18].

The Hyper Suprime-Cam (HSC) collaboration includes the astronomical communities of Japan and Taiwan, and Princeton University. The HSC instrumentation and software were developed by the National Astronomical Observatory of Japan (NAOJ), the Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), the University of Tokyo, the High Energy Accelerator Research Organization (KEK), the Academia Sinica Institute for Astronomy and Astrophysics in Taiwan (ASIAA), and Princeton University. Funding was contributed by the FIRST program from the Japanese Cabinet Office, the Ministry of Education, Culture, Sports, Science and Technology (MEXT), the Japan Society for the Promotion of Science (JSPS), Japan Science and Technology Agency (JST), the Toray Science Foundation, NAOJ, Kavli IPMU, KEK, ASIAA, and Princeton University.

This paper makes use of software developed for the




FIG. 2. Cosmological constraints from CMB lensing and galaxy-galaxy clustering measurements at  $z \sim 4$  (red contours). The blue (green) contours show constraints from the lensing (clustering) probe alone. The degeneracy between  $\sigma_8$  and  $b$  is reduced by the combination of these probes. These constraints are roughly consistent with the *Planck* cosmology [purple cross; 38].

Large Synoptic Survey Telescope. We thank the LSST Project for making their code available as free software at <http://dm.lsst.org>

This paper is based [in part] on data collected at the Subaru Telescope and retrieved from the HSC data archive system, which is operated by Subaru Telescope and Astronomy Data Center (ADC) at NAOJ. Data analysis was in part carried out with the cooperation of Center for Computational Astrophysics (CfCA), NAOJ.

The Pan-STARRS1 Surveys (PS1) and the PS1 public science archive have been made possible through contributions by the Institute for Astronomy, the University of Hawaii, the Pan-STARRS Project Office, the Max Planck Society and its participating institutes, the Max Planck Institute for Astronomy, Heidelberg, and the Max Planck Institute for Extraterrestrial Physics, Garching, The Johns Hopkins University, Durham University, the University of Edinburgh, the Queen’s University Belfast, the Harvard-Smithsonian Center for Astrophysics, the Las Cumbres Observatory Global Telescope Network Incorporated, the National Central University of Taiwan, the Space Telescope Science Institute, the National Aeronautics and Space Administration under grant No. NNX08AR22G issued through the Planetary Science Division of the NASA Science Mission Directorate, the National Science Foundation grant No. AST-1238877, the University of Maryland, Eotvos Lorand University (ELTE), the Los Alamos National Laboratory, and the Gordon and Betty Moore Foundation.

We thank Mathew Madhavacheril, Shuichiro Yokoyama, and Kiyotomo Ichiki for useful discussions. This work was supported in part by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan, and JSPS KAKENHI Grant Numbers JP15H02064, JP19H05100, and JP20H01932.




FIG. 3. *top*: Constraints on  $\sigma_8$  based on 2-point statistics from our CMB lensing and galaxy-galaxy clustering measurements at  $z \sim 4$  (red circle) along with low- $z$  measurements from galaxy surveys. The blue cross denotes the HSC cosmic shear measurement [46], while the green triangle (purple square) represents the combined analysis of cosmic shear, galaxy-galaxy lensing, and projected (redshift-space) galaxy-galaxy clustering conducted with DES (KiDS, 2DFLenS, and BOSS) data [15, 47]. The gray band shows constraints from *Planck* PR3 [38]. *bottom*: Our constraint on  $f\sigma_8(z)$  (red circle) that is obtained by converting the  $\Omega_{m0}$  and  $\sigma_8$  constraints into  $f\sigma_8(z)$ . The constraints at low- $z$  from redshift-space distortion measurements are indicated with the blue cross, green triangle, purple square, orange diamonds, yellow pentagons, brown hexagon, and magenta octagon that correspond to 6dFGRS [48], 2dFGRS [49], GAMA [50], WiggleZ [51], BOSS [52], VIPERS [53], and FastSound [54], respectively.

- 
- [1] R. S. Somerville and R. Davé, Physical Models of Galaxy Formation in a Cosmological Framework, *ARA&A* **53**, 51 (2015), arXiv:1412.2712 [astro-ph.GA].
  - [2] R. H. Wechsler and J. L. Tinker, The Connection Between Galaxies and Their Dark Matter Halos, *ARA&A* **56**, 435 (2018), arXiv:1804.03097 [astro-ph.GA].
  - [3] H. J. McCracken, M. Wolk, S. Colombi, M. Kilbinger, O. Ilbert, S. Peirani, J. Coupon, J. Dunlop, B. Milvang-Jensen, K. Caputi, H. Aussel, M. Béthermin, and O. Le Fèvre, Probing the galaxy-halo connection in UltraVISTA to  $z \sim 2$ , *Mon. Not. Roy. Astron. Soc.* **449**, 901

- (2015), arXiv:1411.4983 [astro-ph.CO].
- [4] P. W. Hatfield, S. N. Lindsay, M. J. Jarvis, B. Häußler, M. Vaccari, and A. Verma, The galaxy-halo connection in the VIDEO survey at  $0.5 < z < 1.7$ , *Mon. Not. Roy. Astron. Soc.* **459**, 2618 (2016), arXiv:1511.05476 [astro-ph.GA].
- [5] S. Ishikawa, N. Kashikawa, J. Toshikawa, M. Tanaka, T. Hamana, Y. Niino, K. Ichikawa, and H. Uchiyama, The Galaxy-Halo Connection in High-redshift Universe: Details and Evolution of Stellar-to-halo Mass Ratios of Lyman Break Galaxies on CFHTLS Deep Fields, *Astrophys. J.* **841**, 8 (2017), arXiv:1612.06869 [astro-ph.GA].
- [6] W. I. Cowley, K. I. Caputi, S. Deshmukh, M. L. N. Ashby, G. G. Fazio, O. Le Fèvre, J. P. U. Fynbo, O. Ilbert, H. J. McCracken, B. Milvang-Jensen, and R. S. Somerville, The Galaxy-Halo Connection for  $1.5 < z < 5$  as Revealed by the Spitzer Matching Survey of the UltraVISTA Ultra-deep Stripes, *Astrophys. J.* **853**, 69 (2018), arXiv:1712.03844 [astro-ph.GA].
- [7] Y. Harikane, M. Ouchi, Y. Ono, S. Saito, P. Behroozi, S. More, K. Shimasaku, J. Toshikawa, Y.-T. Lin, M. Akiyama, J. Coupon, Y. Komiyama, A. Konno, S.-C. Lin, S. Miyazaki, A. J. Nishizawa, T. Shibuya, and J. Silverman, GOLDRUSH. II. Clustering of galaxies at  $z \sim 4-6$  revealed with the half-million dropouts over the  $100 \text{ deg}^2$  area corresponding to  $1 \text{ Gpc}^3$ , *Publ. Astron. Soc. Japan* **70**, S11 (2018), arXiv:1704.06535 [astro-ph.GA].
- [8] A. Leauthaud, J. Tinker, K. Bundy, P. S. Behroozi, R. Massey, J. Rhodes, M. R. George, J.-P. Kneib, A. Benson, R. H. Wechsler, M. T. Busha, P. Capak, M. Cortês, O. Ilbert, A. M. Koekemoer, O. Le Fèvre, S. Lilly, H. J. McCracken, M. Salvato, T. Schrabback, N. Scoville, T. Smith, and J. E. Taylor, New Constraints on the Evolution of the Stellar-to-dark Matter Connection: A Combined Analysis of Galaxy-Galaxy Lensing, Clustering, and Stellar Mass Functions from  $z = 0.2$  to  $z = 1$ , *Astrophys. J.* **744**, 159 (2012), arXiv:1104.0928 [astro-ph.CO].
- [9] J. Coupon, S. Arnouts, L. van Waerbeke, T. Moutard, O. Ilbert, E. van Uitert, T. Erben, B. Garilli, L. Guzzo, C. Heymans, H. Hildebrandt, H. Hoekstra, M. Kilbinger, T. Kitching, Y. Mellier, L. Miller, M. Scodregio, C. Bonnett, E. Branchini, I. Davidzon, G. De Lucia, A. Fritz, L. Fu, P. Hudelot, M. J. Hudson, K. Kuijken, A. Leauthaud, O. Le Fèvre, H. J. McCracken, L. Moscardini, B. T. P. Rowe, T. Schrabback, E. Semboloni, and M. Velander, The galaxy-halo connection from a joint lensing, clustering and abundance analysis in the CFHTLenS/VIPERS field, *Mon. Not. Roy. Astron. Soc.* **449**, 1352 (2015), arXiv:1502.02867 [astro-ph.CO].
- [10] S. Huang, A. Leauthaud, A. Hearin, P. Behroozi, C. Bradshaw, F. Ardila, J. Speagle, A. Tenneti, K. Bundy, J. Greene, C. Sifón, and N. Bahcall, Weak lensing reveals a tight connection between dark matter halo mass and the distribution of stellar mass in massive galaxies, *Mon. Not. Roy. Astron. Soc.* **492**, 3685 (2020), arXiv:1811.01139 [astro-ph.GA].
- [11] E. N. Taylor, M. E. Cluver, A. Duffy, P. Gurri, H. Hoekstra, A. Sonnenfeld, M. N. Bremer, M. M. Brouwer, N. E. Chisari, A. Dvornik, T. Erben, H. Hildebrandt, A. M. Hopkins, L. S. Kelvin, S. Phillipps, A. S. G. Robotham, C. Sifón, M. Vakili, and A. H. Wright, GAMA+KiDS: Empirical correlations between halo mass and other galaxy properties near the knee of the stellar-to-halo mass relation, *Mon. Not. Roy. Astron. Soc.* 10.1093/mnras/staa2648 (2020), arXiv:2006.10040 [astro-ph.GA].
- [12] M. Cacciato, F. C. van den Bosch, S. More, H. Mo, and X. Yang, Cosmological constraints from a combination of galaxy clustering and lensing - III. Application to SDSS data, *Mon. Not. Roy. Astron. Soc.* **430**, 767 (2013), arXiv:1207.0503 [astro-ph.CO].
- [13] S. More, H. Miyatake, R. Mandelbaum, M. Takada, D. N. Spergel, J. R. Brownstein, and D. P. Schneider, The Weak Lensing Signal and the Clustering of BOSS Galaxies. II. Astrophysical and Cosmological Constraints, *Astrophys. J.* **806**, 2 (2015), arXiv:1407.1856 [astro-ph.CO].
- [14] R. Mandelbaum, A. Slosar, T. Baldauf, U. Seljak, C. M. Hirata, R. Nakajima, R. Reyes, and R. E. Smith, Cosmological parameter constraints from galaxy-galaxy lensing and galaxy clustering with the SDSS DR7, *Mon. Not. Roy. Astron. Soc.* **432**, 1544 (2013), arXiv:1207.1120 [astro-ph.CO].
- [15] T. M. C. Abbott, F. B. Abdalla, A. Alarcon, J. Aleksić, S. Allam, S. Allen, A. Amara, J. Annis, J. Asorey, S. Avila, D. Bacon, E. Balbinot, M. Banerji, N. Banik, W. Barkhouse, M. Baumer, E. Baxter, K. Bechtol, M. R. Becker, A. Benoit-Lévy, B. A. Benson, G. M. Bernstein, E. Bertin, J. Blazek, S. L. Bridle, D. Brooks, D. Brout, E. Buckley-Geer, D. L. Burke, M. T. Busha, A. Campos, D. Capozzi, A. Carnero Rosell, M. Carrasco Kind, J. Carretero, F. J. Castander, R. Cawthon, C. Chang, N. Chen, M. Childress, A. Choi, C. Conselice, R. Crittenden, M. Crocce, C. E. Cunha, C. B. D'Andrea, L. N. da Costa, R. Das, T. M. Davis, C. Davis, J. De Vicente, D. L. DePoy, J. DeRose, S. Desai, H. T. Diehl, J. P. Dietrich, S. Dodelson, P. Doel, A. Drlica-Wagner, T. F. Eifler, A. E. Elliott, F. Elsner, J. Elvin-Poole, J. Estrada, A. E. Evrard, Y. Fang, E. Fernandez, A. Ferté, D. A. Finley, B. Flaugher, P. Fosalba, O. Friedrich, J. Frieman, J. García-Bellido, M. Garcia-Fernandez, M. Gatti, E. Gaztanaga, D. W. Gerdes, T. Giannantonio, M. S. S. Gill, K. Glazebrook, D. A. Goldstein, D. Gruen, R. A. Gruendl, J. Gschwend, G. Gutierrez, S. Hamilton, W. G. Hartley, S. R. Hinton, K. Honscheid, B. Hoyle, D. Huterer, B. Jain, D. J. James, M. Jarvis, T. Jeltema, M. D. Johnson, M. W. G. Johnson, T. Kacprzak, S. Kent, A. G. Kim, A. King, D. Kirk, N. Kokron, A. Kovacs, E. Krause, C. Krawiec, A. Kremin, K. Kuehn, S. Kuhlmann, N. Kuropatkin, F. Lacasa, O. Lahav, T. S. Li, A. R. Liddle, C. Lidman, M. Lima, H. Lin, N. MacCrann, M. A. G. Maia, M. Makler, M. Manera, M. March, J. L. Marshall, P. Martini, R. G. McMahon, P. Melchior, F. Menanteau, R. Miquel, V. Miranda, D. Mudd, J. Muir, A. Möller, E. Neilsen, R. C. Nichol, B. Nord, P. Nugent, R. L. C. Ogando, A. Palmese, J. Peacock, H. V. Peiris, J. Peoples, W. J. Percival, D. Petravick, A. A. Plazas, A. Porredon, J. Prat, A. Pujol, M. M. Rau, A. Refregier, P. M. Ricker, N. Roe, R. P. Rollins, A. K. Romer, A. Roodman, R. Rosenfeld, A. J. Ross, E. Rozo, E. S. Rykoff, M. Sako, A. I. Salvador, S. Samuroff, C. Sánchez, E. Sanchez, B. Santiago, V. Scarpine, R. Schindler, D. Scolnic, L. F. Secco, S. Serrano, I. Sevilla-Noarbe, E. Sheldon, R. C. Smith, M. Smith, J. Smith, M. Soares-Santos, F. Sobreira, E. Suchyta, G. Tarle, D. Thomas, M. A. Troxel, D. L. Tucker, B. E. Tucker, S. A. Uddin, T. N.

- Varga, P. Vielzeuf, V. Vikram, A. K. Vivas, A. R. Walker, M. Wang, R. H. Wechsler, J. Weller, W. Wester, R. C. Wolf, B. Yanny, F. Yuan, A. Zenteno, B. Zhang, Y. Zhang, J. Zuntz, and Dark Energy Survey Collaboration, Dark Energy Survey year 1 results: Cosmological constraints from galaxy clustering and weak lensing, *Phys. Rev. D* **98**, 043526 (2018), arXiv:1708.01530 [astro-ph.CO].
- [16] C. Heymans, T. Tröster, M. Asgari, C. Blake, H. Hildebrandt, B. Joachimi, K. Kuijken, C.-A. Lin, A. G. Sánchez, J. L. van den Busch, A. H. Wright, A. Amon, M. Bilicki, J. de Jong, M. Crocce, A. Dvornik, T. Erben, M. C. Fortuna, F. Getman, B. Giblin, K. Glazebrook, H. Hoekstra, S. Joudaki, A. Kannawadi, F. Köhlinger, C. Lidman, L. Miller, N. R. Napolitano, D. Parkinson, P. Schneider, H. Shan, E. A. Valentijn, G. Verdoes Kleijn, and C. Wolf, KiDS-1000 Cosmology: Multi-probe weak gravitational lensing and spectroscopic galaxy clustering constraints, *Astronomy & Astrophysics* **646**, A140 (2021), arXiv:2007.15632 [astro-ph.CO].
- [17] A. G. Riess, S. Casertano, W. Yuan, L. M. Macri, and D. Scolnic, Large Magellanic Cloud Cepheid Standards Provide a 1% Foundation for the Determination of the Hubble Constant and Stronger Evidence for Physics beyond  $\Lambda$ CDM, *Astrophys. J.* **876**, 85 (2019), arXiv:1903.07603 [astro-ph.CO].
- [18] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F. Mota, A. G. Riess, and J. Silk, In the Realm of the Hubble tension – a Review of Solutions, arXiv e-prints, arXiv:2103.01183 (2021), arXiv:2103.01183 [astro-ph.CO].
- [19] H. Aihara, Y. AIsayyad, M. Ando, R. Armstrong, J. Bosch, E. Egami, H. Furusawa, J. Furusawa, A. Goulding, Y. Harikane, C. Hikage, P. T. P. Ho, B.-C. Hsieh, S. Huang, H. Ikeda, M. Imanishi, K. Ito, I. Iwata, A. T. Jaelani, R. Kakuma, K. Kawana, S. Kikuta, U. Kobayashi, M. Koike, Y. Komiyama, X. Li, Y. Liang, Y.-T. Lin, W. Luo, R. Lupton, N. B. Lust, L. A. MacArthur, Y. Matsuoka, S. Mineo, H. Miyatake, S. Miyazaki, S. More, R. Murata, S. V. Namiki, A. J. Nishizawa, M. Oguri, N. Okabe, S. Okamoto, Y. Okura, Y. Ono, M. Onodera, M. Onoue, K. Osato, M. Ouchi, T. Shibuya, M. A. Strauss, N. Sugiyama, Y. Suto, M. Takada, Y. Takagi, T. Takata, S. Takita, M. Tanaka, T. Terai, Y. Toba, H. Uchiyama, Y. Utsumi, S.-Y. Wang, W. Wang, and Y. Yamada, Second data release of the Hyper Suprime-Cam Subaru Strategic Program, *Publ. Astron. Soc. Japan* **71**, 114 (2019), arXiv:1905.12221 [astro-ph.IM].
- [20] J. Bosch, R. Armstrong, S. Bickerton, H. Furusawa, H. Ikeda, M. Koike, R. Lupton, S. Mineo, P. Price, T. Takata, M. Tanaka, N. Yasuda, Y. AIsayyad, A. C. Becker, W. Coulton, J. Coupon, J. Garmilla, S. Huang, K. S. Krughoff, D. Lang, A. Leauthaud, K.-T. Lim, N. B. Lust, L. A. MacArthur, R. Mandelbaum, H. Miyatake, S. Miyazaki, R. Murata, S. More, Y. Okura, R. Owen, J. D. Swinbank, M. A. Strauss, Y. Yamada, and H. Yamanai, The Hyper Suprime-Cam software pipeline, *Publ. Astron. Soc. Japan* **70**, S5 (2018), arXiv:1705.06766 [astro-ph.IM].
- [21] J. Bosch, Y. AIsayyad, R. Armstrong, E. Bellm, H.-F. Chiang, S. Eggl, K. Findeisen, M. Fisher-Levine, L. P. Guy, A. Guyonnet, Ž. Ivezić, T. Jenness, G. Kovács, K. S. Krughoff, R. H. Lupton, N. B. Lust, L. A. MacArthur, J. Meyers, F. Moolekamp, C. B. Morrison, T. D. Morton, W. O’Mullane, J. K. Parejko, A. A. Plazas, P. A. Price, M. L. Rawls, S. L. Reed, P. Schellart, C. T. Slater, I. Sullivan, J. D. Swinbank, D. Taranu, C. Z. Waters, and W. M. Wood-Vasey, An Overview of the LSST Image Processing Pipelines, in *Astronomical Data Analysis Software and Systems XXVII*, Astronomical Society of the Pacific Conference Series, Vol. 523, edited by P. J. Teuben, M. W. Pound, B. A. Thomas, and E. M. Warner (2019) p. 521, arXiv:1812.03248 [astro-ph.IM].
- [22] Y. Ono, M. Ouchi, Y. Harikane, J. Toshikawa, M. Rauch, S. Yuma, M. Sawicki, T. Shibuya, K. Shimasaku, M. Oguri, C. Willott, M. Akhlaghi, M. Akiyama, J. Coupon, N. Kashikawa, Y. Komiyama, A. Konno, L. Lin, Y. Matsuoka, S. Miyazaki, T. Nagao, K. Nakajima, J. Silverman, M. Tanaka, Y. Taniguchi, and S.-Y. Wang, Great Optically Luminous Dropout Research Using Subaru HSC (GOLDRUSH). I. UV luminosity functions at  $z \sim 4-7$  derived with the half-million dropouts on the 100 deg<sup>2</sup> sky, *Publ. Astron. Soc. Japan* **70**, S10 (2018), arXiv:1704.06004 [astro-ph.GA].
- [23] J. Toshikawa, H. Uchiyama, N. Kashikawa, M. Ouchi, R. Overzier, Y. Ono, Y. Harikane, S. Ishikawa, T. Kodama, Y. Matsuda, Y.-T. Lin, M. Onoue, M. Tanaka, T. Nagao, M. Akiyama, Y. Komiyama, T. Goto, and C.-H. Lee, GOLDRUSH. III. A systematic search for proto-clusters at  $z \sim 4$  based on the  $\sim 100$  deg<sup>2</sup> area, *Publ. Astron. Soc. Japan* **70**, S12 (2018), arXiv:1708.09421 [astro-ph.GA].
- [24] H. Aihara, R. Armstrong, S. Bickerton, J. Bosch, J. Coupon, H. Furusawa, Y. Hayashi, H. Ikeda, Y. Kamata, H. Karoji, S. Kawanomoto, M. Koike, Y. Komiyama, D. Lang, R. H. Lupton, S. Mineo, H. Miyatake, S. Miyazaki, T. Morokuma, Y. Obuchi, Y. Oishi, Y. Okura, P. A. Price, T. Takata, M. M. Tanaka, M. Tanaka, Y. Tanaka, T. Uchida, F. Uraguchi, Y. Utsumi, S.-Y. Wang, Y. Yamada, H. Yamanai, N. Yasuda, N. Arimoto, M. Chiba, F. Finet, H. Fujimori, S. Fujimoto, J. Furusawa, T. Goto, A. Goulding, J. E. Gunn, Y. Harikane, T. Hattori, M. Hayashi, K. G. Helminiak, R. Higuchi, C. Hikage, P. T. P. Ho, B.-C. Hsieh, K. Huang, S. Huang, M. Imanishi, I. Iwata, A. T. Jaelani, H.-Y. Jian, N. Kashikawa, N. Katayama, T. Kojima, A. Konno, S. Koshida, H. Kusakabe, A. Leauthaud, C.-H. Lee, L. Lin, Y.-T. Lin, R. Mandelbaum, Y. Matsuoka, E. Medezinski, S. Miyama, R. Momose, A. More, S. More, S. Mukae, R. Murata, H. Murayama, T. Nagao, F. Nakata, M. Niida, H. Niikura, A. J. Nishizawa, M. Oguri, N. Okabe, Y. Ono, M. Onodera, M. Onoue, M. Ouchi, T.-S. Pyo, T. Shibuya, K. Shimasaku, M. Simet, J. Speagle, D. N. Spergel, M. A. Strauss, Y. Sugahara, N. Sugiyama, Y. Suto, N. Suzuki, P. J. Tait, M. Takada, T. Terai, Y. Toba, E. L. Turner, H. Uchiyama, K. Umetsu, Y. Urata, T. Usuda, S. Yeh, and S. Yuma, First data release of the Hyper Suprime-Cam Subaru Strategic Program, *Publ. Astron. Soc. Japan* **70**, S8 (2018), arXiv:1702.08449 [astro-ph.IM].
- [25] Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, R. B. Barreiro, N. Bartolo, S. Basak, K. Benabed, J. P. Bernard, M. Bersanelli, P. Bielewicz, J. J. Bock, J. R. Bond, J. Borrill, F. R. Bouchet, F. Boulanger, M. Bucher, C. Burigana, E. Calabrese, J. F. Cardoso,



- J. Carron, A. Challinor, H. C. Chiang, L. P. L. Colombo, C. Combet, B. P. Crill, F. Cuttaia, P. de Bernardis, G. de Zotti, J. Delabrouille, E. Di Valentino, J. M. Diego, O. Doré, M. Douspis, A. Ducout, X. Dupac, G. Efstathiou, F. Elsner, T. A. Enßlin, H. K. Eriksen, Y. Fantaye, R. Fernandez-Cobos, F. Forastieri, M. Frailis, A. A. Fraisse, E. Franceschi, A. Frolov, S. Galeotta, S. Galli, K. Ganga, R. T. Génova-Santos, M. Gerbino, T. Ghosh, J. González-Nuevo, K. M. Górski, S. Gratton, A. Gruppuso, J. E. Gudmundsson, J. Hamann, W. Handley, F. K. Hansen, D. Herranz, E. Hivon, Z. Huang, A. H. Jaffe, W. C. Jones, A. Karakci, E. Keihänen, R. Keskitalo, K. Kiiveri, J. Kim, L. Knox, N. Krachmalnicoff, M. Kunz, H. Kurki-Suonio, G. Lagache, J. M. Lamarre, A. Lasenby, M. Lattanzi, C. R. Lawrence, M. Le Jeune, F. Levrier, A. Lewis, M. Liguori, P. B. Lilje, V. Lindholm, M. López-Cañiego, P. M. Lubin, Y. Z. Ma, J. F. Macías-Pérez, G. Maggio, D. Maino, N. Mandolesi, A. Mangilli, A. Marcos-Caballero, M. Maris, P. G. Martin, E. Martínez-González, S. Matarrese, N. Mauri, J. D. McEwen, A. Melchiorri, A. Mennella, M. Migliaccio, M. A. Miville-Deschênes, D. Molinari, A. Moneti, L. Montier, G. Morgante, A. Moss, P. Natoli, L. Pagano, D. Paoletti, B. Partridge, G. Patanchon, F. Perrotta, V. Pettorino, F. Piacentini, L. Polastri, G. Polenta, J. L. Puget, J. P. Rachen, M. Reinecke, M. Remazeilles, A. Renzi, G. Rocha, C. Rosset, G. Roudier, J. A. Rubiño-Martín, B. Ruiz-Granados, L. Salvati, M. Sandri, M. Savelainen, D. Scott, C. Sirignano, R. Sunyaev, A. S. Suur-Uski, J. A. Tauber, D. Tavagnacco, M. Tenti, L. Toffolatti, M. Tomasi, T. Trombetti, J. Valiviita, B. Van Tent, P. Vielva, F. Villa, N. Vittorio, B. D. Wandelt, I. K. Wehus, M. White, S. D. M. White, A. Zachei, and A. Zonca, Planck 2018 results. VIII. Gravitational lensing, arXiv e-prints , arXiv:1807.06210 (2018), arXiv:1807.06210 [astro-ph.CO].
- [26] <https://pla.esac.esa.int>.
- [27] J. F. Navarro, C. S. Frenk, and S. D. M. White, The Structure of Cold Dark Matter Halos, *Astrophys. J.* **462**, 563 (1996), arXiv:astro-ph/9508025 [astro-ph].
- [28] J. F. Navarro, C. S. Frenk, and S. D. M. White, A Universal Density Profile from Hierarchical Clustering, *Astrophys. J.* **490**, 493 (1997), arXiv:astro-ph/9611107 [astro-ph].
- [29] M. Takada and B. Jain, The three-point correlation function in cosmology, *Mon. Not. Roy. Astron. Soc.* **340**, 580 (2003), arXiv:astro-ph/0209167 [astro-ph].
- [30] A. R. Duffy, J. Schaye, S. T. Kay, and C. Dalla Vecchia, Dark matter halo concentrations in the Wilkinson Microwave Anisotropy Probe year 5 cosmology, *Mon. Not. Roy. Astron. Soc.* **390**, L64 (2008), arXiv:0804.2486 [astro-ph].
- [31] J. L. Tinker, B. E. Robertson, A. V. Kravtsov, A. Klypin, M. S. Warren, G. Yepes, and S. Gottlöber, The Large-scale Bias of Dark Matter Halos: Numerical Calibration and Model Tests, *Astrophys. J.* **724**, 878 (2010), arXiv:1001.3162 [astro-ph.CO].
- [32] A. Lewis and S. Bridle, Cosmological parameters from CMB and other data: A Monte Carlo approach, *Phys. Rev. D* **66**, 103511 (2002), arXiv:astro-ph/0205436 [astro-ph].
- [33] R. Mandelbaum, H. Miyatake, T. Hamana, M. Oguri, M. Simet, R. Armstrong, J. Bosch, R. Murata, F. Lanusse, A. Leauthaud, J. Coupon, S. More, M. Takada, S. Miyazaki, J. S. Speagle, M. Shirasaki, C. Sifón, S. Huang, A. J. Nishizawa, E. Medezinski, Y. Okura, N. Okabe, N. Czakon, R. Takahashi, W. R. Coulton, C. Hikage, Y. Komiyama, R. H. Lupton, M. A. Strauss, M. Tanaka, and Y. Utsumi, The first-year shear catalog of the Subaru Hyper Suprime-Cam Subaru Strategic Program Survey, *Publ. Astron. Soc. Japan* **70**, S25 (2018), arXiv:1705.06745 [astro-ph.CO].
- [34] R. Mandelbaum, F. Lanusse, A. Leauthaud, R. Armstrong, M. Simet, H. Miyatake, J. E. Meyers, J. Bosch, R. Murata, S. Miyazaki, and M. Tanaka, Weak lensing shear calibration with simulations of the HSC survey, *Mon. Not. Roy. Astron. Soc.* **481**, 3170 (2018), arXiv:1710.00885 [astro-ph.CO].
- [35] D. Foreman-Mackey, D. W. Hogg, D. Lang, and J. Goodman, emcee: The MCMC Hammer, *PASP* **125**, 306 (2013), arXiv:1202.3665 [astro-ph.IM].
- [36] Throughout this paper, we adopt the mode of the posterior distribution to infer the central value of the parameter, and the highest density interval of the posterior to infer the credible interval of the parameter. We use the marginalized posterior distribution when there are multiple fitting parameters.
- [37] T. Baldauf, R. E. Smith, U. Seljak, and R. Mandelbaum, Algorithm for the direct reconstruction of the dark matter correlation function from weak lensing and galaxy clustering, *Phys. Rev. D* **81**, 063531 (2010), arXiv:0911.4973 [astro-ph.CO].
- [38] Planck Collaboration, N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini, A. J. Banday, R. B. Barreiro, N. Bartolo, S. Basak, R. Battye, K. Benabed, J. P. Bernard, M. Bersanelli, P. Bielewicz, J. J. Bock, J. R. Bond, J. Borrill, F. R. Bouchet, F. Boulanger, M. Bucher, C. Burigana, R. C. Butler, E. Calabrese, J. F. Cardoso, J. Carron, A. Challinor, H. C. Chiang, J. Chluba, L. P. L. Colombo, C. Combet, D. Contreras, B. P. Crill, F. Cuttaia, P. de Bernardis, G. de Zotti, J. Delabrouille, J. M. Delouis, E. Di Valentino, J. M. Diego, O. Doré, M. Douspis, A. Ducout, X. Dupac, S. Dusini, G. Efstathiou, F. Elsner, T. A. Enßlin, H. K. Eriksen, Y. Fantaye, M. Farhang, J. Ferguson, R. Fernandez-Cobos, F. Finelli, F. Forastieri, M. Frailis, A. A. Fraisse, E. Franceschi, A. Frolov, S. Galeotta, S. Galli, K. Ganga, R. T. Génova-Santos, M. Gerbino, T. Ghosh, J. González-Nuevo, K. M. Górski, S. Gratton, A. Gruppuso, J. E. Gudmundsson, J. Hamann, W. Handley, F. K. Hansen, D. Herranz, S. R. Hildebrandt, E. Hivon, Z. Huang, A. H. Jaffe, W. C. Jones, A. Karakci, E. Keihänen, R. Keskitalo, K. Kiiveri, J. Kim, T. S. Kisner, L. Knox, N. Krachmalnicoff, M. Kunz, H. Kurki-Suonio, G. Lagache, J. M. Lamarre, A. Lasenby, M. Lattanzi, C. R. Lawrence, M. Le Jeune, P. Lemos, J. Lesgourgues, F. Levrier, A. Lewis, M. Liguori, P. B. Lilje, M. Lilley, V. Lindholm, M. López-Cañiego, P. M. Lubin, Y. Z. Ma, J. F. Macías-Pérez, G. Maggio, D. Maino, N. Mandolesi, A. Mangilli, A. Marcos-Caballero, M. Maris, P. G. Martin, M. Martinelli, E. Martínez-González, S. Matarrese, N. Mauri, J. D. McEwen, P. R. Meinhold, A. Melchiorri, A. Mennella, M. Migliaccio, M. Millea, S. Mitra, M. A. Miville-Deschênes, D. Molinari, L. Montier, G. Morgante, A. Moss, P. Natoli, H. U. Nørgaard-Nielsen, L. Pagano, D. Paoletti, B. Partridge, G. Patanchon, H. V. Peiris, F. Perrotta, V. Pettorino,



- F. Piacentini, L. Polastri, G. Polenta, J. L. Puget, J. P. Rachen, M. Reinecke, M. Remazeilles, A. Renzi, G. Rocha, C. Rosset, G. Roudier, J. A. Rubiño-Martín, B. Ruiz-Granados, L. Salvati, M. Sandri, M. Savelainen, D. Scott, E. P. S. Shellard, C. Sirignano, G. Sirri, L. D. Spencer, R. Sunyaev, A. S. Suur-Uski, J. A. Tauber, D. Tavagnacco, M. Tenti, L. Toffolatti, M. Tomasi, T. Trombetti, L. Valenziano, J. Valiviita, B. Van Tent, L. Vibert, P. Vielva, F. Villa, N. Vittorio, B. D. Wandelt, I. K. Wehus, M. White, S. D. M. White, A. Zachei, and A. Zonca, Planck 2018 results. VI. Cosmological parameters, *Astronomy & Astrophysics* **641**, A6 (2020), arXiv:1807.06209 [astro-ph.CO].
- [39] S. Aiola, E. Calabrese, L. Maurin, S. Naess, B. L. Schmitt, M. H. Abitbol, G. E. Addison, P. A. R. Ade, D. Alonso, M. Amiri, S. Amodeo, E. Angile, J. E. Austermann, T. Baidon, N. Battaglia, J. A. Beall, R. Bean, D. T. Becker, J. R. Bond, S. M. Bruno, V. Calafut, L. E. Campusano, F. Carrero, G. E. Chesmore, H.-m. Cho., S. K. Choi, S. E. Clark, N. F. Cothard, D. Crichton, K. T. Crowley, O. Darwish, R. Datta, E. V. Denison, M. J. Devlin, C. J. Duell, S. M. Duff, A. J. Duivenvoorden, J. Dunkley, R. o. Dünner, T. Essinger-Hileman, M. Fankhanel, S. Ferraro, A. E. Fox, B. Fuzia, P. A. Gallardo, V. Gluscevic, J. E. Golec, E. Grace, M. Gralla, Y. Guan, K. Hall, M. Halpern, D. Han, P. Hargrave, M. Hasselfield, J. M. Helton, S. Henderson, B. Hensley, J. C. Hill, G. C. Hilton, M. Hilton, A. D. Hincks, R. Hložek, S.-P. P. Ho, J. Hubmayr, K. M. Huffenberger, J. P. Hughes, L. Infante, K. Irwin, R. Jackson, J. Klein, K. Knowles, B. Koopman, A. Kosowsky, V. Lakey, D. Li, Y. Li, Z. Li, M. Lokken, T. Louis, M. Lungu, A. MacInnis, M. Madhavacheril, F. Maldonado, M. Mallaby-Kay, D. Marsden, J. McMahon, F. Menanteau, K. Moodley, T. Morton, T. Namikawa, F. Nati, L. Newburgh, J. P. Nibarger, A. Nicola, M. D. Niemack, M. R. Nolta, J. Orlowski-Sherer, L. A. Page, C. G. Pappas, B. Partridge, P. Phakathi, H. Prince, R. Puddu, F. J. Qu, J. Rivera, N. Robertson, F. Rojas, M. Salatino, E. Schaan, A. Schillaci, N. Sehgal, B. D. Sherwin, C. Sierra, J. Sievers, C. Sifon, P. Sikhosana, S. Simon, D. N. Spergel, S. T. Staggs, J. Stevens, E. Storer, D. D. Sunder, E. R. Switzer, B. Thorne, R. Thornton, H. Trac, J. Treu, C. Tucker, L. R. Vale, A. Van Engelen, J. Van Lanen, E. M. Vavagiakis, K. Wagoner, Y. Wang, J. T. Ward, E. J. Wollack, Z. Xu, F. Zago, and N. Zhu, The Atacama Cosmology Telescope: DR4 Maps and Cosmological Parameters, arXiv e-prints , arXiv:2007.07288 (2020), arXiv:2007.07288 [astro-ph.CO].
- [40] O. Darwish, M. S. Madhavacheril, B. Sherwin, S. Aiola, N. Battaglia, J. A. Beall, D. T. Becker, J. R. Bond, E. Calabrese, S. Choi, M. J. Devlin, J. Dunkley, R. Dünner, S. Ferraro, A. E. Fox, P. A. Gallardo, Y. Guan, M. Halpern, D. Han, M. Hasselfield, J. C. Hill, G. C. Hilton, M. Hilton, A. D. Hincks, S.-P. P. Ho, J. Hubmayr, J. P. Hughes, B. J. Koopman, A. Kosowsky, J. Van Lanen, T. Louis, M. Lungu, A. MacInnis, L. Maurin, J. McMahon, K. Moodley, S. Naess, T. Namikawa, L. Newburgh, J. P. Nibarger, M. D. Niemack, L. A. Page, B. Partridge, F. J. Qu, N. Robertson, B. Schmitt, N. Sehgal, C. Sifon, D. N. Spergel, S. Staggs, E. Storer, A. van Engelen, and E. J. Wollack, The Atacama Cosmology Telescope: A CMB lensing mass map over 2100 square degrees of sky and its cross-correlation with BOSS-CMASS galaxies, arXiv e-prints , arXiv:2004.01139 (2020), arXiv:2004.01139 [astro-ph.CO].
- [41] LSST Science Collaboration, P. A. Abell, J. Allison, S. F. Anderson, J. R. Andrew, J. R. P. Angel, L. Armus, D. Arnett, S. J. Asztalos, T. S. Axelrod, S. Bailey, D. R. Ballantyne, J. R. Bankert, W. A. Barkhouse, J. D. Barr, L. F. Barrientos, A. J. Barth, J. G. Bartlett, A. C. Becker, J. Becla, T. C. Beers, J. P. Bernstein, R. Biswas, M. R. Blanton, J. S. Bloom, J. J. Bochanski, P. Boeshaar, K. D. Borne, M. Bradac, W. N. Brandt, C. R. Bridge, M. E. Brown, R. J. Brunner, J. S. Bullock, A. J. Burgasser, J. H. Burge, D. L. Burke, P. A. Cargile, S. Chandrasekharan, G. Chartas, S. R. Chesley, Y.-H. Chu, D. Cinabro, M. W. Claire, C. F. Claver, D. Clowe, A. J. Connolly, K. H. Cook, J. Cooke, A. Cooray, K. R. Covey, C. S. Culliton, R. de Jong, W. H. de Vries, V. P. Debattista, F. Delgado, I. P. Dell’Antonio, S. Dhital, R. Di Stefano, M. Dickinson, B. Dilday, S. G. Djorgovski, G. Dobler, C. Donalek, G. Dubois-Felsmann, J. Durech, A. Eliasdottir, M. Eracleous, L. Eyer, E. E. Falco, X. Fan, C. D. Fassnacht, H. C. Ferguson, Y. R. Fernandez, B. D. Fields, D. Finkbeiner, E. E. Figuera, D. B. Fox, H. Francke, J. S. Frank, J. Frieman, S. Fromenteau, M. Furqan, G. Galaz, A. Gal-Yam, P. Garnavich, E. Gawiser, J. Geary, P. Gee, R. R. Gibson, K. Gilmore, E. A. Grace, R. F. Green, W. J. Gressler, C. J. Grillmair, S. Habib, J. S. Haggerty, M. Hamuy, A. W. Harris, S. L. Hawley, A. F. Heavens, L. Hebb, T. J. Henry, E. Hileman, E. J. Hilton, K. Hoadley, J. B. Holberg, M. J. Holman, S. B. Howell, L. Infante, Z. Ivezic, S. H. Jacoby, B. Jain, R. Jedicke, M. J. Jee, J. Garrett Jernigan, S. W. Jha, K. V. Johnston, R. L. Jones, M. Juric, M. Kaasalainen, Styliani, Kafka, S. M. Kahn, N. A. Kaib, J. Kalirai, J. Kantor, M. M. Kasliwal, C. R. Keeton, R. Kessler, Z. Knezevic, A. Kowalski, V. L. Krabbendam, K. S. Krughoff, S. Kulkarni, S. Kuhlman, M. Lacy, S. Lepine, M. Liang, A. Lien, P. Lira, K. S. Long, S. Lorenz, J. M. Lotz, R. H. Lupton, J. Lutz, L. M. Macri, A. A. Mahabal, R. Mandelbaum, P. Marshall, M. May, P. M. McGehee, B. T. Meadows, A. Meert, A. Milani, C. J. Miller, M. Miller, D. Mills, D. Minniti, D. Monet, A. S. Mukadam, E. Nakar, D. R. Neill, J. A. Newman, S. Nikolaev, M. Nordby, P. O’Connor, M. Oguri, J. Oliver, S. S. Olivier, J. K. Olsen, K. Olsen, E. W. Olszewski, H. Oluseyi, N. D. Padilla, A. Parker, J. Pepper, J. R. Peterson, C. Petry, P. A. Pinto, J. L. Pizagno, B. Popescu, A. Prsa, V. Radcka, M. J. Radford, A. Rasmussen, A. Rau, J. Rho, J. E. Rhoads, G. T. Richards, S. T. Ridgway, B. E. Robertson, R. Roskar, A. Saha, A. Sarajedini, E. Scannapieco, T. Schalk, R. Schindler, S. Schmidt, S. Schmidt, D. P. Schneider, G. Schumacher, R. Scranton, J. Sebag, L. G. Seppala, O. Shemmer, J. D. Simon, M. Sivertz, H. A. Smith, J. Allyn Smith, N. Smith, A. H. Spitz, A. Stanford, K. G. Stassun, J. Strader, M. A. Strauss, C. W. Stubbs, D. W. Sweeney, A. Szalay, P. Szkody, M. Takada, P. Thorman, D. E. Trilling, V. Trimble, A. Tyson, R. Van Berg, D. Vand en Berk, J. VanderPlas, L. Verde, B. Vrtnak, L. M. Walkowicz, B. D. Wandelt, S. Wang, Y. Wang, M. Warner, R. H. Wechsler, A. A. West, O. Wiecha, B. F. Williams, B. Willman, D. Wittman, S. C. Wolff, W. M. Wood-Vasey, P. Wozniak, P. Young, A. Zentner, and H. Zhan, LSST Science Book, Version 2.0, arXiv e-prints , arXiv:0912.0201 (2009), arXiv:0912.0201 [astro-

- ph.IM].
- [42] D. Spergel, N. Gehrels, C. Baltay, D. Bennett, J. Breckinridge, M. Donahue, A. Dressler, B. S. Gaudi, T. Greene, O. Guyon, C. Hirata, J. Kalirai, N. J. Kasdin, B. Macintosh, W. Moos, S. Perlmutter, M. Postman, B. Rauscher, J. Rhodes, Y. Wang, D. Weinberg, D. Benford, M. Hudson, W. S. Jeong, Y. Mellier, W. Traub, T. Yamada, P. Capak, J. Colbert, D. Masters, M. Penny, D. Savransky, D. Stern, N. Zimmerman, R. Barry, L. Bartusek, K. Carpenter, E. Cheng, D. Content, F. Dekens, R. Demers, K. Grady, C. Jackson, G. Kuan, J. Kruk, M. Melton, B. Nemati, B. Parvin, I. Poberezhskiy, C. Peddie, J. Ruffa, J. K. Wallace, A. Whipple, E. Wollack, and F. Zhao, Wide-Field Infrared Survey Telescope-Astrophysics Focused Telescope Assets WFIRST-AFTA 2015 Report, arXiv e-prints, arXiv:1503.03757 (2015), arXiv:1503.03757 [astro-ph.IM].
- [43] P. Ade, J. Aguirre, Z. Ahmed, S. Aiola, A. Ali, D. Alonso, M. A. Alvarez, K. Arnold, P. Ashton, J. Austermann, H. Awan, C. Baccigalupi, T. Baildon, D. Barron, N. Battaglia, R. Battye, E. Baxter, A. Bazarko, J. A. Beall, R. Bean, D. Beck, S. Beckman, B. Beringue, F. Bianchini, S. Boada, D. Boettger, J. R. Bond, J. Borrill, M. L. Brown, S. M. Bruno, S. Bryan, E. Calabrese, V. Calafut, P. Calisse, J. Carron, A. Challinor, G. Chesmore, Y. Chinone, J. Chluba, H.-M. S. Cho, S. Choi, G. Coppi, N. F. Cothard, K. Coughlin, D. Crichton, K. D. Crowley, K. T. Crowley, A. Cukierman, J. M. D’Ewart, R. Dünner, T. de Haan, M. Devlin, S. Dicker, J. Didier, M. Dobbs, B. Dober, C. J. Duell, S. Duff, A. Duivenvoorden, J. Dunkley, J. Dusatko, J. Errard, G. Fabbian, S. Feeney, S. Ferraro, P. Fluxà, K. Freese, J. C. Frisch, A. Frolov, G. Fuller, B. Fuzia, N. Galitzki, P. A. Gallardo, J. Tomas Galvez Gherzi, J. Gao, E. Gawiser, M. Gerbino, V. Gluscevic, N. Goeckner-Wald, J. Golec, S. Gordon, M. Gralla, D. Green, A. Grigorian, J. Groh, C. Groppi, Y. Guan, J. E. Gudmundsson, D. Han, P. Hargrave, M. Hasegawa, M. Hasselfield, M. Hattori, V. Haynes, M. Hazumi, Y. He, E. Healy, S. W. Henderson, C. Hervias-Caimapo, C. A. Hill, J. C. Hill, G. Hilton, M. Hilton, A. D. Hincks, G. Hinshaw, R. Hložek, S. Ho, S.-P. P. Ho, L. Howe, Z. Huang, J. Hubmayr, K. Huffenberger, J. P. Hughes, A. Ijjas, M. Ikape, K. Irwin, A. H. Jaffe, B. Jain, O. Jeong, D. Kaneko, E. D. Karpel, N. Katayama, B. Keating, S. S. Kernasovskiy, R. Keskitalo, T. Kisner, K. Kiuchi, J. Klein, K. Knowles, B. Koopman, A. Kosowsky, N. Krachmalnicoff, S. E. Kuenstner, C.-L. Kuo, A. Kusaka, J. Lashner, A. Lee, E. Lee, D. Leon, J. S. Y. Leung, A. Lewis, Y. Li, Z. Li, M. Limon, E. Linder, C. Lopez-Caraballo, T. Louis, L. Lowry, M. Lungu, M. Madhavacheril, D. Mak, F. Maldonado, H. Mani, B. Mates, F. Matsuda, L. Maurin, P. Mausekopf, A. May, N. McCallum, C. McKenney, J. McMahon, P. D. Meerburg, J. Meyers, A. Miller, M. Mirmelstein, K. Moodley, M. Munchmeyer, C. Munson, S. Naess, F. Nati, M. Navaroli, L. Newburgh, H. N. Nguyen, M. Niemack, H. Nishino, J. Orłowski-Scherer, L. Page, B. Partridge, J. Peloton, F. Perrotta, L. Piccirillo, G. Pisano, D. Poletti, R. Puddu, G. Puglisi, C. Raum, C. L. Reichardt, M. Remazeilles, Y. Rephaeli, D. Riechers, F. Rojas, A. Roy, S. Sadeh, Y. Sakurai, M. Salatino, M. Sathyanarayana Rao, E. Schaan, M. Schmittfull, N. Sehgal, J. Seibert, U. Seljak, B. Sherwin, M. Shimon, C. Sierra, J. Sievers, P. Sikhosana, M. Silva-Feaver, S. M. Simon, A. Sinclair, P. Siritanasak, K. Smith, S. R. Smith, D. Spergel, S. T. Staggs, G. Stein, J. R. Stevens, R. Stompor, A. Suzuki, O. Tajima, S. Takakura, G. Teply, D. B. Thomas, B. Thorne, R. Thornton, H. Trac, C. Tsai, C. Tucker, J. Ullom, S. Vagnozzi, A. van Engelen, J. Van Lanen, D. D. Van Winkle, E. M. Vavagiakis, C. Vergès, M. Vissers, K. Wagoner, S. Walker, J. Ward, B. Westbrook, N. Whitehorn, J. Williams, J. Williams, E. J. Wollack, Z. Xu, B. Yu, C. Yu, F. Zago, H. Zhang, N. Zhu, and Simons Observatory Collaboration, The Simons Observatory: science goals and forecasts, JCAP **2019**, 056 (2019), arXiv:1808.07445 [astro-ph.CO].
- [44] K. Abazajian, G. Addison, P. Adshead, Z. Ahmed, S. W. Allen, D. Alonso, M. Alvarez, A. Anderson, K. S. Arnold, C. Baccigalupi, K. Bailey, D. Barkats, D. Barron, P. S. Barry, J. G. Bartlett, R. Basu Thakur, N. Battaglia, E. Baxter, R. Bean, C. Bebek, A. N. Bender, B. A. Benson, E. Berger, S. Bhimani, C. A. Bischoff, L. Bleem, S. Bocquet, K. Boddy, M. Bonato, J. R. Bond, J. Borrill, F. R. Bouchet, M. L. Brown, S. Bryan, B. Burkhart, V. Buza, K. Byrum, E. Calabrese, V. Calafut, R. Caldwell, J. E. Carlstrom, J. Carron, T. Cecil, A. Challinor, C. L. Chang, Y. Chinone, H.-M. S. Cho, A. Cooray, T. M. Crawford, A. Crites, A. Cukierman, F.-Y. Cyr-Racine, T. de Haan, G. de Zotti, J. Delabrouille, M. Demarteau, M. Devlin, E. Di Valentino, M. Dobbs, S. Duff, A. Duivenvoorden, C. Dvorkin, W. Edwards, J. Eimer, J. Errard, T. Essinger-Hileman, G. Fabbian, C. Feng, S. Ferraro, J. P. Filippini, R. Flauger, B. Flaugher, A. A. Fraise, A. Frolov, N. Galitzki, S. Galli, K. Ganga, M. Gerbino, M. Gilchriese, V. Gluscevic, D. Green, D. Grin, E. Grohs, R. Gualtieri, V. Guarino, J. E. Gudmundsson, S. Habib, G. Haller, M. Halpern, N. W. Halverson, S. Hanany, K. Harrington, M. Hasegawa, M. Hasselfield, M. Hazumi, K. Heitmann, S. Henderson, J. W. Henning, J. C. Hill, R. Hložek, G. Holder, W. Holzappel, J. Hubmayr, K. M. Huffenberger, M. Huffer, H. Hui, K. Irwin, B. R. Johnson, D. Johnstone, W. C. Jones, K. Karkare, N. Katayama, J. Kerby, S. Kernovsky, R. Keskitalo, T. Kisner, L. Knox, A. Kosowsky, J. Kovac, E. D. Kovetz, S. Kuhlmann, C.-l. Kuo, N. Kurita, A. Kusaka, A. Lahteenmaki, C. R. Lawrence, A. T. Lee, A. Lewis, D. Li, E. Linder, M. Loverde, A. Lowitz, M. S. Madhavacheril, A. Mantz, F. Matsuda, P. Mausekopf, J. McMahon, M. McQuinn, P. D. Meerburg, J.-B. Melin, J. Meyers, M. Millea, J. Mohr, L. Moncelsi, T. Mroczkowski, S. Mukherjee, M. Münchmeyer, D. Nagai, J. Nagy, T. Namikawa, F. Nati, T. Natoli, M. Negrello, L. Newburgh, M. D. Niemack, H. Nishino, M. Nordby, V. Novosad, P. O’Connor, G. Obied, S. Padin, S. Pandey, B. Partridge, E. Pierpaoli, L. Pogosian, C. Pryke, G. Puglisi, B. Racine, S. Raghunathan, A. Rahlin, S. Rajagopalan, M. Raveri, M. Reichanadter, C. L. Reichardt, M. Remazeilles, G. Rocha, N. A. Roe, A. Roy, J. Ruhl, M. Salatino, B. Salivanchik, E. Schaan, A. Schillaci, M. M. Schmittfull, D. Scott, N. Sehgal, S. Shandera, C. Sheehy, B. D. Sherwin, E. Shirokoff, S. M. Simon, A. Slosar, R. Somerville, D. Spergel, S. T. Staggs, A. Stark, R. Stompor, K. T. Story, C. Stoughton, A. Suzuki, O. Tajima, G. P. Teply, K. Thompson, P. Timbie, M. Tomasi, J. I. Treu, M. Tristram, G. Tucker, C. Umiltà, A. e. van En-

- gelen, J. D. Vieira, A. G. Vieregg, M. Vogelsberger, G. Wang, S. Watson, M. White, N. Whitehorn, E. J. Wollack, W. L. Kimmy Wu, Z. Xu, S. Yasini, J. Yeck, K. W. Yoon, E. Young, and A. Zonca, CMB-S4 Science Case, Reference Design, and Project Plan, arXiv e-prints , arXiv:1907.04473 (2019), arXiv:1907.04473 [astro-ph.IM].
- [45] M. J. Wilson and M. White, Cosmology with dropout selection: straw-man surveys & CMB lensing, JCAP **2019**, 015 (2019), arXiv:1904.13378 [astro-ph.CO].
- [46] T. Hamana, M. Shirasaki, S. Miyazaki, C. Hikage, M. Oguri, S. More, R. Armstrong, A. Leauthaud, R. Mandelbaum, H. Miyatake, A. J. Nishizawa, M. Simet, M. Takada, H. Aihara, J. Bosch, Y. Komiyama, R. Lupton, H. Murayama, M. A. Strauss, and M. Tanaka, Cosmological constraints from cosmic shear two-point correlation functions with HSC survey first-year data, Publ. Astron. Soc. Japan **72**, 16 (2020), arXiv:1906.06041 [astro-ph.CO].
- [47] S. Joudaki, C. Blake, A. Johnson, A. Amon, M. Asgari, A. Choi, T. Erben, K. Glazebrook, J. Harnois-Déraps, C. Heymans, H. Hildebrandt, H. Hoekstra, D. Klaes, K. Kuijken, C. Lidman, A. Mead, L. Miller, D. Parkinson, G. B. Poole, P. Schneider, M. Viola, and C. Wolf, KiDS-450 + 2dFLenS: Cosmological parameter constraints from weak gravitational lensing tomography and overlapping redshift-space galaxy clustering, Mon. Not. Roy. Astron. Soc. **474**, 4894 (2018), arXiv:1707.06627 [astro-ph.CO].
- [48] F. Beutler, C. Blake, M. Colless, D. H. Jones, L. Staveley-Smith, G. B. Poole, L. Campbell, Q. Parker, W. Saunders, and F. Watson, The 6dF Galaxy Survey:  $z \approx 0$  measurements of the growth rate and  $\sigma_8$ , Mon. Not. Roy. Astron. Soc. **423**, 3430 (2012), arXiv:1204.4725 [astro-ph.CO].
- [49] W. J. Percival, D. Burkey, A. Heavens, A. Taylor, S. Cole, J. A. Peacock, C. M. Baugh, J. Bland-Hawthorn, T. Bridges, R. Cannon, M. Colless, C. Collins, W. Couch, G. Dalton, R. De Propris, S. P. Driver, G. Efstathiou, R. S. Ellis, C. S. Frenk, K. Glazebrook, C. Jackson, O. Lahav, I. Lewis, S. Lumsden, S. Maddox, P. Norberg, B. A. Peterson, W. Sutherland, and K. Taylor, The 2dF Galaxy Redshift Survey: spherical harmonics analysis of fluctuations in the final catalogue, Mon. Not. Roy. Astron. Soc. **353**, 1201 (2004), arXiv:astro-ph/0406513 [astro-ph].
- [50] C. Blake, I. K. Baldry, J. Bland-Hawthorn, L. Christodoulou, M. Colless, C. Conselice, S. P. Driver, A. M. Hopkins, J. Liske, J. Loveday, P. Norberg, J. A. Peacock, G. B. Poole, and A. S. G. Robotham, Galaxy And Mass Assembly (GAMA): improved cosmic growth measurements using multiple tracers of large-scale structure, Mon. Not. Roy. Astron. Soc. **436**, 3089 (2013), arXiv:1309.5556 [astro-ph.CO].
- [51] C. Blake, S. Brough, M. Colless, C. Contreras, W. Couch, S. Croom, T. Davis, M. J. Drinkwater, K. Forster, D. Gilbank, M. Gladders, K. Glazebrook, B. Jelliffe, R. J. Jurek, I. H. Li, B. Madore, D. C. Martin, K. Pimbblet, G. B. Poole, M. Pracy, R. Sharp, E. Wisnioski, D. Woods, T. K. Wyder, and H. K. C. Yee, The WiggleZ Dark Energy Survey: the growth rate of cosmic structure since redshift  $z=0.9$ , Mon. Not. Roy. Astron. Soc. **415**, 2876 (2011), arXiv:1104.2948 [astro-ph.CO].
- [52] S. Alam, M. Ata, S. Bailey, F. Beutler, D. Bizyaev, J. A. Blazek, A. S. Bolton, J. R. Brownstein, A. Burden, C.-H. Chuang, J. Comparat, A. J. Cuesta, K. S. Dawson, D. J. Eisenstein, S. Escoffier, H. Gil-Marín, J. N. Grieb, N. Hand, S. Ho, K. Kinemuchi, D. Kirkby, F. Kitaura, E. Malanushenko, V. Malanushenko, C. Maraston, C. K. McBride, R. C. Nichol, M. D. Olmstead, D. Oravetz, N. Padmanabhan, N. Palanque-Delabrouille, K. Pan, M. Pellejero-Ibanez, W. J. Percival, P. Petitjean, F. Prada, A. M. Price-Whelan, B. A. Reid, S. A. Rodríguez-Torres, N. A. Roe, A. J. Ross, N. P. Ross, G. Rossi, J. A. Rubiño-Martín, S. Saito, S. Salazar-Albornoz, L. Samushia, A. G. Sánchez, S. Satpathy, D. J. Schlegel, D. P. Schneider, C. G. Scóccola, H.-J. Seo, E. S. Sheldon, A. Simmons, A. Slosar, M. A. Strauss, M. E. C. Swanson, D. Thomas, J. L. Tinker, R. Tojeiro, M. V. Magaña, J. A. Vazquez, L. Verde, D. A. Wake, Y. Wang, D. H. Weinberg, M. White, W. M. Wood-Vasey, C. Yèche, I. Zehavi, Z. Zhai, and G.-B. Zhao, The clustering of galaxies in the completed SDSS-III Baryon Oscillation Spectroscopic Survey: cosmological analysis of the DR12 galaxy sample, Mon. Not. Roy. Astron. Soc. **470**, 2617 (2017), arXiv:1607.03155 [astro-ph.CO].
- [53] S. de la Torre, L. Guzzo, J. A. Peacock, E. Branchini, A. Iovino, B. R. Granett, U. Abbas, C. Adami, S. Arnouts, J. Bel, M. Bolzonella, D. Bottini, A. Cappi, J. Coupon, O. Cucciati, I. Davidzon, G. De Lucia, A. Fritz, P. Franzetti, M. Fumana, B. Garilli, O. Ilbert, J. Krywult, V. Le Brun, O. Le Fèvre, D. Maccagni, K. Małek, F. Marulli, H. J. McCracken, L. Moscardini, L. Paioro, W. J. Percival, M. Polletta, A. Pollo, H. Schlagenhafer, M. Scodeggio, L. A. M. Tasca, R. Tojeiro, D. Vergani, A. Zanichelli, A. Burden, C. Di Porto, A. Marchetti, C. Marinoni, Y. Mellier, P. Monaco, R. C. Nichol, S. Phleps, M. Wolk, and G. Zamorani, The VI-MOS Public Extragalactic Redshift Survey (VIPERS) . Galaxy clustering and redshift-space distortions at  $z = 0.8$  in the first data release, Astronomy & Astrophysics **557**, A54 (2013), arXiv:1303.2622 [astro-ph.CO].
- [54] T. Okumura, C. Hikage, T. Totani, M. Tonegawa, H. Okada, K. Glazebrook, C. Blake, P. G. Ferreira, S. More, A. Taruya, S. Tsujikawa, M. Akiyama, G. Dalton, T. Goto, T. Ishikawa, F. Iwamuro, T. Matsubara, T. Nishimichi, K. Ohta, I. Shimizu, R. Takahashi, N. Takato, N. Tamura, K. Yabe, and N. Yoshida, The Subaru FMOS galaxy redshift survey (FastSound). IV. New constraint on gravity theory from redshift space distortions at  $z \sim 1.4$ , Publ. Astron. Soc. Japan **68**, 38 (2016), arXiv:1511.08083 [astro-ph.CO].