

GENERALIZED BRÉZIN–GROSS–WITTEN TAU-FUNCTION AS A HYPERGEOMETRIC SOLUTION OF THE BKP HIERARCHY

ALEXANDER ALEXANDROV

ABSTRACT. In this paper, we prove that the generalized Brézin–Gross–Witten tau-function is a hypergeometric solution of the BKP hierarchy with simple weight generating function. We claim that it describes a spin version of the strictly monotone Hurwitz numbers. A family of the hypergeometric tau-functions of the BKP hierarchy, corresponding to the rational weight generating functions, is investigated. In particular, the cut-and-join operators are constructed, and the explicit description of the BKP Sato Grassmannian points is derived. Representatives of this family can be associated with interesting families of spin Hurwitz numbers including a spin version of the monotone Hurwitz numbers.

MSC 2020 Primary: 37K10; Secondary: 14N10,17B80,81R10.

CONTENTS

1. Introduction	2
1.1. Organization of the paper	3
2. Neutral fermions and boson-fermion correspondence	3
2.1. Neutral fermions	3
2.2. Vertex operators	5
2.3. Boson-fermion correspondence	6
3. $W_{1+\infty}^B$ -algebra	7
3.1. $w_{1+\infty}^B$ -algebra and its central extension	7
3.2. Bases in $w_{1+\infty}^B$ -algebra and its central extension	8
4. Orthogonal Sato Grassmannian	10
4.1. Symmetries of the BKP Sato Grassmannian	11
4.2. KdV vs BKP	13
5. Generalized Brézin–Gross–Witten model	14
5.1. Intersection theory	14
5.2. Cut-and-join operator and BKP tau-function	15
5.3. Schur Q-functions and BKP hierarchy	17
5.4. Schur Q-function expansion of the generalized BGW tau-function	18
6. Higher models	20
6.1. Cut-and-join description	21
6.2. Sato Grassmannian and Kac–Schwarz operators	22
6.3. Examples	24
6.4. Constraints	24
References	25

1. INTRODUCTION

Spin Hurwitz numbers were introduced by Eskin, Okounkov, and Pandharipande [EOP08]. Analogously to the ordinary Hurwitz numbers, they count the ramified coverings, but this time of the surfaces with spin structure. This is an interesting class of enumerative geometry invariants, closely related to the fundamental structures in representation theory and mathematical physics. In particular, the relation between spin Hurwitz numbers and Schur Q -functions was noticed by Gunningham [Gun16]. On the representation theory side the spin Hurwitz numbers are related to the characters of the Sergeev group.

Generating functions of the weighted double spin Hurwitz numbers were identified with the tau-functions of the 2-component BKP integrable hierarchy by Mironov, Morozov and Natanzon [MMN20]. Corresponding class of the BKP tau-functions, so-called *hypergeometric* tau-functions was introduced and investigated by Orlov [Orl03]. In this paper, we introduce and investigate an infinite parametric family of the hypergeometric tau-functions of the 2-component BKP hierarchy. This family is associated with the rational weight generating functions.

The motivation for the investigation of this family is two-fold. On the one hand, the simplest representative of this family is given by the generalized Brézin–Gross–Witten (BGW) tau-function. Relation between the generalized BGW tau-function and the hypergeometric solution of the BKP hierarchy was conjectured in [Ale20a], and its proof is one of the main results of this paper. On the other hand, we claim that this family contains several interesting examples of the generating functions of the weighted spin Hurwitz numbers, including the natural spin analogs of the monotone, strictly monotone, and Bousquet–Mélou–Schaeffer Hurwitz numbers. In particular, the generalized BGW tau-function can be associated with the generating function of the strictly monotone spin Hurwitz numbers. Ordinary strictly monotone Hurwitz numbers describe Grothendieck’s dessins d’enfants and hypermaps, and the spin case should also be related to the interesting invariants of enumerative geometry and combinatorics.

For the ordinary Hurwitz numbers, the rational weight generating functions can be obtained from the combination of the Schur functions with a particular choice of the variables. In the spin case, the correspondence is more complicated and will be considered elsewhere. Therefore, we do not discuss the detailed interpretation of the obtained generating functions in terms of spin Hurwitz numbers and relations between these numbers and intersection theory invariants. Instead in this paper, we focus on the integrable properties of this family. For this purpose we use the standard ingredients of the theory of integrable hierarchies including the free fermions, vertex operators, and the Sato Grassmannian. However, for the BKP and KP hierarchies, corresponding to the spin and ordinary Hurwitz numbers respectively, these ingredients are different, because of the difference between the underlying symmetry algebras, $\mathfrak{go}(\infty)$ and $\mathfrak{gl}(\infty)$.

In particular, we construct the basis for the corresponding points for the BKP Sato Grassmannian and derive associated Kac–Schwarz operators. Moreover, we construct the cut-and-join operators, which, for the polynomial weight generating functions, are the finite degree differential operators. The Kac–Schwarz description allows us to construct the quantum spectral curves and to derive the linear constraints for the tau-functions. We expect that these ingredients will be useful for further investigation of the spin Hurwitz numbers. In particular, they should be useful for the construction of the Chekhov–Eynard–Orantin topological recursion. The description obtained in this paper is the spin analog of the description for the ordinary Hurwitz numbers, considered in [ALS16, AMMN14, KZ15, Zog15].

Notation. In this paper we denote the families of variables or parameters, finite or infinite, by bold symbols. In particular, \mathbf{t} denotes the set of odd variables,

$$(1.1) \quad \mathbf{t} = \{t_1, t_3, t_5, \dots\}.$$

A partition λ is *strict*, if $\lambda_1 > \lambda_2 > \lambda_3 > \dots > \lambda_{\ell(\lambda)} > \lambda_{\ell(\lambda)+1} = 0$, where $\ell(\lambda)$ is the *length* of the partition. We denote the set of strict partitions, including the empty one, by DP.

Acknowledgments. This work was supported by IBS-R003-D1.

1.1. Organization of the paper. In Section 2 we remind the reader the neutral fermion formalism and describe associated vertex operators. Section 3 is devoted to the $w_{1+\infty}^B$ algebra and its central extension, $W_{1+\infty}^B$. In particular, we explicitly describe the convenient bases for these algebras. In Section 4 we outline the Sato Grassmannian description of the BKP hierarchy. In Section 5 we describe the diagonal group element, associated with the generalized BGW tau-function and prove that this tau-function is a hypergeometric solution of the BKP hierarchy. The generalization of this tau-function to the family with rational weight generating functions is investigated in Section 6.

2. NEUTRAL FERMIONS AND BOSON-FERMION CORRESPONDENCE

In this section we remind the reader the neutral fermion formalism and boson-fermion correspondence in the framework of the BKP hierarchy. More details can be found in [DKM81, DJKM82, You89, vdL95, Or103].

2.1. Neutral fermions. Let ϕ_k , $k \in \mathbb{Z}$ be the neutral free fermions satisfying the canonical anticommutation relations

$$(2.1) \quad \{\phi_k, \phi_m\} = (-1)^k \delta_{k+m,0}.$$

Note that $\phi_0^2 = 1/2$. These relations define the Clifford algebra as the associative algebra.

For the vacuum $|0\rangle$ and the co-vacuum $\langle 0|$ vectors, satisfying

$$(2.2) \quad \phi_m |0\rangle = 0, \quad \langle 0| \phi_{-m} = 0, \quad m < 0$$

the elements $\phi_{k_1} \phi_{k_2} \dots \phi_{k_m} |0\rangle$ with $k_1 > k_2 > \dots > k_m \geq 0$ form a basis of a version of infinite wedge space (or neutral fermion Fock space) \mathcal{F}_B ,

$$(2.3) \quad \mathcal{F}_B = \text{span} \{ \phi_{k_1} \phi_{k_2} \dots \phi_{k_m} |0\rangle \mid k_1 > k_2 > \dots > k_m \geq 0 \},$$

and its dual

$$(2.4) \quad \mathcal{F}_B^* = \text{span} \{ \langle 0| \phi_{k_m} \dots \phi_{k_2} \phi_{k_1} \mid k_1 < k_2 < \dots < k_m \leq 0 \}.$$

The space \mathcal{F}_B splits into two subspaces

$$(2.5) \quad \mathcal{F}_B = \mathcal{F}_B^0 \oplus \mathcal{F}_B^1,$$

where \mathcal{F}_B^0 and \mathcal{F}_B^1 denote the subspaces with even and odd numbers of generators respectively. The same decomposition exists for \mathcal{F}_B^* .

There is a nondegenerate bilinear pairing $\mathcal{F}_B \times \mathcal{F}_B^* \rightarrow \mathbb{C}$, and the pairing of $\langle U| \in \mathcal{F}_B^*$ and $|V\rangle \in \mathcal{F}_B$ is denoted by $\langle U|V\rangle$ with

$$(2.6) \quad \langle 0|0\rangle = 1.$$

The *vacuum expectation value* of the Clifford algebra element a is a pairing of $\langle 0|$ and $a|0\rangle$ which is denoted by $\langle 0|a|0\rangle$. It is uniquely defined by the anticommutation relations (2.1), property (2.2), and a relation

$$(2.7) \quad \langle 0| \phi_0 |0\rangle = 0.$$

In particular, if a is an odd element of the Clifford algebra, then $\langle 0|a|0\rangle = 0$. It is easy to see that the bases in (2.3) and (2.4) are orthogonal. Let us focus on the space \mathcal{F}_B^0 . The basis can be labelled by strict partitions $\lambda \in \text{DP}$ in the following way:

$$(2.8) \quad |\lambda\rangle = \begin{cases} \phi_{\lambda_1} \phi_{\lambda_2} \cdots \phi_{\lambda_{\ell(\lambda)}} |0\rangle & \text{for } \ell(\lambda) = 0 \pmod{2}, \\ \sqrt{2} \phi_{\lambda_1} \phi_{\lambda_2} \cdots \phi_{\lambda_{\ell(\lambda)}} \phi_0 |0\rangle & \text{for } \ell(\lambda) = 1 \pmod{2}. \end{cases}$$

For the dual space \mathcal{F}_B^{0*} we have a similar basis. From the anticommutation relations we have:

Proposition 2.1.

$$(2.9) \quad \langle \lambda | \mu \rangle = (-1)^{|\lambda|} \delta_{\lambda, \mu}.$$

It is easy to see that

$$(2.10) \quad \langle 0 | \phi_k \phi_m | 0 \rangle = \delta_{k+m, 0} H[m],$$

where

$$(2.11) \quad H[m] = \begin{cases} 0 & \text{for } m < 0, \\ \frac{1}{2} & \text{for } m = 0, \\ (-1)^m & \text{for } m > 0. \end{cases}$$

Bilinear combinations of neutral fermions $\phi_k \phi_m$ satisfy the commutation relations of the Lie algebra $\mathfrak{go}(\infty)$. Let $(E_{i,j})_{k,l} = \delta_{i,k} \delta_{j,l}$ be the standard basis of the matrix units $\{E_{i,j} | i, j \in \mathbb{Z}\}$. Then $\phi_k \phi_{-m}$ corresponds [DJKM82] to

$$(2.12) \quad F_{k,m} = (-1)^m E_{k,m} - (-1)^k E_{-m,-k}$$

with the commutation relations

$$(2.13) \quad [F_{a,b}, F_{c,d}] = (-1)^b \delta_{b,c} F_{a,d} - (-1)^a \delta_{a+c,0} F_{-b,d} + (-1)^b \delta_{b+d,0} F_{c,-a} - (-1)^a \delta_{a,d} F_{c,b}.$$

For the bilinear combinations of neutral fermions we introduce the *normal ordering* by $:\phi_k \phi_m := \phi_k \phi_m - \langle 0 | \phi_k \phi_m | 0 \rangle$. It is skew-symmetric

$$(2.14) \quad :\phi_k \phi_m := - : \phi_m \phi_k :,$$

in particular, $:\phi_k \phi_k := 0$. The normal ordered quadratic combinations of neutral fermions satisfy the commutation relations of a central extension of the algebra $\mathfrak{go}(\infty)$

$$(2.15) \quad [:\phi_a \phi_b :, : \phi_c \phi_d :] = (-1)^b \delta_{b+c,0} : \phi_a \phi_d : - (-1)^a \delta_{a+c,0} : \phi_b \phi_d : \\ + (-1)^b \delta_{b+d,0} : \phi_c \phi_a : - (-1)^a \delta_{a+d,0} : \phi_c \phi_b : + (\delta_{c,b} \delta_{a,d} - \delta_{a-c,0} \delta_{b-d,0}) ((-1)^a H[b] - (-1)^b H[a]),$$

where $H[a]$ is given by (2.11).

Let us consider the generating function

$$(2.16) \quad \phi(z) = \sum_{k \in \mathbb{Z}} \phi_k z^k.$$

It satisfies the anticommutation relation

$$(2.17) \quad \{\phi(z), \phi(w)\} = \delta(z+w).$$

Here we introduce the delta-function

$$(2.18) \quad \delta(z-w) = \sum_{k \in \mathbb{Z}} \left(\frac{z}{w}\right)^k.$$

It satisfies

$$(2.19) \quad \delta(z-w) f(z) = \delta(z-w) f(w)$$

for any formal series $f(z) \in \mathbb{C}[[z, z^{-1}]]$, and can be represented as

$$(2.20) \quad 2\delta(z+w) = \frac{z-w}{\underline{z}+w} - \frac{z-w}{\underline{w}+z}.$$

Here $\frac{1}{\underline{z}-w}$ denotes the Laurent series expansion in the region $|z| > |w|$,

$$(2.21) \quad \frac{1}{\underline{z}-w} := \frac{1}{z} \sum_{k=0}^{\infty} \left(\frac{w}{z}\right)^k.$$

Quadratic combinations of these generating functions generate a Lie algebra with the following commutation relations

$$(2.22) \quad [\phi_1(z_1)\phi(w_1), \phi(z_2)\phi(w_2)] = \delta(w_1+z_2)\phi(z_1)\phi(w_2) - \delta(z_1+z_2)\phi(w_1)\phi(w_2) \\ + \delta(w_1+w_2)\phi(z_2)\phi(z_1) - \delta(z_1+w_2)\phi(z_2)\phi(w_1).$$

For the normal ordered operator we have

$$(2.23) \quad \phi(z)\phi(w) = : \phi(z)\phi(w) : + \frac{1}{2} \frac{z-w}{\underline{z}+w}.$$

2.2. Vertex operators. For $k \in \mathbb{Z}_{\text{odd}}$ we introduce the *bosonic* operators

$$(2.24) \quad J_k = \frac{1}{2} \sum_{m \in \mathbb{Z}} (-1)^{m+1} : \phi_m \phi_{-m-k} :$$

satisfying a commutation relation of the Heisenberg algebra

$$(2.25) \quad [J_k, J_m] = \frac{k}{2} \delta_{k+m,0}.$$

From (2.2) we have

$$(2.26) \quad J_m |0\rangle = 0, \quad \langle 0| J_{-m} = 0, \quad m > 0.$$

Let us consider the *vertex operator* for the BKP hierarchy introduced in [DJKM82],

$$(2.27) \quad \widehat{V}_B(z) = \exp \left(\sum_{k \in \mathbb{Z}_{\text{odd}}^+} z^k t_k \right) \exp \left(-2 \sum_{k \in \mathbb{Z}_{\text{odd}}^+} \frac{1}{kz^k} \frac{\partial}{\partial t_k} \right).$$

These operators satisfy the anticommutation relations

$$(2.28) \quad \{ \widehat{V}_B(z), \widehat{V}_B(w) \} = 2\delta(z+w)$$

similar to (2.17).

It is convenient to introduce the generating functions of the bosonic operators

$$(2.29) \quad J_+(\mathbf{t}) = \sum_{k \in \mathbb{Z}_{\text{odd}}^+} t_k J_k, \quad J_-(\mathbf{s}) = \sum_{k \in \mathbb{Z}_{\text{odd}}^+} s_k J_{-k}.$$

Then one has

$$(2.30) \quad \widehat{V}_B(z) \langle 0| e^{J_+(\mathbf{t})} = 2 \langle 0| \phi_0 e^{J_+(\mathbf{t})} \phi(z), \\ \widehat{V}_B(z) \langle 0| \phi_0 e^{J_+(\mathbf{t})} = \langle 0| e^{J_+(\mathbf{t})} \phi(z).$$

Let us consider a bilinear combination of the vertex operators \widehat{V}_B ,

$$(2.31) \quad \widehat{V}_B(z, w) = \frac{1}{2} \widehat{V}_B(z) \widehat{V}_B(w).$$

Using anticommutation relation (2.28) it easy to show that the vertex operators $\widehat{V}_B(z, w)$ satisfy the commutation relation, equivalent to one described by (2.22) for the bilinear combinations $\phi(z)\phi(w)$,

$$(2.32) \quad \left[\widehat{V}_B(z_1, w_1), \widehat{V}_B(z_2, w_2) \right] = \delta(w_1 + z_2) \widehat{V}_B(z_1, w_2) \\ - \delta(z_1 + z_2) \widehat{V}_B(w_1, w_2) + \delta(w_1 + w_2) \widehat{V}_B(z_2, z_1) - \delta(z_1 + w_2) \widehat{V}_B(z_2, w_1).$$

These vertex operators, dependent on two parameters, can be represented as

$$(2.33) \quad \widehat{V}_B(z, w) = \frac{1}{2} \frac{z - w}{z + w} \exp \left(\sum_{k \in \mathbb{Z}_{\text{odd}}^+} (z^k + w^k) t_k \right) \exp \left(-2 \sum_{k \in \mathbb{Z}_{\text{odd}}^+} \left(\frac{1}{kz^k} + \frac{1}{kw^k} \right) \frac{\partial}{\partial t_k} \right).$$

It is also convenient to consider a regularized version of the vertex operator, corresponding to $:\phi(z)\phi(-w):$

$$(2.34) \quad \widehat{Y}_B(z, w) = \widehat{V}_B(z, w) - \frac{1}{2} \frac{z - w}{z + w},$$

or, equivalently

$$(2.35) \quad \widehat{Y}_B(z, w) = \frac{1}{2} \left(\widehat{V}_B(z) \widehat{V}_B(w) - \frac{z - w}{z + w} \right).$$

This expression is regular at $z = -w$, moreover, it is antisymmetric with respect to the permutation of z and w ,

$$(2.36) \quad \widehat{Y}_B(z, w) = -\widehat{Y}_B(w, z).$$

From (2.30) it follows that

$$(2.37) \quad \widehat{Y}_B(z, w) \langle 0 | e^{J_+(t)} = \langle 0 | e^{J_+(t)} : \phi(z)\phi(w) : .$$

2.3. Boson-fermion correspondence. For the neutral fermions the boson-fermion correspondence [You89] describes an isomorphism

$$(2.38) \quad \sigma_B^i : \mathcal{F}_B^i \simeq B^{(i)} = \mathbb{C}[[t_1, t_3, t_5, \dots]]$$

for $i = 0, 1$. Here

$$(2.39) \quad \sigma_B^i(|i\rangle) = 1,$$

where we introduce $|1\rangle = \sqrt{2}\phi_0|0\rangle$, and for both $i = 0, 1$ we have

$$(2.40) \quad \sigma_B^i J_{-k} (\sigma_B^i)^{-1} = \frac{k}{2} t_k, \quad \sigma_B^i J_k (\sigma_B^i)^{-1} = \frac{\partial}{\partial t_k}$$

for $k \in \mathbb{Z}_{\text{odd}}^+$. The boson-fermion correspondence is given by

$$(2.41) \quad \sigma_B^i(|a\rangle) = \begin{cases} \langle 1 | e^{J_+(t)} | a \rangle & \text{for } |a\rangle \in \mathcal{F}_B^1, \\ \langle 0 | e^{J_+(t)} | a \rangle & \text{for } |a\rangle \in \mathcal{F}_B^0, \end{cases}$$

where $\langle 1 | = \sqrt{2} \langle 0 | \phi_0$. The boson-fermion correspondence between two different representations of the central extension of the $\mathfrak{go}(\infty)$ algebra is given by

$$(2.42) \quad \sigma_B^i : \phi(z)\phi(w) : (\sigma_B^i)^{-1} = \widehat{Y}_B(z, w).$$

Below we will work only with \mathcal{F}_B^0 component of the fermionic Fock space and its bosonic counterpart. For them we denote the boson-fermion correspondence by σ .

3. $W_{1+\infty}^B$ -ALGEBRA

Content of this section is closely related to the results of van de Leur [vdL95].

3.1. $w_{1+\infty}^B$ -algebra and its central extension. Let us introduce the algebra $W_{1+\infty}^B$. Consider $w_{1+\infty}$, the algebra of diffeomorphisms on the circle,

$$(3.1) \quad w_{1+\infty} = \text{span}\{z^k \partial_z^m \mid k \in \mathbb{Z}, m \in \mathbb{Z}_{\geq 0}\}.$$

Let ι be the anti-involution of $w_{1+\infty}$

$$(3.2) \quad \iota(z) = -z, \quad \iota(z \partial_z) = -z \partial_z.$$

We define

$$(3.3) \quad w_{1+\infty}^B = \{\mathbf{a} \in w_{1+\infty} \mid \iota(\mathbf{a}) = -\mathbf{a}\}.$$

For any $\mathbf{a} \in w_{1+\infty}$ we denote by $\bar{\mathbf{a}}$ the operator, obtained by the sign inversion of z ,

$$(3.4) \quad \bar{\mathbf{a}} = \mathbf{a} \Big|_{z \mapsto -z, \partial_z \mapsto -\partial_z}.$$

Then it is easy to see that

$$(3.5) \quad \iota(\mathbf{a}) = \bar{\mathbf{a}}^*,$$

where $\mathbf{a}^* \in w_{1+\infty}$ is the adjoint operator for which an identity

$$\text{Res}_z (z^{-1} f(z) \mathbf{a} g(z)) = \text{Res}_z (z^{-1} g(z) \mathbf{a}^* f(z))$$

holds for any commuting $f(z)$ and $g(z)$. Here

$$(3.6) \quad \text{Res}_z \sum_{k \in \mathbb{Z}} a_k z^k := a_{-1}.$$

In particular,

$$(3.7) \quad (z^k \partial_z^m)^* = z(-\partial_z)^m z^{k-1}.$$

Definition 1. For any operator $\mathbf{a} \in w_{1+\infty}$ we introduce a differential operator, acting in the bosonic Fock space $\mathbb{C}[[t_1, t_3, t_5 \dots]]$:

$$(3.8) \quad \widehat{W}_{\mathbf{a}}^B = \frac{1}{2} \text{Res}_w w^{-1} \mathbf{a}_w \cdot \widehat{Y}_B(z, w) \Big|_{z=-w}.$$

Here by \mathbf{a}_w we denote the operator \mathbf{a} acting on the space of functions of the variable w .

Lemma 3.1. For $\mathbf{a} \in w_{1+\infty}$

$$(3.9) \quad [\widehat{V}_B(z), \widehat{W}_{\mathbf{a}}^B] = \frac{1}{2} (\mathbf{a}_z - \iota(\mathbf{a}_z)) \cdot \widehat{V}_B(z).$$

Proof. From the anticommutation relation (2.28) we have

$$(3.10) \quad [\widehat{V}_B(z), \widehat{Y}_B(v, w)] = \delta(z+v) \widehat{V}_B(w) - \delta(z+w) \widehat{V}_B(v).$$

Hence,

$$(3.11) \quad \begin{aligned} [\widehat{V}_B(z), \widehat{W}_{\mathbf{a}}^B] &= \frac{1}{2} \text{Res}_w w^{-1} \mathbf{a}_w (\delta(z+v) \widehat{V}_B(w) - \delta(z+w) \widehat{V}_B(v)) \Big|_{v=-w} \\ &= \frac{1}{2} \text{Res}_w w^{-1} (\delta(z-w) \mathbf{a}_w \widehat{V}_B(w) - \delta(z+w) \mathbf{a}_w^* \widehat{V}_B(-w)), \end{aligned}$$

and the statement of the lemma follows from (3.5). \square

Corollary 3.2. For $\mathbf{a} \in w_{1+\infty}^B$

$$(3.12) \quad [\widehat{V}_B(z), \widehat{W}_{\mathbf{a}}^B] = \mathbf{a}_z \cdot \widehat{V}_B(z).$$

From the commutation relation (2.32) and definition of the regularized vertex operator (2.34) we have

$$(3.13) \quad \begin{aligned} & \left[\widehat{Y}_B(z_1, w_1), \widehat{Y}_B(z_2, w_2) \right] = \delta(w_1 + z_2) \widehat{Y}_B(z_1, w_2) \\ & \quad - \delta(z_1 + z_2) \widehat{Y}_B(w_1, w_2) + \delta(w_1 + w_2) \widehat{Y}_B(z_2, z_1) - \delta(z_1 + w_2) \widehat{Y}_B(z_2, w_1) \\ & \quad + \frac{1}{4} \left(\frac{w_1 - z_2}{\underline{w}_1 + z_2} \frac{z_1 - w_2}{\underline{z}_1 + w_2} - \frac{z_1 - z_2}{\underline{z}_1 + z_2} \frac{w_1 - w_2}{\underline{w}_1 + w_2} + \frac{w_2 - w_1}{\underline{w}_2 + w_1} \frac{z_2 - z_1}{\underline{z}_2 + z_1} - \frac{w_2 - z_1}{\underline{w}_2 + z_1} \frac{z_2 - w_1}{\underline{z}_2 + w_1} \right). \end{aligned}$$

Here we use (2.20).

Let

$$(3.14) \quad \mu(\mathbf{a}, \mathbf{b}) = \frac{1}{8} \operatorname{Res}_{w_1} \operatorname{Res}_{w_2} w_1^{-1} w_2^{-1} \frac{w_1 - w_2}{\underline{w}_1 + w_2} (\mathbf{a}_{w_2} \mathbf{b}_{w_1} - \mathbf{a}_{w_1} \mathbf{b}_{w_2}) \frac{w_1 - w_2}{\underline{w}_1 + w_2}.$$

Then using (3.13), from a direct computation we see that operators $\widehat{W}_\mathbf{a}^B$ describe algebra $W_{1+\infty}^B$, a central extension of the algebra $w_{1+\infty}^B$:

Lemma 3.3. For $\mathbf{a}, \mathbf{b} \in w_{1+\infty}^B$

$$(3.15) \quad \left[\widehat{W}_\mathbf{a}^B, \widehat{W}_\mathbf{b}^B \right] = \widehat{W}_{[\mathbf{a}, \mathbf{b}]}^B + \mu(\mathbf{a}, \mathbf{b}).$$

From (2.37) we have

$$(3.16) \quad \widehat{W}_\mathbf{a}^B \cdot \langle 0 | e^{J_+(\mathbf{t})} = \langle 0 | e^{J_+(\mathbf{t})} W_\mathbf{a}^B,$$

where

$$(3.17) \quad W_\mathbf{a}^B = \frac{1}{2} \operatorname{Res}_w w^{-1} : \phi(-w) \mathbf{a}_w \phi(w) :$$

is the bilinear fermionic operator acting on the fermionic Fock space.

3.2. Bases in $w_{1+\infty}^B$ -algebra and its central extension. Let us consider the operators

$$(3.18) \quad \mathbf{w}_{k,m} := -z^{k+m} (\partial_z)^m + (-1)^{k+m} z (\partial_z)^m z^{k+m-1}$$

for $k \in \mathbb{Z}$, $m \in \mathbb{Z}_{\geq 0}$. It is easy to see that $\mathbf{w}_{k,m} \in w_{1+\infty}^B$. Operators with $k+m \in \mathbb{Z}_{\text{odd}}$ constitute a basis in $w_{1+\infty}^B$, for example

$$(3.19) \quad \begin{aligned} \mathbf{w}_{k,0} &= -2z^k, & k \in \mathbb{Z}_{\text{odd}}, \\ \mathbf{w}_{k,1} &= -2z^{k+1} \partial_z - kz^k, & k \in \mathbb{Z}_{\text{even}}, \\ \mathbf{w}_{k,2} &= -2z^{k+2} \partial_z^2 - 2(k+1)z^{k+1} \partial_z - (k+1)kz^k, & k \in \mathbb{Z}_{\text{even}}. \end{aligned}$$

Associated basis of $W_{1+\infty}^B$ consists of operators

$$(3.20) \quad \widehat{W}_{k,m}^B := \widehat{W}_{\mathbf{w}_{k,m}}^B.$$

for $k+m \in \mathbb{Z}_{\text{odd}}$, and a central element.

Let

$$(3.21) \quad \widehat{J}_B(z) = \sum_{k \in \mathbb{Z}_{\text{odd}}^+} \left(kt_k z^{k-1} + 2z^{-k-1} \frac{\partial}{\partial t_k} \right).$$

We introduce the *bosonic normal ordering* $\bullet \dots \bullet$ which puts all t_k to the left of all $\frac{\partial}{\partial t_k}$. Consider the operators

$$(3.22) \quad U_k(z) = \bullet P_k(\widehat{J}_B(z)) \bullet.$$

Here P_m are the Faà di Bruno differential polynomials:

$$(3.23) \quad P_k(\varphi') = e^{-\varphi(z)} \partial_z^k e^{\varphi(z)} = (\partial_z + \varphi')^k,$$

where $\varphi' = \partial_z \varphi(z)$. They have a simple expression in terms of elementary Schur functions p_k ,

$$(3.24) \quad P_k(\varphi') = k! p_k(\varphi^{(j)}(z)/j!),$$

because

$$(3.25) \quad \sum_{k=0}^{\infty} \frac{s^k}{k!} P_k(\partial_z \varphi) = e^{-\varphi(z)} e^{\varphi(z+s)} = e^{\sum_{j=1}^{\infty} \frac{s^j}{j!} \varphi^{(j)}(z)} = \sum_{k=0}^{\infty} s^k p_k(\varphi^{(j)}(z)/j!).$$

Let

$$(3.26) \quad \widehat{\varphi}_B(z) = \sum_{k \in \mathbb{Z}_{\text{odd}}^+} \left(t_k z^k - 2 \frac{1}{k z^k} \frac{\partial}{\partial t_k} \right),$$

then $\widehat{J}_B(z) = \partial_z \widehat{\varphi}_B(z)$ and the vertex operator (2.35) can be expressed as

$$(3.27) \quad \widehat{Y}_B(z, w) = \frac{1}{2} \frac{z-w}{z+w} \bullet e^{\widehat{\varphi}_B(z) + \widehat{\varphi}_B(w)} - 1 \bullet.$$

The following lemma describes operators $\widehat{W}_{k,m}^B$ in terms of the Faà di Bruno polynomials

Lemma 3.4. For $k \in \mathbb{Z}$, $m \in \mathbb{Z}_{\geq 0}$

$$(3.28) \quad \widehat{W}_{k,m}^B = \frac{1}{2} \text{Res}_w w^{k+m-1} \left(\frac{2w}{m+1} U_{m+1}(w) + U_m(w) \right),$$

where we put $U_0(w) = 0$.

Proof. By definition

$$(3.29) \quad \begin{aligned} \widehat{W}_{k,m}^B &= \frac{1}{2} \text{Res}_w w^{-1} \left((-1)^{k+m} w \partial_w^m w^{k+m-1} - w^{k+m} \partial_w^m \right) \widehat{Y}_B(z, w) \Big|_{z=-w} \\ &= \frac{1}{2} \text{Res}_w w^{-1} \left(z^{k+m} \partial_z^m - w^{k+m} \partial_w^m \right) \widehat{Y}_B(z, w) \Big|_{z=-w}. \end{aligned}$$

It is easy to see that for any series $f(z, w) \in \mathbb{C}[[z, w]]$ such that f is antisymmetric, $f(z, w) = -f(w, z)$, one has

$$(3.30) \quad \text{Res}_w w^{-1} z^k \partial_z^m f(z, w) \Big|_{z=-w} = - \text{Res}_w w^{-1} w^k \partial_w^m f(z, w) \Big|_{z=-w}.$$

Since $\widehat{Y}_B(z, w)$ is antisymmetric, (2.36), we conclude that

$$(3.31) \quad \widehat{W}_{k,m}^B = - \text{Res}_w w^{k+m-1} (\partial_w)^m \widehat{Y}_B(z, w) \Big|_{z=-w}.$$

Then

$$(3.32) \quad \begin{aligned} w^k \partial_w^m \widehat{Y}_B(z, w) \Big|_{z=-w} &= w^k \partial_{\Delta}^m \widehat{Y}_B(-w, w + \Delta) \Big|_{\Delta=0} \\ &= -\frac{1}{2} w^k \partial_{\Delta}^m (2w + \Delta) \sum_{j=1}^{\infty} \frac{\Delta^{j-1}}{j!} U_j(w) \Big|_{\Delta=0} \\ &= -\frac{1}{2} w^k \left(\frac{2w}{m+1} U_{m+1}(w) + U_m(w) \right), \end{aligned}$$

where for $m = 0$ we assume $U_0(w) = 0$. This concludes the proof. \square

The *Heisenberg-Virasoro subalgebra* of the algebra $W_{1+\infty}^B$ is generated by the operators

$$(3.33) \quad \widehat{J}_k^B = \begin{cases} 2\frac{\partial}{\partial t_k} & \text{for } k \in \mathbb{Z}_{\text{odd}}^+, \\ -kt_{-k} & \text{for } -k \in \mathbb{Z}_{\text{odd}}^+, \\ 0 & \text{otherwise} \end{cases}$$

and the Virasoro operators

$$(3.34) \quad \widehat{L}_k^B = \begin{cases} \frac{1}{2} \sum_{i+j=k} :\widehat{J}_i^B \widehat{J}_j^B:, & \text{for } k \in \mathbb{Z}_{\text{even}}, \\ 0, & \text{for } k \in \mathbb{Z}_{\text{odd}}. \end{cases}$$

Here the bosonic normal ordering puts all \widehat{J}_m^B with positive m to the right of all \widehat{J}_m^B with negative m . Let us also consider the $W^{(3)}$ -algebra, which includes the generators

$$(3.35) \quad \widehat{M}_k^B = \begin{cases} \frac{1}{3} \sum_{i+j+l=k} :\widehat{J}_i^B \widehat{J}_j^B \widehat{J}_l^B:, & \text{for } k \in \mathbb{Z}_{\text{odd}}, \\ 0, & \text{for } k \in \mathbb{Z}_{\text{even}}. \end{cases}$$

These operators satisfy the following commutation relations

$$(3.36) \quad \begin{aligned} [\widehat{J}_k^B, \widehat{J}_m^B] &= 2\delta_{k+m,0}k, & k, m \in \mathbb{Z}_{\text{odd}}, \\ [\widehat{J}_k^B, \widehat{L}_m^B] &= 2k\widehat{J}_{k+m}^B, & k \in \mathbb{Z}_{\text{odd}}, m \in \mathbb{Z}_{\text{even}}, \\ [\widehat{L}_k^B, \widehat{L}_m^B] &= 2(k-m)\widehat{L}_{k+m}^B + \frac{1}{3}k(k^2-1)\delta_{k,-m}, & k, m \in \mathbb{Z}_{\text{even}}, \\ [\widehat{J}_k^B, \widehat{M}_m^B] &= 4k\widehat{L}_{k+m}^B, & k, m \in \mathbb{Z}_{\text{odd}}, \\ [\widehat{L}_k^B, \widehat{M}_m^B] &= 2(2k-m)\widehat{M}_{k+m}^B + \frac{2}{3}k(k^2-1)\widehat{J}_{k+m}^B, & k \in \mathbb{Z}_{\text{even}}, m \in \mathbb{Z}_{\text{odd}}, \end{aligned}$$

and, except for the case of two operators \widehat{M}_m^B , these operators commute otherwise. A commutator of two \widehat{M}_k^B 's contains the terms of fourth power of the bosonic operators \widehat{J}_m^B , so it can not be represented as a linear combination of \widehat{J}_k^B , \widehat{L}_k^B , and \widehat{M}_k^B .

For operators $\widehat{W}_{k,m}^B$ with $m \leq 2$ we have

$$(3.37) \quad \begin{aligned} \widehat{W}_{k,0}^B &= \widehat{J}_k^B, \\ \widehat{W}_{k,1}^B &= \widehat{L}_k^B - \frac{k}{2}\widehat{J}_k^B, \\ \widehat{W}_{k,2}^B &= \widehat{M}_k^B - (k+1)\widehat{L}_k^B + \frac{(k+1)(2k+1)}{6}\widehat{J}_k^B. \end{aligned}$$

4. ORTHOGONAL SATO GRASSMANNIAN

In this section we briefly summarize some properties of the Sato Grassmannian. It describes the space of solutions of the KP hierarchy [SS83] and its BKP version. We describe the action of the algebra $w_{1+\infty}^B$ on the BKP Sato Grassmannian and clarify the difference between the points of the KP and BKP Sato Grassmannians for the solutions of the KdV hierarchy.

4.1. Symmetries of the BKP Sato Grassmannian. Let us consider the space $H = H_+ \oplus H_-$, where the subspaces

$$(4.1) \quad H_- = z^{-1}\mathbb{C}[[z^{-1}]]$$

and

$$(4.2) \quad H_+ = \mathbb{C}[z]$$

are generated by negative and nonnegative powers of z respectively. Then the Sato Grassmannian Gr consists of all closed linear spaces $\mathcal{W} \in H$, which are compatible with H_+ . Namely, an orthogonal projection $\pi_+ : \mathcal{W} \rightarrow H_+$ should be a Fredholm operator, i.e. both the kernel $\ker \pi_+ \in \mathcal{W}$ and the cokernel $\text{coker} \pi_+ \in H_+$ should be finite-dimensional vector spaces. The Grassmannian Gr consists of components $\text{Gr}^{(k)}$, parametrized by an index of the operator π_+ . Below we use only the component $\text{Gr}^{(0)}$; other components have an equivalent description. The big cell $\text{Gr}_+^{(0)}$ of $\text{Gr}^{(0)}$ is defined by the constraint $\ker \pi_+ = \text{coker} \pi_+ = 0$. We call $\text{Gr}_+^{(0)}$ the *Sato Grassmannian* for simplicity.

A point of the Sato Grassmannian $\mathcal{W} \in \text{Gr}_+^{(0)}$ can be described by an *admissible basis*

$$(4.3) \quad \mathcal{W} = \text{span}_{\mathbb{C}}\{\Phi_1^{\mathcal{W}}, \Phi_2^{\mathcal{W}}, \Phi_3^{\mathcal{W}}, \dots\}.$$

Let

$$(4.4) \quad \Psi = \mathcal{W} \cap 1 + z^{-1}\mathbb{C}[[z^{-1}]].$$

be the *wave function*.

The BKP hierarchy can be represented in terms of tau-function $\tau(\mathbf{t})$ by the Hirota bilinear identity

$$(4.5) \quad \text{Res}_z z^{-1} e^{\sum_{k \in \mathbb{Z}^+_{\text{odd}}} z^k (t_k - t'_k)} \tau(\mathbf{t} - 2[z^{-1}]) \tau(\mathbf{t}' + 2[z^{-1}]) = \tau(\mathbf{t}) \tau(\mathbf{t}').$$

Here \mathbf{t} and \mathbf{t}' are two independent sets of variables. For $f(z) \in H$ and $g(z) \in H$ put

$$(4.6) \quad (f(z), g(z))_B := \text{Res}_z z^{-1} f(z) g(-z).$$

Note that $(\ , \)_B$ is a non-degenerate symmetric bilinear pairing on H . For $\mathcal{W} \in \text{Gr}^{(0)}$ we denote by $\mathcal{W}^{\perp B}$ the orthogonal complement of \mathcal{W} with respect to $(\ , \)_B$. The BKP (or orthogonal) Sato Grassmannian $\text{Gr}_B^{(0)}$ is the subspace of $\text{Gr}^{(0)}$ such that $\mathcal{W}^{\perp B} \subset \mathcal{W}$ and the quotient $\mathcal{W}/\mathcal{W}^{\perp B}$ is one dimensional and is generated by Ψ . Similarly to the KP case, one can consider the big cell of the BKP Sato Grassmannian, $\text{Gr}_{B+}^{(0)} = \text{Gr}_+^{(0)} \cap \text{Gr}_B^{(0)}$. There exists a bijection between the points of the BKP Sato Grassmannian $\mathcal{W} \in \text{Gr}_{B+}^{(0)}$ and the BKP tau-functions with $\tau(\mathbf{0}) = 1$. Below we focus on $\text{Gr}_{B+}^{(0)}$, and put $\tau(\mathbf{0}) = 1$ for simplicity.

Tau-functions of the BKP hierarchy are given by the vacuum expectation values of neutral fermions

$$(4.7) \quad \tau_G(\mathbf{t}) = \langle 0 | e^{J+(\mathbf{t})} G | 0 \rangle.$$

They are labeled by the group element of the centrally extended $\mathfrak{go}(\infty)$ algebra,

$$(4.8) \quad G = \exp \left(\sum_{k,m \in \mathbb{Z}} a_{km} : \phi_k \phi_m : \right).$$

The wave function is equal to the principal specialization of the BKP tau-function

$$(4.9) \quad \Psi(z) = \tau(-2[z^{-1}]) = \tau(2[-z^{-1}]),$$

because the BKP tau-functions depend only on odd times. The *Baker–Akhiezer function*

$$(4.10) \quad \Psi^B(z, \mathbf{t}) = e^{\sum_{k \in \mathbb{Z}^+_{\text{odd}}} t_k z^k} \frac{\tau(\mathbf{t} - 2[z^{-1}])}{\tau(\mathbf{t})}$$

generates the corresponding point of the BKP Sato Grassmannian through the \mathbf{t} series expansion

$$(4.11) \quad \Psi(z, \mathbf{t}) \in \mathcal{W}.$$

Let us label the Baker–Akhiezer function, associated with the tau-function (4.7), by G . Then, from (2.30) we have

$$(4.12) \quad \Psi_G^B(z, \mathbf{t}) = \frac{2 \langle 0 | \phi_0 e^{J_+(\mathbf{t})} \phi(z) G | 0 \rangle}{\tau_G(\mathbf{t})}.$$

For some $\mathbf{a} \in w_{1+\infty}^B$ let us consider

$$(4.13) \quad \tilde{G} = e^{W_{\mathbf{a}}^B} G,$$

where $W_{\mathbf{a}}^B$ is given by (3.17). We assume that \mathbf{a} and G are such that both τ_G and $\tau_{\tilde{G}}$ are well defined. Then from (2.30) for the associated wave function we have

$$(4.14) \quad \begin{aligned} \Psi_{\tilde{G}}^B(z, \mathbf{t}) &= \frac{\widehat{V}_B(z) \langle 0 | e^{J_+(\mathbf{t})} e^{W_{\mathbf{a}}^B} G | 0 \rangle}{\tau_{\tilde{G}}(\mathbf{t})} \\ &= \frac{\widehat{V}_B(z) e^{\widehat{W}_{\mathbf{a}}^B} \langle 0 | e^{J_+(\mathbf{t})} G | 0 \rangle}{\tau_{\tilde{G}}(\mathbf{t})}. \end{aligned}$$

From Corollary 3.2 we have the operator identity

$$(4.15) \quad \widehat{V}_B(z) e^{\widehat{W}_{\mathbf{a}}^B} = e^{\widehat{W}_{\mathbf{a}}^B} \left(e^{-\widehat{W}_{\mathbf{a}}^B} \widehat{V}_B(z) e^{\widehat{W}_{\mathbf{a}}^B} \right) = e^{\widehat{W}_{\mathbf{a}}^B} \left(e^{a z} \widehat{V}_B(z) \right),$$

hence

$$(4.16) \quad \begin{aligned} \Psi_{\tilde{G}}^B(z, \mathbf{t}) &= \frac{e^{\widehat{W}_{\mathbf{a}}^B} e^{a z} \widehat{V}_B(z) \langle 0 | e^{J_+(\mathbf{t})} G | 0 \rangle}{\tau_{\tilde{G}}(\mathbf{t})} \\ &= \frac{e^{\widehat{W}_{\mathbf{a}}^B} \tau_G(\mathbf{t}) e^{a z} \Psi_G^B(z, \mathbf{t})}{\tau_{\tilde{G}}(\mathbf{t})}. \end{aligned}$$

Inverting the operator $e^{\widehat{W}_{\mathbf{a}}^B}$ we have

$$(4.17) \quad e^{a z} \Psi_G^B(z, \mathbf{t}) = \frac{e^{-\widehat{W}_{\mathbf{a}}^B} \tau_{\tilde{G}}(\mathbf{t}) \Psi_{\tilde{G}}^B(z, \mathbf{t})}{\tau_G(\mathbf{t})}.$$

Since $\Psi_{\tilde{G}}^B(z, \mathbf{t}) \in \mathcal{W}_{\tilde{G}}$, we also have

$$(4.18) \quad \frac{e^{-\widehat{W}_{\mathbf{a}}^B} \tau_{\tilde{G}}(\mathbf{t}) \Psi_{\tilde{G}}^B(z, \mathbf{t})}{\tau_G(\mathbf{t})} \in \mathcal{W}_{\tilde{G}}.$$

Therefore,

$$(4.19) \quad e^{\mathbf{a}} \cdot \mathcal{W}_G \subset \mathcal{W}_{\tilde{G}},$$

and inverse is also true

$$(4.20) \quad e^{-\mathbf{a}} \cdot \mathcal{W}_{\tilde{G}} \subset \mathcal{W}_G.$$

Then the following lemma justifies Definition 1:

Lemma 4.1. *For $\mathbf{a} \in w_{1+\infty}^B$ and $\mathcal{W}_G \in \text{Gr}_B^{(0)}$*

$$(4.21) \quad e^{\mathbf{a}} \cdot \mathcal{W}_G = \mathcal{W}_{e^{W_{\mathbf{a}}^B} G}.$$

4.2. KdV vs BKP. In [Ale20b] we prove that after a simple redefinition of times any tau-function of the KdV hierarchy solves the BKP hierarchy:

Theorem 1 ([Ale20b]). *For any KdV tau-function*

$$(4.22) \quad \tau_{\text{BKP}}(\mathbf{t}) = \tau_{\text{KdV}}(\mathbf{t}/2)$$

is a tau-function of the BKP hierarchy.

The KdV hierarchy is a 2-reduction of the KP hierarchy. Let us consider a tau-function of KdV hierarchy τ_{KdV} and the point $\mathcal{W}^{\text{KdV}} \in \text{Gr}^{(0)}$ associated to it as a tau-function of the KP hierarchy. If we consider the same tau-function as a tau-function of the BKP hierarchy, in general it defines another point of the same Sato Grassmannian $\mathcal{W}^{\text{BKP}} \in \text{Gr}_B^{(0)} \subset \text{Gr}^{(0)}$. The reason is that the Baker–Akhiezer functions of two hierarchies have different form.

Indeed, as a tau-function of the KP hierarchy, τ_{KdV} defines the Baker–Akhiezer function by

$$(4.23) \quad \Psi^{\text{KdV}}(z, \mathbf{t}) = e^{\sum_{k \in \mathbb{Z}_{\text{odd}}^+} t_k z^k} \frac{\tau_{\text{KdV}}(\mathbf{t} - [z^{-1}])}{\tau_{\text{KdV}}(\mathbf{t})},$$

where we put $t_{2k} = 0$ for $k > 0$. If we represent it in terms of the corresponding BKP tau-function using Theorem 1, we get

$$(4.24) \quad \Psi^{\text{KdV}}(z, \mathbf{t}/2) = e^{\frac{1}{2} \sum_{k \in \mathbb{Z}_{\text{odd}}^+} t_k z^k} \frac{\tau_{\text{BKP}}(\mathbf{t} - 2[z^{-1}])}{\tau_{\text{BKP}}(\mathbf{t})}.$$

The Baker–Akhiezer function of the same tau-function, considered in the context of the BKP hierarchy, is

$$(4.25) \quad \Psi^{\text{BKP}}(z, \mathbf{t}) = e^{\sum_{k \in \mathbb{Z}_{\text{odd}}^+} t_k z^k} \frac{\tau_{\text{BKP}}(\mathbf{t} - 2[z^{-1}])}{\tau_{\text{BKP}}(\mathbf{t})}.$$

If $\mathcal{W}^{\text{BKP}} \in \text{Gr}_{B^+}^{(0)}$ then at $\mathbf{t} = \mathbf{0}$ these two Baker–Akhiezer functions coincide

$$(4.26) \quad \Phi_1^{\text{KdV}}(z) = \Phi_1^{\text{BKP}}(z) = \tau_{\text{BKP}}(-2[z^{-1}]) =: \Psi(z).$$

However, already the next basis vectors are different:

$$(4.27) \quad \begin{aligned} \Phi_2^{\text{KdV}}(z) &= 2 \frac{\partial}{\partial t_1} \Psi^{\text{KdV}}(z, \mathbf{t}/2) \Big|_{\mathbf{t}=\mathbf{0}} \\ &= z\Psi(z) + 2 \frac{\partial}{\partial t_1} \tau_{\text{BKP}}(\mathbf{t} - 2[z^{-1}]) \Big|_{\mathbf{t}=\mathbf{0}} - 2\Psi(z) \frac{\partial}{\partial t_1} \tau_{\text{BKP}}(\mathbf{t}) \Big|_{\mathbf{t}=\mathbf{0}}, \end{aligned}$$

and

$$(4.28) \quad \begin{aligned} \Phi_2^{\text{BKP}}(z) &= \frac{\partial}{\partial t_1} \Psi^{\text{BKP}}(z, \mathbf{t}/2) \Big|_{\mathbf{t}=\mathbf{0}} \\ &= z\Psi(z) + \frac{\partial}{\partial t_1} \tau_{\text{BKP}}(\mathbf{t} - 2[z^{-1}]) \Big|_{\mathbf{t}=\mathbf{0}} - \Psi(z) \frac{\partial}{\partial t_1} \tau_{\text{BKP}}(\mathbf{t}) \Big|_{\mathbf{t}=\mathbf{0}}. \end{aligned}$$

Moreover, it is clear that this difference, in general, cannot be compensated by Ψ . Hence, in general,

$$(4.29) \quad \mathcal{W}^{\text{KdV}} \neq \mathcal{W}^{\text{BKP}}.$$

Therefore, the Kac–Schwarz algebras of a KdV tau-function, considered in the frameworks of the KP and BKP hierarchies, are different.

5. GENERALIZED BRÉZIN–GROSS–WITTEN MODEL

5.1. Intersection theory. Denote by $\overline{\mathcal{M}}_{g,n}$ the Deligne–Mumford compactification of the moduli space of all compact Riemann surfaces of genus g with n distinct marked points. It is a non-singular complex orbifold of dimension $3g - 3 + n$. It is empty unless the stability condition

$$(5.1) \quad 2g - 2 + n > 0$$

is satisfied.

For each marking index i consider the cotangent line bundle $\mathbb{L}_i \rightarrow \overline{\mathcal{M}}_{g,n}$, whose fiber over a point $[\Sigma, z_1, \dots, z_n] \in \overline{\mathcal{M}}_{g,n}$ is the complex cotangent space $T_{z_i}^* \Sigma$ of Σ at z_i . Let $\psi_i \in H^2(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ denote the first Chern class of \mathbb{L}_i . We consider the intersection numbers

$$(5.2) \quad \langle \tau_{a_1} \tau_{a_2} \cdots \tau_{a_n} \rangle_g := \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{a_1} \psi_2^{a_2} \cdots \psi_n^{a_n}.$$

The integral on the right-hand side of (5.2) vanishes unless the stability condition (5.1) is satisfied, all a_i are non-negative integers, and the dimension constraint

$$(5.3) \quad 3g - 3 + n = \sum_{i=1}^n a_i$$

holds true. Let $T_i, i \geq 0$, be formal variables and let

$$(5.4) \quad \tau_{KW} := \exp \left(\sum_{g=0}^{\infty} \sum_{n=0}^{\infty} \hbar^{2g-2+n} F_{g,n} \right),$$

where

$$(5.5) \quad F_{g,n} := \sum_{a_1, \dots, a_n \geq 0} \langle \tau_{a_1} \tau_{a_2} \cdots \tau_{a_n} \rangle_g \frac{\prod T_{a_i}}{n!}.$$

Witten's conjecture [Wit91], proved by Kontsevich [Kon92], states that the partition function τ_{KW} becomes a tau-function of the KdV hierarchy after the change of variables $T_n = (2n + 1)!! t_{2n+1}$. This is the Kontsevich–Witten tau-function.

In this paper we will focus on a different version of intersection theory on the moduli spaces, which is also governed by the KdV hierarchy, and was recently introduced by Norbury [Nor17]. Namely, he introduced Θ -classes, $\Theta_{g,n} \in H^{4g-4+2n}(\overline{\mathcal{M}}_{g,n})$, and described their intersections with the ψ -classes

$$(5.6) \quad \langle \tau_{a_1} \tau_{a_2} \cdots \tau_{a_n} \rangle_g^\Theta = \int_{\overline{\mathcal{M}}_{g,n}} \Theta_{g,n} \psi_1^{a_1} \psi_2^{a_2} \cdots \psi_n^{a_n}.$$

Again, the integral on the right-hand side vanishes unless the stability condition (5.1) is satisfied, all a_i are non-negative integers, and the dimension constraint

$$(5.7) \quad g - 1 = \sum_{i=1}^n a_i$$

holds true. Consider the generating function of the intersection numbers of Θ -classes and ψ -classes

$$(5.8) \quad F_{g,n}^\Theta = \sum_{a_1, \dots, a_n \geq 0} \frac{\prod T_{a_i}}{n!} \int_{\overline{\mathcal{M}}_{g,n}} \Theta_{g,n} \psi_1^{a_1} \psi_2^{a_2} \cdots \psi_n^{a_n}$$

then, we have a direct analog of the Kontsevich–Witten tau-function:

Theorem ([Nor17]). *Generating function*

$$(5.9) \quad \tau_\Theta = \exp \left(\sum_{g=0}^{\infty} \sum_{n=0}^{\infty} \hbar^{2g-2+n} F_{g,n}^\Theta \right)$$

becomes a tau-function of the KdV hierarchy after the change of variables $T_n = (2n + 1)!! t_{2n+1}$.

Norbury also proved, that τ_Θ is nothing but a tau-function of the Brézin–Gross–Witten (BGW) model

$$(5.10) \quad \tau_\Theta = \tau_{BGW}.$$

We refer the reader to [Nor17, Nor20] for a detailed presentation.

The BGW tau-function can be described by a matrix model. This matrix model description has a natural one-parameter deformation, which we will consider below. Let $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_M)$ be a diagonal matrix. For any function f , dependent on the infinite set of variables $\mathbf{t} = (t_1, t_3, t_5, \dots)$, let

$$(5.11) \quad f([\Lambda^{-1}]) := f(\mathbf{t}) \Big|_{t_k = \frac{1}{k} \text{Tr} \Lambda^{-k}}$$

be the *Miwa parametrization*. The *Generalized Brézin–Gross–Witten model* was introduced in [MMS96] and further investigated in [Ale18]. In the Miwa parametrization it is given by the asymptotic expansion of the matrix integral

$$(5.12) \quad \tau_{BGW}([\Lambda^{-1}], N) := \tilde{\mathcal{C}}^{-1} \int [d\Phi] \exp \left(-\frac{1}{2\hbar} \text{Tr} (\Lambda^2 \Phi + \Phi^{-1} + 2\hbar(N - M) \log \Phi) \right),$$

where one integrates over the normal $M \times M$ matrices.

Remark 5.1. *Let us note that to make the normalization compatible with the one of Norbury, we substitute \hbar of [Ale18] with $2\hbar$.*

For $N = 0$ the generalized BGW tau-function reduces to the original BGW model, $\tau_{BGW}(\mathbf{t}, 0) = \tau_{BGW}(\mathbf{t})$. For arbitrary N , this is a tau-function of the KdV hierarchy. Moreover, the tau-functions for different values of the parameter N (not to be confused with M , the size of the matrices) are related to each other by the MKP hierarchy. The first terms of the series expansion of $\tau_{BGW}(\mathbf{t}, N)$ were described in [Ale18]. The Virasoro constraints for the generalized BGW model can be derived with the help of the Kac–Schwarz approach [Ale18]. These Virasoro constraints lead to the cut-and-join description

$$(5.13) \quad \tau_{BGW}(\mathbf{t}, N) = e^{\hbar \widehat{W}_0(N)} \cdot 1,$$

with the cut-and-join operator

$$(5.14) \quad \widehat{W}_0(N) = \sum_{k, m \in \mathbb{Z}_{\text{odd}}^+} \left(k m t_k t_m \frac{\partial}{\partial t_{k+m-1}} + \frac{1}{2} (k + m + 1) t_{k+m+1} \frac{\partial^2}{\partial t_k \partial t_m} \right) + \left(\frac{1}{8} - \frac{N^2}{2} \right) t_1.$$

5.2. Cut-and-join operator and BKP tau-function. In [Ale20a] it was proven that after a transformation $t_k \mapsto t_k/2$ the cut-and-join operator (5.14) acquires the form

$$(5.15) \quad \widehat{W}_0(N) = \frac{1}{4} \widehat{M}_{-1}^B + \left(\frac{1}{16} - \frac{N^2}{4} \right) \widehat{J}_{-1}^B,$$

where the generators \widehat{J}_{-1}^B and \widehat{M}_{-1}^B are given by (3.33) and (3.35). Hence, it is an element of the BKP symmetry algebra, and $\tau_{BGW}(\mathbf{t}/2, N)$ is a tau-function of the BKP hierarchy.

Remark 5.2. *The BKP integrability also immediately follows from the KdV one, see Theorem 1.*

In the basis (3.37) the cut-and-join operator has the form

$$(5.16) \quad \widehat{W}_0(N) = \frac{1}{4}\widehat{W}_{-1,2}^B + \left(\frac{1}{16} - \frac{N^2}{4}\right)\widehat{W}_{-1,0}^B,$$

and is equal to $\widehat{W}_{\mathfrak{w}_0(N)}^B$ with

$$(5.17) \quad \mathfrak{w}_0(N) = -\frac{1}{2}\left(z\partial_z^2 + \left(\frac{1}{4} - N^2\right)z^{-1}\right) \in w_{1+\infty}^B.$$

In this section we will find another expression for the BKP tau-function $\tau_{BGW}(\mathfrak{t}/2, N)$ in terms of the group elements of the $W_{1+\infty}^B$ algebra.

Let

$$(5.18) \quad F(z; a) := \sum_{k=1}^{\infty} \frac{B_{k+1}(a + 1/2)}{(k+1)kz^k},$$

where $B_k(z)$ are the *Bernoulli polynomials*. This series appears in Stirling's expansion of the gamma function,

$$(5.19) \quad \Gamma(z + 1/2 - a) \sim \sqrt{2\pi}z^{z-a}e^{-z}e^{F(z;a)},$$

valid for large values of $|z|$ with $|\arg(z)| < \pi$. Here a is an arbitrary finite complex number. Let us consider the group operator of the algebra $w_{1+\infty}$

$$(5.20) \quad \mathfrak{0}_B(N) = \exp\left(\log(-N^2)D + 2 \sum_{k \in \mathbb{Z}_{\text{even}}^+} \frac{B_{k+1}(D + 1/2)}{(k+1)kN^k}\right),$$

where

$$(5.21) \quad D := z\partial_z.$$

It can be obtained from the asymptotic expansion of a combination of the gamma-functions

$$(5.22) \quad \mathfrak{0}_B(N) \sim \frac{\Gamma(1/2 + N)}{\Gamma(D + 1/2 + N)} \frac{\Gamma(1/2 - N)}{\Gamma(D + 1/2 - N)}.$$

It is easy to see that

$$(5.23) \quad D^k \in w_{1+\infty}^B, \quad \text{for } k \in \mathbb{Z}_{\text{odd}}^+.$$

Since $B_k(x + 1/2) \in x\mathbb{Q}[x^2]$ for odd k , operator $\log(\mathfrak{0}_B(N))$ is the odd series in D , therefore,

$$(5.24) \quad \log(\mathfrak{0}_B(N)) \in w_{1+\infty}^B.$$

The operator $\mathfrak{w}_0(N)$ can be related to the operator $-z^{-1}/2$ by a conjugation:

Lemma 5.1.

$$(5.25) \quad \mathfrak{w}_0(N) = -\mathfrak{0}_B(N) \frac{z^{-1}}{2} \mathfrak{0}_B(N)^{-1}.$$

Proof. One can rewrite the operator $\mathfrak{w}_0(N)$ in terms of D and z^{-1} ,

$$(5.26) \quad \mathfrak{w}_0(N) = -\frac{z^{-1}}{2} \left(D^2 - D + \frac{1}{4} - N^2\right).$$

Then

$$\begin{aligned}
(5.27) \quad \mathfrak{o}_B(N) \frac{z^{-1}}{2} \mathfrak{o}_B(N)^{-1} &= \frac{z^{-1}}{2} \frac{\Gamma(D + 1/2 + N) \Gamma(D + 1/2 - N)}{\Gamma(D - 1/2 + N) \Gamma(D - 1/2 - N)} \\
&= \frac{z^{-1}}{2} (D - 1/2 + N)(D - 1/2 - N) \\
&= \frac{z^{-1}}{2} \left(D^2 - D + \frac{1}{4} - N^2 \right).
\end{aligned}$$

□

Let us consider the group element of the $W_{1+\infty}^B$ algebra

$$(5.28) \quad \widehat{O}_B(N) := e^{\widehat{W}_{\log(\mathfrak{o}_B(N))}^B}.$$

It is obvious that $\widehat{O}_B(N) \cdot 1 = 1$, then from Lemma 5.1 and cut-and-join representation (5.13) we have

Theorem 2. *Tau-function of the generalized BGW model is a solution of the BKP hierarchy, given by*

$$(5.29) \quad \tau_{BGW}(\mathbf{t}/2, N) = \widehat{O}_B(N) \cdot e^{\frac{\hbar \mathbf{t}_1}{4}}.$$

Using Lemma 4.1 we can get the basis vectors for the point of the BKP Sato Grassmannian $\text{Gr}_{B+}^{(0)}$, associated with $\tau_{BGW}(\mathbf{t}/2, N)$. For the basis vectors normalized by $\Phi_k = z^{k-1}(1 + O(z^{-1}))$ one has

$$\begin{aligned}
(5.30) \quad \Phi_k &= \frac{\Gamma(k - 1/2 + N) \Gamma(k - 1/2 - N)}{\Gamma(D + 1/2 + N) \Gamma(D + 1/2 - N)} \cdot e^{-\frac{\hbar}{2z}} z^{k-1} \\
&= \frac{\Gamma(k - 1/2 + N) \Gamma(k - 1/2 - N)}{\Gamma(D + 1/2 + N) \Gamma(D + 1/2 - N)} \cdot \sum_{m=0}^{\infty} (-\hbar)^m \frac{z^{k-m-1}}{2^m m!} \\
&= \sum_{m=0}^{\infty} (-\hbar)^m \frac{\Gamma(k - 1/2 + N) \Gamma(k - 1/2 - N)}{\Gamma(k - m - 1/2 + N) \Gamma(k - m - 1/2 - N)} \frac{z^{k-m-1}}{2^m m!}.
\end{aligned}$$

Remark 5.3. *Expression for Φ_1 coincides with the expression for the first basis vector of the generalized BGW tau-function, obtained within the framework of KP in [Ale18] see formula (77), if one substitutes \hbar with $2\hbar$ and λ with $-z$. However, the points of the Sato Grassmannian associated with KP and BKP frameworks are different, see Section 4.2.*

5.3. Schur Q-functions and BKP hierarchy. Schur Q-functions were introduced by Schur [Sch11] for the description of the projective representations of the symmetric groups. These functions are labeled by strict partitions $\lambda \in \text{DP}$. The definition and description of the Schur Q-functions can be found in Section 3.8 of Macdonald's book [Mac95]. In many aspects the Schur Q-functions are similar to the ordinary Schur functions. For example, let us mention the *Cauchy formula*

$$(5.31) \quad \sum_{\lambda \in \text{DP}} \frac{Q_\lambda(\mathbf{t}) Q_\lambda(\mathbf{s})}{2^{\ell(\lambda)}} = \exp \left(2 \sum_{k \in \mathbb{Z}_{\text{odd}}^+} k t_k s_k \right),$$

and an analog of the standard hook formula [Sch11]

$$(5.32) \quad Q_\lambda(\delta_{k,1}) = 2^{|\lambda|} \frac{1}{\prod_{j=1}^{\ell(\lambda)} \lambda_j!} \prod_{k < m} \frac{\lambda_k - \lambda_m}{\lambda_k + \lambda_m}.$$

Schur Q-functions are closely related to the BKP hierarchy. Let us consider the class of hypergeometric solutions of the BKP hierarchy introduced by Orlov [Or103]. It was shown by Mironov, Morozov, and Natanzon [MMN20] that the hypergeometric solutions can be interpreted as the generating functions of the spin Hurwitz numbers. Hypergeometric solutions of the 2-component BKP hierarchy are given by the following sums over strict partitions

$$(5.33) \quad \tau(\mathbf{t}, \mathbf{s}) = \sum_{\lambda \in \text{DP}} 2^{-\ell(\lambda)} r_\lambda Q_\lambda(\mathbf{t}/2) Q_\lambda(\mathbf{s}/2),$$

where

$$(5.34) \quad r_\lambda = \prod_{j=1}^{\ell(\lambda)} r(1)r(2) \dots r(\lambda_j)$$

for some function $r(z)$, which we call the *weight generating function*.

Remark 5.4. *Let us note that if $r(k) = 0$ for some $k \in \mathbb{Z}^+$, then the sum in (5.33) is finite and contains only partitions with $\lambda_1 < k$.*

Relation between Schur Q-functions and BKP hierarchy, in particular, is described by the following result of You:

Theorem 3 ([You89]). *For the states (2.8) the boson-fermion correspondence yields*

$$(5.35) \quad \sigma_B^0(|\lambda\rangle) = 2^{-\ell(\lambda)/2} Q_\lambda(\mathbf{t}/2).$$

Schur Q-functions are the polynomial solutions of the BKP hierarchy.

5.4. Schur Q-function expansion of the generalized BGW tau-function. In [Ale20a] we suggest the following:

Conjecture 5.1.

$$(5.36) \quad \tau_{BGW}(\mathbf{t}) = \sum_{\lambda \in \text{DP}} \left(\frac{\hbar}{16} \right)^{|\lambda|} \frac{Q_\lambda(\mathbf{t}) Q_\lambda(\delta_{k,1})^3}{2^{\ell(\lambda)} Q_{2\lambda}(\delta_{k,1})^2}.$$

For the generalized BGW model with arbitrary N in [Ale20a] we conjecture a Schur Q-function expansion, which generalizes (5.36). This expansion is of the hypergeometric type (5.33), and the author conjectured in [Ale20a] that the generalized BGW tau-function, after a simple rescaling of times, is a hypergeometric solution of the BKP hierarchy. We prove this conjecture below, see Theorem 4, and this is one of the most important results of this paper.

It was also noted in [Ale20a], see Remark 3.1, that the analogous to (5.36) Schur Q-function expansion for the Kontsevich–Witten tau-function was established in a series of papers [DFIZ93, J95, MM20]. In [Ale20a] we conjecture that this tau-function is also a hypergeometric solution of the BKP hierarchy. The statement of this conjecture follows from the Schur Q-function expansion of [DFIZ93, J95, MM20] and the relations for the values of the Schur Q-function on special loci proven in [MMNO20] (see also [LY21]).

Let us use Theorem 3 to describe the Schur Q-function expansion of the generalized BGW model. From (3.16) we have

$$(5.37) \quad \widehat{W}_{D^k}^B \cdot \langle 0 | e^{J_+(\mathbf{t})} = \langle 0 | e^{J_+(\mathbf{t})} \frac{1}{2} \sum_{m \in \mathbb{Z}} (-1)^m m^k : \phi_{-m} \phi_m : .$$

Both sides of this identity vanish for even k . For the group operator we have

$$(5.38) \quad \widehat{O}_B(N) \cdot \langle 0 | e^{J_+(\mathbf{t})} = \langle 0 | e^{J_+(\mathbf{t})} O(N),$$

where

$$(5.39) \quad O(N) = \exp \left(\sum_{m \in \mathbb{Z}} (-1)^m \left(\frac{1}{2} \log(-N^2) m + \sum_{k \in \mathbb{Z}_{\text{even}}^+} \frac{B_{k+1}(m+1/2)}{(k+1)kN^k} \right) : \phi_{-m} \phi_m : \right).$$

The tau-function is given by the vacuum expectation value

$$(5.40) \quad \tau_{BGW}(\mathbf{t}/2, N) = \langle 0 | e^{J+(\mathbf{t})} O(N) e^{\frac{\hbar}{2} J_{-1}} | 0 \rangle.$$

Operators $\widehat{W}_{D^k}^B$ are diagonal operators of the $\mathfrak{go}(\infty)$ algebra. For odd k using anti-commutation relations for (2.8) we have

$$(5.41) \quad \frac{1}{2} \sum_{m \in \mathbb{Z}} (-1)^m m^k : \phi_{-m} \phi_m : |\lambda\rangle = - \sum_{j=1}^{\ell(\lambda)} \lambda_j^k |\lambda\rangle,$$

hence, from Theorem 3 for odd k one has

$$(5.42) \quad \widehat{W}_{D^k}^B \cdot Q_\lambda(\mathbf{t}/2) = - \sum_{j=1}^{\ell(\lambda)} \lambda_j^k Q_\lambda(\mathbf{t}/2).$$

Let us introduce

$$(5.43) \quad r^{(N)}(z) = \frac{(2z-1)^2 - 4N^2}{4}.$$

Theorem 2 leads to the proof of Conjecture 2 of [Ale20a]:

Theorem 4. *The partition function of the generalized BGW model in the properly normalized times, $\tau_{BGW}(\mathbf{t}/2, N)$ is a hypergeometric tau-function of the BKP hierarchy (5.33) with $t_k^* = 2\delta_{k,1}$.*

$$(5.44) \quad \tau_{BGW}(\mathbf{t}/2, N) = \sum_{\lambda \in \text{DP}} \left(\frac{\hbar}{4} \right)^{|\lambda|} r_\lambda^{(N)} \frac{Q_\lambda(\mathbf{t}/2) Q_\lambda(\delta_{k,1})}{2^{\ell(\lambda)}}.$$

Proof. Using Cauchy formula (5.31) one gets

$$(5.45) \quad e^{\frac{\hbar t_1}{4}} = \sum_{\lambda \in \text{DP}} \left(\frac{\hbar}{4} \right)^{|\lambda|} \frac{Q_\lambda(\mathbf{t}/2) Q_\lambda(\delta_{k,1})}{2^{\ell(\lambda)}}.$$

From Theorem 2 and formula (5.42) we have

$$(5.46) \quad \begin{aligned} \tau_{BGW}(\mathbf{t}/2, N) &= \widehat{O}_B(N) \cdot \sum_{\lambda \in \text{DP}} \left(\frac{\hbar}{4} \right)^{|\lambda|} \frac{Q_\lambda(\mathbf{t}/2) Q_\lambda(\delta_{k,1})}{2^{\ell(\lambda)}} \\ &= \sum_{\lambda \in \text{DP}} \left(\frac{\hbar}{4} \right)^{|\lambda|} \frac{Q_\lambda(\mathbf{t}/2) Q_\lambda(\delta_{k,1})}{2^{\ell(\lambda)}} \prod_{j=1}^{\ell(\lambda)} \frac{\Gamma(\lambda_j + 1/2 + N) \Gamma(\lambda_j + 1/2 - N)}{\Gamma(1/2 + N) \Gamma(1/2 - N)}. \end{aligned}$$

This formula is equivalent to that of Theorem. □

Let us stress that for the BGW tau-function ($N = 0$) the statement of this theorem reduces to equation (5.36), because from (5.32) we have

$$(5.47) \quad \frac{Q_\lambda(\delta_{k,1})}{Q_{2\lambda}(\delta_{k,1})} = \prod_{j=1}^{\ell(\lambda)} (2\lambda_j - 1)!!.$$

For $N = k + 1/2$ with $k \in \mathbb{Z}_{\geq 0}$ the weight generating function (5.43) vanishes for $z = k + 1$, $r^{(k+1/2)}(k+1) = 0$, hence, in this case the tau-function τ_{BGW} is a polynomial. These polynomial tau-functions are given by certain shifted Schur functions for the triangular partitions [Ale18], which coincide with the shifted Schur Q-functions [Ale20a].

The formula in Theorem 4 is a direct analog of the Schur function expansion of the generating function for the strictly monotone Hurwitz numbers or Grothendieck's dessins d'enfant. We claim that the generalized BGW model is a generating function of the spin version of the disconnected strictly monotone Hurwitz numbers. In this context the cut-and-join description of the generalized BGW model is a direct analog of the Zograf cut-and-join description of Grothendieck's dessins d'enfant [Zog15]. We will discuss the details of this identification with the spin Hurwitz numbers elsewhere.

6. HIGHER MODELS

In this section we discuss a family of hypergeometric tau-functions for the 2-component BKP hierarchy. This family includes the model, considered in the previous section, and can serve as a deep generalization of this model.

First, let us note that the operator $\widehat{O}_B(N)$ is a group operator, therefore it is invertible. Consider

$$(6.1) \quad \tau_{a,b}(\mathbf{t}, \mathbf{s}, \mathbf{u}, \mathbf{w}) = \frac{\widehat{O}_B(u_a) \widehat{O}_B(u_{a-1}) \cdots \widehat{O}_B(u_1)}{\widehat{O}_B(w_b) \widehat{O}_B(w_{b-1}) \cdots \widehat{O}_B(w_1)} \cdot e^{\frac{1}{2} \sum_{k \in \mathbb{Z}_{\text{odd}}^+} k t_k s_k},$$

where u_k and w_k are formal parameters. We will focus on the case when the number of the parameters u_k and w_k is finite, however, many formulas below are valid also for the general case with infinitely many parameters. We regard (6.1) as a formal power series in the variables \mathbf{t} and \mathbf{s} , and parameters \mathbf{u} and \mathbf{w} . By construction, this is a tau-function of 2-component BKP hierarchy. The definition immediately implies the free fermion formula

$$(6.2) \quad \tau_{a,b}(\mathbf{t}, \mathbf{s}, \mathbf{u}, \mathbf{w}) = \langle 0 | e^{J^+(\mathbf{t})} O(u_a) \cdots O(u_1) O^{-1}(w_b) \cdots O^{-1}(w_1) e^{J^-(\mathbf{s})} | 0 \rangle,$$

where the diagonal operators O are given by (5.39).

For $a = 1$, $b = 0$ and $s_j = \delta_{j,1} \hbar/2$ it reduces to the generalized BGW tau-function, considered above

$$(6.3) \quad \tau_{1,0}(\mathbf{t}, \delta_{k,1} \hbar/2, N, 0) = \tau_{BGW}(\mathbf{t}/2, N).$$

In (6.1) we do not introduce the expansion parameter \hbar , it can be restored by rescaling of the variables \mathbf{t} , \mathbf{s} and parameters \mathbf{u} , \mathbf{w} .

Operators $\widehat{O}_B(N_m)$ commute with each other, hence, their order in (6.1) is not important

Lemma 6.1. *For all u_1, u_2*

$$(6.4) \quad [\widehat{O}_B(u_1), \widehat{O}_B(u_2)] = 0.$$

Proof. From (5.28) we see that $\log(\widehat{O}_B(N))$ is a linear combination of $\widehat{W}_{D^k}^B$ with $k \in \mathbb{Z}_{\text{odd}}^+$. Lemma 3.3 implies the identity

$$(6.5) \quad [\widehat{W}_{D^k}^B, \widehat{W}_{D^m}^B] = 0$$

for all $k, m \in \mathbb{Z}_{\text{odd}}^+$, from which the statement of the lemma immediately follows. \square

By analogy with Theorem 4 for the weight generating function, given by (5.43), we have

Proposition 6.2.

$$(6.6) \quad \tau_{a,b}(\mathbf{t}, \mathbf{s}, \mathbf{u}, \mathbf{w}) = \sum_{\lambda \in \text{DP}} \frac{r_\lambda^{(u_1)} r_\lambda^{(u_2)} \dots r_\lambda^{(u_a)}}{r_\lambda^{(w_1)} r_\lambda^{(w_2)} \dots r_\lambda^{(w_b)}} \frac{Q_\lambda(\mathbf{t}/2) Q_\lambda(\mathbf{s}/2)}{2^{-\ell(\lambda)}}$$

Remark 6.1. *It is easy to see that in general (6.1) is not a tau-function of the KdV hierarchy.*

We claim that these tau-functions are the generating functions for the interesting family of the weighted spin Hurwitz numbers, which are the spin analogs to the double weighted Hurwitz numbers with rational weight generating functions, see, e.g., [ACEH20] and references therein. This section contains description of this family of the BKP tau-functions analogous to the description of the corresponding family for the ordinary Hurwitz numbers obtained in [AMMN14, ALS16].

Remark 6.2. *This family of BKP tau-functions for $s_k = \delta_{k,1}$ was introduced by Orlov [Orl03]. He proved, in particular, that this family is related to the hypergeometric solutions of the KP hierarchy by a square root relation. Therefore, the generating functions of the spin Hurwitz numbers (6.1) are related to the corresponding generating functions of the ordinary Hurwitz numbers by a square root relation, completely analogous to the one investigated in [Lee20].*

Remark 6.3. *It would be interesting to construct and investigate the matrix model description of the tau-functions (6.1) similar to the one, constructed in [AC14, AMMN14] for the hypergeometric solutions of the KP hierarchy.*

6.1. Cut-and-join description. Let us derive the cut-and-join description for the tau-functions (6.1). Consider the operator

$$(6.7) \quad \mathbf{w}(\mathbf{u}, \mathbf{w}) = z^{-1} \frac{\prod_{j=1}^a ((D - 1/2)^2 - u_j^2)}{\prod_{j=1}^b ((D - 1/2)^2 - w_j^2)} \in w_{1+\infty}^B.$$

Theorem 5.

$$(6.8) \quad \tau_{a,b}(\mathbf{t}, \mathbf{s}, \mathbf{u}, \mathbf{w}) = e^{-\sum_{k \in \mathbb{Z}_{\text{odd}}^+} s_k \widehat{W}_{\mathbf{w}(\mathbf{u}, \mathbf{w})}^k} \cdot 1$$

Proof. Comparing (3.19), (3.33) and (3.37) we have

$$(6.9) \quad kt_k = -2\widehat{W}_{z^{-k}} \quad \text{for } k \in \mathbb{Z}_{\text{odd}}^+.$$

Therefore

$$(6.10) \quad \tau_{a,b}(\mathbf{t}, \mathbf{s}, \mathbf{u}, \mathbf{w}) = e^{\sum_{j=1}^a \widehat{W}_{\log(0_B(u_j))}^B - \sum_{j=1}^b \widehat{W}_{\log(0_B(w_j))}^B} \cdot e^{-\sum_{k \in \mathbb{Z}_{\text{odd}}^+} s_k \widehat{W}_{z^{-k}}}$$

Repeating the proof of Lemma 5.1 we have

$$(6.11) \quad \frac{\prod_{j=1}^a 0_B(u_j)}{\prod_{j=1}^b 0_B(w_j)} z^{-1} \frac{\prod_{j=1}^b 0_B(w_j)}{\prod_{j=1}^a 0_B(u_j)} = \mathbf{w}(\mathbf{u}, \mathbf{w}),$$

and the statement of the theorem follows from Lemma 3.3. □

Remark 6.4. *From Lemma 3.3 it follows that the operators $\widehat{W}_{\mathbf{w}(\mathbf{u})}^k$ commute with each other*

$$(6.12) \quad [\widehat{W}_{\mathbf{w}(\mathbf{u}, \mathbf{w})}^k, \widehat{W}_{\mathbf{w}(\mathbf{u}, \mathbf{w})}^m] = 0.$$

thus, from Theorem 5 we have the equation

$$(6.13) \quad \frac{\partial}{\partial s_k} \tau_{a,b}(\mathbf{t}, \mathbf{s}, \mathbf{u}, \mathbf{w}) = -\widehat{W}_{\mathbf{w}(\mathbf{u}, \mathbf{w})}^k \cdot \tau_{a,b}(\mathbf{t}, \mathbf{s}, \mathbf{u}, \mathbf{w}).$$

The case $b = 0$ leads to the simple group operators. We expect, that, while the case $b \neq 0$ leads to the more complicated group operators, it still can be described by interesting cut-and-join equations.

By construction, the cut-and-join operator belongs to the the symmetry algebra of the BKP hierarchy. If only a finite number of the variables s_k do not vanish, then the sum $\sum_{k \in \mathbb{Z}_{\text{odd}}^+} s_k \widehat{W}_{\mathbf{w}(\mathbf{u})}^k$ is finite, and for the finite number of parameters \mathbf{u} the operator $\widehat{W}_{\mathbf{w}(\mathbf{u})}^k$ is a finite degree differential operator in variables \mathbf{t} .

Consider the case $s_j = \delta_{j,1} \hbar/2$. It corresponds to the single weighted spin Hurwitz numbers. Then the sum over partitions (6.6) can be expanded as

$$(6.14) \quad \tau_{a,0}(\mathbf{t}, \delta_{j,1} \hbar/2, \mathbf{u}, \mathbf{0}) = \sum_{k=0}^{\infty} \hbar^k \tau_{a,0}^{(k)}(\mathbf{t}, \mathbf{u}),$$

where

$$(6.15) \quad \tau_{a,0}^{(k)}(\mathbf{t}, \mathbf{u}) = \sum_{\lambda \in \text{DP}, |\lambda|=k} r_{\lambda}^{(u_1)} r_{\lambda}^{(u_2)} \dots r_{\lambda}^{(u_a)} \frac{Q_{\lambda}(\mathbf{t}/2) Q_{\lambda}(\delta_{j,1})}{4^{|\lambda|} 2^{-\ell(\lambda)}}.$$

Then the cut-and-join operator provides the recursion in degree of the ramified covering:

$$(6.16) \quad \tau_{a,0}^{(k)}(\mathbf{t}, \mathbf{u}) = -\frac{1}{2k} \widehat{W}_{\mathbf{w}(\mathbf{u})} \cdot \tau_{a,0}^{(k-1)}(\mathbf{t}, \mathbf{u}).$$

We have a similar expansion for $s_j = \delta_{j,k} \hbar/2$ with $k > 1$, it corresponds to the case of orbifold spin Hurwitz numbers.

For the simplest case $a = 1$, $s_j = \delta_{j,1} \hbar/2$ and $u_1 = N$ this operator reduces the the cut-and-join operator (5.16) for the generalized BGW tau-function.

6.2. Sato Grassmannian and Kac–Schwarz operators. Let

$$(6.17) \quad \mathbf{G} = \frac{\prod_{j=1}^a \mathbf{0}_B(u_j)}{\prod_{j=1}^b \mathbf{0}_B(w_j)} e^{-\sum_{k \in \mathbb{Z}_{\text{odd}}^+} s_k z^{-k}}$$

be the group operators of the $w_{1+\infty}^B$ algebra. The point of the Sato Grassmannian, corresponding to the BKP tau-function (6.1) is given by

$$(6.18) \quad \mathcal{W} = \mathbf{G} \cdot \mathcal{W}_0,$$

where $\mathcal{W}_0 = \{1, z, z^2, z^3 \dots\}$ is a point of the Grassmannian for the trivial BKP tau-function $\tau(\mathbf{t}) = 1$.

The normalized basis vectors for the tau-function (6.1) are given by

$$(6.19) \quad \Phi_k = \frac{\prod_{j=1}^b \Gamma(k - 1/2 + w_j) \Gamma(k - 1/2 - w_j)}{\prod_{j=1}^a \Gamma(k - 1/2 + u_j) \Gamma(k - 1/2 - u_j)} \mathbf{G} \cdot z^{k-1}.$$

Let us introduce the operators

$$(6.20) \quad \begin{aligned} \mathbf{p} &:= \mathbf{G} \partial_z \mathbf{G}^{-1}, \\ \mathbf{q} &:= \mathbf{G} z \mathbf{G}^{-1}. \end{aligned}$$

These are the Kac-Schwarz operators

$$(6.21) \quad \mathbf{p} \cdot \mathcal{W} \subset \mathcal{W}, \quad \mathbf{q} \cdot \mathcal{W} \subset \mathcal{W},$$

satisfying the canonical commutation relation

$$(6.22) \quad [\mathbf{p}, \mathbf{q}] = 1.$$

Let us introduce

$$(6.23) \quad \mathbf{R}_m = \frac{\prod_{j=1}^b ((D + 1/2 - m)^2 - w_j^2)}{\prod_{j=1}^a ((D + 1/2 - m)^2 - u_j^2)}.$$

For operators (6.20) we have

$$(6.24) \quad \mathbf{q} = \frac{\prod_{j=1}^a \mathbf{0}_B(u_j)}{\prod_{j=1}^b \mathbf{0}_B(w_j)} \approx \frac{\prod_{j=1}^b \mathbf{0}_B(w_j)}{\prod_{j=1}^a \mathbf{0}_B(u_j)} = z \mathbf{R}_0$$

and

$$(6.25) \quad \begin{aligned} \mathbf{p} &= \frac{\prod_{j=1}^a ((D + 1/2)^2 - u_j^2)}{\prod_{j=1}^b ((D + 1/2)^2 - w_j^2)} \partial_z - \sum_{k \in \mathbb{Z}_{\text{odd}}^+} k s_k \mathbf{q}^{-k-1} \\ &= \mathbf{R}_0^{-1} \partial_z - \sum_{k \in \mathbb{Z}_{\text{odd}}^+} k s_k z^{-k-1} \prod_{m=1}^{k+1} \mathbf{R}_m^{-1}. \end{aligned}$$

By construction for the wave function $\Psi = \Phi_1$ from (6.19) we have

$$(6.26) \quad \mathbf{p} \cdot \Psi(z) = 0.$$

However, it is more convenient to introduce the operator

$$(6.27) \quad \mathbf{c} = \mathbf{q}\mathbf{p} = D - \sum_{k \in \mathbb{Z}_{\text{odd}}^+} k s_k \mathbf{q}^{-k}$$

which also annihilates the wave function, $\mathbf{c} \cdot \Psi = 0$. Let us consider the case when $s_\ell \neq 0$ for some finite odd ℓ and $s_k = 0$ for $k > \ell$. If $b = 0$, then the operator \mathbf{c} is a finite order differential operator with coefficients in $\mathbb{C}[z, z^{-1}]$. If $b > 0$ we introduce

$$(6.28) \quad \mathbf{K} = \prod_{r=1}^{\ell} \prod_{j=1}^b ((D + r - 1/2)^2 - w_j^2).$$

Then for the operator $\mathbf{A} := \mathbf{K}\mathbf{c}$ we have

$$(6.29) \quad \mathbf{A} = D\mathbf{K} - \sum_{k=1}^{\ell} k s_k z^{-k} \prod_{r=1}^{\ell-k} \prod_{j=1}^b ((D + r - 1/2)^2 - w_j^2) \prod_{m=0}^{k-1} \prod_{j=1}^a ((D - m - 1/2)^2 - u_j^2).$$

It is a polynomial of z^{-1} and D , which annihilates the wave function

$$(6.30) \quad \mathbf{A} \cdot \Psi(z) = 0.$$

We can interpret it as the *quantum spectral curve*. For $\mathbf{K} \neq 1$ this operator, in general, is not a Kac-Schwarz operator.

Conjecture 6.1. *The n -point functions for the tau-function (6.1) can be given by the topological recursion, associated with the semi-classical limit of the quantum spectral curve (6.29).*

6.3. Examples. The case of the generalized BGW tau-function, considered in Section 5, corresponds to $a = 1$, $b = 0$, $s_j = \delta_{j,1}\hbar/2$. In this case

$$(6.31) \quad \mathbf{R}_m = \frac{1}{(D + 1/2 - m)^2 - u^2}.$$

The Kac–Schwarz operators, introduced in the previous section, are

$$(6.32) \quad \begin{aligned} \mathbf{p} &= ((D + 1/2)^2 - u^2)\partial_z - \frac{\hbar}{2}z^{-2}\mathbf{R}_1^{-1}\mathbf{R}_2^{-1}, \\ \mathbf{q} &= z\frac{1}{(D + 1/2)^2 - u^2}, \end{aligned}$$

the operator

$$(6.33) \quad \mathbf{c} = D - \frac{\hbar}{2}z^{-1}((D - 1/2)^2 - u^2)$$

coincides with the quantum spectral curve operator \mathbf{A} . The quantum spectral curve equation in this case is closely related to the Bessel equation.

We expect, that the spin Hurwitz numbers interpretation of the case with $a = 0$, $b = 1$ is particularly interesting. Namely, this case can be associated with a spin version of the monotone Hurwitz numbers. Let us consider it in more detail for $s_j = \delta_{j,1}\hbar/2$. Then

$$(6.34) \quad \mathbf{R}_m = (D + 1/2 - m)^2 - w^2,$$

and the Kac–Schwarz operators are

$$(6.35) \quad \begin{aligned} \mathbf{p} &= \frac{1}{(D + 1/2)^2 - w^2}\partial_z - \frac{\hbar}{2}z^{-2}\mathbf{R}_1^{-1}\mathbf{R}_2^{-1}, \\ \mathbf{q} &= z((D + 1/2)^2 - w^2), \end{aligned}$$

and

$$(6.36) \quad \mathbf{c} = D - \frac{\hbar}{2}z^{-1}\frac{1}{(D - 1/2)^2 - w^2}.$$

Then the quantum spectral curve is given by the third order differential operator

$$(6.37) \quad \mathbf{A} = ((D - 1/2)^2 - w^2)D - \frac{\hbar}{2}z^{-1}.$$

6.4. Constraints. We expect that the tau-functions (6.1) satisfy the Virasoro and W-constraints, analogous to ones obtained in [KZ15, AMMN14]. Some of these constraints can be derived using the Kac-Schwarz approach for BKP hierarchy. However, not all of them can be derived this way.

The illustrative example here is the generalized BGW tau-function, considered in the previous section. This model is completely described by the Virasoro constraints, derived in [Ale18] using the Kac-Schwarz description of the KP hierarchy. However, these constraints do not appear in the BKP picture. Hence, for the generalized BGW tau-function, the KP and BKP pictures are complimentary to each other: the Virasoro constraints are natural symmetries of the KP hierarchy, while the cut-and-join operator is an element of the BKP symmetry group. We expect that such interplay between different integrable structures should be common for the enumerative geometry tau-functions.

The simplest constraint for the generalized BGW model, which can be obtained from the BKP approach, is given by

$$(6.38) \quad \widehat{W}_c^B \cdot \tau_{BGW}(\mathbf{t}/2, u) = \mu\tau_{BGW}(\mathbf{t}/2, u)$$

where \mathbf{c} is given by (6.33), and μ is some eigenvalue. From the consideration of this equation at $\mathbf{t} = \mathbf{0}$ it is easy to see that $\mu = 0$. The operator

$$(6.39) \quad \widehat{W}_{\mathbf{c}}^B = -\frac{1}{2}\widehat{W}_{0,1}^B + \frac{\hbar}{4}\widehat{W}_{-1,2}^B + \hbar\left(\frac{1}{16} - \frac{N^2}{4}\right)\widehat{W}_{-1,0}^B$$

describes the cut-and-join equation.

REFERENCES

- [AC14] Jan Ambjørn and Leonid Chekhov. The matrix model for dessins d'enfants. *Ann. Inst. Henri Poincaré D*, 1(3):337–361, 2014.
- [ACEH20] A. Alexandrov, G. Chapuy, B. Eynard, and J. Harnad. Weighted Hurwitz numbers and topological recursion. *Comm. Math. Phys.*, 375(1):237–305, 2020.
- [Ale18] A. Alexandrov. Cut-and-join description of generalized Brezin-Gross-Witten model. *Adv. Theor. Math. Phys.*, 22(6):1347–1399, 2018.
- [Ale20a] Alexander Alexandrov. Intersection numbers on $\overline{\mathcal{M}}_{g,n}$ and BKP hierarchy. *arXiv e-prints*, arXiv:2012.07573, December 2020.
- [Ale20b] Alexander Alexandrov. KdV solves BKP. *arXiv e-prints*, arXiv:2012.10448, December 2020.
- [ALS16] A. Alexandrov, D. Lewanski, and S. Shadrin. Ramifications of Hurwitz theory, KP integrability and quantum curves. *J. High Energy Phys.*, (5):124, front matter+30, 2016.
- [AMMN14] A. Alexandrov, A. Mironov, A. Morozov, and S. Natanzon. On KP-integrable Hurwitz functions. *J. High Energy Phys.*, (11):080, front matter+30, 2014.
- [DFIZ93] P. Di Francesco, C. Itzykson, and J.-B. Zuber. Polynomial averages in the Kontsevich model. *Comm. Math. Phys.*, 151(1):193–219, 1993.
- [DJKM82] Etsuro Date, Michio Jimbo, Masaki Kashiwara, and Tetsuji Miwa. Transformation groups for soliton equations. IV. A new hierarchy of soliton equations of KP-type. *Phys. D*, 4(3):343–365, 1981/82.
- [DKM81] Etsuro Date, Masaki Kashiwara, and Tetsuji Miwa. Transformation groups for soliton equations. II. Vertex operators and τ functions. *Proc. Japan Acad. Ser. A Math. Sci.*, 57(8):387–392, 1981.
- [EOP08] Alex Eskin, Andrei Okounkov, and Rahul Pandharipande. The theta characteristic of a branched covering. *Adv. Math.*, 217(3):873–888, 2008.
- [Gun16] Sam Gunningham. Spin Hurwitz numbers and topological quantum field theory. *Geom. Topol.*, 20(4):1859–1907, 2016.
- [J95] Tadeusz Józefiak. Symmetric functions in the Kontsevich-Witten intersection theory of the moduli space of curves. *Lett. Math. Phys.*, 33(4):347–351, 1995.
- [Kon92] Maxim Kontsevich. Intersection theory on the moduli space of curves and the matrix Airy function. *Comm. Math. Phys.*, 147(1):1–23, 1992.
- [KZ15] Maxim Kazarian and Peter Zograf. Virasoro constraints and topological recursion for Grothendieck’s dessin counting. *Lett. Math. Phys.*, 105(8):1057–1084, 2015.
- [Lee20] Junho Lee. A square root of Hurwitz numbers. *Manuscripta Math.*, 162(1-2):99–113, 2020.
- [LY21] Xiaobo Liu and Chenglang Yang. Schur Q-Polynomials and Kontsevich-Witten Tau Function. *arXiv e-prints*, arXiv:2103.14318, March 2021.
- [Mac95] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [MM20] A. Mironov and A. Morozov. Superintegrability of Kontsevich matrix model. *arXiv e-prints*, arXiv:2011.12917, November 2020.
- [MMN20] A. Mironov, A. Morozov, and S. Natanzon. Cut-and-join structure and integrability for spin Hurwitz numbers. *The European Physical Journal C*, 80(2):97, 2020.
- [MMNO20] A. D. Mironov, A. Yu Morozov, S. M. Natanzon, and A. Yu Orlov. Around spin Hurwitz numbers. *arXiv e-prints*, arXiv:2012.09847, December 2020.
- [MMS96] A. Mironov, A. Morozov, and G. W. Semenov. Unitary matrix integrals in the framework of the generalized Kontsevich model. *Internat. J. Modern Phys. A*, 11(28):5031–5080, 1996.
- [Nor17] Paul Norbury. A new cohomology class on the moduli space of curves. *arXiv e-prints*, arXiv:1712.03662, December 2017.
- [Nor20] Paul Norbury. Enumerative geometry via the moduli space of super Riemann surfaces. *arXiv e-prints*, arXiv:2005.04378, May 2020.

- [Orl03] A. Yu. Orlov. Hypergeometric functions associated with Schur Q -polynomials, and the BKP equation. *Teoret. Mat. Fiz.*, 137(2):253–270, 2003.
- [Sch11] J. Schur. Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene lineare Substitutionen. *J. Reine Angew. Math.*, 139:155–250, 1911.
- [SS83] Mikio Sato and Yasuko Sato. Soliton equations as dynamical systems on infinite-dimensional Grassmann manifold. In *Nonlinear partial differential equations in applied science (Tokyo, 1982)*, volume 81 of *North-Holland Math. Stud.*, pages 259–271. North-Holland, Amsterdam, 1983.
- [vdL95] Johan van de Leur. The Adler-Shiota-van Moerbeke formula for the BKP hierarchy. *J. Math. Phys.*, 36(9):4940–4951, 1995.
- [Wit91] Edward Witten. Two-dimensional gravity and intersection theory on moduli space. In *Surveys in differential geometry (Cambridge, MA, 1990)*, pages 243–310. Lehigh Univ., Bethlehem, PA, 1991.
- [You89] Yuching You. Polynomial solutions of the BKP hierarchy and projective representations of symmetric groups. In *Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988)*, volume 7 of *Adv. Ser. Math. Phys.*, pages 449–464. World Sci. Publ., Teaneck, NJ, 1989.
- [Zog15] Peter Zograf. Enumeration of Grothendieck’s dessins and KP hierarchy. *Int. Math. Res. Not. IMRN*, (24):13533–13544, 2015.

IBS CENTER FOR GEOMETRY AND PHYSICS, POHANG UNIVERSITY OF SCIENCE AND TECHNOLOGY (POSTECH), 77 CHEONGAM-RO, NAM-GU, POHANG, GYEONGBUK, 37673, KOREA
Email address: alexandrovsash at gmail.com