

FANO 3-FOLDS AND DOUBLE COVERS BY HALF ELEPHANTS

LIVIA CAMPO

ABSTRACT. We construct a deformation family for each of the 34 Hilbert series of index 2 Fano 3-folds. In 18 cases we construct two different families, distinguished by the topology of their general members.

1. INTRODUCTION

A Fano 3-fold X is a normal projective 3-dimensional variety with ample anticanonical divisor $-K_X$ and \mathbb{Q} -factorial terminal singularities. The Fano index of X is

$$\iota_X := \max\{q \in \mathbb{Z}_{\geq 1} : -K_X \sim qA \text{ for some } A \in \text{Cl}(X)\}.$$

The vector space structure of Fano 3-folds having Fano index 2 is partially understood: [BS07] give a list of possible Hilbert series of these varieties, although that paper does not confirm the existence of any particular one. We call this list $\mathcal{H}_{\text{BS}}^2$.

1.1. Main Theorem. We consider Fano 3-folds that embed anticanonically in codimension 4 into weighted projective spaces. The aim of this paper is to systematically build at least one deformation family for 34 of the Hilbert series in $\mathcal{H}_{\text{BS}}^2$. The strategy is to retrieve them from just as many codimension 4 Fano 3-folds X of index 1 by performing a quotient by a $\mathbb{Z}/2\mathbb{Z}$ action γ on the ambient spaces of these X .

The construction we achieve is summarised in this diagram:

$$(1.1) \quad \begin{array}{ccc} & \text{codim 4} & \text{codim 3} \\ & X & \xleftarrow{\text{unprojection}} Z \\ \text{index 1} & \downarrow \mathbb{Z}/2\mathbb{Z} \gamma & \downarrow \mathbb{Z}/2\mathbb{Z} \gamma \\ & \tilde{X} & \tilde{Z} \\ \text{index 2} & & \end{array}$$

The Main Theorem we prove is

Theorem 1.1. For each of the 34 power series $\mathcal{P} \in \mathcal{H}_{\text{BS}}^2$ there exists an index 1 Fano 3-fold $X \subset w\mathbb{P}^7$ with a $\mathbb{Z}/2\mathbb{Z}$ action γ such that $\tilde{X} := X/\gamma$ is a quasi-smooth Fano 3-fold with Fano index 2, and $\mathcal{P}_{\tilde{X},A} = \mathcal{P}$ for $A \in |-\frac{1}{2}K_{\tilde{X}}|$. For 18 of the Hilbert series in $\mathcal{H}_{\text{BS}}^2$ there are at least two families.

We produce X by Type I [Pap04, PR04] and Type II unprojections [Pap06] from particular Fano 3-folds Z in codimension 3 and from Fano hypersurfaces respectively. The key is to have X invariant under γ . We discuss how and when this is possible in Sections 4, 5, 7. The details are summarised in Table 1 in 1.3.

Key words and phrases. Fano 3-fold, Fano index, Equivariant unprojection.

The author would like to thank Gavin Brown, Tiago Guerreiro, Stavros Argyrios Papadakis, Kaori Suzuki, and Miles Reid for conversations and comments during the development of this work. The author was supported by EPSRC Doctoral Training Partnership and by EPSRC grant EP/N022513/ held by Alexander Kasprzyk.

1.2. Framework. Some constructions of index 2 Fano 3-folds already exist in the literature. In [PR16] Prokhorov and Reid have constructed one example of an index 2 Fano 3-fold in codimension 3 and one in codimension 4 performing divisorial extractions of curves in \mathbb{P}^3 and in a quadric in \mathbb{P}^4 respectively. Their argument is concluded by running the Sarkisov Program initiated by such extractions. Their construction was further generalised by Ducat in [Duc18], who obtained one new family in codimension 4, and one each in codimension 5 and 6.

There are 35 Hilbert series in $\mathcal{H}_{\text{BS}}^2$ associated to codimension 4 Fano 3-folds with Fano index 2 as in the Graded Ring Database (GRDB) [BK⁺, ABR02], of which each realise one using their constructions. A further case¹ is the Hilbert series of the smooth Fano 3-fold with GRDB ID #41028, constructed by Iskovskih in [Isk77, Isk78, Case 13, Table (6.5)]. We added it to Table 1 for completeness.

Of the 34 Hilbert series relative to codimension 4 index 2 Fano 3-folds constructed with our method, 32 admit a double cover X obtained via Type I unprojection from Z in codimension 3. There are between two and four possible distinguished deformation families of X , associated to just as many formats (Tom and Jerry) of the antisymmetric matrix M defining the equations of Z [BKR12a]. Our construction is applicable to one or more of these deformation families: the number changes case by case, and the full extent of the result is outlined in Table 1. Accordingly, \tilde{X} has one or more distinguished deformation families. We give criteria to find deformation families for \tilde{X} in Sections 4 and 5: fundamentally, most information is included in the geometry of Z .

On the other hand, two codimension 4 index 2 Fano 3-folds in the GRDB have a double cover X obtained via Type II₂ unprojection. Performing the unprojection in this case is more complicated, and we refer to [Pap07, Pap06, Tay20]. The double cover construction still works in this case: this is the content of Section 7.

The Hilbert series with GRDB ID #40933, #40663 are in the overlap with the results of [PR16] and [Duc18]. In particular, Ducat finds two different deformation families of #40663, and we retrieve them here with our method (cf [Duc18, Section 3]).

A different approach to the construction of Fano 3-folds is given by Coughlan and Ducat in [CD20], where the authors employ rank 2 cluster algebras to find families of Fano 3-folds in codimension 4 and 5, also in index 2. In addition, in certain cases they are able to determine different deformation families associated to the same Hilbert series (what they call *cluster formats*). Cluster formats mimic the Tom and Jerry formats of [BKR12a]. Our work finds more deformation families that are not realised by cluster formats, and confirms the results in [CD20] when the cluster formats arise.

1.3. Details of the construction. The following table summarises all possible Tom and Jerry types for index 2 Fano 3-folds in codimension 4. In the last column it also records which of the index 1 Jerry formats of [BKR12b] do not give rise to families in index 2. The fraction notation indicates the centre in X in case of ambiguity. The column $\#_1$ denotes the total number of distinguished deformation families constructed by Theorems 4.3 and 5.1. We write "n/a"

¹The Hilbert series with ID #40367 and #40378 also appear in the GRDB in index 2 and codimension 4. However, the Fano 3-folds associated to these two Hilbert series cannot embed in codimension 4 in the weighted projective space suggested by the Graded Ring Database. Indeed, neither our method nor the method in [CD20] construct them. These two Fano 3-folds may exist embedded in higher codimension.

in the cases in which the Tom and Jerry construction is not applicable. In particular, $\#_{II}$ represents the total number of distinguished deformation families when there is no Type I projection. We construct one in Section 7, but there could possibly be others.

TABLE 1. Deformation families of index 2 Fano 3-folds in codimension 4 and corresponding index 1 double covers

Index 2	Index 1	$\#_I$	Tom	Jerry	J-Failures
39557	327	2	T_3	J_{24}	none
39569	512		n/a: no Type I projection. $\#_{II} \geq 1$		
39576	569	1	T_1	none	J_{25}
39578	574	2	T_1	$J_{24} \bullet_{12}$	J_{45}
39605	869	2	T_4	J_{13}	none
39607	872		n/a: no Type I projection. $\#_{II} \geq 1$		
39660	1158	2	T_5	J_{12}	none
39675	1395	2	T_5	none	J_{12}
39676	1401	1	$\frac{1}{5}: T_2$ $\frac{1}{7}: T_4$	none	$\frac{1}{5}: J_{24} \bullet_{12}, J_{45}$ $\frac{1}{7}: J_{12} \bullet_{15}, J_{24}$
39678	1405	1	T_1	none	none
39890	4810	2	T_3	J_{24}	$J_{14} \bullet_{13}$
39898	4896	2	T_3	J_{24}	$J_{14} \bullet_{13}$
39906	4925	1	$\frac{1}{7}(1, 1, 6): T_2$ $\frac{1}{7}(1, 3, 4): T_1$	none	$\frac{1}{7}(1, 1, 6): J_{34}$ $\frac{1}{7}(1, 3, 4): J_{35}$
39912	4938	1	T_2	none	J_{12}
39913	4939	2	$\frac{1}{5}: T_1$ $\frac{1}{7}: T_2$	$\frac{1}{5}: J_{25} \bullet_{24}$ $\frac{1}{7}: J_{14} \bullet_{13}$	$\frac{1}{5}: J_{35}$ $\frac{1}{7}: J_{24}$
39928	4987	1	T_5	none	J_{12}
39929	5000	1	$\frac{1}{5}: T_2$ $\frac{1}{9}: T_4$	none	$\frac{1}{5}: J_{45}$ $\frac{1}{9}: J_{24}$
39934	5052	2	T_1	$J_{23} \bullet_{13}$	J_{24}
39961	5176	1	$\frac{1}{5}: T_2$ $\frac{1}{7}: T_3$	none	$\frac{1}{5}: J_{35}$ $\frac{1}{7}: J_{25}$
39968	5260	2	T_5	J_{13}	none
39969	5266	2	$\frac{1}{5}: T_3$ $\frac{1}{7}: T_4$	$\frac{1}{5}: J_{24} \bullet_{25}$ $\frac{1}{7}: J_{13} \bullet_{15}$	$\frac{1}{5}: J_{45}$ $\frac{1}{7}: J_{34}$
39970	5279	1	$\frac{1}{3}: T_1$ $\frac{1}{5}(1, 1, 4): T_2$ $\frac{1}{5}(1, 2, 3): T_1$	none	$\frac{1}{3}: J_{45}$ $\frac{1}{5}(1, 1, 4): J_{34}$ $\frac{1}{5}(1, 2, 3): J_{35}$
39991	5516	1	$\frac{1}{3}: T_1$ $\frac{1}{7}: T_3$	none	$\frac{1}{3}: J_{45}$ $\frac{1}{7}: J_{13}$
39993	5519	2	$\frac{1}{3}: T_1$ $\frac{1}{5}: T_2$	$\frac{1}{3}: J_{34}$ $\frac{1}{5}: J_{12}$	$\frac{1}{3}: J_{35}$ $\frac{1}{5}: J_{13}$
40360	10963	2	T_3	J_{24}	$J_{14} \bullet_{13}$
40370	11004	1	T_2	none	J_{12}

40371	11005	2	$\frac{1}{3}: T_1$ $\frac{1}{5}: T_2$	$\frac{1}{3}: J_{25} \bullet_{24}$ $\frac{1}{5}: J_{14} \bullet_{13}$	$\frac{1}{3}: J_{35}$ $\frac{1}{5}: J_{24}$
40399	11104	1	T_5	none	J_{24}
40400	11123	1	$\frac{1}{3}: T_3$ $\frac{1}{5}: T_4$	none	$\frac{1}{3}: J_{24} \bullet_{25}, J_{45}$ $\frac{1}{5}: J_{12} \bullet_{15}, J_{24}$
40407	11222	2	T_1	$J_{23} \bullet_{13}$	J_{24}
40663	16206	2	T_4	J_{23}	$J_{12} \bullet_{15}$
40671	16227	1	T_2	none	J_{12}
40672	16246	2	T_2	$J_{15} \bullet_{14}$	J_{25}
40933	24078	2	T_5	J_{12}	T_1
41028	n/a	n/a			

2. BACKGROUND

We work over the field of complex numbers \mathbb{C} . The lack of structure theorems for Fano 3-folds in codimension greater than 3 has forced the search for new approaches to produce their equations explicitly. Unprojections are a technique to retrieve equations for Fano 3-folds in codimension 4 from Fano 3-folds in lower codimension. They were firstly studied by Kustin and Miller [KM83], and later on by Reid and Papadakis [PR04, Pap04]. There are different kinds of unprojections: the most widely employed are called Type I unprojections.

Type I unprojections are initiated by the following type of data.

- A fixed projective plane $D := \mathbb{P}^2(a, b, c) \subset \mathbb{P}^6(a, b, c, d_1, \dots, d_4)$ with coordinates $x_1, x_2, x_3, y_1, \dots, y_4$ respectively, and defined by the ideal $I_D := \langle y_1, y_2, y_3, y_4 \rangle$.
- A family \mathcal{Z} of codimension 3 Fano 3-folds $Z \subset w\mathbb{P}^6$, each defined by the five maximal pfaffians of a skew-symmetric 5×5 syzygy matrix M whose entries (a_{ij}) have weights

$$\begin{pmatrix} m_{12} & m_{13} & m_{14} & m_{15} \\ & m_{23} & m_{24} & m_{25} \\ & & m_{34} & m_{35} \\ & & & m_{45} \end{pmatrix}.$$

In this context, two kinds of formats arise for M , based on conditions on its entries. These are the so-called *Tom and Jerry formats*. If M is in one of these formats, then $D \subset Z \subset w\mathbb{P}^6$

Definition 2.1 ([BKR12a], Definition 2.2). A 5×5 skew-symmetric matrix M is in Tom_k format if and only if each entry a_{ij} for $i, j \neq k$ is in the ideal I_D .

It is in Jerry_{kl} format if and only if $a_{ij} \in I_D$ for either i or j equals k or l . If M is in Jerry_{kl} format, we call *pivot entry* the entry $a_{kl} \in I_D$.

Recall the definition of Fano 3-fold of *Tom type* (respectively, of *Jerry type*).

Definition 2.2 ([Cam20], Definition 2.2). Let X be a codimension 4 index 1 Fano 3-fold X listed in the table [BKR12b]. We say X is of *Tom (Jerry) type* if it is obtained as Type I unprojection of the codimension 3 pair $Z \supset D$ in a Tom (Jerry) family [BKR12a, PR04]. The image of $D \subset Z$ in X is called *Tom (Jerry) centre*: it is a cyclic quotient singularity $p \in X$. In the unprojection setup $D \subset Z$, D is a complete intersection of four linear forms of weight d_1, \dots, d_4 . Such X of Tom (Jerry) type is said to be *general* if $Z \supset D$ is general in its Tom (Jerry) family.

3. DOUBLE COVERS

Let X be a \mathbb{Q} -factorial terminal Fano 3-fold in codimension 4 having Fano index 1. It is anticanonically embedded in a weighted projective space $\mathbb{P}^7(1, b, c, d_1, \dots, d_4, r)$ with homogeneous coordinates $x_1, x_2, x_3, y_1, \dots, y_4, s$. Since $\iota_X = 1$, we assume that the coordinate x_1 has weight 1. Consider the $\mathbb{Z}/2\mathbb{Z}$ action on $w\mathbb{P}^7$ that changes the sign of x_1 .

$$(3.1) \quad \gamma: (x_1, x_2, x_3, y_1, y_2, y_3, y_4, s) \longmapsto (-x_1, x_2, x_3, y_1, y_2, y_3, y_4, s).$$

First we describe the quotient of the ambient space of X by the $\mathbb{Z}/2\mathbb{Z}$ action γ .

Lemma 3.1. *The $\mathbb{Z}/2\mathbb{Z}$ quotient of $\mathbb{P}^7(1, b, c, d_1, \dots, d_4, r)$ via γ is the weighted projective space $\mathbb{P}^7(2, b, c, d_1, \dots, d_4, r)$ with coordinates $\xi, x_2, x_3, y_1, \dots, y_4, s$ respectively, where $\xi := x_1^2$.*

Proof. Consider the affine patches of $\mathbb{P}^7(1, b, c, d_1, \dots, d_4, r) = \text{Proj } \mathbb{C}(x_1, x_2, x_3, y_1, \dots, y_4, s)$. For instance,

$$\mathcal{U}_{x_2} := \{x_2 \neq 0\} = \text{Spec } \mathbb{C}[x_1, \hat{x}_2, x_3, y_1, \dots, y_4, s]^{\mu_b} \cong \mathbb{A}^7 / \mu_b$$

where μ_b is the finite cyclic group of order b . The affine patches relative to the coordinates x_3, y_1, \dots, y_4, s are analogous. They are invariant under γ if and only if x_1 appears with even powers. Such affine patches in which powers of ξ appear are exactly the affine patches of $\mathbb{P}^7(2, b, c, d_1, \dots, d_4, r)$, with the new coordinate ξ . So,

$$\begin{aligned} \mathbb{P}^7(1, b, c, d_1, \dots, d_4, r) / \gamma &= \text{Proj } \mathbb{C}(x_1, x_2, x_3, y_1, \dots, y_4, s)^\gamma = \text{Proj } \mathbb{C}(x_1^2, x_2, x_3, y_1, \dots, y_4, s) \\ &= \mathbb{P}^7(2, b, c, d_1, \dots, d_4, r). \end{aligned}$$

We identify the homogeneous coordinates of X and \tilde{X} and define $\xi := x_1^2$. \square

Hence, for $X \subset \mathbb{P}^7(1, b, c, d_1, \dots, d_4, r)$ to be invariant under γ , the variable x_1 must appear only with even powers. When this is the case, then the quotient $\tilde{X} := X/\gamma$ sits inside the weighted projective space $\mathbb{P}^7(2, b, c, d_1, \dots, d_4, r)$ and is a Fano 3-fold: its anticanonical divisor $-K_{\tilde{X}}$ is a multiple of an ample divisor (see proof of Lemma 3.2), it has terminal singularities (see Lemma 3.4), it is \mathbb{Q} -factorial.

In this section, let us assume that it is possible to realise $X \subset \mathbb{P}^7(1, b, c, d_1, \dots, d_4, r)$ as invariant under γ . We can therefore draw the following conclusions.

Lemma 3.2. *The Fano 3-fold \tilde{X} has index 2.*

Proof. Consider the quotient map $f: X \rightarrow \tilde{X}$. Then, the anticanonical divisor of X is given by $-K_X = -f^*K_{\tilde{X}} - R$ where R is the ramification divisor. In our case, $-K_X = \{x_1 = 0\} \sim \mathcal{O}(1)$, and the ramification divisor is $R = \{x_1 = 0\}$. Therefore, $-f^*K_{\tilde{X}} = 2\{x_1 = 0\}$. This implies that $-K_{\tilde{X}} = \{\xi = 0\} \sim \mathcal{O}(2)$: thus, \tilde{X} has index 2. \square

Lemma 3.3. *If X is quasismooth, then \tilde{X} is quasismooth.*

Proof. Define the variety V_X as

$$V_X := \{p \in w\mathbb{P}^7 : \text{rk}(J_X|_p) < \text{codim}(X)\}.$$

The condition defining V_X is equivalent to looking at the vanishing locus of all 4×4 minors of the Jacobian matrix J_X of X (see [Har77]). By definition, if V_X is empty, then X is quasismooth. Suppose V_X empty. The difference between the Jacobian matrices J_X and $J_{\tilde{X}}$ lies in the column relative to the derivative by x_1 . Suppose $x_1 \neq 0$; then, the rank of J_X is equal to the rank of

$J_{\tilde{X}}$. Now consider the $\frac{\partial}{\partial x_1}$ column of J_X . For each equation f_i of X , $\frac{\partial f_i}{\partial x_1} = \frac{\partial f_i}{\partial \xi} \frac{\partial \xi}{\partial x_1}$, and $\frac{\partial \xi}{\partial x_1} = 2x_1$. Thus, certain entries of the $\frac{\partial}{\partial x_1}$ column of J_X might vanish for $x_1 = 0$, whereas they would be just constant in $J_{\tilde{X}}$. Thus, for $x_1 = 0$ we have that $\text{rk } J_X \leq \text{rk } J_{\tilde{X}}$; so, \tilde{X} is quasismooth if X is. \square

Lemma 3.4. *The Fano 3-fold \tilde{X} has terminal singularities.*

Proof. The fixed locus of the group action γ is $\text{Fix}(\gamma) = \{x_1 = 0\} \cup \mathbb{P}_{\text{even}}$, where \mathbb{P}_{even} is the weighted projective space defined by the vanishing of all the coordinates of $w\mathbb{P}^7$ with odd weight, except for x_1 . We want to study the intersection $X \cap \text{Fix}(\gamma)$.

Firstly, all the cyclic quotient singularities of X lie at coordinate points different from P_{x_1} . Thus, they all lie inside the locus $\{x_1 = 0\}$. So, $X \cap \{x_1 = 0\} \neq \emptyset$. On the other hand, $X \cap \mathbb{P}_{\text{even}} = \emptyset$, because otherwise X would have non-terminal singularities.

Thus, \tilde{X} has terminal singularities, and they are all fixed by γ . \square

The above lemmas prove the following Theorem, which will be crucial in the rest of this paper to prove a criterion for finding the deformation families of \tilde{X} .

Theorem 3.5. *If $X \subset \mathbb{P}^7(1, b, c, d_1, \dots, d_4, r)$ has an invariant realisation under γ , then \tilde{X} is a terminal \mathbb{Q} -factorial Fano 3-fold. Hence, it is in the Graded Ring Database.*

Proof. By Lemma 3.4, \tilde{X} has only terminal singularities, and has Fano index 2 by Lemma 3.2. Therefore, it is listed in the GRDB. \square

In contrast, our method does not produce the index 2 Fano 3-folds in codimension 3 in the Graded Ring Database. Restrict the action γ in 3.1 to the ambient space $w\mathbb{P}^6$ of Z . With a little abuse of notation, we still call the restriction γ .

Proposition 3.6. *The index 2 Fano 3-fold $\tilde{Z} := Z/\gamma$ in codimension 3 is not terminal.*

Proof. Analogously to Lemma 3.4, the fixed locus of γ is $\text{Fix}(\gamma) = \{x_1 = 0\} \cup \mathbb{P}_{\text{even}}$, for \mathbb{P}_{even} is defined by the vanishing of all the coordinates of $w\mathbb{P}^6$ with odd weight, except for x_1 . We study the intersection $X \cap \text{Fix}(\gamma)$. The proof proceeds similarly to the one of Lemma 3.4. The difference is that $Z \cap \mathbb{P}_{\text{even}} \neq \emptyset$.

Indeed, Z and \mathbb{P}_{even} intersect along $D \subset Z$. Terminal singularities on index 2 Fano 3-folds are of the form $\frac{1}{r}(a, b, r - b)$ with $(a, r) = 1$ and $(b, r) = 1$. Since a is always 2 for index 2, this implies that r must be odd. Hence, either b is odd and $r - b$ is even, or vice versa. Thus in either case, D contains a line constituted by $\frac{1}{2}$ singularities. \square

The phenomenon described in Proposition 3.6 was already anticipated in [PR16, Sections 6.4, 6.5]. There are only two Hilbert series corresponding to terminal index 2 codimension 3 Fano 3-folds: one is smooth, and is constructed by Iskovskih in [Isk77] and [Isk78]. The other one was constructed by Ducat in [Duc18].

We would like to stress that X is general in its Tom or Jerry formats, as in Sections 4, 5 (cf [Cam20, Section 3], [BKR12a, Section 4]). Thus, provided the additional condition of invariance of X under γ , \tilde{X} is general.

4. TOM FAMILIES

Assume that Z is a terminal codimension 3 Fano 3-fold having at least one Type I centre, and let X be its Type I unprojection from the corresponding divisor D .

In this section we show a criterion to determine which Tom formats of Z induce a double cover X for the corresponding \tilde{X} in index 2. This produces deformation families for \tilde{X} . The key point for X of Tom type is in the nodes lying on the divisor $D \subset Z$.

Lemma 4.1. *If a general codimension 3 Fano 3-fold Z in Tom format is invariant under the $\mathbb{Z}/2\mathbb{Z}$ action γ , then the nodes on the divisor $D \subset Z$ are not fixed by γ . Moreover, they are pairwise-identified in the quotient \tilde{Z} , and $\#\{\text{nodes on } Z\} = 2 \cdot \#\{\text{nodes on } \tilde{Z}\}$.*

Proof. From [BKR12a, Theorem 3.2], the nodes of Z only lie on the divisor D . From the proof of Proposition 3.6 we deduce that the nodes are not fixed by γ .

Call $\tilde{D} := D/\gamma$. The equations of the nodes on $\tilde{D} \subset \tilde{Z}$ are given by the 3×3 minors of the Jacobian matrix $J|_{\tilde{Z}}$ restricted to \tilde{D} , i.e. $\wedge^3 \tilde{J}|_{\tilde{D}} = \underline{0}$. Such equations obviously depend on ζ . Lifting these equations to Z , ζ is replaced by x_1^2 . Therefore, the nodes on \tilde{Z} are double the number of nodes on Z .

Thus, the nodes on $D \subset Z$ are pairwise identified by γ in the quotient and the number of nodes on Z is twice the number of nodes on \tilde{Z} . \square

There are either one or two Tom deformation families for each X in index 1 [BKR12a, BKR12b]. For each of the 32 index 1 Fano 3-folds X in the GRDB that are candidates to be double covers of just as many \tilde{X} in index 2, there is exactly one Tom deformation family that realises the double cover. The Tom families coming from codimension 3 Tom formats having odd number of nodes are automatically excluded by Lemma 4.1.

Conversely, it is also possible to prove the following lemma.

Lemma 4.2. *Let Z be a codimension 3 terminal Fano 3-fold of index 1 such that the number of nodes on $D \subset Z$ is even. If a Type I unprojection X of Z has basket of singularities formed only by cyclic quotient singularities with odd order, then it is possible to realise Z as invariant under γ .*

Proof. As shown in the proof of Lemma 4.1, the group action γ identifies the nodes on $D \subset Z$ pairwise. Hence, the number of nodes being even is a necessary condition for the Z to be invariant under γ .

The hypotheses of X having only odd-order cyclic quotient singularities is crucial to produce a terminal codimension 4 index 2 Fano \tilde{X} . Indeed, quotient singularities with even order would make \tilde{X} not terminal (cf proof of Lemma 3.4 and [Suz04, Lemma 1.2 (3)]).

There are no other conditions that Z needs to satisfy in order to be invariant under the action of γ . A detailed explanation on how to exhibit such a realisation of Z explicitly is contained in Section 8. \square

It is important to observe that the codimension 4 index 1 Fano 3-folds that are a double cover of the respective codimension 4 index 2 Fano 3-folds are exactly those having quotient singularities of odd order and such that their corresponding Z in codimension 3 has even number of nodes. Recall from the proof of Lemma 3.4 that $\text{Fix}(\gamma) = \{x_1 = 0\} \cup \mathbb{P}_{\text{even}}$, and that $X \cap \{x_1 = 0\}$ contains all the cyclic quotient singularities of X . Thus, the basket of X is fixed by γ . Therefore, the following situations do not occur: a singularity of even order

$2n$ on X descends to a singularity of order $4n$ on \tilde{X} ; the singularities of even order on X are pairwise identified on \tilde{X} . This therefore proves the following theorem.

Theorem 4.3. *A general codimension 3 terminal Fano 3-fold Z of index 1 in Tom format can be realised as invariant under γ if and only if the number of nodes on $D \subset Z$ is even and the basket of its Type I unprojection X exclusively contains cyclic quotient singularities of odd order.*

For a general treatise of this argument, see [Suz04, Lemma 1.2 (3)].

As summarised in Table 1, each of the 32 codimension 4 index 1 Fano 3-folds that are candidates to be double covers, have one and only one Tom deformation family as described in Theorem 4.3. Except for the Hilbert series with GRDB ID #24078, the only Tom deformation family that admits the quotient by γ is the first Tom family as in [BKR12b]. Instead, the Fano #24078 only admits its second Tom family Tom_5 ; the deformation family Tom_5 of #24078 has Picard rank 2 by [BKQ18].

5. JERRY FAMILIES

In this section we examine codimension 4 terminal Fano 3-folds X of Jerry type, and we determine which Jerry families give rise to a deformation family for \tilde{X} in index 2. What has been discussed for Tom families still holds here, but some care is needed with regards to some features of the Jerry formats.

Suppose that Z is a terminal codimension 3 Fano 3-fold in Jerry format having at least one Type I centre, and call P the weight of the pivot entry. The criterion to determine which Jerry deformation families of X descend to \tilde{X} depends on whether the following condition is satisfied or not.

Condition 5.1. There exists a generator y_k of the ideal I_D whose weight is equal to P .

We therefore have the following theorem.

Theorem 5.1. *A general codimension 3 terminal Fano 3-fold Z of index 1 in Jerry format (except for #24077 Jerry₁₂) can be realised as invariant under γ if and only if the following statements are simultaneously satisfied: the number of nodes on $D \subset Z$ is even; the basket of its Type I unprojection X exclusively contains cyclic quotient singularities of odd order; Condition 5.1 is fulfilled; the weight P of the pivot entry is even.*

Proof. The part of the proof concerning the nodes and the order of the singularities is analogous to the one for Theorem 4.3. This is the first condition that we have to verify when trying to exclude a Jerry format. If the number of nodes on D is even, we proceed with checking the properties of the pivot entry.

The check on the nodes excludes the majority of the failing Jerry formats. Only a few are left, which present different reasons for failure. We expand this in Subsection 5.1 and we show that those formats fall in the phenomenology of [BKR12a, Section 5]. As a consequence, they have non-reduced singularities on D .

The presence of a pivot entry with even weight is crucial for having reduced singularities on D . If P is odd, of the three entries of M that are not in I_D two have odd weight and one has even weight for homogeneity of the maximal pfaffians. Given that we want x_1 to appear only with even powers, we have a further constraint that is often not compatible with the Jerry format, so this forces a 0 in the entry not in I_D for which it is not possible to

find a polynomial not in I_D on the right degree (see Subsection 5.1 below). Furthermore, row/column operations on M show that other entries can be made 0: together with the analysis in [BKR12a, Section 5], this concludes the proof. \square

Remark 5.2. The Fano 3-fold with GRDB ID #24078 of Jerry type is the only family that does not follow Theorem 5.1. Indeed, its codimension 3 source #24077 in Jerry₁₂ format has even number of nodes and satisfies Condition 5.1, but the weight of its pivot is 1, and its basket is $\mathcal{B}_Z := \{\frac{1}{2}(1, 1, 1)\}$. However, employing a code analogous to the ones in Section 8, it is still possible to produce explicit equations for its corresponding index 2 Fano #40933. The ambient space of #24077 is $\mathbb{P}^6(1^6, 2)_{x_i, y}$, and suppose that γ changes sign to the coordinate x_1 , so $\xi := x_1^2$. This Fano 3-fold is peculiar because it is the only index 1 double cover whose ambient space has enough coordinates of weight 1 to fill the entries of M in Jerry₁₂ format appropriately. Suppose $I_D := \langle x_3, x_4, x_5, x_6 \rangle$. The grading of M is the following, and, for $q_1, q_2, q_3 \in I_D$ homogeneous polynomials of degree 1, 2, 2 respectively, it can be filled as

$$\begin{pmatrix} 1 & 1 & 1 & 2 \\ & 1 & 1 & 2 \\ & & 1 & 2 \\ & & & 2 \end{pmatrix} \rightsquigarrow \begin{pmatrix} x_3 & x_4 & x_5 & q_1 \\ & x_6 & q_2 & q_3 \\ & & x_2 & y \\ & & & \xi \end{pmatrix} = M.$$

This shows that only q_2 could be made 0 after row/column operations on M , and that there is no further row/column elimination possible. Thus, #24077 does not fall into the failure description of [BKR12a, Section 5].

5.1. Failure of Jerry formats. We say that a certain format for a codimension 3 Fano 3-fold Z *fails* if the deformation family of Z associated to that format does not enjoy the following properties: terminal, nodal on D with reduced singularities. The reasons for formats' failures is extensively described in [BKR12a, Section 5]. In this paper, we have additional reasons for failure: not having even number of nodes on D , not satisfying Condition 5.1 (in the Jerry case), not having the pivot entry with even weight. The feature that excludes all the failing Tom formats and most of the Jerry formats is the one about the number of nodes. Moreover, except for #24078, all Jerry formats for which Z has even number of nodes on D also satisfy Condition 5.1. Here we want to discuss the Jerry formats for which the conditions 5.1 and the even nodes are satisfied, but do not have the pivot entry with even weight.

There are exactly seven formats that fall in this case. We run an explicit check on all of them using the reasons for failure in [BKR12a, Section 5]. These failing formats all give rise to non-reduced singularities on D . We specify the Type I centre in case of ambiguity, and we refer to [BKR12b] for the grading of the matrix M .

#1401, $\frac{1}{7}(1, 2, 5)$, Jerry₂₄. The ambient space of Z in Jerry₂₄ format is $\mathbb{P}^6(1, 2, 3, 3, 4, 5, 5)$ with homogeneous coordinates $x_1, x_2, y_1, y_2, y_3, y_4, x_3$ respectively, and the ideal I_D is generated by y_1, y_2, y_3, y_4 . The entry a_{13} has degree 3, and we want the coordinate x_1 to appear only with even powers in order to perform the double cover construction in Section 3. Thus, there is no way to fill entry a_{13} with an homogeneous polynomial of degree 3 not in I_D , so a_{13} must be 0. On the other hand, the entries a_{14}, a_{23} must both be equal to y_3 , as there is no other polynomial of degree 4 in I_D in the available coordinates. Thus, the format Jerry₂₄ fails for the reason in [BKR12a, Section 5.2 (4)].

#1401, Jerry₄₅. For Z in Jerry₄₅ the ambient space is $\mathbb{P}^6(1, 2, 3, 3, 4, 5, 7)$ with homogeneous coordinates $x_1, x_2, x_3, y_1, y_2, y_3, y_4$ respectively. The entries a_{12}, a_{13} of M are both equal to x_3 because there is no other polynomial of degree 3 not in the ideal I_D that is only in x_1, x_2, x_3 . In addition, the entry a_{34} can be made 0 via row/column operations on M . Thus, Jerry₄₅ fails because of [BKR12a, Section 5.2 (4)].

#1405, Jerry₂₄. Since there are only two generators of I_D in degree 5, either a_{25} or a_{34} can be made 0 via row/column operations on M . Also $a_{14} = a_{23}$ because there is only one choice for a degree 4 polynomial in I_D . Hence, Jerry₂₄ fails due to [BKR12a, Section 5.2 (4)].

#4987, Jerry₁₂. The entry a_{23} is 0 after row/column operations, and $a_{34} = 0$ because there is no degree 7 polynomial not in I_D . So, Jerry₁₂ fails due to [BKR12a, Section 5.2 (3)].

#11104, Jerry₁₂. The entry a_{23} is 0 after row/column operations, and $a_{34} = 0$ because there is no degree 3 polynomial not in I_D . So, Jerry₁₂ fails due to [BKR12a, Section 5.2 (3)].

#11123, Jerry₁₂•₁₅. The entry a_{15} is already 0 in this format. Moreover, $a_{34} = a_{35}$ because there is only one choice for a degree 3 polynomial in I_D . Thus, Jerry₁₂•₁₅ fails due to [BKR12a, Section 5.2 (4)].

#11123, Jerry₂₄•₂₅. The entry a_{25} is already 0. In addition, a_{15} can be made 0 by row/column operations. Thus, Jerry₂₄•₂₅ fails due to [BKR12a, Section 5.2 (3)].

6. PROOF OF MAIN THEOREM

The proof of Theorem 1.1 relies on the possibility to realise X in codimension 4 and index 1 as invariant under the action γ . The construction of X starts from Z in codimension 3. We gave necessary and sufficient conditions to realise Z as invariant under γ in Theorems 4.3 and 5.1 for both types of deformation families of Z . It is possible to verify that the invariance under γ is maintained to X , that is, performing the unprojection does not affect the γ -invariance.

The unprojection equations for X are calculated from the matrices of coefficients of the ideal generators y_1, \dots, y_4 in the five pfaffian equations of Z . We refer to [Pap04, Section 5] and [Cam20, Appendix] for notation and definitions of these objects. The process continues with taking determinants of certain submatrices of the matrices of coefficients. These determinants enjoy a divisibility property ([Pap04, Lemma 5.3]): for Z in Tom format, the right hand sides of the unprojection equations are divisible by one of the polynomials not in the ideal I_D . The monomials containing ζ in the equations of Z are always considered as coefficients of the generators of I_D , and none of the operations involved in constructing the unprojection equations (i.e. taking coefficients and determinants, divide by a factorizing invariant polynomial) compromises the γ -invariance. A similar argument can be made in the Jerry case.

The discussion about the two double cover Fano 3-folds arising from Type II₂ unprojections is contained in the next Section 7.

7. TYPE II₂ UNPROJECTIONS

The two Hilbert series #39569 and #39607 have candidate double covers, #512 and #872 respectively, that do not have any Type I centre. Indeed, #512 and #872 are obtained as Type II₂ unprojections of the corresponding Fano hypersurfaces Y . The Tom and Jerry construction

is not appropriate in this case, but different deformation families can still arise. For extensive treatise of Type II unprojections, see [Pap08, Pap07, Pap06, Tay20]. In particular, in this Section we follow closely [Pap06] for the general strategy, and [Tay20] for part of the explicit approach. Here we show that Theorem 1.1 still holds for #39569 and #39607.

7.1. **#39607.** The index 1 codimension 4 Fano 3-fold #872 $X \subset \mathbb{P}^7(1, 3^2, 4, 5^2, 6, 7)$ is candidate to be a double cover of #39607 of index 2. Its basket is $\mathcal{B}_X := \{5 \times \frac{1}{3}(1, 1, 2), \frac{1}{5}(1, 1, 4)\}$, and $\frac{1}{5}(1, 1, 4)$ is its Type II₂ centre. It comes from the Type II₂ unprojection of the hypersurface #866 $Y_{15} \subset \mathbb{P}^4(1, 3^2, 4, 5)$ having basket $\mathcal{B}_Y := \{5 \times \frac{1}{3}(1, 1, 2), \frac{1}{4}(1, 1, 3)\}$. Following [Tay20], X is obtained by unprojecting the divisor D defined as the image of $\mathbb{P}^2(1, 1, 4)_{a,b,c}$ inside Y via the embedding

$$\begin{aligned} \phi: \mathbb{P}^2(1, 1, 4) &\longrightarrow \mathbb{P}^4(1, 3, 3, 4, 5) \\ (a, b, c) &\longmapsto (a, b^3, b^3, c, bc) \end{aligned}$$

In this way, $D := \text{Im}(\phi)$ is a divisor inside the general hypersurface $Y_{15} \subset \mathbb{P}^4(1, 3^2, 4, 5)$ with homogeneous coordinates x, u, z, y, v . In addition, $\text{Im}(\phi)$ can be written as the 3×3 minors of the following 3×6 matrix ([Tay20, Example 5.2.2])

$$N = \begin{pmatrix} u & v & 0 & 0 & -x^2z & -yz \\ x^2 & -y & u & v & 0 & 0 \\ 0 & 0 & x^2 & -y & u & v \end{pmatrix}.$$

The equation defining Y_{15} is an homogeneous polynomial of degree 15 of the form

$$(7.1) \quad F = \sum_{i=1}^{20} A_i N_i$$

where N_i is the i -th 3×3 minor of N , and A_i is a general homogeneous polynomial of degree $15 - \deg(N_i)$. Note that here both N_{15} and N_{20} have already degree 15, so A_{15} and A_{20} are both non-zero constants. Since x appears already as a square in N , we only need that x only appears with even powers in the polynomials A_i ; in this way, Y_{15} is invariant under γ .

For Type II₂ unprojections, there will be three new unprojection variables: indeed, the codimension increases by 3. We call them s_1, s_2, s_3 , and they have weights 5, 6, 7 respectively. Finding the equations for X consists in finding the linear and quadratic relations between the unprojection variables s_1, s_2, s_3 (cf [Pap06, Subsection 2.2 and Lemma 2.5]).

Consider the polynomial ring R generated by the coordinates of $\mathbb{P}^4(1, 3^2, 4, 5)$, the monomials $p_1 := -a^2$, $p_2 := c$, and the polynomials A_i (cf [Pap06, Definition of \mathcal{O}_{amb}]). The equations of X are retrieved by looking at the resolution of $R/\langle F = 0 \rangle$, and they consist in sums and products of the coordinates of $\mathbb{P}^4(1, 3^2, 4, 5)$, the monomials $p_1 := -a^2$, $p_2 := c$, and the polynomials A_i . Thus, x still retains its property of appearing only with even powers, and X is defined by 9 equations invariant under γ . So, we can quotient X by the action γ , and obtain \tilde{X} analogously to Section 3. This constructs #39607.

Note that it is possible to check with a Macaulay2 routine that the ring extension $R[s_1, s_2, s_3]$ quotient by the nine equations of X is Gorenstein. We are not aware of any work in the literature regarding the question of whether there are multiple deformation families induced by Type II₂ unprojections, and if there are, how many.

7.2. **#39569.** We proceed analogously for #39569. Its double cover candidate is #512 $X \subset \mathbb{P}^7(1, 3, 5, 6, 7^2, 8, 9)$, whose basket is $\mathcal{B}_X := \{3 \times \frac{1}{3}(1, 1, 2), \frac{1}{5}(1, 2, 3), \frac{1}{7}(1, 1, 6)\}$, and $\frac{1}{7}(1, 1, 6)$ is one of its Type II₂ centres. It is induced by the Type II₂ unprojection of the hypersurface #508 $Y_{21} \subset \mathbb{P}^4(1, 3, 5, 6, 7)$ with basket $\mathcal{B}_Y := \{3 \times \frac{1}{3}(1, 1, 2), \frac{1}{5}(1, 2, 3), \frac{1}{6}(1, 1, 5)\}$. The divisor D is defined as the image of $\mathbb{P}^2(1, 1, 6)_{a,b,c}$ inside Y via the embedding

$$\begin{aligned} \phi: \mathbb{P}^2(1, 1, 6) &\longrightarrow \mathbb{P}^4(1, 3, 5, 6, 7) \\ (a, b, c) &\longmapsto (a, b^3, b^5, c, bc) \end{aligned}$$

The divisor $D := \text{Im}(\phi)$ is given by the vanishing of the 3×3 minors of the 3×6 matrix

$$N = \begin{pmatrix} u & v & -z^2 & 0 & 0 & -yz \\ 0 & -y & u & v & -z^2 & 0 \\ -z & 0 & 0 & -y & u & v \end{pmatrix}.$$

It sits inside the general hypersurface $Y_{21} \subset \mathbb{P}^4(1, 3, 5, 6, 7)$ with homogeneous coordinates x, z, u, y, v respectively. The equation of Y_{21} is again given by (7.1), and we can impose that the general polynomials A_i contain the variable x only with even powers. As before, A_{15} and A_{20} are constants. The three new unprojection variables are s_1, s_2, s_3 with weights 7, 8, 9 respectively. The unprojection of Y and the quotient of X by γ constructs \tilde{X} #39569.

8. EXAMPLES

Here we give two examples of our construction, one of a Tom family and one of a Jerry family. We explicitly construct the deformation families relative to the Hilbert series with GRDB ID #39660. This is $\tilde{X} \subset \mathbb{P}^7(2, 2, 3, 5, 5, 7, 12, 17)$, whose basket of singularities is $\mathcal{B}_{\tilde{X}} := \{\frac{1}{17}(2, 5, 12)\}$. The codimension 4 Fano 3-fold $X \subset \mathbb{P}^7(1, 2, 3, 5, 5, 7, 12, 17)$ in index 1 with Hilbert series #1158 is the candidate to be the double cover for #39660. The coordinates of the ambient space of X are $x, y, z, u_1, u_2, v, w, s$, and its basket of singularities is $\mathcal{B}_X := \{\frac{1}{17}(1, 5, 12)\}$. In this case, the grading of M is

$$\text{wts } M = (m_{ij}) = \begin{pmatrix} 2 & 3 & 5 & 7 \\ & 5 & 7 & 9 \\ & & 8 & 10 \\ & & & 12 \end{pmatrix}.$$

The family of the codimension 3 Fano 3-fold Z with GRDB ID #1157 is composed of 3-folds sitting inside $\mathbb{P}^6(1, 2, 3, 5, 5, 7, 12)$ and whose equations are the five maximal pffians of M with the above grading. The divisor D is $\mathbb{P}^2(2, 5, 12)$. The Tom and Jerry formats of M that admit an embedding $D \subset Z$ are Tom_5 and Jerry_{12} each with 4 and 6 nodes on D respectively. In addition, the pivot entry of the Jerry_{12} format is a_{12} , which has weight 2. Therefore, by Lemmas 4.1, 4.2 we have two possible deformation families for \tilde{X} , one coming from the Tom_5 format, the other from the Jerry_{12} format. It remains to exhibit the equations for these two families. These calculations can be checked using the `tj` package for Magma that can be found on the Graded Ring Database website [BK⁺].

8.1. Finding equations for #39660 of Tom₅ type. Here we want to build \tilde{Z} produced from Z in Tom_5 format. We define the ambient space $\mathbb{P}^6(2, 2, 3, 5, 5, 7, 12)$ with coordinates as above, where x has been replaced by ζ of weight 2. The divisor $D \cong \mathbb{P}^2(2, 5, 12)$ is defined by the vanishing of the coordinates y, z, u_2, v . Here the matrix M in Tom_5 format is filled following

[BKR12a, Section 6.2]. As a rule of thumb, we place homogeneous coordinates in the entries with matching degrees where possible according to the format. The rest of the entries are occupied by general polynomials in the given degrees, still maintaining the format. The matrix can be tidied up by row/column operations. For instance, we can fill the entries of M as follows

$$M = (a_{ij}) = \begin{pmatrix} y & z & u_2 & -v \\ -u_2 & v & -z^3 + \zeta^2 u_1 & \\ & \zeta^3 y + y^4 & -\zeta^5 + u_1^2 & \\ & & w & \end{pmatrix}.$$

The equations of the 3-fold \tilde{Z} are the maximal pfaffians of M , and it is possible to check with a Magma routine that the number of nodes on D is 2, in accordance to Lemma 4.1. The next step is to perform the Type I unprojection from $D \subset \tilde{Z}$ as in [PR04]. The nine equations of \tilde{X} are

$$\begin{aligned} \zeta^3 y^2 + y^5 - u_2^2 - zv &= 0 \\ \zeta^5 y - z^4 + \zeta^2 z u_1 - y u_1^2 - u_2 v &= 0 \\ z^3 u_2 - \zeta^2 u_1 u_2 - v^2 + y w &= 0 \\ \zeta^5 u_2 - \zeta^3 y v - y^4 v - u_1^2 u_2 + z w &= 0 \\ \zeta^3 y z^3 + y^4 z^3 - \zeta^2 y^4 u_1 - \zeta^5 v - z^4 u_1 + \zeta^2 z u_1^2 - y u_1^3 + u_1^2 v - u_1 u_2 v + u_2 w &= 0 \\ \zeta^5 z^3 - \zeta^7 u_1 - z^3 u_1^2 + \zeta^2 u_1^3 - v w + y s &= 0 \\ -\zeta^{10} + 2\zeta^5 u_1^2 + \zeta^3 v^2 + y^3 v^2 - u_1^4 - z s &= 0 \\ \zeta^3 z^3 v + y^3 z^3 v - \zeta^5 u_1 v - \zeta^2 y^3 u_1 v - \zeta^5 w + u_1^2 w + u_2 s &= 0 \\ -\zeta^8 y z^2 - \zeta^5 y^4 z^2 + \zeta^5 z^3 u_1 + \zeta^2 y^3 z^3 u_1 - \zeta^7 u_1^2 - \zeta^4 y^3 u_1^2 \\ + \zeta^3 y z^2 u_1^2 + y^4 z^2 u_1^2 + \zeta^3 z^2 u_2 v + y^3 z^2 u_2 v + w^2 - v s &= 0. \end{aligned}$$

8.2. Finding equations for #39660 of Jerry₁₂ type. In a similar fashion to the Tom case above, here the matrix M in Jerry₁₂ is defined as

$$M = (a_{ij}) = \begin{pmatrix} y & z & u_2 & v \\ -u_2 & v & \zeta^3 z + z^3 & \\ & \zeta^4 + y^4 & \zeta y^4 - u_1^2 & \\ & & w & \end{pmatrix}.$$

The codimension 3 Fano 3-fold Z #1157 satisfies the conditions required in Theorem 5.1. In particular, \tilde{Z} has 3 nodes on D . The equations of \tilde{X} of Jerry₁₂ type with GRDB ID #39660 are

$$\begin{aligned} \zeta^4 y + y^5 - u_2^2 - zv &= 0 \\ \zeta y^5 - \zeta^3 z^2 - z^4 - y u_1^2 - u_2 v &= 0 \\ \zeta^3 z u_2 + z^3 u_2 - v^2 - y w &= 0 \\ \zeta y^4 u_2 - \zeta^4 v - y^4 v - u_1^2 u_2 - z w &= 0 \\ \zeta^7 z + \zeta^3 y^4 z + \zeta^4 z^3 + y^4 z^3 - \zeta y^4 v + u_1^2 v - u_2 w &= 0 \\ \zeta^7 u_2 + \zeta^4 z^2 u_2 - \zeta^3 z u_1^2 - z^3 u_1^2 + v w + y s &= 0 \\ \zeta y^4 u_1^2 - \zeta y^3 u_2 v + y^3 v^2 - \zeta^4 w - u_1^4 - z s &= 0 \end{aligned}$$

$$\begin{aligned} \zeta^{11} + \zeta^7 y^4 + \zeta^8 z^2 + \zeta^4 y^4 z^2 + \zeta^3 y^3 z v + y^3 z^3 v - \zeta y^3 v^2 - u_1^2 w + u_2 s &= 0 \\ \zeta^7 u_1^2 + \zeta^3 y^4 u_1^2 - \zeta^4 y^3 u_2^2 + \zeta^4 z^2 u_1^2 + y^4 z^2 u_1^2 - \zeta y^3 z^2 u_2^2 + \zeta^3 y^3 u_2 v + y^3 z^2 u_2 v + w^2 - v s &= 0. \end{aligned}$$

REFERENCES

- [ABR02] S. Altınok, G. Brown, and M. Reid. Fano 3-folds, K3 surfaces and graded rings. In *Topology and geometry: commemorating SISTAG*, volume 314 of *Contemp. Math.*, pages 25–53. Amer. Math. Soc., Providence, RI, 2002.
- [BK⁺] G. Brown, A. M. Kasprzyk, et al. Graded Ring Database. *Online. Access via <http://www.grdb.co.uk>*.
- [BKQ18] G. Brown, A. M. Kasprzyk, and M. I. Qureshi. Fano 3-folds in $\mathbb{P}^2 \times \mathbb{P}^2$ format, Tom and Jerry. *Eur. J. Math.*, 4(1):51–72, 2018.
- [BKR12a] G. Brown, M. Kerber, and M. Reid. Fano 3-folds in codimension 4, Tom and Jerry. Part I. *Compos. Math.*, 148(4):1171–1194, 2012.
- [BKR12b] G. Brown, M. Kerber, and M. Reid. Tom and Jerry table, part of "Fano 3-folds in codimension 4, Tom and Terry. Part I". *Compositio Mathematica*, 148(4):1171–1194, 2012.
- [BS07] G. Brown and K. Suzuki. Fano 3-folds with divisible anticanonical class. *Manuscripta Math.*, 123(1):37–51, 2007.
- [Cam20] L. Campo. Sarkisov links for index 1 fano 3-folds in codimension 4. *arXiv preprint arXiv:2011.12209*, 2020.
- [CD20] S. Coughlan and T. Ducat. Constructing Fano 3-folds from cluster varieties of rank 2. *Compos. Math.*, 156(9):1873–1914, 2020.
- [Duc18] T. Ducat. Constructing Q-Fano 3-folds à la Prokhorov & Reid. *Bull. Lond. Math. Soc.*, 50(3):420–434, 2018.
- [Har77] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [Isk77] V. A. Iskovskih. Fano threefolds. I. *Izv. Akad. Nauk SSSR Ser. Mat.*, 41(3):516–562, 717, 1977.
- [Isk78] V. A. Iskovskih. Fano threefolds. II. *Izv. Akad. Nauk SSSR Ser. Mat.*, 42(3):506–549, 1978.
- [KM83] A. R. Kustin and M. Miller. Constructing big Gorenstein ideals from small ones. *J. Algebra*, 85(2):303–322, 1983.
- [Pap04] S. A. Papadakis. Kustin-Miller unprojection with complexes. *J. Algebraic Geom.*, 13(2):249–268, 2004.
- [Pap06] S. A. Papadakis. Type II unprojection. *J. Algebraic Geom.*, 15(3):399–414, 2006.
- [Pap07] S. A. Papadakis. Towards a general theory of unprojection. *J. Math. Kyoto Univ.*, 47(3):579–598, 2007.
- [Pap08] S. A. Papadakis. The equations of type II_1 unprojection. *J. Pure Appl. Algebra*, 212(10):2194–2208, 2008.
- [PR04] S. A. Papadakis and M. Reid. Kustin-Miller unprojection without complexes. *J. Algebraic Geom.*, 13(3):563–577, 2004.
- [PR16] Y. Prokhorov and M. Reid. On Q-Fano 3-folds of Fano index 2. In *Minimal models and extremal rays (Kyoto, 2011)*, volume 70 of *Adv. Stud. Pure Math.*, pages 397–420. Math. Soc. Japan, [Tokyo], 2016.
- [Suz04] K. Suzuki. On Fano indices of Q-Fano 3-folds. *Manuscripta Math.*, 114(2):229–246, 2004.
- [Tay20] R. Taylor. *Type II unprojections, Fano threefolds and codimension four constructions*. PhD thesis, University of Warwick, 2020.

SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF NOTTINGHAM, UNIVERSITY PARK, NOTTINGHAM, NG7 2RD, UNITED KINGDOM

Email address: livia.campo@nottingham.ac.uk