

A peek outside our Universe

Enrique Gaztañaga & Pablo Fosalba

Institute of Space Sciences (ICE, CSIC), 08193 Barcelona, Spain

Institut d'Estudis Espacials de Catalunya (IEEC), 08034 Barcelona, Spain

gaztanaga@gmail.com (corresponding) & fosalba@gmail.com

Essay written for the Gravity Research Foundation 2021 Awards for Essays on Gravitation. April 2, 2021

According to General Relativity (GR) a universe with a cosmological constant, Λ , like ours, is trapped inside an event horizon $r < \sqrt{3/\Lambda}$. What is outside? We show, using Israel (1967) junction conditions, that there could be a different universe outside. Our Universe looks like a Black Hole for an outside observer. Outgoing radial null geodesics can not escape our universe, but incoming photons can enter and leave an imprint on our CMB sky. We present a picture of such a fossil record from the analysis of CMB maps that agrees with the Black Hole universe predictions but challenge our understanding of the origin of the primordial universe.

I. INTRODUCTION

An Event Horizon (EH), r_* , or trapped surface, for a given observer can be defined as the distance beyond which this observer will never see: $r < r_*$. The most famous EH is of course that of a Black Hole (BH) of mass M . Dimensional analysis tell us that a relativistic (c) gravitational (G) system of mass M has an associated EH: $r_* \simeq GM/c^2$. The same dimensional analysis indicates that a cosmological constant, Λ , is associated to a relativistic EH: $r_* \simeq 1/\sqrt{\Lambda c^2}$ which, in principle, is independent of G . We will show first how these EHs appear as solutions to Einstein's field equations and next how the different EHs are related in our Universe expansion.

The most general form for a metric with spherical symmetry in proper coordinates $x^\mu = (t, r, \delta, \theta)$ with $c = 1$ is [1]:

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -A(t, r) dt^2 + B(t, r) dr^2 + r^2 d\Omega^2 \quad (1)$$

where we have introduced: $d\Omega^2 = \cos^2 \delta d\theta^2 + d\delta^2$. Einstein's field equations for empty space $\rho = p = \Lambda = 0$ results in the Schwarzschild (SCHW) metric:

$$ds^2 = -\left[1 - \frac{2GM}{r}\right] dt^2 + \frac{dr^2}{1 - \frac{2GM}{r}} + r^2 d\Omega^2 \quad (2)$$

where M can be interpreted as a singular point mass at $r = 0$. The EH at $r_* = 2GM$ prevent us from seeing such naked singularity. Outgoing null radial geodesics cannot leave the interior of r_* , while incoming ones can cross inside r_* .

The solution for the above metric Eq.1 with $\rho = p = M = 0$, but $\Lambda \neq 0$ is the deSitter (deS) metric:

$$ds^2 = -\left[1 - r^2 H_\Lambda^2\right] dt^2 + \frac{dr^2}{1 - r^2 H_\Lambda^2} + r^2 d\Omega^2 \quad (3)$$

which is also static and has an EH, or trapped surface, at $r_* = 1/H_\Lambda$ where $H_\Lambda^2 \equiv 8\pi G \rho_\Lambda/3$ and $\rho_\Lambda = \Lambda/(8\pi G) + \rho_{vac}$. We include a constant $\rho_{vac} = V = V(\phi) = -p_{vac}$ to account for vacuum energy (or the potential of a trapped scalar field) which is physically degenerate with Λ . The inside of $r_* = 1/H_\Lambda$ is causally disconnected and corresponds to an EH: radial null events ($ds^2 = 0$) connecting $(0, r_0)$ with (t, r) take $t = \infty$ to reach $r = r_*$ from any point inside.

A. The FLRW metric & Λ

The Friedmann-Lemaitre-Robertson-Walker (FLRW) flat ($k = 0$) metric in spherical comoving coordinates $(\tau, \chi, \delta, \theta)$, corresponds to an homogeneous and isotropic space-time:

$$ds^2 = -d\tau^2 + a(\tau)^2 [d\chi^2 + \chi^2 d\Omega^2] \quad (4)$$

The scale factor, $a(\tau)$, describes the expansion/contraction as a function of time. For a comoving observer and a perfect fluid, the field equations reduce to:

$$3H^2 \equiv \left(\frac{\dot{a}}{a}\right)^2 = 8\pi G(\rho_m a^{-3} + \rho_R a^{-4} + \rho_\Lambda) \quad (5)$$

where ρ_m and ρ_R are the matter and radiation density today ($a = 1$) and $\rho_\Lambda = -p_\Lambda$ is the effective cosmological constant density introduced in deS metric. Given ρ and p at some time, we can find $a = a(\tau)$ and determine the metric in Eq.4. Observations show that the expansion rate today is dominated by ρ_Λ . This indicates that the FLRW metric describes the interior of a trapped surface of size $r_* = 1/H_\Lambda$, like deS metric. In fact, both metrics are equivalent in that regime [2].

Interpreting Λ as a boundary term to the GR equations, and imposing the so-called zero action principle, $r_* = 1/H_\Lambda$ can be given in terms of ordinary contributions to the energy density, $\rho_\Lambda = \rho_m/2 + \rho_R > [3]$.

B. SCH-FLRW perturbation

The Schwarzschild (SCHW) metric is commonly used to describe the outside of BHs or stars and it should be understood as a perturbation inside a larger background. An example of this is the SCHW-deS metric, where ds^2 equals:

$$-\left[1 - \frac{2GM}{r} - r^2 H_\Lambda^2\right] dt^2 + \frac{dr^2}{1 - \frac{2GM}{r} - r^2 H_\Lambda^2} + r^2 d\Omega^2 \quad (6)$$

which corresponds to a solution with both non-zero Λ and M . More generally, the FLRW metric is the background to the SCHW solution [4]. For large r we recover the deS metric, as in Eq.6 above, which is equivalent to the FLRW metric dominated by Λ [2]. For $r < 1/H_\Lambda$ we recover SCHW.

II. OUTSIDE OUR FLRW UNIVERSE

In proper coordinates $r = a\chi$ the FLRW metric with $H = H(\tau)$ is also trapped inside the same EH as the deS one: $r_* = 1/H_\Lambda$ because $H(\tau) > H_\Lambda$. We can see this by considering outgoing radial null geodesics in the FLRW metric (Eq.4):

$$r_{out} = a(\tau) \int_\tau^\infty \frac{d\tau'}{a(\tau')} = a \int_a^\infty \frac{d \ln a'}{a' H(a')} < \frac{1}{H_\Lambda} = r_* \quad (7)$$

which shows that signals can not escape from the inside to the outside of the EH. But incoming radial null geodesics $a(\tau) \int_0^\tau \frac{d\tau'}{a(\tau')}$ could be larger than r_* if we look back in time for long enough. This shows that observers living in the interior are trapped inside the EH, but they can, in principle, observe what happened outside.

What is outside $r_* = 1/H_\Lambda$ in the FLRW metric? The FLRW comoving coordinates (τ, χ) can be matched to the SCHW proper coordinates (t, r) . The joint metric, is what [3, 5] call a BH.u solution. The particular case where the inside is deS (and the outside SCHW) is called BH.fv (where fv stand for false vacuum). The joint BH.u metric is also a solution to Einstein's field equations. To prove this we just need to show that the junction follows the Israel matching conditions [6]. The two metrics can be matched on a timelike hypersurface Σ of constant χ :

$$ds_\Sigma^2 = h_{ab} dy^a dy^b = -d\tau^2 + a^2(\tau) \chi_\S^2 d\Omega^2 \quad (8)$$

and the extrinsic curvature K at Σ is the same in both sides. The matching conditions $h_- = h_+$ and $K_+ = K_-$ reduced to [5]

$$r = R(\tau) = a(\tau) \chi_\S ; \quad \dot{R}^2 = R^2 H^2 = \frac{R_*}{R} \quad (9)$$

where $R_* \equiv 2GM$. Staring from small a , as we increase τ , both R and $\dot{R} = HR$ grow until we reach $HR = c = 1$ which corresponds to the event horizon $R_* = 2GM = 1/H_\Lambda$. It takes $t = \infty$ in the SCHW time to asymptotically reach R_* . This proves that the joint metric BH.u is also a solution to the Einstein's field equations with no surface terms in the junction [6]. This is equivalent to stating that the Λ term corresponds to a trapped surface $R_* = 1/H_\Lambda$ in the FLRW metric which matches the EH of a BH. Generalization to $k \neq 1$ is straightforward (see §12.5.1 in [1] or [7]).

Recall that the SCHW metric is a perturbation of a larger FLRW metric, i.e, a BH-like metric embedded in a background described by the FLRW metric. This means that outside $r_* = 1/H_\Lambda$ we just have another FLRW metric, like in Matryoshka doll. From outside, the inner FLRW metric looks like a BH. There could be many other BH.u inside and outside $r_* = 1/H_\Lambda$, so the structure could be better described by a fractal.

A. Causal structure

In the FLRW Universe, the Hubble Horizon r_H is defined as $r_H = c/H$. Scales larger than r_H cannot evolve because the time a perturbation takes to travel that distance is larger




Figure 1. Proper coordinate: $R = a(\tau)\chi$ in units of c/H_0 as a function of cosmic time a . The Hubble horizon c/H (blue continuous line), is compared to the observable universe $a\chi_0$ after inflation (dashed line) and the primordial causal boundary $\chi_\S = c/(a_i H_i)$ (dot-dashed red line). Larger scales (green shading) are causally disconnected, smaller scales (yellow shading) are dynamically frozen. After inflation c/H grows again. At $a \approx 1$ the Hubble horizon reaches our event horizon $R_* = c/H_\Lambda$. At CMB last scattering we can observe both frozen and causally disconnected perturbations.

than the expansion time. This means that $r > r_H$ scales are "frozen out" (structure can not evolve) and are causally disconnected from the rest. Thus, c/H represents a dynamical causal horizon that is evolving (blue line in Fig.1).

We can sketch the evolution of our universe in Fig.1. A primordial field ψ settles or fluctuates into a false (or slow rolling) vacuum which will create a BH.fv with a junction Σ in Eq.8 where the causal boundary is fixed in comoving coordinates and corresponds to the particle horizon during inflation $\chi_\S = c/(a_i H_i)$ or the Hubble horizon when inflation begins. The size $R = a(\tau)\chi_\S$ of this vacuum grows and asymptotically tends to $R_* = c/H$ following Eq.9 with $H = H_i$. The inside of this BH will be expanding exponentially $a = e^{\tau H_i}$ while the Hubble horizon is fixed $1/H_i$. When this inflation ends [8] vacuum energy excess converts into matter and radiation (reheating). This results in BH.u, where the infinitesimal Hubble horizon starts to grow following the standard Big Bang evolution. The observable universe (or particle horizon) after inflation, χ_0 , is:

$$\chi_0 = \chi_0(a) = \int_{a_e}^a \frac{d \ln a'}{a' H(a')} = \chi_0(1) - \bar{\chi}(a), \quad (10)$$

where a_e is the scale factor when inflation ends. For $\Omega_\Lambda \approx 0.7$, the particle horizon today is $\chi_0(1) \approx 3.26c/H_0$ and $\bar{\chi}(a) = \int_a^1 d \ln a' / (a' H)$ is the radial lookback time, which for a flat universe agrees with the comoving angular diameter distance, $d_A = \bar{\chi}$. The observable universe becomes larger than R_* when $a > 1$, as shown in Fig.1 (compare dotted and dashed lines). This shows that, observers like us, living in the interior of the BH universe, are trapped inside R_* but can nevertheless observe what happened outside. We can estimate χ_\S from $\rho_\Lambda = \langle \rho_m/2 + \rho_R \rangle$ (see §I.A), where the average is in the




Figure 2. Map of the best-fit values of the dark-energy density Ω_Λ across the celestial sphere estimated from partial sky (discs) measurements of the Planck CMB maps. The large grey circles delimit areas across the sky with significantly different values of Ω_Λ .

lightcone inside χ_\S . For $\Omega_\Lambda \approx 0.7$ [3] found: $\chi_\S \approx 3.34c/H_0$ which is close to χ_O today. Imagine that Ω_Λ is caused by some Dark Energy and has nothing to do with χ_\S . We still have that $\chi_\S \lesssim \chi_O$, because otherwise χ_\S would have crossed $RH = 1$ early on, resulting in smaller χ_O than measured (see Fig.1).

Thus, at the time of CMB last scattering (when $d_A \approx \chi_O$), χ_\S corresponds to an angle $\theta = \chi_\S/d_A \lesssim 1 \text{ rad} \approx 60 \text{ deg}$. So we can actually observe scales larger than χ_\S . Scales that are not causally connected! This could be related to the so-called CMB anomalies (i.e, apparent deviations with respect to simple predictions from Λ CDM, see [9] and references therein), or the parameter tensions in measurements from vastly different cosmic scales or times [10].

B. A peek outside

A recent analysis of the Planck temperature anisotropy data, [11], shows that the distribution of best-fit dark-energy density Ω_Λ exhibits three distinct regions across the CMB sky (marked by the three large grey circles in Fig.2). These regions have a radius ranging from 40 to 70 degrees. The size of these structures are in agreement with the scale of the causal boundary, χ_\S , for Λ dominated universes. As shown in [11], the size of each of these regions is correlated with the mean value of Ω_Λ over that portion of the sky, in good agreement with the BH.u prediction. The same large-scale anisotropic patterns are observed for the distribution of other basic Λ CDM cosmological parameters. This represents a very significant break-down of the main hypothesis of the Big Bang model: the assumption that the universe is isotropic on large scales. The observed anisotropy has a tiny probability (of order 10^{-9}) of being a Gaussian statistical fluctuation of an otherwise isotropic universe [11].

In summary, Fig.2 shows that: a) regions with a given value of Ω_Λ have a corresponding angular size θ that agree with the BH.u prediction (see Fig.31 in [11]), b) causally disconnected regions of the sky (larger than χ_\S) fit very well the same physical model of acoustic oscillations, c) the background is similar, but with parameters that are significantly different across disconnected regions. This suggests that the underlying physical mechanism sourcing the observed anisotropy encompasses scales beyond our causal universe. This is in apparent tension with simple models of inflation (as source of perturbation for the largest scales) and opens the door to revisiting our basic understanding of the origin of the primordial universe.

Acknowledgments: This work has been supported by spanish MINECO grants PGC2018-102021, ESP2017-89838-C3-1-R, and EU grants LACEGAL 734374 and EWC 776247 with ERDF funds, and grant 2017-SGR-885 of the Generalitat de Catalunya.

-
- [1] T. Padmanabhan, *Gravitation*, Cambridge Univ. Press (2010).
 - [2] A. Mitra, Interpretational conflicts between the static and non-static forms of de Sitter metric, *Nature Sci.Rep.* **2**, 923 (2012).
 - [3] E. Gaztañaga, The cosmological constant as a zero action boundary, *MNRAS* **502**, 436 (2021); E. Gaztanaga, The size of our causal Universe, *MNRAS* **494**, 2766 (2020).
 - [4] N. Kaloper, M. Kleban, and D. Martin, McVittie’s legacy: Black holes in an expanding universe, *Phys. Rev. D* **81**, 104044 (2010).
 - [5] E. Gaztanaga, Inside a Black Hole: the illusion of a Big Bang, hal-03106344, darkcosmos.com (2021).
 - [6] W. Israel, Singular hypersurfaces and thin shells in general relativity, *Nuovo Cimento B Serie* **48**, 463 (1967).
 - [7] W. M. Stuckey, The observable universe inside a black hole, *American Journal of Physics* **62**, 788 (1994), see author corrected version in ResearchGate.
 - [8] A. A. Starobinskiĭ, Spectrum of relict gravitational radiation and the early state of the universe, *Soviet J. of Exp. and Th. Physics Letters* **30**, 682 (1979); A. H. Guth, Inflationary universe: A possible solution to the horizon and flatness problems, *Phys. Rev. D* **23**, 347 (1981); A. D. Linde, A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems, *Physics Letters B* **108**, 389 (1982); A. Albrecht and P. J. Steinhardt, Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking, *Phys. Rev. Lett.* **48**, 1220 (1982).
 - [9] Planck Collaboration, Planck 2018 results. VII. Isotropy and statistics of the CMB, *A&A* **641**, A7 (2020); D. J. Schwarz, C. J. Copi, D. Huterer, and G. D. Starkman, CMB anomalies after Planck, *Classical and Quantum Gravity* **33**, 184001 (2016).
 - [10] Planck Collaboration, Planck 2018 results. VI. Cosmological parameters, *A&A* **641**, A6 (2020); A. G. Riess, The expansion of the Universe is faster than expected, *Nature Reviews Physics* **2**, 10 (2019).
 - [11] P. Fosalba and E. Gaztanaga, Explaining Cosmological Anisotropy: Evidence for Causal Horizons from CMB data, *MNRAS submitted* (2020), arXiv:2011.00910.