

Renormalization group analysis on emergence of higher rank symmetry and higher moment conservation

Hongchao Li^{1,2,*} and Peng Ye^{1,†}

¹*School of Physics and State Key Laboratory of Optoelectronic Materials and Technologies, Sun Yat-sen University, Guangzhou, 510275, China*

²*Department of Modern Physics, University of Science and Technology of China, Hefei, 230026, China*

Higher rank symmetry and higher moment conservation have been drawn considerable attention from, e.g., subdiffusive transport to fracton topological order. In this paper, we perform a one-loop renormalization group (RG) analysis and show how these phenomena emerge at low energies. We consider a d -dimensional model of interacting bosons of d components. At higher-rank-symmetric points with conserved angular moments, the a -th bosons have kinetic energy only along the \hat{x}^a direction. Therefore, the symmetric points look highly anisotropic and fine-tuned. By studying RG in a wide vicinity of the symmetric points, we find that symmetry-disallowed kinetic terms tend to be irrelevant within the perturbative regime, which potentially leads to emergent higher-rank symmetry and higher-moment conservation at the deep infrared limit. While non-perturbative analysis is called for in the future, by regarding higher-rank symmetry as an emergent phenomenon, the RG analysis presented in this paper holds alternative promise for realizing higher-rank symmetry and higher-moment conservation in experimentally achievable systems.

I. INTRODUCTION

The celebrated Noether's theorem [1] relates a conservation law to an underlying continuous symmetry. For example, in a $U(1)$ -symmetric Hamiltonian of bosons, bosonic operator $\hat{\Phi}(\mathbf{x})$ is changed to $e^{i\theta}\hat{\Phi}(\mathbf{x})$ under symmetry transformations where the real parameter θ doesn't depend on coordinate $\mathbf{x} = (x^1, x^2, \dots, x^d)$ in d -dimensional space. By means of Noether's theorem, one can show that the total boson number, i.e., $\int d^d x \rho(\mathbf{x})$, is a conserved quantity, where particle number density $\rho(\mathbf{x}) = \hat{\Phi}^\dagger(\mathbf{x})\hat{\Phi}(\mathbf{x})$. Apparently, $\rho(\mathbf{x})$ is just the zeroth order of conventional multipole expansions:

$$\rho, \rho \cdot \mathbf{x}, \rho x^a x^b, \dots \quad (1)$$

in a standard electromagnetism textbook [2]. In a particle-number-conserving system, *higher moment conservation*, e.g., conservation of dipoles and quadrupoles, is in principle allowed. Furthermore, if the density is vector-like with multi-components, denoted as $\boldsymbol{\rho} = (\rho^1, \rho^2, \dots, \rho^d)$, then we can define another set of multipole expansions:

$$\boldsymbol{\rho}, \boldsymbol{\rho} \cdot \mathbf{x}, \sum_{e,f=1}^d \epsilon^{ab\dots ef} \rho^e x^f, \dots \quad (2)$$

where the third one is *angular moment*. $\epsilon^{ab\dots}$ is Levi-Civita symbol. If $d = 2, 3$, it can be rewritten in a compact form: $\boldsymbol{\rho} \times \mathbf{x}$. In $d = 2$, $\boldsymbol{\rho} \times \mathbf{x} = \rho^1 x^2 - \rho^2 x^1$.

Indeed, recently we have been witnesses to ongoing research progress on higher-moment conservation and the

associated higher-rank version of global symmetry [3–9], especially in the field of fracton physics [3–69]. Some typical examples of research include subdiffusive transport at late times, non-ergodicity, Hilbert space fragmentation, and spontaneous symmetry breaking [31, 33, 41, 68–74]. In a simple scalar theory, the associated higher-rank symmetry transformations are parametrized by $\theta(\mathbf{x})$ that is a polynomial function of \mathbf{x} [8]. Inspired by the conventional correspondence between global symmetry and gauge symmetry, upon “gauging” higher-rank symmetry, higher-rank gauge fields can be obtained [5]. Here, the gauge fields are usually higher-rank symmetric tensor fields, which leads to generalized Maxwell equations [6] and exotic theory of spin systems in Yb-based breathing pyrochlores [75].

As a nontrivial consequence of higher moment conservation, the mobility of particles is inevitably restricted, either partially or completely. For example, it is quite intuitive that dipole conservation strictly forbids a single particle motion along all spatial directions. Such particles are called “fractons” or 0-dimensional particles [3, 4]. Similarly, one can define *lineons* (1-dimensional particle) that are movable within a stack of parallel straight lines and *planons* (2-dimensional particle) that are movable within a stack of parallel planes. Regarding these strange particles as bosons, we can consider their Bose-Einstein condensation, such that the spontaneous breaking of higher-rank symmetry occurs. As a result, a class of exotic quantum phases of matter dubbed *fractonic superfluids* [68, 69] is formed. In Ref. [69], a convenient notation $d\text{SF}^i$ was introduced to denote d -dimensional fractonic superfluids with i -dimensional particle condensation, e.g., $d\text{SF}^0$ with condensed fractons and $d\text{SF}^1$ with condensed lineons. The conventional superfluid phase corresponds to $d\text{SF}^d$ where bosons can freely move.

Higher-rank symmetric microscopic models often look quite unrealistic, highly anisotropic and fine-tuned [5]. For example, Hamiltonian does not has the usual kinetic

* Present address: Department of Physics, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

† yepeng5@mail.sysu.edu.cn

energy term [68]. And interaction is delicately designed [69]. However, as we've known, in many condensed matter systems, symmetry has been found to be significantly enhanced at low energies. For example, Lorentz symmetry emerge in graphene which is microscopically built by non-relativistic electrons. Thus, one may wonder whether it is possible that long-wavelength low-energy limit will conserve higher moments and respect higher-rank symmetry as an emergent phenomenon.

For this purpose, we may apply the traditional theoretical approach: renormalization group (RG). If there exists a phase region such that all models in the region can flow to symmetric points, we can regard higher-rank symmetry as an *emergent symmetry*. Theoretically, one advantage of such an emergent higher-rank symmetry is its robustness against symmetry-breaking perturbation. Practically, we expect that such a scenario holds promise for *more flexible realization* of exotic higher-rank symmetry and higher-moment conservation in both theoretical and experimental studies.

In this paper, we identify such a wide phase region that supports emergent higher-rank symmetry and conservation of *angular moments*, i.e., $\int d^2x \boldsymbol{\rho} \times \mathbf{x} = \int d^2x (\rho^1 x^2 - \rho^2 x^1)$ for a two-component boson fields in two dimensions. We start with a two-dimensional many-boson system in the normal state (i.e. without lineon condensation) of fractonic superfluids (denoted as 2SF¹). The Hamiltonian is a symmetric point in the parameter space where the a -th ($a = 1, 2$) component bosons only have kinetic terms along a -th axis (dubbed “*diagonal*” kinetic terms). There also exists a weak inter-component scattering term allowed by higher-rank symmetry. We shall perform a RG analysis in the vicinity of the symmetric point by adding symmetry-disallowed kinetic terms (dubbed “*off-diagonal*” kinetic terms) as a perturbation. The one-loop calculation of β -function shows that there exists a finite phase region (Fig. 4) where off-diagonal kinetic terms tend to be irrelevant under RG iteration. In other words, the high-energy model, which is not symmetric but more realistic and less fine-tuned, has a tendency to flow to the symmetric point. As a result, higher-rank symmetry as well as conservation of angular moments emerges.

The remainder of this paper is organized as follows. In Sec. II, we introduce the d -component bosonic systems and its higher-rank symmetry. In Sec. III, we discuss the scaling and Feynmann rules of the d -component bosonic systems. Further, we figure out the β -functions of parameters in the systems with renormalization group (RG) analysis and depict the global phase diagram. In Sec. IV, we summarize and provide our prediction on conditions of possible realization of the systems with higher-rank symmetry.

II. MODEL AND SYMMETRY

The symmetric point Hamiltonian for the d -component bosonic systems in real space [69] is given by $\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_1$ with

$$\mathcal{H}_0 = \sum_{a=1}^d \left[\frac{t_a}{2} (\partial_a \hat{\Phi}_a^\dagger) (\partial_a \hat{\Phi}_a) - \mu \hat{\Phi}_a^\dagger \hat{\Phi}_a \right], \quad (3)$$

$$\mathcal{H}_1 = \frac{1}{2} \sum_{a \neq b} K_{ab} (\hat{\Phi}_a^\dagger \partial_a \hat{\Phi}_b^\dagger + \hat{\Phi}_b^\dagger \partial_b \hat{\Phi}_a^\dagger) (\hat{\Phi}_a \partial_a \hat{\Phi}_b + \hat{\Phi}_b \partial_b \hat{\Phi}_a), \quad (4)$$

where \mathcal{H}_0 is free Hamiltonian and \mathcal{H}_1 is interaction Hamiltonian. Here $t_a = m_a^{-1}$ stands for the inverse of mass along the a -th direction and μ stands for the chemical potential. $\hat{\Phi}_a^\dagger(\mathbf{x})$ and $\hat{\Phi}_a(\mathbf{x})$ are respectively creation and annihilation operators of d -component bosons, and satisfy the bosonic commutation relations. The interaction strength K_{ab} is a symmetric matrix with vanishing diagonal elements, i.e., $K_{aa} = 0, K_{ab} = K_{ba}$. Each term in \mathcal{H} is invariant under both the conventional global symmetry transformations ($\hat{\Phi}_a \rightarrow e^{i\theta_a} \hat{\Phi}_a$, $\theta_a \in \mathbb{R}$) and higher-rank symmetry transformations:

$$(\hat{\Phi}_a, \hat{\Phi}_b) \longrightarrow (\hat{\Phi}_a e^{i\lambda_{ab} x^b}, \hat{\Phi}_b e^{i\lambda_{ba} x^a}) \quad (5)$$

for each pair (ϕ_a, ϕ_b) with $\lambda_{ab} = -\lambda_{ba} \in \mathbb{R}$. According to the Noether's theorem, the conventional global $U(1)$ symmetry and the higher-rank symmetry correspond to conserved total charge (particle number) \hat{Q}^a and conserved total angular moments \hat{Q}^{ab} ($\hat{Q}^{ab} = -\hat{Q}^{ba}$) [1, 5, 69]:

$$\hat{Q}^a = \int d^d x \hat{\rho}^a, \quad \hat{Q}^{ab} = \int d^d x (\hat{\rho}^a x^b - \hat{\rho}^b x^a). \quad (6)$$

Here $\hat{\rho}^a = \hat{\Phi}_a^\dagger \hat{\Phi}_a$. Intuitively, the conserved quantities \hat{Q}^{ab} enforce that a single a -th component boson can only move along the a -th direction. More explanation on the classical mechanical consequence of the conservation is available in Appendix A and Ref. [69].

In the coherent-state path-integral formulation with imaginary time, the Lagrangian density \mathcal{L} can be written as $\mathcal{L} = \phi_a^* \partial_\tau \phi_a + \mathcal{H}$ with action $S = \int d\tau d^d x \mathcal{L}$. Here the bosonic fields $\phi_a = \phi_a(\mathbf{x}, \tau) \in \mathbb{C}$ are the eigenvalues of coherent-state operators $\hat{\Phi}_a(\mathbf{x}, \tau)$. With the Fourier transformation of the coherent state: $\phi_a(\tau, \mathbf{r}) = \frac{1}{\sqrt{\beta}} \sum_n \int \frac{d^d k}{(2\pi)^d} e^{i\mathbf{k} \cdot \mathbf{r} - i\omega_n \tau} \phi_a(i\omega_n, \mathbf{k})$ and its complex conjugate: $\phi_a^*(\tau, \mathbf{r}) = \frac{1}{\sqrt{\beta}} \sum_n \int \frac{d^d k}{(2\pi)^d} e^{-i\mathbf{k} \cdot \mathbf{r} + i\omega_n \tau} \phi_a^*(i\omega_n, \mathbf{k})$ for $a = 1, 2, \dots, d$ with $\beta = \frac{1}{k_B T}$ where T represents the temperature of bosons. In the frequency-momentum space,

$$S = \sum_{a,k} (-i\omega_n + \xi_a) \phi_a^* \phi_a + \sum_{\substack{1,2,3,4 \\ a \neq b}} \frac{K_{ab}}{2\beta} (k_2^a + k_1^b) (k_3^a + k_4^b) \cdot \phi_{1a}^* \phi_{2b}^* \phi_{3b} \phi_{4a} \delta(1+2-3-4). \quad (7)$$




FIG. 1. The left one shows the bare interaction vertex of d -component bosons. Here the solid lines are Feynman propagators and the vertex is the coefficient: $\frac{1}{2\beta} K_{ab}(k_2^a + k_1^b)(k_3^a + k_4^b)\delta(1+2-3-4)$. The right one represents the Feynman loop diagram of the correction for the parameter \mathcal{T}_1 .

In the first term on the r.h.s., the simplified notation ϕ_a stands for $\phi_a(i\omega_n, \mathbf{k})$ which is the frequency-momentum image of the field $\phi_a(\tau, \mathbf{r})$. ω_n is a bosonic Matsubara frequency and \mathbf{k} is a momentum vector: $\mathbf{k} = (k^1, k^2, \dots, k^d)$. We use k^a to denote the a -th spatial component of momentum vector \mathbf{k} . $\sum_{\mathbf{k}}$ stands for $\sum_{\omega_n} \int \frac{d^d k}{(2\pi)^d}$. In the second term, since there are four pairs of frequency and momentum, we introduce a new notation ϕ_{ia} to compactly represent $\phi_a(i\omega_{n_i}, \mathbf{k}_i)$ where the label $i = 1, 2, \dots, 4$. k_i^a stands for the a -th spatial component of momentum vector \mathbf{k}_i . The sum $\sum_{1,2,3,4}$ denotes $\sum_{i=1}^4 \sum_{\omega_{n_i}} \int \frac{d^d k_i}{(2\pi)^d}$. Other notations like $\sum_{1,2}, \sum_2, \dots$ in the forthcoming texts are defined in the similar way. Besides, the kinetic energy with momentum \mathbf{k} is defined as: $\xi_a = \frac{1}{2} t_a (k^a)^2 - \mu$ (In the following text, we began to introduce anisotropic kinetic energy). For momentum \mathbf{k}_i , the associated kinetic energy is $\xi_{ia} = \frac{1}{2} \sum_{b=1}^d t_{ab} (k_i^b)^2 - \mu$. Last, we use $\delta(1+2-3-4)$ to represent $\delta_{(n_1+n_2), (n_3+n_4)} (2\pi)^d \delta(\mathbf{k}_1 + \mathbf{k}_2 - \mathbf{k}_3 - \mathbf{k}_4)$. $\delta_{(n_1+n_2), (n_3+n_4)}$ is a dimensionless Kronecker symbol.

Before moving forward, we perturb the Lagrangian density by adding small ‘‘off-diagonal’’ kinetic terms that break higher-rank symmetry. They are the terms deviating the model from the symmetric point. As such, kinetic terms of both directions are present, which can be written as $\sum_{a,b=1}^d \left[\frac{t_{ab}}{2} (\partial_b \phi_a^*) (\partial_b \phi_a) \right] = \sum_{a=1}^d \left[\frac{t_{aa}}{2} (\partial_a \phi_a^*) (\partial_a \phi_a) \right] + \sum_{a \neq b} \left[\frac{t_{ab}}{2} (\partial_b \phi_a^*) (\partial_b \phi_a) \right]$. The kinetic parameter t_a is rewritten as t_{aa} for the notational convenience. Those off-diagonal kinetic terms with nonzero t_{ab} ($a \neq b$) manifestly break higher-rank symmetry. Similarly, we can also understand these parameters as inverse of mass of field configuration ϕ_a along other directions other than the a -th one: $t_{ab} = 1/m_{ab}$. In this way, the kinetic energy with momentum \mathbf{k} can be redefined as: $\xi_a = \frac{1}{2} \sum_{a'=1}^d t_{aa'} (k^{a'})^2 - \mu$.

III. RENORMALIZATION GROUP ANALYSIS

A. Scaling and Feynman rules

We consider $d = 2$. We set the restriction of field configuration ϕ_a as: $\sum_{b=1}^2 \frac{t_{ab}}{t_a} (k^b)^2 \leq \Lambda^2$ since our total kinetic energy is given by $\sum_b \frac{t_{ab}}{t_a} (k^b)^2$ with surfaces of equal energy: $\sum_b \frac{t_{ab}}{t_a} (k^b)^2 = \kappa^2$. Here κ is an arbitrary constant with the momentum dimension and Λ is the cut-off of momentum here. The high-energy part corresponds to: $\sqrt{\sum_{b=1}^2 \frac{t_{ab}}{t_a} (k^b)^2} \in [\Lambda/s, \Lambda]$, where the scaling parameter $s > 1$ and sends \mathbf{k} to \mathbf{k}/s . We also define $s = e^l$ with $l > 0$. We consider the free part of the action $\sum_n \int \frac{d^2 k}{(2\pi)^2} \phi_a^* \left(-i\omega_n + \sum_{b=1}^2 \frac{t_{ab}(k^b)^2}{2} \right) \phi_a$. In the following text, we also call it fast modes and denote it with \rangle . It is noticed that we do not take the part related to chemical potential into account since it must be relevant if we choose the kinetic part to be marginal. Suppose the scaling dimension of ϕ is Δ_ϕ , and we can assume the change of temperature and frequencies are described as: $T \rightarrow s^{-z} T$, and $\omega_n \rightarrow s^{-z} \omega_n$, $n \in \mathbb{Z}$ respectively. So the free part is scaled to $\sum_n \int \frac{d^2 k}{(2\pi)^2} \phi_a^* \left(-i\omega_n s^{-z} + \sum_{b=1}^2 \frac{t_{ab}(k^b)^2}{2} s^{-2} \right) \phi_a s^{2\Delta_\phi - 2}$. To fix the momentum dependence, we need $\Delta_\phi = \frac{d}{2} + 1 = 2$ and $z = 2$. In $d = 2$ case considered here, the interaction matrix is simply determined by a single parameter, i.e., $K_{12} = K_{21} := \mathcal{K}$. For further simplifying calculation for $d = 2$, we assume $t_1 := t_{11} = t_2 := t_{22} := \mathcal{T}_0$, $t_{12} = t_{21} := \mathcal{T}_1$. There are two small parameters compared to \mathcal{T}_0 . The first one is symmetry-disallowed off-diagonal kinetic parameter \mathcal{T}_1 which is marginal. It can also be seen as a perturbation relative to the interaction parameter $\mathcal{K}\Lambda^2$. The second one is the irrelevant interaction parameter $\mathcal{K}\Lambda^2$ considered as an infinitesimal quantity compared with diagonal kinetic parameter \mathcal{T}_0 . Hence, the following calculation will be proceeded in the perturbative regime: $\mathcal{T}_0 \gg \mathcal{K}\Lambda^2 \gg \mathcal{T}_1$, where Λ is the momentum cut-off.

Next, we write Feynman rules. The bare Feynman propagators are given by: $\langle \phi_{ia}^* \phi_{jb} \rangle = (i\omega_{n_i} - \xi_{ia})^{-1} \delta_{ab} \delta_{ij}$ with path-integral quantization. Here $\xi_{ia} = \frac{\sum_{a'=1}^d t_{aa'} (k_i^{a'})^2}{2} - \mu$ stands for the kinetic energy of ϕ_{ia} [see definition below Eq. (7)]. In addition, the interaction vertex is drawn in Fig. 1. This diagram represents the interaction term, i.e. the second term in Eq. (7), where solid lines represent the bosonic fields and vertex stands for the interaction coefficient. In the RG procedure to be performed, we apply the standard cumulant expansion and relate the mean of the exponential to the exponential of the means [76, 77]: $\langle e^{-S_{int}} \rangle = e^{-\langle S_{int} \rangle + \frac{1}{2} (\langle S_{int}^2 \rangle - \langle S_{int} \rangle^2) + \dots}$. Here ϕ_{\rangle} and ϕ_{\langle} respectively correspond to the fast modes and slow modes of bosonic fields and S_{int} stands for interaction part of action. In this case, the notation $\langle \rangle_{\rangle}$ is to take the average over the fast modes. Therefore, after

calculating the average on fast modes, we have the form of the effective action. $S_{eff}[\phi_{<}] = S_0[\phi_{<}] + S'_{int}[\phi_{<}] = S_0[\phi_{<}] + \langle S_{int} \rangle_{>}[\phi_{<}] - \frac{1}{2}(\langle S_{int}^2 \rangle_{>} - \langle S_{int} \rangle_{>}^2)[\phi_{<}]$. The form of $\langle S_{int} \rangle_{>}$ contributes to the β -functions of kinetic parameters, also named first-order correction. Further, $\frac{1}{2}(\langle S_{int}^2 \rangle_{>} - \langle S_{int} \rangle_{>}^2)[\phi_{<}]$ leads to the β -functions of the elements in the K matrix.

B. One-loop correction to \mathcal{T}_1

The interaction part of action reads:

$$S_{int} = \frac{1}{2\beta} \sum_{1,2,3,4} \sum_{a \neq b} K_{ab}(k_2^a + k_1^b)(k_3^a + k_4^b) \phi_{1a}^* \phi_{2b}^* \phi_{3b} \phi_{4a} \delta(1+2-3-4). \quad (8)$$

$$\begin{aligned} S'_{int} &= \frac{1}{2\beta'} s^{-z} \sum_{i=1}^4 \sum_{\omega'_{n_i}} \int \frac{d^d k'_i}{(2\pi)^d} s^{-4d} K_{ab}((k'_2)^a + (k'_1)^b)((k'_3)^a + (k'_4)^b) s^{-2} \\ &\quad \cdot \phi_a^*(i\omega'_{n_1}, \mathbf{k}'_1) \phi_b^*(i\omega'_{n_2}, \mathbf{k}'_2) \phi'_b(i\omega'_{n_3}, \mathbf{k}'_3) \phi'_a(i\omega'_{n_4}, \mathbf{k}'_4) s^{4\Delta_\phi} \delta_{(n'_1+n'_2), (n'_3+n'_4)} (2\pi)^d \delta(\mathbf{k}'_1 + \mathbf{k}'_2 - \mathbf{k}'_3 - \mathbf{k}'_4) s^d \\ &= \frac{1}{2\beta'} s^{2-z-d} \sum_{i=1}^4 \sum_{\omega'_{n_i}} \int \frac{d^d k'_i}{(2\pi)^d} \sum_{a \neq b} K_{ab}((k'_2)^a + (k'_1)^b)((k'_3)^a + (k'_4)^b) \\ &\quad \cdot \phi_a^*(i\omega'_{n_1}, \mathbf{k}'_1) \phi_b^*(i\omega'_{n_2}, \mathbf{k}'_2) \phi'_b(i\omega'_{n_3}, \mathbf{k}'_3) \phi'_a(i\omega'_{n_4}, \mathbf{k}'_4) \delta_{(n'_1+n'_2), (n'_3+n'_4)} (2\pi)^d \delta(\mathbf{k}'_1 + \mathbf{k}'_2 - \mathbf{k}'_3 - \mathbf{k}'_4). \end{aligned} \quad (9)$$

Here the dimensionful δ -function of momenta is also scaled [$\delta(\mathbf{k}') = s^{-d}\delta(\mathbf{k})$] but the Kronecker symbol of Matsubara frequencies is dimensionless and invariant upon scaling. Therefore, the scaling of K_{ab} is given by $K'_{ab} = K_{ab}s^{-d}$. It illustrates that the K matrix at tree level is irrelevant in perturbative RG.

Then, let us consider the first-order correction to \mathcal{T}_1 : $\langle S_{int} \rangle_{>}$ contributing to the correction of kinetic parameters.

$$\langle S_{int} \rangle_{>} = \left\langle \frac{1}{2\beta} \sum_{1,2,3,4} \sum_{a \neq b} K_{ab}(k_2^a + k_1^b)(k_3^a + k_4^b) \cdot \phi_{1a}^* \phi_{2b}^* \phi_{3b} \phi_{4a} \delta(1+2-3-4) \right\rangle_{>}. \quad (10)$$

To proceed further, we should split each momentum integration into slow part ($<$) and fast part ($>$). In this way, the integration of four momenta ($\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3, \mathbf{k}_4$) is split to $2^4 = 16$ combinations. According to Wick's theorem, for formulas of bare propagators, we conclude that either ϕ_{1a}^* and ϕ_{4a} must be paired or ϕ_{2b}^* and ϕ_{3b} must be paired. Other contractions vanish due to $K_{ab} = 0$ when $a = b$. Therefore, in 16 combinations, only the following two are non-vanishing:

- **Case-I:** The momenta carried by ϕ_{1a}^* and ϕ_{4a} , i.e.,

Below we determine the scaling of the K matrix. To be specific, we define the new momentum and frequency as: $\mathbf{k}' = s\mathbf{k}$, $\omega'_n = s^z\omega_n$ or $T' = s^z T$ with $z = 2$. We define the rescaled field: $\phi'_a(i\omega'_n, \mathbf{k}') = s^{-\Delta_\phi} \phi_a(i\omega'_n s^{-z}, \frac{\mathbf{k}'}{s})$ with $\Delta_\phi = \frac{d}{2} + 1$. Then the scaling of parameters K_{ab} can be determined. With the form of interacting action (8), we obtain:

\mathbf{k}_1 and \mathbf{k}_4 , are fast momenta¹. $\mathbf{k}_1 = \mathbf{k}_4$ and $\omega_{n_1} = \omega_{n_4}$ required by the formula of bare propagator;

- **Case-II:** The momenta carried by ϕ_{2b}^* and ϕ_{3b} , i.e., \mathbf{k}_2 and \mathbf{k}_3 , are fast momenta. $\mathbf{k}_2 = \mathbf{k}_3$ and $\omega_{n_2} = \omega_{n_3}$ required by the formula of bare propagator.

By further considering $\delta(1+2-3-4)$, we have:

- **Case-I:** The momenta carried by ϕ_{1a}^* and ϕ_{4a} , i.e., \mathbf{k}_1 and \mathbf{k}_4 , are fast momenta. $\mathbf{k}_1 = \mathbf{k}_4$, $\omega_{n_1} = \omega_{n_4}$, $\mathbf{k}_2 = \mathbf{k}_3$, and $\omega_{n_2} = \omega_{n_3}$;
- **Case-II:** The momenta carried by ϕ_{2b}^* and ϕ_{3b} , i.e., \mathbf{k}_2 and \mathbf{k}_3 , are fast momenta. $\mathbf{k}_2 = \mathbf{k}_3$ and $\omega_{n_2} = \omega_{n_3}$, $\mathbf{k}_1 = \mathbf{k}_4$, and $\omega_{n_1} = \omega_{n_4}$.

These two contractions correspond to the Feynman diagram in Fig. 1. We firstly focus on case-I. When 1,4 are fast modes, the momentum conservation and Feynman rules will tell us: $\mathbf{k}_2 = \mathbf{k}_3, \mathbf{k}_1 = \mathbf{k}_4; \omega_{n_2} = \omega_{n_3}, \omega_{n_1} = \omega_{n_4}$. Since we focus on the two-dimensional case ($d = 2$), we set all off-diagonal K -matrix elements

¹ By fast (slow) momentum, we mean that the kinetic energy corresponding to the momentum is large (small).

as: $K_{12} = K_{21} := \mathcal{K}$. For further simplifying calculation, we assume $t_1 := t_{11} = t_2 := t_{22} := \mathcal{T}_0$, $t_{12} = t_{21} := \mathcal{T}_1$. In this way, $\langle S_{int} \rangle^I$ is given by:

$$\begin{aligned}
\langle S_{int} \rangle^I &= \frac{1}{2\beta} \sum_{a \neq b} \sum_{1,2} \mathcal{K}(k_2^a + k_1^b)^2 \langle \phi_{1a}^* \phi_{1a} \rangle \phi_{2b}^* \phi_{2b} \\
&= \frac{1}{2\beta} \sum_{a \neq b} \sum_{1,2} \mathcal{K}(k_2^a + k_1^b)^2 \frac{1}{i\omega_{n_1} - \xi_{1a}} \phi_{2b}^* \phi_{2b} \\
&= -\frac{1}{2} \sum_{a \neq b} \sum_2 \int_{>} \frac{d^2 k_1}{(2\pi)^2} \mathcal{K}(k_2^a + k_1^b)^2 f_B(\xi_{1a}) \phi_{2b}^* \phi_{2b} \\
&= -\frac{1}{2} \sum_{a \neq b} \sum_2 \int_{>} \frac{d^2 k_1}{(2\pi)^2} \mathcal{K} [(k_2^a)^2 + 2k_2^a k_1^b + (k_1^b)^2] \\
&\quad \cdot f_B(\xi_{1a}) \phi_{2b}^* \phi_{2b}, \tag{11}
\end{aligned}$$

where \mathbf{k}_1 and \mathbf{k}_2 correspond to fast and slow momenta respectively. The bosonic Matsubara summation is applied: $\frac{1}{\beta} \sum_{\omega_n} \frac{1}{i\omega_n - \xi_{1a}} = -f_B(\xi_{1a}) = -\frac{1}{e^{\beta\xi_{1a}} - 1}$. And the propagator $\langle \phi_{1a}^* \phi_{1a} \rangle$ is given by: $\langle \phi_{1a}^* \phi_{1a} \rangle = \frac{1}{i\omega_{n_1} - \xi_{1a}}$ with ξ_{1a} ($a = 1, b = 2$ or $a = 2, b = 1$):

$$\xi_{1a} := \frac{\mathcal{T}_0(k_1^a)^2 + \mathcal{T}_1(k_1^b)^2}{2} - \mu. \tag{12}$$

In the same way, we can give the form of case-II:

$$\begin{aligned}
\langle S_{int} \rangle^{II} &= \frac{1}{2\beta} \sum_{a \neq b} \sum_{1,2} \mathcal{K}(k_2^a + k_1^b)^2 \langle \phi_{2b}^* \phi_{2b} \rangle \phi_{1a}^* \phi_{1a} \\
&= \frac{1}{2\beta} \sum_{a \neq b} \sum_{1,2} \mathcal{K}(k_2^a + k_1^b)^2 \frac{1}{i\omega_{n_2} - \xi_{2b}} \phi_{1a}^* \phi_{1a} \\
&= -\frac{1}{2} \sum_{a \neq b} \sum_1 \int_{>} \frac{d^2 k_2}{(2\pi)^2} \mathcal{K}(k_2^a + k_1^b)^2 f_B(\xi_{2b}) \phi_{1a}^* \phi_{1a} \\
&= -\frac{1}{2} \sum_{a \neq b} \sum_1 \int_{>} \frac{d^2 k_2}{(2\pi)^2} \mathcal{K} [(k_2^a)^2 + 2k_2^a k_1^b + (k_1^b)^2] \\
&\quad \cdot f_B(\xi_{2b}) \phi_{1a}^* \phi_{1a}.
\end{aligned}$$

By observing these two actions, we find they are completely equivalent to each other by exchanging the indices ($a \leftrightarrow b, 1 \leftrightarrow 2$). Then we focus on the case-I and multiply it by 2, namely:

$$\begin{aligned}
\langle S_{int} \rangle &= \langle S_{int} \rangle^I + \langle S_{int} \rangle^{II} = 2\langle S_{int} \rangle^I, \\
&= -\sum_{a \neq b} \sum_2 \int_{>} \frac{d^2 k_1}{(2\pi)^2} \mathcal{K} [(k_2^a)^2 + 2k_2^a k_1^b + (k_1^b)^2] \\
&\quad \cdot f_B(\xi_{1a}) \phi_{2b}^* \phi_{2b} = \langle S_{int} \rangle^{(1)} + \langle S_{int} \rangle^{(2)} + \langle S_{int} \rangle^{(3)},
\end{aligned}$$

where we arrange all terms into three parts:

$$\langle S_{int} \rangle^{(1)} := -\sum_{a \neq b} \sum_2 \mathcal{K}(k_2^a)^2 \int_{>} \frac{d^2 k_1}{(2\pi)^2} f_B(\xi_{1a}) \phi_{2b}^* \phi_{2b}, \tag{13}$$

$$\langle S_{int} \rangle^{(2)} := -2 \sum_{a \neq b} \sum_2 \mathcal{K} k_2^a \int_{>} \frac{d^2 k_1}{(2\pi)^2} k_1^b f_B(\xi_{1a}) \phi_{2b}^* \phi_{2b}, \tag{14}$$

$$\langle S_{int} \rangle^{(3)} := -\sum_{a \neq b} \sum_2 \mathcal{K} \int_{>} \frac{d^2 k_1}{(2\pi)^2} f_B(\xi_{1a}) (k_1^b)^2 \phi_{2b}^* \phi_{2b}. \tag{15}$$

The fast momentum $\mathbf{k}_1 = (k_1^1, k_1^2)$ takes values in the elliptic shell (Λ is the momentum cut-off and the RG flow parameter $s = e^l$ with $s \rightarrow 1$ and $l \rightarrow 0$):

$$\left(\frac{\Lambda}{s}\right)^2 < (k_1^a)^2 + \frac{\mathcal{T}_1}{\mathcal{T}_0} (k_1^b)^2 < \Lambda^2 \tag{16}$$

which defines the domain of integral $\int_{>}$. Here we should focus on the integrated domain of fast momenta. It is only determined by the fields carrying those fast momenta. To be specific, we consider the fast momentum \mathbf{k}_i carried by the fast mode ϕ_{ia} or ϕ_{ia}^* with the label $i = 1, 2, \dots, 4$ defined above. Its integrated domain is given by $(\Lambda/s)^2 < (k_i^a)^2 + \frac{\mathcal{T}_1}{\mathcal{T}_0} (k_i^b)^2 < \Lambda^2$ for $b \neq a$. This conclusion will be used in the following text. To compute $\langle S_{int} \rangle^{(1)}$ in Eq. (13), we may introduce a new momentum $\tilde{\mathbf{k}}$: $\tilde{k}^a = k_1^a, \tilde{k}^b = \sqrt{\frac{\mathcal{T}_1}{\mathcal{T}_0}} k_1^b$. Then, Eq. (16) is changed to: $(\frac{\Lambda}{s})^2 < (\tilde{k}^a)^2 + (\tilde{k}^b)^2 < \Lambda^2$, i.e., $(\frac{\Lambda}{s})^2 < \tilde{\mathbf{k}}^2 < \Lambda^2$. And Eq. (12) is changed to:

$$\xi_{1a} = \frac{\mathcal{T}_0 [(\tilde{k}^a)^2 + (\tilde{k}^b)^2]}{2} - \mu = \frac{\mathcal{T}_0 \tilde{\mathbf{k}}^2}{2} - \mu. \tag{17}$$

Using the new momentum variables, we can work out the integral $\int_{>}$ in $\langle S_{int} \rangle^{(1)}$ in Eq. (13) ($s-1$ is small enough):

$$\begin{aligned}
&\int_{>} \frac{d^2 k_1}{(2\pi)^2} f_B(\xi_{1a}) \\
&= \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \int_{>} \frac{d^2 \tilde{k}}{(2\pi)^2} f_B(\xi_{1a}) \\
&= \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \frac{1}{(2\pi)^2} \int_{\Lambda/s}^{\Lambda} 2\pi |\tilde{\mathbf{k}}| d|\tilde{\mathbf{k}}| \frac{1}{e^{\beta\xi_{1a}} - 1} \\
&= \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \frac{1}{(2\pi)^2} \int_{(\Lambda/s)^2}^{\Lambda^2} \frac{2\pi}{2} d(|\tilde{\mathbf{k}}|^2) \frac{1}{e^{\beta\xi_{1a}} - 1} \\
&\approx \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \frac{1}{(2\pi)^2} \frac{2\pi}{2} \frac{1}{e^{\beta\xi_{0\Lambda}} - 1} \left[\Lambda^2 - \left(\frac{\Lambda}{s}\right)^2 \right] \\
&\approx \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \frac{1}{(2\pi)^2} \frac{2\pi}{2} \frac{1}{e^{\beta\xi_{0\Lambda}} - 1} 2\Lambda^2 l = \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \frac{1}{2\pi} \frac{1}{e^{\beta\xi_{0\Lambda}} - 1} \Lambda^2 l,
\end{aligned}$$

where we have considered infinitesimal $s - 1$ and the definition of $s = e^l$. And, $\xi_{0\Lambda} := \frac{\mathcal{T}_0}{2}\Lambda^2 - \mu$. As a result,

$$\langle S_{int} \rangle^{(1)} = -S_2 f_B(\xi_{0\Lambda}) \frac{\Lambda^2}{(2\pi)^2} \sum_{a \neq b} \sum_2 \mathcal{K}(k_2^a)^2 \phi_{2b}^* \phi_{2b} l, \quad (18)$$

where $S_2 := 2\pi \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}}$ and $f_B(\xi_{0\Lambda}) := \frac{1}{e^{\beta \xi_{0\Lambda}} - 1}$. Below, we will use dl to replace l since it is infinitesimal.

The second one $\langle S_{int} \rangle^{(2)}$ in Eq. (14) vanishes since it is an odd function of the integrated momentum \mathbf{k}_1 . The third term in Eq. (15) is directly connected to correction of the chemical potential. We just briefly give our results here for it is not our focus:

$$\langle S_{int} \rangle^{(3)} = -\frac{1}{2} \sum_{a \neq b} \sum_2 \Lambda^4 \frac{S_2}{(2\pi)^2} f_B(\xi_{a\Lambda}) K \phi_{2b}^* \phi_{2b} \left(1 - \frac{1}{s}\right).$$

Therefore, the only term that can renormalize \mathcal{T}_1 is $\langle S_{int} \rangle^{(1)}$ in Eq. (18). The bare off-diagonal kinetic term is given by $\sum_{a \neq b} \sum_k \frac{\mathcal{T}_1}{2} (k_2^a)^2 \phi_{2b}^* \phi_{2b}$ in the frequency-momentum space. Hence, we have the β -function of the parameter \mathcal{T}_1 by referencing the effective action: $S_{eff}[\phi_{<}] = S_0[\phi_{<}] + \langle S_{int} \rangle[\phi_{<}] - \frac{1}{2} (\langle S_{int}^2 \rangle - \langle S_{int} \rangle^2)[\phi_{<}]$.

$$\frac{d\mathcal{T}_1}{dl} = -S_2 \mathcal{K} f_B(\xi_{a\Lambda}) \frac{\Lambda^2}{(2\pi)^2} = -\sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \mathcal{K} f_B(\xi_{a\Lambda}) \frac{\Lambda^2}{2\pi}. \quad (19)$$

This corresponds to Fig. 1. Besides, by referencing the term containing the chemical potential, we have the β -function of the chemical potential μ : $\frac{d\mu}{dl} = 2\mu - \frac{1}{2} \Lambda^4 \frac{S_2}{(2\pi)^2} f_B(\xi_{a\Lambda}) \mathcal{K}$ with the term 2μ originating from the contribution from the tree-level diagrams.

C. Vertex correction to \mathcal{K}

We then turn to the vertex correction with all the contractions contained in $\langle S_{int}^2 \rangle$:

$$\begin{aligned} \langle S_{int}^2 \rangle &= \frac{1}{4\beta^2} \sum_{a \neq b} \sum_{a' \neq b'} \sum_{1,2,3,4} \sum_{1',2',3',4'} \\ &K_{ab} K_{a'b'} \langle \phi_{1a}^* \phi_{2b}^* \phi_{3b} \phi_{4a} \phi_{1'a'}^* \phi_{2'b'}^* \phi_{3'b'} \phi_{4'a'} \rangle \\ &\cdot (k_2^a + k_1^b)(k_3^a + k_4^b)(k_{2'}^{a'} + k_{1'}^{b'})(k_{3'}^{a'} + k_{4'}^{b'}) \\ &\cdot \delta(1+2-3-4)\delta(1'+2'-3'-4') \\ &= \langle S_{int}^2 \rangle^{(1)} + \langle S_{int}^2 \rangle^{(2)} \end{aligned} \quad (20)$$

Since we only consider the loop-diagram contribution, only two kinds of contractions will not vanish under calculation. The first term $\langle S_{int}^2 \rangle^{(1)}$ can be concluded as four cases:

- **Case 1:** The momenta $\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3'$ and \mathbf{k}_4' are fast momenta. $\mathbf{k}_1 = \mathbf{k}_3'$, $\omega_{n_1} = \omega_{n_3'}$; $\mathbf{k}_2 = \mathbf{k}_4'$ and $\omega_{n_2} = \omega_{n_4'}$.




FIG. 2. The Feynman diagram of the first type of contraction in the vertex correction contributes to the correction of the parameter \mathcal{K} . It corresponds to the **Case 1-Case 4** below. For simplicity, we here show one of the related Feynman diagrams. By exchanging the symmetric indices, we can obtain the other three possible diagrams.




FIG. 3. The Feynman diagrams of the second type of contraction in the vertex correction contribute to the correction of other terms like $g|\phi_a|^4$. They correspond to the **Case 5-Case 12** below. For simplicity, we here show two of them. By exchanging the symmetric indices, we can have the other six possible diagrams.

- **Case 2:** The momenta $\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3'$ and \mathbf{k}_4' are fast momenta. $\mathbf{k}_1 = \mathbf{k}_4'$, $\omega_{n_1} = \omega_{n_4'}$; $\mathbf{k}_2 = \mathbf{k}_3'$ and $\omega_{n_2} = \omega_{n_3'}$.
- **Case 3:** The momenta $\mathbf{k}_1', \mathbf{k}_2', \mathbf{k}_3$ and \mathbf{k}_4 are fast momenta. $\mathbf{k}_1' = \mathbf{k}_3$, $\omega_{n_1'} = \omega_{n_3}$; $\mathbf{k}_2' = \mathbf{k}_4$ and $\omega_{n_2'} = \omega_{n_4}$.
- **Case 4:** The momenta $\mathbf{k}_1', \mathbf{k}_2', \mathbf{k}_3$ and \mathbf{k}_4 are fast momenta. $\mathbf{k}_1' = \mathbf{k}_4$, $\omega_{n_1'} = \omega_{n_4}$; $\mathbf{k}_2' = \mathbf{k}_3$ and $\omega_{n_2'} = \omega_{n_3}$.

These four cases correspond to the Feynman diagram in Fig. 2. Similarly, it can also be proved that these four cases are equal to one another by exchanging the indices. We directly take the **Case 1** as an example and multiply it by 4 for simplicity. By further considering the delta functions in vertex correction, the relationships between momenta and frequencies are given by: $\mathbf{k}_1 = \mathbf{k}_3', \mathbf{k}_2 = \mathbf{k}_4', \mathbf{k}_1 + \mathbf{k}_2 = \mathbf{k}_3 + \mathbf{k}_4 = \mathbf{k}_1' + \mathbf{k}_2' = \mathbf{k}_3' + \mathbf{k}_4'$; $\omega_{n_1} = \omega_{n_3'}, \omega_{n_2} = \omega_{n_4'}, \omega_{n_1} + \omega_{n_2} = \omega_{n_3} + \omega_{n_4} = \omega_{n_1'} + \omega_{n_2'} = \omega_{n_3'} + \omega_{n_4'}$. Hence, with the formula (20), we obtain that

$$\begin{aligned}
\langle S_{int}^2 \rangle_{>}^{(1)} &= \frac{1}{\beta^2} \sum_{a \neq b} \sum_{1,2,3,4} \sum_{1',2'} \mathcal{K}^2(k_2^a + k_1^b)^2 (k_3^a + k_4^b) (k_2^{a'} + k_1^{b'}) \frac{1}{i\omega_{n_1} - \xi_{1a}} \frac{1}{i\omega_{n_2} - \xi_{2b}} \phi_{1'a}^* \phi_{2'b}^* \phi_{3b} \phi_{4a} \times \\
&\quad \delta(1' + 2' - 3 - 4) \delta(1' + 2' - 2 - 1) \\
&= \frac{1}{\beta^2} \sum_{a \neq b} \sum_{2,3,4} \sum_{1',2'} \mathcal{K}^2(k_2^a - k_2^b + k_1^b + k_2^b)^2 (k_3^a + k_4^b) (k_2^{a'} + k_1^{b'}) \times \\
&\quad \frac{1}{i\omega_{n_1} + i\omega_{n_2} - i\omega_{n_2} - \xi_{1'+2'-2,a}} \cdot \frac{1}{i\omega_{n_2} - \xi_{2b}} \phi_{1'a}^* \phi_{2'b}^* \phi_{3b} \phi_{4a} \delta(1' + 2' - 3 - 4) \\
&= -\frac{1}{\beta^2} \sum_{a \neq b} \int \frac{d^2 k_2}{(2\pi)^2} \sum_{3,4} \sum_{1',2'} \mathcal{K}^2(k_2^a - k_2^b + k_1^b + k_2^b)^2 (k_3^a + k_4^b) (k_2^{a'} + k_1^{b'}) \\
&\quad \sum_{n_2} \frac{1}{i\omega_{n_2} - i\omega_{n_1} - i\omega_{n_2'} + \xi_{1'+2'-2,a}} \cdot \frac{1}{i\omega_{n_2} - \xi_{2b}} \phi_{1'a}^* \phi_{2'b}^* \phi_{3b} \phi_{4a} \delta(1' + 2' - 3 - 4), \tag{21}
\end{aligned}$$

where we introduce the notation: $\xi_{1'+2'-2,a} = \frac{\sum_{a'=1}^2 t_{aa'} (k_1^{a'} + k_2^{a'} - k_2^a)^2}{2} - \mu$. This expression contains a sum of Matsubara frequencies n_2 : $\sum_{n_2} \frac{1}{i\omega_{n_2} - i\omega_{n_1} - i\omega_{n_2'} + \xi_{1'+2'-2,a}} \cdot \frac{1}{i\omega_{n_2} - \xi_{2b}}$. By applying the formula: $\frac{1}{\beta} \sum_n \frac{1}{i\omega_n - \xi_1} \frac{1}{i\omega_n - \xi_2} = -\frac{f_B(\xi_1) - f_B(\xi_2)}{\xi_1 - \xi_2} = \frac{2\pi n}{\beta}$, for Matsubara frequencies for bosons: $\omega_n = \frac{2\pi n}{\beta}$, we can obtain the result with substitutions: $\xi_1 = i\omega_{n_1} + i\omega_{n_2'} - \xi_{1'+2'-2,a}$ and $\xi_2 = \xi_{2b}$. The sum of n_2 can be rewritten as:

$$\begin{aligned}
&\frac{1}{\beta} \sum_{n_2} \frac{1}{i\omega_{n_2} - i\omega_{n_1} - i\omega_{n_2'} + \xi_{1'+2'-2,a}} \cdot \frac{1}{i\omega_{n_2} - \xi_{2b}} \\
&= -\frac{f_B(\xi_{2b}) - f_B(-\xi_{1'+2'-2,a})}{\xi_{1'+2'-2,a} + \xi_{2b} - i\omega_{n_1} - i\omega_{n_2'}} \\
&= -\frac{f_B(\xi_{2b}) + f_B(\xi_{1'+2'-2,a}) + 1}{\xi_{1'+2'-2,a} + \xi_{2b} - i\omega_{n_1} - i\omega_{n_2'}}, \tag{22}
\end{aligned}$$

where we apply $\frac{f_B(i\omega_{n_1} + i\omega_{n_2} - \xi)}{\exp[2\pi i(n_1 + n_2) - \beta\xi] - 1} = \frac{1}{\exp[-\beta\xi] - 1} = f_B(-\xi) = \frac{1}{e^{-\beta\xi} - 1} = -1 - \frac{1}{e^{\beta\xi} - 1} = -1 - f_B(\xi)$. Replacing the sum in (21) with (22), $\langle S_{int}^2 \rangle_{>}^{(1)}$ can be rewritten as:

$$\begin{aligned}
\langle S_{int}^2 \rangle_{>}^{(1)} &= \frac{1}{\beta} \sum_{a \neq b} \sum_{3,4} \sum_{1',2'} \mathcal{K}^2(k_3^a + k_4^b) (k_2^a + k_1^b) \\
&\quad \cdot \phi_{1'a}^* \phi_{2'b}^* \phi_{3b} \phi_{4a} \delta(1' + 2' - 3 - 4) \\
&\quad \cdot \int \frac{d^2 k_2}{(2\pi)^2} \frac{f_B(\xi_{2b}) + f_B(\xi_{1'+2'-2,a}) + 1}{\xi_{1'+2'-2,a} + \xi_{2b} - i\omega_{n_1} - i\omega_{n_2'}} \\
&\quad \cdot (k_2^a - k_2^b + k_1^b + k_2^b)^2 \\
&\approx \frac{1}{\beta} \sum_{a \neq b} \sum_{3,4} \sum_{1',2'} \mathcal{K}^2(k_3^a + k_4^b) (k_2^{a'} + k_1^{b'}) \\
&\quad \cdot \phi_{1'a}^* \phi_{2'b}^* \phi_{3b} \phi_{4a} \delta(1' + 2' - 3 - 4) \\
&\quad \int \frac{d^2 k_2}{(2\pi)^2} \frac{f_B(\xi_{2b}) + f_B(\xi_{2a}) + 1}{\xi_{2a} + \xi_{2b} - i\omega_{n_1} - i\omega_{n_2'}} (k_2^a - k_2^b)^2,
\end{aligned}$$

where the momenta $\mathbf{k}_{1'}$ and $\mathbf{k}_{2'}$ can be ignored in the expression since $\phi_{1'a}$ and $\phi_{2'b}$ are both slow modes. The integrated region of the momentum \mathbf{k}_2 in $\langle S_{int}^2 \rangle_{>}^{(1)}$ is given by: $(\Lambda/s)^2 \leq \sum_{a(a \neq b)} (k_2^b)^2 + \frac{T_1}{T_0} (k_2^a)^2 \leq \Lambda^2$ since the momentum \mathbf{k}_2 is carried by the field $\phi_{2'b}^*$. With substitutions $\tilde{k}^b = k_2^b$, $\tilde{k}^a = \sqrt{\frac{T_1}{T_0}} k_2^a$, we limit our integrated region on $(\Lambda/s)^2 \leq \tilde{\mathbf{k}}^2 \leq \Lambda^2$. Therefore, the contribution from the first type of contraction is given by:

$$\langle S_{int}^2 \rangle_{>}^{(1)} = \frac{A}{\beta} \sum_{a \neq b} \sum_{1'2'34} \mathcal{K}^2(k_3^a + k_4^b) (k_2^a + k_1^b) \cdot \phi_{1'a}^* \phi_{2'b}^* \phi_{3b} \phi_{4a} \delta(1' + 2' - 3 - 4) dl, \tag{23}$$

where the parameter A takes the form of:

$$\begin{aligned}
A &= \frac{1}{dl} \int_{>} \frac{d^2 k_2}{(2\pi)^2} \frac{f_B(\xi_{2a}) + f_B(\xi_{2b}) + 1}{\xi_{2a} + \xi_{2b} - \omega_{1'} - \omega_{2'} - i0^+} (k_2^a - k_2^b)^2 \\
&\approx \frac{1}{dl} \int_{>} \frac{d^2 k_2}{(2\pi)^2} \frac{f_B(\xi_{2a}) + f_B(\xi_{2b}) + 1}{\xi_{2a} + \xi_{2b}} (k_2^a - k_2^b)^2, \tag{24}
\end{aligned}$$

where we use analytic continuation here and replace the imaginary frequencies with $\omega_{1'}$ and $\omega_{2'}$. It is necessarily assume that the frequencies and momenta on the external lines can be ignored compared with $\xi_{0\Lambda}$ and Λ . It is obvious that the expression (24) is positive. In other words, $A > 0$. More calculations on A are shown in Appendix B. By comparing the form of (23) with the bare interaction vertex, we find that it will correct the interaction parameter \mathcal{K} effectively. According to the effective action: $S_{eff} = S_0 + \langle S_{int} \rangle - \frac{1}{2} (\langle S_{int}^2 \rangle - \langle S_{int} \rangle^2)$, we have the form of the β -function of parameter \mathcal{K} :

$$\frac{d\mathcal{K}}{dl} = -\mathcal{K}^2 \times A - 2\mathcal{K}, \tag{25}$$

where the term $-2\mathcal{K}$ originates from the rescaling of slow momenta carried by the slow modes in (9) above. We

have the general form of $\mathcal{K}(l)$ and $\mathcal{T}_1(l)$:

$$\mathcal{K}(l) = \frac{2/A}{\frac{\mathcal{K}(0)+2/A}{\mathcal{K}(0)}e^{2l} - 1}, \quad (26)$$

$$\mathcal{T}_1(l) = \left[\mathcal{T}_1(0)^{3/2} + \sqrt{\mathcal{T}_0} \frac{3\Lambda^2}{4A\pi} f_B(\xi_{0\Lambda}) \cdot \ln\left(\frac{2}{-\mathcal{K}(0)Ae^{-2l} + (2 + \mathcal{K}(0)A)}\right) \right]^{2/3}, \quad (27)$$

with the separatrix:

$$\mathcal{T}_1(0) = (f_B(\xi_{0\Lambda})\sqrt{\mathcal{T}_0} \frac{3\Lambda^2}{4A\pi} \ln(\frac{2 + \mathcal{K}(0)A}{2}))^{2/3}. \quad (28)$$

The second type of contraction $\langle S_{int}^2 \rangle_{>}^{(2)}$ contains over-all eight cases presented below.

- **Case 5:** The momenta \mathbf{k}_1 , \mathbf{k}_3 , \mathbf{k}_2' and \mathbf{k}_4' are fast momenta. $\mathbf{k}_1 = \mathbf{k}_4'$, $\omega_{n_1} = \omega_{n_4'}$; $\mathbf{k}_3 = \mathbf{k}_2'$ and $\omega_{n_3} = \omega_{n_2'}$.
- **Case 6:** The momenta \mathbf{k}_1 , \mathbf{k}_4 , \mathbf{k}_1' and \mathbf{k}_4' are fast momenta. $\mathbf{k}_1 = \mathbf{k}_4'$, $\omega_{n_1} = \omega_{n_4'}$; $\mathbf{k}_4 = \mathbf{k}_1'$ and $\omega_{n_4} = \omega_{n_1'}$.
- **Case 7:** The momenta \mathbf{k}_1 , \mathbf{k}_3 , \mathbf{k}_1' and \mathbf{k}_3' are fast momenta. $\mathbf{k}_1 = \mathbf{k}_3'$, $\omega_{n_1} = \omega_{n_3'}$; $\mathbf{k}_3 = \mathbf{k}_1'$ and $\omega_{n_3} = \omega_{n_1'}$.
- **Case 8:** The momenta \mathbf{k}_1 , \mathbf{k}_4 , \mathbf{k}_2' and \mathbf{k}_3' are fast momenta. $\mathbf{k}_1 = \mathbf{k}_3'$, $\omega_{n_1} = \omega_{n_3'}$; $\mathbf{k}_4 = \mathbf{k}_2'$ and $\omega_{n_4} = \omega_{n_2'}$.
- **Case 9:** The momenta \mathbf{k}_2 , \mathbf{k}_3 , \mathbf{k}_2' and \mathbf{k}_3' are fast momenta. $\mathbf{k}_2 = \mathbf{k}_3'$, $\omega_{n_2} = \omega_{n_3'}$; $\mathbf{k}_3 = \mathbf{k}_2'$ and $\omega_{n_3} = \omega_{n_2'}$.
- **Case 10:** The momenta \mathbf{k}_2 , \mathbf{k}_4 , \mathbf{k}_1' and \mathbf{k}_3' are fast momenta. $\mathbf{k}_2 = \mathbf{k}_3'$, $\omega_{n_2} = \omega_{n_3'}$; $\mathbf{k}_4 = \mathbf{k}_1'$ and $\omega_{n_4} = \omega_{n_1'}$.
- **Case 11:** The momenta \mathbf{k}_2 , \mathbf{k}_3 , \mathbf{k}_1' and \mathbf{k}_4' are fast momenta. $\mathbf{k}_2 = \mathbf{k}_4'$, $\omega_{n_2} = \omega_{n_4'}$; $\mathbf{k}_3 = \mathbf{k}_1'$ and $\omega_{n_3} = \omega_{n_1'}$.
- **Case 12:** The momenta \mathbf{k}_2 , \mathbf{k}_4 , \mathbf{k}_2' and \mathbf{k}_3' are fast momenta. $\mathbf{k}_2 = \mathbf{k}_3'$, $\omega_{n_2} = \omega_{n_3'}$; $\mathbf{k}_4 = \mathbf{k}_2'$ and $\omega_{n_4} = \omega_{n_2'}$.

The Feynman diagrams related are presented in Fig. 3. After integrating the fast momenta, all of these terms will be turned into terms with constant coefficients. Therefore, integration from the two connected diagrams will be explained as corrections of other possible terms such as $g|\phi_a|^4$. We do not care about this parameter for it does no contribution to the parameters \mathcal{K} and \mathcal{T}_1 . For simplicity, we do not present more details here.




FIG. 4. RG flow for $\mathcal{T}_1'(l) = \mathcal{T}_1(l) \left(f_B(\xi_{0\Lambda}) \sqrt{\mathcal{T}_0} \frac{3\Lambda^2}{4A\pi} \right)^{-2/3}$ and $\mathcal{K}'(l) = A\mathcal{K}(l)/2$. Here the red line is a separatrix below which higher-rank symmetry emerges. The parameters $(\mathcal{T}_1'(0), \mathcal{K}'(0))$ of these lines are (from above to below): $(2.57 \times 10^{-3}, 10^{-4})$, $(2.43 \times 10^{-3}, 10^{-4})$, $(2.30 \times 10^{-3}, 10^{-4})$, $(2.16 \times 10^{-3}, 10^{-4})$, $(2.01 \times 10^{-3}, 10^{-4})$, $(1.86 \times 10^{-3}, 10^{-4})$, and $(1.70 \times 10^{-3}, 10^{-4})$.

D. Global phase diagram

In summary, we can see the first-order corrections only correct the kinetic terms on directions other than the a -th one of field configurations ϕ_a [see definition below Eq. (7)]. In this way, the kinetic energy along the a -th axis will not be influenced by the contraction. We can directly prove that the higher-order correction will still contribute nothing to the parameter \mathcal{T}_0 . Hence, we have the β -function for parameter \mathcal{T}_1 corresponding to Fig. 1, given by Eq. (19). We can safely come to the conclusion that if $\mathcal{K} > 0$, the parameter t_1 will reduce in the RG flow. This requires the parameter \mathcal{K} to be positive. Only with positive \mathcal{K} can parameter \mathcal{T}_1 flow to zero in the RG analysis. Nevertheless, at this step, it is insufficient to tell whether the parameter \mathcal{T}_1 is irrelevant here since the elements in the matrix \mathcal{K} will also flow to zero. What we need is further calculation on vertex correction, which has been given in Eq. (25).

A numerical result of beta functions (19) and (25) is shown in Fig. 4, where the red line is the separatrix (28). In the region below the line, all $\mathcal{T}_1(0)$ will flow to zero in RG flow, indicating the emergence of higher-rank symmetry in low-energy physics since those terms violating the higher-rank symmetry will vanish in the low-energy physics. In addition, after comparing the parameters $\mathcal{T}_1(0)$ and $\mathcal{K}(0)\Lambda^2$ near the separatrix, we have $\mathcal{T}_1(0) \ll \mathcal{K}(0)\Lambda^2 \ll \mathcal{T}_0$ with the approximation $\frac{\xi_{0\Lambda}}{2k_B T} \gg 1$, which can be realized in cold-atom systems. This tells us our calculation is consistent with our initial assumption above.

From the phase diagram, we can safely come to the conclusion that the higher-rank symmetry will emerge after integrating the high-energy modes since off-diagonal kinetic parameters are just perturbations to the system. However, it should be noted that our RG analysis is only

valid near the initial point $l = 0$. We here also do not take higher-order terms into account. Therefore, our calculation and results may be invalidated when $l \rightarrow \infty$. The reason we do not just finish our RG analysis on the axis: $\mathcal{T}_1(0) = 0$ is we only consider the lower-order corrections. If we start with an initial point on the horizontal axis: $\mathcal{T}_1(0) = 0$, we have $S_2 \rightarrow \infty$ in $\frac{d\mathcal{T}_1}{dl}$, which requires more higher-order terms since they contain expressions with higher order of S_2 . We have to take all of them into account for accuracy, which is beyond the current perturbation calculations. Although our calculation cannot take all the higher-order corrections into account, it also exhibits the tendency of RG flow giving the accurate prediction near the region with $l = 0$. Alternatively, the exact flat “band” along one direction for a boson, in analogy to bosonic Landau level, leads to strong correlation effect even though interaction is not large. We also comment that the perturbation calculations fail completely if we instead study models [68] with conserved dipole moments, where bosons are fractons that are immobile towards all directions. In such models, the Hamiltonian is intrinsically non-Gaussian with no kinetic terms at all. But analytic difficulty becomes much smaller when fracton condensation is considered as what Ref. [68] did.

Above discussion is all about the β -functions of parameters of 2-component bosons in two-dimensional space. Similarly, we can extend the case to the d -dimensional space with $d > 2$. We here give the forms of β -functions of d -component bosonic systems for reference. The β -functions of t_{ab} ($a \neq b$) and K_{ab} ($a \neq b$) is given by: $\frac{dt_{ab}}{dl} = -S_d K_{ab} f_B(\xi_{0\Lambda}) \frac{\Lambda^d}{(2\pi)^d}$, $\frac{dK_{ab}}{dl} = -A_d K_{ab}^2 - dK_{ab}$, where S_d represents the volume of d -dimensional ellipsoid: $\sum_{b=1}^d \frac{t_{ab}}{t_a}(k^b)^2 = 1$ and $A_d = \frac{1}{2\beta l} \int_{P_d} \frac{d^d k}{(2\pi)^d} \frac{f_B(\xi_{ka}) + f_B(\xi_{kb}) + 1}{\xi_{ka} + \xi_{kb}} (k^a - k^b)^2$ with integrated domain $P_d: (\Lambda e^{-l})^2 \leq \sum_{b=1}^d \frac{t_{ab}}{t_a}(k^b)^2 \leq \Lambda^2$. They give the same type of RG flow phase diagram as Fig. 4.

IV. SUMMARY AND OUTLOOK

In this paper, we study how higher-rank symmetry (5) and angular momentum conservation (6) emerge at low energies through a RG analysis. In other words, we identify them as emergent phenomena rather than strict properties of microscopic models. A phase diagram is given in Fig. 4, in which a wide parameter region is found to support emergent phenomena. Despite the limitation of one-loop perturbation techniques, we argue that emergence occurs in the deep infrared regime. On the other hand, by regarding higher-rank symmetry as emergent symmetry, our work opens a door to new way of thinking on realization of such unconventional symmetry and higher moment conservation in more realistic models, e.g., in simple frustrated spin models near symmetric points. Recently, some higher-moment conserving 1D spin systems have been found to support anomalously slow, subdiffusive late-time transport [74, 78]. Thus, it is interesting

to ask how *emergent* conservation of angular moments affect the late-time transport. Besides, the Hamiltonian can be reformulated on a square lattice. In an optical lattice of cold-atom experiments, one may choose lattice constant $l = 700nm$, $\Lambda \sim \frac{1}{l} = 1.43 \times 10^7 m^{-1}$, $m_0 = \frac{1}{t_0} = 8.22 \times 10^{-34} kg$, $T = 10K$. The condition $\frac{t_0 \Lambda^2 - 2\mu}{2k_B T} \sim 10^3$ where the chemical potential is negative indicates the possibility for simulating the 2-component bosons on the optical lattice. For accuracy and extension, it will be interesting to further consider the correction from higher-order term with more loops in the Feynman diagram. Finally, it will be interesting to study symmetry-protected topological phases (SPTs) with such emergent higher-rank symmetry.

ACKNOWLEDGEMENTS

Discussions with Ruizhi Liu and Yixin Xu are acknowledged. This work of both authors was done in Guangzhou South Campus of Sun Yat-sen University (SYSU) with full financial support from SYSU talent plan, Guangdong Basic and Applied Basic Research Foundation under Grant No. 2020B1515120100, and National Natural Science Foundation of China Grant (No. 11847608 & No. 12074438).

Appendix A: On how the angular momentum conservation affects single particle motion

In the main text, we introduce the following conserved quantities ($a, b = 1, 2, \dots, d; a < b$) in d -dimensional space:

$$\hat{Q}^a = \int d^d x \hat{\rho}^a, \quad \hat{Q}^{ab} = \int d^d x (\hat{\rho}^a x^b - \hat{\rho}^b x^a). \quad (A1)$$

For each a , the conservation of \hat{Q}^a requires that all a -th component bosons are always in the d -dimensional space. The conserved quantities \hat{Q}^{ab} leads to the mobility restriction: a single a -th component boson is only allowed to move along a -th directions. To be more specific, we consider $d = 2$ and the following three quantities are conserved:

$$\hat{Q}^1 = \int d^d x \hat{\rho}^1, \quad \hat{Q}^2 = \int d^d x \hat{\rho}^2, \quad \hat{Q}^{12} = \int d^d x (\hat{\rho}^1 x^2 - \hat{\rho}^2 x^1). \quad (A2)$$

\hat{Q}^1 and \hat{Q}^2 are the usual conserved charge (particle number) of bosons of the 1st and 2nd components, respectively. The conserved quantity \hat{Q}^{12} is the total angular moments formed by bosons. Suppose $\hat{Q}^1 = N_1$ and $\hat{Q}^2 = N_2$. We can use Dirac function to express $\hat{\rho}^1$ and $\hat{\rho}^2$ in the eigenbasis:

$$\rho^1(\mathbf{x}) = \sum_i^{N_1} \delta(\mathbf{x} - \mathbf{x}_i), \quad \rho^2(\mathbf{x}) = \sum_j^{N_2} \delta(\mathbf{x} - \mathbf{x}_j). \quad (A3)$$

It means that bosons of the 1st (2nd) component are located in \mathbf{x}_i (\mathbf{x}_j) with $i = 1, 2, 3, \dots, N_1$ ($j = 1, 2, 3, \dots, N_2$). Then, \mathcal{Q}^{12} are reduced to:

$$\mathcal{Q}^{12} = \sum_i^{N_1} x_i^2 - \sum_j^{N_2} x_j^1. \quad (\text{A4})$$

From this expression, one can conclude that, if we move a 1st component boson, in order to keep \mathcal{Q}^{12} invariant, the boson is only allowed to be movable in the 1st direction such that its coordinate x^2 is unchanged. Of course, we can *collectively* move bosons of both components, such that the change in $\sum_i^{N_1} x_i^2$ can be canceled out by the change in $\sum_j^{N_2} x_j^1$ such that \mathcal{Q}^{12} is unaltered. This scenario is beyond the single particle movement and gives nontrivial effects when inter-component interaction (K -term) is involved.

For simplicity, we will approximately use $\mathcal{T}_1(0)$ to replace \mathcal{T}_1 below for convenience. Besides, we have $k_2^b = \Lambda \sin \theta$ and $k_2^a = \Lambda \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \cos \theta$ in the integrated domain. Hence, the kinetic energy ξ_{2a} and ξ_{2b} can be approximately considered as $\xi_{1\Lambda} = \xi_{0\Lambda} (\frac{\mathcal{T}_0}{\mathcal{T}_1} \cos^2 \theta + \frac{\mathcal{T}_1}{\mathcal{T}_0} \sin^2 \theta)$ and $\xi_{0\Lambda}$ separately if we assume the chemical potential is small enough: $|\mu| \ll \frac{t_0 \Lambda^2}{2}$. In this way, the parameter A can be rewritten as:

$$\begin{aligned} A &\approx \frac{1}{2dl} \int_0^{2\pi} \frac{d\theta}{(2\pi)^2} (\Lambda^2 - (\frac{\Lambda}{s})^2) \frac{f_B(\xi_{1\Lambda}) + f_B(\xi_{0\Lambda}) + 1}{\xi_{1\Lambda} + \xi_{0\Lambda}} (\Lambda \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \cos \theta - \Lambda \sin \theta)^2 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \\ &= \Lambda^4 \int_0^{2\pi} \frac{d\theta}{(2\pi)^2} \frac{f_B(\xi_{1\Lambda}) + f_B(\xi_{0\Lambda}) + 1}{\xi_{1\Lambda} + \xi_{0\Lambda}} (\sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \cos \theta - \sin \theta)^2 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \\ &= \Lambda^4 \int_0^{2\pi} \frac{d\theta}{(2\pi)^2} \frac{f_B(\xi_{1\Lambda}) + f_B(\xi_{0\Lambda}) + 1}{\xi_{1\Lambda} + \xi_{0\Lambda}} (\sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \cos \theta - \sin \theta)^2 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \\ &\approx \Lambda^4 \int_0^{2\pi} \frac{d\theta}{(2\pi)^2} \frac{1}{(\xi_{0\Lambda} + \xi_{1\Lambda}) (e^{\xi_{0\Lambda} (\frac{\mathcal{T}_0}{\mathcal{T}_1} \cos^2 \theta + \frac{\mathcal{T}_1}{\mathcal{T}_0} \sin^2 \theta) / k_B T} - 1)} \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \\ &\quad + \Lambda^4 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \int_0^{2\pi} \frac{d\theta}{(2\pi)^2} \frac{f_B(\xi_{0\Lambda}) + 1}{\xi_{1\Lambda} + \xi_{0\Lambda}} (\sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \cos \theta - \sin \theta)^2 \\ &\approx \frac{4\Lambda^2}{\mathcal{T}_0} \int_{\frac{\pi}{2} - \mathcal{T}'_1}^{\frac{\pi}{2} + \mathcal{T}'_1} \frac{d\theta}{(2\pi)^2} \frac{2k_B T}{\mathcal{T}_0 \Lambda^2 \mathcal{T}'_1 \sin^2 \theta} \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} + \Lambda^4 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \int_0^{2\pi} \frac{d\theta}{(2\pi)^2} \frac{(\sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \cos \theta - \sin \theta)^2 (1 + f_B(\xi_{0\Lambda}))}{\xi_{0\Lambda} (1 + \frac{\mathcal{T}_0}{\mathcal{T}_1} \cos^2 \theta + \frac{\mathcal{T}_1}{\mathcal{T}_0} \sin^2 \theta)}, \quad (\text{B2}) \end{aligned}$$

with $\mathcal{T}'_1 = \frac{\mathcal{T}_1}{\mathcal{T}_0} \ll 1$. Here we approximately consider that $\frac{\mathcal{T}_1 \Lambda^2}{2k_B T} \ll 1$. This expression contains two integrals. The first one includes a trick. Since $\mathcal{T}'_1 \ll 1$, the first integral becomes nontrivial only around the two points: $\theta = \frac{\pi}{2}$ and $\theta = \frac{3\pi}{2}$, which give the same value. Therefore, we change our integrated region as: $\int_0^{2\pi} \rightarrow 2 \int_{\frac{\pi}{2} - \mathcal{T}'_1}^{\frac{\pi}{2} + \mathcal{T}'_1}$. In addition, we have $\xi_{1\Lambda} = \frac{\mathcal{T}_1 \Lambda^2}{2} \ll \xi_{0\Lambda}$ at $\theta = \frac{\pi}{2}$ and $\theta = \frac{3\pi}{2}$. In this way, the first integral given by: $\frac{4\Lambda^2}{\mathcal{T}_0} \int_{\frac{\pi}{2} - \mathcal{T}'_1}^{\frac{\pi}{2} + \mathcal{T}'_1} \frac{d\theta}{(2\pi)^2} \frac{2k_B T}{\mathcal{T}_0 \Lambda^2 \mathcal{T}'_1 \sin^2 \theta} \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}}$ can

Appendix B: Details of the parameter A

Here we focus on the specific value of the parameter A by taking some approximations here. By referencing the definition of $\tilde{\mathbf{k}}$ above, we have:

$$\begin{aligned} A &= \frac{1}{dl} \int_{>} \frac{d^2 \tilde{\mathbf{k}}}{(2\pi)^2} \frac{f_B(\xi_{2a}) + f_B(\xi_{2b}) + 1}{\xi_{2a} + \xi_{2b}} (k_2^a - k_2^b)^2 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \\ &= \frac{1}{dl} \int_{>} \frac{|\tilde{\mathbf{k}}| d|\tilde{\mathbf{k}}| d\theta}{(2\pi)^2} \frac{f_B(\xi_{2a}) + f_B(\xi_{2b}) + 1}{\xi_{2a} + \xi_{2b}} (k_2^a - k_2^b)^2 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \\ &= \frac{1}{dl} \int_{>} \frac{d|\tilde{\mathbf{k}}|^2 d\theta}{2(2\pi)^2} \frac{f_B(\xi_{2a}) + f_B(\xi_{2b}) + 1}{\xi_{2a} + \xi_{2b}} (k_2^a - k_2^b)^2 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}}. \quad (\text{B1}) \end{aligned}$$

be figured out to be: $\frac{4\Lambda^2}{\mathcal{T}_0} \frac{4k_B T}{(2\pi)^2 \mathcal{T}_0 \Lambda^2} \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}}$. The second one $\Lambda^4 \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \int_0^{2\pi} \frac{d\theta}{(2\pi)^2} \frac{(\sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1}} \cos \theta - \sin \theta)^2 (1 + f_B(\xi_{0\Lambda}))}{\xi_{0\Lambda} (1 + \frac{\mathcal{T}_0}{\mathcal{T}_1} \cos^2 \theta + \frac{\mathcal{T}_1}{\mathcal{T}_0} \sin^2 \theta)}$ can be proved to be $\frac{\Lambda^2}{\pi \mathcal{T}_0} \frac{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}}{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)} + 1} (1 + f_B(\xi_{0\Lambda})) \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}}$. Combined

with these two results, the expression of A is given by:

$$A = \frac{\Lambda^2}{\pi\mathcal{T}_0} \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \left(\frac{4k_B T}{\pi\mathcal{T}_0\Lambda^2} + \frac{\frac{\mathcal{T}_0}{\mathcal{T}_1}}{\frac{\mathcal{T}_0}{\mathcal{T}_1} + 1} (1 + f_B(\xi_{0\Lambda})) \right) \\ \approx \frac{\Lambda^2}{\pi\mathcal{T}_0} \sqrt{\frac{\mathcal{T}_0}{\mathcal{T}_1(0)}} \left(\frac{4k_B T}{\pi\mathcal{T}_0\Lambda^2} + 1 + f_B(\xi_{0\Lambda}) \right) > 0, \quad (\text{B3})$$

-
- [1] E. Noether, “Invariante variationsprobleme,” *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse* **1918**, 235–257 (1918).
- [2] David J Griffiths, *Introduction to electrodynamics; 4th ed.* (Pearson, Boston, MA, 2013) re-published by Cambridge University Press in 2017.
- [3] Rahul M. Nandkishore and Michael Hermele, “Fractons,” *Annual Review of Condensed Matter Physics* **10**, 295–313 (2019), arXiv:1803.11196 [cond-mat.str-el].
- [4] Michael Pretko, Xie Chen, and Yizhi You, “Fracton phases of matter,” *International Journal of Modern Physics A* **35**, 2030003 (2020), arXiv:2001.01722 [cond-mat.str-el].
- [5] Michael Pretko, “The fracton gauge principle,” *Phys. Rev. B* **98**, 115134 (2018).
- [6] Michael Pretko, “Generalized electromagnetism of subdimensional particles: A spin liquid story,” *Phys. Rev. B* **96**, 035119 (2017).
- [7] Michael Pretko, “Subdimensional particle structure of higher rank $u(1)$ spin liquids,” *Phys. Rev. B* **95**, 115139 (2017).
- [8] Andrey Gromov, “Towards classification of fracton phases: The multipole algebra,” *Phys. Rev. X* **9**, 031035 (2019).
- [9] Nathan Seiberg and Shu-Heng Shao, “Exotic $U(1)$ Symmetries, Duality, and Fractons in 3+1-Dimensional Quantum Field Theory,” *SciPost Phys.* **9**, 46 (2020).
- [10] Sagar Vijay, Jeongwan Haah, and Liang Fu, “A new kind of topological quantum order: A dimensional hierarchy of quasiparticles built from stationary excitations,” *Phys. Rev. B* **92**, 235136 (2015).
- [11] Sagar Vijay, Jeongwan Haah, and Liang Fu, “Fracton topological order, generalized lattice gauge theory, and duality,” *Phys. Rev. B* **94**, 235157 (2016).
- [12] Abhinav Prem, Jeongwan Haah, and Rahul Nandkishore, “Glassy quantum dynamics in translation invariant fracton models,” *Phys. Rev. B* **95**, 155133 (2017).
- [13] Claudio Chamon, “Quantum glassiness in strongly correlated clean systems: An example of topological overprotection,” *Phys. Rev. Lett.* **94**, 040402 (2005).
- [14] Wilbur Shirley, Kevin Slagle, and Xie Chen, “Foliated fracton order from gauging subsystem symmetries,” *SciPost Phys.* **6**, 41 (2019).
- [15] Han Ma, Ethan Lake, Xie Chen, and Michael Hermele, “Fracton topological order via coupled layers,” *Phys. Rev. B* **95**, 245126 (2017).
- [16] Jeongwan Haah, “Local stabilizer codes in three dimensions without string logical operators,” *Phys. Rev. A* **83**, 042330 (2011).
- [17] Daniel Bulmash and Maissam Barkeshli, “Gauging fractons: Immobile non-Abelian quasiparticles, fractals, and position-dependent degeneracies,” *Phys. Rev. B* **100**, 155146 (2019), arXiv:1905.05771 [cond-mat.str-el].
- [18] Abhinav Prem and Dominic Williamson, “Gauging permutation symmetries as a route to non-Abelian fractons,” *SciPost Physics* **7**, 068 (2019), arXiv:1905.06309 [cond-mat.str-el].
- [19] D. Bulmash and M. Barkeshli, “Generalized $U(1)$ Gauge Field Theories and Fractal Dynamics,” arXiv e-prints (2018), arXiv:1806.01855 [cond-mat.str-el].
- [20] Kevin T. Tian, Eric Samperton, and Zhenghan Wang, “Haah codes on general three-manifolds,” *Annals of Physics* **412**, 168014 (2020), arXiv:1812.02101 [quant-ph].
- [21] Yizhi You, Daniel Litinski, and Felix von Oppen, “Higher-order topological superconductors as generators of quantum codes,” *Phys. Rev. B* **100**, 054513 (2019), arXiv:1810.10556 [cond-mat.str-el].
- [22] Han Ma, Michael Hermele, and Xie Chen, “Fracton topological order from the higgs and partial-confinement mechanisms of rank-two gauge theory,” *Phys. Rev. B* **98**, 035111 (2018).
- [23] Kevin Slagle and Yong Baek Kim, “Fracton topological order from nearest-neighbor two-spin interactions and dualities,” *Phys. Rev. B* **96**, 165106 (2017).
- [24] Gábor B. Halász, Timothy H. Hsieh, and Leon Balents, “Fracton topological phases from strongly coupled spin chains,” *Phys. Rev. Lett.* **119**, 257202 (2017).
- [25] Kevin T. Tian and Zhenghan Wang, “Generalized Haah Codes and Fracton Models,” arXiv e-prints, arXiv:1902.04543 (2019), arXiv:1902.04543 [quant-ph].
- [26] Wilbur Shirley, Kevin Slagle, and Xie Chen, “Foliated fracton order from gauging subsystem symmetries,” *SciPost Phys.* **6**, 41 (2019).
- [27] Wilbur Shirley, Kevin Slagle, and Xie Chen, “Fractional excitations in foliated fracton phases,” *Annals of Physics* **410**, 167922 (2019), arXiv:1806.08625 [cond-mat.str-el].
- [28] Kevin Slagle, David Aasen, and Dominic Williamson, “Foliated Field Theory and String-Membrane-Net Condensation Picture of Fracton Order,” *SciPost Phys.* **6**, 43 (2019).
- [29] Wilbur Shirley, Kevin Slagle, Zhenghan Wang, and Xie Chen, “Fracton models on general three-dimensional manifolds,” *Phys. Rev. X* **8**, 031051 (2018).
- [30] Abhinav Prem, Sheng-Jie Huang, Hao Song, and Michael Hermele, “Cage-net fracton models,” *Phys. Rev. X* **9**, 021010 (2019).
- [31] Shriya Pai, Michael Pretko, and Rahul M. Nandkishore, “Localization in fractonic random circuits,” *Phys. Rev. X* **9**, 021003 (2019).
- [32] Shriya Pai and Michael Pretko, “Dynamical Scar States

- in Driven Fracton Systems,” *Phys. Rev. Lett.* **123**, 136401 (2019), [arXiv:1903.06173 \[cond-mat.stat-mech\]](#).
- [33] Pablo Sala, Tibor Rakovszky, Ruben Verresen, Michael Knap, and Frank Pollmann, “Ergodicity Breaking Arising from Hilbert Space Fragmentation in Dipole-Conserving Hamiltonians,” *Physical Review X* **10**, 011047 (2020), [arXiv:1904.04266 \[cond-mat.str-el\]](#).
- [34] Ajesh Kumar and Andrew C. Potter, “Symmetry-enforced fractonicity and two-dimensional quantum crystal melting,” *Phys. Rev. B* **100**, 045119 (2019).
- [35] Leo Radzihovsky and Michael Hermele, “Fractons from Vector Gauge Theory,” *Phys. Rev. Lett.* **124**, 050402 (2020), [arXiv:1905.06951 \[cond-mat.str-el\]](#).
- [36] Arpit Dua, Isaac H. Kim, Meng Cheng, and Dominic J. Williamson, “Sorting topological stabilizer models in three dimensions,” *Phys. Rev. B* **100**, 155137 (2019), [arXiv:1908.08049 \[quant-ph\]](#).
- [37] Andrey Gromov, “Chiral topological elasticity and fracton order,” *Phys. Rev. Lett.* **122**, 076403 (2019).
- [38] Jeongwan Haah, *Lattice quantum codes and exotic topological phases of matter*, Ph.D. thesis, California Institute of Technology (2013).
- [39] Yizhi You, Trithep Devakul, S. L. Sondhi, and F. J. Burnell, “Fractonic Chern-Simons and BF theories,” *Physical Review Research* **2**, 023249 (2020), [arXiv:1904.11530 \[cond-mat.str-el\]](#).
- [40] John Sous and Michael Pretko, “Fractons from polarons and hole-doped antiferromagnets: Microscopic realizations,” arXiv e-prints, [arXiv:1904.08424 \(2019\)](#), [arXiv:1904.08424 \[cond-mat.str-el\]](#).
- [41] Vedika Khemani, Michael Hermele, and Rahul Nandkishore, “Localization from hilbert space shattering: From theory to physical realizations,” *Phys. Rev. B* **101**, 174204 (2020).
- [42] Juven Wang and Kai Xu, “Higher-Rank Tensor Field Theory of Non-Abelian Fracton and Embeddon,” arXiv e-prints, [arXiv:1909.13879 \(2019\)](#), [arXiv:1909.13879 \[hep-th\]](#).
- [43] Juven Wang and Shing-Tung Yau, “Non-Abelian gauged fracton matter field theory: Sigma models, superfluids, and vortices,” *Physical Review Research* **2**, 043219 (2020), [arXiv:1912.13485 \[cond-mat.str-el\]](#).
- [44] Shriya Pai and Michael Pretko, “Fractonic line excitations: An inroad from three-dimensional elasticity theory,” *Phys. Rev. B* **97**, 235102 (2018).
- [45] Michael Pretko and Rahul M. Nandkishore, “Localization of extended quantum objects,” *Phys. Rev. B* **98**, 134301 (2018).
- [46] Dominic J. Williamson, Zhen Bi, and Meng Cheng, “Fractonic matter in symmetry-enriched $u(1)$ gauge theory,” *Phys. Rev. B* **100**, 125150 (2019).
- [47] Arpit Dua, Dominic J. Williamson, Jeongwan Haah, and Meng Cheng, “Compactifying fracton stabilizer models,” *Phys. Rev. B* **99**, 245135 (2019).
- [48] Bowen Shi and Yuan-Ming Lu, “Deciphering the nonlocal entanglement entropy of fracton topological orders,” *Phys. Rev. B* **97**, 144106 (2018).
- [49] Hao Song, Abhinav Prem, Sheng-Jie Huang, and M. A. Martin-Delgado, “Twisted fracton models in three dimensions,” *Phys. Rev. B* **99**, 155118 (2019).
- [50] Han Ma and Michael Pretko, “Higher-rank deconfined quantum criticality at the lifshitz transition and the exciton bose condensate,” *Phys. Rev. B* **98**, 125105 (2018).
- [51] Juven Wang, Kai Xu, and Shing-Tung Yau, “Higher-Rank Non-Abelian Tensor Field Theory: Higher-Moment or Subdimensional Polynomial Global Symmetry, Algebraic Variety, Noether’s Theorem, and Gauge,” arXiv e-prints, [arXiv:1911.01804 \(2019\)](#), [arXiv:1911.01804 \[hep-th\]](#).
- [52] Kevin Slagle, “Foliated Quantum Field Theory of Fracton Order,” arXiv e-prints, [arXiv:2008.03852 \(2020\)](#), [arXiv:2008.03852 \[hep-th\]](#).
- [53] Dominic J. Williamson and Trithep Devakul, “Type-II fractons from coupled spin chains and layers,” arXiv e-prints, [arXiv:2007.07894 \(2020\)](#), [arXiv:2007.07894 \[cond-mat.str-el\]](#).
- [54] Pranay Gorantla, Ho Tat Lam, Nathan Seiberg, and Shu-Heng Shao, “More Exotic Field Theories in 3+1 Dimensions,” arXiv e-prints, [arXiv:2007.04904 \(2020\)](#), [arXiv:2007.04904 \[cond-mat.str-el\]](#).
- [55] Dung Xuan Nguyen, Andrey Gromov, and Sergej Moroz, “Fracton-elasticity duality of two-dimensional superfluid vortex crystals: defect interactions and quantum melting,” arXiv e-prints, [arXiv:2005.12317 \(2020\)](#), [arXiv:2005.12317 \[cond-mat.quant-gas\]](#).
- [56] Michael Pretko, S. A. Parameswaran, and Michael Hermele, “Odd Fracton Theories, Proximate Orders, and Parton Constructions,” arXiv e-prints, [arXiv:2004.14393 \(2020\)](#), [arXiv:2004.14393 \[cond-mat.str-el\]](#).
- [57] Dominic J. Williamson and Meng Cheng, “Designer non-Abelian fractons from topological layers,” arXiv e-prints, [arXiv:2004.07251 \(2020\)](#), [arXiv:2004.07251 \[cond-mat.str-el\]](#).
- [58] Nathan Seiberg and Shu-Heng Shao, “Exotic \mathbb{Z}_N Symmetries, Duality, and Fractons in 3+1-Dimensional Quantum Field Theory,” arXiv e-prints, [arXiv:2004.06115 \(2020\)](#), [arXiv:2004.06115 \[cond-mat.str-el\]](#).
- [59] David T. Stephen, José Garre-Rubio, Arpit Dua, and Dominic J. Williamson, “Subsystem symmetry enriched topological order in three dimensions,” *Physical Review Research* **2**, 033331 (2020), [arXiv:2004.04181 \[cond-mat.str-el\]](#).
- [60] Andrey Gromov, Andrew Lucas, and Rahul M. Nandkishore, “Fracton hydrodynamics,” *Physical Review Research* **2**, 033124 (2020), [arXiv:2003.09429 \[cond-mat.str-el\]](#).
- [61] Juven Wang, “Non-Liquid Cellular States,” arXiv e-prints, [arXiv:2002.12932 \(2020\)](#), [arXiv:2002.12932 \[cond-mat.str-el\]](#).
- [62] Wilbur Shirley, “Fractonic order and emergent fermionic gauge theory,” arXiv e-prints, [arXiv:2002.12026 \(2020\)](#), [arXiv:2002.12026 \[cond-mat.str-el\]](#).
- [63] David Aasen, Daniel Bulmash, Abhinav Prem, Kevin Slagle, and Dominic J. Williamson, “Topological Defect Networks for Fractons of all Types,” arXiv e-prints, [arXiv:2002.05166 \(2020\)](#), [arXiv:2002.05166 \[cond-mat.str-el\]](#).
- [64] Xiao-Gang Wen, “A systematic construction of gapped non-liquid states,” arXiv e-prints, [arXiv:2002.02433 \(2020\)](#), [arXiv:2002.02433 \[cond-mat.str-el\]](#).
- [65] Ting Fung Jeffrey Poon and Xiong-Jun Liu, “Quantum phase transition of fracton topological orders,” arXiv e-prints, [arXiv:2001.05937 \(2020\)](#), [arXiv:2001.05937 \[cond-mat.str-el\]](#).
- [66] Meng-Yuan Li and Peng Ye, “Fracton physics of spatially extended excitations,” *Phys. Rev. B* **101**, 245134 (2020).
- [67] Meng-Yuan Li and Peng Ye, “Fracton physics of spatially extended excitations. II. Polynomial ground

- state degeneracy of exactly solvable models,” (2021), [arXiv:2104.05735 \[cond-mat.str-el\]](#).
- [68] Jian-Keng Yuan, Shuai A. Chen, and Peng Ye, “Fractional superfluids,” *Physical Review Research* **2**, 023267 (2020), [arXiv:1911.02876 \[cond-mat.str-el\]](#).
- [69] Shuai A. Chen, Jian-Keng Yuan, and Peng Ye, “Fractional superfluids. ii. condensing subdimensional particles,” *Phys. Rev. Research* **3**, 013226 (2021).
- [70] Sanjay Moudgalya, B. Andrei Bernevig, and Nicolas Regnault, “Quantum many-body scars in a Landau level on a thin torus,” *Phys. Rev. B* **102**, 195150 (2020).
- [71] S. R. Taylor, M. Schulz, F. Pollmann, and R. Moessner, “Experimental probes of stark many-body localization,” *Phys. Rev. B* **102**, 054206 (2020).
- [72] Sanjay Moudgalya, Abhinav Prem, Rahul Nandkishore, Nicolas Regnault, and B. Andrei Bernevig, “Thermalization and its absence within Krylov subspaces of a constrained Hamiltonian,” (2019), [arXiv:1910.14048 \[cond-mat.str-el\]](#).
- [73] Tibor Rakovszky, Pablo Sala, Ruben Verresen, Michael Knap, and Frank Pollmann, “Statistical localization: From strong fragmentation to strong edge modes,” *Phys. Rev. B* **101**, 125126 (2020).
- [74] Johannes Feldmeier, Pablo Sala, Giuseppe De Tomasi, Frank Pollmann, and Michael Knap, “Anomalous diffusion in dipole- and higher-moment-conserving systems,” *Phys. Rev. Lett.* **125**, 245303 (2020).
- [75] Han Yan, Owen Benton, L. D. C. Jaubert, and Nic Shannon, “Rank-2 $u(1)$ spin liquid on the breathing pyrochlore lattice,” *Phys. Rev. Lett.* **124**, 127203 (2020).
- [76] R. Shankar, “Renormalization-group approach to interacting fermions,” *Reviews of Modern Physics* **66**, 129–192 (1994).
- [77] Zohar Komargodski and Adam Schwimmer, “On renormalization group flows in four dimensions,” *Journal of High Energy Physics* **2011** (2011), [10.1007/jhep12\(2011\)099](#).
- [78] Sanjay Moudgalya, Abhinav Prem, David A. Huse, and Amos Chan, “Spectral statistics in constrained many-body quantum chaotic systems,” (2021), [arXiv:2009.11863 \[cond-mat.stat-mech\]](#).