

Heterogeneous Dense Subhypergraph Detection

Mingao Yuan¹ and Zuofeng Shang²

¹Department of Statistics, North Dakota State University, e-mail: mingao.yuan@ndsu.edu

²Department of Mathematical Sciences, New Jersey Institute of Technology, e-mail: zshang@njit.edu

Abstract: We study the problem of testing the existence of a heterogeneous dense subhypergraph. The null hypothesis corresponds to a heterogeneous Erdős-Rényi uniform random hypergraph and the alternative hypothesis corresponds to a heterogeneous uniform random hypergraph that contains a dense subhypergraph. We establish detection boundaries when the edge probabilities are known and construct an asymptotically powerful test for distinguishing the hypotheses. We also construct an adaptive test which does not involve edge probabilities, and hence, is more practically useful.

MSC2020 subject classifications: Primary 62G10; Secondary 05C80,05C65.

Keywords and phrases: detection boundary, heterogeneous uniform hypergraph, dense subhypergraph, asymptotically powerful test.

1. Introduction

Suppose $(\mathcal{V}, \mathcal{E})$ is an *undirected* m -uniform hypergraph on $N := |\mathcal{V}|$ vertices with an edge set \mathcal{E} . Each edge in \mathcal{E} contains exactly m distinct vertices. Without loss of generality, assume $\mathcal{V} = [N] := \{1, 2, \dots, N\}$. The adjacency tensor is an m -dimensional 0-1 symmetric array $A \in (\{0, 1\}^N)^{\otimes m}$ such that $A_{i_1 i_2 \dots i_m} = 1$ if and only if $\{i_1, i_2, \dots, i_m\} \in \mathcal{E}$. This implies that $A_{i_1 i_2 \dots i_m} = 0$ if $\{i_1, i_2, \dots, i_m\}$ contains identical indexes, i.e., no self-loops are existent. Symmetry is defined as $A_{i_1 i_2 \dots i_m} = A_{j_1 j_2 \dots j_m}$ if i_1, i_2, \dots, i_m is a permutation of j_1, j_2, \dots, j_m . Let $p = \{p_{i_1 i_2 \dots i_m} \in [0, 1] : 1 \leq i_1 < i_2 < \dots < i_m \leq N\}$ be a collection of edge-specific probability values. Let $\mathcal{H}_m(N, p)$ denote the Erdős-Rényi m -uniform random hypergraph (see [28] for $m = 2$). Equivalently, $A_{i_1 i_2 \dots i_m}, 1 \leq i_1 < i_2 < \dots < i_m \leq N$, are independent Bernoulli variables with

$$\mathbb{P}(A_{i_1 i_2 \dots i_m} = 1) = p_{i_1 i_2 \dots i_m}. \quad (1)$$

For a positive integer $n < N$, a subset $S \subset \mathcal{V}$ with $|S| = n$ and a scalar $\rho_S > 1$, let $\mathcal{H}_m(N, p; n, \rho_S)$ denote an m -uniform random hypergraph with a dense subhypergraph S . Equivalently, $A_{i_1 i_2 \dots i_m}, 1 \leq i_1 < i_2 < \dots < i_m \leq N$, are independent Bernoulli variables such that

$$\mathbb{P}(A_{i_1 i_2 \dots i_m} = 1) = \begin{cases} p_{i_1 i_2 \dots i_m} \rho_S, & i_1, i_2, \dots, i_m \in S, \\ p_{i_1 i_2 \dots i_m}, & \text{otherwise.} \end{cases} \quad (2)$$

The assumption $\rho_S > 1$ implies that the vertices within S are more possibly connected, so S can be viewed as an underlying dense subhypergraph. Since the Bernoulli probabilities in (1) and (2) are edge-specific, models $\mathcal{H}_m(N, p)$ and $\mathcal{H}_m(N, p; n, \rho_S)$ are both heterogeneous. Given A , we are interested in the following hypothesis testing problem:

$$\begin{aligned} (\text{null hypothesis}) H_0: & \quad A \sim \mathcal{H}_m(N, p) \\ (\text{alternative hypothesis}) H_1: & \quad \text{there exists an } S \subset \mathcal{V} \text{ with } |S| = n \text{ and a } \rho_S > 1 \\ & \quad \text{such that } A \sim \mathcal{H}_m(N, p; n, \rho_S). \end{aligned} \quad (3)$$

The null hypothesis in (3) says that A follows an Erdős-Rényi m -uniform random hypergraph. The alternative hypothesis says that A follows an m -uniform random hypergraph with an underlying dense subhypergraph. When $m = 2$, [5] derive detection boundaries for testing (3). There is a lack of literature dealing with (3) for arbitrary m which will be investigated in this paper.

Given an observed hypergraph with an adjacency tensor A , a statistical test T for testing (3) is a function of A such that $T = 1$ if H_0 is rejected and $T = 0$ otherwise. Define the risk of T as

$$\phi_N(T) = \mathbb{P}_0(T = 1) + \max_{|S|=n} \mathbb{P}_S(T = 0).$$

Here \mathbb{P}_0 and \mathbb{P}_S are the probability measures under H_0 and H_1 , respectively. If $\phi_N(T) \rightarrow 0$ (or $\phi_N(T) \rightarrow 1$), we say that the test T is asymptotically powerful (or asymptotically powerless). In this paper, we provide conditions under which all tests for (3) are asymptotically powerless. We also provide conditions under which (3) is asymptotically distinguishable. As an initial stage, we consider the case of known p and construct an asymptotically powerful test statistic. We then move forward to the more realistic unknown p scenario and construct an adaptive test statistic. Our work is a hypergraphic extension of [5]. There are rich literature on classic homogeneous sub(hyper)graph detection or testing in which $p_{i_1 i_2 \dots i_m}$ is constant. For instance, [24, 19, 32, 11, 18, 13, 23, 31] proposed various detection algorithms. In stochastic block models, various algorithms for detecting the underlying communities have been proposed in [15, 16, 3, 9, 12, 21, 22, 26, 27, 20, 14, 1, 2], and [30] established sharp boundaries for testing the existence of communities. More references could be found in the survey paper [6]. Nonetheless, problem (3) is more challenging than the above homogeneous scenarios due to degree heterogeneity.

The rest of the paper proceeds as follows. In Section 2, we derive detection boundaries with all model parameters being known. In Section 3, we construct an adaptive test to address the unknown edge probabilities. All additional proofs are deferred to Section 4.

2. Detection boundary when p is known

In this section, we derive detection boundaries for testing (3) when p is known. Assume that $n, N \rightarrow \infty$ and

$$\max_{S \subset \mathcal{V}, |S|=n} \max_{\{i_1, \dots, i_m\} \subset S} \rho_S^2 p_{i_1 i_2 \dots i_m} = o(1). \quad (4)$$

Condition (4) implies that the hypergraph is suitably sparse. Let $A_D = \sum_{i_1, \dots, i_m \in D} A_{i_1 i_2 \dots i_m}$ which is the number of edges of the subhypergraph restricted to the vertex set $D \subset \mathcal{V}$. Under H_0 , the edge rate is $\mu_{D,m} = \mathbb{E}_0[A_D]/|D|^{(m)}$ with $|D|^{(m)} := \binom{|D|}{m}$. Here \mathbb{E}_0 is the expectation under H_0 . Denote $x \wedge y = \min\{x, y\}$, $x \vee y = \max\{x, y\}$ and $[x]_+ = \max\{x, 0\}$. Let

$$h(x) = (x + 1) \log(x + 1) - x, \quad \forall x \in [0, 1]$$

which is related to the Kullback-Leibler divergence between two Bernoulli distributions. Consider the following two scenarios regarding $n, N, m, \mu_{D,m}$.

Scenario 1. There exist $\delta \in (0, 0.5)$ and a positive sequence $\gamma_N = o(1)$ such that $n = O(N^{0.5-\delta})$ and

$$\max_{S \subset \mathcal{V}, |S|=n} \max_{D \subset S, |D| < n/(N/n)^{\gamma_N}} \frac{|D|^{(m)} N \mu_{D,m}}{N^{(m)} |D| \mu_{S,m}} \leq \delta, \quad (5)$$

$$\max_{S \subset \mathcal{V}, |S|=n} \frac{1}{\mu_{S,m}} = o\left(\frac{n^{m-1}}{\log \frac{N}{n}}\right). \quad (6)$$

Scenario 2. Suppose $\log n = o(\log N)$ and

$$\max_{S \subset \mathcal{V}, |S|=n} \frac{1}{\mu_{S,m}} = o\left(\frac{\log \frac{N}{n}}{\log n}\right) = o(1). \quad (7)$$

The two scenarios above impose suitable assumptions on $n, N, m, \mu_{D,m}$. For instance, (5) says that the ratio of $\mu_{D,m}$ to $\mu_{S,m}$ is well controlled when $D \subset S$ has small cardinality, i.e., a small $D \subset S$ with large node degrees is nonexistent. Hence, (5) could be considered as a measure of heterogeneity. Condition (6) or (7) requires the edge density in the dense subhypergraph being not too small, which implies that the underlying subhypergraph has enough signal to be detected. The conditions degenerate to [5] when $m = 2$. For $m \geq 3$ and $\mu_{S,m} = \mu_{S,2}$, condition (6) is weaker than $m = 2$. The theorem below provides a circumstance that the hypotheses in (3) are asymptotically indistinguishable, which is a generalization of [5] to arbitrary fixed $m \geq 3$. The indistinguishable regions for $m \geq 3$ and $m = 2$ are significantly different and the proof is technically much more involved.

Theorem 2.1. *Under either Scenario I or Scenario II, all tests are asymptotically powerless if there exists a constant $\epsilon \in (0, 1)$ such that*

$$\max_{\substack{S \subset \mathcal{V} \\ |S|=n}} \max_{D \subset S} \frac{|D|^{(m)} \mu_{D,m} h(\rho_S - 1)}{|D| \log \frac{N}{|D|}} \leq 1 - \epsilon. \quad (8)$$

Condition (8) is equivalent to that for any $S \subset \mathcal{V}$ with $|S| = n$,

$$h(\rho_S - 1) \leq \frac{1 - \epsilon}{\max_{D \subset S} \frac{|D|^{(m)} \mu_{D,m}}{|D| \log \frac{N}{|D|}}}. \quad (9)$$

Theorem 2.1 says that if ρ_S is close to one uniformly for S of cardinality n , then the hypotheses in (3) are indistinguishable. When $m = 2$, (8) degenerates to condition (5) in [5]. To gain more insights about how (8) varies along with m , we suppose $\mu_{D,m} = \mu_{D,2}$ for $m \geq 3$, i.e., the edge rates are constant along with m . Since $|D|^{(m)}$ increases in m , the RHS of (9) decreases in m , i.e., the range of ρ_S becomes smaller. This implies that testing (3) for general $m \geq 3$ becomes easier than $m = 2$. Proof of Theorem 2.1 proceeds by showing that the hypotheses in (3) are asymptotically mutually contiguous under (8) which requires analyzing the likelihood ratio statistic.

Proof of Theorem 2.1. For $i_1 < i_2 < \dots < i_m$, denote $\mathbf{i}_m = \{i_1, \dots, i_m\} \subset D$ and denote $p_{i_1 \dots i_m}$ as $p_{\mathbf{i}_m}$. Let $\theta_{\mathbf{i}_m}(q) = \log \frac{q(1-p_{\mathbf{i}_m})}{p_{\mathbf{i}_m}(1-q)}$ for any $q \in (0, 1)$, and $\Lambda_{\mathbf{i}_m}(\theta) = \log(1-p_{\mathbf{i}_m} + p_{\mathbf{i}_m} e^\theta)$. Let $H_p(q)$ be the Kullback-Leibler divergence from $\text{Bern}(q)$ to $\text{Bern}(p)$ defined as $H_p(q) = q \log \frac{q}{p} + (1-q) \log \frac{1-q}{1-p}$, for $p, q \in (0, 1)$. For a given subset $S \subset [N]$ with $|S| = n$, the likelihood ratio statistics based on (3) is equal to

$$L_S = \frac{\prod_{\substack{i_1 < \dots < i_m \\ \mathbf{i}_m \in S}} (\rho_S p_{\mathbf{i}_m})^{A_{\mathbf{i}_m}} (1 - \rho_S p_{\mathbf{i}_m})^{1-A_{\mathbf{i}_m}}}{\prod_{\substack{i_1 < \dots < i_m \\ \mathbf{i}_m \in S}} (p_{\mathbf{i}_m})^{A_{\mathbf{i}_m}} (1 - p_{\mathbf{i}_m})^{1-A_{\mathbf{i}_m}}} = \exp \left\{ \sum_{\mathbf{i}_m \in S} [A_{\mathbf{i}_m} \theta_{\mathbf{i}_m}(\rho_S p_{\mathbf{i}_m}) - \Lambda_{\mathbf{i}_m}(\theta_{\mathbf{i}_m}(\rho_S p_{\mathbf{i}_m}))] \right\}.$$

Then it's easy to express the unconditional likelihood ratio statistic as $L = \binom{N}{n}^{-1} \sum_{|S|=n} L_S$. We adopt the likelihood ratio truncation skill used in [4, 7] to get $\tilde{L} = \binom{N}{n}^{-1} \sum_{|S|=n} L_S I_{\Gamma_S}$, where I_E is an indicator function for event E and

$$\Gamma_S = \left\{ \sum_{\mathbf{i}_m \in D} A_{\mathbf{i}_m} \theta_{\mathbf{i}_m}(\rho_S p_{\mathbf{i}_m}) \leq \sum_{\mathbf{i}_m \in D} \zeta_D \theta_{\mathbf{i}_m}(\rho_S p_{\mathbf{i}_m}), \text{ all } D \in E_S \right\},$$

with ζ_D provided in Lemma 4.1. We will proceed by showing $\mathbb{E}_0 \tilde{L} = 1 + o(1)$ and $\mathbb{E}_0 \tilde{L}^2 \leq 1 + o(1)$, where \mathbb{E}_0 denotes expectation under H_0 .

We begin with the first-order moment. It is easy to verify $\mathbb{E}_0 \tilde{L} = \mathbb{P}_S \Gamma_S$. Note that $\theta_{\mathbf{i}_m}(\rho_S p_{\mathbf{i}_m}) \sim \log \rho_S$ uniformly for all $\mathbf{i}_m \in S$ with $|S| = n$ by a similar proof of equation (54) in [5]. By Bennett's inequality, it follows that

$$1 - \mathbb{P}_S \Gamma_S = \mathbb{P}_S \Gamma_S^c \leq \sum_{D \in E_S} \mathbb{P} \left(\sum_{\mathbf{i}_m \in D} A_{\mathbf{i}_m} > \zeta_D \sum_{\mathbf{i}_m \in D} p_{\mathbf{i}_m} \right) \leq \sum_{D \in E_S} \exp \left(-\rho_S \sum_{\mathbf{i}_m \in D} p_{\mathbf{i}_m} h \left(\frac{\zeta_D}{\rho_S} - 1 \right) \right).$$

By Lemma 4.2, we get

$$\begin{aligned} 1 - \min_{S \subset \mathcal{V}, |S|=n} \mathbb{P}_S \Gamma_S &\leq \max_{S \subset \mathcal{V}, |S|=n} \sum_{D \in E_S} \exp \left(-\rho_S \sum_{\mathbf{i}_m \in D} p_{\mathbf{i}_m} h \left(\frac{\zeta_D}{\rho_S} - 1 \right) \right) \\ &\leq \max_{S \subset \mathcal{V}, |S|=n} \sum_{k=m}^n \binom{N}{n} \exp \left(-\rho_S \sum_{\mathbf{i}_m \in D} p_{\mathbf{i}_m} h \left(\frac{\zeta_D}{\rho_S} - 1 \right) \right) \\ &\leq \max_{S \subset \mathcal{V}, |S|=n} \sum_{k=m}^n \exp \left(k \log \frac{ne}{k} - \rho_S \sum_{\mathbf{i}_m \in D} p_{\mathbf{i}_m} h \left(\frac{\zeta_D}{\rho_S} - 1 \right) \right) \\ &\leq \sum_{k=m}^n e^{-k(c_n - 1)} = o(1). \end{aligned}$$

Hence, $\mathbb{E}_0 \tilde{L} = 1 + o(1)$.

We next consider the second-order moment. For $S_1, S_2 \subset \mathcal{V}$ with $|S_1| = |S_2| = n$, let $D = S_1 \cap S_2$. Clearly, we have the following

$$\begin{aligned} \mathbb{E}_0 \tilde{L}^2 &= \binom{N}{n}^{-2} \sum_{|S_1|=n, |S_2|=n} \mathbb{E}_0 L_{S_1} I_{\Gamma_{S_1}} L_{S_2} I_{\Gamma_{S_2}} \\ &= \mathbb{E}_0 \left(I_{\Gamma_{S_1}} I_{\Gamma_{S_2}} \exp \left(\sum_{\mathbf{i}_m \in D} A_{\mathbf{i}_m} [\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m}) + \theta_{\mathbf{i}_m}(\rho_{S_2} p_{\mathbf{i}_m})] - \Lambda_{\mathbf{i}_m}(\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m})) - \Lambda_{\mathbf{i}_m}(\theta_{\mathbf{i}_m}(\rho_{S_2} p_{\mathbf{i}_m})) \right) \right) \\ &\leq \mathbb{E} \left[I_{D \in E_{S_1}} \mathbb{E}_0 \left(I_{\Gamma_{S_1}} \exp \left(\sum_{\mathbf{i}_m \in D} 2A_{\mathbf{i}_m} [\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m}) - 2\Lambda_{\mathbf{i}_m}(\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m}))] \right) \right) \right] \\ &\quad + \mathbb{E} \left[I_{D \notin E_{S_1}} \mathbb{E}_0 \left(I_{\Gamma_{S_1}} \exp \left(\sum_{\mathbf{i}_m \in D} 2A_{\mathbf{i}_m} [\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m}) - 2\Lambda_{\mathbf{i}_m}(\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m}))] \right) \right) \right] \\ &= R_1 + R_2. \end{aligned}$$

We shall show $R_1 = o(1)$ and $R_2 \leq 1 + o(1)$. Since $p_{\mathbf{i}_m} = o(1)$, we have

$$\Delta_{\mathbf{i}_m}^{(1)} := \log \left(1 + \frac{p_{\mathbf{i}_m}(\rho_{S_1} - 1)^2}{1 - p_{\mathbf{i}_m}} \right) \leq p_{\mathbf{i}_m}(\rho_{S_1} - 1)^2.$$

Besides,

$$\mathbb{P}(|D| = k) \leq \exp \left(-k \left(\log \frac{Nk}{n^2} + O(1) \right) \right).$$

Consequently, by the definition of E_{S_1} one has

$$\begin{aligned} R_2 &\leq \mathbb{E} \left[I_{D \notin E_{S_1}} \exp \left(\sum_{\mathbf{i}_m \in D} \Delta_{\mathbf{i}_m}^{(1)} \right) \right] \\ &\leq \mathbb{E} \left[I_{D \notin E_{S_1}} \exp \left(\left(1 - \frac{\epsilon}{2}\right) |D| \left(\log \frac{N|D|}{n^2} - \log \log \frac{N}{n} \right) \right) \right] \\ &\leq 1 + \sum_{k=m}^n \exp \left(k \left(\log \frac{Nk}{n^2} - \log \log \frac{N}{n} \right) - k \left(\log \frac{Nk}{n^2} + O(1) \right) \right) = 1 + o(1). \end{aligned}$$

Note that $\frac{\log \zeta_D}{\log \rho_{S_1}} \sim \frac{\theta_{\mathbf{i}_m}(\zeta_D p_{\mathbf{i}_m})}{\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m})}$ uniformly for $\mathbf{i}_m \in D$, $D \in E_{S_1}$ and $|S_1| = n$. For any $x \in [0, 1]$ and $D \in E_{S_1}$, we have

$$R_1 \leq \mathbb{E} \left[I_{D \in E_{S_1}} \mathbb{E}_0 \left(\exp \left(\sum_{\mathbf{i}_m \in D} 2\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m}) (x A_{\mathbf{i}_m} + (1-x) \zeta_D p_{\mathbf{i}_m}) - 2\Lambda_{\mathbf{i}_m}(\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m})) \right) \right) \right],$$

which is minimized at $x = \frac{\theta_{\mathbf{i}_m}(\zeta_D p_{\mathbf{i}_m})}{2\theta_{\mathbf{i}_m}(\rho_{S_1} p_{\mathbf{i}_m})} \sim \frac{\log \zeta_D}{2 \log \rho_{S_1}}$ by (50) in [5]. Plugging this into the above equation yields

$$R_1 \leq \mathbb{E} \left[I_{D \in E_{S_1}} \mathbb{E}_0 \exp \left(\sum_{\mathbf{i}_m \in D} \Delta_{\mathbf{i}_m}^{(2)} \right) \right] \leq \sum_{k=m}^n \exp(k(-2c_n + O(1))) = o(1),$$

where $\Delta_{\mathbf{i}_m}^{(2)} = H_{p_{\mathbf{i}_m}}(\zeta_D p_{\mathbf{i}_m}) - 2H_{\rho_{S_1} p_{\mathbf{i}_m}}(\zeta_D p_{\mathbf{i}_m})$, the second inequality follows from the proof of Theorem 1 in [5] and the last step follows from Lemma 4.2. Then the proof is complete. \square

Next, we shall show that the condition (8) in Theorem 2.1 is also necessary for indistinguishability. Define the hypergraphic scan statistic as

$$T_n = \max_{D \subset \mathcal{V}, |D| \leq n} T_D, \quad \text{where } T_D = \frac{|D|^{(m)}}{|D|} \cdot \frac{\mu_{D,m} h \left(\left[\frac{A_D}{\mathbb{E}_0[A_D]} - 1 \right]_+ \right)}{\log \frac{N}{|D|}}. \quad (10)$$

In the above, $\mathbb{E}_0[A_D] = \sum_{\mathbf{i}_m \in D} p_{\mathbf{i}_m}$ which is available given that p is known. For any $S \subset \mathcal{V}$ with $|S| = n$, let

$$D_S^* = \arg \max_{D \subset S} \frac{|D|^{(m)}}{|D|} \cdot \frac{\mu_{D,m}}{\log \frac{N}{|D|}}.$$

Theorem 2.2. *Suppose $n = o(N)$ and $\rho_S \mathbb{E}_0[A_{D_S^*}] \rightarrow \infty$. The scan test T_n is asymptotically powerful if there exists a constant $\epsilon \in (0, 1)$ such that*

$$\min_{\substack{S \subset \mathcal{V} \\ |S|=n}} \max_{D \subset S} \frac{|D|^{(m)}}{|D|} \cdot \frac{\mu_{D,m} h(\rho_S - 1)}{\log \frac{N}{|D|}} \geq 1 + \epsilon. \quad (11)$$

Theorem 2.2 can be proved similarly as Theorem 2 in [5]. Specifically, it applies the Bennett's inequality to show that the γ_N -risk of the scan test T_n tends to zero. Theorems 2.1 and 2.2 together depict a detection boundary for testing (3).

3. An adaptive test

In practice, p is often unknown so the test statistic T_n may not be applicable. Instead, we will propose a new test which is adaptive to the value of p . This problem is challenging since p contains $N^{(m)}$ unknown parameters and estimation of these large amount of parameters seems infeasible. To reduce the amount of over-parametrization, consider a special case that $p_{i_1 \dots i_m} = \prod_{k=1}^m W_{i_k}$ for an unknown vector $W = (W_1, \dots, W_N)$

with $W_i \in [0, 1]$. When $m = 2$, this is just the rank-1 model studied in [5]. Moreover, assume that

$$\left(\frac{W_{max}}{W_{min}}\right)^m = o\left(n^{\frac{m}{m+1}} \wedge \left[W_{min}^m \left(\frac{N}{n}\right)^{m-1-\delta_m}\right]\right), \quad (12)$$

where $\delta_m = 0$ for even m and $\delta_m \in (0, 1)$ for odd m , $W_{max} = \max\{W_1, \dots, W_N\}$ and $W_{min} = \min\{W_1, \dots, W_N\}$. Note that the RHS of (12) converges to zero faster when m is even. Condition (12) accommodates heterogeneity in the hypergraph. To see this, consider $n = \sqrt{N}$ and $W_i = \left(\frac{i}{N}\right)^{\frac{1}{\kappa(m+1)}}$ in which $\kappa \geq 4$ is a constant. Since $\sum_{i=1}^N W_i \asymp \frac{k(m+1)}{k(m+1)+1}N$, it is easy to verify that the average degree of node N (of order N^{m-1}) is approximately $N^{\frac{1}{\kappa(m+1)}}$ times of the average degree of node 1 (of order $N^{m-1-\frac{1}{\kappa(m+1)}}$).

When the edge probability p is unknown, we need to modify the scan test. Essentially, we have to estimate $\mathbb{E}_0[A_D]$ for any subset $D \subset \mathcal{V}$ by $\widehat{p}_{D,m}$ defined as follows:

$$\widehat{p}_{D,m} = \frac{1}{2^m} \left(A_{\mathcal{V}}^{\frac{1}{m}} - (A_{\mathcal{V}} - 2A_{D,D^c})^{\frac{1}{m}} \right)^m,$$

where

$$A_{D,D^c} = \frac{1}{m!} \sum_{k=1}^{t_m} \binom{m}{2k-1} \sum_{\substack{\{i_1, \dots, i_{2k-1}\} \subset D^c \\ \{i_{2k}, \dots, i_m\} \subset D}} A_{i_1 \dots i_m},$$

$t_m = \frac{m}{2}$ for even m , and $t_m = \frac{m+1}{2}$ for odd m , and $D^c = \mathcal{V} - D$. Define

$$\widehat{p}_{D,m}^* = \widehat{p}_{D,m} \vee \frac{|D|^m}{N^{m-1}} \log^{2m} \frac{N}{|D|}.$$

We then propose the following modified scan test statistic:

$$\widehat{T}_n = \max_{D \subset \mathcal{V}, |D| \leq n} \widehat{T}_D, \quad \widehat{T}_D = \frac{\widehat{p}_{D,m}^* h\left(\left[\frac{A_D}{\widehat{p}_{D,m}^*} - 1\right]_+\right)}{|D| \log \frac{N}{|D|}}. \quad (13)$$

Note that \widehat{T}_n does not involve p and hence is adaptive. Theorem 3.1 shows that \widehat{T}_n is asymptotically powerful under the condition (11).

Theorem 3.1. *Suppose $n = o(N)$ and $\rho_S \mathbb{E}_0[A_{D_S}^*] \rightarrow \infty$. If (11) holds, then the modified scan test \widehat{T}_n is asymptotically powerful.*

Based on Theorem 3.1, the modified scan test \widehat{T}_n still can achieve the detection boundary in Theorem 2.2. Moreover, the rate $n^{\frac{m}{m+1}}$ is optimal under Scenario I and Scenario 2 and it can't be improved.

The main ingredient in the proof of Theorem 3.1 is to show that $\widehat{p}_{D,m}$ can accurately estimate $\mathbb{E}_0[A_D]$, hence, \widehat{T}_n will perform similarly as T_n . Before proving Theorem 3.1, let us provide a quick sketch on the estimation accuracy of $\widehat{p}_{D,m}$. Under assumption (12), it can be shown that for any $D \subset \mathcal{V}$, $\sum_{i \in \mathcal{V}-D} W_i - \sum_{i \in D} W_i$ is non-negative for large N . Therefore, we have

$$2 \sum_{i \in D} W_i = \sum_{i \in \mathcal{V}} W_i - \left(\sum_{i \in \mathcal{V}-D} W_i - \sum_{i \in D} W_i \right) = \left[\left(\sum_{i \in \mathcal{V}} W_i \right)^m \right]^{\frac{1}{m}} - \left[\left(\sum_{i \in \mathcal{V}-D} W_i - \sum_{i \in D} W_i \right)^m \right]^{\frac{1}{m}}.$$

Note that

$$\left(\sum_{i \in \mathcal{V}} W_i \right)^m = m! \mathbb{E}_0 A_{\mathcal{V}} + \sum_{\substack{|\{i_1, i_2, \dots, i_m\}| \leq m-1 \\ \{i_1, \dots, i_m\} \subset \mathcal{V}}} W_{i_1} \cdots W_{i_m}$$

and, by binomial formula,

$$\begin{aligned}
& \left(\sum_{i \in \mathcal{V}-D} W_i - \sum_{i \in D} W_i \right)^m \\
&= \left(\sum_{i \in \mathcal{V}-D} W_i + \sum_{i \in D} W_i \right)^m - 2 \sum_{k=1}^{t_m} \binom{m}{2k-1} \left(\sum_{i \in \mathcal{V}-D} W_i \right)^{2k-1} \left(\sum_{i \in D} W_i \right)^{m-2k+1} \\
&= m! \mathbb{E}_0 A_{\mathcal{V}} + \sum_{\substack{|\{i_1, i_2, \dots, i_m\}| \leq m-1 \\ \{i_1, \dots, i_m\} \subset \mathcal{V}}} W_{i_1} \cdots W_{i_m} - 2 \sum_{k=1}^{t_m} \binom{m}{2k-1} \left(\sum_{i \in \mathcal{V}-D} W_i \right)^{2k-1} \left(\sum_{i \in D} W_i \right)^{m-2k+1}.
\end{aligned}$$

By law of large number, it can be shown that $A_{\mathcal{V}} = (1 + o_P(1)) \mathbb{E}_0[A_{\mathcal{V}}]$, and for any $D \subset \mathcal{V}$, $A_{D, D^c} = (1 + o_P(1)) \mathbb{E}_0[A_{D, D^c}]$. By assumption (12), it can be shown that, for any $D \subset \mathcal{V}$,

$$\mathbb{E}_0[A_D] \gg \frac{1}{m!} \sum_{\substack{|\{i_1, i_2, \dots, i_m\}| \leq m-1 \\ \{i_1, \dots, i_m\} \subset D}} W_{i_1} \cdots W_{i_m},$$

which leads to that

$$\begin{aligned}
\left(\sum_{i \in \mathcal{V}} W_i \right)^m &= (1 + o_P(1)) m! A_{\mathcal{V}}, \\
\left(\sum_{i \in \mathcal{V}-D} W_i - \sum_{i \in D} W_i \right)^m &= (1 + o_P(1)) m! A_{\mathcal{V}} - 2(1 + o_P(1)) m! A_{D, D^c}.
\end{aligned}$$

Hence, we get

$$\begin{aligned}
\mathbb{E}_0[A_D] &= \sum_{\substack{i_1 < i_2 < \dots < i_m \\ \{i_1, \dots, i_m\} \subset D}} W_{i_1} \cdots W_{i_m} \\
&= \frac{1}{2^m m!} \left(2 \sum_{i \in D} W_i \right)^m - \frac{1}{m!} \sum_{\substack{|\{i_1, i_2, \dots, i_m\}| \leq m-1 \\ \{i_1, \dots, i_m\} \subset D}} W_{i_1} \cdots W_{i_m} \\
&= \frac{1 + o_P(1)}{2^m} \left(A_{\mathcal{V}}^{\frac{1}{m}} - (A_{\mathcal{V}} - 2A_{D, D^c})^{\frac{1}{m}} \right)^m - o(\mathbb{E}_0[A_D]) \\
&= (1 + o_P(1)) \widehat{p}_{D, m} - o(\mathbb{E}_0[A_D]).
\end{aligned} \tag{14}$$

Therefore, $(1 + o_P(1)) \widehat{p}_{D, m} = (1 + o_P(1)) \mathbb{E}_0[A_D]$, i.e., $\widehat{p}_{D, m}$ is proven a good estimator of $\mathbb{E}_0[A_D]$.

Proof of Theorem 3.1. Firstly, we control the type I error. Note that for $a \geq b$, $ah \left(\left[\frac{a}{b} - 1 \right]_+ \right) \leq bh \left(\left[\frac{a}{b} - 1 \right]_+ \right)$. We only need to prove

$$\max_{\frac{1}{m+1} \leq |D| \leq n} \frac{\mathbb{E}_0[A_D]}{\widehat{p}_{D, m}^*} \leq 1 + o_P(1). \tag{15}$$

Define $\mathcal{D} = \left\{ D \subset \mathcal{V} : \frac{1}{m+1} \leq |D| \leq n, \widehat{p}_{D, m}^* \leq \mathbb{E}_0[A_D] \right\}$. It suffices to prove (15) for $D \in \mathcal{D}$. By the definition of $\widehat{p}_{D, m}^*$, we have

$$\frac{|D|^m}{N^{m-1}} \log^{2m} \frac{N}{|D|} \leq \widehat{p}_{D, m}^* \leq \mathbb{E}_0[A_D] \leq \left(\sum_{i \in D} W_i \right)^m,$$

which implies $\sum_{i \in D} W_i \geq \frac{|D|}{N^{\frac{m-1}{m}}} \log^2 \frac{N}{|D|}$. Besides, by assumption (12), we have $W_{\min} \geq \frac{1}{N^{\frac{m-1-\delta_m}{m}}}$. Hence,

$$\begin{aligned} \mathbb{E}_0[A_{D,D^c}] &= \frac{1}{m!} \sum_{k=1}^{t_m} \binom{m}{2k-1} \left(\sum_{i \in \mathcal{V}-D} W_i \right)^{2k-1} \left(\sum_{i \in D} W_i \right)^{m-2k+1} \\ &\geq \frac{1}{m!} \sum_{k=1}^{t_m} \binom{m}{2k-1} (N-|D|)^{2k-1} W_{\min}^{2k-1} \left(\sum_{i \in D} W_i \right)^{m-2k+1} \\ &\geq \frac{1}{m!} \sum_{k=1}^{t_m} \binom{m}{2k-1} \frac{(N-|D|)^{2k-1}}{N^{\frac{m-1-\delta_m}{m}(2k-1)}} \frac{\left(|D| \log^2 \frac{N}{|D|} \right)^{m-2k+1}}{N^{\frac{m-1}{m}(m-2k+1)}}. \end{aligned} \quad (16)$$

For even m and a constant $c_1 > 0$, using the last term $k = \frac{m}{2}$ in (16), we have

$$\mathbb{E}_0[A_{D,D^c}] \geq c_1 |D| \log^2 \frac{N}{|D|}.$$

For odd m and a constant $c_2 > 0$, using the last term $k = \frac{m+1}{2}$ in (16), we have

$$\mathbb{E}_0[A_{D,D^c}] \geq c_2 N^{1+\delta_m} \geq c_2 |D| \log^2 \frac{N}{|D|}.$$

Take $c = \min\{c_1, c_2\}$. For a constant $c_0 > 0$, by Bennett's inequality we get

$$\begin{aligned} &\mathbb{P} \left(\min_{D \in \mathcal{D}} A_{D,D^c} - \mathbb{E}_0[A_{D,D^c}] \leq -\sqrt{2c \left(\frac{1}{c} + c_0 \right) \mathbb{E}_0[A_{D,D^c}] |D| \log \frac{N}{|D|}} \right) \\ &\leq \sum_{k=m}^n \sum_{|D|=k} \exp \left(-\mathbb{E}_0[A_{D,D^c}] h \left(\sqrt{\frac{2c \left(\frac{1}{c} + c_0 \right) |D| \log \frac{N}{|D|}}{\mathbb{E}_0[A_{D,D^c}]}} \right) \right) \\ &\leq \sum_{k=m}^n \sum_{|D|=k} \exp \left(-c \left(\frac{1}{c} + c_0 \right) |D| \log \frac{N}{|D|} \right) \\ &\leq \sum_{k=m}^n \exp \left(k \log \frac{Ne}{n} \right) \exp \left(-c \left(\frac{1}{c} + c_0 \right) k \log \frac{N}{k} \right) = o(1). \end{aligned}$$

Here, we used the fact that $h(x) \sim \frac{x^2}{2}$ for $x = o(1)$. Consequently, we have

$$A_{D,D^c} = (1 + o_p(1)) \mathbb{E}_0[A_{D,D^c}],$$

uniformly for $D \in \mathcal{D}$. Obviously,

$$A_{\mathcal{V}} = (1 + o_P(1)) \mathbb{E}_0[A_{\mathcal{V}}].$$

By Lemma 4.6, we obtain

$$\frac{\widehat{p}_{D,m}^*}{\mathbb{E}_0[A_{\mathcal{V}}]} \geq \frac{\widehat{p}_{D,m}}{\mathbb{E}_0[A_{\mathcal{V}}]} = 1 + o_p(1).$$

Then by the proof of Theorem 2.2, the type I error goes to zero.

Next, we control type II error. Obviously, we have

$$A_{D^*,D^{*c}} = (1 + o_p(1)) \mathbb{E}_1[A_{D^*,D^{*c}}], \quad A_{\mathcal{V}} = (1 + o_P(1)) \mathbb{E}_1[A_{\mathcal{V}}].$$

By assumption (12), since $\rho_S W_{\min}^m \leq 1$, then $\rho_S \ll \left(\frac{W_{\min}}{W_{\max}} \right)^m \left(\frac{N}{n} \right)^{m-1-\delta_m}$. Then

$$1 \leq \frac{\mathbb{E}_1[A_{D^*,D^{*c}}]}{\mathbb{E}_0[A_{D^*,D^{*c}}]} \leq 1 + \frac{\mathbb{E}_1[A_{D^*,C-D^*}]}{\mathbb{E}_0[A_{D^*,D^{*c}}]} = 1 + \rho_S \frac{\mathbb{E}_0[A_{D^*,C-D^*}]}{\mathbb{E}_0[A_{D^*,D^{*c}}]}.$$

For even m , we have

$$\rho_S \frac{\mathbb{E}_0[A_{D^*, C-D^*}]}{\mathbb{E}_0[A_{D^*, D^{*c}}]} = O\left(\rho_S \frac{|D^*|(n - |D^*|)^{m-1} \vee |D^*|^m \frac{W_{max}^m}{W_{min}^m}}{|D^*|(N - |D^*|)^{m-1}}\right) = o(1).$$

For odd m , we have

$$\rho_S \frac{\mathbb{E}_1[A_{D^*, C-D^*}]}{\mathbb{E}_0[A_{D^*, D^{*c}}]} = O\left(\rho_S \frac{(n - |D^*|)^m \vee |D^*|^m \frac{W_{max}^m}{W_{min}^m}}{(N - |D^*|)^m}\right) = o(1).$$

Hence, $\mathbb{E}_1[A_{D^*, D^{*c}}] = (1 + o(1))\mathbb{E}_0[A_{D^*, D^{*c}}]$. Similarly we can get $\mathbb{E}_1[A_{\mathcal{V}}] = (1 + o(1))\mathbb{E}_0[A_{\mathcal{V}}]$. Consequently, we have

$$A_{D^*, D^{*c}} = (1 + o_P(1))\mathbb{E}_0[A_{D^*, D^{*c}}], \quad A_{\mathcal{V}} = (1 + o_P(1))\mathbb{E}_0[A_{\mathcal{V}}].$$

By Lemma 4.6, one has $\widehat{p}_{D^*, m} = (1 + o_P(1))\mathbb{E}_0[A_{D^*}]$. Hence,

$$\widehat{p}_{D^*, m}^* = (1 + o_P(1))\mathbb{E}_0[A_{D^*}] \vee \frac{|D^*|^m \log^{2m} \frac{N}{|D^*|}}{N^{m-1}}.$$

If $\widehat{p}_{D^*, m}^* = (1 + o_P(1))\mathbb{E}_0[A_{D^*}]$, the proof is the same as Theorem 2.2.

Next, we assume $\widehat{p}_{D^*, m}^* = \frac{|D^*|^m \log^{2m} \frac{N}{|D^*|}}{N^{m-1}} > \mathbb{E}_0[A_{D^*}]$. Note that $h^{-1}(x) \geq \sqrt{x}$. Then condition (??) implies that

$$\rho_S > \sqrt{\frac{N^{m-1}}{|D^*|^{m-1} \log^{2m-1} \frac{N}{|D^*|}} \frac{1}{|D^*|}}.$$

Recall that $W_{min} \geq \frac{1}{N^{\frac{m-1-\delta m}{m}}}$. Hence,

$$\frac{\mathbb{E}_1[A_{D^*}]}{\widehat{p}_{D^*, m}^*} \geq \frac{\rho_S |D^*|^m W_{min}^m}{|D^*|^m \log^{2m} \frac{N}{|D^*|}} \geq \frac{\rho_S}{\log^{2m} \frac{N}{|D^*|}} \rightarrow \infty.$$

Since $A_{D^*} = (1 + o_P(1))\mathbb{E}_1[A_{D^*}]$ and $h(x-1) \sim x \log x$ for $x \rightarrow \infty$, we have

$$\begin{aligned} \widehat{p}_{D^*, m}^* h\left(\left[\frac{A_{D^*}}{\widehat{p}_{D^*, m}^*} - 1\right]_+\right) &\geq \widehat{p}_{D^*, m}^* h\left(\left[\frac{\mathbb{E}_1[A_{D^*}]}{\widehat{p}_{D^*, m}^*} - 1\right]_+\right) \\ &\geq \mathbb{E}_1[A_{D^*}] \log \frac{\mathbb{E}_1[A_{D^*}]}{\widehat{p}_{D^*, m}^*} \\ &\geq \mathbb{E}_1[A_{D^*}] \log \frac{\rho_S}{\log^{2m} \frac{N}{|D^*|}} \\ &\geq \mathbb{E}_1[A_{D^*}] \log \rho_S = \mathbb{E}_0[A_{D^*}] h(\rho_S - 1) \\ &\geq (1 + \epsilon) |D^*| \log \frac{N}{|D^*|}. \end{aligned}$$

Proof is complete. □

4. Proof of additional lemmas

In this section, we prove the lemmas. For a subset $S \subset \mathcal{V}$, define

$$E_S = \left\{ D \subset S : (\rho_S - 1)^2 \mathbb{E}_0[A_D] > \left(1 - \frac{\epsilon}{2}\right) |D| \left(\log \frac{N|D|}{n^2} - \log \log \frac{N}{n} \right) \right\}.$$

The following preliminary lemmas can be found in [5].

Lemma 4.1. *Under the conditions of Theorem 2.1, for any $S \subset \mathcal{V}$ with $|S| = n$ and $D \in E_S$, there is a unique number $\zeta_D \geq 1$ satisfying*

$$(1 + \epsilon)\mathbb{E}_0[A_D]h(\zeta_D - 1) = |D| \log \frac{N}{|D|},$$

and $\theta_{i_1 \dots i_m}(\zeta_{DP_{i_1 \dots i_m}}) \leq 2\theta_{i_1 \dots i_m}(\rho_{SP_{i_1 \dots i_m}})$ for $i_1, \dots, i_m \in D$.

Lemma 4.2. *Under the conditions of Theorem 2.1, we have*

$$c_n := \min_{S \subset \mathcal{V}, |S|=n} \min_{D \in E_S} \left((1 - \epsilon) \frac{|D|^{(m)}}{|D|} \rho_S \mu_{D,m} h \left(\frac{\zeta_D}{\rho_S} - 1 \right) - \log \frac{n}{|D|} \right) \rightarrow \infty.$$

Lemma 4.3. *Under the conditions of Theorem 2.1, for any $S \subset \mathcal{V}$ with $|S| = n$ and $D \in E_S$, we have*

$$\frac{\log \frac{n}{|D|}}{\log \frac{N}{n}} (\log \rho_S \vee 1) = o(1), \quad \frac{\log \frac{N}{n}}{\log \rho_S} \rightarrow \infty.$$

Proof of Lemma 4.3. Firstly, we consider Scenario I. For any $S \subset \mathcal{V}$ with $|S| = n$, define $\eta_S \geq \rho_S$ such that

$$\frac{n^{(m)} \mu_{S,m} h(\eta_S - 1)}{n \log \frac{N}{n}} = 1 - \frac{2\epsilon}{3}.$$

Hence, by (6), we get

$$h(\eta_S - 1) < \frac{n \log \frac{N}{n}}{n^{(m)} \mu_{S,m}} = o(1),$$

which implies $\eta_S = 1 + o(1)$. Hence, $h(\eta_S - 1) \sim \frac{(\eta_S - 1)^2}{2}$. If $|D| < \left(\frac{n}{\frac{N}{n}}\right)^{\frac{1}{\gamma_N}}$, then we have

$$\begin{aligned} (\rho_S - 1)^2 \frac{\mathbb{E}_0[A_D]}{|D|} &\leq (\eta_S - 1)^2 \frac{|D|^{(m)}}{|D|} \mu_{D,m} \\ &\leq \delta (\eta_S - 1)^2 \frac{|S|^{(m)}}{|S|} \mu_{S,m} \\ &= \left(1 - \frac{2\epsilon}{3}\right) \delta (\eta_S - 1)^2 \frac{\log \frac{N}{n}}{h(\eta_S - 1)} \\ &= \left(1 - \frac{2\epsilon}{3}\right) 2\delta \log \frac{N}{n} \\ &\leq \left(1 - \frac{\epsilon}{2}\right) \left(\log \frac{N|D|}{n^2} - \log \log \frac{N}{n}\right). \end{aligned}$$

This is a contradiction to the fact that $D \in E_S$. As a result, we have $|D| \geq \left(\frac{n}{\frac{N}{n}}\right)^{\frac{1}{\gamma_N}}$. Then the desired result follows.

Under Scenario II, the proof is almost the same as in [5]. We omit it here. \square

Lemma 4.4. *Under the assumption (12), we have $|D^*| \geq n^{\frac{1}{m+1}}$.*

Proof of Lemma 4.4. Note that the function $f(x) = (x - 1) \cdots (x - m + 1) / (\log N - \log x)$ is increasing for $x \ll N$. Suppose $|D| < n^{\frac{1}{m+1}}$ for any $D \subset \mathcal{V}$, then we have

$$\begin{aligned} \frac{|D|^{(m)}}{|D|} \frac{\mu_{D,m}}{\log \frac{N}{|D|}} &\leq \frac{|D|^{(m)} W_{max}^m}{|D| \log \frac{N}{|D|}} \leq \frac{|n^{\frac{1}{m+1}}|^{(m)} W_{max}^m}{|n^{\frac{1}{m+1}}| \log \frac{N}{|n^{\frac{1}{m+1}}|}} \\ &< \frac{|n^{\frac{1}{m+1}}|^{(m)} n^{\frac{m}{m+1}} W_{min}^m}{|n^{\frac{1}{m+1}}| \log \frac{N}{|n^{\frac{1}{m+1}}|}} \leq \frac{|n^{\frac{1}{m+1}}|^{(m)} n^{\frac{m+m}{m+1}} W_{min}^m}{|n^{\frac{1+m}{m+1}}| \log \frac{N}{|S|}} \\ &\leq \frac{|S|^{(m)} W_{min}^m}{|S| \log \frac{N}{|S|}} \leq \frac{|S|^{(m)} \mu_{S,m}}{|S| \log \frac{N}{|S|}}. \end{aligned}$$

Hence, by the definition of D^* , we get $|D^*| \geq n^{\frac{1}{m+1}}$. \square

Lemma 4.5. Suppose $n, k \geq 2$ are fixed integers. Let $f(x_1, \dots, x_n) = \frac{\sum_{i=1}^n x_i^k}{(\sum_{i=1}^n x_i)^k}$ be a function on $x_i \in [a, b], i = 1, 2, \dots, n$ and $0 < a < b < 1$. Then

$$\max_{x_i \in [a, b], i=1, \dots, n} f(x_1, \dots, x_n) \leq \frac{x_0(a^k - b^k) + nb^k}{[x_0(a - b) + nb]^k} \leq \frac{1}{(k-1)n^{k-1}} \sum_{t=1}^{k-1} \left(\frac{b}{a}\right)^t,$$

where

$$x_0 = \frac{(a^k - b^k)nb - kn(a - b)b^k}{(k-1)(a-b)(a^k - b^k)}.$$

Proof of Lemma 4.5. Consider $f(x_1, \dots, x_n)$ as a function of x_1 with the rest arguments fixed. The derivative with respect to x_1 is equal to

$$f_{x_1} = \frac{k \sum_{i=2}^n x_i}{(\sum_{i=1}^n x_i)^{k+1}} \left(x_1^{k-1} - \left[\left(\frac{\sum_{i=2}^n x_i^k}{\sum_{i=2}^n x_i} \right)^{\frac{1}{k-1}} \right]^{k-1} \right).$$

When $x_1 > \left(\frac{\sum_{i=2}^n x_i^k}{\sum_{i=2}^n x_i} \right)^{\frac{1}{k-1}}$, $f(x_1, \dots, x_n)$ is increasing as a function of x_1 . When $x_1 < \left(\frac{\sum_{i=2}^n x_i^k}{\sum_{i=2}^n x_i} \right)^{\frac{1}{k-1}}$, $f(x_1, \dots, x_n)$ is decreasing as a function of x_1 . Hence, we get

$$\max_{x_1 \in [a, b]} f(x_1, \dots, x_n) = \frac{a^k + \sum_{i=2}^n x_i^k}{(a + \sum_{i=2}^n x_i)^k} \vee \frac{b^k + \sum_{i=2}^n x_i^k}{(b + \sum_{i=2}^n x_i)^k}.$$

Repeating this procedure for each x_2, \dots, x_n , we conclude

$$\max_{x_i \in [a, b], i=1, \dots, n} f(x_1, \dots, x_n) = \max_{0 \leq x \leq n, x: \text{integer}} \frac{xa^k + (n-x)b^k}{[xa + (n-x)b]^k} \leq \max_{0 \leq x \leq n} \frac{x(a^k - b^k) + nb^k}{[x(a-b) + nb]^k}.$$

Let $g(x) = \frac{x(a^k - b^k) + nb^k}{[x(a-b) + nb]^k}$ for $0 \leq x \leq n$. The derivative of $g(x)$ is

$$g_x = \frac{(k-1)(a-b)(a^k - b^k)}{[x(a-b) + nb]^{k+1}} (x_0 - x).$$

When $x < x_0$, $g(x)$ is increasing. When $x > x_0$, the $g(x)$ is decreasing. Hence, $\max_{x \in [0, n]} g(x) = g(x_0)$. Note that

$$\begin{aligned} x_0(a^k - b^k) + nb^k &= \frac{(a^k - b^k)nb - kn(a-b)b^k}{(k-1)(a-b)} + nb^k \\ &= \frac{n(a^{k-1}b + a^{k-2}b^2 + \dots + ab^{k-1} + b^k) - nb^k}{k-1} \\ &= \frac{na^k}{k-1} \sum_{t=1}^{k-1} \left(\frac{b}{a}\right)^t, \\ x_0(a-b) + nbk &= \frac{knb}{k-1} - \frac{knb^k}{(k-1)(a^{k-1} + a^{k-2}b + \dots + b^{k-1})} \\ &= \frac{kna}{k-1} \left(\frac{b}{a} - \frac{\left(\frac{b}{a}\right)^k}{\sum_{t=0}^{k-1} \left(\frac{b}{a}\right)^t} \right) \geq \frac{kna}{k-1} \left(1 - \frac{1}{k} \right) = na. \end{aligned}$$

Hence, we have

$$g(x_0) \leq \frac{1}{n^k a^k} \frac{na^k}{k-1} \sum_{t=1}^{k-1} \left(\frac{b}{a}\right)^t = \frac{1}{(k-1)n^{k-1}} \sum_{t=1}^{k-1} \left(\frac{b}{a}\right)^t.$$

\square

Lemma 4.6. *Under the assumption (12), if*

$$A_{\mathcal{V}} = (1 + o_P(1))\mathbb{E}_0[A_{\mathcal{V}}],$$

$$A_{D, D^c} = (1 + o_P(1))\mathbb{E}_0[A_{D, D^c}],$$

uniformly for all $D \subset \mathcal{V}$ with $n^{\frac{1}{m+1}} \leq |D| \leq n$, then

$$\widehat{p}_{D, m} = (1 + o_p(1))\mathbb{E}_0[A_D],$$

uniformly for all $D \subset \mathcal{V}$ with $n^{\frac{1}{m+1}} \leq |D| \leq n$. Under H_1 , the above result still holds.

Proof of Lemma 4.4. Define a function $f(x, y) = \left(x^{\frac{1}{m}} - (x - 2y)^{\frac{1}{m}}\right)^m$ for $x \geq 2y$. At a fixed point (a, b) , the Taylor expansion is

$$f(x, y) = f(a, b) + \frac{\partial f}{\partial x}(x^*, y^*)(x - a) + \frac{\partial f}{\partial y}(x^*, y^*)(y - b), \quad (17)$$

where x^* is between a and x , y^* is between b and y and

$$\begin{aligned} \frac{\partial f}{\partial x}(x, y) &= \left(x^{\frac{1}{m}} - (x - 2y)^{\frac{1}{m}}\right)^{m-1} \left(x^{\frac{1-m}{m}} - (x - 2y)^{\frac{1-m}{m}}\right), \\ \frac{\partial f}{\partial x}(x, y) &= \frac{2 \left(x^{\frac{1}{m}} - (x - 2y)^{\frac{1}{m}}\right)^m}{\left(x^{\frac{1}{m}} - (x - 2y)^{\frac{1}{m}}\right) (x - 2y)^{1-\frac{1}{m}}}. \end{aligned}$$

Under the assumption (12), $\left(\frac{W_{max}}{W_{min}}\right)^m \ll \left(\frac{N}{n}\right)^{m-1-\delta_m} W_{min}^m \ll \left(\frac{N}{n}\right)^{m-1-\delta_m}$. Then

$$\frac{\mathbb{E}_0[A_{D, D^c}]}{\mathbb{E}_0[A_{\mathcal{V}}]} \leq \frac{\sum_{k=1}^t \binom{m}{2k-1} (N-n)^{2k-1} n^{m-2k+1} W_{max}^m}{N^m W_{min}^m} \ll \left(\frac{N}{n}\right)^{m-1-\delta_m} = o(1). \quad (18)$$

Hence, we get $f(\mathbb{E}_0[A_{\mathcal{V}}], \mathbb{E}_0[A_{D, D^c}]) = (1 + o(1))\mathbb{E}_0[A_{\mathcal{V}}]$ uniformly in D . For $(x, y) = (A_{\mathcal{V}}, A_{D, D^c})$ and $(a, b) = (\mathbb{E}_0[A_{\mathcal{V}}], \mathbb{E}_0[A_{D, D^c}])$, we also have

$$\frac{\partial f}{\partial x}(x^*, y^*) = 1 + o_p(1), \quad \frac{\partial f}{\partial y}(x^*, y^*) = 2 + o_p(1).$$

By (17), it's easy to check

$$\frac{2^m \widehat{p}_{D, m}}{\mathbb{E}_0[A_{\mathcal{V}}]} = \frac{f(A_{\mathcal{V}}, A_{D, D^c})}{f(\mathbb{E}_0[A_{\mathcal{V}}], \mathbb{E}_0[A_{D, D^c}])(1 + o(1))} = 1 + o_p(1).$$

It suffices to show $\mathbb{E}_0[A_{\mathcal{V}}] = 2^m \mathbb{E}_0[A_D]$. To this end, by (14) and (18), we only need to prove

$$\mathbb{E}_0[A_D] \gg \frac{1}{m!} \sum_{\substack{|\{i_1, i_2, \dots, i_m\}| \leq m-1 \\ \{i_1, \dots, i_m\} \subset D}} W_{i_1} \cdots W_{i_m}, \quad (19)$$

for any $|D| \geq n^{\frac{1}{m+1}}$. Note that

$$\mathbb{E}_0[A_D] = \frac{1}{m!} \left(\sum_{i \in D} W_i \right)^m - \frac{1}{m!} \sum_{\substack{|\{i_1, i_2, \dots, i_m\}| \leq m-1 \\ \{i_1, \dots, i_m\} \subset D}} W_{i_1} \cdots W_{i_m}.$$

Obviously, $\sum_{\substack{|\{i_1, i_2, \dots, i_m\}| \leq m-1 \\ \{i_1, \dots, i_m\} \subset D}} W_{i_1} \cdots W_{i_m}$ can be written as a summation of products of $\sum_{i \in D} W_i^k$, $k = 1, 2, \dots, m$. By Lemma 4.5 and assumption (12), for the term $\sum_{i \in D} W_i^k$ with $k \geq 2$, we have

$$\frac{\sum_{i \in D} W_i^k}{\left(\sum_{i \in D} W_i\right)^k} = O\left(\frac{1}{|D|^{k-1}} \left(\frac{W_{max}}{W_{min}}\right)^{k-1}\right) = O\left(\frac{o(n^{\frac{k-1}{m+1}})}{n^{\frac{k-1}{m+1}}}\right) = o(1).$$

Hence, (19) holds. Under H_1 , the result can be similarly proved. \square

References

- [1] Ahn, K., Lee, K. and Suh, C. (2016). Community recovery in hypergraphs, *54th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*. DOI: 10.1109/ALLERTON.2016.7852294.
- [2] Ahn, K., Lee, K. and Suh, C. (2018). Hypergraph Spectral Clustering in the Weighted Stochastic Block Model. *IEEE Journal of Selected Topics in Signal Processing* 12(5), 2018.
- [3] Angelini, M., Caltagirone, F., Krzakala, F. and Zdeborova, L. (2015). Spectral detection on sparse hypergraphs. *Allerton Conference on Communication, Control, and Computing*, 66-73.
- [4] Arias-Castro, E. and Verzelen, N. (2014). Community detection in dense random networks. *Annals of Statistics*, 42, 3: 940-969.
- [5] Bogerd, K., Castro, R., Hofstad, R. and Verzelen, N. (2019). Detecting a planted community in an inhomogeneous random graph. *Bernoulli*, **27**, 1159–1188.
- [6] Bi, X., Tang, X., Yuan, Y., Zhang, Y. and Qu, A. (2021). Tensors in statistics. *Annual Review of Statistics and Its Application*, 8.
- [7] Butucea, C. and Ingster, Y. (2013). Detection of a sparse submatrix of a high-dimensional noisy matrix. *Bernoulli*, 195B, 2652-2688.
- [8] Buló S. and Pelillo, M. (2009). New Bounds on the Clique Number of Graphs Based on Spectral Hypergraph Theory. *Learning and Intelligent Optimization. LION 2009*, 45-58.
- [9] Bolla, M. (1993). Spectra, euclidean representations and clusterings of hypergraphs. *Discrete Mathematics*, **117(1)**, 19-39.
- [10] Bollobás, B. and Erdős, P. (1976). Cliques in random graphs. *Mathematical Proceedings of the Cambridge Philosophical Society*, **80**, 419-427.
- [11] Bhangale, A., Gandhi, R. and Kortsarz, G. (2017). Improved approximation algorithm for the dense-3-subhypergraph problem. <https://arxiv.org/abs/1704.08620>
- [12] Chertok, M. and Keller, Y. (2010). Efficient high order matching. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **32(12)**, 2205-2215.
- [13] Chlamtac, E., M. Dinitz, C. Konrad, G. Kortsarz, and G. Rabanca. The Densest k-Subhypergraph problem. *SIAM Journal on Discrete Mathematics*. 32 (2):1458–77.
- [14] Florescu, L. and Perkins, W. (2016). Spectral thresholds in the bipartite stochastic block model. *29th Annual Conference on Learning Theory*, 49: 943-959.
- [15] Ghoshdastidar, D. and Dukkipati, A. (2014). Consistency of spectral partitioning of uniform hypergraphs under planted partition model. *Advances in Neural Information Processing Systems (NIPS)*, 397-405.
- [16] Ghoshdastidar, D. and Dukkipati A. (2017). Consistency of spectral hypergraph partitioning under planted partition model. *The Annals of Statistics*, **45(1)**, 289-315.
- [17] Hall, P. and Heyde, C. C. (2014). *Martingale limit theory and its application*. Academic press.
- [18] Hu, S., Wu, X. and Chan, T-H. (2017). Maintaining densest subsets efficiently in evolving hypergraphs *Proceedings of the 2017 ACM on Conference on Information and Knowledge Management*, 929-938, Singapore.
- [19] Kim, S. 2011. Higher-order correlation clustering for image segmentation. *Advances in Neural Information Processing Systems* 1530–8.
- [20] Ke, Z., Shi, F. and Xia, D. (2020). Community Detection for Hypergraph Networks via Regularized Tensor Power Iteration. <https://arxiv.org/pdf/1909.06503.pdf>.

- [21] Kim, C., Bandeira, A. and Goemans, M. (2017). Community detection in hypergraphs, spiked tensor models, and sum-of-squares. *2017 International Conference on Sampling Theory and Applications (SampTA)*, 124-128.
- [22] Lin, C., Chien, I. and Wang, I. (2017). On the fundamental statistical limit of community detection in random hypergraphs. *Information Theory (ISIT), 2017 IEEE International Symposium*, 2178-2182.
- [23] Luo, Y. and Zhang, A. (2020). Open problem: average-case hardness of hypergraphic planted clique detection. *Proceedings of Machine Learning Research*, 1-4, 2020.
- [24] Liu, H., Jan, L. and Yan, S. (2015). Dense subgraph partition of positive hypergraphs. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **37**, 3:541-554.
- [25] Peng, X. (2016). The Ramsey number of generalized loose paths in hypergraphs. *Discrete Mathematics* 339 (2):539-46.
- [26] Rodriguez, J. A. (2009). Laplacian eigenvalues and partition problems in hypergraphs. *Applied Mathematics Letters*, **22(6)**, 916-921.
- [27] Rota Bulò, S. and Pelillo, M. (2013). A game-theoretic approach to hypergraph clustering. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, **35(6)**, 1312-1327.
- [28] Soderberg, B. (2002). General formalism for inhomogeneous random graphs. *Phys. Rev. E*. 66,066121.
- [29] Verzelen, N., Arias-Castro, E. (2015). Community detection in sparse random networks. *Ann. Appl. Probab.*, **25**, 6:3465-3510.
- [30] Yuan, M., Liu, R., Feng, Y. and Shang, Z. (2018). Testing community structures for hypergraphs. arXiv preprint [arXiv:1810.04617](https://arxiv.org/abs/1810.04617).
- [31] Yuan, M. and Shang, Z. (2021). Sharp detection boundaries on testing dense subhypergraph. <https://arxiv.org/abs/2101.04584>
- [32] Zhou, D., Huang, J. and Scholkopf, B. (2006). Learning with hyper graphs: Clustering, classification, and embedding. *Advances in Neural Information Processing Systems*. 6:1601-8.