

# Dynamic Properties of Two-Dimensional Latticed Holographic System

Peng Liu <sup>1\*</sup> and Jian-Pin Wu <sup>2,3†</sup>

<sup>1</sup> *Department of Physics and Siyuan Laboratory,  
Jinan University, Guangzhou 510632, P.R. China*

<sup>2</sup> *Center for Gravitation and Cosmology,  
College of Physical Science and Technology,  
Yangzhou University, Yangzhou 225009, China*

<sup>3</sup> *School of Aeronautics and Astronautics,  
Shanghai Jiao Tong University, Shanghai 200240, China*

## Abstract

We study the anisotropic properties of dynamical quantities: direct current (DC) conductivity and butterfly velocity. The anisotropy plays a crucial role in determining the phase structure of the two-lattice system. Even a small deviation from isotropy can lead to distinct phase structures, as well as the IR fixed points of our holographic systems. In particular, for anisotropic cases, the most important property is that the IR fixed point can be non- $\text{AdS}_2 \times \mathbb{R}^2$  even for metallic phases. As that of a one-lattice system, the butterfly velocity can also diagnose the quantum phase transition (QPT) in this two-dimensional anisotropic latticed system.

---

\*Electronic address: [phylp@email.jnu.edu.cn](mailto:phylp@email.jnu.edu.cn)

†Electronic address: [jianpinwu@yzu.edu.cn](mailto:jianpinwu@yzu.edu.cn)

## Contents

<b>I. Introduction</b>	2
<b>II. Holographic Q-lattice model</b>	4
<b>III. Anisotropy effect on dynamical quantities</b>	5
A. The DC conductivity	5
B. The butterfly velocity and QPT	9
C. The scaling behavior of the butterfly	11
<b>IV. Discussion</b>	16
<b>Acknowledgments</b>	17
<b>References</b>	17

## I. INTRODUCTION

Anisotropy is universal and results in many rich phenomena in nature, such as magnetic systems, latticed systems, and so on [1]. In some strongly correlated systems, the anisotropy is associated with many dynamical measures, for example, the transport properties. The effects of anisotropy on dynamical measures for strongly correlated systems are useful for practical uses and are worthy of further investigation. Strongly correlated systems, however, are long-standing hard problems in physics. Gauge/gravity duality can bring novel insight and understanding into the strongly correlated systems and offer a good platform to study the anisotropy of dynamical quantities in strongly correlated systems.

Gauge/gravity duality has been proved powerful tools to study strongly correlated systems and dynamical properties [2–6]. Anisotropy is also ubiquitous in holographic systems, such as systems with lattices, anisotropic axions, massive gravity, and so on [7–10]. All these models realize the anisotropy by explicitly breaking the isotropic symmetry. The anisotropy can also be introduced by spontaneous symmetry breaking, such as holographic charge density wave models [11, 12].

Due to the breaking of translational symmetry from lattices, the DC conductivity is

finite. In [13], a novel technique to calculate the DC conductivity from lattice systems has been developed. Using these techniques, a simple analytic expression for the DC conductivity of a large class of holographic lattice models can be obtained, see [14–16] and therein. In particular, in terms of the relation between the DC conductivity and the temperature, it is convenient to work out the (metal or insulating) phase of the system, see for instances [17–27].

The butterfly effect ubiquitously exists in holographic systems and is depicted by a shock-wave solution of a black hole near the horizon [28–32]. Since the Lyapunov exponent in a holographic system saturates the bound of chaos [33], the butterfly velocity is an important dynamical quantity to signal some characteristics of a holographic system. In particular, the butterfly velocity is proposed as the characteristic quantity of a holographic system [32, 34–36]. In addition, it has been found that the butterfly velocity can characterize the phase transition [24, 37].

This leads to three interesting possibilities: the system is a metallic phase in any direction, the system is a metallic phase in one direction and an insulating phase in another direction, and the system is an insulating phase in any direction. We are particularly concerned about the quantum phase transitions corresponding to these three different states and their relationship with anisotropy.

In this paper, we want to study DC conductivity and butterfly velocity over an anisotropic background from Q-lattice. Most of the previous work only studied anisotropy with the one-dimensional lattice. Here, we shall derive the expressions of the DC conductivity and the butterfly velocity in the holographic system with the two-dimensional lattice. And it would be crucial to check, if those behaviors in the holographic system with one-dimensional lattice are still valid for two-dimensional cases. In this paper, we shall especially examine that how MIT is affected by the anisotropy and to what extent the characterization of the phase transition by butterfly velocity is universal. We organize this paper into three parts: we introduce the 2-dimensional Q-lattice model in II; then we study the anisotropic dynamical quantities, including the DC conductivity and the butterfly velocity, and the related properties, in III; we conclude and discuss in IV.

## II. HOLOGRAPHIC Q-LATTICE MODEL

The holographic Q-lattice model is a concise realization of the periodic structure. Previous holographic lattice models, such as the ionic lattices model and the scalar lattices model, introduce spatially periodic structures on scalar fields or the chemical potential (see [7] for a recent review). The resultant equations of motion are a set of highly nonlinear partial differential equations, which pose a challenge for numerical solutions. By contrast, the Q-lattice model introduces a complex scalar field, which results in only ordinary differential equations. Therefore, the Q-lattice model is an easier realization of lattice structures. The Q-lattice model is useful in modeling the Mott insulator and MIT [21, 22].

The Lagrangian of the Q-lattice model is [13, 20, 38, 39],

$$\mathcal{L} = R + 6 - \frac{1}{2}F^{ab}F_{ab} + \partial_a\Phi_1^\dagger\partial^a\Phi_1 + \partial_a\Phi_2^\dagger\partial^a\Phi_2 + m_1^2|\Phi_1|^2 + m_2^2|\Phi_2|^2. \quad (1)$$

System (1) can be solved with ansatz,

$$ds^2 = \frac{1}{z^2} \left[ -(1-z)p(z)U dt^2 + \frac{dz^2}{(1-z)p(z)U} + V_1 dx^2 + V_2 dy^2 \right], \quad (2)$$

$$A = \mu(1-z)adt, \quad \Phi_1 = e^{i\tilde{k}_1x} z^{3-\Delta_1}\phi_1, \quad \Phi_2 = e^{i\tilde{k}_2y} z^{3-\Delta_2}\phi_2$$

where  $p(z) = 1 + z + z^2 - \mu^2 z^3/2$  and  $\Delta_{1,2} = 3/2 \pm (9/4 + m_{1,2}^2)^{1/2}$ . For concreteness, we set  $m_1^2 = m_2^2 = m^2 = -2$ , therefore we have  $\Delta_1 = \Delta_2 = \Delta = 2$ . The horizon and the boundary locate at  $z = 1$  and  $z = 0$  respectively. The  $A_a$  is the Maxwell field, and  $\phi_{1,2}$  is the complex scalar field mimicking the lattice structures. Consequently, the functions to solve are  $(a, \phi, U, V_1, V_2)$ .

To solve the system (1), we need to specify the boundary conditions and system parameters. We set  $a(0) = 1$ , then  $A_t(0) = \mu$  becomes the chemical potential of the dual system. The boundary condition  $\tilde{\lambda}_{1,2} = \phi_{1,2}(0)$  is the strength of the lattice deformation, and  $\tilde{k}_{1,2}$  is the wave vector of the periodic structure. The asymptotic AdS<sub>4</sub> requires that  $U(0) = 1, V_1(0) = 1, V_2(0) = 1$ . Other boundary conditions at the horizon ( $z = 1$ ) can be fixed by regularity. The Hawking temperature reads  $\tilde{T} = (6 - \mu^2)U(1)/(8\pi)$ . The black brane solutions can be categorized by 3 dimensionless parameters  $(\tilde{T}, \lambda_1, \lambda_2, k_1, k_2) \equiv \left( \frac{\tilde{T}}{\mu}, \frac{\tilde{\lambda}_1}{\mu}, \frac{\tilde{\lambda}_2}{\mu}, \frac{\tilde{k}_1}{\mu}, \frac{\tilde{k}_2}{\mu} \right)$ , where we adopt the chemical potential  $\mu$  as the scaling unit.

### III. ANISOTROPY EFFECT ON DYNAMICAL QUANTITIES

#### A. The DC conductivity

The DC conductivity  $\sigma_{\text{DC}}$  can be obtained with horizon data. For an anisotropic background, the DC conductivity is anisotropic as well. Previous studies on anisotropy focused only on two distinct directions. To reveal more detailed anisotropic behavior of the DC conductivity, we study the DC conductivity along an arbitrary direction, i.e., the anisotropic DC conductivity.

In Fig. 1, we can find that the upper left plot and the upper right plot correspond to insulating phase and metallic phases, respectively. For the lower row plots, the system is insulating in  $x$ -direction, while metallic in  $y$ -direction. Also, the intersection point in both plots of the lower row suggests that  $\sigma_x = \sigma_y$  at certain temperature, which results in isotropic  $\sigma$ .

To obtain the anisotropic DC conductivity, we study the linear responses behaviors of the current

$$J_i = R_{ij}E_j, \quad (3)$$

where  $J_i$ ,  $E_j$ ,  $R_{ij}$  are current, electric field and response matrix, respectively. Following the method in [13] (also see [14, 18]), we can calculate the DC conductivity  $\sigma_{x\text{DC}}, \sigma_{y\text{DC}}$  along the  $x, y$  direction, respectively

$$\sigma_{x\text{DC}} = R_{11} = \left( \sqrt{\frac{V_2}{V_1}} + \frac{\mu^2 a^2 \sqrt{V_1 V_2}}{k_1^2 \phi_1^2} \right) \Big|_{z=1}, \quad (4)$$

$$\sigma_{y\text{DC}} = R_{22} = \left( \sqrt{\frac{V_1}{V_2}} + \frac{\mu^2 a^2 \sqrt{V_1 V_2}}{k_2^2 \phi_2^2} \right) \Big|_{z=1}, \quad (5)$$

with the off-diagonal components  $R_{12} = R_{21} = 0$ . Therefore the angle-dependent DC conductivity reads,

$$\begin{aligned} \sigma_{\text{DC}}(\theta) &= |J^a|/|E^a| = \sqrt{J^a J_a / (E^b E_b)} = \sqrt{R_{ij} E_j R_{ik} E_k / (E_s E_s)} \\ &= \sqrt{(R_{11}^2 E_1^2 + R_{22}^2 E_2^2) / (E_1^2 + E_2^2)} = \sqrt{\sigma_{x\text{DC}}^2 \cos^2 \theta + \sigma_{y\text{DC}}^2 \sin^2 \theta}. \end{aligned} \quad (6)$$

Once the DC conductivity is at hand, we can define the metallic phase as  $\sigma'_{\text{DC}}(T) < 0$  and the insulating phase as  $\sigma'_{\text{DC}}(T) > 0$ . From this definition, we can see that as long as the system is in the metallic phase in one direction, the system is always in the metallic phase in any direction except the direction perpendicular to it where the system is in insulating




FIG. 1:  $\sigma_{DC}$  as the function of  $T$  for different wave number and angle  $\theta$ . The intersection point is the place where  $\sigma_x = \sigma_y$ .

phase (See Fig. 2). Moreover, the insulating phase in any direction requires the system to be insulating phase along the direction of two lattices.

Now we study the DC conductivity as the function of the temperature  $T$  for different angles  $\theta$ . For simplicity, we take  $\lambda_1 = \lambda_2 = 2$  here and the anisotropy is realized by setting  $k_1 \neq k_2$ . We find that when the system is insulating or metallic along both  $x$  and  $y$  directions, the system is also insulating or metallic for arbitrary angles (see the plots above in FIG.1). However, if the system is insulating along  $x$  direction but metallic along  $y$  direction, some interesting phenomena emerge. We summarize the observations as follows.

- When  $\theta$  is small, the system is insulating as expected (see the plots below in FIG.1 and FIG.2).




FIG. 2:  $\sigma_{DC}$  as the function of  $T$  for  $\lambda_1 = \lambda_2 = 2$ ,  $k_1 = 0.03$  and  $k_2 = 1.2$  in the region of small  $\theta$ .

- As  $\theta$  increases, it is observed that the DC conductivity goes down as the temperature decreases, which seems to be an insulating state, but after reaching its minimum it turns to rise with the decrease of the temperature, which is metallic behavior. A similar phenomenon is also observed in the holographic doped Mott system studied in [21].
- When  $\theta$  becomes larger, the system becomes metallic, which is also as expected (see the plots below in FIG.1).

The above phenomena match the analysis on the metallic in an arbitrary direction in the previous paragraph.

Compared with the one-dimensional lattice system, the extra lattice can not only realize the phase transition in the other direction, but also affect the phase of the previous lattice system. Next, we shall study the MIT phase diagram. In our previous works [20], MIT phase diagram  $(\lambda_1, k_1)$  for  $k_2 = 0$  has been worked out. Here, we want to study the effect from the lattice along  $y$  direction on the phase diagram  $(\lambda_1, k_1)$ . So we plot the phase diagram  $(\lambda_1, k_1)$  for different  $k_2$  at  $T = 0.001$  (FIG.3). The properties are summarized as follows.

- For fixed  $\lambda_1$ , the state of system changes from metal to insulator as  $k_1$  decreases for different  $k_2$ . It is similar to that for  $k_2 = 0$ , which has been studied in [20].
- For  $k_2 \neq 0$ , with the increase of  $k_2$ , the critical line goes up, which means that the phase transition from metal to insulator happen at larger  $k_1$  ( $\lambda_1$  is fixed). It is because




FIG. 3: MIT phase diagram  $(\lambda_1, k_1)$  for  $\lambda_2 = 2$  and different  $k_2$  at  $T = 0.001$ . The phase above the critical line is the metallic phase and the one below the critical line is the insulating phase.

the phase transition from insulator to metal happens along  $y$  direction as  $k_2$  increases, and when  $k_2$  is further tuned larger, the system along  $y$  direction exhibits more obvious metallic behavior.

- $k_2 = 0$  sets the upper bound of the critical line in phase diagram  $(\lambda_1, k_1)$ . As  $k_2$  increases, this critical line goes up and approaches this upper bound.

To understand the above phenomena, we implement the mode analysis for  $\text{AdS}_2 \times \mathbb{R}^2$  on this two-lattice system.

$$\begin{aligned}
 U &= 6r^2 (1 + u_1 r^\delta), & V_1 &= v_{10} (1 + v_{11} r^\delta), & V_2 &= v_{20} (1 + v_{21} r^\delta), \\
 a &= 2\sqrt{3}r (1 + a_1 r^\delta), & \phi_1 &= e^{ik_1 x_1} \varphi_1 r^\delta, & \phi_2 &= e^{ik_2 x_1} \varphi_2 r^\delta.
 \end{aligned}
 \tag{7}$$

We find that the mode expansions of these two lattice fields are independent from each other. Specifically, we find the scaling dimensions of the two lattice fields read

$$\delta_{\varphi_1} = -\frac{1}{2} + \frac{1}{2\sqrt{3}} \sqrt{3 + 2m^2 + 2e^{-2v_{10}} k_1^2}, \quad \delta_{\varphi_2} = -\frac{1}{2} + \frac{1}{2\sqrt{3}} \sqrt{3 + 2m^2 + 2e^{-2v_{20}} k_2^2}.
 \tag{8}$$

Therefore, the lattice will deform away the  $\text{AdS}_2 \times \mathbb{R}^2$  as long as either of,

$$(e^{-v_{10}} k_1)^2 > -m^2, \quad (e^{-v_{20}} k_2)^2 > -m^2,
 \tag{9}$$

is satisfied.

From the (9) we can see that, for small values of  $k_2$  the IR fixed point is no longer  $\text{AdS}_2 \times \mathbb{R}^2$ , and hence it will lay significant effects on the phase structure of the first lattice. However, we also find that when  $k_2$  is large, the effect of the second lattice on the first lattice disappears. As long as  $k_2$  is large enough, the second lattice is irrelevant, and the system is in deep metallic phase. In the dual picture, large  $k_2$  means small lattice spacing. The electron can easily move between different sites, no localization happens, and the system will naturally behave more like a non-latticed system. Holographically, this behavior still asks for an analytical understanding.

## B. The butterfly velocity and QPT

In this section, we study the anisotropic butterfly velocity. In our previous works [24, 37], the anisotropic butterfly velocity has been explored in special case of  $\lambda_2 = 0$  and  $k_2 = 0$ . Here, we shall turn on the lattice along  $y$  direction and see how it affects the butterfly velocity.

In original coordinate  $(t, z, x, y)$  the butterfly velocity is anisotropic. Since the butterfly velocity only depends on the horizon data, therefore, one may implement a convenient coordinate transformation  $\tilde{y} = \sqrt{\frac{V_{yy}}{V_{xx}}}y$ , and in the new coordinate  $(t, z, x, \tilde{y})$ , the butterfly velocity  $v_B$  is isotropic again. When going back to the original coordinate, the angle-dependent butterfly velocity  $\mathbf{v}_B(\theta)$  can be obtained as,

$$\mathbf{v}_B(\theta) = v_B \sqrt{\cos^2(\theta) + \frac{V_{xx} \sin^2(\theta)}{V_{yy}}}. \quad (10)$$

From Eq.(10), we can infer that the period of  $\mathbf{v}_B(\theta)$  is  $\pi$  and  $\mathbf{v}_B(\pi/2 - \theta) = \mathbf{v}_B(\pi/2 + \theta)$  and so we shall only focus on the angle range  $\theta \in [0, \pi/2]$  here. We show  $\mathbf{v}_B(\theta)$  as the function of  $\theta$  at  $\lambda_1 = \lambda_2 = 2$ ,  $k_1 = 0.03$  with different  $k_2$  in FIG.4, from which we can see that for  $k_2 < k_1$ ,  $\mathbf{v}_B(\theta)$  monotonically increases as  $\theta$  increases (left plot in FIG.4), but for  $k_2 > k_1$ , it monotonically decreases as  $\theta$  increase (right plot in FIG.4)<sup>1</sup>. Therefore, we can conclude that the lattice suppresses the butterfly velocity. It is no question that when

---

<sup>1</sup> Without loss of generality, we shall fix  $\lambda_1 = \lambda_2 = 2$  in the subsequent study and only change  $k_1$  and  $k_2$  to implement the anisotropy geometry.




FIG. 4:  $v_B(\theta)$  as the function at  $k_1 = 0.03$  and  $T = 0.0001$  with different  $k_2$ .

$k_1 = k_2$ ,  $v_B(\theta)$  is a constant independent of  $\theta$  (green line in left plot in FIG.4). In addition, when  $k_2 \gg k_1$ ,  $v_B(\theta)$  also is approximately a constant (green line in right plot in FIG.4).




FIG. 5:  $\partial_{k_1} v_B$  as the function of  $k_1$  at  $T = 0.001$ . The red dashed line represents the position of the critical point while the green dashed line denotes the position of the local maximum of  $\partial_{k_1} v_B$ .

We also study the behavior of butterfly velocity  $v_B$  along  $x$  direction near the QPT. FIG.5 shows  $\partial_{k_1} v_B$  as the function of  $k_1$  at  $T = 0.001$  with different  $k_2$ . We find that even when the lattice effect along  $y$  direction is introduced,  $v_B$  along  $x$  direction can diagnose the QPT along  $x$  direction. It indicates there is a rapid change of  $v_B$  as well as the local extremes of  $\partial_k v_B$  near the QCPs.

The essence of the above phenomenon is that the occurrence of quantum phase transition is usually accompanied by the change of IR fixed point. The power-law relationship between butterfly velocity and temperature at zero temperature limit is directly determined by IR fixed point. Therefore, the first derivative of the butterfly velocity with the system parameters will show an obvious peak behavior when it crosses the fixed point.

Notice that the butterfly velocity presents a discontinuity in the first-order derivative in the process of holographic superconductivity phase transition [37]. It is worth noting that the phenomena in quantum phase transition are different from this, and the mechanisms behind them are also completely different. The superconductivity phase transition will be accompanied by condensation, and the background geometry of the system will certainly be modified. From the perturbation analysis, it can be seen that the order of the perturbation received by the metric is twice that of the condensation field. Therefore, it can be concluded that the butterfly velocity in [37] has an obvious scaling law at the critical point, and its critical exponent is 1. The MIT phase transition discussed in this paper is driven by the change of IR fixed point, and there is no emergence of the condensation. Therefore, the power-law dependence of butterfly velocity on temperature can well reflect this point.

Next, we want to study the behavior of  $v_B$  in zero-temperature limit.

### C. The scaling behavior of the butterfly

In Ref. [34], they deduce that metallic phases in one-dimensional Q-lattice model always correspond to the well-known  $\text{AdS}_2 \times \mathbb{R}^2$  IR geometry, on which  $v_B \sim T^{1/2}$ . And then, in our previous work [24], we find that for one-dimensional Q-lattice model, the scaling of  $v_B$  with temperature is  $v_B \sim T^\alpha$  in both metallic and insulating phases, but the exponent  $\alpha$  approaches 1 (in fact it is slightly larger than 1) in insulating phase, which is different from that in metallic phase. It indicates the theory flows to different IR fixed points for metallic and insulating phases, respectively.

We also assume that in two-dimensional Q-lattice model the scaling of  $v_B$  follows the same behavior as  $v_B \sim T^\alpha$ . In addition, we also assume  $V_{xx}/V_{yy} \sim T^\beta$ . Then,  $\mathbf{v}_B(\theta)$  becomes,

$$\mathbf{v}_B(\theta) \sim T^\alpha \sqrt{\cos^2(\theta) + \sin^2(\theta)T^\beta} \quad (11)$$




FIG. 6:  $T\mathbf{v}'_B/\mathbf{v}_B$  *v.s.*  $T$  for different phases over isotropy background ( $k_1 = k_2 = 1.5, 2$  corresponds to metallic phases and  $k_1 = k_2 = 0.1, 0.3, 0.5$  corresponds to insulating phases).

Therefore we have the temperature power exponent as

$$T\mathbf{v}'_B/\mathbf{v}_B \sim \frac{2\alpha \cos^2(\theta) + (2\alpha + \beta) \sin^2(\theta)T^\beta}{2(\cos^2(\theta) + \sin^2(\theta)T^\beta)}, \quad (12)$$

where  $\prime$  denotes the derivative with respect to the temperature  $T$ . Explicitly, we can work out the temperature power exponent for some special  $\theta$  as

$$T\mathbf{v}'_B/\mathbf{v}_B \sim \begin{cases} \alpha & \theta = 0 \\ \alpha + \frac{\beta}{2} & \theta = \pi/2, \\ \alpha + \frac{\beta T^\beta}{2T^\beta + 2} & \theta = \pi/4 \end{cases}, \quad (13)$$

Now, we begin to numerically study the behavior of  $\mathbf{v}_B$  in zero temperature limit. And then we also try to understand the corresponding behavior based on the above assumption. FIG.6 exhibits  $T\mathbf{v}'_B/\mathbf{v}_B$  as the function of  $T$  for different phases over isotropy background. We clearly see that for metallic phases,  $\alpha = 1/2$ . As expected it is the same as the metallic phase of one-dimensional Q-lattice model. This suggests that the metallic phases for the isotropic 2-lattice system are essentially the same as that of the 1-lattice system. This is as expected since adding an extra copy in a perpendicular direction shall not change the IR geometry too much. However, for insulating phases,  $\alpha$  approaches 1 but is slightly smaller than 1. It indicates that the insulating phase over isotropy two-dimensional Q-lattice model

flows a novel IR fixed point, which is different from the well-known  $\text{AdS}_2 \times \mathbb{R}^2$  IR geometry as expected, is also different from that of one-dimensional Q-lattice model where the  $\alpha$  approaches a fixed value slightly larger than unity. Given the mild difference for insulating phases, the insulating phases indeed converge to certain fixed values.




FIG. 7:  $T\mathbf{v}'_B/\mathbf{v}_B$  *v.s.*  $T$  over anisotropy background for  $k_1 = 1.5$  and  $k_2 = 1.5, 1.2, 1.1$ , respectively. The system is metallic phase in any direction.




FIG. 8:  $T\mathbf{v}'_B/\mathbf{v}_B$  *v.s.*  $T$  over anisotropy background for  $k_1 = 1.5$  and  $k_2 = 1$ . Notice that this system is still metallic phase in any direction.

Then we study the anisotropy two-dimensional Q-lattice model. The first case is that the system is a metallic phase in any direction. FIG.7 shows  $T\mathbf{v}'_B/\mathbf{v}_B$  *v.s.*  $T$  over anisotropy background for  $k_1 = 1.5$  and  $k_2 = 1.5, 1.2, 1.1$ , respectively. First, we see that for fixed

lattice parameter,  $T\mathbf{v}'_B/\mathbf{v}_B$  tends to converge to a fixed value down to ultra low temperature  $T = 10^{-7}$  for different angle  $\theta$ . This actually reflects a fact that  $V_{xx}/V_{yy}$  is regular in zero temperature limit when both directions are in metallic phase. The angular independence of the  $T\mathbf{v}'_B/\mathbf{v}_B$  means that  $\lim_{T \rightarrow 0} \beta \rightarrow 0$ , from (12) we find that the  $\mathbf{v}_B(\theta)$  converges to a fixed value along any direction. Secondly, for different lattice parameters,  $T\mathbf{v}'_B/\mathbf{v}_B$  tends to converge to a different value in the zero temperature limit, which indicates that the theory flows to different IR fixed points. This is a key difference from the case with a single lattice, where the metallic phases are shown related only to the  $\text{AdS}_2 \times \mathbb{R}^2$  IR fixed points.

However, when the anisotropy becomes larger (here, we take  $k_1 = 1.5$  and  $k_2 = 1$ ), we observe an novel phenomenon that, at least down to the  $T \sim 10^{-9}$ ,  $v_B$  does no longer follow the behavior  $v_B \sim T^\alpha$  (see FIG.8). Notice that this system is still metallic phase in any direction. On the other hand, we also observe that the  $Tv'_{Bx}/v_{Bx}$  is smaller than that of  $Tv'_{By}/v_{By}$ , while the  $Tv'_{B,\theta=\pi/4}/v_{B,\theta=\pi/4}$  converges to that of  $Tv'_{Bx}/v_{Bx}$ . This suggests that in the limit of zero temperature,  $\beta$  does not tend to vanishing and we conclude that  $\beta > 0$ .

Another anisotropic case is that the system is in metallic phase in  $x$  direction and insulating phase in  $y$  direction. The behaviors for this case are shown in FIG.9. We see that along  $x$  direction,  $T\mathbf{v}'_B/\mathbf{v}_B$  converges to a fixed value being smaller than unity. While along  $y$  direction,  $T\mathbf{v}'_B/\mathbf{v}_B$  converges to a fixed value being larger than unity. Notice that, the  $T\mathbf{v}'_{Bx}/\mathbf{v}_{Bx}$  is smaller than that of  $T\mathbf{v}'_{By}/\mathbf{v}_{By}$ , while the  $T\mathbf{v}'_{B,\theta=\pi/4}/\mathbf{v}_{B,\theta=\pi/4}$  seems converge to that of  $T\mathbf{v}'_{By}/\mathbf{v}_{By}$ . This situation is delicate, since  $T\mathbf{v}'_{Bx}/\mathbf{v}_{Bx} < \mathbf{v}'_{By}/\mathbf{v}_{By}$  suggests  $\beta > 0$ , while  $T\mathbf{v}'_{B,\theta=\pi/4}/\mathbf{v}_{B,\theta=\pi/4}$  converging to  $T\mathbf{v}'_{Bx}/\mathbf{v}_{Bx}$  suggests  $\beta < 0$ . This seemingly contradicted conclusion is on account of the temperature so far is not low enough. Closer examination of the zoom in inset in FIG.9 will reveal that,  $T\mathbf{v}'_{B,\theta=\pi/4}/\mathbf{v}_{B,\theta=\pi/4}$  starts to deviate from  $T\mathbf{v}'_{By}/\mathbf{v}_{By}$ , and it will converge to  $T\mathbf{v}'_{Bx}/\mathbf{v}_{Bx}$  in even lower temperature. In this case, we still have that  $\beta > 0$ .

Next, we elaborate on the anisotropic case where the system is insulating phase in any direction (FIG.10).  $T\mathbf{v}'_B/\mathbf{v}_B$  converges to different fixed value equaling to or being smaller than unity for different angle  $\theta$ . The  $T\mathbf{v}'_{Bx}/\mathbf{v}_{Bx}$  and  $T\mathbf{v}'_{By}/\mathbf{v}_{By}$  converge to slightly different values, while  $T\mathbf{v}'_{B,\theta=\pi/4}/\mathbf{v}_{B,\theta=\pi/4}$  converges to that of  $T\mathbf{v}'_{By}/\mathbf{v}_{By}$ . This means that  $V_{xx}/V_{yy} \sim T^\beta$  with  $\beta < 0$ , i.e., the  $\lim_{T \rightarrow 0} V_{xx}/V_{yy} \rightarrow \infty$ . This is different from the case where the system is in metallic phase in any direction.

Before closing this subsection, we would like to give some comments. First, according to




FIG. 9:  $T v'_B / v_B$  *v.s.*  $T$  over anisotropy background for  $k_1 = 0.5$  and  $k_2 = 0.1$ . The system is in metallic phase in  $x$  direction and insulating phase in  $y$  direction.




FIG. 10:  $T v'_B / v_B$  *v.s.*  $T$  over anisotropy background for  $k_1 = 0.1$  and  $k_2 = 0.01$ . The system is insulating phase in any direction.

the properties of the DC conductivity with the temperature, the anisotropic system can be categorized into three classes:

- Case 1: The system is metallic phase in any direction.
- Case 2: The system is in metallic phase in  $x$  direction and insulating phase in  $y$  direction.
- Case 3: The system is insulating phase in any direction.

Further, by using the scaling behavior of the butterfly, we can meticulously depict this anisotropic system by the parameter  $\beta$ .

- When the system is metallic phase in any direction (case 1),  $\beta = 0$  for mildly anisotropic lattice.
- When further increasing the anisotropy,  $\beta > 0$ . This phenomenon persists from case 1 to case 2.
- When the system is insulating phase in any direction,  $\beta < 0$ .

To summarize, for anisotropic cases, the most important property is that the IR fixed point can be non-AdS<sub>2</sub> × ℝ<sup>2</sup> even for metallic phases. Also, we observed some patterns of  $\alpha$  and  $\beta$  for different cases. This may help explain the bizarre IR fixed point.

#### IV. DISCUSSION

In this paper, we have studied the anisotropy effect on the dynamical quantities, including the DC conductivity and the butterfly velocity. The spatial anisotropy is realized in Q-lattice model by setting the wave number  $k_1 \neq k_2$ . We reveal the phase structure by DC conductivity. The anisotropy lays a significant effect on the phase structures, that we divide into three classes: metallic, metallic and insulating in different directions, insulating.

When the anisotropy is introduced, MIT is still universal. But MIT phase diagram along  $x$  direction is greatly affected by the lattice parameters along  $y$  direction. An interesting result is found that  $k_2 = 0$  sets the upper bound of the critical line in the phase diagram  $(\lambda_1, k_1)$ . As  $k_2$  increases, this critical line goes up and approaches this upper bound. That is to say, the phase structure is novel that one lattice can affect the other lattice. We provided several understandings on that, including mode analysis.

The butterfly velocity is also angle-dependent when the anisotropy is introduced. We further confirm the result that the lattice suppresses the butterfly velocity. It has been observed in our previous work [24], in which the lattice along  $y$  direction is turned off. In addition, we also confirm the robustness of the butterfly as the diagnostic of the QPT in an anisotropic system.

For anisotropic cases, the most important property is that the IR fixed point can be non-AdS<sub>2</sub> × ℝ<sup>2</sup> even for metallic phases. Also, we observed some patterns of  $\alpha$  and  $\beta$  for different cases. This may help explain the bizarre IR fixed point.

### Acknowledgments

Peng Liu would like to thank Yun-Ha Zha for her kind encouragement during this work. This work is supported by the Natural Science Foundation of China under Grant No. 11847055, 11805083, 11775036. J. P. Wu is also supported by Top Talent Support Program from Yangzhou University.

- 
- [1] Glotzer, Sharon C., and Michael J. Solomon. “Anisotropy of building blocks and their assembly into complex structures.” *Nature materials* 6.8 (2007): 557.
  - [2] J. M. Maldacena, “The Large N limit of superconformal field theories and supergravity,” *Adv. Theor. Math. Phys.* **2**, 231 (1998) [hep-th/9711200].
  - [3] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, “Gauge theory correlators from noncritical string theory,” *Phys. Lett. B* **428**, 105 (1998) [hep-th/9802109].
  - [4] E. Witten, “Anti-de Sitter space and holography,” *Adv. Theor. Math. Phys.* **2**, 253 (1998) [hep-th/9802150].
  - [5] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, “Large N field theories, string theory and gravity,” *Phys. Rept.* **323**, 183 (2000) [hep-th/9905111].
  - [6] S. A. Hartnoll, A. Lucas and S. Sachdev, “Holographic quantum matter,” arXiv:1612.07324 [hep-th].
  - [7] Y. Ling, “Holographic lattices and metal-insulator transition,” *Int. J. Mod. Phys. A* **30** (2015) no.28&29, 1545013.
  - [8] L. Q. Fang, X. H. Ge, J. P. Wu and H. Q. Leng, “Anisotropic Fermi surface from holography,” *Phys. Rev. D* **91**, no. 12, 126009 (2015) [arXiv:1409.6062 [hep-th]].
  - [9] I. Aref’eva and K. Rannu, “Holographic Anisotropic Background with Confinement-Deconfinement Phase Transition,” *JHEP* **1805**, 206 (2018) [arXiv:1802.05652 [hep-th]].
  - [10] H. S. Jeong, Y. Ahn, D. Ahn, C. Niu, W. J. Li and K. Y. Kim, “Thermal diffusivity and

- butterfly velocity in anisotropic Q-Lattice models,” *JHEP* **1801**, 140 (2018) [arXiv:1708.08822 [hep-th]].
- [11] A. Donos and J. P. Gauntlett, ”Holographic charge density waves,” *Phys. Rev. D* **87**, no. 12, 126008 (2013) doi:10.1103/PhysRevD.87.126008 [arXiv:1303.4398 [hep-th]].
- [12] Y. Ling, C. Niu, J. Wu, Z. Xian and H. b. Zhang, ”Metal-insulator Transition by Holographic Charge Density Waves,” *Phys. Rev. Lett.* **113**, 091602 (2014) [arXiv:1404.0777 [hep-th]].
- [13] A. Donos and J. P. Gauntlett, ”Novel metals and insulators from holography,” *JHEP* **1406**, 007 (2014) [arXiv:1401.5077 [hep-th]].
- [14] M. Blake and A. Donos, ”Quantum Critical Transport and the Hall Angle,” *Phys. Rev. Lett.* **114**, no. 2, 021601 (2015) [arXiv:1406.1659 [hep-th]].
- [15] A. Donos and J. P. Gauntlett, ”Thermoelectric DC conductivities from black hole horizons,” *JHEP* **1411**, 081 (2014) [arXiv:1406.4742 [hep-th]].
- [16] Z. Zhou, J. P. Wu and Y. Ling, ”DC and Hall conductivity in holographic massive Einstein-Maxwell-Dilaton gravity,” *JHEP* **1508**, 067 (2015) [arXiv:1504.00535 [hep-th]].
- [17] A. Donos, B. Goutřaux and E. Kiritsis, ”Holographic Metals and Insulators with Helical Symmetry,” *JHEP* **1409**, 038 (2014) [arXiv:1406.6351 [hep-th]].
- [18] Y. Ling, P. Liu, J. P. Wu and Z. Zhou, ”Holographic Metal-Insulator Transition in Higher Derivative Gravity,” *Phys. Lett. B* **766**, 41 (2017) [arXiv:1606.07866 [hep-th]].
- [19] J. P. Wu, ”Transport phenomena and Weyl correction in effective holographic theory of momentum dissipation,” *Eur. Phys. J. C* **78**, no. 4, 292 (2018) [arXiv:1902.03225 [hep-th]].
- [20] Y. Ling, P. Liu, C. Niu, J. P. Wu and Z. Y. Xian, ”Holographic Entanglement Entropy Close to Quantum Phase Transitions,” *JHEP* **1604**, 114 (2016) [arXiv:1502.03661 [hep-th]].
- [21] Y. Ling, P. Liu, C. Niu and J. P. Wu, ”Building a doped Mott system by holography,” *Phys. Rev. D* **92**, no. 8, 086003 (2015) [arXiv:1507.02514 [hep-th]].
- [22] Y. Ling, P. Liu and J. P. Wu, ”A novel insulator by holographic Q-lattices,” *JHEP* **1602**, 075 (2016) [arXiv:1510.05456 [hep-th]].
- [23] Y. Ling, P. Liu and J. P. Wu, ”Characterization of Quantum Phase Transition using Holographic Entanglement Entropy,” *Phys. Rev. D* **93**, no. 12, 126004 (2016) [arXiv:1604.04857 [hep-th]].
- [24] Y. Ling, P. Liu and J. P. Wu, ”Holographic Butterfly Effect at Quantum Critical Points,” *JHEP* **1710**, 025 (2017) [arXiv:1610.02669 [hep-th]].

- [25] M. Baggioli and O. Pujolas, “Electron-Phonon Interactions, Metal-Insulator Transitions, and Holographic Massive Gravity,” *Phys. Rev. Lett.* **114**, no. 25, 251602 (2015) [arXiv:1411.1003 [hep-th]].
- [26] M. Baggioli and O. Pujolas, “On holographic disorder-driven metal-insulator transitions,” *JHEP* **1701**, 040 (2017) [arXiv:1601.07897 [hep-th]].
- [27] M. Baggioli and O. Pujolas, “On Effective Holographic Mott Insulators,” *JHEP* **1612**, 107 (2016) [arXiv:1604.08915 [hep-th]].
- [28] D. A. Roberts and B. Swingle, “Lieb-Robinson Bound and the Butterfly Effect in Quantum Field Theories,” *Phys. Rev. Lett.* **117**, no. 9, 091602 (2016) [arXiv:1603.09298 [hep-th]].
- [29] S. H. Shenker and D. Stanford, “Black holes and the butterfly effect,” *JHEP* **1403**, 067 (2014) [arXiv:1306.0622 [hep-th]].
- [30] S. H. Shenker and D. Stanford, “Multiple Shocks,” *JHEP* **1412**, 046 (2014) [arXiv:1312.3296 [hep-th]].
- [31] S. H. Shenker and D. Stanford, “Stringy effects in scrambling,” *JHEP* **1505**, 132 (2015) [arXiv:1412.6087 [hep-th]].
- [32] M. Blake, “Universal Charge Diffusion and the Butterfly Effect in Holographic Theories,” *Phys. Rev. Lett.* **117**, no. 9, 091601 (2016) [arXiv:1603.08510 [hep-th]].
- [33] J. Maldacena, S. H. Shenker and D. Stanford, “A bound on chaos,” *JHEP* **1608**, 106 (2016) [arXiv:1503.01409 [hep-th]].
- [34] M. Blake, “Universal Diffusion in Incoherent Black Holes,” *Phys. Rev. D* **94**, no. 8, 086014 (2016) [arXiv:1604.01754 [hep-th]].
- [35] A. Lucas and J. Steinberg, “Charge diffusion and the butterfly effect in striped holographic matter,” *JHEP* **1610**, 143 (2016) [arXiv:1608.03286 [hep-th]].
- [36] S. A. Hartnoll, “Theory of universal incoherent metallic transport,” *Nature Phys.* **11**, 54 (2015) [arXiv:1405.3651 [cond-mat.str-el]].
- [37] Y. Ling, P. Liu and J. P. Wu, “Note on the butterfly effect in holographic superconductor models,” *Phys. Lett. B* **768**, 288 (2017) [arXiv:1610.07146 [hep-th]].
- [38] A. Donos and J. P. Gauntlett, “Holographic Q-lattices,” *JHEP* **1404**, 040 (2014) [arXiv:1311.3292 [hep-th]].
- [39] Y. Ling, P. Liu, C. Niu, J. P. Wu and Z. Y. Xian, “Holographic Superconductor on Q-lattice,” *JHEP* **1502**, 059 (2015) [arXiv:1410.6761 [hep-th]].