

Variations on the Post Correspondence Problem for free groups^{*}

Laura Ciobanu^[0000-0002-9451-1471] and Alan D. Logan^[0000-0003-1767-6798]

Heriot-Watt University, Edinburgh EH14 4AS, Scotland
{L.Ciobanu,A.Logan}@hw.ac.uk

Abstract. The Post Correspondence Problem is a classical decision problem about equalisers of free monoid homomorphisms. We prove connections between several variations of this classical problem, but in the setting of free groups and free group homomorphisms. Among other results, and working under certain injectivity assumptions, we prove that computing the rank of the equaliser of a pair of free group homomorphisms can be applied to computing a basis of this equaliser, and also to solve the “generalised” Post Correspondence Problem for free groups.

Keywords: Post Correspondence Problem, free group, rational constraint.

1 Introduction

In this article we connect several variations of the classical Post Correspondence Problem in the setting of free groups. The problems we consider have been open since the 1980s, and understanding how they relate and compare to their analogues in free monoids could bring us closer to their resolution. All problems are defined in Table 1, while their status in free groups and monoids is given in Table 2. However, three of these problems deserve proper introductions.

The first problem we consider is the Post Correspondence Problem (PCP) for free groups. This is completely analogous to the classical Post Correspondence Problem, which is about free monoids rather than free groups, and has had numerous applications in mathematics and computer science [11]. The PCP for other classes of groups has been successfully studied (see for example [17, Theorem 5.8]), but it remains open for free groups, where it is defined as follows. Let Σ and Δ be two alphabets, let $g, h : F(\Sigma) \rightarrow F(\Delta)$ be two group homomorphisms from the free group over Σ to the free group over Δ , and store this data in a four-tuple $I = (\Sigma, \Delta, g, h)$, called an *instance* of the PCP. The PCP is the decision problem:

Given $I = (\Sigma, \Delta, g, h)$, is there $x \in F(\Sigma) \setminus \{1\}$ such that $g(x) = h(x)$?

That is, if we consider the *equaliser* $\text{Eq}(g, h) = \{x \in F(\Sigma) \mid g(x) = h(x)\}$ of g and h , the PCP asks if $\text{Eq}(g, h)$ is non-trivial. Determining the decidability of this problem is an important question [6, Problem 5.1.4] [17, Section 1.4].

^{*} Supported by EPSRC grant EP/R035814/1

Our second problem asks not just about the triviality of $\text{Eq}(g, h)$, but for a finite description of it. We write PCP^{inj} (see Table 1) for the PCP with at least one map injective, in which case $\text{Eq}(g, h)$ is finitely generated [10] and a finite description relates to bases: The *Basis Problem* (BP) takes as input an instance $I = (\Sigma, \Delta, g, h)$ of the PCP^{inj} and outputs a basis for $\text{Eq}(g, h)$. In Section 7 we show that the BP is equivalent to the *Rank Problem* (RP), which seeks the number of elements in the basis, and was asked by Stallings in 1984. Recent results settle the BP for certain classes of free group maps [2–4, 8], but despite this progress its solubility remains open in general. The analogous problem for free monoids, which we call the *Algorithmic Equaliser Problem* (AEP) (see [4, page 2]) because it aims to describe the equaliser in terms of automata rather than bases, is insoluble [14, Theorem 5.2].

Our third problem is the *generalised Post Correspondence Problem* (GPCP), which is an important generalisation of the PCP for both free groups and monoids from 1982 [7]. For group homomorphisms $g, h : F(\Sigma) \rightarrow F(\Delta)$ and fixed elements u_1, u_2, v_1, v_2 of $F(\Delta)$, an instance of the GPCP is an 8-tuple $I_{\text{GPCP}} = (\Sigma, \Delta, g, h, u_1, u_2, v_1, v_2)$ and the GPCP itself is the decision problem:

Given $I_{\text{GPCP}} = (\Sigma, \Delta, g, h, u_1, u_2, v_1, v_2)$,
is there $x \in F(\Sigma) \setminus \{1\}$ such that $u_1g(x)u_2 = v_1h(x)v_2$?


Table 1: Summary of certain decision problems related to the PCP

Problems (for free groups)	Fixed: finite alphabets Σ and Δ and free groups $F(\Sigma), F(\Delta)$. Input: homomorphisms $g, h : F(\Sigma) \rightarrow F(\Delta)$ Additional input for GPCP: $u_1, u_2, v_1, v_2 \in F(\Delta)$ Additional input for $\text{PCP}_{\mathcal{R}}$: rational set $\mathcal{R} \subseteq F(\Sigma)$ Additional input for $\text{PCP}_{\mathcal{EL}}$: $a, b \in \Sigma^{\pm 1}, \Omega \subset \Sigma$
	Is it decidable whether:
PCP	there exists $x \in F(\Sigma) \setminus \{1\}$ s.t. $g(x) = h(x)$?
GPCP	there exists $x \in F(\Sigma) \setminus \{1\}$ s.t. $u_1g(x)u_2 = v_1h(x)v_2$?
$\text{PCP}_{\mathcal{R}}$	there exists $x \neq 1$ in \mathcal{R} s.t. $g(x) = h(x)$?
$\text{PCP}_{\mathcal{EL}}$	there exists $x \in F(\Sigma) \setminus \{1\}$ s.t. $g(x) = h(x)$ and x decomposes as a freely reduced word ayb for some $y \in F(\Omega)$
$\text{PCP}^{(\neg \text{inj}, \neg \text{inj})}$	PCP with neither g , nor h injective
$\text{PCP}^{(\neg \text{inj}, \text{inj})}$	PCP with exactly one of g, h injective
$\text{PCP}^{(\text{inj}, \text{inj})}$	PCP with both g, h injective
PCP^{inj}	$\text{PCP}^{(\neg \text{inj}, \text{inj})} \cup \text{PCP}^{(\text{inj}, \text{inj})}$ (i.e. PCP with at least one of g, h injective)
PCP^{CI}	PCP with g, h s.t. $g(y) \neq u^{-1}h(y)u$ for all $u \in F(\Delta), y \in F(\Sigma) \setminus \{1\}$
$\text{PCP}^{\text{inj} + \text{CI}}$	$\text{PCP}^{\text{inj}} \cup \text{PCP}^{\text{CI}}$
$\text{PCP}(n)$	$\text{PCP}(n)$ for alphabet size $ \Sigma = n$
GPCP variants	$\text{GPCP}^{(\neg \text{inj}, \neg \text{inj})}, \text{GPCP}^{\text{inj}}, \text{GPCP}^{\text{inj} + \text{CI}}, \text{GPCP}(n)$, etc. analogue to PCP variants

For free monoids, the PCP is equivalent to the GPCP [11, Theorem 8]. The corresponding connection for free groups is more complicated, and explaining this connection is the main motivation of this article. In particular, the GPCP for free groups is known to be undecidable [17, Corollary 4.2] but this proof does not imply that the PCP for free groups is undecidable (because of injectivity issues; see Section 2). In Theorem 4 we connect the PCP with the GPCP via a sequence of implications, and require at least one map to be injective.

Main theorem Theorem A summarises the connections proven in this paper (arrows are labeled by the section numbers where the implications are proven), and Section 8 brings all the results together. Note that asking for both maps to be injective refines the results in this theorem, as does restricting the size of the source alphabet Σ (see Theorem 9).

Theorem A (Theorem 9) *In finitely generated free groups the following implications hold.*



This theorem implies that Stallings' Rank Problem is of central importance, as we have the chain: $\text{RP} \Rightarrow \text{GPCP}^{\text{inj}} \Rightarrow \text{PCP}$.

Rational constraints The implication $\text{BP} \Rightarrow \text{GPCP}^{\text{inj}}$ above is surprising because the inputs to the problems are very different. The proof of this implication uses the PCP^{inj} with a certain rational constraint, namely the problem $\text{PCP}_{\mathcal{E}\mathcal{L}}^{\text{inj}}$ (see Table 1). The relationship between the GPCP and the $\text{PCP}_{\mathcal{E}\mathcal{L}}$ still holds if neither map is injective. As the GPCP for free groups is undecidable in general, this connection yields the following result.

Theorem B (Theorem 3) *There exists a rational constraint \mathcal{R} such that the $\text{PCP}_{\mathcal{R}}$ is undecidable.*

Random maps and generic behaviour. A different perspective on the PCP and its variations is to consider the behaviour of these problems when the pairs of homomorphisms are picked randomly (while the two alphabets $\Delta = \{x_1, \dots, x_m\}$ and $\Sigma = \{y_1, \dots, y_k\}$, and ambient free groups $F(\Delta)$ and $F(\Sigma)$ remain fixed). Any homomorphism is completely determined by how it acts on the generators, and so picking g and h randomly is to be interpreted as picking $(g(x_1), \dots, g(x_m))$ and $(h(x_1), \dots, h(x_m))$ as random tuples of words in $F(\Sigma)$ (see Section 6 for

details). There is a vast literature (see for example [13]) on the types of objects or behaviours which appear with probability 0, called *negligible*, or with probability 1, called *generic*, in infinite groups. In this spirit, the *generic* PCP refers to the PCP applied to a generic set (of pairs) of maps, that is, a set of measure 1 in the set of all (pairs of) maps, and we say that the generic PCP is decidable if the PCP is decidable for ‘almost all’ instances, that is, for a set of measure 1 of pairs of homomorphisms.

In Section 6 we describe the setup used to count pairs of maps and compute probabilities, and show that among all pairs of maps g, h , the property of being *conjugacy inequivalent* (that is, for every $u \in F(\Delta)$ there is no $x \neq 1$ in $F(\Sigma)$ such that $g(x) = u^{-1}h(x)u$; defined in Table 1 as PCP^{CI} and GPCP^{CI}) occurs with probability 1; that is, conjugacy inequivalent maps are generic:

Theorem C (Theorem 5) *Instances of the $\text{PCP}^{\text{inj} + \text{CI}}$ are generic instances of the PCP. That is, with probability 1, a pair of maps is conjugacy inequivalent.*

Theorem C shows that the implication $\text{PCP} \Rightarrow \text{GPCP}^{\text{inj} + \text{CI}}$ in Theorem A is the generic setting, and hence for ‘almost all maps’ we have $\text{PCP} \Leftrightarrow \text{GPCP}$.

We conclude the introduction with a summary of the status the PCP and its variations for free monoids and groups. We aim to study the computational complexity of these problems and how this complexity behaves with respect to the implications proved in this paper in future work.

Table 2: Status of results for free monoids and free groups

Problems	In free monoids	References for free monoids	In free groups	References for free groups
general PCP	undecidable	[19]	unknown	
general AEP / BP	undecidable	[14, Theorem 5.2]	unknown	
$\text{PCP}^{(\neg \text{inj}, \neg \text{inj})}$	undecidable	[19]	decidable	Lemma 1
PCP^{inj}	undecidable	[15]	unknown	
GPCP	undecidable	[11, Theorem 8]	undecidable	[17, Corollary 4.2]
$\text{GPCP}^{(\neg \text{inj}, \neg \text{inj})}$	undecidable	[11, Theorem 8]	undecidable	Lemma 2
GPCP^{inj}	undecidable	[15]	unknown	
$\text{GPCP}^{\text{inj} + \text{CI}}$	N/A		unknown	
$\text{PCP}_{\mathcal{R}}$	undecidable	[19]	undecidable	Theorem B
$\text{PCP}^{\text{inj} + \text{CI}}$	N/A		unknown	
generic PCP	decidable	[9, Theorem 4.4]	decidable	[5]

2 Non-injective maps

In this section we investigate the $\text{PCP}^{(\neg \text{inj}, \neg \text{inj})}$ and $\text{GPCP}^{(\neg \text{inj}, \neg \text{inj})}$, which are the PCP and GPCP under the assumption that both maps are non-injective.

The PCP for non-injective maps We first prove that the $\text{PCP}^{(\neg \text{inj}, \neg \text{inj})}$ is trivially decidable, with the answer always being “yes”.

Lemma 1. *If $g, h : F(\Sigma) \rightarrow F(\Delta)$ are both non-injective homomorphisms then $\text{Eq}(g, h)$ is non-trivial.*

Proof. We prove that $\ker(g) \cap \ker(h)$ is non-trivial, which is sufficient. Let $c \in \ker(g)$ and $d \in \ker(h)$ be non-trivial elements. If $\langle c, d \rangle \cong \mathbb{Z} = \langle x \rangle$, there exist integers k, l such that $c = x^k$ and $d = x^l$. Then $g(x^{kl}) = 1 = h(x^{kl})$ so $x^{kl} \in \ker(g) \cap \ker(h)$ with x^{kl} non-trivial, as required. If $\langle c, d \rangle \not\cong \mathbb{Z}$ then $g([c, d]) = 1 = h([c, d])$, so $[c, d] \in \ker(g) \cap \ker(h)$ with $[c, d]$ non-trivial, as required. \square

As we can algorithmically determine if a free group homomorphism is injective (e.g. via Stallings’ foldings), Lemma 1 gives us the following:

Proposition 1. $\text{PCP} \Leftrightarrow \text{PCP}^{(\neg \text{inj}, \neg \text{inj})}$

The GPCP for non-injective maps Myasnikov, Nikolaev and Ushakov defined the PCP and GPCP for general groups in [17]. Due to this more general setting their formulation is slightly different to ours but, from a decidability point of view, the problems are equivalent for free groups. They proved that the GPCP is undecidable for free groups; however, we now dig into their proof and observe that it assumes both maps are non-injective. Therefore, GPCP^{inj} remains open.

Lemma 2. *The $\text{GPCP}^{(\neg \text{inj}, \neg \text{inj})}$ is undecidable.*

Proof. Let H be a group with undecidable word problem and let $\langle \mathbf{x} \mid \mathbf{r} \rangle$ be a presentation of H . Set $\Delta := \mathbf{x}$, define

$$\Sigma := \{(x, 1) \mid x \in \mathbf{x}\} \cup \{(1, x^{-1}) \mid x \in \mathbf{x}\} \cup \{(R, 1) \mid x \in \mathbf{r}\} \cup \{(1, R^{-1}) \mid R \in \mathbf{r}\}$$

and define $g : (p, q) \mapsto p$ and $h : (p, q) \mapsto q$. Note that neither g nor h is injective, as if $R \in \mathbf{r}$ then $g(R, 1)$ may be realised as the image of a word over $\{(x, 1) \mid x \in \mathbf{x}\} \cup \{(1, x^{-1}) \mid x \in \mathbf{x}\}$, and analogously for $h(1, R)$. Taking $w \in F(\Delta)$, the instance $(\Sigma, \Delta, g, h, w, 1, 1, 1)$ of the $\text{GPCP}^{(\neg \text{inj}, \neg \text{inj})}$ has a solution if and only if the word w defines the identity of H [17, Proof of Proposition 4.1]. As H has undecidable word problem it follows that the $\text{GPCP}^{(\neg \text{inj}, \neg \text{inj})}$ is undecidable. \square

3 The GPCP and extreme-letter restrictions

In this section we connect the GPCP and the PCP under a certain rational constraint. This connection underlies Theorem B, as well as the implications $\text{BP} \Rightarrow \text{GPCP}^{\text{inj}}$ and $\text{PCP} \Rightarrow \text{GPCP}^{\text{inj} + \text{Cl}}$ in Theorem A.

Our results here, as in much of the rest of the paper, are broken down in terms of injectivity, and also alphabet sizes; understanding for which sizes of

alphabet Σ the classical Post Correspondence Problem is decidable/undecidable is an important research theme [7, 11, 18].

For an alphabet Σ , let Σ^{-1} be the set of formal inverses of Σ , and write $\Sigma^{\pm 1} = \Sigma \cup \Sigma^{-1}$. For example, if $\Sigma = \{a, b\}$ then $\Sigma^{\pm 1} = \{a, b, a^{-1}, b^{-1}\}$.

The *extreme-letter-restricted Post Correspondence Problem for free groups* $\text{PCP}_{\mathcal{E}\mathcal{L}}$ (see also Table 1) is the following problem: Let $g, h : F(\Sigma) \rightarrow F(\Delta)$ be two group homomorphisms, $\Omega \subset \Sigma$ a set, and $a, b \in \Sigma^{\pm 1}$ two letters; an instance of the $\text{PCP}_{\mathcal{E}\mathcal{L}}$ is a 6-tuple $I_{\text{PCP}_{\mathcal{E}\mathcal{L}}} = (\Sigma, \Delta, g, h, a, \Omega, b)$ and the $\text{PCP}_{\mathcal{E}\mathcal{L}}$ itself is:

- Given $I_{\text{PCP}_{\mathcal{E}\mathcal{L}}} = (\Sigma, \Delta, g, h, a, \Omega, b)$, is there $x \in F(\Sigma) \setminus \{1\}$ such that:
- $g(x) = h(x)$, and
 - x decomposes as a freely reduced word ayb for some $y \in F(\Omega)$?

Connecting the GPCP and the PCP. We start with an instance $I_{\text{GPCP}} = (\Sigma, \Delta, g, h, u_1, u_2, v_1, v_2)$ of the GPCP and consider the instance

$$I_{\text{PCP}} = (\Sigma \sqcup \{B, E\}, \Delta \sqcup \{B, E, \#\}, g', h')$$

of the PCP, where g' and h' are defined as follows.

$$g'(z) := \begin{cases} \#^{-1}g(z)\# & \text{if } z \in \Sigma \\ B\#u_1\# & \text{if } z = B \\ \#^{-1}u_2\#E & \text{if } z = E \end{cases} \quad h'(z) := \begin{cases} \#h(z)\#^{-1} & \text{if } z \in \Sigma \\ B\#v_1\#^{-1} & \text{if } z = B \\ \#v_2\#E & \text{if } z = E \end{cases}$$

Injectivity is preserved through this construction since $\text{rk}(\text{Im}(g)) + 2 = \text{rk}(\text{Im}(g'))$: this can be seen via Stallings' foldings, or directly by noting that the image of g' restricted to $F(\Sigma)$ is isomorphic to $\text{Im}(g)$, that B only occurs in $g'(d)$ and E only occurs in $g'(e)$. Analogously, h' is an injection if and only if h is, as again $\text{rk}(\text{Im}(h)) + 2 = \text{rk}(\text{Im}(h'))$. Thus we get:

Lemma 3. *The map g' is injective if and only if g is, and the map h' is injective if and only if h is.*

We now connect the solutions of I_{GPCP} to those of I_{PCP} .

Lemma 4. *A word $y \in F(\Sigma)$ is a solution to I_{GPCP} if and only if the word ' ByE ' is a solution to I_{PCP} .*

Proof. Starting with y being a solution to I_{GPCP} , we obtain the following sequence of equivalent identities:

$$\begin{aligned} u_1g(y)u_2 &= v_1h(y)v_2 \\ B\#(u_1g(y)u_2)\#E &= B\#(v_1h(y)v_2)\#E \\ B\#u_1\# \cdot \#^{-1}g(y)\# \cdot \#^{-1}u_2\#E &= B\#v_1\#^{-1} \cdot \#h(y)\#^{-1} \cdot \#v_2\#E \\ g'(B)g'(y)g'(E) &= h'(B)h'(y)h'(E) \\ g'(ByE) &= h'(ByE). \end{aligned}$$

Therefore ByE is a solution to I_{PCP} , so the claimed equivalence follows. \square

Theorem 1 includes conditions on injectivity and on alphabet sizes; see Table 1 for definitions.

Theorem 1. *The following hold in a finitely generated free group.*

1. $\text{PCP}_{\mathcal{E}\mathcal{L}}^{(\neg\text{inj},\text{inj})}(n+2) \Rightarrow \text{GPCP}^{(\neg\text{inj},\text{inj})}(n)$
2. $\text{PCP}_{\mathcal{E}\mathcal{L}}^{(\text{inj},\text{inj})}(n+2) \Rightarrow \text{GPCP}^{(\text{inj},\text{inj})}(n)$

Proof. Let I_{GPCP} be an instance of the GPCP^{inj} , and construct from it the instance $I_{\text{PCP}_{\mathcal{E}\mathcal{L}}} = (\Sigma \sqcup \{B, E\}, \Delta \sqcup \{B, E, \#\}, g', h', B, \Sigma, E)$ of the $\text{PCP}_{\mathcal{E}\mathcal{L}}$, which is the instance I_{PCP} defined above under the constraint that solutions have the form ByE for some $y \in F(\Sigma)$.

By Lemma 3, $I_{\text{PCP}_{\mathcal{E}\mathcal{L}}}$ is an instance of the $\text{PCP}_{\mathcal{E}\mathcal{L}}^{(\neg\text{inj},\text{inj})}(n+2)$ if and only if I_{GPCP} is an instance of $\text{GPCP}^{(\neg\text{inj},\text{inj})}(n)$, and similarly for $\text{PCP}_{\mathcal{E}\mathcal{L}}^{(\text{inj},\text{inj})}(n+2)$ and $\text{GPCP}^{(\text{inj},\text{inj})}(n)$. The result then follows from Lemma 4. \square

The above does not prove that $\text{PCP}^{\text{inj}} \Leftrightarrow \text{GPCP}^{\text{inj}}$, because I_{PCP} might have solutions of the form BxB^{-1} or $E^{-1}xE$. For example, if we let $I_{\text{GPCP}} = (\{a\}, \{a, c, d\}, g, h, c, \epsilon, \epsilon, d)$ with $g(a) = a$ and $h(a) = cac^{-1}$, then there is no $x \in F(a)$ such that $cg(a) = h(a)d$, but defining g', h' as above then $BaB^{-1} \in \text{Eq}(g', h')$. In Section 6.2 we consider maps where such solutions are impossible, and there the equivalence $\text{PCP}^{\text{inj}} \Leftrightarrow \text{GPCP}^{\text{inj}}$ does hold.

4 The PCP under rational constraints

For an alphabet A , a language $L \subseteq A^*$ is *regular* if there exists some finite state automaton over A which accepts exactly the words in L . Let $\pi : (\Sigma^{\pm 1})^* \rightarrow F(\Sigma)$ be the natural projection map. A subset $R \subseteq F(\Sigma)$ is *rational* if $R = \pi(L)$ for some regular language $L \subseteq (\Sigma^{\pm 1})^*$.

In this section we consider the $\text{PCP}_{\mathcal{R}}^{\text{inj}}$, which is the PCP^{inj} subject to the rational constraint \mathcal{R} . We prove that the $\text{PCP}_{\mathcal{R}}^{\text{inj}}$ under any rational constraint \mathcal{R} can be solved via the Basis Problem (BP) (so $\text{BP} \Rightarrow \text{PCP}_{\mathcal{R}}^{\text{inj}}$ from Theorem A). We later apply this to prove $\text{BP} \Rightarrow \text{GPCP}^{\text{inj}}$ from Theorem A, as the $\text{PCP}_{\mathcal{E}\mathcal{L}}^{\text{inj}}$ from Section 3 is simply the PCP^{inj} under a specific rational constraint.

Theorem 2. *The following hold in a finitely generated free group.*

1. $\text{BP}^{(\neg\text{inj},\text{inj})}(n) \Rightarrow \text{PCP}_{\mathcal{R}}^{(\neg\text{inj},\text{inj})}(n)$
2. $\text{BP}^{(\text{inj},\text{inj})}(n) \Rightarrow \text{PCP}_{\mathcal{R}}^{(\text{inj},\text{inj})}(n)$

Proof. Let g, h be homomorphisms from $F(\Sigma)$ to $F(\Delta)$ such that at least one of them is injective. Their equaliser $\text{Eq}(g, h)$ is a finitely generated subgroup of $F(\Sigma)$ [10], so $\text{Eq}(g, h)$ is a rational set (see for example Section 3.1 in [1]).

As the Basis Problem is soluble, we can compute a basis for $\text{Eq}(g, h)$. This is equivalent to finding a finite state automaton \mathcal{A} (called a “core graph” in the literature on free groups; see [12]) that accepts the set $\text{Eq}(g, h)$.

Let \mathcal{R} be a rational set in $F(\Sigma)$. The $\text{PCP}_{\mathcal{R}}$ for g and h is equivalent to determining if there exists any non-trivial $x \in \mathcal{R} \cap \text{Eq}(g, h)$. Since the intersection of two rational sets is rational, and an automaton recognising this intersection is computable by the standard product construction of automata, one can determine whether $\mathcal{R} \cap \text{Eq}(g, h)$ is trivial or not, and thus solve $\text{PCP}_{\mathcal{R}}$. \square

The following theorem immediately implies Theorem B, as restrictions on the maps do not affect the rational constraints.

Theorem 3 (Theorem B). $\text{PCP}_{\mathcal{E}\mathcal{L}}^{(\neg \text{inj}, \neg \text{inj})}$ is undecidable in free groups.

Proof. We have $\text{PCP}_{\mathcal{E}\mathcal{L}}^{(\neg \text{inj}, \neg \text{inj})} \Rightarrow \text{GPCP}^{(\neg \text{inj}, \neg \text{inj})}$ by Lemmas 3 and 4. The result follows as $\text{GPCP}^{(\neg \text{inj}, \neg \text{inj})}$ is undecidable by Lemma 2. \square

5 Main results, part 1

Here we combine results from the previous sections to prove certain of the implications in Theorem A. The implications we prove refine Theorem A, as they additionally contain information on alphabet sizes and on injectivity.

Theorem 4. *The following hold in finitely generated free groups.*

1. $\text{BP}^{(\neg \text{inj}, \text{inj})}(n+2) \Rightarrow \text{GPCP}^{(\neg \text{inj}, \text{inj})}(n) \Rightarrow \text{PCP}^{(\neg \text{inj}, \text{inj})}(n)$
2. $\text{BP}^{(\text{inj}, \text{inj})}(n+2) \Rightarrow \text{GPCP}^{(\text{inj}, \text{inj})}(n) \Rightarrow \text{PCP}^{(\text{inj}, \text{inj})}(n)$

Proof. As $\text{PCP}_{\mathcal{E}\mathcal{L}}^{(\neg \text{inj}, \text{inj})}$ is an instance of the PCP^{inj} under a rational constraint, Theorem 2 gives us that $\text{BP}^{(\neg \text{inj}, \text{inj})}(n+2) \Rightarrow \text{PCP}_{\mathcal{E}\mathcal{L}}^{(\neg \text{inj}, \text{inj})}(n+2)$, while Theorem 1 gives us that $\text{PCP}_{\mathcal{E}\mathcal{L}}^{(\neg \text{inj}, \text{inj})}(n+2) \Rightarrow \text{GPCP}^{(\neg \text{inj}, \text{inj})}(n)$, and the implication $\text{GPCP}^{(\neg \text{inj}, \text{inj})}(n) \Rightarrow \text{PCP}^{(\neg \text{inj}, \text{inj})}(n)$ is obvious as instances of the PCP are instances of the GPCP but with empty constants u_i, v_i . Sequence (1), with one map injective, therefore holds, while the proof of sequence (2) is identical. \square

Removing the injectivity assumptions gives the following corollary; the implications $\text{BP} \Rightarrow \text{GPCP}^{\text{inj}} \Rightarrow \text{PCP}$ of Theorem A follow immediately.

Corollary 1. $\text{BP}(n+2) \Rightarrow \text{GPCP}^{\text{inj}}(n) \Rightarrow \text{PCP}(n)$

Proof. Theorem 4 gives that $\text{BP}(n+2) \Rightarrow \text{GPCP}^{\text{inj}}(n) \Rightarrow \text{PCP}^{\text{inj}}(n)$, while $\text{PCP}(n) \Leftrightarrow \text{PCP}^{\text{inj}}(n)$ by Proposition 1. \square

6 Conjugacy inequivalent maps

In this section we prove genericity results and give conditions under which the PCP implies the GPCP. In particular, we prove Theorem C, and we prove the implication $\text{PCP} \Rightarrow \text{GPCP}^{\text{inj} + \text{CI}}$ from Theorem A.

A pair of maps $g, h : F(\Sigma) \rightarrow F(\Delta)$ is said to be *conjugacy inequivalent* if for every $u \in F(\Delta)$ there does not exist any non-trivial $x \in F(\Sigma)$ such that $g(x) =$

$u^{-1}h(x)u$ (see Table 1). For example, if the images of $g, h : F(\Sigma) \rightarrow F(\Delta)$ are *conjugacy separated*, that is, if $\text{Im}(g) \cap u^{-1} \text{Im}(h)u$ is trivial for all $u \in F(\Delta)$, then g and h are conjugacy inequivalent. We write $\text{PCP}^{\text{inj} + \text{CI}} / \text{GPCP}^{\text{inj} + \text{CI}}$ for those instances of the $\text{GPCP}^{\text{inj}} / \text{PCP}^{\text{inj}}$ where the maps are conjugacy inequivalent.

6.1 Random maps and genericity

Here we show that among all pairs of maps $g, h : F(\Sigma) \rightarrow F(\Delta)$, the property of being conjugacy inequivalent occurs with probability 1; that is, conjugacy inequivalent maps are *generic*.

Theorem 5 (Theorem C). *Instances of the $\text{PCP}^{\text{inj} + \text{CI}}$ are generic instances of the PCP. That is, with probability 1, a pair of maps is conjugacy inequivalent.*

Before we prove the theorem, we need to describe the way in which probabilities are computed. We fix the two alphabets $\Delta = \{x_1, \dots, x_m\}$ and $\Sigma = \{y_1, \dots, y_k\}$, $m \leq k$, and ambient free groups $F(\Delta)$ and $F(\Sigma)$, and pick g and h randomly by choosing $(g(x_1), \dots, g(x_m))$ and $(h(x_1), \dots, h(x_m))$ independently at random, as tuples of words of length bounded by n in $F(\Sigma)$. If \mathcal{P} is a property of tuples (or subgroups) of $F(\Sigma)$, we say that *generically many* tuples (or finitely generated subgroups) of $F(\Sigma)$ satisfy \mathcal{P} if the proportion of m -tuples of words of length $\leq n$ in $F(\Sigma)$ which satisfy \mathcal{P} (or generate a subgroup satisfying \mathcal{P}), among all possible m -tuples of words of length $\leq n$, tends to 1 when n tends to infinity.

Proof. Let $n > 0$ be an integer, and let (a_1, \dots, a_m) and (b_1, \dots, b_m) be two tuples of words in $F(\Sigma)$ satisfying length inequalities $|a_i| \leq n$ and $|b_i| \leq n$ for all i . We let the maps $g, h : F(\Sigma) \rightarrow F(\Delta)$ that are part of an instance of PCP be defined as $g(x_i) = a_i$ and $h(x_i) = b_i$, and note that the images $\text{Im}(g)$ and $\text{Im}(h)$ in $F(\Delta)$ are subgroups generated by (a_1, \dots, a_m) and (b_1, \dots, b_m) , respectively.

We claim that among all $2m$ -tuples $(a_1, \dots, a_m, b_1, \dots, b_m)$ with $|a_i|, |b_i| \leq n$, a proportion of them tending to 1 as $n \rightarrow \infty$ satisfy (1) the subgroups $L = \langle a_1, \dots, a_m \rangle$ and $K = \langle b_1, \dots, b_m \rangle$ are both of rank m , and (2) for every $u \in F(\Delta)$ we have $L^u \cap K = \{1\}$. Claim (1) is equivalent to g, h being generically injective, and follows from [16], while claim (2) is equivalent to $\text{Im}(g)^u \cap \text{Im}(h) = \{1\}$ for every $u \in F(\Delta)$, which implies g and h are generically conjugacy separated, and follows from [5, Theorem 1]. More specifically, [5, Theorem 1] proves that for any tuple (a_1, \dots, a_m) , ‘almost all’ (precisely computed) tuples (b_1, \dots, b_m) , with $|b_i| \leq n$, give subgroups $L = \langle a_1, \dots, a_m \rangle$ and $K = \langle b_1, \dots, b_m \rangle$ with trivial pullback, that is, for every $u \in F(\Delta)$, $K^u \cap L = \{1\}$. Going over all (a_1, \dots, a_m) with $|a_i| \leq n$ and counting the tuples (b_1, \dots, b_m) (as in [5]) satisfying property (2) gives the genericity result for all $2m$ -tuples. \square

6.2 The GPCP for conjugacy inequivalent maps

We now prove that the PCP implies the $\text{GPCP}^{\text{inj} + \text{CI}}$ and hence that, generically, the PCP implies the GPCP. Recall that if I_{GPCP} is a specific instance of

the GPCP we can associate to it a specific instance $I_{\text{PCP}} = (\Sigma \sqcup \{B, E\}, \Delta \sqcup \{B, E, \#\}, g', h')$, as in Section 3. We start by classifying the solutions to I_{PCP} .

Lemma 5. *Let I_{GPCP} be an instance of the GPCP^{inj}, with associated instance I_{PCP} of the PCP^{inj}. Every solution to I_{PCP} is a product of solutions of the form $(BxE)^{\pm 1}$, $E^{-1}xE$ and BxB^{-1} , for $x \in F(\Sigma)$.*

Lemma 5 is proven in the appendix. We now have:

Theorem 6. *Let $I_{\text{GPCP}} = (\Sigma, \Delta, g, h, u_1, u_2, v_1, v_2)$ be an instance of the GPCP^{inj}, such that there is no non-trivial $x \in F(\Sigma)$ with $u_1g(x)u_1^{-1} = v_1h(x)v_1^{-1}$ or $u_2^{-1}g(x)u_2 = v_2^{-1}h(x)v_2$. Then I_{GPCP} has a solution (possibly trivial) if and only if the associated instance I_{PCP} of the PCP^{inj} has a non-trivial solution.*

Proof. By Lemma 4, if I_{GPCP} has a solution then I_{PCP} has a non-trivial solution. For the other direction, note that the assumptions in the theorem are equivalent to I_{GPCP} having no solutions of the form BxB^{-1} or $E^{-1}xE$, and so by Lemma 5, every non-trivial solution to I_{GPCP} has the form $Bx_1E \cdots Bx_nE$ for some $x_i \in F(\Sigma)$. The Bx_iE subwords block this word off into chunks, and we see that each such word is a solution to I_{PCP} . By Lemma 4, each x_i is a solution to I_{GPCP} . Hence, if I_{PCP} has a non-trivial solution then I_{GPCP} has a solution. \square

Theorem 6 depends both on the maps g and h and on the constants u_i, v_i . The definition of conjugacy inequivalent maps implies that the conditions of Theorem 6 hold always, independent of the u_i, v_i . We therefore have:

Theorem 7. *The following hold in finitely generated free groups.*

1. $\text{PCP}^{(-\text{inj}, \text{inj})}(n+2) \Rightarrow \text{GPCP}^{(-\text{inj}, \text{inj})+\text{CI}}(n)$
2. $\text{PCP}^{(\text{inj}, \text{inj})}(n+2) \Rightarrow \text{GPCP}^{(\text{inj}, \text{inj})+\text{CI}}(n)$

Removing the injectivity assumptions gives the following corollary; the implication $\text{PCP} \Rightarrow \text{GPCP}^{\text{inj}+\text{CI}}$ of Theorem A follow immediately.

Corollary 2. $\text{PCP}(n+2) \Rightarrow \text{GPCP}^{\text{inj}+\text{CI}}(n)$

Proof. Theorem 7 gives us that $\text{PCP}^{\text{inj}}(n+2) \Rightarrow \text{GPCP}^{\text{inj}+\text{CI}}(n)$, while the $\text{PCP}(n)$ and $\text{PCP}^{\text{inj}}(n)$ are equivalent by Proposition 1. \square

7 The Basis Problem and Stallings' Rank Problem

In this section we link the Basis Problem to Stallings' Rank Problem. Clearly the Basis Problem solves the Rank Problem, as the rank is simply the size of the basis. We prove that these problems are equivalent, with Lemma 6 providing the non-obvious direction of the equivalence. Combining this equivalence with Corollary 1 gives: $\text{RP} \Rightarrow \text{GPCP}^{\text{inj}} \Rightarrow \text{PCP}$.

The proof of Lemma 6 is based on the following construction of Goldstein–Turner, which they used to prove that $\text{Eq}(g, h)$ is finitely generated [10]. Let

$F = F(\Delta)$ be a finitely generated free group, let H be a subgroup with basis $\mathcal{N} = \{\alpha_1, \dots, \alpha_k\}$ of F , where $\alpha_i \in F$, and let $\phi : H \rightarrow F$ be a homomorphism. The *derived graph* D_ϕ of ϕ relative to \mathcal{N} is the graph with vertex set F , and directed edges labeled by elements of \mathcal{N} according to the rule: if $u = \phi(\alpha_i)v\alpha_i^{-1}$ then connect u to v by an edge with initial vertex u , end vertex v , and label α_i . We shall use the fact that a word $w(\alpha_1, \dots, \alpha_k)$ is fixed by ϕ if and only if the path which starts at the vertex ϵ (the identity of F) and is labeled by $w(\alpha_1, \dots, \alpha_k)$ is a loop, so also ends at ϵ .

Lemma 6. *There exists an algorithm with input an instance $I = (\Sigma, \Delta, g, h)$ of the PCP^{inj} and the rank $\text{rk}(\text{Eq}(g, h))$ of the equaliser of g and h , and output a basis for $\text{Eq}(g, h)$.*

The following shows that Stallings' Rank Problem is equivalent to the BP.

Theorem 8. *The following hold in finitely generated free groups.*

1. $\text{BP}^{(\neg \text{inj}, \text{inj})}(n+2) \Leftrightarrow \text{RP}^{(\neg \text{inj}, \text{inj})}(n)$
2. $\text{BP}^{(\text{inj}, \text{inj})}(n+2) \Leftrightarrow \text{RP}^{(\text{inj}, \text{inj})}(n)$

Proof. Let I_{PCP} be an instance of the PCP^{inj} . As the rank of a free group is precisely the size of some (hence any) basis for it, if we can compute a basis for $\text{Eq}(g, h)$ then we can compute the rank of $\text{Eq}(g, h)$. On the other hand, by Lemma 6 if we can compute the rank of $\text{Eq}(g, h)$ then we can compute a basis of $\text{Eq}(g, h)$. \square

We therefore have:

Corollary 3. $\text{BP}(n) \Leftrightarrow \text{RP}(n)$

8 Main results, part 2

We now combine results from the previous sections to the following result, from which Theorem A follows immediately.

Theorem 9. *In finitely generated free groups the following implications hold.*

$$\begin{array}{ccccccc}
 & & \text{RP}(n) & & & & \\
 & & \Updownarrow & & & & \\
 & & \text{BP}(n) & \Longrightarrow & \text{GPCP}^{\text{inj}}(n) & \Longrightarrow & \text{PCP}(n-2) & \Longrightarrow & \text{GPCP}^{\text{inj}+\text{CI}}(n) \\
 & & \Downarrow & & & & \\
 & & \text{PCP}_{\mathcal{R}}^{\text{inj}}(n) & & & &
 \end{array}$$

Proof. The proof is a summary of the results already established in the rest of the paper, and we give a schematic version of it here.

$\text{RP}(n) \Leftrightarrow \text{BP}(n)$ holds by Corollary 3.

$\text{BP}(n) \Rightarrow \text{PCP}_{\mathcal{R}}^{\text{inj}}(n)$ holds by Theorem 2.

$\text{BP}(n) \Rightarrow \text{GPCP}^{\text{inj}}(n) \Rightarrow \text{PCP}(n)$ holds by Corollary 1.

$\text{PCP}(n-2) \Rightarrow \text{GPCP}^{\text{inj}+\text{CI}}(n)$ holds by Corollary 2. \square

References

1. Bartholdi, L., Silva, P.V.: Rational subsets of groups (2010), <https://arxiv.org/pdf/1012.1532.pdf>
2. Bogopolski, O., Maslakova, O.: An algorithm for finding a basis of the fixed point subgroup of an automorphism of a free group. *Internat. J. Algebra Comput.* **26**(1), 29–67 (2016)
3. Ciobanu, L., Logan, A.: Fixed points and stable images of endomorphisms for the free group of rank two. *arXiv:2009.04937* (2020)
4. Ciobanu, L., Logan, A.: The Post correspondence problem and equalisers for certain free group and monoid morphisms. In: 47th International Colloquium on Automata, Languages, and Programming (ICALP 2020). pp. 120:1–120:16 (2020)
5. Ciobanu, L., Martino, A., Ventura, E.: The generic Hanna Neumann Conjecture and Post Correspondence Problem (2008), <http://www-eupm.upc.es/~ventura/ventura/files/31t.pdf>
6. Diekert, V., Kharlampovich, O., Lohrey, M., Myasnikov, A.: Algorithmic problems in group theory. Dagstuhl seminar report 19131 (2019)
7. Ehrenfeucht, A., Karhumäki, J., Rozenberg, G.: The (generalized) Post correspondence problem with lists consisting of two words is decidable. *Theoret. Comput. Sci.* **21**(2), 119–144 (1982)
8. Feighn, M., Handel, M.: Algorithmic constructions of relative train track maps and CTs. *Groups Geom. Dyn.* **12**(3), 1159–1238 (2018)
9. Gilman, R., Miasnikov, A.G., Myasnikov, A.D., Ushakov, A.: Report on generic case complexity (2007), <https://arxiv.org/pdf/0707.1364v1.pdf>
10. Goldstein, R.Z., Turner, E.C.: Fixed subgroups of homomorphisms of free groups. *Bull. London Math. Soc.* **18**(5), 468–470 (1986)
11. Harju, T., Karhumäki, J.: Morphisms. In: Handbook of formal languages, Vol. 1, pp. 439–510. Springer, Berlin (1997)
12. Kapovich, I., Myasnikov, A.: Stallings foldings and subgroups of free groups. *J. Algebra* **248**(2), 608–668 (2002)
13. Kapovich, I., Myasnikov, A., Schupp, P., Shpilrain, V.: Generic-case complexity, decision problems in group theory, and random walks. *J. Algebra* **264**(2), 665–694 (2003)
14. Karhumäki, J., Saarela, A.: Noneffective regularity of equality languages and bounded delay morphisms. *Discrete Math. Theor. Comput. Sci.* **12**(4), 9–17 (2010)
15. Lecerf, Y.: Réursive insolubilité de l'équation générale de diagonalisation de deux monomorphismes de monoïdes libres $\varphi x = \psi x$. *C. R. Acad. Sci. Paris* **257**, 2940–2943 (1963)
16. Martino, A., Turner, E., Ventura, E.: The density of injective endomorphisms of a free group (2006), <http://www-eupm.upc.es/~ventura/ventura/files/23t.pdf>
17. Myasnikov, A., Nikolaev, A., Ushakov, A.: The Post correspondence problem in groups. *J. Group Theory* **17**(6), 991–1008 (2014)
18. Neary, T.: Undecidability in binary tag systems and the post correspondence problem for five pairs of words. In: 32nd International Symposium on Theoretical Aspects of Computer Science, LIPIcs. Leibniz Int. Proc. Inform., vol. 30, pp. 649–661. Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern (2015)
19. Post, E.L.: A variant of a recursively unsolvable problem. *Bull. Amer. Math. Soc.* **52**, 264–268 (1946)

Appendix: Additional material to Sections 6 & 7

Proof (Lemma 5). Let x' be a solution to I_{PCP} , and decompose it as a freely reduced word

$$x_0\alpha_1x_1\alpha_2\cdots x_{n-1}\alpha_nx_n$$

for $x_i \in F(\Sigma)$ and $\alpha_i \in \{B, E\}^{\pm 1}$.

We shall refer to the letters $B^{\pm 1}, E^{\pm 1}$ as *markers*. Indeed, they act as “separators” in the word x' , because in the definitions of g' and h' the letter B only occurs in $g'(B)$ and $h'(B)$, and the letter E only occurs in $g'(E)$ and $h'(E)$; thus either the $B^{\pm 1}$ and $E^{\pm 1}$ terms in x' are preserved under application of g' and h' , or two of these letters cancel in the image. Therefore, the parts of each of $g'(\alpha_i x_i \alpha_{i+1})$ and $h'(\alpha_i x_i \alpha_{i+1})$ lying between the markers must be equal, or otherwise there exist some indices i, j such that both $g'(\alpha_i x_i \alpha_{i+1})$ and $h'(\alpha_j x_j \alpha_{j+1})$ are contained in $F(\Delta)$: this collapsing must happen in each image as the numbers of $B^{\pm 1}$ and $E^{\pm 1}$ terms must be equal in both images. By inspecting the maps, we see that this collapsing requires $x_i \in \ker(g')$ and $\alpha_i = \alpha_{i+1}^{-1}$, and $x_j \in \ker(h')$ and $\alpha_j = \alpha_{j+1}^{-1}$. However, by Lemma 3 only one of g' or h' is injective, g' say, so then x_i is trivial, and so $\alpha_i x_i \alpha_{i+1}$ is empty, contradicting the decomposition being freely reduced. Therefore, we have that for all $1 \leq i < n$ the parts of each of $g'(\alpha_i x_i \alpha_{i+1})$ and $h'(\alpha_i x_i \alpha_{i+1})$ lying between the markers $B^{\pm 1}, E^{\pm 1}$ are equal. This also implies that the parts of $g'(x_0 \alpha_1)$ and $h'(x_0 \alpha_1)$ lying before the markers are equal.

We next prove that x_0 is empty, $\alpha_1 \in \{B, E^{-1}\}$, $\alpha_2 \in \{B^{-1}, E\}$ (so in particular, non-empty), and that $\alpha_1 x_1 \alpha_2$ is a solution to I_{PCP} .

Suppose x_0 is non-empty. Then either $g'(x_0 \alpha_1)$ starts with $\#^{-1}$ or E (when u_2 is empty), or $x_0 \in \ker g'$ and the image starts with B or E^{-1} , while either $h'(x_0 \alpha_1)$ starts with $\#$ or B^{-1} (when v_1 is empty), or $x_0 \in \ker h'$ and the image starts with B or E^{-1} . As one map is injective, by Lemma 3, we have that $g'(x_0 \alpha_1)$ and $h'(x_0 \alpha_1)$ start with different letters, contradicting the above paragraph. Hence, x_0 is empty. We then have that $\alpha_1 \in \{B, E^{-1}\}$, as this is the only way that the parts of $g'(\alpha_1)$ and $h'(\alpha_1)$ lying before the marker can be equal.

Suppose α_2 is empty, so $x' = \alpha_1 x_1$. Then as x'^{-1} is also a solution to I_{PCP} , we also have that x_1 is empty, and so $x' = \alpha_1 \in \{B, E^{-1}\}$. However, neither B nor E^{-1} is a solution to I_{PCP} , a contradiction. Hence, α_2 is non-empty. Moreover, $\alpha_2 \in \{B^{-1}, E\}$, as otherwise the parts of each of $g'(\alpha_1 x_1 \alpha_2)$ and $h'(\alpha_1 x_1 \alpha_2)$ lying between the markers are non-equal (for example, $g'(Bx_1B) = B\#u_1g(x_1)B\cdots$ while $h'(Bx_1B) = B\#v_1h(x_1)\#^{-1}B\cdots$). Finally, under these restrictions on α_1 and α_2 , and because the parts of $g'(\alpha_1 x_1 \alpha_2)$ and $h'(\alpha_1 x_1 \alpha_2)$ lying between the markers are equal, we get that $\alpha_1 x_1 \alpha_2$ is a solution to I_{PCP} of the form $(Bx_1E)^{\pm 1}, E^{-1}x_1E$, or Bx_1B^{-1} for $x_1 \in F(\Sigma)$.

We can now prove the result: the product $(\alpha_1 x_1 \alpha_2)^{-1} x'$ is also a solution to I_{PCP} , and it decomposes as $x_2 \alpha_3 \cdots x_{n-1} \alpha_n x_n$. Applying the above argument, we get that x_2 is empty and that $\alpha_3 x_3 \alpha_4$ is a solution to I_{PCP} of the form $(Bx_3E)^{\pm 1}, E^{-1}x_3E$, or Bx_3B^{-1} for $x_3 \in F(\Sigma)$. Repeatedly reducing the solution like this,

we see that x' decomposes as $x''x_n$ where x'' is a product of solutions of the form $(BxE)^{\pm 1}$, $E^{-1}xE$, and BxB^{-1} for $x \in F(\Sigma)$. The result then follows as x_n is empty, which can be seen by applying the above argument to the solution x'^{-1} , which decomposes as $x_n\alpha_{n-1}\cdots$. \square

Proof (Lemma 6). Suppose without loss of generality that g is injective, and consider the homomorphism $\phi = h \circ g^{-1} : \text{Im}(g) \rightarrow F(\Delta)$. Then $g(\text{Eq}(g, h))$ is precisely the fixed subgroup $\text{Fix}(\phi) = \{x \in \text{Im}(g) \mid \phi(x) = x\}$, and as g is injective it restricts to an isomorphism between these two subgroups. We therefore give an algorithm which takes as input $I = (\Sigma, \Delta, g, h)$ and the rank of $\text{Eq}(g, h)$ and outputs a basis \mathcal{B} for $\text{Fix}(\phi)$; this is sufficient as $g^{-1}(\mathcal{B})$ is then an algorithmically computable basis for $\text{Eq}(g, h)$. If $\text{rk}(\text{Eq}(g, h)) = 0$ then the basis is the empty set, so we may assume $\text{rk}(\text{Eq}(g, h)) \geq 1$ (this reduction is not necessary, but not doing so introduces certain subtleties).

Let Γ_ϕ be the union of those loops in D_ϕ which contain no degree-1 vertices, but contain the vertex ϵ ; this is simply the component of the core graph of D_ϕ which contains the specified vertex. As labels of loops in D_ϕ correspond to elements of $\text{Fix}(\phi)$, any basis of the fundamental group $\pi_1(\Gamma_\phi)$ corresponds to a basis of $\text{Fix}(\phi)$. Now, a basis for $\pi_1(\Gamma_\phi)$ can be computed via standard algorithms (see for example [12, Propositions 6.7]), and therefore to prove the result we only need to construct the graph Γ_ϕ .

Start with vertex set V consisting of a single vertex ϵ corresponding to the empty word. Now enter a loop, terminating when $|E| - |V| = \text{rk}(\text{Eq}(g, h)) - 1$, as follows: For each $v \in V$ and each $i \in \{1, \dots, k\}$, add an edge to E starting at v and ending at the vertex corresponding to the element $\phi(\alpha_i)g\alpha_i^{-1}$ of $F(\Sigma)$; if there is no such vertex in V then first add one to V . When the loop terminates, prune the resulting graph to obtain a graph Γ'_ϕ by iteratively removing all degree-1 vertices.

Note that $\text{rk}(\text{Eq}(g, h))$ is known, so we can determine if a graph satisfies $|E| - |V| = \text{rk}(\text{Eq}(g, h)) - 1$; in particular, at each iteration in the looping procedure we know whether to continue or to terminate the loop.

There are two things to prove. Firstly, that the looping procedure terminates (and so the above is actually an algorithm), and secondly that the terminating graph Γ'_ϕ is in fact the graph Γ_ϕ . So, note that the procedure constructs *some* subgraph of D_ϕ which contains the empty word as a vertex, which has no degree-1 vertices, and which satisfies $\#\text{edges} - \#\text{vertices} = \text{rk}(\text{Eq}(g, h)) - 1$, or no such subgraph exists. Now, the graph Γ_ϕ satisfies these conditions (because its fundamental group has rank $\text{rk}(\text{Eq}(g, h))$), so the procedure does terminate. Moreover, every loop in D_ϕ which contains the empty word is a loop in Γ_ϕ , and so the subgraph constructed is in fact a subgraph of Γ_ϕ . The result then follows because no proper subgraph of Γ_ϕ satisfies $\#\text{edges} - \#\text{vertices} = \text{rk}(\text{Eq}(g, h)) - 1$ (subgraphs in fact satisfy $\#\text{edges} - \#\text{vertices} \leq \text{rk}(\text{Eq}(g, h)) - 1$). \square