

# Thermal fluctuations of a stellar system

Jun Yan Lau<sup>1,2\*</sup> & James Binney<sup>2†</sup>

<sup>1</sup>*UCL Mullard Space Sciences Laboratory, Holmbury St Mary, Surrey RH5 6NT*

<sup>2</sup>*Rudolf Peierls Centre for Theoretical Physics, Clarendon Laboratory, Parks Road, Oxford, OX1 3PU, UK*

16 April 2021

## ABSTRACT

The excursions of star clusters and galaxies around statistical equilibria are studied. For an ergodic model with monotone decreasing DF Antonov’s Hermitian operator on six-dimensional phase space has the normal modes as its eigenfunctions. The excitation energy of the system is just the sum of the (positive) energies associated with each normal mode. The positivity of modal energies opens the way to modelling the thermal properties of clusters in close analogy with those of crystals. Formulae are given for the DFs of modes, which are of the type first described by van Kampen rather than Landau. Each mode comprises the response of non-resonant stars to driving by the gravitational field of stars on a group of resonant tori. The structure of each mode is sensitive to the degree of self gravity. The emergence of global distortions in N-body models when particles are started from an analytical equilibrium is explained in terms of the interplay of normal modes.

**Key words:** Galaxy: kinematics and dynamics – galaxies: kinematics and dynamics – methods: analytical

## 1 INTRODUCTION

Galaxies and star clusters are in approximate states of equilibrium and have for decades been fitted to models in which the distribution function  $f(\mathbf{x}, \mathbf{v})$  of their constituent particles (stars, dark-matter particles) are steady-state solutions of the collisionless Boltzmann equation (CBE). The advent of massive simulations of galaxy formation (Laporte et al. 2019) and detailed data from the Gaia mission (Gaia Collaboration & Brown 2018) and large integral field units such as MUSE (e.g. Vitral & Mamon 2021) have stimulated interest in non-equilibrium features of galaxies, especially the Milky Way (Antoja et al. 2018).

An obvious source of disequilibrium is Poisson noise: stellar systems contain only a finite number of stars and in principle their observables depend on the precise locations of stars, and thus fluctuate as stars move hither and thither. Since a significant fraction of the light from a stellar population is contributed by the small fraction of stars that are nearing the ends of their lives, Poisson noise is particularly pronounced in surface brightness, and a widely used technique for estimating the distances to early-type galaxies revolves around quantifying fluctuations in surface brightness (Tonry & Schneider 1988). The long-term evolution of star clusters has long been attributed to the fluctuations in their gravitational fields that accompany Poisson fluctua-

tions in stellar density. In the classical theory of such evolution, the field fluctuations are computed under neglect of the system’s self-gravity. Recently it has become apparent that this assumption can be profoundly misleading. Heyvaerts (2010) and Chavanis (2012) developed a formalism, the Balescu-Lenard (BL) equation, for computing how stars diffuse through phase space, in which it is possible to take self-gravity into effect. Fouvy et al. (2015) applied this theory to razor-thin discs and showed that in these systems self-gravity accelerates the rate of diffusion by more than two orders of magnitude. In addition to accelerating the diffusion, self-gravity radically changes the way diffusion occurs by enhancing large-scale fluctuations much more than small-scale ones, so diffusion is dominated by a small number of global modes of oscillation, which have quite well defined frequencies. Stars which resonate with these global oscillations dominate the diffusion.

Hamilton et al. (2018) applied the BL equation to spherical star clusters and found that even in these systems global modes excited by Poisson noise generate a significant diffusive flux through phase space. Lau & Binney (2019) explored stellar diffusion in N-body models of clusters in which forces were computed by direct summation. The initial conditions were obtained by sampling the analytic DF of a steady-state model cluster. Hence only Poisson noise separated their initial conditions from perfect equilibrium. They found that the amplitudes of dipole and quadrupole distortions of the cluster grew steadily for several crossing times. Hence their initial conditions were symmetric and smooth to an extent

\* E-mail: jun.lau.20@ucl.ac.uk

† E-mail: binney@physics.ox.ac.uk

that no real cluster will ever be. Work is currently underway to check that clusters observed by Gaia show similar dipole and quadrupole distortions. Non-equilibrium structures were first detected more than 40 years ago (Malin & Carter 1980) but have invariably been modelled in terms of test particles rather than collective oscillations of the host system.

Perhaps the most exciting single discovery made in the Gaia DR2 data is the phase spiral that Antoja et al. (2018) uncovered in the distribution of stars in the  $(z, v_z)$  plane. The spiral is surely a symptom of a macroscopic oscillation of the disc that has a significant component in the  $z$  direction. If we had a credible dynamical model of this oscillation, we would be able to extract from the Gaia data information about the structure of the disc and the agent [likely the Sagittarius dwarf galaxy (Binney & Schönrich 2018; Laporte et al. 2019; Bland-Hawthorn & Tepper-Garcia 2020)] that excited it. Unfortunately, the disc’s self-gravity certainly plays an important role in the oscillation, and there’s little prospect of adequately modelling the disc’s oscillation until we have a better understanding of the global oscillations of stellar systems. This paper aims to take a small step towards that understanding.

Poisson noise is for us the natural starting point because it sets a lower limit on fluctuations. Such noise can also be called thermal noise because fluctuations in the pressure on a small disc or the current in a small circuit also arise because gases comprise a finite number of molecules and circuits contain a finite number of electrons. The theory of thermal fluctuations in gases and condensed matter is well understood, so it makes sense to consider thermal noise in stellar systems in this context.

The thermal properties of solids require a completely different approach to that used to understand the thermal properties of dilute gases: whereas in a dilute gas we may treat each molecule as an independent system, in a solid molecules lack independence because they are tightly coupled to one another. Following Debye the independent elements in thermal theory become collective oscillations of the entire system: phonons. At sufficiently high temperatures, each phonon carries  $k_B T$  of thermal energy on average. Since the density of phonon states  $g(E)$  rises rapidly with increasing wavenumber, most thermal energy is invested in phonons with wavelengths comparable to the inter-molecular spacing and we arrive at results that are similar to those for an ideal gas except at low temperatures, when high-frequency modes are exponentially suppressed and mesoscopic modes dominate.

The stars of a cluster are strongly coupled by gravity, so they cannot be considered independent. Moreover, the work of Hamilton et al. (2018) and Lau & Binney (2019) suggests that a cluster’s system-scale collective oscillations are thermally excited to significant amplitudes, so we should approach the thermal properties of a cluster in terms of normal modes regardless of temperature. Specifically, we should express the system’s thermal energy as a sum of the energies of its normal modes,

$$E = \sum_{\beta} |c_{\beta}|^2 E^{(\beta)}, \quad (1)$$

with each amplitude  $c_{\beta}$  being a random variable with variance  $k_B T / E^{(\beta)}$ .

Section 2 introduces basic concepts and establishes notation. Section 3 introduces the Hermitian operator on phase space whose eigenfunctions are the required normal modes of the cluster. We show that the energies of normal modes are additive as equation (1) requires, and give a very simple expression for a mode’s energy in terms of its DF. This expression implies that the energy of oscillatory modes is positive. We show also that the operator gives rise to a slightly different conserved quantity that provides a means to establish stability. In Section 4 we obtain an expression for the DFs of the normal modes. This contains a free function and parameters that can be computed from the free function by matrix algebra. We investigate the way in which the structures of normal modes depends on the extent to which a system is self-gravitating and use the understanding thus gained to explain the emergence of system-scale distortions in N-body realisations of clusters. We re-express a mode’s energy in terms of the free function and the potential that the mode generates. In Section 5 we argue that van Kampen modes rest on conceptual foundations which are as solid as those that are generally accepted as secure in other branches of physics.

In Section 6 we stress the importance of the concept of particle dressing in stellar dynamics as in other branches of physics, and discuss the role that van Kampen modes play in dressing. We also consider the prospect for using van Kampen modes to extend conventional statistical mechanics to stellar systems. Section 7 sums up and indicates two directions for future work.

## 2 MATHEMATICAL BACKGROUND

Here we introduce essential mathematical tools and establish our notation. We focus on stable ergodic clusters, that is systems with unperturbed DFs  $f_0(H)$  where

$$H(\mathbf{w}'), \equiv \frac{1}{2}v'^2 + \Phi(\mathbf{x}') \quad (2)$$

is the Hamiltonian of a single particle in the gravitational potential  $\Phi(\mathbf{x})$ . A necessary and sufficient condition for such a system to be stable is that the derivative  $f'_0 < 0$  at all energies (Antonov 1961).

### 2.1 Variable degree of self gravity

In the following it proves helpful to be able to consider self-gravity to be a variable  $\xi$  that runs from zero (stars move in the fixed potential of a specified density distribution) to unity (stars experience only their gravitational attraction to the other stars). It is straightforward to set up a numerical experiment for any given value of  $\xi$  by sampling an analytic density distribution in the usual way and taking the force on each star to be  $\xi$  times the force returned by an N-body solver plus  $(1 - \xi)$  times the force provided by the analytic density.

### 2.2 Angle-action variables

The role that Cartesian variables play for homogeneous systems is played for spheroidal systems by angle-action variables  $(\boldsymbol{\theta}, \mathbf{J})$ . The actions  $J_i$  are constants of motion while their conjugate variables, the angles  $\theta_i$ , increase linearly

in time, so  $\boldsymbol{\theta}(t) = \boldsymbol{\theta}(0) + \boldsymbol{\Omega}t$ . The particles' Hamiltonian  $H(\mathbf{x}, \mathbf{v})$  is a function  $H(\mathbf{J})$  of the actions only and the frequencies  $\Omega_i$  that control the rates of increase of the angles are given by  $\boldsymbol{\Omega} = \partial H / \partial \mathbf{J}$ . Angle-action variables are canonical, so the volume element of phase space  $d^6\mathbf{w} = d^3\mathbf{x}d^3\mathbf{v} = d^3\boldsymbol{\theta}d^3\mathbf{J}$  and Poisson brackets can be computed as

$$[f, g] = \sum_i \left( \frac{\partial f}{\partial \theta_i} \frac{\partial g}{\partial J_i} - \frac{\partial f}{\partial J_i} \frac{\partial g}{\partial \theta_i} \right). \quad (3)$$

Functions on phase space can be expressed as Fourier series:

$$h(\mathbf{w}) = \sum_{\mathbf{n}} h_{\mathbf{n}}(\mathbf{J}) e^{i\mathbf{n}\cdot\boldsymbol{\theta}}; \quad h_{\mathbf{n}}(\mathbf{J}) = \int \frac{d^3\boldsymbol{\theta}}{(2\pi)^3} e^{-i\mathbf{n}\cdot\boldsymbol{\theta}} h(\mathbf{w}). \quad (4)$$

Note that for real  $h$ ,  $h_{-\mathbf{n}} = h_{\mathbf{n}}^*$ .

### 2.3 Potential-density pairs

Unfortunately, while the potential  $\Phi(\mathbf{x})$  is a function of only  $\mathbf{x}$ , it becomes a function of both  $\boldsymbol{\theta}$  and  $\mathbf{J}$ . So while angle-action variables make dynamics trivial, they seriously complicate the solution of Poisson's equation. Following Kalnajs (1976) this difficulty is finessed by introducing a basis of biorthogonal potential-density pairs. That is, a set of pairs  $(\rho^{(\alpha)}, \Phi^{(\alpha)})$  such that

$$4\pi G\rho^{(\alpha)} = \nabla^2\Phi^{(\alpha)} \quad \text{and} \quad \int d^3\mathbf{x} \Phi^{(\alpha)*} \rho^{(\alpha')} = -\mathcal{E}\delta_{\alpha\alpha'}, \quad (5)$$

where  $\mathcal{E}$  is an arbitrary constant with the dimensions of energy. Given a density distribution  $\rho(\mathbf{x})$ , we expand it in the basis

$$\rho(\mathbf{x}) = \sum_{\alpha} A_{\alpha} \rho^{(\alpha)}(\mathbf{x}) \quad \Leftrightarrow \quad \Phi(\mathbf{x}) = \sum_{\alpha} A_{\alpha} \Phi^{(\alpha)}(\mathbf{x}), \quad (6)$$

where

$$A_{\alpha} = -\frac{\xi}{\mathcal{E}} \int d^6\mathbf{w} \Phi^{(\alpha)*}(\mathbf{x}) f(\mathbf{w}). \quad (7)$$

If  $\rho$  and  $\Phi$  are time-dependent, the  $A_{\alpha}$  become time-dependent. From equations (5) and (6) one can obtain an expression for  $\Phi$  in terms of  $\rho$ . Comparison of this relation with Poisson's integral, yields

$$\frac{G}{|\mathbf{x}' - \mathbf{x}|} = \frac{1}{\mathcal{E}} \sum_{\alpha} \Phi^{(\alpha)}(\mathbf{x}) \Phi^{(\alpha)*}(\mathbf{x}'). \quad (8)$$

## 3 ANTONOV'S OPERATOR K

We now derive for stable ergodic clusters the Hermitian operator  $K$  introduced by Antonov (1961). The true normal modes of the system are eigenfunctions of  $K$  with non-negative eigenvalues that prove to be the squares of the modes' (real) frequencies. Following Antonov (1961) we split the perturbed DF  $f_1$  into parts that are even and odd in  $\mathbf{v}$ :

$$f_1(\mathbf{x}, \mathbf{v}) = f_+(\mathbf{x}, \mathbf{v}) + f_-(\mathbf{x}, \mathbf{v}) \quad (9)$$

where

$$f_{\pm}(\mathbf{x}, \mathbf{v}) \equiv \frac{1}{2}[f_1(\mathbf{x}, \mathbf{v}) \pm f_1(\mathbf{x}, -\mathbf{v})]. \quad (10)$$

In the absence of a perturbation,  $f_-$  vanishes, so this part of the DF isolates the effect of the perturbation. On the other

hand the perturbation changes the potential only through  $f_+$ .<sup>1</sup>

We define an inner product on the space of DFs by

$$\langle g|f \rangle \equiv - \int d^6\mathbf{w} \frac{g^* f}{f'_0(H)}, \quad (11)$$

where  $H(\mathbf{J})$  is the unperturbed Hamiltonian and the leading minus reflects the fact that  $f'_0 < 0$ . Since  $d^6\mathbf{w} f$  has dimensions of mass and  $f/f'_0$  has dimensions of  $v^2$ ,  $\langle g|f \rangle$  has the dimensions of  $Mv^2$ , i.e., energy. When we Fourier expand the DFs we find

$$\langle g|f \rangle = -(2\pi)^3 \int \frac{d^3\mathbf{J}}{f'_0} \sum_{\mathbf{k}} g_{\mathbf{k}}^* f_{\mathbf{k}} \quad (12)$$

Notice that

$$\langle f_1|f_1 \rangle = \langle f_-|f_- \rangle + \langle f_+|f_+ \rangle. \quad (13)$$

$H(\mathbf{J})$  is even in  $\mathbf{v}$  and the Poisson bracket operator is odd in  $\mathbf{v}$ , so our division (9) of  $f$  splits the linearised CBE,  $\partial_t f_1 + [f_1, H] + [f_0, \Phi_1] = 0$ , into two equations

$$\frac{\partial f_+}{\partial t} = -[f_-, H]; \quad \frac{\partial f_-}{\partial t} = -[f_+, H] + [\Phi_1, f_0]. \quad (14)$$

Now  $[\Phi_1, f_0] = f'_0(H)[\Phi_1, H]$ , so

$$\begin{aligned} \frac{\partial}{\partial t} [\Phi_1, f_0](\mathbf{w}) &= f'_0(H) \left[ \frac{\partial \Phi_1}{\partial t}, H \right] \\ &= -f'_0(H) \left[ \int d^6\mathbf{w}' \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|} \frac{\partial f_+(\mathbf{w}')}{\partial t}, H(\mathbf{w}) \right]. \end{aligned}$$

We differentiate the second of equations (14) wrt  $t$  and use the first equation to eliminate  $\partial_t f_+$  from the rhs to obtain

$$\begin{aligned} \frac{\partial^2 f_-}{\partial t^2} &= [[f_-, H], H] \\ &+ f'_0(H) \left[ \int d^6\mathbf{w}' \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|} [f_-(\mathbf{w}'), H(\mathbf{w}')], H(\mathbf{w}) \right] \end{aligned} \quad (16)$$

This equation is of the form

$$\frac{\partial^2 f_-}{\partial t^2} = -K f_-, \quad (17)$$

where the operator

$$\begin{aligned} K &\equiv -[[f_-, H], H] \\ &- f'_0(H) \left[ \int d^6\mathbf{w}' \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|} [f_-(\mathbf{w}'), H(\mathbf{w}')], H(\mathbf{w}) \right] \end{aligned} \quad (18)$$

In terms of angle-action coordinates  $(\boldsymbol{\theta}, \mathbf{J})$ ,

$$[f(\boldsymbol{\theta}, \mathbf{J}), H(\mathbf{J})] = \boldsymbol{\Omega} \cdot \frac{\partial f}{\partial \boldsymbol{\theta}}, \quad (19)$$

so  $K$  can be written

$$\begin{aligned} K &= - \left( \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \right)^2 f_- \\ &- f'_0(H) \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \int d^6\mathbf{w}' \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|} \boldsymbol{\Omega}' \cdot \frac{\partial f_-(\mathbf{w}')}{\partial \boldsymbol{\theta}'}. \end{aligned} \quad (20)$$

<sup>1</sup> In Dirac's seminal textbook, he argues that the second-order Klein-Gordon equation cannot stand in for the Schrödinger equation, which is first-order in time. So he factorises the Klein-Gordon operator into two first-order operators, by splitting the wavefunction into four parts. Antonov proceeded in the opposite direction: by splitting the DF he derived two first-order operators and then combined them to obtain a second-order equation.

Inserting equation (8) into (20) yields

$$K = - \left( \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \right)^2 f_- - f'_0(H) \frac{\xi}{\xi} \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \\ \times \int d^6 \mathbf{w}' \sum_{\alpha} \Phi^{(\alpha)}(\mathbf{x}) \Phi^{(\alpha)*}(\mathbf{x}') \boldsymbol{\Omega}' \cdot \frac{\partial f_-(\mathbf{w}')}{\partial \boldsymbol{\theta}'}. \quad (21)$$

At this point it's convenient to define

$$j_{\alpha}[f_1](t) \equiv -i \frac{\xi}{\xi} \int d^6 \mathbf{w} \Phi^{(\alpha)*}(\mathbf{x}) \boldsymbol{\Omega} \cdot \frac{\partial f_-(\mathbf{w})}{\partial \boldsymbol{\theta}}, \quad (22)$$

because it allows us to write equation (21) in the form

$$K = - \left( \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \right)^2 f_- - i f'_0(H) \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \sum_{\alpha} \Phi^{(\alpha)}(\mathbf{x}) j_{\alpha}[f_-]. \quad (23)$$

$j_{\alpha}$  is a functional of  $f_1$  rather than just  $f_-$  because it can also be computed from  $f_+$ : eliminating  $f_-$  using the first of equations (14) and equation (19) we obtain

$$j_{\alpha}[f_1](t) = i \frac{\xi}{\xi} \int d^6 \mathbf{w} \Phi^{(\alpha)*}(\mathbf{x}) \frac{\partial f_+}{\partial t}. \quad (24)$$

Since by equation (7) the coefficient  $A_{\alpha}$  of the potential/density expansion is a linear functional of  $f_1$ , and  $A[f_-] = 0$ , the derivative of  $f_+$  in equation (24) can be replaced by a derivative of  $A_{\alpha}$  to yield

$$j_{\alpha}[f_1](t) = -i \frac{\partial A_{\alpha}}{\partial t}. \quad (25)$$

### 3.1 Energy of a disturbance

When we use equation (8) to eliminate  $|\mathbf{x} - \mathbf{x}'|$  from equation (5.130) of Binney & Tremaine (2008), we find that the energy associated with a linearised disturbance is

$$E[f_1] = \frac{1}{2} \left\{ \int \frac{d^6 \mathbf{w}}{|f_0|} f_1^2 - \frac{\xi}{\xi} \sum_{\alpha} \left| \int d^6 \mathbf{w} \Phi^{(\alpha)}(\mathbf{x}) f_1(\mathbf{w}) \right|^2 \right\} \\ = \frac{1}{2} \left\{ \langle f_- | f_- \rangle + \langle f_+ | f_+ \rangle - \frac{\xi}{\xi} \sum_{\alpha} |A_{\alpha}[f_+]|^2 \right\}. \quad (26)$$

In the first line of this equation, the first and second terms on the right quantify the potential and kinetic energies of the perturbation, respectively. Hence the inner product  $\langle f_1 | f_1 \rangle$  gives twice a perturbation's kinetic energy. In the last term,  $A_{\alpha}$  is proportional to the degree of self-gravity  $\xi$ , so overall the final term is proportional to  $\xi$  as it should be.

In Appendix A we show that  $K$  is Hermitian and that

$$\langle f_- | K | f_- \rangle = |\omega^2| \left\{ \langle f_+ | f_+ \rangle - \xi \sum_{\alpha} |A_{\alpha}[f_+]|^2 \right\}. \quad (27)$$

When we use this equation to simplify equation (26), we discover the energy associated with an eigenfunction of  $K$  identically vanishes in the unstable case  $\omega^2 < 0$ , and in the stable case is

$$E[f_1] = \langle f_- | f_- \rangle = -(2\pi)^3 \int \frac{d^3 \mathbf{J}}{f'_0} \sum_{\mathbf{k}} |f_{\mathbf{k}-}|^2, \quad (28)$$

where  $f_{\mathbf{k}-}$  denotes the  $\mathbf{k}$  component of  $f_-$ . The right side of equation (28) is inherently positive and vanishes only if  $f_-$  vanishes. Moreover, by equation (14)  $\omega f_+ = \mathbf{k} \cdot \boldsymbol{\Omega} f_-$ , so  $f_+$  must also vanish if  $f_-$  does. Hence the energy of every mode of a stable system is positive.

Since  $K$  is Hermitian it has a complete set of orthogonal eigenfunctions. Expressing an arbitrary disturbance as a linear combination of eigenfunctions  $f_- = \sum_{\beta} c_{\beta} f_-^{(\beta)}$  and inserting this expansion into equation (28), we conclude that the disturbance's energy is the sum of the energies of its component eigenfunctions

$$E[f_1] = \sum_{\beta} |c_{\beta}|^2 \langle f_-^{(\beta)} | f_-^{(\beta)} \rangle \quad (29)$$

as equation (1) postulates.

These results establish that no mode of an ergodic system has negative energy; oscillatory modes have positive energy and growing/decaying modes (if any) have zero energy. Equation (26), from which we started, does not make evident the non-negativity of energies.

### 3.2 Antonov's conserved quantity

The rather involved argument just given starts from equation (26) for the energy of a disturbance, which Binney & Tremaine (2008) derive by considering the work that must be done to initiate the disturbance. A much simpler argument based on the Hermitian nature of  $K$  yields the closely related conserved quantity,

$$\tilde{E} = \frac{1}{2} \left( \langle \dot{f}_- | \dot{f}_- \rangle + \langle f_- | K | f_- \rangle \right). \quad (30)$$

Indeed,

$$2 \frac{d\tilde{E}}{dt} = \langle \ddot{f}_- | \dot{f}_- \rangle + \langle \dot{f}_- | \ddot{f}_- \rangle + \langle \dot{f}_- | K | f_- \rangle + \langle f_- | K | \dot{f}_- \rangle = 0. \quad (31)$$

An eigenfunction of  $K$  with eigenvalue  $\omega^2$  has  $\tilde{E} = \omega^2 \langle f_- | f_- \rangle$ , so in this case conservation of  $\tilde{E}$  implies conservation of  $E = \tilde{E}/\omega^2$ .  $\tilde{E}$  does not have the dimensions of energy, however.

Since  $\langle \dot{f}_- | \dot{f}_- \rangle > 0$ , instability, and thus systematic growth of  $f_-$ , is excluded by conservation of  $\tilde{E}$  unless  $\langle f_- | K | f_- \rangle < 0$  for some function  $f_-(\mathbf{x}, \mathbf{v})$ . In fact positivity of  $\langle f_- | K | f_- \rangle$  for all functions in the natural space is a necessary and sufficient condition for stability (Laval et al. 1965; Kulsrud & Mark 1970).

## 4 DFS OF VAN KAMPEN MODES

If the system is stable, the Hamiltonian's time-reversal invariance ensures that  $\omega$  is real because the existence of exponentially decaying solutions would imply the existence of growing solutions. van Kampen (1955) applied these arguments to an electrostatic plasma and deduced some properties of the normal modes of a plasma, which are known as van Kampen modes. The corresponding modes of a stellar system have received little attention, although Vandervoort (2003) derived some of their properties. We now examine the van Kampen modes of a stellar system in their role as eigenfunctions of the operator  $K$ .

### 4.1 Operators that commute with $K$

Finding the eigenfunctions of  $K$  is facilitated by identifying operators that commute with  $K$  and seeking eigenfunctions of  $K$  that are also eigenfunctions of these operators.  $K$  is

the operator associated with the time-translation invariance of the underlying equilibrium. The Hamiltonian  $H$  is invariant under increments in the angle variables  $\theta_i$  but in the presence of self gravity ( $\xi > 0$ ),  $K$  does not share the invariance with respect to increments in  $\theta_1$  and  $\theta_2$  conjugate to the radial action  $J_r$  and the modulus  $L \equiv |\mathbf{L}|$  of the angular momentum vector  $\mathbf{L}$  because  $\Phi_1(\mathbf{x})$  lacks this invariance. Fortunately,  $K$  is always invariant under increments of the angle variable  $\theta_3$  conjugate to  $L_z$ . To see this it is best to return to the definition of  $K$  in equation (18). Since  $[H, L_k] = 0$ , we have

$$[Kf_-, L_z] = -[[f_-, L_z], H], H] - f'_0(H) \times \left[ \int d^6 \mathbf{w}' \left[ \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|}, L_z \right] [f_-(\mathbf{w}'), H(\mathbf{w}'), H(\mathbf{w}')] \right] \quad (32)$$

The operator  $[\cdot, L_z]$  rotates the orbital plane on which  $\mathbf{x}$  lies (by incrementing the argument of the ascending node  $\Omega$ ). The operator  $[\cdot, L_z + L'_z]$ , where  $L'_z$  operates on  $\mathbf{x}'$ , rotates  $\mathbf{x}$  and  $\mathbf{x}'$  through the same angle, so<sup>2</sup>

$$[|\mathbf{x}' - \mathbf{x}|, L_z + L'_z] = 0. \quad (33)$$

Hence taking advantage of the fact that for any  $f, g, h$ ,  $\int d^6 \mathbf{w} [f, g]h = \int d^6 \mathbf{w} f[g, h]$ , we have

$$\begin{aligned} & \int d^6 \mathbf{w}' \left[ \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|}, L_z \right] [f_-(\mathbf{w}'), H(\mathbf{w}')] \\ &= - \int d^6 \mathbf{w}' \left[ \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|}, L'_z \right] [f_-(\mathbf{w}'), H(\mathbf{w}')] \\ &= \int d^6 \mathbf{w}' \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|} \left[ [f_-(\mathbf{w}'), H(\mathbf{w}'), L'_z] \right] \\ &= \int d^6 \mathbf{w}' \frac{\xi G}{|\mathbf{x}' - \mathbf{x}|} \left[ [f_-(\mathbf{w}'), L'_z], H(\mathbf{w}')] \right]. \quad (34) \end{aligned}$$

When this result is used in equation (32), we obtain

$$[Kf_-, L_z] = K[f_-, L_z], \quad (35)$$

so  $K$  commutes with the operator  $[\cdot, L_z]$ . From these commutations it follows that the van Kampen modes provide representations of the group of translations around tori that is generated by  $[\cdot, L_z]$ . This compact Abelian group has only one-dimensional irreps, which can be reduced to multiplication by  $e^{i\alpha}$ . Hence the van Kampen modes can be indexed by the integer  $n_3$  associated with  $\theta_3$ .

## 4.2 Derivation of the DF

Bearing in mind that for an eigenfunction equation (25) yields  $j_\alpha[f_1] = -\omega A_\alpha$ , from equation (23) we have for an eigenfunction

$$-\omega^2 f_- = \left( \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \right)^2 f_- - i\omega f'_0(H) \sum_\alpha A_\alpha \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \Phi^{(\alpha)}(\mathbf{x}). \quad (36)$$

Now we apply the derivative  $\boldsymbol{\Omega} \cdot \partial / \partial \boldsymbol{\theta}$  to both sides and

<sup>2</sup> The operator  $[\cdot, L]$  rotates the orbit within the orbital plane while holding constant  $\theta_1$ , so in the Keplerian case it rotates the ellipse rather than moving the star along its ellipse. If  $\mathbf{w}$  and  $\mathbf{w}'$  lie on the same orbital plane, rotating both orbits within the plane will leave  $|\mathbf{x}' - \mathbf{x}|$  invariant, but if the points lie on different planes,  $|\mathbf{x}' - \mathbf{x}|$  will not be invariant. Hence  $[|\mathbf{x}' - \mathbf{x}|, L] \neq 0$ .

use equations (14) and (19) to eliminate  $f_-$  in favour of  $f_+$ . Collecting terms with  $f_+$  on the left we then have

$$\left\{ \omega^2 + \left( \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \right)^2 \right\} f_+ = f'_0(H) \sum_\alpha A_\alpha \left( \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \right)^2 \Phi^{(\alpha)}(\mathbf{x}). \quad (37)$$

Fourier decomposed in angle variables this becomes

$$\left\{ \omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2 \right\} f_{\mathbf{n}+} = -f'_0(H) (\mathbf{n} \cdot \boldsymbol{\Omega})^2 \sum_\alpha A_\alpha \Phi_{\mathbf{n}}^{(\alpha)}. \quad (38)$$

This equation is analogous to the standard equation for the Laplace-transformed DF  $\bar{f} \equiv \int_0^t dt' f e^{-pt'}$ ,

$$(p + \mathbf{i} \mathbf{n} \cdot \boldsymbol{\Omega}) \bar{f}_{\mathbf{n}}(\mathbf{J}, p) = -f'_0(H) \mathbf{n} \cdot \boldsymbol{\Omega} \sum_\alpha \bar{A}_\alpha(p) \Phi_{\mathbf{n}}^{(\alpha)} + f_{\mathbf{n}}(\mathbf{J}, 0) \quad (39)$$

in that it relates the DF to the driving potential, but there are two significant differences:

1 The rhs of equation (38) doesn't contain an initial condition analogous to  $\hat{f}_{\mathbf{n}}(\mathbf{J}, 0)$  on the rhs of (39). It's not there because we are seeking a normal mode rather than the solution to an initial-value problem.

2 Equation (38) starts with a factor  $\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2$  while equation (39) starts with  $p + \mathbf{i} \mathbf{n} \cdot \boldsymbol{\Omega}$ . Before we divide by this factor we must recognise that when it vanishes, which for a range of frequencies it will over 2d resonant surfaces in action space,  $f_{\mathbf{n}+}$  is unconstrained. This being so, after we've divided by  $\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2$ , we should add a function that's non-zero only on resonant surfaces. Then we have

$$\begin{aligned} -f_{\mathbf{n}+}(\mathbf{J}) &= f'_0(H) \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2}{\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2} \sum_\alpha A_\alpha \Phi_{\mathbf{n}}^{(\alpha)}(\mathbf{J}) \\ &+ g_{\mathbf{n}}(\mathbf{J}) \delta(\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2), \quad (40) \end{aligned}$$

where  $g_{\mathbf{n}}$  is an arbitrary function. Adding this term makes it possible for  $f_{\mathbf{n}+}$  to take whatever value  $g_{\mathbf{n}}$  specifies on the resonant surfaces. (The values taken by  $g_{\mathbf{n}}$  off resonant surfaces are immaterial.)

Multiplying equation (40) by  $\int d^6 \mathbf{w} e^{i\mathbf{n} \cdot \boldsymbol{\theta}} \Phi^{(\alpha')*}(\mathbf{x})$  and summing over  $\mathbf{n}$ , we get

$$\begin{aligned} \frac{\mathcal{E}}{\xi} A_{\alpha'} &= \int d^3 \mathbf{J} d^3 \boldsymbol{\theta} \sum_{\mathbf{n}} e^{i\mathbf{n} \cdot \boldsymbol{\theta}} \Phi^{(\alpha')*}(\mathbf{x}) \\ &\times \left\{ f'_0(H) \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 \sum_\alpha A_\alpha \Phi_{\mathbf{n}}^{(\alpha)}}{\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2} + g_{\mathbf{n}}(\mathbf{J}) \delta(\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2) \right\} \\ &= (2\pi)^3 \mathcal{P} \int d^3 \mathbf{J} \sum_{\mathbf{n}} \left\{ f'_0 \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2}{\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2} \sum_\alpha A_\alpha \Phi_{\mathbf{n}}^{(\alpha')*} \Phi_{\mathbf{n}}^{(\alpha)} \right. \\ &\left. + \Phi_{\mathbf{n}}^{(\alpha')*} g_{\mathbf{n}}(\mathbf{J}) \delta(\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2) \right\}, \quad (41) \end{aligned}$$

where the integral over  $\mathbf{J}$  is a principal value in the sense that actions at which  $\mathbf{n} \cdot \boldsymbol{\Omega} = \pm\omega$  are to be excluded and the large values of the integrand as such points are approached largely cancel during integration. Equation (41) has the form

$$\sum_\alpha M_{\alpha' \alpha} A_\alpha = -B_{\alpha'}, \quad (42)$$

where<sup>3</sup>

$$\begin{aligned}
M_{\alpha'\alpha}(\omega) &\equiv \delta_{\alpha'\alpha} \\
&- \frac{(2\pi)^3 \xi}{\mathcal{E}} \mathcal{P} \int d^3 \mathbf{J} f'_0(H) \sum_{\mathbf{n}} \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2}{\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2} \sum_{\alpha} \Phi_{\mathbf{n}}^{(\alpha')*} \Phi_{\mathbf{n}}^{(\alpha)} \\
B_{\alpha'} &\equiv - \frac{(2\pi)^3 \xi}{\mathcal{E}} \int d^3 \mathbf{J} \sum_{\mathbf{n}} \Phi_{\mathbf{n}}^{(\alpha')*}(\mathbf{J}) g_{\mathbf{n}}(\mathbf{J}) \delta(\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2).
\end{aligned} \tag{43}$$

In a conventional normal-mode analysis we derive an equation<sup>4</sup> which is homogeneous in the disturbance's amplitude (here  $A_{\alpha}$ ) with the consequence that non-trivial solutions exist only when the determinant of a matrix (here  $\mathbf{M}$ ) vanishes. The dispersion relation of Landau modes is the condition for  $|\mathbf{M}|$  to vanish. Our recognition that there can be non-trivial distributions of stars on resonant tori gives rise to non-vanishing  $B_{\alpha}$ , and thus causes  $A_{\alpha}$  to satisfy an inhomogeneous equation that can always be satisfied, so there is no dispersion relation. In the case of a stable system, the frequencies of all Landau modes lie below the real axis so  $\mathbf{M}(\omega)$  has an inverse for any real  $\omega$ . Consequently, for any  $B_{\alpha}$  there is always a unique corresponding  $A_{\alpha}$ . Hence given  $\omega$  and  $\mathbf{n}$ , we can determine the  $A_{\alpha}$  for any function  $g(\mathbf{J})$  on the surface  $\mathbf{n} \cdot \boldsymbol{\Omega} = \omega$ . By equation (6), these coefficients describe the spatial structure that  $g$  generates:

$$\Phi_A(\mathbf{x}) = \sum_{\alpha} A_{\alpha} \Phi^{(\alpha)}(\mathbf{x}) \quad ; \quad \rho_A(\mathbf{x}) = \sum_{\alpha} A_{\alpha} \rho^{(\alpha)}(\mathbf{x}). \tag{44}$$

The  $B_{\alpha}$  turn out to be the analogous expansion coefficients of the gravitationally active part of the density

$$\rho_B \equiv \xi \int d^3 \mathbf{v} \sum_{\mathbf{n}} g_{\mathbf{n}}(\mathbf{J}) e^{i\mathbf{n} \cdot \boldsymbol{\theta}} \delta(\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2). \tag{45}$$

Indeed,

$$\begin{aligned}
-\frac{1}{\mathcal{E}} \int d^3 \mathbf{x} \Phi^{(\alpha)*}(\mathbf{x}) \rho_B &= -\frac{\xi}{\mathcal{E}} \sum_{\mathbf{nn}'} \int d^3 \mathbf{J} \int d^3 \boldsymbol{\theta} \Phi_{\mathbf{n}'}^{(\alpha)*}(\mathbf{J}) e^{-i\mathbf{n}' \cdot \boldsymbol{\theta}} \\
&\quad \times g_{\mathbf{n}}(\mathbf{J}) e^{i\mathbf{n} \cdot \boldsymbol{\theta}} \delta(\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2) \\
&= B_{\alpha}.
\end{aligned} \tag{46}$$

If  $\omega^2 < 0$  (an unstable mode),  $B_{\alpha} = 0$  because the argument of the delta-function in the definition of  $B_{\alpha}$  cannot vanish. So given  $\omega^2 < 0$  an associated DF can be found only

<sup>3</sup> Our matrix  $\mathbf{M}$  is analogous to the matrix  $\boldsymbol{\epsilon}$  of Hamilton et al. (2018) rather than their  $\mathbf{M} = \mathbf{I} - \boldsymbol{\epsilon}$ . The integrand in our  $\mathbf{M}$  differs from that of Hamilton et al. (2018) in that frequencies occur squared because we are working with an operator that is second-order rather than first-order in time. The polarisation and response operators defined in Chapter 5 of Binney & Tremaine (2008) also involve frequencies rather than their squares. Similar operators can be derived from our  $\mathbf{M}$  by decomposition of our integral by partial fractions.

<sup>4</sup> Specifically,

$$\begin{aligned}
\bar{A}_{\alpha}(p) &= - \frac{(2\pi)^3 \xi}{\mathcal{E}} \int d^3 \mathbf{J} \sum_{\mathbf{n}} f'_0(H) \frac{\mathbf{in} \cdot \boldsymbol{\Omega}}{p + \mathbf{in} \cdot \boldsymbol{\Omega}} \\
&\quad \times \sum_{\alpha'} \bar{A}_{\alpha'}(p) [\Phi^{(\alpha)}(\mathbf{n}, \mathbf{J})]^{*} \Phi^{(\alpha')}(\mathbf{n}, \mathbf{J}).
\end{aligned}$$

if  $|\mathbf{M}| = 0$ . Consequently, to prove the system's stability it suffices to show that  $|\mathbf{M}|$  has no zeroes on the imaginary axis. In any case, the normal-mode frequencies are confined to a continuum of real values with the possible addition of discrete pure imaginary values.

### 4.3 Modes and dressing

To understand the physical reality that underlies this mathematics, consider that in the absence of self gravity ( $\xi = 0$ ), a non-trivial distribution of stars with respect to  $\boldsymbol{\theta}$  on resonant surfaces will generate oscillations in the density at frequency  $\omega$  that will persist for ever. The  $\delta$ -function component of the DF (40) of a van Kampen mode represents this phenomenon, and the  $B_{\alpha}$  quantify the spatial form of this driving structure. When  $\xi > 0$ , these oscillations affect the dynamics of all stars, including non-resonant stars. The regular part of the mode's DF (40) describes these sympathetic oscillations of non-resonant stars. The  $A_{\alpha}$  quantify the spatial structure of this ‘‘dressed’’ response to the driver  $g$ .

The functions  $g_{\mathbf{n}}$  are arbitrary, so we may consider the case in which  $g_{\mathbf{n}}$  vanishes for all vectors but one,  $\mathbf{N}$ . Consider now the effect of multiplying equation (40) by  $\int d^6 \mathbf{w} e^{i\mathbf{n} \cdot \boldsymbol{\theta}} \Phi^{(\alpha')*}(\mathbf{x})$  as in the derivation of equation (41) but now *not* summing over  $\mathbf{n}$ . Then we have

$$\begin{aligned}
C_{\alpha'\mathbf{n}} &= (2\pi)^3 \mathcal{P} \int d^3 \mathbf{J} \Phi_{\mathbf{n}}^{(\alpha')*} \left\{ f'_0 \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2}{\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2} \sum_{\alpha} A_{\alpha} \Phi_{\mathbf{n}}^{(\alpha)} \right. \\
&\quad \left. + g_{\mathbf{n}}(\mathbf{J}) \delta(\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2) \right\},
\end{aligned} \tag{47}$$

where

$$C_{\alpha'\mathbf{n}} \equiv - \int d^6 \mathbf{w} \Phi^{(\alpha)*}(\mathbf{x}) e^{i\mathbf{n} \cdot \boldsymbol{\theta}} f_{\mathbf{n}+}, \tag{48}$$

so

$$A_{\alpha} = \frac{\xi}{\mathcal{E}} \sum_{\mathbf{n}} C_{\alpha\mathbf{n}} \tag{49}$$

When  $\mathbf{n} \neq \mathbf{N}$ , the right side of equation (47) has only the term proportional to  $\sum_{\alpha} A_{\alpha} \Phi_{\mathbf{n}}^{(\alpha)}$ , so would vanish when  $\xi$  and therefore  $A_{\alpha}$  vanish. That is, without self-gravity,  $C_{\alpha\mathbf{n}} \neq 0$  only for  $\mathbf{n} = \mathbf{N}$ ; in this case the van Kampen modes could be labelled by  $\mathbf{N}$ . In the presence of self-gravity, we have no reason to expect  $C_{\alpha\mathbf{n}}$  to vanish for  $\mathbf{n} \neq \mathbf{N}$  because equation (49) includes a contribution to  $A_{\alpha}$  from  $C_{\alpha\mathbf{N}} \neq 0$ . Self-gravity has this impact because it stops  $K$  commuting with the operators  $[\cdot, J_r]$  and  $[\cdot, L]$  as discussed above. From the fact that  $K$  does commute with the operator  $[\cdot, L_z]$  it follows that even when  $\xi > 0$ ,  $A_{\alpha\mathbf{n}} = 0$  unless  $n_3 = N_3$ .

When a Landau mode is weakly damped,  $|\mathbf{M}| = 0$  at a frequency  $\omega_0$  that lies just below the real axis, and in consequence  $|\mathbf{M}|$  is small on the real axis just above this zero. In view of equation (42),  $|\mathbf{A}|/|\mathbf{B}|$  will be large in these circumstances. That is, van Kampen modes with frequencies close to  $\Re(\omega_0)$  are ‘‘heavily dressed’’. This is the true significance of Landau modes. Put differently, while van Kampen modes exist at any  $\omega$ , they have a bigger footprint in action space at frequencies that lie close to those of Landau modes.

Whereas a gas ball has at most a finite number of normal modes at a countable number of frequencies, a cluster

has an infinite number of normal modes at every frequency. This difference is a consequence of the likely completeness of normal modes in each system (Section 5) and the fact that much more information is required to specify the DF of a cluster than the state of a gas ball: the disturbance has adiabatically deformed the latter from its equilibrium, so the velocity distribution remains Maxwellian and one only has to specify a mean and dispersion at each location  $\mathbf{x}$ . In a cluster we need to specify the DF in six-dimensional phase space.

Whereas the Hermiticity of  $K$  ensures that van Kampen modes for different  $\omega$  are orthogonal, it falls to us to select from all modes for any given  $\omega$  a complete set of mutually orthogonal modes. That is, we have to select a set of functions  $g(\mathbf{J})$  that generate modes that are orthogonal in the sense  $\langle g_i | g_j \rangle = \delta_{ij}$ . Since  $K$  commutes with  $[\cdot, L_z]$ , we can require modes to be eigenfunctions of this Hermitian operator, and our task is reduced to finding an orthogonal set of vectors whose components are indexed by  $n_1$  and  $n_2$ .

#### 4.4 Energy of a van Kampen mode

Equation (28) says that the energy of a van Kampen mode is just the norm of the odd part of its DF, and equation (40) gives the even part of a mode's DF from which the odd part follows trivially. So the natural next step is to substitute from the second of these equations into the first and express the mode's energy in terms of its parameters  $g_{\mathbf{n}}$  and its potential  $\Phi[f]$ . This exercise proves long, and is made intricate by the singular denominator  $\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2$  in equation (40) for the DF.

In Appendix B we compute  $\langle \tilde{f} | f \rangle$  for modes that have a common frequency  $\omega$  ( $\langle \tilde{f} | f \rangle$  inevitably vanishes if the modes have different frequencies). The result is

$$\langle f_- | \tilde{f}_- \rangle = -(2\pi)^3 \delta(\omega^2 - \tilde{\omega}^2) \int d^3\mathbf{J} \sum_{n_1, n_2} \left( \pi^2 \omega^4 f'_0 \Phi_{\mathbf{n}}^*[f] \Phi_{\mathbf{n}}[\tilde{f}] + \frac{g_{\mathbf{n}}^* \tilde{g}_{\mathbf{n}}}{f'_0} \right) \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2), \quad (50)$$

A pleasing feature of this expression is that it has no reference to potential/density basis functions. The factor  $\delta(\omega^2 - \tilde{\omega}^2)$  on the rhs reflects the orthogonality of modes of different frequencies that follows from the Hermiticity of  $K$ .<sup>5</sup> A remarkable feature of equations (50) is that its action-space integral is confined to the resonant tori, even though the modes very much involve non-resonant stars.

When we set  $\tilde{f} = f$ , the inner product is divergent for finite  $g$  because finite  $g$  generates an  $f_+$  that diverges as the resonant surface in action space is approached, and the divergences on opposite sides of the surface do not cancel because energy density is proportional to  $f^2$ . This result signals that we can have only an infinitesimal number of stars on any resonant surface; that is,  $g$  should be replaced by  $g d\omega$ . Since  $f$  and  $\Phi[f]$  are linear functions of  $g$ , this implies replacement of  $f$  by  $f d\omega$  with  $f$  still related to  $g$  by equation (40), and replacement of  $\Phi[f]$  by  $\Phi[f] d\omega$ . This done, we

obtain the excess energy associated with the perturbation  $f$  by integrating equation (50) over  $\omega$  and  $\tilde{\omega}$ :

$$\begin{aligned} E &= \sum_{n_1, n_2} \int d\omega d\tilde{\omega} \langle \rho_{f_-} | \rho_{\tilde{f}_-} \rangle \\ &= - \sum_{n_1, n_2} \int \frac{d\omega}{2\omega} \int d^3\mathbf{J} \left( \pi^2 \omega^4 f'_0 |\Phi_{\mathbf{n}}[f]|^2 + \frac{|g_{\mathbf{n}}|^2}{f'_0} \right) \\ &\quad \times \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2) \\ &= -\frac{1}{4} \sum_{n_1, n_2} \int d^3\mathbf{J} \left( \pi^2 (\mathbf{n} \cdot \boldsymbol{\Omega})^2 f'_0 |\Phi_{\mathbf{n}}[f]|^2 + \frac{|g_{\mathbf{n}}|^2}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 f'_0} \right) \quad (51) \end{aligned}$$

This is the energy associated with just one mode per frequency, with the integral over  $|\mathbf{J}|$  being effectively a sum over frequencies. To get the whole thermal energy of a cluster, we must sum over the different modes that exist at a given frequency. This sum will be over  $n_3$ , which labels modes with different azimuthal density distributions, and an additional number associated with different choices of the function  $g_{\mathbf{n}}(\mathbf{J})$ , which is effectively a function of the single variable that fixes position along the line  $\mathbf{n} \cdot \boldsymbol{\Omega} = \omega$  in  $(L, J_r)$  space.

There is a striking similarity between the first term in the expression (51) for  $E$  and the expression  $\rho_E = \frac{1}{2} \omega^2 \epsilon_0 A^2$  for the energy density contributed by an electromagnetic wave with vector-potential amplitude  $A$ . In the electromagnetic case one factor of  $\omega$  arises from the quantisation condition  $E = \hbar\omega$  and the other arises from the canonical momentum  $\omega \mathbf{A}$  of the field  $\mathbf{A}$ . The second term in equation (51) has a different structure, however, and this term is arguably more important than the first because  $\Phi[f]$  is driven by  $g$ .

A natural question to ask is why the coefficient of  $|\Phi_{\mathbf{n}}[f]|^2$  in equation (51) is positive, given that gravitational potential energy is inherently negative. The explanation must be that this term encapsulates all the energy, kinetic as well as potential, that's tied up in the disturbance in non-resonant stars that is excited by the resonant stars. In the absence of self-gravity, the non-resonant stars are not disturbed, so this contribution to the energy vanishes with  $\Phi[f]$ .

The second term in the integrand of equation (51) is essentially set by Poisson noise and can be considered a given, while the first term depends on the system's dynamics. The ratio of the two terms is proportional to  $(\mathbf{n} \cdot \boldsymbol{\Omega})^4 f_0'^2$ , so the relative contributions to  $E$  from the resonant driver  $g$  and the non-resonant response  $\Phi$  are sensitive to this factor. In principle  $|\mathbf{n} \cdot \boldsymbol{\Omega}|$  can be made as small as we please at given  $\mathbf{J}$ , but only by going to large  $|\mathbf{n}|$ , and at large  $|\mathbf{n}|$  the projection of  $g_{\mathbf{n}}(\mathbf{J})$  into real space (measured by  $B_\alpha$ ) will be small and thus the response (measured by  $A_\alpha$  and  $\Phi[f]$ ) will be small. Hence the first term in the integrand of equation (51) will be significant only for small  $|\mathbf{n}|$ . The fundamental dipole mode most obviously satisfies this criterion.

This consideration draws attention to short vectors  $\mathbf{n}$  that make  $|\mathbf{n} \cdot \boldsymbol{\Omega}|$  small (if it is small on any torus, it will be small on many tori) because for these vectors the noise component  $g_{\mathbf{n}}$  generates a response at the least cost in energy, so the response is likely to be large. We saw above, moreover, that the response is enhanced when  $\mathbf{n} \cdot \boldsymbol{\Omega}$  is close to the real part of the frequency of a weakly damped Landau mode, because then  $\det(\mathbf{M})$  is small. The fundamental

<sup>5</sup> The appearance of  $\omega^2$  rather than  $\omega$  reflects time-reversal symmetry. From a practical perspective, it ensures that real time dependence ( $\cos \omega t$ ) is possible.

dipole mode has been shown to be weakly damped in typical models (Weinberg 1994; Saha 1991; Hamilton et al. 2018).

#### 4.5 Initialisation of N-body models

Suppose we set up a cluster by randomly sampling an analytic DF. When the selection is complete, the actual DF will differ from the analytic DF by virtue of Poisson noise, so the  $g_{\mathbf{n}}$  will be non-zero and the cluster's van Kampen modes will be excited. The coefficients  $B_{\alpha}$  that quantify the noisiness of the density distribution are unambiguously fixed by the Monte-Carlo selection, but the coefficients  $A_{\alpha}$  that are generated from them via equation (42) will vary with the degree of self-gravity  $\xi$ . Hence the DFs of the modes that sum to the sampled phase-space distribution will depend on  $\xi$ , but their sum must produce the DF sampled regardless of  $\xi$ . When  $\xi = 1$ , the modes are heavily dressed and yet produce the same small (Poisson) fluctuations in density as in the case  $\xi = 0$  of vanishing self-gravity because the contributions of different modes cancel to a considerable extent. This cancellation is particularly pronounced in the case of low-order modes (small  $|\mathbf{n}|$ ), and it occurs because when  $\xi = 1$  the phases of modes are correlated, whereas when  $\xi = 0$  they are probably uncorrelated.

Once we start moving the stars with  $\xi$  set to unity, the phase differences between modes with different frequencies will tend towards uniform distribution in  $(0, 2\pi)$  and cancellations between perturbations to the density will diminish. Consequently, the heavily dressed individual modes will become manifest and the system will become less spherical as Lau & Binney (2019) found empirically. By contrast, when stars are moved in the analytic potential, the initially uniformly distributed phases of the modes obtained with  $\xi = 0$  remain uniformly distributed and no significant change in the density fluctuations will be observed.

The larger the value of  $\xi$ , the larger will be the values of  $A_{\alpha}$  that are generated by the given  $B_{\alpha}$ , so the greater will be the departures from spherical symmetry once the phases of modes have decorrelated.

To obtain a self-consistent realisation of a self-gravitating system one needs to excite the modes for  $\xi = 1$  with random phases, and it is not clear how this can be done without computing the system's modes.

## 5 COMPLETENESS OF VAN KAMPEN MODES

We have defined the van Kampen modes of a stellar system to be the eigenfunctions of Antonov's Hermitian operator  $K$ . In quantum mechanics it is conventional to assume that the eigenfunctions of any Hermitian operator form a complete set although proof of completeness requires the operator to be bounded (e.g. Dieudonné 1969, §11.5), which some operators of physical interest are not. Similarly, much of condensed-matter physics relies on Bloch's theorem that there is a complete set of stationary states for an electron in a crystal that have wavefunctions of the form  $\psi(\mathbf{x}) = e^{i\mathbf{k}\cdot\mathbf{x}}u_p(\mathbf{x})$  with  $u_p(\mathbf{x}+\mathbf{a}) = u_p(\mathbf{x})$  for any lattice vector  $\mathbf{a}$ . The standard derivation of Bloch's theorem (e.g. Elliott & Dawber 1989, §14.4) starts from the observation that if  $\psi(\mathbf{x})$  is a stationary state, then so is  $\psi(\mathbf{x} + \mathbf{a})$ . Hence the

stationary states of a given energy provide a representation of the Abelian translation group. Such groups only have one-dimensional irreducible representations, so the action of the group can be reduced to multiplication by  $e^{i\mathbf{k}\cdot\mathbf{a}}$ . That is, any functions providing a representation of the translation group can be reduced to functions satisfying  $\psi(\mathbf{x} + \mathbf{a}) = e^{i\mathbf{k}\cdot\mathbf{a}}\psi(\mathbf{x})$ , a relation that is clearly satisfied by  $\psi(\mathbf{x}) = e^{i\mathbf{k}\cdot\mathbf{x}}u_p(\mathbf{x})$ .

Analogously, we might argue that the time-translation invariance of  $\partial_t^2 + K$  implies that if  $f(\mathbf{w}, t)$  satisfies  $(\partial_t^2 + K)f = 0$ , then so does  $f(\mathbf{w}, t + \tau)$  and it follows that any set of solutions can be reduced to ones that satisfy  $f(\mathbf{w}, t + \tau) = e^{-i\omega\tau}f(\mathbf{w}, t)$ . Then setting  $t$  to zero we infer that solutions of the form  $f(\mathbf{w}, \tau) = e^{-i\omega\tau}f(\mathbf{w})$  are complete.

The above arguments for the completeness of Bloch waves and van Kampen modes are open to the objection that the theorem regarding the decomposition of representations into irreducible representations requires the group to be compact, which translation groups are not.<sup>6</sup> Hence, the completeness of van Kampen modes cannot be rigorously established by the group-theoretic argument, although similar arguments are widely accepted in physics.

Case (1959) established the completeness of the van Kampen modes of an electrostatic plasma by direct demonstration that any DF  $f(\mathbf{w})$  can be written as a sum of van Kampen modes. It seems likely that a similar demonstration can be accomplished for ergodic stellar systems but this exercise lies beyond the scope of this paper.

## 6 DISCUSSION

### 6.1 Particle dressing

The concept of particle dressing has been central to high-energy physics for over half a century, but has been slow to catch on in stellar dynamics. In galactic dynamics it can be traced at least as far back as Julian & Toomre (1966), who showed that a gas cloud in a galactic disc would attract an entourage of passing stars  $\sim$ ten times more massive than itself. Toomre & Kalnajs (1991) showed that individual disc stars also enhance their masses tenfold by attracting an entourage of other stars. Sellwood & Carlberg (2014) showed that large-amplitude spiral structure emerges from Poisson noise through successive spiral instabilities, but this important process was only firmly connected to particle dressing by Fouvry et al. (2015). Hamilton (2021) made the connection between the BL equation and particle dressing beautifully clear via Rostoker's principle: that it is permissible to compute the effects of discreteness as from the interaction of uncorrelated but dressed particles (Rostoker 1964). Here we have interpreted a van Kampen mode as the result of dressing not one star but an ensemble of stars on a group of resonant tori.

Given that the CBE is the first equation in the BBGKY

<sup>6</sup> The reduction theorem applies only to unitary representations, which associates every group member  $g$  with a unitary operator  $T_g$  on a vector space. If a group is compact, Maschke's operator  $S^2 = \sum_g T_g^\dagger T_g$  can be used to establish that any representation is isomorphic to a unitary representation. In the non-compact case the sum over  $g$  is ill-defined.

hierarchy of equations with the two-particle correlation function set to zero, the importance of dressing for the structure of van Kampen modes may seem paradoxical. The CBE is a mean-field approximation akin to the Weiss theory of magnetism, and embraces correlations that are induced by perturbing fields. Hence it embraces the dressing of resonant tori involved in van Kampen modes. Moreover, when a simulation is started, its DF is inevitably perturbed from the underlying analytic DF and thus its van Kampen modes are excited.

## 6.2 van Kampen vs Landau modes

Normal modes are perhaps the most important single tool in theoretical physics – quantum field theory has even taught us to see particles as excitations of normal modes of the vacuum. Modal analyses have played a significant role in stellar dynamics since the seminal work of Kalnajs (1965), Toomre (Toomre 1964, 1981) and later the prescient work of Weinberg (Weinberg 1993, 1994, 1998, 2001), but in all these studies the modes considered were those of Landau. These ‘modes’ lack key properties of true modes: (i) completeness in the sense that any initial condition can be expressed as a linear combination of modes, and (ii) additivity of energies. These two properties are essential for the use of modes in physics and engineering outside stellar dynamics. The contents pages of the two volumes of Fridman & Polyachenko (1984) explain the focus on Landau modes: the community wanted to establish which equilibrium models are stable, rather than to investigate, as we do, the excursions that stable systems make around equilibrium.

The procedure by which Landau modes are constructed (e.g. Binney & Tremaine 2008, §5.2) is strongly time-asymmetric and as a consequence extracts from a system with time-symmetric dynamics a solution that is strongly marked by the arrow of time: if the DF of a decaying Landau mode at a late time were modified by mapping  $\mathbf{v} \rightarrow -\mathbf{v}$  and  $t \rightarrow -t$  and the system then allowed to evolve under the CBE, the amplitude of the disturbance would grow for only a finite time before decaying away because the original decaying solution was only valid for  $t > 0$ .

Doubt is sometimes cast on the physical standing of van Kampen modes because their DFs contain a delta-function. Actually this feature is a natural consequence of their forming a continuum. Testable predictions of van Kampen modes will always emerge after integration over  $\omega$ , just as in the familiar quantum-mechanical treatment of radiative transitions sensible results emerge only after integration over the frequency of the electromagnetic field,<sup>7</sup> and the delta-functions will disappear in the process. Landau modes are superpositions of van Kampen modes, and they decay as their constituent van Kampen modes drift apart in phase (Case 1959). If you run a decayed Landau mode back in time, the phases move back into alignment for a finite time before drifting apart, so the disturbance grows for only a limited time.

<sup>7</sup> When deriving Fermi’s golden rule, one analogously integrates over the energies of final states.

## 6.3 Thermodynamics of star clusters

When a stellar system is born, its van Kampen modes are assigned particular amplitudes and phases. At birth the phases may be highly correlated, but they will decorrelate on a dynamical timescale. This decorrelation may manifest itself through the emergence of system-scale fluctuations in the density, wandering of the point of highest central density, etc (Lau & Binney 2019; Heggie et al. 2020). On a longer timescale non-linear terms in the CBE will mediate exchanges of energy between modes. We know that the amplitudes of modes are invariant at linear order, and it is likely that they evolve at quadratic order in the perturbations. The 2-body timescale is precisely the timescale associated with terms of quadratic order (Chavanis 2012), so van Kampen modes probably exchange energy on the 2-body timescale.

There is no reason to believe that when a cluster is first realised the amplitudes of its van Kampen modes conform to the Gibbs distribution. We expect exchanges of energy between modes to drive the distribution towards the Gibbs distribution, and the consequential equipartition of energy between modes. A seductive programme of work is to assume equipartition of energy and random phases between the modes of a particular mean-field model and to compare the observables predicted thus with observational data and N-body simulations. The success of such predictions is an open question because the timescale on which modes are exchanging energy and equipartition can be approached is probably the same timescale on which core collapse and evaporation are changing the mean-field model, and thus the modes between which energy is distributed.

In this regard it is instructive to compare the applicability of thermodynamics to star clusters and to classical systems that also have access to states of very high entropy, for example a mixture of two parts hydrogen and one part oxygen, or a diamond, both of which have higher entropy states (water vapour and graphite) that can only be reached by climbing over a significant energy barrier. On account of this barrier, a hydrogen/oxygen mixture and a diamond will extensively explore the configurations accessible with thermal energy regardless of the existence of states of much higher entropy. A stellar system is not denied access to states of high energy by an energy barrier but by a glacial rate of energy transport.

Nevertheless, computing the thermodynamics of a cluster is feasible because the positivity of model energy ensures that the cluster’s constant-energy surfaces have finite volume  $\Omega$ , and it would be very interesting to examine its predictions.

In such a theory the DF would itself be a random variable in addition to the coordinates of stars, which are the random variables whose probability distribution the DF specifies. Testable predictions would emerge from the theory as double expectations: first  $\langle \mathcal{O} \rangle_f = \int d^6\mathbf{w} f(\mathbf{w}) \mathcal{O}(\mathbf{w})$  and then an average of these averages weighted by the probability of each DF  $f$ . Hence a prerequisite of the theory is the ability to assign probabilities to DFs in a rational way. In particular, the probability assigned to a group of DFs must remain unchanged as a cluster evolves under the CBE. In standard statistical mechanics the analogous requirement, that a probability density on phase space be invariant under Hamiltonian evolution, is satisfied by making a priori

probability proportional to the measure of phase-space volume  $d^N q d^N p$  that canonical coordinates  $(q, p)$  deliver. In a companion paper we extend this idea to the space of distribution functions by defining canonical coordinates for this space. It turns out that the energy of a van Kampen mode then takes the form of a sum of Hamiltonians of simple-harmonic oscillators.

#### 6.4 Prior work

To our knowledge the van Kampen modes of a stellar system have previously been considered only by Vandervoort (2003), who followed van Kampen (1955) in deriving the modes directly from the CBE. Antonov (1961) obtained the second-order, Hermitian differential operator  $K$  by splitting the DF into parts even and odd in  $\mathbf{v}$ , but he was focused on proving his stability principle and didn't show that the eigenfunctions of  $K$  are the van Kampen modes. He didn't take advantage of angle-action variables or compute excitation energies. Polyachenko et al. (2021) discussed the relation of Landau and van Kampen modes in the context of the periodic cube, though principally in the unphysical case that the cube's mass exceeds the Jeans mass so the system is unstable. Their work makes very clear that Landau 'modes' lack essential properties of modes, and also illustrates what a treacherous arena the complex plane is: physically ill-motivated changes in the contour of integration over velocity give rise to solutions with radically different properties. In particular they show that Landau's choice of contour breaks the time-reversal symmetry of the underlying problem.

Polyachenko et al. (2021) restrict the term 'van Kampen mode' to modes with real frequencies and speak of the wider class of modes discussed here as 'true modes'. The distinction is only relevant in unstable systems. Then, as we showed in Section 3, the spectrum of  $K$  comprises a section of the real axis and discrete points on the imaginary axis at  $\pm iy$ . The principal point made by Polyachenko et al. (2021) is that the unstable Jeans mode at  $+iy$  is accompanied by a decaying mode at  $-iy$ . This is a trivial consequence of time-reversibility but Polyachenko et al. (2021) show that some effort is required to explain why Landau's analysis misses this mode.

## 7 CONCLUSIONS

As the completeness and precision of astronomical data grow, the oscillations of stellar systems around equilibrium configurations will increase in observational significance. The natural way to produce theoretical predictions of these phenomena is to adapt the techniques of statistical mechanics to stellar dynamics. This paper takes a step in this direction for ergodic stellar systems by focusing attention on the van Kampen modes of stellar systems, which have hitherto been eclipsed by Landau modes.

We showed for the first time that van Kampen modes of an ergodic system are the eigenfunctions of Antonov's second-order Hermitian operator on phase space. In consequence, the modes of a stellar system are either purely sinusoidal or exponentially growing; there are no over-stable

modes or modes comprising decaying oscillations. The frequencies of oscillating modes from a continuum, and the DFs of these modes contain delta-functions which disappear when testable predictions are extracted by integrating over frequencies. Any exponentially growing/decaying modes are isolated in frequency space and their DFs do not contain delta-functions. The energy of an oscillating mode is just the norm of the odd part of its DF. From this it followed that these modes have positive energy. The energy of a growing/decaying mode is identically zero.

We interpreted van Kampen modes as dressed sets of resonant tori. How heavily they are dressed increases with the extent to which the system is self-gravitating, and with proximity in frequency space to a pole of the response matrix  $\mathbf{M}(\omega)$  – the poles of this matrix are each associated with a Landau 'mode'. In the absence of self gravity, the structure of van Kampen modes is very simple, but in the general case the structure is buried in matrix algebra. The general outlines of the modes are nonetheless clear. At every frequency  $\omega$  there are many modes that could be indexed by (i) the integer vector  $\mathbf{n}$  for which  $\mathbf{n} \cdot \boldsymbol{\Omega} = \omega$  for some resonant actions  $\mathbf{J}$ , and (ii) by additional indices that enumerate members of a complete set of functions  $g_{\mathbf{n}}(\mathbf{J})$  on the resonant tori. Thus a star cluster has many more modes than the equivalent gas ball. This fact reflects the large amount of information required to specify a DF rather than the density and pressure in a ball of gas.

When a model cluster is realised, its van Kampen modes acquire non-zero amplitudes by virtue of Poisson noise. The phases of modes evolve on a dynamical timescale while their amplitudes probably evolve on the 2-body timescale. Consequently, when there is an early phase of relaxation in the evolution of a simulated cluster in which system-scale distortions emerge. Consideration of the way modes depend of the degree of self-gravity explains why system-scale distortions are less prominent in simulations that are less self-gravitating.

The positivity of the energies of van Kampen modes opens the door to the application of standard statistical physics to stellar systems: while in the long term systems will drift through core collapse and evaporation to states of ever higher entropy, in the medium term disturbed systems may relax to distributions of energy among van Kampen modes that maximise entropy.

Landau's treatment of perturbations is arguably unhelpful. It involves the choice of a contour of integration that breaks time-reversal symmetry. In any analysis a key role is played by a response operator like that,  $\mathbf{M}(\omega)$ , defined by equation (43). The self-consistent response of the system is given by the inverse of this operator along the real axis. Landau modes are associated with zeros of  $|\mathbf{M}|$ . If the system is unstable, there are pairs of zeros on the positive and negative imaginary axis, and these zeros are associated with growing and decaying van Kampen modes of zero energy. If the system is stable, all zeros lie below the real axis, and these zeros are not associated with true modes. Traditionally they have been associated with 'Landau modes' but we argue that this association is unhelpful because the corresponding dynamics lacks all the properties that make modes central tools of theoretical physics. Zeros that lie near the real axis are important because they imply that  $|\mathbf{M}|$  is small, and therefore the response is massive, along a section

of the real axis. In this section of the real axis, van Kampen modes are heavily dressed and therefore exceptionally important dynamically.

Two directions for future work beckon. One is to discover how to extend the results of this paper to non-ergodic systems. The first step would be to consider DFs of the form  $f(H, L)$  [or equivalently  $f(J_r, L)$ ]. If that were successful, one might proceed to the general case of  $f(\mathbf{J})$ . The immediate issue is of course the ubiquity in the formulae above of  $f'_0$ .

Another beckoning direction is practical implementation of the formulae: decompose likely disturbances  $f_1(\mathbf{w})$  into modes at different degrees of self-gravity  $\xi$ ; finding the distribution  $f_1(\mathbf{w})$  that maximises entropy for given energy of disturbance. A major issue here is choosing an orthogonal basis of modes for each given frequency by choosing a basis of DFs  $g(\mathbf{w})$  on the resonant tori for that frequency. A first step in this direction is probably best taken in the context of the periodic cube, where the angle-action variables and potential theory are both trivial and aligned. An investigation of this type will indicate whether van Kampen modes will prove to be practical tools for the analysis of astronomical data, or are so intractable that, like partition functions, their principal value lies in the insight they give to how macroscopic phenomena emerge from micro-physics.

## ACKNOWLEDGEMENTS

We thank an anonymous referee for an exceptionally careful reading and numerous constructive suggestions, and we thank J. Magorrian and B. Kocsis for helpful comments on early drafts. Our discussions of the completeness of van Kampen modes benefited from discussions with U. Tillmann. Jun Yan Lau gratefully acknowledges support from University College London's Overseas and Graduate Research Scholarships. James Binney is supported by the UK Science and Technology Facilities Council under grant number ST/N000919/1 and by the Leverhulme Trust through an Emeritus Fellowship.

## DATA AVAILABILITY

No new data was generated or analysed in support of this research.

## REFERENCES

- Antoja T. et al., 2018, *Nat*, 561, 360  
 Antonov V. A., 1961, *SovA*, 4, 859  
 Barnes J., Goodman J., Hut P., 1986, *ApJ*, 300, 112  
 Binney J., Schönrich R., 2018, *MNRAS*, 481, 1501  
 Binney J., Tremaine S., 2008, *Galactic Dynamics: Second Edition*. Princeton University Press  
 Bland-Hawthorn J., Tepper-Garcia T., 2020, arXiv e-prints, arXiv:2009.02434  
 Case K. M., 1959, *Annals of Physics*, 7, 349  
 Chavanis P.-H., 2012, *Physica A*, 391, 3680  
 Dieudonné J., 1969, *Foundations of modern analysis*. Academic Press

- Elliott J. P., Dawber P. G., 1989, *Symmetry in physics*. Macmillan  
 Fouvry J. B., Pichon C., Magorrian J., Chavanis P. H., 2015, *A&A*, 584, A129  
 Fridman A. M., Polyachenko V. L., 1984, *Physics of gravitating systems*, vols I and II. Springer  
 Gaia Collaboration, Brown A. G. A. e. a., 2018, *A&A*, 616, A1  
 Hamilton C., 2021, *MNRAS*, 501, 3371  
 Hamilton C., Fouvry J.-B., Binney J., Pichon C., 2018, *MNRAS*, 481, 2041  
 Heggie D. C., Breen P. G., Varri A. L., 2020, *MNRAS*, 492, 6019  
 Heyvaerts J., 2010, *MNRAS*, 407, 355  
 Julian W. H., Toomre A., 1966, *ApJ*, 146, 810  
 Kalnajs A. J., 1965, PhD thesis, Harvard University.  
 Kalnajs A. J., 1976, *ApJ*, 205, 745  
 Kulsrud R. M., Mark J. W. K., 1970, *ApJ*, 160, 471  
 Laporte C. F. P., Minchev I., Johnston K. V., Gómez F. A., 2019, *MNRAS*, 485, 3134  
 Lau J. Y., Binney J., 2019, *MNRAS*, 490, 478  
 Laval G., Mercier C., Pellat R., 1965, *Nucl.Fus.*, 5, 156  
 Magorrian J., 2006, *MNRAS*, 373, 425  
 Malin D. F., Carter D., 1980, *Nat*, 285, 643  
 Morrison P. J., 1980, *Phys.Lett.A*, 80, 383  
 Morrison P. J., Shadwick B. A., 1994, *Acta Physica Polonica A*, 85, 759  
 Polyachenko E. V., Shukhman I. G., Borodina O. I., 2021, *MNRAS*, 503, 660  
 Rostoker N., 1964, *Physics of Fluids*, 7, 479  
 Saha P., 1991, *MNRAS*, 248, 494  
 Sellwood J. A., Carlberg R. G., 2014, *ApJ*, 785, 137  
 Tonry J., Schneider D. P., 1988, *AJ*, 96, 807  
 Toomre A., 1964, *ApJ*, 139, 1217  
 Toomre A., 1981, in *Structure and Evolution of Normal Galaxies*, Fall S. M., Lynden-Bell D., eds., pp. 111–136  
 Toomre A., Kalnajs A. J., 1991, in *Dynamics of Disc Galaxies*, Sundelius B., ed., p. 341  
 van Kampen N. G., 1955, *Physica*, 21, 949  
 Vandervoort P. O., 2003, *MNRAS*, 339, 537  
 Vitral E., Mamon G. A., 2021, *A&A*, 646, A63  
 Weinberg M. D., 1993, *ApJ*, 410, 543  
 Weinberg M. D., 1994, *ApJ*, 421, 481  
 Weinberg M. D., 1998, *MNRAS*, 297, 101  
 Weinberg M. D., 2001, *MNRAS*, 328, 311

## APPENDIX A: MATRIX ELEMENTS OF ANTONOV'S OPERATOR

Here we compute a general matrix element  $\langle \tilde{f}_- | K | f_- \rangle$  of Antonov's operator, where  $f$  and  $\tilde{f}$  any two DFs. Equation (23) yields

$$\begin{aligned} \langle \tilde{f}_- | K | f_- \rangle &= \int \frac{d^6 \mathbf{w}}{f_0^*} \tilde{f}_-^* \left( \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \right)^2 f_- \\ &+ i \frac{\xi}{\mathcal{E}} \int d^6 \mathbf{w} \tilde{f}_-^* \boldsymbol{\Omega} \cdot \frac{\partial}{\partial \boldsymbol{\theta}} \sum_{\alpha} \Phi^{(\alpha)}(\mathbf{x}) j_{\alpha}[f_1]. \quad (\text{A1}) \end{aligned}$$

Since we can write  $d^6 \mathbf{w} = d^3 \mathbf{J} d^3 \boldsymbol{\theta}$ , we can shift the deriva-

tives wrt  $\boldsymbol{\theta}$  around by partial integration and obtain

$$\begin{aligned}
 \langle \tilde{f}_- | K | f_- \rangle &= - \int \frac{d^6 \mathbf{w}}{f'_0} \left( \boldsymbol{\Omega} \cdot \frac{\partial \tilde{f}_-^*}{\partial \boldsymbol{\theta}} \right) \left( \boldsymbol{\Omega} \cdot \frac{\partial f_-}{\partial \boldsymbol{\theta}} \right) \\
 &\quad - i \int d^6 \mathbf{w} \sum_{\alpha} \Phi^{(\alpha)}(\mathbf{x}) \boldsymbol{\Omega} \cdot \frac{\partial \tilde{f}_-^*}{\partial \boldsymbol{\theta}} j_{\alpha}[f_1] \\
 &= - \int \frac{d^6 \mathbf{w}}{f'_0} \left( \boldsymbol{\Omega} \cdot \frac{\partial \tilde{f}_-^*}{\partial \boldsymbol{\theta}} \right) \left( \boldsymbol{\Omega} \cdot \frac{\partial f_-}{\partial \boldsymbol{\theta}} \right) \\
 &\quad - i \frac{\xi}{\xi} \sum_{\alpha} j_{\alpha}^*[\tilde{f}_1] j_{\alpha}[f_1]. \tag{A2}
 \end{aligned}$$

The symmetry of the rhs wrt  $f, \tilde{f}$  implies that  $K$  is Hermitian. Since  $K$  is Hermitian, all its eigenvalues  $\omega^2$  are real and  $K$ 's eigenfunctions are either sinusoidal or show pure exponential growth/decay.

When we set  $\tilde{f} = f$ , we obtain

$$\langle f_- | K | f_- \rangle = \int \frac{d^6 \mathbf{w}}{|f'_0|} \left| \boldsymbol{\Omega} \cdot \frac{\partial f_-}{\partial \boldsymbol{\theta}} \right|^2 - \frac{\xi}{\xi} \sum_{\alpha} |j_{\alpha}[f_1]|^2. \tag{A3}$$

It is interesting to express the rhs of equation (A3) in terms of  $f_+$  using equations (14), (19) and (25). The result is

$$\langle f_- | K | f_- \rangle = \int \frac{d^6 \mathbf{w}}{|f'_0|} \left| \frac{\partial f_+}{\partial t} \right|^2 - \frac{\xi}{\xi} \sum_{\alpha} \left| \frac{\partial A_{\alpha}[f_+]}{\partial t} \right|^2. \tag{A4}$$

In the case that  $f_-$  is an eigenfunction of  $K$  with eigenvalue  $\omega^2$ , we can replace time derivatives by  $-i\omega$  (with  $\omega$  potentially pure imaginary), and equation (A4) becomes

$$\begin{aligned}
 \langle f_- | K | f_- \rangle &= \omega^2 \langle f_- | f_- \rangle \\
 &= |\omega^2| \left\{ \int \frac{d^6 \mathbf{w}}{|f'_0|} |f_+|^2 - \xi \sum_{\alpha} |A_{\alpha}[f_+]|^2 \right\} \\
 &= |\omega^2| \left\{ \langle f_+ | f_+ \rangle - \xi \sum_{\alpha} |A_{\alpha}[f_+]|^2 \right\}. \tag{A5}
 \end{aligned}$$

## APPENDIX B: INNER PRODUCT OF VAN KAMPEN MODES

Here we compute the inner product of two van Kampen modes of the same frequency. We have

$$\begin{aligned}
 \langle f_- | \tilde{f}_- \rangle &= - \int \frac{d^6 \mathbf{w}}{f'_0} f_-^* \tilde{f}_- \\
 &= -(2\pi)^3 \int \frac{d^3 \mathbf{J}}{f'_0} \sum_{\mathbf{n}} f_{\mathbf{n}-}^* \tilde{f}_{\mathbf{n}-}. \tag{B1}
 \end{aligned}$$

Hence  $\langle f_- | \tilde{f}_- \rangle = 0$  unless there is some vector  $\mathbf{n}$  at which both  $f_{\mathbf{n}-}$  and  $\tilde{f}_{\mathbf{n}-}$  are non-zero. This fact confirms that in the absence of self-gravity, when  $f_{\mathbf{n}-}$  vanishes if  $g_{\mathbf{n}}$  vanishes, the sought-after basis modes for a given frequency are indexed by  $\mathbf{n}$  in the sense that the Fourier expansions of their DFs contain only  $\pm \mathbf{n}$ . When  $\xi > 0$ ,  $f_{\mathbf{n}'-}$  is expected to be non-zero when  $g_{\mathbf{n}'} = 0$  providing  $g_{\mathbf{n}} \neq 0$  for a vector  $\mathbf{n}$  such that  $n_3 = n'_3$ .

Equation (40) gives  $f_+$  for a van Kampen mode, and we have seen that in the case of a mode  $|\mathbf{n} \cdot \boldsymbol{\Omega}| f_{\mathbf{n}-} = \omega f_{\mathbf{n}+}$ ,

so

$$\begin{aligned}
 \langle f_- | \tilde{f}_- \rangle &= -(2\pi)^3 \int \frac{d^3 \mathbf{J}}{f'_0} \sum_{n_1, n_2} \frac{\omega \tilde{\omega}}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2} f_{\mathbf{n}+}^* \tilde{f}_{\mathbf{n}+} \\
 &= -(2\pi)^3 \omega \tilde{\omega} \mathcal{P} \int \frac{d^3 \mathbf{J}}{f'_0} \sum_{n_1, n_2} \frac{1}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2} \\
 &\quad \times \left( \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 f'_0}{\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2} \sum_{\alpha} A_{\alpha}^* \Phi_{\mathbf{n}}^{(\alpha)*}(\mathbf{J}) + g_{\mathbf{n}}^*(\mathbf{J}) \delta(\omega^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2) \right) \\
 &\quad \times \left( \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 f'_0}{\tilde{\omega}^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2} \sum_{\alpha} \tilde{A}_{\alpha} \Phi_{\mathbf{n}}^{(\alpha)}(\mathbf{J}) + \tilde{g}_{\mathbf{n}}(\mathbf{J}) \delta(\tilde{\omega}^2 - (\mathbf{n} \cdot \boldsymbol{\Omega})^2) \right), \tag{B2}
 \end{aligned}$$

where quantities associated with the mode  $\tilde{f}$  are marked by tildes. When we multiply out the big brackets, we get a term with two denominators of the form  $(\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2$ . The integral over these is to be interpreted as a principal value, that is by excluding points at which the denominator vanishes. We use the identity

$$\begin{aligned}
 \mathcal{P} \frac{1}{x-x_1} \mathcal{P} \frac{1}{x-x_2} &= \mathcal{P} \frac{1}{x_1-x_2} \left( \mathcal{P} \frac{1}{x-x_1} - \mathcal{P} \frac{1}{x-x_2} \right) \\
 &\quad + \pi^2 \delta(x-x_1) \delta(x_1-x_2) \tag{B3}
 \end{aligned}$$

to rewrite this term as

$$\begin{aligned}
 \int d^3 \mathbf{J} \dots &= \mathcal{P} \int \frac{d^3 \mathbf{J}}{f'_0} \sum_{n_1, n_2} (\mathbf{n} \cdot \boldsymbol{\Omega} f'_0)^2 \sum_{\alpha\beta} A_{\alpha}^* \tilde{A}_{\beta} \Phi_{\mathbf{n}}^{(\alpha)*} \Phi_{\mathbf{n}}^{(\beta)} \\
 &\quad \times \left\{ \frac{1}{\omega^2 - \tilde{\omega}^2} \left( \frac{1}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2} - \frac{1}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \tilde{\omega}^2} \right) \right. \\
 &\quad \left. + \pi^2 \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2) \delta(\omega^2 - \tilde{\omega}^2) \right\}. \tag{B4}
 \end{aligned}$$

The cross terms in the product of equation (B2) can be written

$$\begin{aligned}
 \int d^3 \mathbf{J} \dots &= \int \frac{d^3 \mathbf{J}}{f'_0} \frac{f'_0}{\omega^2 - \tilde{\omega}^2} \sum_{n_1, n_2} \\
 &\quad \times \left( \sum_{\alpha} A_{\alpha}^* \Phi_{\mathbf{n}}^{(\alpha)*} \tilde{g}_{\mathbf{n}} \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \tilde{\omega}^2) \right. \\
 &\quad \left. - \sum_{\beta} \tilde{A}_{\beta} \Phi_{\mathbf{n}}^{(\beta)} g_{\mathbf{n}}^* \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2) \right), \tag{B5}
 \end{aligned}$$

while the term involving  $g_{\mathbf{n}}^* \tilde{g}_{\mathbf{n}}$  can be written

$$\int d^3 \mathbf{J} \dots = \int \frac{d^3 \mathbf{J}}{f'_0} \sum_{n_1, n_2} \frac{g_{\mathbf{n}}^* \tilde{g}_{\mathbf{n}}}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2} \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2) \delta(\omega^2 - \tilde{\omega}^2). \tag{B6}$$

Adding these fragments together and reinstating the prefac-

tor in equation (B2) we have

$$\begin{aligned}
 \langle f_- | \tilde{f}_- \rangle &= -(2\pi^3)\omega\tilde{\omega} \mathcal{P} \int \frac{d^3\mathbf{J}}{f'_0} \sum_{n_1, n_2} \left\{ \frac{f'_0}{\omega^2 - \tilde{\omega}^2} \left[ \right. \right. \\
 &\times \left( \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 f'_0}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2} \sum_{\alpha} A_{\alpha}^* \Phi_{\mathbf{n}}^{(\alpha)*} + g_{\mathbf{n}}^* \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2) \right) \sum_{\beta} \tilde{A}_{\beta} \Phi_{\mathbf{n}}^{(\beta)} \\
 &- \left. \left. \left( \frac{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 f'_0}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \tilde{\omega}^2} \sum_{\beta} \tilde{A}_{\beta} \Phi_{\mathbf{n}}^{(\beta)} + \tilde{g}_{\mathbf{n}} \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \tilde{\omega}^2) \right) \sum_{\alpha} A_{\alpha}^* \Phi_{\mathbf{n}}^{(\alpha)*} \right] \right. \\
 &+ \left( \pi^2 (\mathbf{n} \cdot \boldsymbol{\Omega} f'_0)^2 \sum_{\alpha\beta} A_{\alpha}^* \tilde{A}_{\beta} \Phi_{\mathbf{n}}^{(\alpha)*} \Phi_{\mathbf{n}}^{(\beta)} + \frac{g_{\mathbf{n}}^* \tilde{g}_{\mathbf{n}}}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2} \right) \\
 &\quad \left. \times \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2) \delta(\omega^2 - \tilde{\omega}^2) \right\} \\
 &= -(2\pi)^3 \frac{\omega\tilde{\omega}}{\omega^2 - \tilde{\omega}^2} \int d^3\mathbf{J} \sum_{n_1, n_2} \left( f_{\mathbf{n}+}^* \sum_{\beta} \tilde{A}_{\beta} \Phi_{\mathbf{n}}^{(\beta)} \right. \\
 &\quad \left. - \tilde{f}_{\mathbf{n}+} \sum_{\alpha} A_{\alpha}^* \Phi_{\mathbf{n}}^{(\alpha)*} \right) \\
 &- (2\pi)^3 \omega\tilde{\omega} \int d^3\mathbf{J} \sum_{n_1, n_2} \left( \pi^2 (\mathbf{n} \cdot \boldsymbol{\Omega})^2 f'_0 \sum_{\alpha\beta} A_{\alpha}^* \tilde{A}_{\beta} \Phi_{\mathbf{n}}^{(\alpha)*} \Phi_{\mathbf{n}}^{(\beta)} \right. \\
 &\quad \left. + \frac{g_{\mathbf{n}}^* \tilde{g}_{\mathbf{n}}}{(\mathbf{n} \cdot \boldsymbol{\Omega})^2 f'_0} \right) \delta((\mathbf{n} \cdot \boldsymbol{\Omega})^2 - \omega^2) \delta(\omega^2 - \tilde{\omega}^2). \quad (\text{B7})
 \end{aligned}$$

Now

$$\begin{aligned}
 A_{\alpha} &= \int d^6\mathbf{w} \Phi^{(\alpha)*} f_+ \\
 &= \int d^3\mathbf{J} d^3\boldsymbol{\theta} \sum_{\mathbf{n}} \Phi_{\mathbf{n}}^{(\alpha)*} e^{-i\mathbf{n} \cdot \boldsymbol{\theta}} f_+ = \sum_{\mathbf{n}} \Phi_{\mathbf{n}}^{(\alpha)*} f_{\mathbf{n}+}, (\text{B8})
 \end{aligned}$$

so the first integral in our final expression for  $\langle f_- | \tilde{f}_- \rangle$  vanishes because its integrand is  $\sum_{\alpha} (\tilde{A}_{\alpha} A_{\alpha}^* - A_{\alpha}^* \tilde{A}_{\alpha})$ . With some further simplifications we can write the inner product in the form given by equation (50).