

THE AMPLITUDE OF MOTZKIN PATHS










HELMUT PRODINGER

ABSTRACT. The amplitude of Motzkin paths was recently introduced, which is basically twice the height. We analyze this parameter using generating functions.

1. INTRODUCTION

A Motzkin path consists of up-steps, down-steps, and horizontal steps, see sequence A091965 in [7] and the references given there. As Dyck paths, they start at the origin and end, after n steps again at the x -axis, but are not allowed to go below the x -axis. The height of a Motzkin path is the highest y -coordinate that the path reaches. The average height of such random paths of length n was considered in an early paper [6], it is asymptotically given by $\sqrt{\frac{\pi n}{3}}$.

In the recent paper [2] an interesting new concept was introduced: the *amplitude*. It is basically twice the height, but with a twist. If there exists a horizontal step on level h , which is the height, the amplitude is $2h + 1$, otherwise it is $2h$. To clarify the concept, we created a list of all 9 Motzkin paths of length 4 with height and amplitude given.

Motzkin path	horizontal on maximal level	height	amplitude
	Yes	0	1
	No	1	2
	No	1	2
	Yes	1	3
	Yes	1	3
	Yes	1	3
	No	1	2
	No	1	2
	No	2	4

2010 *Mathematics Subject Classification.* 05A15.

Key words and phrases. Motzkin paths, height, amplitude, generating functions, asymptotics.

Now let $N^{\leq h}(z)$ be the generating function of Motzkin paths of height $\leq h$, but where horizontal steps on level h are forbidden. The system to compute this is quite similar:

$$\begin{pmatrix} 1-z & -z & 0 & \dots & & \\ -z & 1-z & -z & 0 & \dots & \\ 0 & -z & 1-z & -z & 0 & \dots \\ \ddots & \ddots & \ddots & & & \\ & & & -z & \mathbf{1} & \end{pmatrix} \begin{pmatrix} g_0 \\ g_1 \\ g_2 \\ \vdots \\ g_h \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

The only difference in the matrix is the entry in the last row, written in boldface. Let D_n^* be the determinant of this system matrix, with n rows and columns. Again,

$$N^{\leq h}(z) = g_0 = \frac{D_h^*}{D_{h+1}^*}.$$

Expanding the matrix along the last row, we find

$$D_n^* = D_{n-1} - z^2 D_{n-2} = \frac{1}{1-v} \frac{1-v^{2n+1}}{(1+v+v^2)^n}$$

and

$$N^{\leq h}(z) = g_0 = \frac{D_h^*}{D_{h+1}^*} = (1+v+v^2) \frac{1-v^{2h+1}}{1-v^{2h+3}}.$$

Now let us consider

$$M^{\leq h}(z) - N^{\leq h}(z).$$

There is obviously a lot of cancellation going on. The objects which are still counted, have height $= h$, and *have* horizontal steps on level h . That is one of the two quantities that we wanted to compute, and we get

$$\begin{aligned} \text{Horiz}_h(z) &= (1+v+v^2) \frac{1-v^{2h+2}}{1-v^{2h+4}} - (1+v+v^2) \frac{1-v^{2h+1}}{1-v^{2h+3}} \\ &= (1+v+v^2)(1-v^{-2}) \left[\frac{v^{2h+4}}{1-v^{2h+4}} - \frac{v^{2h+3}}{1-v^{2h+3}} \right]. \end{aligned}$$

Similarly, considering $N^{\leq h}(z) - M^{\leq h-1}(z)$, we find that only objects are counted that have height $= h$, and *no* horizontal steps on level h . Thus

$$\begin{aligned} \text{No-Horiz}_h(z) &= (1+v+v^2) \left[\frac{1-v^{2h+1}}{1-v^{2h+3}} - \frac{1-v^{2h}}{1-v^{2h+2}} \right] \\ &= (1+v+v^2)(1-v^{-2}) \left[\frac{v^{2h+3}}{1-v^{2h+3}} - \frac{v^{2h+2}}{1-v^{2h+2}} \right]. \end{aligned}$$

As a check, we get

$$\begin{aligned} \text{Horiz}_h(z) + \text{No-Horiz}_h(z) &= (1 + v + v^2)(1 - v^{-2}) \left[\frac{v^{2h+4}}{1 - v^{2h+4}} - \frac{v^{2h+2}}{1 - v^{2h+2}} \right] \\ &= M^{\leq h}(z) - M^{\leq h-1}(z) = M^{=h}(z). \end{aligned}$$

3. EXPLICIT ENUMERATIONS

All our generating functions contain the term

$$(1 + v + v^2)(1 - v^{-2}) \frac{v^{2h+a}}{1 - v^{2h+a}} = (1 + v + v^2)(1 - v^{-2}) \sum_{k \geq 1} v^{k(2h+a)}$$

for various values of a . We show how to compute the coefficient of z^n in this. It will be done using contour integration. The contour is a small circle in the z -plane or v -plane.

$$\begin{aligned} [z^n](1 + v + v^2)(1 - v^{-2}) \frac{v^{2h+a}}{1 - v^{2h+a}} &= \frac{1}{2\pi i} \oint \frac{dz}{z^{n+1}} (1 + v + v^2)(1 - v^{-2}) \sum_{k \geq 1} v^{k(2h+a)} \\ &= \frac{1}{2\pi i} \oint \frac{dv(1 - v^2)}{(1 + v + v^2)^2} \frac{(1 + v + v^2)^{n+1}}{v^{n+1}} (1 + v + v^2)(1 - v^{-2}) \sum_{k \geq 1} v^{k(2h+a)} \\ &= - \sum_{k \geq 1} [v^{n+2-k(2h+a)}] (1 - v^2)^2 (1 + v + v^2)^n \\ &= - \sum_{k \geq 1} \left[\binom{n, 3}{n+2-k(2h+a)} - 2 \binom{n, 3}{n-k(2h+a)} + \binom{n, 3}{n-2-k(2h+a)} \right]. \end{aligned}$$

In this computation, we used the notion of *trinomial* coefficients:

$$\binom{n, 3}{k} = [t^k](1 + t + t^2)^n.$$

4. THE AVERAGE AMPLITUDE

Here we compute:

$$\begin{aligned} &\sum_{h \geq 0} (2h + 1) \text{Horiz}_h(z) + \sum_{h \geq 1} (2h) \text{No-Horiz}_h(z) \\ &= (1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 0} (2h + 1) \left[\frac{v^{2h+4}}{1 - v^{2h+4}} - \frac{v^{2h+3}}{1 - v^{2h+3}} \right] \\ &\quad + (1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 0} (2h) \left[\frac{v^{2h+3}}{1 - v^{2h+3}} - \frac{v^{2h+2}}{1 - v^{2h+2}} \right] \\ &= -(1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 0} \frac{v^{2h+3}}{1 - v^{2h+3}} \end{aligned}$$

$$\begin{aligned}
& + (1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 0} (2h + 1) \frac{v^{2h+4}}{1 - v^{2h+4}} \\
& - (1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 0} (2h + 2) \frac{v^{2h+4}}{1 - v^{2h+4}} \\
& = -(1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 0} \frac{v^{2h+3}}{1 - v^{2h+3}} - (1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 1} \frac{v^{2h+2}}{1 - v^{2h+2}} \\
& = -(1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 0} \frac{v^{2h+1}}{1 - v^{2h+1}} + (1 + v + v^2)(1 - v^{-2}) \frac{v}{1 - v} \\
& - (1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 1} \frac{v^{2h}}{1 - v^{2h}} + (1 + v + v^2)(1 - v^{-2}) \frac{v^2}{1 - v^2} \\
& = -(1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 0} \frac{v^{2h+1}}{1 - v^{2h+1}} \\
& - (1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 1} \frac{v^{2h}}{1 - v^{2h}} - \frac{(1 + 2v)(1 + v + v^2)}{v} \\
& = -(1 + v + v^2)(1 - v^{-2}) \sum_{h \geq 1} \frac{v^h}{1 - v^h} - \frac{(1 + 2v)(1 + v + v^2)}{v}.
\end{aligned}$$

To find asymptotics from here, we need the local expansion of this generating function around $v \sim 1$, or, equivalently, $z \sim \frac{1}{3}$. See [3] for more explanations of how this method works; it also involves singularity analysis of generating functions [4]. While this combined approach of Mellin transform and singularity analysis has been used for more than 30 years, we would like to cite [5], where many technical details have been worked out in a similar instance. For example, the expansion of the sum that we need is derived there. We combine all the expansions:

$$\left(6(1 - v) + \dots\right) \left(-\frac{\log(1 - v)}{1 - v} + \frac{\gamma}{1 - v} + \dots\right) - 9 + 6(1 - v) + \dots = -6 \log(1 - v) + \dots$$

Translating this into the z -world means $1 - v \sim \sqrt{3}\sqrt{1 - 3z}$, and then

$$[z^n] \left(-6 \log(1 - v)\right) \sim [z^n] \left(-6 \log \sqrt{1 - 3z}\right) = 3 \frac{3^n}{n} = \frac{3^{n+1}}{n}.$$

The total number of Motzkin paths is

$$[z^n] \frac{1 - z - \sqrt{1 - 2z - 3z^2}}{2z^2} \sim [z^n] \left(3 - 3\sqrt{3}\sqrt{1 - 3z}\right) \sim 3\sqrt{3} \frac{3^n}{2\sqrt{\pi}n^{3/2}}.$$

Taking quotients, we get the average amplitude of random Motzkin paths of length n , which is asymptotic to

$$2\sqrt{\frac{\pi n}{3}}.$$

This is about twice as much as the average height of random Motzkin paths, which is what we expected.

Now we consider

$$\begin{aligned} \sum_{h \geq 0} \text{No-Horiz}_h(z) &= (1+v+v^2)(1-v^{-2}) \sum_{h \geq 0} \left[\frac{v^{2h+3}}{1-v^{2h+3}} - \frac{v^{2h+2}}{1-v^{2h+2}} \right] \\ &= (1+v+v^2)(1-v^{-2}) \sum_{h \geq 1} \left[\frac{v^{2h-1}}{1-v^{2h-1}} - \frac{v^{2h}}{1-v^{2h}} \right] + \frac{(1+v)(1+v+v^2)}{v} \\ &= (1+v+v^2)(1-v^{-2}) \sum_{h \geq 1} \frac{v^h}{1-v^h} (-1)^{h-1} + \frac{(1+v)(1+v+v^2)}{v} \end{aligned}$$

This needs to be expanded about $v = 1$:

$$\begin{aligned} (1+v+v^2)(1-v^{-2}) &\sim -6(1-v) - 3(1-v)^2 + \dots; \\ \frac{(1+v)(1+v+v^2)}{v} &\sim 6 - 3(1-v) + \dots; \end{aligned}$$

for the remaining sum we need the Mellin transform [3]. Set $v = e^{-t}$, and transform

$$\mathcal{M} \sum_{h \geq 1} \frac{v^h}{1-v^h} (-1)^{h-1} = \mathcal{M} \sum_{h, k \geq 1} e^{-thk} (-1)^{h-1} = \Gamma(s) \zeta(s)^2 (1-2^{1-s}).$$

By the Mellin inversion formula:

$$\sum_{h \geq 1} \frac{v^h}{1-v^h} (-1)^{h-1} = \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \Gamma(s) \zeta(s)^2 (1-2^{1-s}) t^{-s} ds.$$

The line of integration will be shifted to the left, and the collected residues constitute the expansion about $t = 0$:

$$\sum_{h \geq 1} \frac{v^h}{1-v^h} (-1)^{h-1} \sim \frac{\log 2}{t} - \frac{1}{4} + \frac{t}{48} \sim \frac{\log 2}{1-v} - \frac{\log 2}{2} - \frac{1}{4} + \dots.$$

Combining,

$$\sum_{h \geq 0} \text{No-Horiz}_h(z) \sim 6 - 6 \log 2 - \frac{3}{2}(1-v) + \dots.$$

But

$$\frac{1-z-\sqrt{1-2z-3z^2}}{2z^2} = 1+v+v^2 \sim 3-3(1-v) + \dots.$$

Comparing the coefficients of $1 - v$, we find that asymptotically about half of the Motzkin paths belong to the ‘No-horizontal’ and about half of the Motzkin paths belong to the ‘horizontal’ class. Again, this is intuitively clear, once one sees the generating functions of both families.

REFERENCES

- [1] Louis Comtet. *Advanced combinatorics*. D. Reidel Publishing Co., Dordrecht, enlarged edition, 1974. The art of finite and infinite expansions.
- [2] Julien Courtiel, Andrew Elvey Price, and Irène Marcovici. Bijections between walks inside a triangular domain and Motzkin paths of bounded amplitude. *The electronic journal of combinatorics*, 28(2)(2):#P2.6h, 2021.
- [3] Philippe Flajolet, Xavier Gourdon, and Philippe Dumas. Mellin transforms and asymptotics: harmonic sums. volume 144, pages 3–58. 1995. Special volume on mathematical analysis of algorithms.
- [4] Philippe Flajolet and Andrew Odlyzko. Singularity analysis of generating functions. *SIAM J. Discrete Math.*, 3(2):216–240, 1990.
- [5] Clemens Heuberger, Helmut Prodinger, and Stephan Wagner. The height of multiple edge plane trees. *Aequationes Math.*, 90(3):625–645, 2016.
- [6] Helmut Prodinger. The average height of a stack where three operations are allowed and some related problems. *J. Combin. Inform. System Sci.*, 5(4):287–304, 1980.
- [7] Neil J. A. Sloane. The on-line encyclopedia of integer sequences (<https://oeis.org/>). *Notices Amer. Math. Soc.*, 65(9):1062–1074, 2018.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STELLENBOSCH 7602, STELLENBOSCH, SOUTH AFRICA AND DEPARTMENT OF MATHEMATICS AND MATHEMATICAL STATISTICS, UMEA UNIVERSITY, 907 36 UMEA, SWEDEN

Email address: `hproding@sun.ac.za`