

SUPERDENSITY OF NON-INTEGRABLE BILLIARDS AND DIOPHANTINE APPROXIMATION

J. BECK AND W.W.L. CHEN

ABSTRACT. Consider a finite polysquare or square tiled region, a connected, but not necessarily simply-connected, polygonal region tiled with aligned unit squares. Using ideas from diophantine approximation, we prove that a half-infinite billiard orbit in such a region is superdense, a best possible form of time-quantitative density, if and only if the initial slope of the orbit is a badly approximable number.

As the traditional approach to questions of density and uniformity via ergodic theory depends on results such as Birkhoff's ergodic theorem which are essentially time-qualitative in nature and do not appear to lead naturally to time-quantitative statements, we appeal to a non-ergodic approach that is based largely on number theory and combinatorics. In particular, we use the famous 3-distance theorem in diophantine approximation combined with an iterative process.

This paper improves on an earlier result of the authors and Yang [3] where it is shown that badly approximable numbers that satisfy a quite severe technical restriction on the digits of their continued fractions lead to superdensity. Here we overcome this technical impediment.

1. INTRODUCTION

Over the last 40 years or so, the subject of polygonal billiards has become a very active and fruitful field of research, following the pioneering results of Katok and Zemlyakov in 1975 concerning density, and of Kerckhoff, Masur and Smillie in 1986 concerning uniform distribution.

We say that a polygon P is *rational* if the angle at every vertex of P is a rational multiple of π .

Theorem (Katok–Zemlyakov [8]). *Apart from a countable set of initial directions, any half-infinite billiard orbit in a rational polygon P is dense in P unless it hits a vertex of P and becomes undefined.*

Theorem (Kerckhoff–Masur–Smillie [9]). *For almost all initial directions, any half-infinite billiard orbit in a rational polygon P is uniformly distributed in P unless it hits a vertex of P and becomes undefined.*

We restrict our attention to questions on density and the special case when the region under consideration is what we call here a finite *polysquare region*, or a finite *square tiled region* in the terminology of dynamical systems. More precisely, this is an arbitrary connected, but not necessarily simply-connected, polygon P on the plane which is tiled with unit squares, assumed to be closed, that we call the *atomic squares* of P , and which satisfies the following conditions:

- (i) Any two atomic squares in P either are disjoint, or intersect at a single point, or have a common edge.
- (ii) Any two atomic squares in P are joined by a chain of atomic squares where any two neighbors in the chain have a common edge.

2010 *Mathematics Subject Classification.* 11K38, 37E35.

Key words and phrases. geodesics, billiards, density.

Note that P may have *holes*, and we also allow *whole barriers* which are horizontal or vertical *walls* that consist of one or more boundary edges of atomic squares. For example, the polysquare region in the picture on the left in Figure 1.1 has 5 atomic squares, whereas the polysquare region in the picture on the right in Figure 1.1 has 32 atomic squares, 2 holes as well as 3 horizontal walls and 4 vertical walls.




Figure 1.1: some polysquare-shaped billiard tables

Furthermore, a finite polysquare region can be converted to a finite *polysquare surface*, or *square tiled surface*, by pairwise identification of the boundary horizontal edges and pairwise identification of the boundary vertical edges.

A very natural question that arises from the result of Katok and Zemlyakov concerns those initial directions where the billiard orbits exhibit the best form of time-quantitative density which we call *superdensity*.

We say that a half-infinite billiard orbit $\mathcal{B}(t)$, $t \geq 0$, equipped with the usual arc-length parametrization, in a finite polysquare region P is superdense in P if there exists an absolute constant $C_1 = C_1(P; \mathcal{B}) > 0$ such that, for every integer $n \geq 1$, the initial segment $\mathcal{B}(t)$, $0 \leq t \leq C_1 n$, of the orbit gets $(1/n)$ -close to every point of P .

This concept of superdensity is a best possible form of time-quantitative density, in the sense that the linear length $C_1 n$ cannot be replaced by any sublinear length $o(n)$ as $n \rightarrow \infty$. For a simple proof of this; see [3, Section 6.1].

In this paper, a billiard *always* means the ideal case of a point-billiard that exhibits the simplest elastic collision with the boundary or a wall, where the angle of incidence equals the angle of reflection, as illustrated in Figure 1.1.

Apart from the work of Katok and Zemlyakov, our study is also motivated by the following integrable special case when the polysquare region under consideration is the unit square. It is well known that the distribution of the irrational rotation sequence $n\alpha \bmod 1$, $n = 1, 2, 3, \dots$, is intimately related to the distribution of half-infinite geodesics of slope α on the unit torus $[0, 1)^2$. Using this link, an old result of Khinchin [11, Theorem 26] then implies that a half-infinite geodesic is superdense on the unit torus $[0, 1)^2$ if and only if the slope is a badly approximable number. The first result concerning superdensity of a billiard orbit, namely the special case of the square billiard, follows almost immediately from this.

Theorem (Khinchin). *Any half-infinite billiard orbit is superdense in the unit square if and only if the initial slope of the orbit is a badly approximable number.*

The missing link from the geodesic on the unit torus to the billiard orbit in the unit square is provided by König and Szücs [12] and often known as the unfolding of

the square billiard. Reflecting the unit square across a vertical edge and a horizontal edge, we obtain a 4-fold covering of the unit square.

In Figure 1.2, the 2×2 torus in the picture on the right is a 4-fold covering of the unit square in the picture on the left, with the covering map

$$(x, y) \rightarrow (\min\{2 - x, x\}, \min\{2 - y, y\}), \quad (x, y) \in [0, 2)^2,$$

where $0 \leq \{z\} < 1$ denotes the fractional part of a real number z . Clearly this covering map projects a geodesic on the 2×2 torus $[0, 2)^2$ to a billiard orbit in the unit square $[0, 1]^2$. Since we can tile the whole plane with 2×2 squares, the König–Szücs construction has the remarkable property that it unfolds the zig-zag billiard trajectory in the square $[0, 1]^2$ to a straight line on the plane.




Figure 1.2: unfolding of the square billiard

Remark. The statement and an alternative proof of Khinchin’s result can be found in [3, Lemma 6.1.1].

Combining the König–Szücs unfolding argument with some well-known results of Weyl [20], Hardy and Littlewood [6, 7], Ostrowski [14] and Khinchin [10, 11] concerning geodesics on the unit torus, we obtain a relatively simple and almost complete theory of the square billiard. We may say therefore that the square billiard is essentially fully understood. The same holds for the rectangle billiard where unfolding works equally well.

The purpose of this paper is to establish the following more general result.

Theorem 1. *Let P be an arbitrary finite polysquare region. Then any half-infinite billiard orbit that does not hit a vertex of P is superdense in P if and only if the initial slope of the orbit is a badly approximable number.*

Theorem 1 connects number theory to physics, since billiard clearly represents a mechanical system.

The main difficulty of the generalization from the unit square to an arbitrary finite polysquare region is that billiard flow in many of these finite polysquare regions has orbit-singularities, which lead to unpredictable *slow chaos*. Such dynamical systems are called *non-integrable*. They are in sharp contrast to billiard flow in the unit square which is singularity-free, and exhibits well predictable long-term behavior.

Unfolding works perfectly well for billiards in the whole class of rational polygons. Every rational polygon billiard has a finite *covering space*, which is a flat surface with 1-direction geodesic flow, and the covering map projects this flow to billiard flow in the polygon. This covering space is obtained by similar iterated unfoldings of the polygon, and the necessary boundary pairings, to make this a closed surface, come from identifying parallel edges. For the details, see [13].

We illustrate this general covering space construction technique in the special case of billiards in finite polysquare regions. In Figure 1.3, the picture on the left

shows billiard in a finite polysquare region, and the picture on the right shows its corresponding 4-copy construction as a consequence of König–Szücs unfolding. This is a polysquare surface, where edge identifications are given by translation. The same 4-copy construction technique works for billiard in any finite polysquare region.




Figure 1.3: unfolding of a polysquare-shaped billiard

Needless to say, the concept of superdensity for a geodesic on a polysquare surface can be defined in a way analogous to superdensity of a billiard orbit in a polysquare region.

We shall prove the following more general result which clearly implies Theorem 1.

Theorem 2. *Let \mathcal{P} be an arbitrary finite polysquare surface. A half-infinite geodesic that does not hit a vertex of \mathcal{P} is superdense on \mathcal{P} if and only if the slope of the geodesic is a badly approximable number.*

Recall that an irrational number α , with continued fraction

$$\alpha = [a_0; a_1, a_2, a_3, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}, \quad (1.1)$$

is said to be badly approximable if there exists a constant A such that the continued fraction digits $a_i \leq A$ for every $i = 0, 1, 2, 3, \dots$.

In a recent series of papers [1, 2, 3, 4], the authors and their co-authors are able to establish a number of results which describe the long-term *time-quantitative* behavior in many flat dynamical systems. In particular, a weaker form of Theorem 2 is established, where it is shown that superdensity follows if the slope α given by (1.1) satisfies the additional technical requirement that the digits $a_0, a_1, a_2, a_3, \dots$ are all integer multiples of the street-LCM of the finite polysquare surface under consideration. Here the street-LCM of a finite polysquare surface is the lowest common multiple of the lengths of the horizontal and the vertical streets of the surface. While this may not cover all badly approximable slopes, it nevertheless gives an uncountable set of slopes which guarantee superdensity. In Theorem 2, we remove this technical impediment.

Theorem 2 is an *if and only if* type result, where one of the two implications is a straightforward corollary of Khinchin’s theorem. Indeed, 1-direction geodesic flow on a finite polysquare surface *modulo one* becomes 1-direction geodesic flow on the unit torus $[0, 1)^2$, which implies that a superdense geodesic must have a badly approximable slope. The hard task is to prove the converse, that every badly approximable slope leads to superdensity.

The proof of Theorem 2 here, however, is completely different from our earlier technique. However, they share a common characteristic, in that neither is based on the traditional application of ergodic theory in the earlier study of density and uniformity in such dynamical systems. Indeed, the traditional approach is based on application of results such as Birkhoff's ergodic theorem which are essentially time-qualitative in nature. Lacking an error term, they do not appear to lead naturally to time-quantitative statements. Instead, we appeal to a non-ergodic approach, and base our arguments on number theory and combinatorics.

We thus have two methods to prove superdensity. They have different advantages, and they are not comparable. For instance, the shortline method developed in [3] works beyond billiards in polysquare regions, and can give superdensity for billiard in any regular polygon. We do not see how to we can establish such a result with the method of this paper.

Our proof here of Theorem 2 is not simple. We shall therefore illustrate the ideas by first studying the special case of the L-surface, which is arguably the simplest non-integrable polysquare surface.

The picture on the left in Figure 1.4 shows the L-shape region composed of 3 unit squares. Furthermore, it shows the L-graph, which is an undirected planar graph with 8 vertices, 5 horizontal edges and 5 vertical edges. Using this picture and identifying the edges, we can reduce the number of vertices. Identifying the edges h_1 , we see that $A = G$ and $B = F$. Identifying the edges h_2 , we see that $B = E$ and $C = D$. Identifying the edges v_1 , we see that $A = C$ and $H = D$. Identifying the edges v_2 , we see that $H = E$ and $G = F$. Thus $A = G = F = B = E = H = D = C$, so that all the vertices are identified with each other, and we have essentially only 1 vertex. This single vertex is a split singularity point of the geodesic flow, explaining why it is a *non-integrable* system; see the two geodesics in the picture on the right in Figure 1.4.




Figure 1.4: the L-surface and two geodesics

Furthermore, the surface has 3 faces and, after identification, 6 edges, so the Euler characteristic is $\chi = 1 - 6 + 3 = -2$. The genus g is obtained from the well known formula $g = 1 - (\chi/2) = 1 + 1 = 2$, and so the surface is homeomorphic to a 2-holed torus using the classification theorem of closed surfaces. We call this the *L-surface*. It is equipped with a flat metric, and the curvature is zero on every square face. The two geodesics in the picture on the right in Figure 1.4 illustrate why the vertex E , and hence every other vertex, is a split singularity of the geodesic flow. The L-surface is classified as a Riemann surface with a singular point.

As an analog of geodesics on the square torus, we can consider geodesics on the L-surface. It is customary to look at a geodesic from a dynamic viewpoint. We

thus assume that a particle moves on the geodesic with unit speed, so that time equals distance. If a geodesic on the L-surface has irrational slope and never hits the singular point, then in the L-shape it is represented as a union of infinitely many *parallel* line segments. A geodesic is uniquely determined by one of its points and its constant velocity vector, just like a geodesic on the square torus.

We shall first prove the following special case of Theorem 2.

Theorem 3. *If a half-infinite geodesic that does not hit a vertex of the L-surface has a slope that is a badly approximable number, then it is superdense on the L-surface.*

We can summarize the proof of Theorem 3, and hence also Theorem 2, in a nutshell as a *careful* use of the classical tool of continued fractions.

Before we begin our proof of Theorem 3, however, we make some comments.

First of all, the situation is completely different if we consider uniform distribution instead of density. Corresponding to superdensity, we can define superuniformity as the best form of time-quantitative uniformity, when the relevant discrepancy is of logarithmic size compared to the length of the geodesic. For half-infinite geodesics on the unit torus $[0, 1)^2$, the result of Khinchin says that a geodesic on the unit torus is superuniform if and only if its slope is a badly approximable number. This, however, does not remain the case if we consider half-infinite geodesics on the L-surface. While any half-infinite geodesic with a badly approximable slope is uniformly distributed on the L-surface, the rate of convergence to uniformity can be vastly different for two distinct badly approximable slopes. For example, it is shown in [1, 2] that a half-infinite geodesic of slope $\alpha = \sqrt{2}$ on the L-surface is superuniform, whereas a half-infinite geodesic of slope $\alpha = 1 + \sqrt{2}$ on the L-surface exhibits discrepancy greater than random square-root size.

Next, it is not difficult to visualize from Figure 1.5 that geodesic flow on the boundary surface of the unit cube leads to 4-direction geodesic flow on a polysquare surface with a cross-shaped net, obtained by *cutting* and *flattening* the boundary surface of the cube.




Figure 1.5: geodesic flow on the boundary surface of the cube

The 4-direction geodesic flow on this surface with cross-shaped net can then be converted to 1-direction geodesic flow on a different type of 4-copy construction, as shown in Figure 1.6. To get a 1-direction linear flow, we take 4 *rotated* copies of the usual cross-shaped net of the boundary surface of the cube, and glue them together by translation of the corresponding edges in different copies. We thus obtain a flat *translation surface*.

Note the difference between the two 4-copy constructions in Figures 1.3 and 1.6. In the former we reflect on a side, which changes the orientation. In the latter we rotate by a right angle, which preserves the orientation.




Figure 1.6: translation surface as a 4-fold covering of the boundary surface of the cube

The unit cube is the simplest example of a finite polycube region.

By a finite polycube region, we mean an arbitrary connected, but not necessarily simply-connected, polyhedron region P in 3-space which is tiled with unit cubes, assumed to be closed, that we call the *atomic cubes* of P , and which satisfies the following conditions:

- (i) Any two atomic cubes in P either are disjoint, or intersect at a single point, or have a single common edge, or have a common face.
- (ii) Any two atomic cubes in P are joined by a chain of atomic cubes where any two neighbors in the chain have a common face.

Assume further that P is simply connected. We now consider geodesic flow on the boundary surface of P . By gluing together the 4 rotated copies of the surface that are obtained by cutting and flattening the boundary surface of P , we convert the problem of geodesic flow on the boundary surface of P to a problem of 1-direction geodesic flow on a suitable finite polysquare surface. We thus have the following analog of Theorem 1.

Theorem 4. *Let \mathcal{P} be the boundary surface of an arbitrary finite simply connected polycube region. Then any half-infinite geodesic that does not hit a vertex of \mathcal{P} is superdense on \mathcal{P} if and only if the slope of the geodesic is a badly approximable number.*

Finally, we have said earlier that rectangle billiard is basically well understood. How about billiard in a right triangle which is *half* of a rectangle? It turns out that in the general case where the angles of the triangle can be irrational multiples of π , virtually nothing is known.

Consider billiard in an arbitrary right triangle T . We can ask many questions. Does there exist a half-infinite billiard orbit that is dense in T ? If so, can one find an explicit half-infinite billiard orbit that is dense in T ? Does there exist a slope such that every half-infinite billiard orbit with this initial slope that does not hit a vertex of T is dense in T ? If so, is it true that every half-infinite billiard orbit with almost any initial slope that does not hit a vertex of T is dense in T ?

Our inability to find answers to these questions illustrates the lack of methods that enable us to handle geodesic flow on *infinite* flat surfaces. Indeed, in the case where the acute angle of the right triangle T is an irrational multiple of π , iterated unfolding reduces billiard orbits in T to geodesic flow on an *infinite* flat surface. Unfortunately we know much, much less about infinite flat surfaces than about finite flat surfaces.

2. SOME PREREQUISITES

Without loss of generality, we assume that the slope of the geodesic is greater than 1. Suppose that the geodesic has slope $1/\alpha$, where $0 < \alpha < 1$ is irrational.

Our first tool is an interval exchange transformation $T = T_\alpha$ which encodes the information concerning the particular order with which a geodesic of slope $1/\alpha$ keeps hitting the horizontal edges h_1, h_2, h_3 .

The first step involves identifying the horizontal edges h_1, h_2, h_3 with unit intervals by making use of the correspondences

$$h_1 = [0, 1), \quad h_2 = [1, 2), \quad h_3 = [2, 3),$$

perhaps somewhat abusing notation, as shown in Figure 2.1.




Figure 2.1: representing horizontal edges of the L-surface by intervals

We now consider the piecewise linear map $T = T_\alpha$ defined according to Figure 2.2.




Figure 2.2: the interval exchange transformation $T = T_\alpha$

This is called the interval exchange transformation. More precisely, we have

$$T([0, 1 - \alpha)) = [2 + \alpha, 3), \quad T([1 - \alpha, 1)) = [1, 1 + \alpha), \quad (2.1)$$

$$T([1, 2 - \alpha)) = [1 + \alpha, 2), \quad T([2 - \alpha, 2)) = [2, 2 + \alpha), \quad (2.2)$$

$$T([2, 3 - \alpha)) = [\alpha, 1), \quad T([3 - \alpha, 3)) = [0, \alpha), \quad (2.3)$$

where, for instance, $T([0, 1 - \alpha)) = [2 + \alpha, 3)$ describes the $1/\alpha$ flow mapping the part $[0, 1 - \alpha)$ in $h_1 = [0, 1)$ to the part $[2 + \alpha, 3)$ in $h_3 = [2, 3)$ linearly in the form

$$Tx = 2 + \alpha + x, \quad x \in [0, 1 - \alpha),$$

and similarly for the rest in (2.1)–(2.3).

The novelty is that T acts on the longer interval $[0, 3)$ instead of the unit interval $[0, 1)$, but if we consider T modulo 1, then it acts simply as an α -shift, or irrational rotation, in the unit interval.

We next consider our main idea, which involves continued fractions. Consider an irrational number

$$\alpha = [a_0; a_1, a_2, a_3, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}, \quad (2.4)$$

where $a_0 \geq 0$ and $a_i \geq 1$, $i = 1, 2, 3, \dots$, are integers. The rational numbers

$$\frac{p_k}{q_k} = \frac{p_k(\alpha)}{q_k(\alpha)} = [a_0; a_1, \dots, a_k], \quad k = 0, 1, 2, 3, \dots, \quad (2.5)$$

are the k -convergents of α . It is well known that they give rise to the best rational approximations of the irrational number α , and we have

$$\frac{p_0}{q_0} < \frac{p_2}{q_2} < \frac{p_4}{q_4} < \dots < \alpha < \dots < \frac{p_5}{q_5} < \frac{p_3}{q_3} < \frac{p_1}{q_1}. \quad (2.6)$$

Let $\|y\|$ denote the distance of a real number y from the nearest integer. We shall make use of the fact that for an irrational number α the sequence

$$\min_{1 \leq k \leq n} \|k\alpha\|, \quad n = 1, 2, 3, \dots,$$

is well described by the continued fraction expansion of α .

For every $k = 0, 1, 2, 3, \dots$, we have

$$\|q\alpha\| \geq \|q_k\alpha\|, \quad 1 \leq q < q_{k+1}, \quad (2.7)$$

$$\|q_{k+1}\alpha\| < \|q_k\alpha\|,$$

as well as

$$\frac{1}{q_{k+1} + q_k} \leq \|q_k\alpha\| \leq \frac{1}{q_{k+1}}. \quad (2.8)$$

Indeed, the sequences p_k and q_k , $k = 0, 1, 2, 3, \dots$, are given by the initial values

$$p_0 = a_0, \quad p_1 = a_1 a_0 + 1, \quad q_0 = 1, \quad q_1 = a_1,$$

and the recurrence relations

$$p_{k+1} = a_{k+1} p_k + p_{k-1}, \quad q_{k+1} = a_{k+1} q_k + q_{k-1}, \quad k \geq 1. \quad (2.9)$$

We also have

$$p_{k-1} q_k - q_{k-1} p_k = (-1)^k, \quad k \geq 1.$$

On the other hand, using (2.6) and (2.9), it is easy to show that

$$\|q_{k+1}\alpha\| + a_{k+1} \|q_k\alpha\| = \|q_{k-1}\alpha\|. \quad (2.10)$$

We need the following result.

Theorem (3-distance theorem). *Consider the $n + 1$ numbers $0, \alpha, 2\alpha, 3\alpha, \dots, n\alpha$ modulo 1 in the unit torus/circle $[0, 1)$, leading to an $(n + 1)$ -partition. This partition exhibits at most 3 different distances between consecutive points. Furthermore, every positive integer n can be expressed uniquely in the form*

$$n = \mu q_k + q_{k-1} + r, \quad \text{with } 1 \leq \mu \leq a_{k+1} \text{ and } 0 \leq r < q_k,$$

in terms of the continued fraction (2.4) of α and its convergents (2.5), with the convention that $q_{-1} = 0$. Then

- (i) the distance $\|q_k \alpha\|$ shows up precisely $n + 1 - q_k$ times;
- (ii) the distance $\|q_{k-1} \alpha\| - \mu \|q_k \alpha\|$ shows up precisely $r + 1$ times; and
- (iii) the distance $\|q_{k-1} \alpha\| - (\mu - 1) \|q_k \alpha\|$ shows up precisely $q_k - r - 1$ times.

This surprising geometric fact, formulated as a conjecture by Steinhaus, has many proofs, by Sós [16, 17], Swierczkowski [19], Surányi [18], Halton [5] and Slater [15], with others published more recently.

3. PROOF OF THEOREM 3

Given an integer $k \geq 1$, let $\mathcal{A}_k(\alpha)$ denote the partition of the unit torus/circle $[0, 1)$ with $q_{k+1} = q_{k+1}(\alpha)$ division points $\{q\alpha\}$, $-1 \leq q \leq q_{k+1} - 2$, where $\{x\}$ denotes the fractional part of a real number x . Note that the choices $q = -1, 0$ in $\{q\alpha\}$ represent the *dangerous* endpoints of the special intervals $[0, 1 - \alpha)$ and $[1 - \alpha, 1)$ in (2.1). These are the two *singularities* of the interval exchange transformation T restricted to the interval $0 \leq x < 1$, in the sense that both 0 and $T(1 - \alpha) = 1$ represent the split singularity of the L-surface.

A consequence of the special choice $n = q_{k+1} - 1$ is that the 3-distance theorem simplifies to a 2-distance theorem. This in turn leads to some very useful information concerning the distances between the consecutive points of the q_{k+1} -partition $\mathcal{A}_k(\alpha)$ of the unit torus/circle $[0, 1)$. Indeed, using the second recurrence relation in (2.9), we have

$$n = q_{k+1} - 1 = a_{k+1} q_k + q_{k-1} - 1 = \mu q_k + q_{k-1} + r,$$

with $\mu = a_{k+1} - 1$ and $r = q_k - 1$. Since $q_k - r - 1 = 0$, it follows from the 3-distance theorem that there are only two distances

$$\|q_k \alpha\| \quad \text{and} \quad \|q_{k-1} \alpha\| - (a_{k+1} - 1) \|q_k \alpha\| = \|q_{k+1} \alpha\| + \|q_k \alpha\|, \quad (3.1)$$

in view of (2.10).

It follows immediately from (2.7) that one of the neighbors of 0 in the partition $\mathcal{A}_k(\alpha)$ is $\{q_k \alpha\}$ which clearly has distance $\|q_k \alpha\|$ from 0 in the unit torus/circle. Since α is irrational, the other neighbor of 0 in the partition $\mathcal{A}_k(\alpha)$ must have distance $\|q_{k+1} \alpha\| + \|q_k \alpha\|$ from 0 in the unit torus/circle. Simple calculation then shows that it is $\{((a_{k+1} - 1)q_k + q_{k-1})\alpha\}$. Thus the two neighbors

$$\{q_k \alpha\} \quad \text{and} \quad \{((a_{k+1} - 1)q_k + q_{k-1})\alpha\}$$

of 0 in the partition $\mathcal{A}_k(\alpha)$ exhibit the two gaps in (3.1) in some order. Similarly, the two neighbors

$$\{(q_k - 1)\alpha\} \quad \text{and} \quad \{((a_{k+1} - 1)q_k + q_{k-1} - 1)\alpha\}$$

of $1 - \alpha = \{-\alpha\}$ in the partition $\mathcal{A}_k(\alpha)$ exhibit the same two gaps in (3.1) in the same order.

The union of the left and right neighborhoods of 0 in the partition $\mathcal{A}_k(\alpha)$ has the form

$$B(0) = (-d^*, d^{**}), \quad (3.2)$$

and the union of the left and right neighborhoods of $1 - \alpha = \{-\alpha\}$ in the partition $\mathcal{A}_k(\alpha)$ has a similar form

$$B(-1) = (1 - \alpha - d^*, 1 - \alpha + d^{**}), \quad (3.3)$$

due to the same order, where

$$\{d^*, d^{**}\} = \{\|q_k\alpha\|, \|q_{k+1}\alpha\| + \|q_k\alpha\|\}, \quad (3.4)$$

but we have not specified which one is which. We refer to $B(0)$ and $B(-1)$ as the buffer zones of the singularities 0 and $1 - \alpha$ respectively in the partition $\mathcal{A}_k(\alpha)$.

We consider the special intervals

$$J_k(q) = J(\alpha; k; q) = [\{q\alpha\} - d^{**}, \{q\alpha\} + d^*], \quad 1 \leq q \leq q_{k+1} - 2. \quad (3.5)$$

Note that these short special intervals have three crucial properties:

- (i) They completely cover the two long intervals $(0, 1 - \alpha)$ and $(1 - \alpha, 1)$.
- (ii) They avoid the singularities 0, 1 and $1 - \alpha$, in view of (3.2)–(3.4).
- (iii) Any two intervals in $(0, 1 - \alpha)$ or in $(1 - \alpha, 1)$ arising from neighboring partition points exhibit *substantial overlapping*. More precisely, if $1 \leq q', q'' \leq q_{k+1} - 2$ are two integers such that $\{q'\alpha\}$ and $\{q''\alpha\}$ are neighboring points in the partition $\mathcal{A}_k(\alpha)$, so that both points are in the interval $(0, 1 - \alpha)$ or both points are in the interval $(1 - \alpha, 1)$, then

$$\text{length}(J_k(q') \cap J_k(q'')) \geq \min\{d^*, d^{**}\} = \|q_k\alpha\|. \quad (3.6)$$

Since the length of $J_k(q)$ is $2\|q_k\alpha\| + \|q_{k+1}\alpha\|$, the trivial upper bound

$$2\|q_k\alpha\| + \|q_{k+1}\alpha\| < 3\|q_k\alpha\|$$

and (3.6) together justify the term *substantial overlapping*.

Since T acts on the interval/circle $[0, 3)$, for every interval $J_k(q)$, $1 \leq q \leq q_{k+1} - 2$, given by (3.5), we define its *3-copy extension* $J_k(q; 3)$ by

$$J_k(q; 3) = J_k(q) \cup (1 + J_k(q)) \cup (2 + J_k(q)) \subset [0, 3), \quad (3.7)$$

a union of $J_k(q)$ with two of its translates.

After our preparation, we are now ready to study an orbit. Let $\mathcal{L}_\alpha(S; t)$, $t \geq 0$, be a parametrized half-infinite geodesic with initial point S and slope $1/\alpha$, under the usual arc-length parametrization.

Let M be *large*, and consider the initial segment $\mathcal{L}_\alpha(S; t)$, $0 \leq t \leq M$, of length M , which we denote by $(\mathcal{L}_\alpha; M)$. Suppose that

$$0 \leq t_1 < t_2 < t_3 < \dots < t_m \leq M, \quad (3.8)$$

where

$$t_{i+1} - t_i = \sqrt{1 + \alpha^2}, \quad 1 \leq i \leq m - 1, \quad (3.9)$$

is the sequence of time instances t when the initial segment $\mathcal{L}_\alpha(S; t)$, $0 \leq t \leq M$, intersects the union $h_1 \cup h_2 \cup h_3 = [0, 3)$ of the 3 horizontal edges of the L-surface in Figure 2.1. For notational simplicity let

$$y_i = \mathcal{L}_\alpha(S; t_i) \in [0, 3), \quad 1 \leq i \leq m, \quad (3.10)$$

denote these intersection points.

Using the interval exchange transformation $T = T_\alpha : [0, 3) \rightarrow [0, 3)$, we see that any two time-consecutive intersection points are governed by the simple relation

$$T(y_i) = y_{i+1}, \quad 1 \leq i \leq m - 1. \quad (3.11)$$

Lemma 3.1. *Suppose that $J_k(\ell_1)$ is a special interval of the form (3.5), and there exists $r(\ell_1) \in \{0, 1, 2\}$ such that*

$$\{y_i : 1 \leq i \leq m\} \cap (r(\ell_1) + J_k(\ell_1)) = \emptyset, \quad (3.12)$$

where $\{y_i : 1 \leq i \leq m\}$ is the set of intersection points defined in (3.10). Then for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$, we have

$$\{y_i : q_{k+1} \leq i \leq m - q_{k+1}\} \cap T^h(r(\ell_1) + J_k(\ell_1)) = \emptyset. \quad (3.13)$$

Proof. Since $q_{k+1} \geq 1$ and $m - q_{k+1} \leq m$, it follows trivially from (3.12) that (3.13) holds for $h = 0$.

Combining (3.11) and (3.12), we see that

$$\{y_i : 2 \leq i \leq m\} \cap T(r(\ell_1) + J_k(\ell_1)) = \emptyset.$$

Iterating this argument, we see that for every integer $1 \leq h \leq q_{k+1} - 2 - \ell_1$, we have

$$\{y_i : 1 + h \leq i \leq m\} \cap T^h(r(\ell_1) + J_k(\ell_1)) = \emptyset.$$

Since $q_{k+1} \geq q_{k+1} - 1 - \ell_1 \geq h + 1$ and $m - q_{k+1} \leq m$, it follows that (3.13) holds for every integer $1 \leq h \leq q_{k+1} - 2 - \ell_1$.

For every negative integer $1 - \ell_1 \leq h \leq -1$, using the inverse transformation T^{-1} , combining (3.11) and (3.12), and iterating, we have

$$\{y_i : 1 \leq i \leq m + h\} \cap T^h(r(\ell_1) + J_k(\ell_1)) = \emptyset.$$

Since $q_{k+1} \geq 1$ and $m + h \geq m + 1 - \ell_1 \geq m + 1 - q_{k+1} \geq m - q_{k+1}$, it follows that (3.13) holds for every integer $1 - \ell_1 \leq h \leq -1$.

The proof of the lemma is now complete. \square

Remark. We often refer to the deduction of (3.13) from (3.12) as a T -power extension argument.

For notational convenience, for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$, we write

$$T^h(r(\ell_1) + J_k(\ell_1)) = r(\ell_1 + h) + J_k(\ell_1 + h). \quad (3.14)$$

Note that (3.14) defines $r(q)$ for every integer $1 \leq q \leq q_{k+1} - 2$. Furthermore, combining (3.13) and (3.14), we have

$$\{y_i : q_{k+1} \leq i \leq m - q_{k+1}\} \cap (r(q) + J_k(q)) = \emptyset \quad (3.15)$$

for every integer $1 \leq q \leq q_{k+1} - 2$.

Suppose that $I_0 \subset [0, 3)$ is $(\mathcal{L}_\alpha; M)$ -free, so that

$$\{y_i : 1 \leq i \leq m\} \cap I_0 = \emptyset,$$

where $\{y_i : 1 \leq i \leq m\}$ is the set of intersection points defined in (3.10). Let k be an integer, and suppose that $J_k(\ell_1)$ is a special interval of the form (3.5), and there exists $r(\ell_1) \in \{0, 1, 2\}$ such that

$$r(\ell_1) + J_k(\ell_1) \subset I_0. \quad (3.16)$$

Then (3.12) holds.

Remark. We shall later choose an *optimal* value of k for which (3.16) holds.

We distinguish a few cases according to the special relations between various sets of intersection points and various special intervals. We take advantage of the substantial overlapping of the short special intervals $J_k(q)$ defined by (3.5).

Recall that if $1 \leq q', q'' \leq q_{k+1} - 2$ are two integers such that $\{q'\alpha\}$ and $\{q''\alpha\}$ are neighboring points in the partition $\mathcal{A}_k(\alpha)$, so that both points are in the interval

$(0, 1 - \alpha)$ or both points are in the interval $(1 - \alpha, 1)$, then combining (2.8) and (3.6), we have

$$\text{length}(J_k(q') \cap J_k(q'')) \geq \|q_k \alpha\| \geq \frac{1}{q_{k+1} + q_k} > \frac{1}{2q_{k+1}}.$$

Recall also from (2.8), (3.4) and (3.5) that

$$\text{length}(J_k(q)) = 2\|q_k \alpha\| + \|q_{k+1} \alpha\| < 3\|q_k \alpha\| \leq \frac{3}{q_{k+1}},$$

so that

$$\text{length}(J_{k+8}(q)) < \frac{3}{q_{k+9}}. \quad (3.17)$$

On the other hand, a trivial deduction from (2.9) gives

$$q_{k+2} = a_{k+2}q_{k+1} + q_k \geq q_{k+1} + q_k = a_{k+1}q_k + q_{k-1} + q_k \geq 2q_k,$$

so that iterating this a few times, we conclude that

$$q_{k+9} \geq 2q_{k+7} \geq 4q_{k+5} \geq 8q_{k+3} \geq 16q_{k+1}. \quad (3.18)$$

Combining (3.17) and (3.18), we conclude that the intersection $J_k(q') \cap J_k(q'')$ must contain a special interval of the type $J_{k+8}(q)$ for some $1 \leq q \leq q_{k+9} - 2$. We split the argument into two complementary cases.

Case 1A. The following *intersection property* holds. For every

$$J_{k+8}(\ell) \subset J_k(q) \quad \text{and} \quad r \in \{0, 1, 2\},$$

with $r \neq r(q)$ given by (3.14), we have

$$\{y_i : q_{k+1} \leq i \leq m - q_{k+1}\} \cap (r + J_{k+8}(\ell)) \neq \emptyset. \quad (3.19)$$

Lemma 3.2. *Case 1A is impossible.*

Case 1B. There exist

$$J_{k+8}(\ell_2) \subset J_k(q^+) \quad \text{and} \quad r_1 \in \{0, 1, 2\},$$

with $r_1 \neq r(q^+)$ given by (3.14), such that

$$\{y_i : q_{k+1} \leq i \leq m - q_{k+1}\} \cap (r_1 + J_{k+8}(\ell_2)) = \emptyset. \quad (3.20)$$

Since $J_{k+8}(\ell_2) \subset J_k(q^+)$, it clearly follows from (3.15) that

$$\{y_i : q_{k+1} \leq i \leq m - q_{k+1}\} \cap (r(q^+) + J_{k+8}(\ell_2)) = \emptyset. \quad (3.21)$$

Since (3.20) and (3.21) are analogs of (3.12), the T -power expansion argument in the proof of Lemma 3.1 shows that for every integer $1 - \ell_2 \leq h \leq q_{k+9} - 2 - \ell_2$, we have

$$\{y_i : q_{k+1} + q_{k+9} \leq i \leq m - q_{k+1} - q_{k+9}\} \cap T^h(r_2 + J_{k+8}(\ell_2)) = \emptyset, \quad (3.22)$$

$$\{y_i : q_{k+1} + q_{k+9} \leq i \leq m - q_{k+1} - q_{k+9}\} \cap T^h(r(q^+) + J_{k+8}(\ell_2)) = \emptyset. \quad (3.23)$$

For notational convenience, for every integer $1 - \ell_2 \leq h \leq q_{k+9} - 2 - \ell_2$, we write

$$T^h(r(q^+) + J_{k+8}(\ell_2)) = r^*(\ell_2 + h) + J_{k+8}(\ell_2 + h), \quad (3.24)$$

$$T^h(r_2 + J_{k+8}(\ell_2)) = r^{**}(\ell_2 + h) + J_{k+8}(\ell_2 + h). \quad (3.25)$$

Then combining (3.22)–(3.25), we have

$$\{y_i : q_{k+1} + q_{k+9} \leq i \leq m - q_{k+1} - q_{k+9}\} \cap (r^*(q) + J_{k+8}(q)) = \emptyset, \quad (3.26)$$

$$\{y_i : q_{k+1} + q_{k+9} \leq i \leq m - q_{k+1} - q_{k+9}\} \cap (r^{**}(q) + J_{k+8}(q)) = \emptyset, \quad (3.27)$$

for every integer $1 \leq q \leq q_{k+9} - 2$. Clearly

$$r^*(q) \neq r^{**}(q), \quad 1 \leq q \leq q_{k+9} - 2.$$

We now split Case 1B into two complementary cases.

Case 2A. The following *intersection property* holds. For every

$$J_{k+16}(\ell) \subset J_{k+8}(q) \quad \text{and} \quad r \in \{0, 1, 2\},$$

with $r \neq r^*(q), r^{**}(q)$ given by (3.24) and (3.25), we have

$$\{y_i : q_{k+1} + q_{k+9} \leq i \leq m - q_{k+1} - q_{k+9}\} \cap (r + J_{k+16}(\ell)) \neq \emptyset.$$

Lemma 3.3. *Case 2A is impossible.*

Case 2B. There exist

$$J_{k+16}(\ell_3) \subset J_{k+8}(q^{++}) \quad \text{and} \quad r_2 \in \{0, 1, 2\},$$

with $r_2 \neq r^*(q^{++}), r^{**}(q^{++})$ given by (3.24) and (3.25), such that

$$\{y_i : q_{k+1} + q_{k+9} \leq i \leq m - q_{k+1} - q_{k+9}\} \cap (r_2 + J_{k+16}(\ell_3)) = \emptyset. \quad (3.28)$$

Lemma 3.4. *If Case 2B holds, then*

$$m \leq 2q_{k+1} + 2q_{k+9} + 2q_{k+17} + 6. \quad (3.29)$$

Proof of Theorem 3. Suppose that $I_0 \subset [0, 3)$ is $(\mathcal{L}_\alpha; M)$ -free, so that

$$\{y_i : 1 \leq i \leq m\} \cap I_0 = \emptyset.$$

Let $J_k(\ell_1) = J(\alpha; k; \ell_1)$ be the *longest* special interval of the form (3.5) such that

$$r(\ell_1) + J_k(\ell_1) \subset I_0 \quad \text{for some } \ell_1 \text{ and } r(\ell_1) \in \{0, 1, 2\}.$$

Then

$$\text{length}(I_0) < 4(\|q_{k-1}\alpha\| + \|q_k\alpha\|), \quad (3.30)$$

since otherwise there exists a longer special interval

$$r(\ell) + J_{k-1}(\ell) \subset I_0 \quad \text{for some } \ell \text{ and } r(\ell) \in \{0, 1, 2\},$$

a contradiction. Combining (3.30) with (2.8), we have

$$\text{length}(I_0) < \frac{4}{q_k} + \frac{4}{q_{k+1}} < \frac{8}{q_k}. \quad (3.31)$$

On the other hand, it follows from (3.8) and (3.9) that

$$M \leq (m+1)\sqrt{1+\alpha^2}. \quad (3.32)$$

Also, in view of Lemmas 3.2–3.4, it is clear that the bound (3.29) holds. Finally, recall that $0 < \alpha < 1$ and

$$\alpha = [a_1, a_2, a_3, \dots] = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}$$

is badly approximable, so there exists a constant A such that the continued fraction digits $a_i \leq A$ for every $i = 1, 2, 3, \dots$. It follows from (2.9) that

$$q_{k+1} < q_{k+9} < q_{k+17} \leq (A+1)^{17}q_k. \quad (3.33)$$

Combining (3.29), (3.32) and (3.33), we see that

$$M \leq (2q_{k+1} + 2q_{k+9} + 2q_{k+17} + 7)\sqrt{1+\alpha^2} < 7(A+1)^{17}q_k\sqrt{2}. \quad (3.34)$$

It now follows from (3.31) and (3.34) that a geodesic segment $\mathcal{L}_\alpha(S; t)$, $0 \leq t \leq M$, of length $M = 7(A+1)^{17}q_k\sqrt{2}$ must intersect every subinterval I of $h_1 \cup h_2 \cup h_3$ with

$\text{length}(I) = 8/q_k$. Since the product $M \text{length}(I) = 56(A+1)^{17}\sqrt{2}$ is a constant independent of k , this establishes superdensity of the half-infinite geodesic. \square

4. PROOF OF LEMMAS 3.2–3.4

Before we present the proof of our main lemmas, we begin by investigating a simple situation which serves to illustrate our method.

Simple Case. There exist integers $r^*(\ell_1)$ and $r^{**}(\ell_1)$ such that

- (1) $r(\ell_1), r^*(\ell_1), r^{**}(\ell_1)$ form a permutation of $0, 1, 2$;
- (2) $\{y_i : 1 \leq i \leq m\} \cap (r^*(\ell_1) + J_k(\ell_1)) = \emptyset$; and
- (3) $\{y_i : 1 \leq i \leq m\} \cap (r^{**}(\ell_1) + J_k(\ell_1)) = \emptyset$.

Lemma 4.1. *If the Simple Case holds, then $m \leq 2q_{k+1} + 6$.*

Proof. For notational simplicity, we write

$$Q(k; m) = \{y_i : q_{k+1} \leq i \leq m - q_{k+1}\}. \quad (4.1)$$

Since the properties (2) and (3) in the Simple Case are analogs of (3.12), we can repeat the T -power extension argument in Lemma 3.1 and conclude that for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$, we have

$$Q(k; m) \cap T^h(r^*(\ell_1) + J_k(\ell_1)) = \emptyset, \quad (4.2)$$

$$Q(k; m) \cap T^h(r^{**}(\ell_1) + J_k(\ell_1)) = \emptyset. \quad (4.3)$$

As in (3.14), we write

$$T^h(r^*(\ell_1) + J_k(\ell_1)) = r^*(\ell_1 + h) + J_k(\ell_1 + h), \quad (4.4)$$

$$T^h(r^{**}(\ell_1) + J_k(\ell_1)) = r^{**}(\ell_1 + h) + J_k(\ell_1 + h), \quad (4.5)$$

for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$. Note that (4.4) and (4.5) define $r^*(q)$ and $r^{**}(q)$ respectively for every integer $1 \leq q \leq q_{k+1} - 2$. Recall next that $r(q)$ is defined by (3.14). Indeed, using the notation (3.14), it is easy to check that the assertion (3.13) for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$ implies that

$$Q(k; m) \cap (r(\ell_1 + h) + J_k(\ell_1 + h)) = \emptyset \quad (4.6)$$

for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$.

Combining (4.2)–(4.6) for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$, we deduce that for every integer $1 \leq q \leq q_{k+1} - 2$, we have

$$Q(k; m) \cap (r^*(q) + J_k(q)) = \emptyset, \quad (4.7)$$

$$Q(k; m) \cap (r^{**}(q) + J_k(q)) = \emptyset, \quad (4.8)$$

$$Q(k; m) \cap (r(q) + J_k(q)) = \emptyset. \quad (4.9)$$

Also, in view of the property (1) in the Simple Case, it is clear that $r(q), r^*(q), r^{**}(q)$ form a permutation of $0, 1, 2$ for every integer $1 \leq q \leq q_{k+1} - 2$.

By definition, the 3-copy extensions $J_k(q; 3)$, $1 \leq q \leq q_{k+1} - 2$, give rise to 6 continuous chains of overlapping intervals in the torus/circle $[0, 3)$ such that the 6 chains completely cover the 6 intervals

$$(0, 1 - \alpha), \quad (1 - \alpha, 1), \quad (1, 2 - \alpha), \quad (2 - \alpha, 2), \quad (2, 3 - \alpha), \quad (3 - \alpha, 3),$$

and there are only 6 points in $[0, 3)$ that are not covered by the 6 chains, namely

$$0, \quad 1 - \alpha, \quad 1, \quad 2 - \alpha, \quad 2, \quad 3 - \alpha.$$

Combining (4.7)–(4.9) for every integer $1 \leq q \leq q_{k+1} - 2$, we deduce that the set $Q(k; m)$ is not covered by the 6 chains. Indeed, if $m \geq 2q_{k+1} + 7$, then the set (4.1)

has at least 7 distinct elements, which is more than 6, giving rise to a contradiction. We conclude therefore that, under the conditions of the Simple Case, we must have $m \leq 2q_{k+1} + 6$, and this completes the proof. \square

Proof of Lemma 3.2. Again, for notational simplicity, we use (4.1).

Our first step is to prove that, under the condition of Case 1A, any two neighboring 3-copy extensions $J_k(q'; 3)$ and $J_k(q''; 3)$ are *synchronized* in the following precise sense: For each $r \in \{0, 1, 2\}$, we have

$$Q(k; m) \cap (r + J_k(q')) = \emptyset \quad \text{if and only if} \quad Q(k; m) \cap (r + J_k(q'')) = \emptyset. \quad (4.10)$$

To establish this, we consider two cases.

Suppose first that $r = r(q')$. Using the notation (3.14), it follows from (3.13) that

$$Q(k; m) \cap (r(q') + J_k(q')) = \emptyset. \quad (4.11)$$

Assume on the contrary that

$$Q(k; m) \cap (r(q') + J_k(q'')) \neq \emptyset. \quad (4.12)$$

Then it follows from (3.15) that $r(q'') \neq r(q')$. On the other hand, we know that the intersection $J_k(q') \cap J_k(q'')$ must contain a special interval of the type $J_{k+8}(j_0)$ for some j_0 , so

$$r(q') + J_{k+8}(j_0) \subset (r(q') + J_k(q')) \cap (r(q') + J_k(q'')). \quad (4.13)$$

Since $r(q') \neq r(q'')$ and $J_{k+8}(j_0) \subset J_k(q'')$, the condition of Case 1A is satisfied with $q = q''$, and (3.19) becomes

$$Q(k; m) \cap (r(q') + J_{k+8}(j_0)) \neq \emptyset. \quad (4.14)$$

But (4.11) and (4.13) contradict (4.14), and so (4.12) fails. Thus our claim (4.10) holds in this case.

Suppose next that $r \neq r(q')$. As before, we use the fact that the intersection $J_k(q') \cap J_k(q'')$ must contain a special interval of the type $J_{k+8}(j_0)$ for some j_0 , so

$$r + J_{k+8}(j_0) \subset (r + J_k(q')) \cap (r + J_k(q'')). \quad (4.15)$$

Since $r \neq r(q')$, the condition of Case 1A is satisfied with $q = q'$, and (3.19) becomes

$$Q(k; m) \cap (r + J_{k+8}(j_0)) \neq \emptyset. \quad (4.16)$$

It now follows from (4.15) and (4.16) that

$$Q(k; m) \cap (r + J_k(q')) \neq \emptyset \quad \text{and} \quad Q(k; m) \cap (r + J_k(q'')) \neq \emptyset,$$

so that our claim (4.10) holds also in this case.

By definition, the 3-copy extensions $J_k(q; 3)$, $1 \leq q \leq q_{k+1} - 2$, give rise to 6 continuous chains of overlapping intervals in the torus/circle $[0, 3)$ such that the 6 chains completely cover the 6 intervals

$$(0, 1 - \alpha), \quad (1 - \alpha, 1), \quad (1, 2 - \alpha), \quad (2 - \alpha, 2), \quad (2, 3 - \alpha), \quad (3 - \alpha, 3). \quad (4.17)$$

The synchronization property we have just established now implies that each of the 6 long special intervals in (4.17) satisfies one of the following two properties. Either such a long special interval is *disjoint* from the set $Q(k; m)$, or the set $Q(k; m)$ is *dense* in such a long special interval, in the precise sense that every subinterval of length $1/q_{k+8}$ contains a point from the set $Q(k; m)$.

Moreover, it is not difficult to show that precisely 2 of the 6 long special intervals in (4.17) are disjoint from the set $Q(k; m)$. To see this, choose two integers q' and q'' satisfying $1 \leq q', q'' \leq q_{k+1} - 2$ such that

$$J_k(q') \subset (0, 1 - \alpha) \quad \text{and} \quad J_k(q'') \subset (1 - \alpha, 1). \quad (4.18)$$

Then it follows from (3.13) and (3.14) that

$$Q(k; m) \cap (r(q') + J_k(q')) = \emptyset \quad \text{and} \quad Q(k; m) \cap (r(q'') + J_k(q'')) = \emptyset. \quad (4.19)$$

Now write

$$\mathcal{I}_1 = r(q') + (0, 1 - \alpha) \quad \text{and} \quad \mathcal{I}_2 = r(q'') + (1 - \alpha, 1). \quad (4.20)$$

The synchronization property and (4.18)–(4.20) now imply that

$$Q(k; m) \cap \mathcal{I}_1 = \emptyset \quad \text{and} \quad Q(k; m) \cap \mathcal{I}_2 = \emptyset.$$

Note that the union $\mathcal{I}_1 \cup \mathcal{I}_2$ modulo 1 is precisely the unit interval $[0, 1)$.

Now \mathcal{I}_1 and \mathcal{I}_2 are 2 of the 6 long special intervals in (4.17). Let \mathcal{I}_j , $3 \leq j \leq 6$, denote the remaining long special intervals in (4.17). The condition of Case 1A now implies that these 4 intervals are not disjoint from $Q(k; m)$, so that $Q(k; m)$ is dense in each of them. Each T -image $T(\mathcal{I}_j)$, $j = 1, 2$, has at most 1 common point with the set $Q(k; m)$. This is a contradiction, since the union $T(\mathcal{I}_1) \cup T(\mathcal{I}_2)$ has a *substantial* intersection with the union $\mathcal{I}_3 \cup \mathcal{I}_4 \cup \mathcal{I}_5 \cup \mathcal{I}_6$, which implies that it must have a *substantial* intersection with the set $Q(k; m)$, much more than at most 2 elements. Thus Case 1A is impossible, and this completes the proof. \square

Proof of Lemma 3.3. For notational simplicity, we write

$$Q^*(k; m) = \{y_i : q_{k+1} + q_{k+9} \leq i \leq m - q_{k+1} - q_{k+9}\}. \quad (4.21)$$

We can proceed along similar lines as in the first part of the proof of Lemma 3.2 for Case 1A, and show that any two neighboring 3-copy extensions $J_{k+8}(q'; 3)$ and $J_{k+8}(q''; 3)$ are *synchronized* in the following precise sense: For each $r \in \{0, 1, 2\}$, we have

$$Q^*(k; m) \cap (r + J_{k+8}(q')) = \emptyset \quad \text{if and only if} \quad Q^*(k; m) \cap (r + J_{k+8}(q'')) = \emptyset.$$

By definition, the 3-copy extensions $J_{k+8}(q; 3)$, $1 \leq q \leq q_{k+9} - 2$, give rise to 6 continuous chains of overlapping intervals in the torus/circle $[0, 3)$ such that the 6 chains completely cover the 6 intervals (4.17). The synchronization property now implies that each of the 6 long special intervals in (4.17) satisfies one of the following two properties. Either such a long special interval is *disjoint* from the set $Q^*(k; m)$, or the set $Q^*(k; m)$ is *dense* in such a long special interval, in the precise sense that every subinterval of length $1/q_{k+16}$ contains a point from the set $Q^*(k; m)$.

Moreover, it is not difficult to show that precisely 4 of the 6 long special intervals in (4.17) are disjoint from the set $Q^*(k; m)$. To see this, choose two integers q' and q'' satisfying $1 \leq q', q'' \leq q_{k+9} - 2$ such that

$$J_{k+8}(q') \subset (0, 1 - \alpha) \quad \text{and} \quad J_{k+8}(q'') \subset (1 - \alpha, 1). \quad (4.22)$$

Then it follows from (3.26) and (3.27) that

$$Q^*(k; m) \cap (r^*(q') + J_{k+8}(q')) = \emptyset, \quad Q^*(k; m) \cap (r^*(q'') + J_{k+8}(q'')) = \emptyset, \quad (4.23)$$

$$Q^*(k; m) \cap (r^{**}(q') + J_{k+8}(q')) = \emptyset, \quad Q^*(k; m) \cap (r^{**}(q'') + J_{k+8}(q'')) = \emptyset. \quad (4.24)$$

Now write

$$\mathcal{I}_1 = r^*(q') + (0, 1 - \alpha), \quad \mathcal{I}_2 = r^*(q'') + (1 - \alpha, 1), \quad (4.25)$$

$$\mathcal{I}_3 = r^{**}(q') + (0, 1 - \alpha), \quad \mathcal{I}_4 = r^{**}(q'') + (1 - \alpha, 1). \quad (4.26)$$

The synchronization property and (4.22)–(4.26) now imply that

$$\begin{aligned} Q^*(k; m) \cap \mathcal{I}_1 &= \emptyset, & Q^*(k; m) \cap \mathcal{I}_2 &= \emptyset, \\ Q^*(k; m) \cap \mathcal{I}_3 &= \emptyset, & Q^*(k; m) \cap \mathcal{I}_4 &= \emptyset. \end{aligned}$$

Note that the union $\mathcal{I}_1 \cup \mathcal{I}_2 \cup \mathcal{I}_3 \cup \mathcal{I}_4$ modulo 1 is precisely the unit interval $[0, 1]$ twice.

Now $\mathcal{I}_1, \mathcal{I}_2, \mathcal{I}_3, \mathcal{I}_4$ are 4 of the 6 long special intervals in (4.17). Let \mathcal{I}_j , $j = 5, 6$, denote the remaining long special intervals in (4.17). The condition of Case 2A now implies that these 2 intervals are not disjoint from $Q^*(k; m)$, so that $Q^*(k; m)$ is dense in each of them. Each T -image $T(\mathcal{I}_j)$, $j = 1, \dots, 4$, has at most 1 common point with the set $Q^*(k; m)$. This is a contradiction, since the union

$$T(\mathcal{I}_1) \cup T(\mathcal{I}_2) \cup T(\mathcal{I}_3) \cup T(\mathcal{I}_4)$$

has a *substantial* intersection with the union $\mathcal{I}_5 \cup \mathcal{I}_6$, which implies that it must have a *substantial* intersection with the set $Q^*(k; m)$, much more than at most 4 elements. Thus Case 2A is impossible, and this completes the proof. \square

Proof of Lemma 3.4. For notational simplicity, we define $Q^*(k; m)$ by (4.21), and write

$$Q^{**}(k; m) = \{y_i : q_{k+1} + q_{k+9} + q_{k+17} \leq i \leq m - q_{k+1} - q_{k+9} - q_{k+17}\}.$$

Since (3.28) is an analog of (3.12), we can repeat the T -power extension argument in Lemma 3.1 and conclude that for every integer $1 - \ell_3 \leq h \leq q_{k+17} - 2 - \ell_3$, we have

$$Q^{**}(k; m) \cap T^h(r_2 + J_{k+16}(\ell_3)) = \emptyset. \quad (4.27)$$

Since $J_{k+16}(\ell_3) \subset J_{k+8}(q^{++})$, it follows from (3.26) and (3.27) that

$$Q^*(k; m) \cap (r^*(q^{++}) + J_{k+16}(\ell_3)) = \emptyset, \quad (4.28)$$

$$Q^*(k; m) \cap (r^{**}(q^{++}) + J_{k+16}(\ell_3)) = \emptyset. \quad (4.29)$$

Next, note that (4.28) and (4.29) are also analogs of (3.12), so again we can repeat the T -power extension argument in Lemma 3.1 and conclude that for every integer $1 - \ell_3 \leq h \leq q_{k+17} - 2 - \ell_3$, we have

$$Q^{**}(k; m) \cap T^h(r^*(q^{++}) + J_{k+16}(\ell_3)) = \emptyset, \quad (4.30)$$

$$Q^{**}(k; m) \cap T^h(r^{**}(q^{++}) + J_{k+16}(\ell_3)) = \emptyset. \quad (4.31)$$

Now, for every integer $1 - \ell_3 \leq h \leq q_{k+17} - 2 - \ell_3$, we write

$$T^h(r^*(q^{++}) + J_{k+16}(\ell_3)) = r^{(0)}(\ell_3 + h) + J_{k+16}(\ell_3 + h), \quad (4.32)$$

$$T^h(r^{**}(q^{++}) + J_{k+16}(\ell_3)) = r^{(1)}(\ell_3 + h) + J_{k+16}(\ell_3 + h), \quad (4.33)$$

$$T^h(r_2 + J_{k+16}(\ell_3)) = r^{(2)}(\ell_3 + h) + J_{k+16}(\ell_3 + h). \quad (4.34)$$

Clearly it follows from the assumption of Case 2B that $r^{(0)}(q), r^{(1)}(q), r^{(2)}(q)$ form a permutation of $0, 1, 2$ for every integer $1 \leq q \leq q_{k+17} - 2$. Combining (4.27) and

(4.30)–(4.34), we have

$$Q^{**}(k; m) \cap (r^{(0)}(q) + J_{k+16}(q)) = \emptyset, \quad (4.35)$$

$$Q^{**}(k; m) \cap (r^{(1)}(q) + J_{k+16}(q)) = \emptyset, \quad (4.36)$$

$$Q^{**}(k; m) \cap (r^{(2)}(q) + J_{k+16}(q)) = \emptyset, \quad (4.37)$$

for every integer $1 \leq q \leq q_{k+17} - 2$.

Note now that (4.35)–(4.37) are similar to (4.7)–(4.9) in the proof of Lemma 4.1, so we now mimic the last part of that proof.

By definition, the 3-copy extensions $J_{k+16}(q; 3)$, $1 \leq q \leq q_{k+17} - 2$, give rise to 6 continuous chains of overlapping intervals in the torus/circle $[0, 3)$ such that the 6 chains completely cover the 6 intervals

$$(0, 1 - \alpha), \quad (1 - \alpha, 1), \quad (1, 2 - \alpha), \quad (2 - \alpha, 2), \quad (2, 3 - \alpha), \quad (3 - \alpha, 3),$$

and there are only 6 points in $[0, 3)$ that are not covered by the 6 chains, namely

$$0, \quad 1 - \alpha, \quad 1, \quad 2 - \alpha, \quad 2, \quad 3 - \alpha.$$

Combining (4.35)–(4.37) for every integer $1 \leq q \leq q_{k+17} - 2$, we deduce that the set $Q^{**}(k; m)$ is not covered by the 6 chains. Indeed, if $m \geq 2q_{k+1} + 2q_{k+9} + 2q_{k+17} + 7$, then the set $Q^{**}(k; m)$ has at least 7 distinct elements, which is more than 6, giving rise to a contradiction. We conclude therefore that, under the conditions of Case 2B, we must have $m \leq 2q_{k+1} + 2q_{k+9} + 2q_{k+17} + 6$, and this completes the proof. \square

5. PROOF OF THEOREM 2

Consider now an arbitrary finite polysquare surface \mathcal{P} . We are now in a position to complete the proof of Theorem 2, and show that if the slope of a half-infinite geodesic on \mathcal{P} is a badly approximable number, then the geodesic is superdense on \mathcal{P} .

The proof is a fairly straightforward adaptation of the proof of Theorem 3, apart from the observation that the number of cases we need to consider is a function of the number of square faces of \mathcal{P} , and so can be arbitrarily large.

Without loss of generality, we assume as before that the slope of the half-infinite geodesic is greater than 1. Suppose that it has slope $1/\alpha$, where $0 < \alpha < 1$ is irrational.

Our first step is to generalize the interval exchange transformation $T = T_\alpha$ defined in Section 2. Suppose that the polysquare surface \mathcal{P} has s square faces. Each square face has a top horizontal edge and a bottom horizontal edge. Each top horizontal edge is identified with a unique bottom horizontal edge, and these give rise to s horizontal edges h_1, h_2, \dots, h_s . We now identify these horizontal edges with unit intervals by making use of the correspondences

$$h_1 = [0, 1), \quad h_2 = [1, 2), \quad \dots, \quad h_s = [s - 1, s).$$

We now consider the piecewise linear map $T = T_\alpha$ defined according to the analog of Figure 2.2 that corresponds to \mathcal{P} . More precisely, for each integer $1 \leq j \leq s$, there exist unique integers $0 \leq j', j'' \leq s - 1$ such that

$$T([j - 1, j - \alpha)) = [j' + \alpha, j' + 1) \quad \text{and} \quad T([j - \alpha, j)) = [j'', j'' + \alpha).$$

Indeed, it is not difficult to see that the map $T : [0, s) \rightarrow [0, s)$ is one-to-one and onto. While T acts on the longer interval $[0, s)$ instead of the unit interval $[0, 1)$, if we consider T modulo 1, then it acts simply as an α -shift, or irrational rotation, in the unit interval.

As before, our main idea involves continued fractions, in particular, the special case of the 3-distance theorem which becomes a 2-distance theorem. Indeed, we can repeat our discussion at the beginning of Section 3 *verbatim* up to the end of the paragraph preceding (3.7). In particular, for any integer $1 \leq q \leq q_{k+1} - 2$, we can define the special interval $J_k(q)$ as in (3.5). Then analogous to (3.7), for any integer $1 \leq q \leq q_{k+1} - 2$, we define its s -copy extension $J_k(q; s)$ by

$$J_k(q; s) = J_k(q) \cup (1 + J_k(q)) \cup \dots \cup ((s-1) + J_k(q)) \subset [0, s),$$

a union of $J_k(q)$ with $s-1$ of its translates.

Let $\mathcal{L}_\alpha(t) = \mathcal{L}_\alpha(\mathcal{P}; S; t)$, $t \geq 0$, be a parametrized half-infinite geodesic on \mathcal{P} with initial point S and slope $1/\alpha$, under the usual arc-length parametrization.

Let M be *large*, and consider the initial segment $\mathcal{L}_\alpha(t)$, $0 \leq t \leq M$, of length M , which we denote by $(\mathcal{L}_\alpha; M)$. Suppose that

$$0 \leq t_1 < t_2 < t_3 < \dots < t_m \leq M, \quad (5.1)$$

where

$$t_{i+1} - t_i = \sqrt{1 + \alpha^2}, \quad 1 \leq i \leq m-1, \quad (5.2)$$

is the sequence of time instances t when the initial segment $\mathcal{L}_\alpha(t)$, $0 \leq t \leq M$, intersects the union $h_1 \cup \dots \cup h_s = [0, s)$ of the s horizontal edges of the polysquare surface \mathcal{P} . For notational simplicity let

$$y_i = \mathcal{L}_\alpha(t_i) \in [0, s), \quad 1 \leq i \leq m, \quad (5.3)$$

denote these intersection points.

Using the interval exchange transformation $T = T_\alpha : [0, s) \rightarrow [0, s)$, we see that any two time-consecutive intersection points are governed by the simple relation

$$T(y_i) = y_{i+1}, \quad 1 \leq i \leq m-1.$$

We have the following analog of Lemma 3.1 which is easily established by the T -power extension argument.

Lemma 5.1. *Suppose that $J_k(\ell_1)$ is a special interval of the form (3.5), and there exists $r(\ell_1) \in \{0, 1, \dots, s-1\}$ such that*

$$\{y_i : 1 \leq i \leq m\} \cap (r(\ell_1) + J_k(\ell_1)) = \emptyset, \quad (5.4)$$

where $\{y_i : 1 \leq i \leq m\}$ is the set of intersection points defined in (5.3). Then for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$, we have

$$\{y_i : q_{k+1} \leq i \leq m - q_{k+1}\} \cap T^h(r(\ell_1) + J_k(\ell_1)) = \emptyset. \quad (5.5)$$

For notational convenience, for every integer $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$, we write

$$T^h(r(\ell_1) + J_k(\ell_1)) = r_1^{(0)}(\ell_1 + h) + J_k(\ell_1 + h). \quad (5.6)$$

Note that (5.6) defines $r(q)$ for every integer $1 \leq q \leq q_{k+1} - 2$. Furthermore, combining (5.5) and (5.6), we have

$$\{y_i : q_{k+1} \leq i \leq m - q_{k+1}\} \cap (r_1^{(0)}(q) + J_k(q)) = \emptyset \quad (5.7)$$

for every integer $1 \leq q \leq q_{k+1} - 2$.

Suppose that $I_0 \subset [0, s)$ is $(\mathcal{L}_\alpha; M)$ -free, so that

$$\{y_i : 1 \leq i \leq m\} \cap I_0 = \emptyset,$$

where $\{y_i : 1 \leq i \leq m\}$ is the set of intersection points defined in (5.3). Let k be an integer, and suppose that $J_k(\ell_1)$ is a special interval of the form (3.5), and there exists $r(\ell_1) \in \{0, 1, \dots, s-1\}$ such that

$$r(\ell_1) + J_k(\ell_1) \subset I_0. \quad (5.8)$$

Then (5.4) holds. As before, we shall later choose an *optimal* value of k for which (5.8) holds.

Again, we distinguish a few cases according to the special relations between various sets of intersection points and various special intervals. We take advantage of the substantial overlapping of the short special intervals $J_k(q)$ defined by (3.5).

Recall that if $1 \leq q', q'' \leq q_{k+1} - 2$ are two integers such that $\{q'\alpha\}$ and $\{q''\alpha\}$ are neighboring points in the partition $\mathcal{A}_k(\alpha)$, then the intersection $J_k(q') \cap J_k(q'')$ must contain a special interval of the type $J_{k+8}(q)$ for some $1 \leq q \leq q_{k+9} - 2$. We split the argument into two complementary cases.

Write

$$Q(k; m) = \{y_i : q_{k+1} \leq i \leq m - q_{k+1}\}.$$

Case 1A. The following *intersection property* holds. For every

$$J_{k+8}(\ell) \subset J_k(q) \quad \text{and} \quad r \in \{0, 1, \dots, s-1\},$$

with $r \neq r_1^{(0)}(q)$ given by (5.6), we have

$$Q(k; m) \cap (r + J_{k+8}(\ell)) \neq \emptyset.$$

Case 1B. There exist

$$J_{k+8}(\ell_2) \subset J_k(q^{(1)}) \quad \text{and} \quad r_1 \in \{0, 1, \dots, s-1\},$$

with $r_1 \neq r_1^{(0)}(q^{(1)})$ given by (5.6), such that

$$Q(k; m) \cap (r_1 + J_{k+8}(\ell_2)) = \emptyset. \quad (5.9)$$

Since $J_{k+8}(\ell_2) \subset J_k(q^{(1)})$, it clearly follows from (5.7) that

$$Q(k; m) \cap (r_1^{(0)}(q^{(1)}) + J_{k+8}(\ell_2)) = \emptyset. \quad (5.10)$$

Since (5.9) and (5.10) are analogs of (5.4), the T -power expansion argument shows that for every integer $1 - \ell_2 \leq h \leq q_{k+9} - 2 - \ell_2$, we have

$$Q^{(1)}(k; m) \cap T^h(r_1 + J_{k+8}(\ell_2)) = \emptyset, \quad (5.11)$$

$$Q^{(1)}(k; m) \cap T^h(r_1^{(0)}(q^{(1)}) + J_{k+8}(\ell_2)) = \emptyset, \quad (5.12)$$

where, corresponding to $Q^*(k; m)$ in Section 4, we write

$$Q^{(1)}(k; m) = \{y_i : q_{k+1} + q_{k+9} \leq i \leq m - q_{k+1} - q_{k+9}\}.$$

For notational convenience, for every integer $1 - \ell_2 \leq h \leq q_{k+9} - 2 - \ell_2$, we write

$$T^h(r_1^{(0)}(q^{(1)}) + J_{k+8}(\ell_2)) = r_2^{(0)}(\ell_2 + h) + J_{k+8}(\ell_2 + h), \quad (5.13)$$

$$T^h(r_1 + J_{k+8}(\ell_2)) = r_2^{(1)}(\ell_2 + h) + J_{k+8}(\ell_2 + h). \quad (5.14)$$

Then combining (5.11)–(5.14), we have

$$Q^{(1)}(k; m) \cap (r_2^{(0)}(q) + J_{k+8}(q)) = \emptyset, \quad (5.15)$$

$$Q^{(1)}(k; m) \cap (r_2^{(1)}(q) + J_{k+8}(q)) = \emptyset, \quad (5.16)$$

for every integer $1 \leq q \leq q_{k+9} - 2$. Clearly

$$r_2^{(0)}(q) \neq r_2^{(1)}(q), \quad 1 \leq q \leq q_{k+9} - 2.$$

We now split Case 1B into two complementary cases.

Case 2A. The following *intersection property* holds. For every

$$J_{k+16}(\ell) \subset J_{k+8}(q) \quad \text{and} \quad r \in \{0, 1, \dots, s-1\},$$

with $r \neq r_2^{(0)}(q), r_2^{(1)}(q)$ given by (5.13) and (5.14), we have

$$Q^{(1)}(k; m) \cap (r + J_{k+16}(\ell)) \neq \emptyset.$$

Case 2B. There exist

$$J_{k+16}(\ell_3) \subset J_{k+8}(q^{(2)}) \quad \text{and} \quad r_2 \in \{0, 1, \dots, s-1\},$$

with $r_2 \neq r_2^{(0)}(q^{(2)}), r_2^{(1)}(q^{(2)})$ given by (5.13) and (5.14), such that

$$Q^{(1)}(k; m) \cap (r_2 + J_{k+16}(\ell_3)) = \emptyset. \quad (5.17)$$

Since $J_{k+16}(\ell_3) \subset J_{k+8}(q^{(2)})$, it clearly follows from (5.15) and (5.16) that

$$Q^{(1)}(k; m) \cap (r_2^{(0)}(q^{(2)}) + J_{k+16}(\ell_3)) = \emptyset, \quad (5.18)$$

$$Q^{(1)}(k; m) \cap (r_2^{(1)}(q^{(2)}) + J_{k+16}(\ell_3)) = \emptyset. \quad (5.19)$$

Since (5.17)–(5.19) are analogs of (5.4), the T -power expansion argument shows that for every integer $1 - \ell_3 \leq h \leq q_{k+17} - 2 - \ell_3$, we have

$$Q^{(2)}(k; m) \cap T^h(r_2 + J_{k+16}(\ell_3)) = \emptyset, \quad (5.20)$$

$$Q^{(2)}(k; m) \cap T^h(r_2^{(0)}(q^{(2)}) + J_{k+16}(\ell_3)) = \emptyset, \quad (5.21)$$

$$Q^{(2)}(k; m) \cap T^h(r_2^{(1)}(q^{(2)}) + J_{k+16}(\ell_3)) = \emptyset, \quad (5.22)$$

where, corresponding to $Q^{**}(k; m)$ in Section 4, we write

$$Q^{(2)}(k; m) = \{y_i : q_{k+1} + q_{k+9} + q_{k+17} \leq i \leq m - q_{k+1} - q_{k+9} - q_{k+17}\}.$$

For notational convenience, for every integer $1 - \ell_3 \leq h \leq q_{k+17} - 2 - \ell_3$, we write

$$T^h(r_2^{(0)}(q^{(2)}) + J_{k+16}(\ell_3)) = r_3^{(0)}(\ell_3 + h) + J_{k+16}(\ell_3 + h), \quad (5.23)$$

$$T^h(r_2^{(1)}(q^{(2)}) + J_{k+16}(\ell_3)) = r_3^{(1)}(\ell_3 + h) + J_{k+16}(\ell_3 + h), \quad (5.24)$$

$$T^h(r_2 + J_{k+16}(\ell_3)) = r_3^{(2)}(\ell_3 + h) + J_{k+16}(\ell_3 + h). \quad (5.25)$$

Then combining (5.20)–(5.25), we have

$$Q^{(2)}(k; m) \cap (r_3^{(0)}(q) + J_{k+16}(q)) = \emptyset,$$

$$Q^{(2)}(k; m) \cap (r_3^{(1)}(q) + J_{k+16}(q)) = \emptyset,$$

$$Q^{(2)}(k; m) \cap (r_3^{(2)}(q) + J_{k+16}(q)) = \emptyset,$$

for every integer $1 \leq q \leq q_{k+17} - 2$. Clearly

$$r_3^{(0)}(q), r_3^{(1)}(q), r_3^{(2)}(q) \text{ are distinct, } \quad 1 \leq q \leq q_{k+17} - 2.$$

We now split Case 2B into two complementary cases.

Case 3A. The following *intersection property* holds. For every

$$J_{k+24}(\ell) \subset J_{k+16}(q) \quad \text{and} \quad r \in \{0, 1, \dots, s-1\},$$

with $r \neq r_3^{(0)}(q), r_3^{(1)}(q), r_3^{(2)}(q)$ given by (5.23)–(5.25), we have

$$Q^{(2)}(k; m) \cap (r + J_{k+24}(\ell)) \neq \emptyset.$$

Case 3B. There exist

$$J_{k+24}(\ell_4) \subset J_{k+8}(q^{(3)}) \quad \text{and} \quad r_3 \in \{0, 1, \dots, s-1\},$$

with $r_3 \neq r_3^{(0)}(q), r_3^{(1)}(q), r_3^{(2)}(q)$ given by (5.23)–(5.25), such that

$$Q^{(2)}(k; m) \cap (r_3 + J_{k+24}(\ell_4)) = \emptyset.$$

Suppose that $1 \leq \tau \leq s-2$. Assume that for every integer $1 \leq q \leq q_{k+8\tau-7} - 2$, there are distinct integers

$$r_\tau^{(0)}(q), \dots, r_\tau^{(\tau-1)}(q) \in \{0, 1, \dots, s-1\} \quad (5.26)$$

such that for every integer $1 \leq j \leq \tau$,

$$Q^{(\tau-1)}(k; m) \cap (r_\tau^{(j-1)}(q) + J_{k+8\tau-8}(q)) = \emptyset \quad (5.27)$$

for every integer $1 \leq q \leq q_{k+8\tau-7} - 2$, where

$$Q^{(\tau-1)}(k; m) = \left\{ y_i : \sum_{u=1}^{\tau} q_{k+8u-7} \leq i \leq m - \sum_{u=1}^{\tau} q_{k+8u-7} \right\}.$$

Assume further that we have two complementary cases.

Case τ A. The following *intersection property* holds. For every

$$J_{k+8\tau}(\ell) \subset J_{k+8\tau-8}(q) \quad \text{and} \quad r \in \{0, 1, \dots, s-1\},$$

with $r \neq r_\tau^{(0)}(q), \dots, r_\tau^{(\tau-1)}(q)$ given by (5.26), we have

$$Q^{(\tau-1)}(k; m) \cap (r + J_{k+8\tau}(\ell)) \neq \emptyset.$$

Case τ B. There exist

$$J_{k+8\tau}(\ell_{\tau+1}) \subset J_{k+8\tau-8}(q^{(\tau)}) \quad \text{and} \quad r_\tau \in \{0, 1, \dots, s-1\},$$

with $r_\tau \neq r_\tau^{(0)}(q), \dots, r_\tau^{(\tau-1)}(q)$ given by (5.26), such that

$$Q^{(\tau-1)}(k; m) \cap (r_\tau + J_{k+8\tau}(\ell_{\tau+1})) = \emptyset. \quad (5.28)$$

Since $J_{k+8\tau}(\ell_{\tau+1}) \subset J_{k+8\tau-8}(q^{(\tau)})$, it clearly follows from (5.27) that for every integer $1 \leq j \leq \tau$,

$$Q^{(\tau-1)}(k; m) \cap (r_\tau^{(j-1)}(q^{(\tau)}) + J_{k+8\tau}(\ell_{\tau+1})) = \emptyset. \quad (5.29)$$

Since (5.28) and (5.29) are analogs of (5.4), the T -power expansion argument shows that for every integer $1 - \ell_{\tau+1} \leq h \leq q_{k+8\tau+1} - 2 - \ell_{\tau+1}$, we have

$$Q^{(\tau)}(k; m) \cap T^h(r_\tau + J_{k+8\tau}(\ell_{\tau+1})) = \emptyset, \quad (5.30)$$

as well as

$$Q^{(\tau)}(k; m) \cap T^h(r_\tau^{(j-1)}(q^{(\tau)}) + J_{k+8\tau}(\ell_{\tau+1})) = \emptyset \quad (5.31)$$

for every integer $1 \leq j \leq \tau$, where

$$Q^{(\tau)}(k; m) = \left\{ y_i : \sum_{u=1}^{\tau+1} q_{k+8u-7} \leq i \leq m - \sum_{u=1}^{\tau+1} q_{k+8u-7} \right\}.$$

For notational convenience, for every integer $1 - \ell_{\tau+1} \leq h \leq q_{k+8\tau+1} - 2 - \ell_{\tau+1}$, we write

$$T^h(r_\tau^{(j-1)}(q^{(\tau)}) + J_{k+8\tau}(\ell_{\tau+1})) = r_{\tau+1}^{(j-1)}(\ell_{\tau+1} + h) + J_{k+8\tau}(\ell_{\tau+1} + h) \quad (5.32)$$

for every integer $1 \leq j \leq \tau$, and also write

$$T^h(r_\tau + J_{k+8\tau}(\ell_{\tau+1})) = r_{\tau+1}^{(\tau)}(\ell_{\tau+1} + h) + J_{k+8\tau}(\ell_{\tau+1} + h). \quad (5.33)$$

Then combining (5.30)–(5.33), we have, for every integer $1 \leq j \leq \tau + 1$,

$$Q^{(\tau)}(k; m) \cap (r_{\tau+1}^{(j-1)}(q) + J_{k+8\tau}(q)) = \emptyset$$

for every integer $1 \leq q \leq q_{k+8\tau+1} - 2$. Clearly

$$r_{\tau+1}^{(0)}(q), \dots, r_{\tau+1}^{(\tau)}(q) \text{ are distinct, } 1 \leq q \leq q_{k+8\tau+1} - 2.$$

We now split Case τB into two complementary cases.

Case $(\tau + 1)A$. The following *intersection property* holds. For every

$$J_{k+8\tau+8}(\ell) \subset J_{k+8\tau}(q) \quad \text{and} \quad r \in \{0, 1, \dots, s-1\},$$

with $r \neq r_{\tau+1}^{(0)}(q), \dots, r_{\tau+1}^{(\tau)}(q)$ given by (5.32) and (5.33), we have

$$Q^{(\tau)}(k; m) \cap (r + J_{k+8\tau+8}(\ell)) \neq \emptyset.$$

Case $(\tau + 1)B$. There exist

$$J_{k+8\tau+8}(\ell_{\tau+2}) \subset J_{k+8\tau}(q^{(\tau+1)}) \quad \text{and} \quad r_{\tau+1} \in \{0, 1, \dots, s-1\},$$

with $r_{\tau+1} \neq r_{\tau+1}^{(0)}(q), \dots, r_{\tau+1}^{(\tau)}(q)$ given by (5.32) and (5.33), such that

$$Q^{(\tau)}(k; m) \cap (r_{\tau+1} + J_{k+8\tau+8}(\ell_{\tau+2})) = \emptyset.$$

In particular, if $\tau = s - 2$, we have the following final case.

Case $(s - 1)B$. There exist

$$J_{k+8s-8}(\ell_s) \subset J_{k+8s-16}(q^{(s-1)}) \quad \text{and} \quad r_{s-1} \in \{0, 1, \dots, s-1\},$$

with $r_{s-1} \neq r_{s-1}^{(0)}(q^{(s-1)}), \dots, r_{s-1}^{(s-2)}(q^{(s-1)})$ given by (5.32) and (5.33) in the special case $\tau = s - 2$, such that

$$Q^{(s-2)}(k; m) \cap (r_{s-1} + J_{k+8s-8}(\ell_s)) = \emptyset. \quad (5.34)$$

Lemma 5.2. *For every $\tau = 1, \dots, s - 1$, Case τA is impossible.*

Lemma 5.3. *If Case $(s - 1)B$ holds, then*

$$m \leq 2s + 2 \sum_{u=1}^s q_{k+8u-7}. \quad (5.35)$$

Before we prove Lemmas 5.2 and 5.3, we first complete the proof of Theorem 2.

Proof of Theorem 2. Suppose that $I_0 \subset [0, s)$ is $(\mathcal{L}_\alpha; M)$ -free, so that

$$\{y_i : 1 \leq i \leq m\} \cap I_0 = \emptyset.$$

Let $J_k(\ell_1) = J(\alpha; k; \ell_1)$ be the *longest* special interval of the form (3.5) such that

$$r(\ell_1) + J_k(\ell_1) \subset I_0 \quad \text{for some } \ell_1 \text{ and } r(\ell_1) \in \{0, 1, \dots, s-1\}.$$

Then we can show as before that

$$\text{length}(I_0) < \frac{8}{q_k}. \quad (5.36)$$

On the other hand, it follows from (5.1) and (5.2) that

$$M \leq (m + 1)\sqrt{1 + \alpha^2}. \quad (5.37)$$

Also, in view of Lemmas 5.2 and 5.3, it is clear that the bound (5.35) holds. Finally, the inequalities (3.33) are replaced by the inequalities

$$q_{k+1} < q_{k+9} < \dots < q_{k+8s-7} \leq (A + 1)^{8s-7} q_k. \quad (5.38)$$

Combining (5.35), (5.37) and (5.38), we see that

$$M \leq \left(2s + 1 + 2 \sum_{u=1}^s q_{k+8u-7} \right) \sqrt{1 + \alpha^2} < (4s + 1)(A + 1)^{8s-7} q_k \sqrt{2}. \quad (5.39)$$

It now follows from (5.36) and (5.39) that a geodesic segment $\mathcal{L}_\alpha(t)$, $0 \leq t \leq M$, of length $M = (4s + 1)(A + 1)^{8s-7} q_k \sqrt{2}$ must intersect every subinterval I of $h_1 \cup \dots \cup h_s$ with $\text{length}(I) = 8/q_k$. Since the product $M \text{length}(I)$ is a constant independent of k , this establishes superdensity of the half-infinite geodesic. \square

It remains to prove Lemmas 5.2 and 5.3.

Proof of Lemma 5.2. We can proceed along similar lines as in the first part of the proof of Lemma 3.2, and show that any two neighboring s -copy extensions $J_{k+8\tau-8}(q'; s)$ and $J_{k+8\tau-8}(q''; s)$ are *synchronized* in the following precise sense: For each $r \in \{0, 1, \dots, s-1\}$, we have

$$Q^{(\tau-1)}(k; m) \cap (r + J_{k+8\tau-8}(q')) = \emptyset$$

if and only if

$$Q^{(\tau-1)}(k; m) \cap (r + J_{k+8\tau-8}(q'')) = \emptyset.$$

By definition, the s -copy extensions $J_{k+8\tau-8}(q; s)$, $1 \leq q \leq q_{k+8\tau-7} - 2$, give rise to $2s$ continuous chains of overlapping intervals in the torus/circle $[0, s)$ such that the $2s$ chains completely cover the $2s$ intervals

$$[0, 1 - \alpha), \quad [1 - \alpha, 1), \quad \dots, \quad [s - 1, s - \alpha), \quad [s - \alpha, s). \quad (5.40)$$

The synchronization property now implies that each of the $2s$ long special intervals in (5.40) satisfies one of the following two properties. Either such a long special interval is *disjoint* from the set $Q^{(\tau-1)}(k; m)$, or the set $Q^{(\tau-1)}(k; m)$ is *dense* in such a long special interval, in the precise sense that every subinterval of length $1/q_{k+8s}$ contains a point from the set $Q^{(\tau-1)}(k; m)$.

Moreover, it is not difficult to show that precisely 2τ of the $2s$ long special intervals in (5.40) are disjoint from the set $Q^{(\tau-1)}(k; m)$. To see this, choose two integers q' and q'' satisfying $1 \leq q', q'' \leq q_{k+8\tau-7} - 2$ such that

$$J_{k+8\tau-8}(q') \subset (0, 1 - \alpha) \quad \text{and} \quad J_{k+8\tau-8}(q'') \subset (1 - \alpha, 1). \quad (5.41)$$

Then it follows from (5.27) that for every integer $1 \leq j \leq \tau$,

$$Q^{(\tau-1)}(k; m) \cap (r_\tau^{(j-1)}(q') + J_{k+8\tau-8}(q')) = \emptyset, \quad (5.42)$$

$$Q^{(\tau-1)}(k; m) \cap (r_\tau^{(j-1)}(q'') + J_{k+8\tau-8}(q'')) = \emptyset. \quad (5.43)$$

For every integer $1 \leq j \leq \tau$, now write

$$\mathcal{I}_{2j-1} = r_\tau^{(j-1)}(q') + (0, 1 - \alpha) \quad \text{and} \quad \mathcal{I}_{2j} = r_\tau^{(j-1)}(q'') + (1 - \alpha, 1). \quad (5.44)$$

The synchronization property and (5.41)–(5.44) now imply that for every integer $1 \leq j \leq \tau$, we have

$$Q^{(\tau-1)}(k; m) \cap \mathcal{I}_{2j-1} = \emptyset \quad \text{and} \quad Q^{(\tau-1)}(k; m) \cap \mathcal{I}_{2j} = \emptyset.$$

Note that the union $\mathcal{I}_1 \cup \dots \cup \mathcal{I}_{2\tau}$ modulo 1 is precisely the unit interval $[0, 1)$ taken τ times.

Now $\mathcal{I}_1, \dots, \mathcal{I}_{2\tau}$ are 2τ of the $2s$ long special intervals in (5.40). Let the remaining long special intervals in (5.40) be denoted by \mathcal{I}_{2j-1} and \mathcal{I}_{2j} , $\tau < j \leq s$. The condition of Case τA now implies that these $2s - 2\tau$ intervals are not disjoint from $Q^{(\tau-1)}(k; m)$, so that $Q^{(\tau-1)}(k; m)$ is dense in each of them.

Each T -image $T(\mathcal{I}_{2j-1})$ and $T(\mathcal{I}_{2j})$, $1 \leq j \leq \tau$, has at most 1 common point with the set $Q^{(\tau-1)}(k; m)$. This is a contradiction, since the union

$$T(\mathcal{I}_1) \cup \dots \cup T(\mathcal{I}_{2\tau})$$

has a *substantial* intersection with the union $\mathcal{I}_{2\tau+1} \cup \dots \cup \mathcal{I}_{2s}$, which implies that it must have a *substantial* intersection with the set $Q^{(\tau-1)}(k; m)$, much more than at most 2τ elements. Thus Case τA is impossible, and this completes the proof. \square

Proof of Lemma 5.3. Since (5.34) is an analog of (5.4), we can repeat the T -power extension argument and conclude that for every integer $1 - \ell_s \leq h \leq q_{k+8s-7} - 2 - \ell_s$, we have

$$Q^{(s-1)}(k; m) \cap T^h(r_{s-1} + J_{k+8s-8}(\ell_s)) = \emptyset. \quad (5.45)$$

Since $J_{k+8s-8}(\ell_s) \subset J_{k+8s-16}(q^{(s-1)})$, it follows from (5.27) with $\tau = s - 1$ that for every integer $1 \leq j \leq s - 1$,

$$Q^{(s-2)}(k; m) \cap (r_{s-1}^{(j-1)}(q^{(s-1)}) + J_{k+8s-8}(\ell_s)) = \emptyset. \quad (5.46)$$

Next, note that (5.46) are also analogs of (5.4), so again we can repeat the T -power extension argument and conclude that for every integer $1 \leq j \leq s - 1$, and for every integer $1 - \ell_s \leq h \leq q_{k+8s-7} - 2 - \ell_s$, we have

$$Q^{(s-1)}(k; m) \cap T^h(r_{s-1}^{(j-1)}(q^{(s-1)}) + J_{k+8s-8}(\ell_s)) = \emptyset. \quad (5.47)$$

Now, for every integer $1 - \ell_s \leq h \leq q_{k+8s-7} - 2 - \ell_s$, we write

$$T^h(r_{s-1}^{(j-1)}(q^{(s-1)}) + J_{k+8s-8}(\ell_s)) = r^{(j-1)}(\ell_s + h) + J_{k+8s-8}(\ell_s + h) \quad (5.48)$$

for every integer $1 \leq j \leq s - 1$, and also write

$$T^h(r_{s-1} + J_{k+8s-8}(\ell_s)) = r^{(s-1)}(\ell_s + h) + J_{k+8s-8}(\ell_s + h). \quad (5.49)$$

Clearly it follows from the assumption of Case $(s - 1)B$ that $r^{(0)}(q), \dots, r^{(s-1)}(q)$ form a permutation of $0, 1, \dots, s - 1$ for every integer $1 \leq q \leq q_{k+8s-7} - 2$. Combining (5.45) and (5.47)–(5.49), we have, for every integer $1 \leq j \leq s$,

$$Q^{(P-1)}(k; m) \cap (r^{(j-1)}(q) + J_{k+8s-8}(q)) = \emptyset \quad (5.50)$$

for every integer $1 \leq q \leq q_{k+8s-7} - 2$.

Note now that (5.50) are similar to (4.7)–(4.9) in the proof of Lemma 4.1, so we now mimic the last part of that proof.

By definition, the s -copy extensions $J_{k+8s-8}(q; s)$, $1 \leq q \leq q_{k+8s-7} - 2$, give rise to $2s$ continuous chains of overlapping intervals in the torus/circle $[0, s)$ such that the $2s$ chains completely cover the $2s$ intervals

$$[0, 1 - \alpha), \quad [1 - \alpha, 1), \quad \dots, \quad [s - 1, s - \alpha), \quad [s - \alpha, s),$$

and there are only $2s$ points in $[0, s)$ that are not covered by the $2s$ chains, namely

$$j - 1, \quad j - \alpha, \quad 1 \leq j \leq s.$$

Combining (5.50) for every integer $1 \leq q \leq q_{k+8s-7} - 2$, we deduce that the set $Q^{(s-1)}(k; m)$ is not covered by the $2s$ chains. Indeed, if

$$m \geq 2s + 1 + 2 \sum_{u=1}^s q_{k+8u-7},$$

then the set $Q^{(s-1)}(k; m)$ has at least $2s + 1$ distinct elements, which is more than $2s$, giving rise to a contradiction. We conclude therefore that, under the conditions of Case $(s - 1)B$, we must have

$$m \leq 2s + 2 \sum_{u=1}^s q_{k+8u-7},$$

and this completes the proof. \square

REFERENCES

- [1] J. Beck, M. Donders, Y. Yang. Quantitative behavior of non-integrable systems (I). *Acta Math. Hungar.* **161** (2020), 66–184.
- [2] J. Beck, M. Donders, Y. Yang. Quantitative behavior of non-integrable systems (II). *Acta Math. Hungar.* **162** (2020), 220–324.
- [3] J. Beck, W.W.L. Chen, Y. Yang. Quantitative behavior of non-integrable systems (III). arxiv.org/abs/2006.06213, 93 pp.
- [4] J. Beck, W.W.L. Chen, Y. Yang. Quantitative behavior of non-integrable systems (IV). arxiv.org/abs/2012.12038, 120 pp.
- [5] J.H. Halton. The distribution of the sequence $\{n\xi\}$ ($n = 0, 1, 2, \dots$). *Proc. Cambridge Philos. Soc.* **61** (1965), 665–670.
- [6] G.H. Hardy, J.E. Littlewood. Some problems of diophantine approximation: the lattice points of a right-angled triangle I. *Proc. London Math. Soc.* **3** (1920), 15–36.
- [7] G.H. Hardy, J.E. Littlewood. Some problems of diophantine approximation: the lattice points of a right-angled triangle II. *Abh. Math. Sem. Univ. Hamburg* **1** (1922), 212–249.
- [8] A. Katok, A. Zemlyakov. Topological transitivity of billiards in polygons. *Math. Notes* **18** (1975), 760–764.
- [9] S. Kerckhoff, H. Masur, J. Smillie. Ergodicity of billiard flows and quadratic differentials. *Ann. of Math.* **124** (1986), 293–311.
- [10] A.Ya. Khinchin. Einige Sätze über Kettenbrüche, mit Anwendungen auf die Theorie der Diophantischen Approximationen. *Math. Ann.* **92** (1924), 115–125.
- [11] A.Ya. Khinchin. *Continued Fractions* (Dover, 1997).
- [12] D. König, A. Szücs. Mouvement d’un point abandonné à l’intérieur d’un cube. *Rend. Circ. Mat. Palermo* **36** (1913), 79–90.
- [13] H. Masur, S. Tabachnikov. Rational billiards and flat surfaces. *Handbook of Dynamical Systems*, vol. 1A (B. Hasselblatt, A. Katok, eds.), pp. 1017–1089 (Elsevier, 2006).
- [14] A. Ostrowski. Bemerkungen zur Theorie der Diophantischen Approximationen. *Abh. Math. Sem. Univ. Hamburg* **1** (1922), 77–98.
- [15] N.B. Slater. Gap and steps for the sequence $n\theta \bmod 1$. *Proc. Cambridge Philos. Soc.* **63** (1967), 1115–1123.
- [16] V.T. Sós. On the theory of diophantine approximations. *Acta Math. Acad. Sci. Hungar.* **8** (1957), 461–472.
- [17] V.T. Sós. On the distribution mod 1 of the sequence $n\alpha$. *Ann. Univ. Sci. Budapest Eötvös Sect. Math.* **1** (1958), 127–134.
- [18] J. Surányi. Über die Anordnung der Vielfachen einer reellen Zahl mod 1. *Ann. Univ. Sci. Budapest Eötvös Sect. Math.* **1** (1958), 107–111.
- [19] S. Swierczkowski. On successive settings of an arc on the circumference of a circle. *Fund. Math.* **46** (1959), 187–189.
- [20] H. Weyl. Über die Gleichverteilung von Zahlen mod. Eins. *Math. Ann.* **77** (1916), 313–352.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, HILL CENTER FOR THE MATHEMATICAL SCIENCES, PISCATAWAY NJ 08854, USA

Email address: jbeck@math.rutgers.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, MACQUARIE UNIVERSITY, SYDNEY NSW 2109, AUSTRALIA

Email address: william.chen@mq.edu.au