

# A computational proof of the existence of the Dual Isogeny

Markos Karameris<sup>1</sup>[0000-0001-7009-5285]

National Kapodistrian University of Athens

**Abstract.** For  $E$  an elliptic curve over a perfect field  $K$ , we present a proof of the existence of the dual isogeny  $\hat{\phi}$  using computational methods linked to Vélú's formulae instead of the standard Galois correspondence method.

**Keywords:** Elliptic Curves · Isogenies · Vélú's formula · Alternative Proof

## 1 Preliminaries and Notation

Let  $E(K)$  denote an elliptic curve over a perfect field  $K$ . For simplicity we assume that  $\text{char}(K) \neq 2$  or  $3$ . We will mostly preserve the notation of [1].

To a finite subgroup of  $E$  we can associate an isogeny  $\phi : E \rightarrow E'$ . This isogeny is not uniquely determined but if we require the isogeny to be normalized then we do get a unique morphism.

**Definition 1.** *If  $\omega'$  is the invariant differential of  $E'$  then the pullback  $\phi^*(\omega')$  is a holomorphic differential of  $E$  and since the space  $\Omega_E$  is 1-dimensional we have  $\phi^*(\omega') = c\omega$  for some  $c \in K$ . We call  $\phi$  normalized iff  $c = 1$ .*

The proof of the existence is presented in [1] in two parts:

**Proposition 1.** *Let  $\phi : E \rightarrow E_1$  and  $\psi : E \rightarrow E_2$  be two isogenies with  $\phi$  seperable and  $\ker(\phi) \subset \ker(\psi)$ . Then there exists a unique isogeny  $\lambda : E_1 \rightarrow E_2$  with  $\psi = \lambda \circ \phi$ .*

The theorem is then proved in Theorem 6.1 using the above proposition for  $\psi = [m]$ . Notice that requiring the isogeny to be seperable is natural as an inseperable isogeny cannot be normal. The primary tools we will be using are Vélú's formulae: Let  $G$  be a subgroup of  $E(K)$  and write  $P = (x_P, y_P)$  then as in [2] the formula we will be using is:

$$\phi(P) = (x(P) + \sum_{Q \in G \neq O} x(P+Q) - x(Q), y(P) + \sum_{Q \in G \neq O} y(P+Q) - y(Q)), P \notin G \quad (1)$$

where we also set  $\phi(P) = O, \forall P \in G$  by definition.

*Remark 1.* The above definition is invariant under the action of  $G$  and it's kernel is exactly  $G$  as intended. It can only be used to describe a normalized isogeny  $\phi$  which follows from the uniqueness property.

## 2 Existence of the Dual Isogeny

The first result we will prove is the same as that of Proposition 1 but in the case of normalized elliptic curve isogenies  $\phi, \psi$ .

**Proposition 2.** *Let  $\phi : E \rightarrow E_1$  and  $\psi : E \rightarrow E_2$  be two normalized isogenies with  $\ker(\phi) \subset \ker(\psi)$ . Then there exists a unique isogeny  $\lambda : E_1 \rightarrow E_2$  with  $\psi = \lambda \circ \phi$ .*

*Proof.* If  $\phi : E \rightarrow E_1$  is normalized then we have that  $\phi(P) = (\sum_{Q \in G'}(x_{P+Q} - x_Q), \sum_{Q \in G'}(y_{P+Q} - y_Q))$ . But then  $\lambda(P) = (\sum_{g \in G/G'}(x_{P+\phi(g)} - \phi_x(g)), \sum_{g \in G/G'}(y_{P+\phi(g)} - \phi_y(g)))$ . We observe that:  $x_{\phi(P)+\phi(g)} = x_{\phi(P+g)}$  and similarly for  $y$  we get:  $\lambda \circ \phi(P) = (\sum_{g \in G/G'}(\phi_x(P+g) - \phi_x(g)), \sum_{g \in G/G'}(\phi_y(P+g) - \phi_y(g))) = (\sum_{g \in G/G'}(\sum_{Q \in G'}(x_{P+g+Q} - x_{g+Q})), \sum_{g \in G/G'}(\sum_{Q \in G'}(y_{P+g+Q} - y_{g+Q}))) = (\sum_{Q \in G}(x_{P+Q} - x_Q), \sum_{Q \in G}(y_{P+Q} - y_Q))$  (for a homomorphism  $h$  with normal subgroup  $G'$ ,  $\sum_{g \in G} h(g) = \sum_{g \in G/G'} \sum_{Q \in G} h(g+Q)$ ). And thus we obtain a normal isogeny with kernel  $G$  which is also unique as it is normalized with  $\lambda \circ \phi(P) = \psi(P)$ . The fact  $\lambda$  is an isogeny  $E_1 \rightarrow E_2$  is immediate  $\lambda(O) = \lambda(\phi(O)) = \psi(O) = O$  and  $\lambda(E_2) = \lambda(\phi(E_1)) = \psi(E_1) = E_3$  as  $\phi, \psi$  are onto.

So far we restricted our results to normal isogenies however:

**Proposition 3.** *For every seperable isogeny  $\phi : E \rightarrow E'$  there exists an isomorphism  $i : E' \rightarrow E''$  such that  $i \circ \phi$  is normalized*

*Proof.* for a proof see [3].

We are now ready to prove the original statement of Proposition 1. for seperable isogenies.

*Proof.* Let  $i_\phi, i_\psi$  be the corresponding isomorphisms such that  $\phi' = i_\phi \circ \phi$  and  $\psi' = i_\psi \circ \psi$  are both normalized. Then by Proposition 2 there is a  $\lambda$  such that  $\psi' = \lambda \circ \phi'$ . Then  $i_\psi^{-1} \circ \lambda \circ i_\phi$  is the required isogeny  $E_1 \rightarrow E_2$ .

*Remark 2.* The fact that we require both  $\phi$  and  $\psi$  to be seperable does not cause any problems as in the case of inseperable  $\psi$  we can simply write  $\psi = \pi^n \circ \psi_{sep}$  where  $\pi$  is the Frobenius endomorphism (this is possible because we are in a perfect field or even over a finite field of characteristic  $p$  essentially). Applying our result to  $\phi, \psi_{sep}$  we get an isogeny  $\lambda : \lambda \circ \phi = \psi_{sep} \implies \pi^n \circ \lambda \circ \phi = \psi$ . The required isogeny is thus  $\lambda' = \pi^n \circ \lambda$ .

The above remark settles Proposition 1.

### 3 Computational Aspects

The proposed approach also offers a way to compute the dual isogeny directly using Vélú's formulae. Let  $\phi : E \rightarrow E/G$  be an isogeny over  $K$ , then the required steps are:

1. compute  $\phi_{sep}$  and  $n \in \mathbb{N}$  such that  $\phi = \phi_{sep} \circ \pi^n$
2. compute  $\hat{\pi}$  via the  $[p]$  endomorphism
3. compute  $[m]_{sep}$  and  $e \in \mathbb{N}$  such that  $[m] = \pi^e \circ [m]_{sep}$
4. normalize  $\phi_{sep}$  using  $i_\phi$  into  $\phi_{norm}$
5. normalize  $[m]_{sep}$  using  $i_{[m]}$  into  $[m]_{norm}$
6. using Vélú's formulae this gives  $\lambda : \lambda \circ \phi_{norm} = [m]_{norm}$
7. set  $\widehat{\phi_{sep}} = i_{[m]}^{-1} \circ \pi^e \circ \lambda \circ i_\phi$ , the dual isogeny is  $\widehat{\phi} = \hat{\pi}^n \circ \widehat{\phi_{sep}}$

Separating the separable and inseparable part is a straightforward procedure: writing  $\phi$  in terms of rational functions it simply means finding the greatest power of  $p$  dividing all the exponents. The second part is also immediate from 1, simply write  $[p] = [p]_{sep} \circ \pi^k$  and take  $\hat{\pi} = [p]_{sep} \circ \pi^{k-1}$ . The only remaining step is the normalization process which we will describe explicitly.

*Remark 3.* Given an isogeny  $\phi : E \rightarrow E'$ , we are looking for an isomorphism  $i_\phi : E' \rightarrow E''$  such that  $(i_\phi \circ \phi)^*(\omega') = \omega$ . Writing  $\omega = \frac{dx}{2y+a_1x+a_3}$ ,  $\omega' = \frac{dx}{2y+a_1'x+a_3'}$  and  $\phi(x, y) = (r(x), s(x)y + z(x))$  in rational form, abusing notation we get the equation  $\frac{d(i_\phi(r(x)))}{2i_\phi(s(x)y+z(x))+a_1'i_\phi(r(x))+a_3'} = \frac{dx}{2y+a_1x+a_3}$

We can tackle this problem by noting that  $i_\phi(x, y) = (u^2x + t_1, u^3y + t_2u^2x + t_3)$  for some  $t_i \in K$  which leads us to the following lemma:

**Lemma 1.** *Let  $i_\phi : E \rightarrow E'$  be an isomorphism as above, then  $i_\phi^*(\omega') = \frac{1}{u}\omega$*

*Proof.* The proof is a direct computation along with the observation that the nominator has a term of the form  $u^3$  in the denominator and  $u^2$  in the nominator.

Using the Lemma above we immediately obtain the desired result:

- for the isogeny  $[m]_{sep}$  we simply set  $u = (\frac{m}{p^e})^{-1}$  where  $p^{e+1} \nmid m$
- for  $\phi_{sep}$  we compute  $\phi_{sep}^*(\omega') = c\omega$  and set  $u = c^{-1}$

Combining these results enables us to compute explicitly the dual isogeny without resorting to the usual method of computing via the preimage of the arbitrary isogeny  $\phi$ .

## References

1. J. Silverman, The Arithmetic of Elliptic Curves. 2nd edn. Springer, San Fransisco (2008)
2. Josep M. Miret, Ramiro Moreno and Anna Rio, Generalizations of Vélus formulae for isogenies between elliptic curves, Publ. Mat. (2007)
3. Algorithms for Algebraic Curves, <https://www.math.u-bordeaux.fr/~jcouveig/cours/course5bis.pdf>. Last accessed 8 April 2021