

HENSEL MINIMALITY II: MIXED CHARACTERISTIC AND A DIOPHANTINE APPLICATION

RAF CLUCKERS, IMMANUEL HALUPCZOK, SILVAIN RIDEAU-KIKUCHI,
AND FLORIS VERMEULEN

ABSTRACT. We extend the results of [8] on Hensel minimality to include also the mixed characteristic case. This completes our axiomatic framework for tame non-archimedean geometry over Henselian valued fields of characteristic zero. We prove a Jacobian property, a strong form of Taylor approximations of definable functions, respresendency results and cell decomposition, all under 1-h-minimality. We obtain a diophantine application of counting rational points of bounded height on Hensel minimal curves.

CONTENTS

1. Introduction	1
2. Hensel minimality in characteristic zero (including mixed characteristic)	3
3. Geometry of definable sets in 1-h-minimal structures	20
4. Diophantine application for 1-h-minimal transcendental curves	26
5. Some questions	29
References	31

1. INTRODUCTION

1.1. For a long time one has been looking for adequate analogues of o-minimality in other settings than real closed fields. Several notions have been put forward,

2010 *Mathematics Subject Classification.* Primary 03C99, 14G05; Secondary 03C65, 12J20, 11G50, 11D88, 03C98, 14E18, 41A58.

Key words and phrases. Non-archimedean geometry, tame geometry on Henselian valued fields, analogues to o-minimality, model theory of valued fields, cell decomposition, Taylor approximation, Lipschitz continuity, rational points of bounded height, parametrizations.

The author R. C. was partially supported by the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) with ERC Grant Agreement nr. 615722 MOTMELSUM, by KU Leuven IF C14/17/083, and thanks the Labex CEMPI (ANR-11-LABX-0007-01). The author I. H. was partially supported by the research training group *GRK 2240: Algebro-Geometric Methods in Algebra, Arithmetic and Topology* and by the individual research grant No. 426488848, both funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation). The author S. R. was partially supported by GeoMod AAPG2019 (ANR-DFG), Geometric and Combinatorial Configurations in Model Theory. The author F. V. was partially supported by KU Leuven IF C14/17/083, and partially by F.W.O. Flanders (Belgium) with grant number 11F1921N.

each with certain strengths and weaknesses. In Part I of this work, [8], we have put forward a notion for tame geometry on non-archimedean values fields, called Hensel minimality, both simple and strong and providing a common variant for the settings from e.g. [13, 15, 12, 14, 9]. It is similar to o-minimality [10, 18] both in its simplicity and its strong consequences. In Part I we have focused on the equicharacteristic zero case. In this sequel, we focus on the mixed characteristic case, and we give a diophantine application similar to the results by Pila-Wilkie [17] in the o-minimal case, but for the moment this application is for curves only.

In the mixed characteristic case, there are several candidate adaptations to the notion of Hensel minimality from [8], based on coarsenings, compactness, or on a more literal adaptation of the definitions for the equi-characteristic zero case. We treat several of these variants, and in a most important case we show that they are equivalent. From this equivalence we derive the geometric results of [8] also in the mixed characteristic case, with in particular some strong results on Taylor approximation of definable functions.

Central in the definition of Hensel minimality from [8] is the condition that unary definable subsets X of the valued field K can be prepared by finite sets C (think of the role of the finite set of endpoints of intervals in o-minimality for preparing unary sets in \mathbb{R}). In addition, one imposes conditions on the parameters over which C is definable. (In the o-minimal case, this comes for free.) This is mimicked in Definition 2.2.1. It is surprising and rewarding that such basic conditions suffice to understand so much about the geometry of definable sets and functions.

1.2. Let us give a short global overview of the paper. In Section 2 we introduce several variants of Hensel minimality (for both the mixed and the equi-characteristic cases), and we prove that all of them are equivalent in the key case of 1-h-minimality (Theorem 2.2.7). We do this for a large part by classical strategies: coarsening of valuations of mixed characteristic to equi-characteristic zero, and model theoretic compactness. A combination of these classical strategies with new geometric and model theoretic arguments lead to our strongest results, like Theorem 3.1.2 on Taylor approximation. It is precisely this result Theorem 3.1.2 on Taylor approximation that plays a key role in our diophantine application Theorem 4.1.6, where we estimate the number of rational points of bounded height on transcendental curves. This is the Hensel minimal analogue of point counting on transcendental curves case as treated by Bombieri and Pila [4] and o-minimally by Pila-Wilkie [17], and, it is the axiomatic analogue of point counting on subanalytic sets as studied in [6] and [7].

In Section 2.6, we develop some resplendency results analogous to the ones from [8, Section 4]; they are used to show the equivalences of Theorem 2.2.7. It follows from the equivalence with item (4) in Theorem 2.2.7 that 1-h-minimality is preserved under coarsening the valuation; such a coarsening result was previously known only for ω -h-minimality by [8, Corollary 4.2.4].

One of the big open challenges in the current framework is to push the diophantine application further towards arbitrary dimension, and thus to get a full, Hensel

minimal analogue of the results by Pila-Wilkie [17]. Other main challenges are further developments of the geometry, like t-stratifications building on and extending [8, Section 5.5]. It may also be interesting to compare the notions of Hensel minimal structures with distal expansions on valued fields as studied recently in [1]. We conclude the paper with some related open questions.

2. HENSEL MINIMALITY IN CHARACTERISTIC ZERO (INCLUDING MIXED CHARACTERISTIC)

In this whole section we fix a theory \mathcal{T} of valued fields of characteristic zero in a language \mathcal{L} expanding the language $\mathcal{L}_{\text{val}} = \{+, \cdot, \mathcal{O}_K\}$ of valued fields. Note that \mathcal{T} is allowed to be non-complete, and that each model K of \mathcal{T} is a valued field of characteristic zero, which includes both possibilities that K has mixed characteristic or equicharacteristic zero.

In this section we give four alternative definitions of 1-h-minimality for \mathcal{T} (see Definitions 2.2.1, 2.2.3, 2.2.4, 2.2.6), and we show that they are equivalent in Theorem 2.2.7. To keep the generality of [8], we will treat more generally ℓ -h-minimality for $\ell \geq 0$ either an integer or equal to ω . The first definition is a close adaptation of the main notion of Hensel minimality of [8, Definition 2.3.3]. Definitions 2.2.3 and 2.2.4 are based on coarsenings, and Definition 2.2.6 corresponds to the criterion for 1-h-minimality from [8, Theorem 2.9.1]. Since at least for $\ell = 1$ these notions coincide, we simply will call them 1-h-minimality.

2.1. Basic terminology. We use terminology and concepts from [8]. Let us quickly recall some of these. By a valued field we mean a non-trivially a valued field, i.e., the field of fractions of a valuation ring which is not a field itself. Any valued field K is a structure in the language $\mathcal{L}_{\text{val}} = \{+, \cdot, \mathcal{O}_K\}$ of valued fields, where $+$ and \cdot are addition and multiplication on K , and where \mathcal{O}_K is the valuation ring of K . The maximal ideal of \mathcal{O}_K is denoted by \mathcal{M}_K . We use multiplicative notation for the value group which we denote by Γ_K^\times and we write $\Gamma_K := \Gamma_K^\times \cup \{0\}$. We write $|\cdot|: K \rightarrow \Gamma_K$ for the valuation map. By an open ball we mean a ball of the form $B_{<\lambda}(a) = \{x \in K \mid |x - a| < \lambda\}$ for some λ in Γ_K^\times and some a in K . We define $\text{rad}_{\text{op}} B_{<\lambda}(a) = \lambda$ to be the open radius of such a ball. Similarly, a closed ball is a set of the form $B_{\leq\lambda}(a) = \{x \in K \mid |x - a| \leq \lambda\}$. Its closed radius is $\text{rad}_{\text{cl}} B_{\leq\lambda}(a) = \lambda$.

For any proper ideal I of the valuation ring \mathcal{O}_K , we write RV_I for the disjoint union of $\{0\}$ with the quotient group $K^\times/1+I$, and $\text{rv}_I: K \rightarrow \text{RV}_I$ for the quotient map extended by sending 0 to 0. When I is the open ball $\{x \in K \mid |x| < \lambda\}$ for some $\lambda \leq 1$ in Γ_K^\times , we simply write RV_λ and rv_λ instead of RV_I and rv_I , and we write RV and rv instead of RV_1 and rv_1 .

We will sometimes also write RV_K for RV and similarly $\text{RV}_{K,\lambda}$ for RV_λ if multiple fields are under consideration.

We recall some key notions from [8]:

2.1.1. Definition. Let $\lambda \leq 1$ be an element of Γ_K^\times .

- (1) Given an arbitrary set $X \subset K$ and a finite non-empty set $C \subset K$, we say that X is λ -prepared by C if the condition whether some $x \in K$ lies in X depends only on the tuple $(\text{rv}_\lambda(x - c))_{c \in C}$.
- (2) We say that a ball $B \subset K$ is λ -next to an element $c \in K$ if

$$B = \{x \in K \mid \text{rv}_\lambda(x - c) = \xi\}$$

for some (nonzero) element ξ of RV_λ .

- (3) We say that a ball $B \subset K$ is λ -next to finite non-empty set $C \subset K$ if B equals $\bigcap_{c \in C} B_c$ with B_c a ball λ -next to c for each $c \in C$.

We recall some notation from [8, Section 6].

2.1.2. Definition (Equi-characteristic 0 coarsening). Given a model $K \models \mathcal{T}$, we write $\mathcal{O}_{K,\text{ecc}}$ for the smallest subring of K containing \mathcal{O}_K and \mathbb{Q} and we let $|\cdot|_{\text{ecc}}: K \rightarrow \Gamma_{K,\text{ecc}}$ be the corresponding valuation. (Thus, $|\cdot|_{\text{ecc}}$ is the finest coarsening of $|\cdot|$ which has equi-characteristic 0; note that $|\cdot|_{\text{ecc}}$ can be a trivial valuation on K .) If $|\cdot|_{\text{ecc}}$ is a nontrivial valuation (i.e., $\mathcal{O}_{K,\text{ecc}} \neq K$), then we also use the following notation: $\text{rv}_{\text{ecc}}: K \rightarrow \text{RV}_{\text{ecc}}$ is the leading term map with respect to $|\cdot|_{\text{ecc}}$; given $\lambda \in \Gamma_{K,\text{ecc}}$, $\text{rv}_\lambda: K \rightarrow \text{RV}_\lambda$ is the leading term map with respect to λ ; and \mathcal{L}_{ecc} is the expansion of \mathcal{L} by a predicate for $\mathcal{O}_{K,\text{ecc}}$. More generally, for any non-trivial coarsening $|\cdot|_c$ of the valuation on K , write \mathcal{L}_c for the expansion of \mathcal{L} by a predicate for the valuation ring for $|\cdot|_c$.

2.2. Equivalent definitions. There are several natural notions of Hensel minimality in mixed characteristic. We give four possible definitions. Theorem 2.2.7 states that these are all equivalent, in the case of 1-h-minimality.

We first adapt the main definition of Hensel minimality from [8, Definition 2.3.3] to include the mixed characteristic case. Recall that \mathcal{T} is a theory of valued fields of characteristic zero in a language $\mathcal{L} \supset \mathcal{L}_{\text{val}}$.

2.2.1. Definition. Let $\ell \geq 0$ be either an integer or ω . Say that \mathcal{T} is ℓ - h^{mix} -minimal if for each model K of \mathcal{T} , for each integer $n \geq 1$, each $\lambda \leq 1$ in Γ_K^\times , each $A \subset K$, each finite $A' \subset \text{RV}_\lambda$ of cardinality $\#A' \leq \ell$, and each $(A \cup \text{RV}_{|n|} \cup A')$ -definable set $X \subset K$, there exists an integer $m \geq 1$ such that X is $|m|\lambda$ -prepared by a finite A -definable set $C \subset K$.

Clearly, if all models K of \mathcal{T} are of equicharacteristic zero, then \mathcal{T} is ℓ - h^{mix} -minimal if and only if \mathcal{T} is ℓ - h -minimal in the sense of [8, Definition 2.3.3]. In the mixed characteristic case, one is obliged to take the valuation of integers into account in Definition 2.2.1. The general philosophy is that λ -preparation in equi-characteristic 0 becomes $\lambda \cdot |m|$ -preparation in mixed characteristic, for some integer $m \geq 1$, as the following example illustrates.

2.2.2. Example. The set X of cubes in the 3-adic numbers \mathbb{Q}_3 cannot be 1-prepared by any finite set C , since each of the infinitely many disjoint balls $27^r(1+3\mathbb{Z}_p)$, $r \in \mathbb{Z}$, contains both, cubes and non-cubes. However, X is a union of fibers of the map $\text{rv}_{|3|}: \mathbb{Q}_3 \rightarrow \text{RV}_{|3|}$, so it is $|3|$ -prepared by the set $\{0\}$.

The following two definitions use equicharacteristic zero coarsenings of the valuation, with notation from Definition 2.1.2.

2.2.3. Definition. Let $\ell \geq 0$ be either an integer or ω . We say that \mathcal{T} is ℓ - h^{ecc} -minimal if for every model $K \models \mathcal{T}$ the following holds: If the valuation $|\cdot|_{\text{ecc}}$ on K is non-trivial, then the \mathcal{L}_{ecc} -theory of K , when considered as a valued field with the valuation $|\cdot|_{\text{ecc}}$, is ℓ - h -minimal in the sense of [8, Definition 2.3.3].

One can also require every equi-characteristic zero coarsening to be ℓ - h -minimal, leading to the following definition.

2.2.4. Definition. Let $\ell \geq 0$ be either an integer or ω . We say that \mathcal{T} is ℓ - h^{coars} -minimal if for every model $K \models \mathcal{T}$ and each non-trivial equicharacteristic coarsening $|\cdot|_c$ of the valuation on K , the \mathcal{L}_c -theory of K , when considered as a valued field with the valuation $|\cdot|_c$, is ℓ - h -minimal in the sense of [8, Definition 2.3.3].

2.2.5. Remark. For theories of fields of equi-characteristic zero valued fields, ℓ - h^{ecc} -minimality is clearly equivalent to ℓ - h -minimality in the sense of [8, Definition 2.3.3], for each $\ell \geq 0$. For more subtle reasons, ω - h^{coars} -minimality is equivalent to ω - h^{ecc} -minimality by [8, Corollary 4.2.4]. For $\ell = 1$, we will show the analogue of this (and more) in Theorem 2.2.7.

By the usual play of compactness, preparation results that hold in each model of \mathcal{T} also hold uniformly for all models of \mathcal{T} , and results in equi-characteristic can be transferred to mixed characteristic. We will give examples in the proofs for Corollary 2.3.8 and Corollary 2.5.5 to illustrate how compactness is used for these purposes.

The criterion from [8, Theorem 2.9.1] for 1- h -minimality in equi-characteristic zero leads to the following alternative definition in mixed characteristic.

2.2.6. Definition. Let \mathcal{T} be a theory of valued fields of characteristic 0 (with any residue characteristic) in a language \mathcal{L} expanding the language of valued fields. Let $f : K \rightarrow K$ be A -definable, for some set $A \subset K \cup \text{RV}_{|n|}$ for some positive integer n . We define the following two properties:

(T1^{mix}) There exists a finite A -definable $C \subset K$ and a positive integer m such that for any ball B $|m|$ -next to C there exists $\mu_B \in \Gamma_K$ such that for $x, y \in B$ we have

$$\mu_B |m| \cdot |x - y| \leq |f(x) - f(y)| \leq \mu_B \left| \frac{1}{m} \right| \cdot |x - y|.$$

(T2) The set $\{y \in K \mid f^{-1}(y) \text{ is infinite}\}$ is finite.

We say that \mathcal{T} satisfies (T1,T2) if for all f , A and n as above, the two conditions (T1^{mix}) and (T2) hold.

We now have several variants of 1- h -minimality in mixed characteristic. The main result of this section is the following, stating that all of the above definitions agree.

2.2.7. Theorem. *The following are equivalent, for a theory \mathcal{T} of valued fields of characteristic zero (possibly of mixed characteristic) in a language \mathcal{L} containing \mathcal{L}_{val} .*

- (1) \mathcal{T} is 1- h^{ecc} -minimal.
- (2) \mathcal{T} is 1- h^{mix} -minimal.
- (3) \mathcal{T} satisfies (T1, T2).
- (4) \mathcal{T} is 1- h^{coars} -minimal.

Therefore we will call this common notion simply 1-h-minimality.

We will prove Theorem 2.2.7 in Section 2.7. Until then, we continue distinguishing between the four notions.

2.3. Basic results under 1- h^{mix} -minimality. In short, all results of [8] up to [8, Lemma 2.8.5] hold with only minor changes. If no proof is given, then the proof is identical to the cited result from [8].

2.3.1. Lemma. *Assume that \mathcal{T} is 0- h^{mix} -minimal. The following results are true for any model K of \mathcal{T} .*

- (1) (Adding constants [8, Lemma 2.4.1]). *If $A \subset K \cup \text{RV}_{|n|}^{\text{eq}}$ then $\text{Th}_{\mathcal{L}(A)}(K)$ is 1- h^{mix} -minimal. (And similarly, if \mathcal{T} is ℓ - h^{mix} -minimal, then $\text{Th}_{\mathcal{L}(A)}(K)$ is ℓ - h^{mix} -minimal for any ℓ and any such A .)*
- (2) (Preparation is first order [8, Lemma 2.4.2]). *If X_q, C_q are \emptyset -definable families of subsets of K with q running over an \emptyset -definable Q in an arbitrary imaginary sort, and C_q is finite for all q , then the set of $(q, \lambda) \in Q \times \Gamma_K^\times$ with $\lambda \leq 1$ such that C_q λ -prepares X_q is \emptyset -definable.*
- (3) (\exists^∞ -elimination [8, Lemma 2.5.2]). *Every infinite \emptyset -definable $X \subset K$ contains an open ball.*
- (4) (Finite sets are RV-parametrized [8, Lemma 2.5.3]). *If $C_q \subset K$ is an \emptyset -definable family of finite sets, for q in some arbitrary \emptyset -definable imaginary sort Q , then there is a \emptyset -definable family of injections $f_q : C_q \rightarrow \text{RV}_{|n|}^k$ (for some $k \geq 0$ and $n \geq 1$).*
- (5) (λ -next balls as unions of fibres [8, Lemma 2.5.4]). *Let $C \subset K$ be a finite \emptyset -definable set. Then for any $\lambda \leq 1$, there is a \emptyset -definable map $f : K \rightarrow \text{RV}_{|n|}^k \times \text{RV}_{|m|\lambda}$ (for some k, n and m) such that every ball λ -next to C is a union of fibres of f . If moreover, $\lambda = |n'|$ for some integer $n' > 0$ then we can ensure that for some integer $p > 0$ any ball $|p|$ -next to C is contained in a fibre of f .*

In (1) of Lemma 2.3.1, by $\text{RV}_{|n|}^{\text{eq}}$, we mean imaginary sorts of the form $(\text{RV}_{|n|})^m / \sim$, for some m and some \emptyset -definable equivalence relations \sim .

Proof. All statements are straight forward adaptations of the corresponding proofs in [8], with the following modifications:

- (4) As in [8], we assume that $\#C_q$ does not depend on q and we set $a_q := \frac{1}{\#C_q} \sum_{x \in C_q} x$. Using this, we define the map $\hat{f}_q : C_q \rightarrow \text{RV}_{|n|}, x \mapsto \text{rv}_{|n|}(x - a_q)$. If we take n to be a multiple of $\#C_q$, this map is not constant on C_q . Then we finish as in [8], namely treating each fiber of \hat{f}_q by induction on $\#C_q$. (We take the final n to be a multiple of all cardinalities $\#C_q$ appearing during this process.)

(5) We follow [8]. For $x \in K$ let $\mu(x) = \min\{|x - c| \mid c \in C\}$ and define

$$C(x) = \{c \in C \mid |x - c| = \mu(x)\},$$

$$a(x) = \frac{1}{\#C(x)} \sum_{c \in C(x)} c.$$

The map $a : K \rightarrow K$ has finite image, so by (4) we can find an injection $\alpha : \text{im } a \rightarrow \text{RV}_{|n|}^k$. Let $m = \max\{\#C(x) \mid x \in K\}$ and define the map f as

$$f(x) := \begin{cases} (\alpha(a(x)), \text{rv}_{|m|\lambda}(x - a(x))) & \text{if } |x - a(x)| \geq \mu(x)\lambda|m| \\ (\alpha(a(x)), \text{rv}_{|m|\lambda}(0)) & \text{if } |x - a(x)| < \mu(x)\lambda|m|. \end{cases}$$

If x and x' are in the same ball λ -next to C then $C(x) = C(x')$ and so $a(x) = a(x')$. If x and x' are in the same fibre of f and we are in the first case then the fact that $\text{rv}_{|m|\lambda}(x - a(x)) = \text{rv}_{|m|\lambda}(x' - a(x'))$ implies that $\text{rv}_\lambda(x - c) = \text{rv}_\lambda(x' - c)$ for any $c \in C$. Thus any fibre of f is contained in a ball λ -next to C .

If $\lambda = |n'|$ is an integer, then one can take $p = n' \cdot m$. Then f will be constant on any ball $|p|$ -next to C . \square

The following proposition states that we can also prepare families, similarly to [8, Proposition 2.6.2]. That proposition will also be needed for 0-h^{mix}-minimality, so we formulate it more generally.

2.3.2. Proposition (Preparing families). *Assume ℓ -h^{mix}-minimality for \mathcal{T} and some $\ell \geq 0$. Let K be a model of \mathcal{T} . Let $A \subset K$ and let*

$$W \subset K \times \text{RV}_{|n|}^k \times \bigcup_{\lambda \leq 1} \text{RV}_\lambda^\ell$$

be A -definable for some integers k and $n \geq 1$. Then there exists a finite A -definable set C and a positive integer m such that for any $\lambda \leq 1$ and any ball B which is $|m|\lambda$ -next to C , the set $W_{x,\lambda} := \{(\xi, \xi') \in \text{RV}_{|n|}^k \times \text{RV}_\lambda^\ell \mid (x, \xi, \xi') \in W\}$ is independent of x as x runs over B .

This also implies the following corollaries. Denote by RV_\bullet the disjoint union of all RV_λ for $\lambda \leq 1$.

2.3.3. Corollary. *Assume 1-h^{mix}-minimality for \mathcal{T} . The following hold for any model K of \mathcal{T} .*

- (1) (*RV-unions stay finite* [8, Corollary 2.6.7]). *Let $W \subset K \times \text{RV}_\bullet^k$ be \emptyset -definable such that W_ξ is finite for any $\xi \in \text{RV}_\bullet^k$. Then the union $\bigcup_\xi W_\xi$ is also finite.*
- (2) (*Finite image in K* [8, Corollary 2.6.8]). *The image of any \emptyset -definable $f : \text{RV}_\bullet^k \rightarrow K$ is finite.*
- (3) (*Removing RV-parameters* [8, Corollary 2.6.10]). *Let C be a finite $A \cup \text{RV}_{|n|}^{\text{eq}}$ -definable set, for some $A \subset K$ and some integer $n > 0$. Then there exists a finite A -definable set C' containing C .*

To obtain (T1,T2) from 1-h^{mix}-minimality for \mathcal{T} , we follow [8, Section 2.8]. Here we have to be slightly more careful in our formulations and proofs. First, we set some notation.

2.3.4. Definition. If B is an open ball in K , and λ is in Γ_K^\times , $\lambda \leq 1$, then a λ -shrinking of B is an open ball $B' \subset B$ with

$$\text{rad}_{\text{op}} B' = \lambda \text{rad}_{\text{op}} B.$$

2.3.5. Lemma. Assume 1-h^{mix}-minimality for \mathcal{T} . Let K be a model of \mathcal{T} and let $f : K \rightarrow K$ be an \emptyset -definable function. Then the following hold.

- (1) (Basic preservation of dimension [8, Lemma 2.8.1]). The set of $y \in K$ for which $f^{-1}(y)$ is infinite, is finite.
- (2) (Piecewise constant or injective [8, Lemma 2.8.2]). There exists a finite \emptyset -definable set C and a positive integer m such that on any ball B $|m|$ -next to C , f is either constant or injective.
- (3) (Images of most balls are almost balls [8, Lemma 2.8.3]). There exists a finite \emptyset -definable set C and positive integers m, n such that for any ball B contained in a ball $|m|$ -next to C , either $f(B)$ is a singleton, or for any $y \in f(B)$, there are open balls B', B'' for which $y \in B' \subset f(B) \subset B''$ and

$$\text{rad}_{\text{op}} B' \geq |n| \text{rad}_{\text{op}} B''.$$

- (4) (Preservation of scaling factor [8, Lemma 2.8.4]). Let B be \mathcal{M}_K . Suppose that there are α, β in Γ_K^\times , $\alpha < 1$, such that for every open ball $B' \subset B$ of radius α , the image $f(B')$ is contained in an open ball of radius β . Assume moreover that there is an integer p and open balls B', B'' such that $B' \subset f(B) \subset B''$ and

$$\text{rad}_{\text{op}} B' \geq |p| \text{rad}_{\text{op}} B''.$$

Then $f(B)$ is contained in an open ball of radius at most $\frac{\beta}{|n|\alpha}$ for some positive integer n .

Proof. The proofs of the first two statements are identical to the equicharacteristic zero case.

So we wish to prove the third statement. Use (2) to obtain a finite \emptyset -definable set C and an integer m such that on any ball $|m|$ -next to C , f is either constant or injective. Let $W \subset K \times (\Gamma_K^\times)^2$ consist of those (x, λ, μ) with $\mu \leq 1$ for which either $f(B_{<\lambda}(x))$ is a singleton, or for every $y \in f(B)$ there are open balls B', B'' with $y \in B' \subset f(B_{<\lambda}(x)) \subset B''$ and

$$\text{rad}_{\text{op}} B' \geq \mu \text{rad}_{\text{op}} B''.$$

We may enlarge C and m , such that C also $|m|$ -prepares this set W . Use Proposition 2.3.2 to obtain a finite \emptyset -definable D and a positive integer n such that for any ball B $\lambda|n|$ -next to C , the set $f(B)$ is $\lambda|n|$ -prepared by D . This can be done by appealing to Lemma 2.3.1 to obtain a \emptyset -definable family, parametrized over $\text{RV}_{|p|}^k \times \text{RV}_{\lambda|m'|}$, of sets such that any ball $\lambda|m|$ -next to C is a union of elements of the family. For $\lambda \in \Gamma_K^\times$, $\lambda \leq 1$, use Lemma 2.3.1 to get a map $g_\lambda : K \rightarrow \text{RV}_{|p|}^k \times \text{RV}_{\lambda|n''|}$ such

that any fibre is contained in a ball $\lambda|n|$ -next to D . We may assume that g_λ is an \emptyset -definable family of maps, with a parameter λ . Let C' be a finite \emptyset -definable set which $\lambda|n'|$ -prepares every map $g_\lambda \circ f$. This can again be done using Proposition 2.3.2.

Now let B be a ball contained in a ball B_1 $|m|$ -next to C . Then B is $\lambda|m|$ -next to C , for some $\lambda \in \Gamma_K^\times$, $\lambda \leq 1$. Since C' is finite we can find some $|m'|$ -shrinking B'_1 of B_1 which is disjoint from C' . After a translation, we obtain an $|m'|$ -shrinking of B which is contained in B'_1 . Call this ball B' , it is $\lambda|mm'|$ -next to C . Since B'_1 is disjoint from C' , we have that B' is contained in a ball λ -next to C' . Hence we can find a further $|n'|$ -shrinking of B' , say B'' , which is contained in a ball $\lambda|n'|$ -next to C' . By definition of C' this means that $g_\lambda \circ f$ is constant on B'' . In other words, $f(B'')$ is contained in a ball $\lambda|n|$ -next to D . On the other hand, B'' is $\lambda|mm'n'|$ -next to C so that $f(B'')$ is $\lambda|nm'n'|$ -prepared by D . Thus for any $y \in f(B'')$ there are open balls $\tilde{B} \subset f(B'') \subset \bar{B}$ such that $y \in \tilde{B}$ and

$$\text{rad}_{\text{op}} \tilde{B} \geq |m'n'| \text{rad}_{\text{op}} \bar{B}.$$

If $B'' = B_\mu(x)$ then $(x, \mu, |m'n'|) \in W$. Hence by the choice of C , we find that for any ball B contained in the ball B_1 $|m|$ -next to C , there exists a positive integer p such that any $|p|$ -shrinking of B has the desired property (here $p = m'n'$). We now make p independent of B by using compactness. If no p would work for every such ball, we may use compactness to find a ball B for which no p works. But this contradicts the above statement. Thus any ball B contained in a ball $|mp|$ -next to C has the desired property.

The proof of the final statement is a straightforward adaptation of the cited result in equicharacteristic zero. \square

Finally, we can prove an approximate valuative Jacobian property in mixed characteristic. The lemma and its proof are similar to [8, Lemma 2.8.5] and is sharpened to an actual Jacobian property below in Corollary 3.1.3. Note that the sharpened version is obtained only using a huge detour, involving approximations by second degree Taylor polynomials. We do not see a more direct proof of this sharpened version.

2.3.6. Lemma (Approximate valuative Jacobian property). *Assume 1 - h^{mix} -minimality for \mathcal{T} . Let K be a model of \mathcal{T} and let $f : K \rightarrow K$ be an A -definable function, for some $A \subset K \cup \text{RV}_{|n|}$. Then there exists a finite A -definable set C and a positive integer m such that for every ball B $|m|$ -next to C there exists a $\mu_B \in \Gamma_K$ such that for all $x, y \in B$ we have*

$$\mu_B |m| |x - y| \leq |f(x) - f(y)| \leq \mu_B \left| \frac{1}{m} \right| |x - y|.$$

Proof. We may assume that $A = \emptyset$ by Lemma 2.3.1. Using Lemma 2.3.5, we can find a finite \emptyset -definable set C_0 and a positive integer m such that

- f is constant or injective on balls $|m|$ -next to C_0 , and

- if B is contained in a ball $|m|$ -next to C_0 then either $f(B)$ is a singleton, or there are open balls $B' \subset f(B) \subset B''$ such that the radii of B' and B'' differ by at most $|m|$.

For an open ball $B_{<\alpha}(x)$ contained in a ball $|m|$ -next to C on which f is injective, define $\mu(x, \alpha)$ to be the set of $\mu \in \Gamma_K^\times$ for which $f(B)$ is contained in an open ball of radius μ and contains an open ball of radius $|m|\mu$. Note that pointwise, $|m|\mu(x, \alpha) \leq \mu(x, \alpha)$. Also define

$$s(x, \alpha) = \{\mu/\alpha \mid \mu \in \mu(x, \alpha)\}.$$

Claim: there exists a finite \emptyset -definable set C and a positive integer p such that

- (1) Lemma 2.3.5(3) holds for f with respect to C and p ,
- (2) $\mu(x, \alpha)$ and $s(x, \alpha)$ are independent of x as x runs over a ball $|p|$ -next to C ,
- (3) if $B_{<\alpha}(x)$ and $B_\beta(y)$ are open balls contained in the same ball $|p|$ -next to C and $\alpha \leq \beta$, then

$$s(y, \beta) \leq s(x, \alpha)/|p|$$

pointwise.

To prove the claim, let $W \subset K \times (\Gamma_K^\times)^2$ consist of those (x, λ, μ) such that either f is constant on $B_{<\lambda}(x)$, or $f(B_{<\lambda}(x))$ is contained in a ball B' of radius μ , and contains a ball of radius $|m|\mu$. Let C_1 $|n|$ -prepare this set W . We can assume that $|n| \leq |m|$. We put $C = C_0 \cup C_1$. Let B be $|n|$ -next to C . If f is constant on B then there is nothing to check, so we can assume that f is injective on B . By preparation, $\mu(x, \alpha)$ and $s(x, \alpha)$ are constant for $x \in B$ and $\alpha \leq \text{rad}_{\text{op}} B$, so we write simply $\mu(\alpha)$ and $s(\alpha)$. Fix $\mu \in \mu(\alpha)$ and take $\alpha \leq \beta \leq \text{rad}_{\text{op}} B$. Then any ball of radius α inside B has image under f contained in a ball of radius at most μ . Hence, by a rescaled version of Lemma 2.3.5(4), there exists an integer n such that

$$|m|\mu(\beta) \leq \frac{\mu(\alpha) \cdot \beta}{|n| \cdot \alpha}.$$

But this means precisely that

$$s(\beta) \leq s(\alpha)/|mn|,$$

proving the claim. This will suffice to prove one part of the inequality. For the other inequality, we apply a similar reasoning to f^{-1} as follows.

Denote by Y the set of $y \in K$ for which $f^{-1}(y)$ is finite. This is a cofinite \emptyset -definable set in K , by Lemma 2.3.1 and 2.3.5. Use Lemma 2.3.1 to obtain a \emptyset -definable family of injections

$$h_y : f^{-1}(y) \rightarrow \text{RV}_{|n_0|}^k,$$

for $y \in Y$. For $\eta \in \text{RV}_{|n_0|}^k$ define

$$\begin{aligned} Y_\eta &= \{y \in Y \mid \eta \in \text{im } h_y\}, \\ g_\eta : Y_\eta &\rightarrow K : y \mapsto h_y^{-1}(\eta). \end{aligned}$$

Then we have that

$$\bigcup_{y \in Y} \{y\} \times f^{-1}(y) = \bigsqcup_{\eta \in \text{RV}_{|n_0|}^k} \text{graph}(g_\eta).$$

Now apply the claim above to every function g_η to obtain a finite η -definable set D_η and an integer p_η such that the claim holds for g_η with respect to D_η and p_η . We may moreover assume that D_η $|p_\eta|$ -prepares Y_η . By compactness, we may take $p = p_\eta$ independent of η and $(D_\eta)_\eta$ a \emptyset -definable family. By Corollary 2.3.3 the union $D = \cup_\eta D_\eta$ is a finite \emptyset -definable set.

Using Lemma 2.3.1, take a \emptyset -definable function $\chi : K \rightarrow \text{RV}_{|n_1|}^{k'}$ such that any ball $|p|$ -next to D is a union of fibres of χ , and every ball $|p'|$ -next to D is contained in a fibre of χ . Now enlarge C and m , so that C also $|m|$ -prepares the family of sets $f^{-1}(\chi^{-1}(\eta))$ for $\eta \in \text{RV}_{|n_1|}^{k'}$. We claim that then for some integer q , if $B_{<\alpha}(x)$ and $B_\beta(y)$ are open balls contained in the same ball $|q|$ -next to C and $\alpha \leq \beta$, then

$$s(y, \alpha) \leq s(x, \beta)/|q|$$

pointwise.

So suppose that we have open balls B_1, B with $\text{rad}_{\text{op}} B_1 = \alpha \leq \text{rad}_{\text{op}} B = \beta$ which are contained in the same ball $|m|$ -next to C . If f is constant on B then we are done, so assume that f is injective on B . We may assume that $B_1 \subset B$ since $\mu(x, \alpha)$ is independent of x as x runs over B . Let $B' \subset f(B) \subset B''$ be open balls whose radii differ by at most $|m|$. By definition of D , $f(B)$ is contained in a ball $|p|$ -next to D . Therefore, perhaps after shrinking B'' , we can assume that also B'' is contained in a ball $|p|$ -next to D . Let $\eta \in \text{RV}_{|n_0|}^k$ be such that $g_\eta(B'') \subset B$ and note that there is exactly one such η . Using the claim applied to g_η , take open balls $\tilde{B}' \subset g_\eta(B') \subset \overline{B}'$ such that their radii differ by at most $|p|$ and do the same with open balls $\tilde{B}'' \subset g_\eta(B'') \subset \overline{B}''$. Again by the claim applied to g_η we obtain that

$$\frac{\text{rad}_{\text{op}} \overline{B}''}{\text{rad}_{\text{op}} B''} \leq \frac{1}{|p|} \frac{\text{rad}_{\text{op}} \overline{B}'}{\text{rad}_{\text{op}} B'}.$$

Since the radii of B'' and B' differ by at most $|m|$ this implies that

$$\text{rad}_{\text{op}} \overline{B}'' \leq \frac{1}{|mp|} \text{rad}_{\text{op}} \overline{B}' \leq \frac{1}{|mp^2|} \text{rad}_{\text{op}} \tilde{B}'.$$

We use exactly the same reasoning for the smaller ball B_1 to find open balls $B'_1, B''_1, \tilde{B}'_1, \dots$ with the same properties as above. Then

$$\begin{aligned} \frac{\text{rad}_{\text{op}} B}{\text{rad}_{\text{op}} B'} &\leq \frac{\text{rad}_{\text{op}} \overline{B}''}{\text{rad}_{\text{op}} B'} \leq \frac{1}{|mp^2|} \frac{\text{rad}_{\text{op}} \overline{B}'}{\text{rad}_{\text{op}} B'} \leq \frac{1}{|mp^3|} \frac{\text{rad}_{\text{op}} \overline{B}'_1}{\text{rad}_{\text{op}} B_1} \\ &\leq \frac{1}{|mp^4|} \frac{\text{rad}_{\text{op}} \tilde{B}'_1}{\text{rad}_{\text{op}} B'_1} \leq \frac{1}{|mp^4|} \frac{\text{rad}_{\text{op}} B_1}{\text{rad}_{\text{op}} B'_1}, \end{aligned}$$

where we have applied the claim to g_η and the open balls $B'_1 \subset B'$. Therefore, we have that

$$\frac{\text{rad}_{\text{op}} B''_1}{\text{rad}_{\text{op}} B_1} \leq \frac{1}{|mp^4|} \frac{\text{rad}_{\text{op}} B''}{\text{rad}_{\text{op}} B}.$$

But this implies precisely that for $q = m^2 p^4$ we have

$$s(\alpha) \leq s(\beta)/|q|,$$

proving the claim.

Now take x, y in the same ball B $|q|$ -next to C . Denote by β the open radius of B . Let $\mu \in \mu(x, \beta)$, so that $f(B)$ is contained in a ball of radius μ and contains a ball of radius $|m|\mu$. If $\alpha > |x - y|$ then x and y are contained in an open ball of radius α . If $\mu' \in \mu(x, \alpha)$ then $f(x), f(y)$ are contained in an open ball of radius μ' . Thus

$$|f(x) - f(y)|/\alpha < \mu'/\alpha \leq \frac{1}{|q|} \mu/\beta.$$

Since this holds for any $\alpha > |x - y|$ this gives that

$$|f(x) - f(y)| \leq \frac{1}{|q|} \frac{\mu}{\beta} |x - y|.$$

For the other inequality, let $\alpha \leq |x - y|$ and denote by B' the open ball of radius α around x . By the injectivity of f on B , $f(y)$ is not in $f(B')$. But $f(B')$ contains an open ball of radius $|m|\mu'$ and so

$$|f(x) - f(y)|/\alpha \geq |m|\mu'/\alpha \geq |mq|\mu/\beta.$$

Again, since this holds for any $\alpha \leq |x - y|$ we conclude that

$$|f(x) - f(y)| \geq |mq| \frac{\mu}{\beta} |x - y|.$$

This proves the Lemma. □

We now obtain already a part of Theorem 2.2.7.

2.3.7. Corollary. *Any 1- h^{mix} -minimal theory \mathcal{T} satisfies (T1, T2).*

Proof. This is direct from Item 1 from Lemma 2.3.5 and Lemma 2.3.6. □

As a sample use of compactness, we give an adaptation of Proposition 2.3.2, where the point is that the integer m can be taken uniformly over all models K of the (possibly non-complete) theory \mathcal{T} .

2.3.8. Corollary (of Proposition 2.3.2). *Assume that \mathcal{T} is ℓ - h^{mix} -minimal for some $\ell \geq 0$, and suppose that ϕ is an \mathcal{L} -formula such that for every model $K \models \mathcal{T}$, $W_K := \phi(K)$ is a subset of $K \times \text{RV}_{|n|}^k \times \text{RV}_{\lambda_K}^\ell$ for some k, n , and some $\lambda_K \leq 1$ in Γ_K^\times . Then there exists an \mathcal{L} -formula ψ and an integer $m \geq 1$ such that for every model $K \models \mathcal{T}$, $C_K := \psi(K)$ is a finite subset of K which $\lambda_K \cdot |m|$ -prepares W_K in the following sense: For every ball $B \subset K$ which is $\lambda_K \cdot |m|$ -next to C_K , the fiber*

$$W_{K,x} := \{\xi \in \text{RV}_{|n|}^k \times \text{RV}_{\lambda_K}^\ell \mid (x, \xi) \in W_K\}$$

does not depend on x when x runs over B .

Proof. Let ϕ be given as in the statement. Whether a pair (m, ψ) works as desired in a model K can be expressed by an \mathcal{L} -sentence, by Lemma 2.3.1. By compactness and ℓ - h^{mix} -minimality, we deduce that there exist finitely many pairs (m_i, ψ_i) which cover all models. We may furthermore assume that the sets $\psi_i(K)$ are finite for each model K . We are done by letting m be the least common multiple of the m_i (so that $|m| \leq |m_i|$ for each i) and ψ be the disjunction of the ψ_i . \square

2.4. Basic results under (T1,T2).

2.4.1. **Lemma.** *Assume that \mathcal{T} satisfies (T1,T2). Let K be a model of \mathcal{T} . Then*

- (1) *(RV $_\lambda$ -unions stay finite [8, Lemma 2.9.4]). If $C_\xi \subset K$ is a definable family (with parameters) of finite sets, parametrized by $\xi \in \text{RV}_\lambda^k$, then $\bigcup_\xi C_\xi$ is still finite.*
- (2) *(Eliminating RV-parameters [8, Lemma 2.9.5]). If $f : K \rightarrow K$ is A -definable for some $A \subset K \cup \text{RV}_{|n|}$ then we can find a finite ($A \cap K$)-definable set C and an integer m such that f satisfies property (T1 $^{\text{mix}}$) with respect to C and m . In particular, \mathcal{T} is 0- h^{mix} -minimal.*

Proof. The proof goes as in [8]. \square

Item 2 of Lemma 2.4.1 implies that any theory \mathcal{T} satisfying (T1,T2) is already 0^{mix} - h -minimal. In particular, we can use Proposition 2.3.2 with $\ell = 0$ and Lemma 2.3.1.

The following is a first adaptation of [8, Lemma 2.9.6]. A more precise adaptation will be given in Corollary 3.1.3.

2.4.2. **Lemma** (Images of balls). *Assume that \mathcal{T} satisfies (T1,T2). Let $f : K \rightarrow K$ be A -definable, for some $A \subset K \cup \text{RV}_{|n|}$. Then there exists a finite A -definable C and a positive integer m such that (T1 $^{\text{mix}}$) holds for f and C and m and such that the following holds for any ball B $|m|$ -next to C . Let $B' \subset B$ be an open ball and let μ_B be as in (T1 $^{\text{mix}}$). Then for every $x \in B'$ there are open balls B_1, \tilde{B}_1 such that $f(x) \in \tilde{B}_1 \subset f(B') \subset B_1$ and moreover*

$$|m| \mu_B \text{rad}_{\text{op}}(B') \leq \text{rad}_{\text{op}}(\tilde{B}_1) \leq \text{rad}_{\text{op}}(B_1) \leq \left| \frac{1}{m} \right| \mu_B \text{rad}_{\text{op}}(B').$$

Proof. Using Lemma 2.3.1 we may as well assume that $A = \emptyset$. Take a finite \emptyset -definable set C and an integer m such that (T1 $^{\text{mix}}$) holds for f with respect to C and m . Let $\chi : K \rightarrow \text{RV}_{|p|}^k$ be a \emptyset -definable map coming from Lemma 2.3.1(5) for C, m . So every ball $|m|$ -next to C is a union of fibres of χ and every fibre of χ contains a ball $|n'|$ -next to C . Let D be a finite \emptyset -definable set $|n|$ -preparing the family $(f(\chi^{-1}(\eta)))_{\eta \in \text{RV}_{|p|}^k}$, here we use Proposition 2.3.2. Let $\psi : K \rightarrow \text{RV}_{|p'|}^{k'}$ be a \emptyset -definable map such that every ball $|n|$ -next to D is a union of fibres of ψ and every fibre of ψ contains a ball $|n'|$ -next to D . Finally, use Proposition 2.3.2 again to $|q|$ -prepare the family $(f^{-1}(\psi^{-1}(\eta)))_{\eta \in \text{RV}_{|p'|}^{k'}}$ with a finite \emptyset -definable set C_0 .

We claim that $C' = C \cup C_0$ suffices. So let B' be a ball $|q|$ -next to C' and let B be the ball $|m|$ -next to C containing B' . We can assume that $\mu_B \neq 0$, for else f is constant on B . Fix any open ball B'' in B' and let $x, y \in B''$. Then

$$|f(x) - f(y)| \leq \frac{\mu_B}{|m|} |x - y| < \frac{\mu_B}{|m|} \text{rad}_{\text{op}} B''.$$

Therefore, if we denote by B''_1 the open ball of radius $\frac{\mu_B}{|m|} \text{rad}_{\text{op}} B''$ around $f(x)$ then $f(B'')$ is contained in B''_1 . By definition of C_0 , $f(B')$ is contained in a ball B_1 $|n|$ -next to D . On the other hand, $f(B)$ is $|n|$ -prepared by D . Since $B_1 \cap f(B) \neq \emptyset$ we have

$$f(B'') \subset f(B) \subset B_1 \subset f(B).$$

Using property (T1^{mix}) we see that

$$\text{rad}_{\text{op}} B_1 \geq |m| \mu_B \text{rad}_{\text{op}} B''.$$

Now take $z \in K$ such that $|f(x) - z| < |m| \mu_B \text{rad}_{\text{op}} B''$. Then we have $z \in B_1 \subset f(B)$ and so there is some $x' \in B$ with $f(x') = z$. Applying (T1^{mix}) one more time yields that

$$|x - x'| \leq \frac{1}{|m| \mu_B} |f(x) - f(x')| < \text{rad}_{\text{op}} B''.$$

We conclude that $f(B'')$ contains the open ball of radius $|m| \mu_B \text{rad}_{\text{op}} B''$ around $f(x)$. \square

2.5. Basic results under 1-h^{ecc}-minimality. We provide the tools necessary to transfer preparation results to mixed characteristic, starting from 1-h^{ecc}-minimality. As a key technical result, we give the mixed characteristic versions of [8, Corollary 2.6.6] about preparation of families.

2.5.1. Notation. The notion of balls λ -next to a finite set $C \subset K$ now has different meanings for $|\cdot|$ and for $|\cdot|_{\text{ecc}}$, with notation from Definition 2.1.2. To make clear which of the valuations we mean, we either write $|1|$ -next or $|1|_{\text{ecc}}$ -next (instead of just 1-next).

2.5.2. Remark. Suppose that $|\cdot|_{\text{ecc}}$ is non-trivial on K . For any $x, x' \in K$, we have

$$(2.5.1) \quad |x|_{\text{ecc}} \leq |x'|_{\text{ecc}} \iff \exists m \in \mathbb{N}_{\geq 1} : |m \cdot x| \leq |x'|,$$

and given a finite set $C \subset K$, the points x and x' lie in the same ball $|1|_{\text{ecc}}$ -next to C if and only if for every integer $m \geq 1$, they lie in the same ball $|m|$ -next to C .

Assuming 1-h^{ecc}-minimality of K as an \mathcal{L}_{ecc} -structure, we will be able to find finite \mathcal{L}_{ecc} -definable sets. To get back to the smaller language \mathcal{L} , we will use the following lemma:

2.5.3. Lemma (From \mathcal{L}_{ecc} -definable to \mathcal{L} -definable). *Suppose that \mathcal{T} is 1-h^{ecc}-minimal. Let K be a model of \mathcal{T} and suppose that K is \aleph_0 -saturated and strongly \aleph_0 -homogeneous as an \mathcal{L}^{eq} -structure. (Note that this in particular implies that $|\cdot|_{\text{ecc}}$ is non-trivial.) Then, any finite \mathcal{L}_{ecc} -definable set $C \subset K$ is already \mathcal{L} -definable.*

2.5.4. *Remark.* It is a standard result that every model K has an elementary extension satisfying the properties of the lemma: Indeed, any model which is special in the sense of [19, Definition 6.1.1] is strongly \aleph_0 -homogeneous by [19, Theorem 6.1.6], and it is easy to construct \aleph_0 -saturated special models.

Proof of Lemma 2.5.3. It suffices to prove that for any $a \in C$, all realizations of $p := \text{tp}_{\mathcal{L}}(a/\emptyset)$ lie in C ; indeed, by \aleph_0 -saturation, this then implies that p is algebraic (using that C is finite), and hence isolated by some formula $\phi_p(x)$, and hence C is defined by the disjunction of finitely many such $\phi_p(x)$.

So now suppose for contradiction that there exist $a \in C$ and $a' \in K \setminus C$ which have the same \mathcal{L} -type over \emptyset . Then by our homogeneity assumption, there exists an \mathcal{L} -automorphism of K sending a to a' (and hence not fixing C setwise). Such an automorphism also preserves \mathcal{O}_{ecc} and hence is an \mathcal{L}_{ecc} -automorphism, but this contradicts C being \mathcal{L}_{ecc} -definable. \square

We obtain yet another part of Theorem 2.2.7.

2.5.5. **Corollary.** *Assume that \mathcal{T} is 1- h^{ecc} -minimal. Then \mathcal{T} is 1- h^{mix} -minimal.*

Proof. By Remark 2.5.4, we may consider a model K of \mathcal{T} which is sufficiently saturated and sufficiently homogeneous (as in Lemma 2.5.3). Consider an integer n , $\lambda \in \Gamma_K^\times$, $A \subset K$, $\xi \in \text{RV}_\lambda$, $\eta \in \text{RV}_{|n|}^k$ for some integers $k, n > 0$, and an $\mathcal{L}(A \cup \{\eta, \xi\})$ -definable set $X \subset K$. We have to show that X can be $|m| \cdot \lambda$ -prepared by some finite $\mathcal{L}(A)$ -definable set C for some integer $m > 0$.

Let λ_{ecc} be the image of λ in $\Gamma_{K, \text{ecc}}$. Since $B_{<\lambda_{\text{ecc}}}(1) \subset B_{<\lambda}(1)$, we have a canonical surjection $\text{RV}_{\lambda_{\text{ecc}}} \rightarrow \text{RV}_\lambda$. Similarly, there is a canonical surjection $\text{RV}_{\text{ecc}} \rightarrow \text{RV}_{|n|}$. We fix any preimage $(\xi_{\text{ecc}}, \eta_{\text{ecc}}) \in \text{RV}_{\lambda_{\text{ecc}}} \times \text{RV}_{\text{ecc}}^k$ of (ξ, η) , so that X is $\mathcal{L}(A \cup \{\xi_{\text{ecc}}, \eta_{\text{ecc}}\})$ -definable. By 1- h -minimality for the \mathcal{L}_{ecc} -structure on K , there exists a finite $\mathcal{L}_{\text{ecc}}(A)$ -definable set C such that for every pair x, x' in the same ball λ_{ecc} -next to C , we have $x \in X \iff x' \in X$. By Lemma 2.5.3, C is already $\mathcal{L}(A)$ -definable; we claim that it is as desired. Suppose for contradiction that there exists no m as in the corollary, i.e., for every integer $m \geq 1$, there exists a pair of points $(x, x') \in K^2$ which lie in the same ball $\lambda \cdot |m|$ -next to C such that $x \in X$ but $x' \notin X$. By \aleph_0 -saturation (in the language \mathcal{L}), we find a single pair $(x, x') \in K^2$ of points with $x \in X$ but $x' \notin X$ and which lie in the same ball $\lambda \cdot |m|$ -next to C for every $m \geq 1$. The latter implies that x and x' lie in the same ball λ_{ecc} -next to C (by Remark 2.5.2), so we get a contradiction to our choice of C . \square

2.6. **Resplendency.** In order to prove that (3) implies (1) in Theorem 2.2.7, we will need a way to add the coarsened valuation ring to the language. This is made possible via a mixed characteristic resplendency result, as in [8, Section 4].

Fix a \emptyset -definable ideal I of \mathcal{O}_K which is neither $\{0\}$ nor equal to \mathcal{O}_K . Call a language \mathcal{L}' an $\bigcup_n \text{RV}_{nI}$ -expansion of \mathcal{L} if \mathcal{L}' is obtained from \mathcal{L} by adding (any) predicates which live on Cartesian products of the (imaginary) definable sets RV_{nI} for some $n > 0$. (Recall that RV_I has been defined in 2.1.)

In the following, we assume that K is κ -saturated for some $\kappa > |\mathcal{L}|$; we call a set “small” if its cardinality is less than κ , and “large” otherwise.

2.6.1. Definition (Mixed- I -preparation). We say that K has *mixed- I -preparation* if for every integer n , every set $A \subset K$ and every $(A \cup \text{RV}_{nI})$ -definable subset $X \subset K$, there exists a finite A -definable set $C \subset K$ and an integer m such that X is mI -prepared by C .

We say that K has *resplendent mixed- I -preparation* if for every set $A \subset K$, for every $\bigcup_n \text{RV}_{nI}$ -expansion \mathcal{L}' of \mathcal{L} , and for every $\mathcal{L}'(A)$ -definable subset $X \subset K$, there exists an integer m and a finite $\mathcal{L}(A)$ -definable set $C \subset K$ such that X is mI -prepared by C .

Clearly, resplendent mixed- I -preparation implies mixed- I -preparation.

We consider the language \mathcal{L} as having the sorts K and RV_{nI} for each $n > 0$. Let moreover \mathcal{L}_I be the language with the same sorts, but consisting only of the additive group on K and the maps rv_{nI} .

2.6.2. Lemma (Preparation and partial isomorphisms). *Let $A \leq K$ be a small \mathbb{Q} -sub-vector space. The following are equivalent:*

- (i) *Any $\mathcal{L}(A \cup \bigcup_n \text{RV}_{nI})$ -definable set $X \subset K$ can be mI -prepared, for some integer m , by some finite set $C \subset A$.*
- (ii) *For every small subset $A_2 \subset K$, $c_1, c_2 \in K$ and all (potentially large) sets $B_1, B_2 \subset \bigcup_n \text{RV}_{nI}$ with $\bigcup_n \text{rv}_{nI}(\langle A, c_1 \rangle) \subset B_1$, if $f: AB_1c_1 \rightarrow A_2B_2c_2$ is a partial \mathcal{L}_I -isomorphism sending c_1 to c_2 whose restriction $f|_{AB_1}$ is a partial elementary \mathcal{L} -isomorphism, then the entire f is a partial elementary \mathcal{L} -isomorphism.*
- (iii) *For all $c_1, c_2 \in K$ and all (potentially large) sets $B \subset \text{RV}_{nI}$ containing $\bigcup_n \text{rv}_{nI}(\langle A, c_1 \rangle)$, any partial $\mathcal{L}_I(A \cup B)$ -isomorphism $f: \{c_1\} \rightarrow \{c_2\}$, is a partial elementary $\mathcal{L}(A \cup B)$ -isomorphism.*

The proof is the same as [8, Lemma 4.1.14], but let us give the proof for completeness.

Proof. (i) \Rightarrow (iii): Let f be as in (iii). We have to check that for every $\mathcal{L}(A \cup B)$ -definable set $X \subset K$, $c_1 \in X$ if and only if $c_2 \in X$. By (i), there exists a finite $C \subset A$ and an integer m such that X is mI -prepared by C . Since f is an $\mathcal{L}_I(A \cup B)$ -isomorphism and B contains $\text{rv}_{mI}(\langle A, c_1 \rangle)$, for all $a \in C$ and all $r \geq 1$, we have

$$\text{rv}_{mI}(c_2 - a) = \text{rv}_{mI}(f(c_1) - a) = f(\text{rv}_{mI}(c_1 - a)) = \text{rv}_{mI}(c_1 - a).$$

Since X is I -prepared by C , it follows that $c_1 \in X$ if and only if $c_2 \in X$.

(iii) \Rightarrow (ii): Let f be as in (ii). Since f is \mathcal{L} -elementary if and only if its restriction to every finite domain is, we may assume B_i small. Using the assumption that $f|_{AB_1}$ is \mathcal{L} -elementary, we can extend $(f|_{AB_1})^{-1}$ \mathcal{L} -elementarily to some g defined at c_2 . Let $c'_1 := g(c_2)$. Then $g \circ f: \{c_1\} \rightarrow \{c'_1\}$ is a partial $\mathcal{L}_I(A \cup B_1)$ -isomorphism. Since $\bigcup_n \text{rv}_{nI}(\langle A, c_1 \rangle) \subset B_1$, it follows by (iii) that $g \circ f$ is an elementary \mathcal{L} -isomorphism. As g is also \mathcal{L} -elementary, so is f .

(ii) \Rightarrow (i): Let X be as in (i), and let $B \subset \bigcup_n \text{RV}_{nI}$ be a finite subset such that X is $\mathcal{L}(A \cup B)$ -definable. Let \mathcal{L}'_I de note the expansion of \mathcal{L}_I by the full $\mathcal{L}(A)$ -induced structure on $\bigcup_n \text{RV}_{nI}$.

Consider any $c_1, c_2 \in K$ which have the same $\text{qf-}\mathcal{L}'_I$ -type over $A \cup B$. Then the map $f: c_1 \rightarrow c_2$ is an $\mathcal{L}'_I(A \cup B)$ -isomorphism and extends to $f: AB_1c_1 \rightarrow A_2B_2c_2$, where $B_i := B \cup \bigcup_n \text{rv}_{nI}(\langle A, c_i \rangle)$. By definition of \mathcal{L}'_I , the restriction $f|_{AB_1}$ is \mathcal{L} -elementary, so by (ii), the map f is \mathcal{L} -elementary. Since moreover f is the identity on $A \cup B$, this implies that c_1 and c_2 have the same \mathcal{L} -type over $A \cup B$.

We just proved that the $\mathcal{L}(A \cup B)$ -type of any element $c \in K$ is implied by its $\text{qf-}\mathcal{L}'_I(A \cup B)$ -type. By a classical compactness argument (cf. the proof of [19, Theorem 3.2.5]), it follows that any $\mathcal{L}(A \cup B)$ -formula in one valued field variable is equivalent to a quantifier free $\mathcal{L}'_I(A \cup B)$ -formula. In particular, this applies to our set X . Since $\mathcal{L}'_I(B)$ is an $\bigcup_n \text{RV}_{nI}$ -expansion of \mathcal{L}_I , X is indeed mI -prepared by some finite $C \subset A$. \square

We can now prove, exactly as in [8, Lemma 4.1.16]:

2.6.3. Lemma (Back and forth over $\bigcup_n \text{RV}_{nI}$). *Suppose that K has mixed- I -preparation. Then the set of partial elementary \mathcal{L} -isomorphisms $f: \bigcup_n \text{RV}_{nI} \cup A_1 \rightarrow \bigcup_n \text{RV}_{nI} \cup A_2$ (where A_1, A_2 run over all small subsets of K) has the back-and-forth, i.e., given such an f and a $c_1 \in K \setminus A_1$, f can be extended to a partial elementary \mathcal{L} -isomorphism on $\bigcup_n \text{RV}_{nI} \cup A_1 \cup \{c_1\}$.*

And finally, as in [8, Proposition 4.1.7]:

2.6.4. Proposition. *The following are equivalent:*

- (i) $\text{Th}(K)$ has mixed- I -preparation.
- (ii) $\text{Th}(K)$ has resplendent mixed- I -preparation.

We now get the analogue of [8, Theorem 4.1.19] (about $\bigcup_n \text{RV}_n$ -expansions preserving Hensel minimality), where the case with $\ell = 1$ is replaced by conditions (T1,T2). Once we will have proved Theorem 2.2.7 in full, we recover the case $\ell = 1$ as well.

- 2.6.5. Proposition.**
- (i) *If \mathcal{T} is ℓ - h^{mix} -minimal for some $\ell \in \{0, \omega\}$, then any $\bigcup_n \text{RV}_n$ -expansion also is ℓ - h^{mix} -minimal.*
 - (ii) *If \mathcal{T} satisfies (T1,T2), then any $(\bigcup_n \text{RV}_{|n|})$ -expansion of \mathcal{T} also satisfies (T1,T2).*

Proof. The proof is exactly analogous to [8, Theorem 4.1.19]: (i) follows easily from Proposition 2.6.4. The proof of (ii) also uses Proposition 2.6.4, and in addition Lemma 2.4.1, Proposition 2.3.2 in the $\ell = 0$ case (this is allowed by the remark after Lemma 2.4.1) and that finite sets can be RV -parametrized. \square

2.7. Proof of Theorem 2.2.7. We are now ready to complete the proof of Theorem 2.2.7.

Proof of Theorem 2.2.7. (1) implies (2). This is Corollary 2.5.5.

(2) implies (3). This is Corollary 2.3.7.

(3) implies (4). Let $|\cdot|_c$ be a nontrivial equicharacteristic zero coarsening of the valuation $|\cdot|$.

The coarsened valuation ring $\mathcal{O}_{K,c}$ is a pullback of some subset of RV . Hence by Item (ii) of Proposition 2.6.5, the theory of K in $\mathcal{L}' = \mathcal{L} \cup \{\mathcal{O}_{K,c}\}$ still satisfies (T1,T2). We work in this language \mathcal{L}' .

We will use [8, Theorem 2.9.1]. So let $f : K \rightarrow K$ be A -definable for some $A \subset K \cup \text{RV}_c$. By adding parameters from K to the language we may as well assume that $A \subset \text{RV}_c$. We will find a finite \emptyset -definable C which \mathfrak{m}_c -prepares f . We induct on $r = \#A$. The case $r = 0$ is clear. So suppose that $r > 0$. Let $A = A' \cup \{\zeta\}$ and consider an A' -definable family $(f_y)_y$ for $y \in K$ with $f_y = f$ if $\text{rv}_c y = \zeta$. Here we use the fact that \mathfrak{m}_c is \emptyset -definable. Let $Y = \text{rv}_c^{-1}(\zeta)$. By induction on r there exist for every y a finite y -definable C_y \mathfrak{m}_c -preparing f_y . Fix a y -definable injection $h_y : C_y \rightarrow \text{RV}_{|n_0|}^k$ for some integer n_0 . Such a map exists by Lemma 2.3.1(4) which holds under (T1,T2) by Lemma 2.4.1. By compactness we may assume that both C_y and h_y form \emptyset -definable families with a parameter y . For $\eta \in \text{RV}_{n_0}^k$ consider

$$Y_\eta = \{y \in K \mid \eta \in \text{im}(h_y)\}$$

$$g_\eta : Y_\eta \rightarrow K : y \mapsto h_y^{-1}(\eta).$$

Both of these are η -definable and via compactness we may assume that the entire family is \emptyset -definable with a parameter. Note that

$$\bigcup_{y \in K} \{y\} \times C_y = \bigsqcup_{\eta \in \text{RV}_{n_0}^k} \text{graph}(g_\eta).$$

Let D_η be a finite η -definable set $|m_0|$ -preparing Y_η and such that g_η (extended by zero) satisfies Lemma 2.4.2 with respect to D_η and m_0 . This is possible by Lemma 2.4.2. Moreover take D_η to \mathfrak{m}_c -prepare the $(A' \cup \{\eta\})$ -definable

$$\tilde{f}_\eta : K \rightarrow K : y \mapsto f_y(g_\eta(y)).$$

This is via induction on r together with the fact that we can add η to the language and preserve (T1, T2). By compactness, we have without loss that D_η forms a \emptyset -definable family, with a parameter η , and that m_0 does not depend on η . Now put $D = \{0\} \cup \bigcup_{\eta \in \text{RV}_{|n_0|}^k} D_\eta$, which is a finite \emptyset -definable set by Lemma 2.3.1.

If $D \cap Y \neq \emptyset$ then $C = \bigcup_{y \in D} C_y$ is a finite \emptyset -definable set which \mathfrak{m}_c -prepares f . So suppose that $D \cap Y = \emptyset$. Since $0 \in D$ and as Y is an rv_c -fibre, we have that Y is \mathfrak{m}_c -next to D .

Use Lemma 2.3.1 to find integers p, k' and a \emptyset -definable map

$$\chi : K \rightarrow \text{RV}_{|p|}^{k'}$$

such that any ball $|m_0|$ -next to D is a union of fibres of χ . Use Proposition 2.3.2 to find a finite \emptyset -definable set C and a positive integer n' such that C $|n'|$ -prepares every set $g_\eta(\chi^{-1}(\xi))$ for any $\eta \in \text{RV}_{|n_0|}^k$ and any $\xi \in \text{RV}_{|p|}^{k'}$ for which $\chi^{-1}(\xi) \subset Y_\eta$. Then

we have that for any ball B $|m_0|$ -next to D with $B \subset Y_\eta$ that $g_\eta(B)$ is $|n'|$ -prepared by C . We will prove that C \mathfrak{m}_c -prepares f .

We claim that C \mathfrak{m}_c -prepares $g_\eta(Y)$ for any $\eta \in \text{RV}_{n_0}^k$ with $Y \subset Y_\eta$. To this end, we will show that for any $\lambda \in \Gamma_K^\times$, $\lambda \leq 1$, if B_0 is a ball $|m_0|\lambda$ -next to D then $g_\eta(B_0)$ is $|m_0^2 n' \lambda|$ -prepared by C whenever $B \subset Y_\eta$. So let B_0 be $|m_0|\lambda$ -next to D for some $\lambda \in \Gamma_K^\times$, $\lambda \leq 1$ and let B be the ball $|m_0|$ -next to D containing B_0 . Suppose that $B \subset Y_\eta$ and let μ_B be as in (T1,T2) for g_η on B . By Lemma 2.4.2, for any $x \in B_0$ the open ball of radius $|m_0|\mu_B \text{rad}_{\text{op}}(B_0)$ around $g_\eta(x)$ is contained in $g_\eta(B)$. By the same Lemma, $g_\eta(B)$ is contained in an open ball of radius $|1/m_0|\mu_B \text{rad}_{\text{op}}(B)$. Now let B'_0 be the ball $|m_0^2 n' \lambda|$ -next to C containing $g_\eta(x)$ and let B' be the ball $|n'|$ -next to C containing B'_0 . Since $g_\eta(B)$ is $|n'|$ -prepared by C , $B' \subset g_\eta(B)$. Hence

$$\text{rad}_{\text{op}} B'_0 = |m_0^2| \lambda \text{rad}_{\text{op}} B' \leq |m_0| \lambda \mu_B \text{rad}_{\text{op}} B = |m_0| \mu_B \text{rad}_{\text{op}} B_0.$$

Therefore $B'_0 \subset g_\eta(B_0)$ and we conclude that C $|m_0^2 n' \lambda|$ -prepares $g_\eta(B)$.

We now prove the claim that C \mathfrak{m}_c -prepares $g_\eta(Y)$ for any $\eta \in \text{RV}_{n_0}^k$ with $Y \subset Y_\eta$. Fix such an η and recall that Y is \mathfrak{m}_c -next to D . Let B_0 be the ball $|m_0|$ -next to D containing Y . By property T1, g_η is either injective or constant on B_0 . If g_η is constant on B_0 there is nothing to show, so assume that g_η is injective on B_0 . Let B' be a ball \mathfrak{m}_c -next to C . If $B' \cap g_\eta(Y) = \emptyset$ then there is nothing to show. So we may assume that $g_\eta(y)$ and B' are not disjoint. Assume that the claim is false, so there exists some $z \in B' \setminus g_\eta(Y)$. Now B' is contained in a ball $|n|$ -next to C , which is in turn contained in $g_\eta(B_0)$ by construction of C . By injectivity of g_η on B_0 , there is a unique $x \in B_0$ with $g_\eta(x) = z$. We necessarily have that $x \notin Y$. Take an open ball B_1 around Y not containing x . This ball is $|m_0|\lambda$ -next to D for some $\lambda \leq 1$ in Γ_K^\times . Since Y is \mathfrak{m}_c -next to D we have that the image of $|m_0|\lambda$ is 1 under the map $\Gamma_K^\times \rightarrow \Gamma_{K,c}^\times$. By the previous paragraph, $g_\eta(B_1)$ is $|m_0^2 n' \lambda|$ -prepared by C . Now similarly, $|m_0^2 n' \lambda|$ becomes 1 in $\Gamma_{K,c}^\times$ and so $B' \subset g_\eta(B_1)$. But then $z \in g_\eta(B_1)$, contrary to assumption.

We now prove that C \mathfrak{m}_c -prepares f . Suppose the result is false, and let B be a ball \mathfrak{m}_c -next to C on which the result does not hold. So there exist distinct $x, x', x'' \in B$ and $\mu_1, \mu_2 \in \Gamma_{K,c}$ with $\mu_1 \neq \mu_2$ and

$$\begin{aligned} |f(x) - f(x')|_c &= \mu_1 |x - x'|_c \\ |f(x) - f(x'')|_c &= \mu_2 |x - x''|_c. \end{aligned}$$

Assume that $|x - x'|_c \geq |x - x''|_c$ and let B_1 be the smallest closed ball containing x, x' for $|\cdot|_c$. Fix a $y_0 \in Y$. Since C_{y_0} \mathfrak{m}_c -prepares $f_{y_0} = f$ and since $x, x', x'' \in B_1$ we have $C_{y_0} \cap B_1 \neq \emptyset$. So let $\eta \in \text{RV}^k$ be such that $g_\eta(y_0) \in B_1$. Recall that Y is a maximal ball disjoint from D for the coarsened norm. By our construction of D , g_η is defined on all of Y . We claim that since D \mathfrak{m}_c -prepares \tilde{f}_η , $g_\eta(Y)$ cannot contain all of B_1 . Indeed, suppose to the contrary that $g_\eta(y) = x, g_\eta(y') = x'$ and

$g_\eta(y'') = x''$ for some $y, y', y'' \in Y$. Then there are $\mu_{g_\eta}, \mu_{\tilde{f}_\eta} \in \Gamma_{K,c}$ such that

$$\begin{aligned} |f(x) - f(x')|_c &= \mu_1 |x - x'|_c = \mu_1 \mu_{g_\eta} |y - y'|_c \\ &= |\tilde{f}_\eta(y) - \tilde{f}_\eta(y')|_c = \mu_{\tilde{f}_\eta} |y - y'|_c. \end{aligned}$$

Hence $\mu_1 = \mu_{\tilde{f}_\eta} / \mu_{g_\eta}$. But exactly the same reasoning works for μ_2 , contrary to our assumption that μ_1 and μ_2 are distinct. Thus $g_\eta(Y)$ cannot contain all of B_1 .

Therefore

$$\text{rad}_{\text{op,ecc}}(g_\eta(Y)) \leq \text{rad}_{\text{cl,ecc}}(B_1) = |x - x'|_c.$$

Now C \mathbf{m}_c -prepares $g_\eta(Y)$ so there exists some $c \in C$ with

$$|g_\eta(y_0) - c|_c \leq \text{rad}_{\text{op,ecc}}(g_\eta(Y)) \leq |x - x'|_c.$$

So we have

$$|x - c|_c \leq \max\{|x - g_\eta(y_0)|_c, |g_\eta(y_0) - c|_c\} \leq |x - x'|_c.$$

But this would mean that x and x' are not in the same ball \mathbf{m}_c -next to C , contradiction.

(4) **implies (1)**. Trivial. □

3. GEOMETRY OF DEFINABLE SETS IN 1-H-MINIMAL STRUCTURES

In this section we develop our main geometrical results under 1-h-minimality, often using the equivalences of Theorem 2.2.7.

3.1. Taylor approximation. Results which simply state that every definable set/function of a certain kind has some nice (language-independent) properties immediately follow for valued fields of mixed characteristic via coarsenings; in particular, we have a good dimension theory, similar to [8, Section 5.3]. More generally we have the following:

3.1.1. Proposition (Language-independent properties). *Suppose that \mathcal{T} is 1-h-minimal. Then the conclusions of the following results hold for any model K of \mathcal{T} : [8, Theorem 5.1.1] (almost everywhere continuity), [8, Theorem 5.1.5] (almost everywhere C^k), [8, Proposition 5.3.4] (properties of dimension).*

Proof. We may work in a model $K \models \mathcal{T}$ such that $|\cdot|_{\text{ecc}}$ is non-trivial. Then every \mathcal{L} -definable object is in particular \mathcal{L}_{ecc} -definable, so all of the above equi-characteristic 0 results apply to the definable objects in question and yield the desired mixed-characteristic result, except in the case of [8, Item 3] of [8, Proposition 5.3.4]. (Concerning [8, Theorem 5.1.1] and [8, Theorem 5.1.5], note that $|\cdot|$ and $|\cdot|_{\text{ecc}}$ induce the same topology and hence equivalent notions of continuity and derivatives.)

Now [8, Item 3] of [8, Proposition 5.3.4] (definability of dimension) can easily be reproved directly in \mathcal{L} , using Lemma 2.3.1 and that 0-dimensional is equivalent to finite. □

Now we find back the precise analogue of [8, Theorem 3.2.2] on Taylor approximation.

3.1.2. Theorem (Taylor approximation). *Suppose that \mathcal{T} is 1-h-minimal and let K be a model of \mathcal{T} . Let $f: K \rightarrow K$ be an A -definable function, for $A \subset K \cup \text{RV}_{|n|}$, and let $r \in \mathbb{N}$ be given. Then there exists a finite $A \cap K$ -definable set C and an integer $m \geq 1$ such that for every ball B $|m|$ -next to C , f is $(r + 1)$ -fold differentiable on B , $|f^{(r+1)}|$ is constant on B , and we have:*

$$(3.1.1) \quad |f(x) - T_{f,x_0}^{\leq r}(x)| = \left| \frac{1}{(r+1)!} f^{(r+1)}(x_0) \cdot (x - x_0)^{r+1} \right|$$

for every $x_0, x \in B$.

Proof. As usual, we may simply assume that $A = \emptyset$, since we can add parameters from $K \cup \text{RV}_{|n|}$ to \mathcal{L} and preserve 1-h-minimality. We first prove a slightly weaker version of this result, where the conclusion states that

$$(3.1.2) \quad |f(x) - T_{f,x_0}^{\leq r}(x)| \leq \left| \frac{1}{m} \cdot f^{(r+1)}(x_0) \cdot (x - x_0)^{r+1} \right|.$$

Proceeding as in the proof of Corollary 2.5.5, we assume that K is sufficiently saturated and sufficiently homogeneous and we use [8, Theorem 3.2.2] and Lemma 2.5.3 to find a finite \mathcal{L} -definable set C such that for x_0, x in the same ball $|1|_{\text{ecc}}$ -next to C , we have

$$(3.1.3) \quad |f(x) - T_{f,x_0}^{\leq r}(x)|_{\text{ecc}} \leq |f^{(r+1)}(x_0) \cdot (x - x_0)^{r+1}|_{\text{ecc}}.$$

Differentiability of f away from C is clear. Now suppose that there exists no $m \geq 1$ satisfying the condition involving (3.1.2). As before, we then use \aleph_0 -saturation to find a pair $(x_0, x) \in K^2$ of points which lie in the same ball $|1|_{\text{ecc}}$ -next to C and such that (3.1.1) fails for every m . The latter means that (3.1.3) fails for x_0, x , so we have a contradiction to our choice of C .

For the weakened theorem, it remains to ensure that $|f^{(r+1)}|$ is constant on balls $|m|$ -next to C . By applying Proposition 2.3.2 to the graph of $x \mapsto \text{rv}(f^{(r+1)}(x))$, we can enlarge C so that this works for balls $|m'|$ -next to C , for some $m' \geq 1$. Now the weaker statement holds using $m \cdot m'$ (since $|m \cdot m'| \leq |m|, |m'|$).

We now explain how to get the statement with $(r + 1)!$ instead of m . Go through the proof of [8, Corollary 3.2.5] (for $r + 1$) to obtain a finite set C , together with positive integers n, n' such that for any ball B $|n|$ -next to C and any $x_0 \in B$ we have

$$|f^{(r+2)}(x_0)| \text{rad}_{\text{op}} B \leq \left| \frac{1}{n'} \right| |f^{(r+1)}(x_0)|.$$

Now at the cost of enlarging n (and making B smaller) we obtain

$$|f^{(r+2)}(x_0)| \text{rad}_{\text{op}} B \leq |m| |f^{(r+1)}(x_0)|.$$

Thus we obtain [8, Corollary 3.2.5] exactly, namely, on the ball B we have for $x, x_0 \in B$ that

$$|f(x) - T_{f,x_0}^{\leq r+1}(x)| \leq \frac{|f^{(r+1)}(x_0)(x - x_0)^{r+2}|}{|(r+1)!| \text{rad}_{\text{op}}(B)}.$$

But since $|x - x_0| < \text{rad}_{\text{op}}(B)$ we get that

$$|f(x) - T_{f,x_0}^{\leq r+1}(x)| < |f^{(r+1)}(x_0)(x - x_0)^{r+1}| / |(r+1)!|.$$

Now apply the ultrametric triangle inequality to get

$$\begin{aligned} |f(x) - T_{f,x_0}^{\leq r}(x)| &\leq \max\{|f(x) - T_{f,x_0}^{r+1}(x)|, \left| \frac{f^{(r+1)}(x_0)}{(r+1)!} (x - x_0)^{r+1} \right|\} \\ &= \left| \frac{1}{(r+1)!} \right| \cdot |f^{(r+1)}(x_0)(x - x_0)^{r+1}|. \quad \square \end{aligned}$$

Surprisingly, this implies an exact version of property (T1^{mix}) and of Lemma 2.3.6, namely, an exact valuative Jacobian property similar to [8, Corollary 3.1.6], and, we get an ever finer version with rv in Corollary 3.1.4.

3.1.3. Corollary (Valuative Jacobian property). *Let \mathcal{T} be a 1-h-minimal theory. Let K be a model of \mathcal{T} and let $f : K \rightarrow K$ be A -definable, for some $A \subset K \cup \text{RV}_{|n|}$. Then there exists a finite \emptyset -definable set C and an integer m such that for any ball B $|m|$ -next to C , f is differentiable on B , $|f'|$ is constant on B , and, for all $x, x' \in B$ one has*

$$|f(x) - f(x')| = |f'(x)| \cdot |x - x'|.$$

Moreover, we can ensure that if $B' \subset B$ is an open ball then $f(B')$ is either a singleton or an open ball of radius

$$|f'(x)| \cdot \text{rad}_{\text{op}} B'.$$

Proof. Apply Theorem 3.1.2 for $r = 0$ for the first statement. For the moreover part, follow the proof of [8, Lemma 2.9.6]. \square

The following finer variant with rv instead of the valuation is useful for motivic integration, for the change of variables formulas and for Fourier transforms.

3.1.4. Corollary (Jacobian property). *Let \mathcal{T} be a 1-h-minimal theory. Let K be a model of \mathcal{T} and let $f : K \rightarrow K$ be A -definable, for some $A \subset K \cup \text{RV}_{|n|}$. Then there exists a finite \emptyset -definable set C and an integer m such that for every $\lambda \in \Gamma_K^\times$, $\lambda \leq 1$ and every ball $|m|\lambda$ -next to C we have:*

- (1) *The derivative f' exists on B and $\text{rv}_\lambda \circ f'$ is constant on B .*
- (2) *For any $x, x' \in B$ we have $\text{rv}_\lambda \left(\frac{f(x) - f(x')}{x - x'} \right) = \text{rv}_\lambda(f')$.*
- (3) *If $B' \subset B$ is any open ball, then $f(B')$ is either a point or an open ball.*

Proof. Identical to [8, Corollary 3.2.7]. \square

3.2. Algebraic Skolem functions. We show that as in the equicharacteristic zero case, one can obtain $\text{acl} = \text{dcl}$ by an $\bigcup_n \text{RV}_n$ -expansion. Recall that a theory satisfies $\text{acl} = \text{dcl}$ if for any model K and any subset $A \subset K$ we have $\text{acl}_K(A) = \text{dcl}_K(A)$. Equivalently, this means that algebraic Skolem functions exist in any model of the theory.

3.2.1. Lemma. *Suppose that $\text{Th}(K)$ is 0-h-minimal (possibly of mixed characteristic) and that for any n and any set $A \subset \text{RV}_{|n|}$ we have $\text{acl}_{\text{RV}_{|n|}}(A) = \text{dcl}_{\text{RV}_{|n|}}(A)$. Then for any set $A \subset K$ we have $\text{acl}_K(A) = \text{dcl}_K(A)$.*

Proof. Identical to [8, Lemma 4.3.1]. □

3.2.2. Proposition. *Suppose that $\text{Th}(K)$ is 1-h-minimal of possibly mixed characteristic in a language \mathcal{L} . Then there exists a $\cup_n \text{RV}_n$ -expansion \mathcal{L}' of \mathcal{L} such that $\text{Th}_{\mathcal{L}'}(K)$ is still 1-h-minimal and for any model K' of $\text{Th}_{\mathcal{L}'}(K)$ and any subset $A \subset K'$ we have*

$$\text{acl}_{\mathcal{L}', K'}(A) = \text{dcl}_{\mathcal{L}', K'}(A).$$

Proof. Identical to [8, Proposition 4.3.3]. □

By an $\cup_n \text{RV}_n$ -expansion of the language as in Proposition 3.2.2 we can thus always obtain $\text{acl} = \text{dcl}$. The following lemma provides a tool for returning to the original language.

3.2.3. Lemma (Undoing RV-expansions). *Suppose that $\text{Th}_{\mathcal{L}}(K)$ is 0-h-minimal and that \mathcal{L}' is an $\text{RV}_{|m|}$ -expansion of \mathcal{L} , for some integer $m > 0$. Let $\chi': K^n \rightarrow \text{RV}_{|N'|}^{k'}$ be an \mathcal{L}' -definable map (for some positive integers k' and N'). Then there exists an \mathcal{L} -definable map $\chi: K^n \rightarrow \text{RV}_{|N|}^k$ (for some k and N) such that χ' factors over χ , i.e. $\chi' = g \circ \chi$, for some function $g: \text{RV}_{|N|}^k \rightarrow \text{RV}_{|N'|}^{k'}$ (which is automatically \mathcal{L}' -definable).*

Proof. The existence of χ which factors over χ' follows by induction on n , Propositions 2.3.2 and 2.6.5, and (5) of Lemma 2.3.1, similarly as in the proof for [8, Lemma 4.3.4]. □

3.3. Cell decomposition and (higher dimensional) Jacobian property. As in [8, Section 5.2], we approach the cell decomposition results in a simpler way than usual (compared to e.g. [16]), by imposing $\text{acl} = \text{dcl}$ in K , or in other words, that we have algebraic Skolem functions in K . This is harmless by Proposition 3.2.2 and Lemma 3.2.3. In fact, the assumption that acl equals dcl does more than just simplify the arguments: it also allows one to formulate stronger results like piecewise Lipschitz continuity results as in Theorem 3.3.4. This is similar to [7], where this condition is also used for similar reasons.

In the following, for $m \leq n$, we denote the projection $K^n \rightarrow K^m$ to the first m coordinates by $\pi_{\leq m}$, or also by $\pi_{< m+1}$.

3.3.1. Definition (Cells, twisted boxes). Fix any parameter set $A \subset K^{\text{eq}}$ and consider a non-empty A -definable set $X \subset K^n$ for some n , an integer N , and, for $i = 1, \dots, n$, values j_i in $\{0, 1\}$ and A -definable functions $c_i: \pi_{< i}(X) \rightarrow K$. Then X is called an A -definable cell of depth $|N|$, with center tuple $c = (c_i)_{i=1}^n$ and of cell-type $j = (j_i)_{i=1}^n$ if it is of the form

$$X = \{x \in K^n \mid (\text{rv}_{|N|}(x_i - c_i(x_{< i})))_{i=1}^n \in R\},$$

for a (automatically A -definable) set

$$R \subset \prod_{i=1}^n (j_i \cdot \text{RV}_{|N|}^\times),$$

where $x_{<i} = \pi_{<i}(x)$ and where $0 \cdot \text{RV}_{|N|}^\times = \{0\} \subset \text{RV}_{|N|}$, and $1 \cdot \text{RV}_{|N|}^\times = \text{RV}_{|N|} \setminus \{0\}$.

If X is such a cell, then, for any $r \in R$, the subset

$$\{x \in K^n \mid \text{rv}_{|N|}(x_i - c_i(x_{<i}))_{i=1}^n = r\},$$

of X is called a *twisted box* of the cell X . We also call X itself a twisted box if it is a cell consisting of a single twisted box (i.e., if R is a singleton).

As in [8, Section 5.2], we first state the simplest version of cell decomposition and then formulate stronger versions as addenda:

3.3.2. Theorem (Cell decomposition). *Suppose that acl equals dcl in $\text{Th}(K)$, and that $\text{Th}(K)$ is 1- h -minimal. Consider a \emptyset -definable set $X \subset K^n$ for some n . Then there exists a cell decomposition of X , namely, a partition of X into finitely many \emptyset -definable cells A_ℓ of some depth $N > 0$.*

1. **Addendum** (Preparation of $\text{RV}_{|M|}$ -sets). *On top of the assumptions from Theorem 3.3.2, let also a \emptyset -definable set $P \subset X \times \text{RV}_{|M|}^k$ be given for some k and $M > 0$. We consider P as the function sending $x \in X$ to the fiber $P_x := \{\xi \in \text{RV}_{|M|}^k \mid (x, \xi) \in P\}$.*

Then the cells A_ℓ from Theorem 3.3.2 can be taken such that moreover P (seen as function) is constant on each twisted box of each cell A_ℓ .

2. **Addendum** (Continuous functions). *On top of the assumptions from Theorem 3.3.2 (with Addendum 1, if desired), suppose that finitely many \emptyset -definable functions $f_j: X \rightarrow K$ are given. Then the A_ℓ can be taken such that moreover, the restriction $f_j|_{A_\ell}$ of each function f_j to each cell A_ℓ is continuous, and, that each component $c_i: \pi_{<i}(A_\ell) \rightarrow K$ of the center tuple of A_ℓ is continuous.*

In the one-dimensional case, we can prepare the domain and the image of the functions in a compatible way:

3. **Addendum** (Compatible preparation of domain and image). *Under the assumptions of Addendum 2, if $n = 1$, we may moreover impose that there exists an integer M such that for each ℓ and each j , $f_j|_{A_\ell}$ is either constant or injective, $f_j(A_\ell)$ is a \emptyset -definable cell of depth N and for every twisted box R of A_ℓ , there are twisted boxes $S \subset S'$ of $f_j(A_\ell)$ of depth NM respectively N such that*

$$S \subset f_j(R) \subset S'.$$

We recall the usual notion of Lipschitz continuity.

3.3.3. Definition (Lipschitz continuity). For a valued field K and an element λ in its value group Γ_K^\times , a function $f: X \subset K^n \rightarrow K^m$ is called Lipschitz continuous with Lipschitz constant λ if for all x and x' in X one has

$$|f(x) - f(x')| \leq \lambda|x - x'|,$$

where the norm of tuples is, as usual, the sup-norm. We call such f shortly λ -Lipschitz.

Call f locally λ -Lipschitz, if for each $x \in X$ there is an open neighborhood U of x such that the restriction of f to U is λ -Lipschitz.

4. Addendum (Lipschitz centers). *Theorem 3.3.2 (with Addenda 1 and 2 if desired), is also valid in the following variant: Instead of imposing that A_ℓ itself is a cell in the sense of Definition 3.3.1, we only impose that $\sigma_\ell(A_\ell)$ is a cell, for some coordinate permutation $\sigma_\ell: K^n \rightarrow K^n$. With this extra freedom given by the choice of σ_ℓ , we can moreover ensure that there is an integer $M > 0$ such that $\sigma_\ell(A_\ell)$ is of cell-type $(1, \dots, 1, 0, \dots, 0)$ and that each component c_i of the center tuple $(c_i)_i$ of $\sigma_\ell(A_\ell)$ is $|1/M|$ -Lipschitz.*

Closely related to that addendum, we also have the following reformulation of the piecewise continuity result of [7] in the Hensel minimal setting.

3.3.4. Theorem (Piecewise Lipschitz continuity). *Suppose that acl equals dcl in $\text{Th}(K)$ and that $\text{Th}(K)$ is 1-h-minimal. Consider a \emptyset -definable set $X \subset K^n$ for some n and a \emptyset -definable function $f: X \rightarrow K$. Suppose that f is locally 1-Lipschitz. Then there exist an integer $M > 0$ and a finite partition of X into \emptyset -definable sets A_ℓ such that the restriction of f to A_ℓ is $1/|M|$ -Lipschitz, for each ℓ .*

3.3.5. Definition (Supremum Jacobian property). Let integers $m > 0$ and $n \geq 0$ be given. For $X \subset K^n$ open and $f: X \rightarrow K$ a function, we say that f has the $|m|$ -supremum Jacobian Property on X if f is C^1 on X , and f is either constant on X , or, for every x and y in X with $x \neq y$ we have:

$$(3.3.1) \quad |(\text{grad } f)(x) - (\text{grad } f)(y)| < |m \cdot (\text{grad } f)(y)|$$

and

$$(3.3.2) \quad |f(x) - f(y) - (\text{grad } f)(y) \cdot (x - y)| < |m \cdot (\text{grad } f)(y)| \cdot |x - y|.$$

In the definition, as usual, one considers $(\text{grad } f)(y)$ as a matrix with a single row with entries $(\partial f / \partial x_i)(y)$, which is multiplied with the column vector $x - y$.

3.3.6. Theorem (Sup-Jac-preparation). *Suppose that $\text{Th}(K)$ is 1-h-minimal. Let integers $m > 0$ and $n \geq 0$ be given. For every \emptyset -definable function $f: K^n \rightarrow K$, there exists a \emptyset -definable map $\chi: K^n \rightarrow \text{RV}_{|N|}^k$ (for some integers $k \geq 0$ and $N \geq 1$) such that for each fiber F of χ which is n -dimensional, F is open and f restricted to F has the $|m|$ -supremum Jacobian property on F .*

Proof of Theorems 3.3.2, 3.3.4 and 3.3.6 and the addenda to cell decomposition. Let us first explain how compactness works for showing Theorem 3.3.6. Let f be given as in Theorem 3.3.6. We may suppose that K is saturated when we prove existence of χ . Let \mathcal{L}_{ecc} be the RV-enrichment of \mathcal{L} obtained by adding to Γ_K the cut given by the maximal ideal of $\mathcal{O}_{K, \text{ecc}}$. Then, by compactness and saturatedness, there exists an \mathcal{L}_{ecc} -definable map χ as desired by the theorem, by [8, Theorem 5.4.10]. Now, by Lemma 3.2.3, we may assume that χ is already \mathcal{L} -definable and we are done for

Theorem 3.3.6. The proofs of Theorems 3.3.2 and 3.3.4 and the addenda are direct adaptations of the proofs of [8, Theorems 5.2.4 and 5.2.8]. \square

4. DIOPHANTINE APPLICATION FOR 1-H-MINIMAL TRANSCENDENTAL CURVES

In this section we give a diophantine application of Hensel minimality by bounding the number of rational points of bounded height on transcendental curves that are definable in Hensel minimal structures, in the spirit of [4], [17], [6] and [7]. In more detail, our Diophantine application for curves resembles the results for curves from Bombieri-Pila [4], but allow for an axiomatic setting, analogous to the results by Pila-Wilkie [17] (where \mathfrak{o} -minimality is used). It extends [6] and [7] to our axiomatic setting of Hensel minimality, but only in the curve case. In [6] and [7] the analytic (and subanalytic) situation is treated. These approaches all use Taylor approximation to finite order. In the one-dimensional case, Theorem 3.1.2 is strong enough as Taylor approximation, but in higher dimensions new and extra work is needed which seems quite hard at the moment. See Section 5.5 for a discussion on the difficulties to extend the axiomatic approach to higher dimensions.

Note that the finiteness result of [3] for the number of polynomials of bounded degree on the transcendental part of non-archimedean analytic and subanalytic sets X in K^n with $K = \mathbb{C}((t))$ no longer needs to hold in general 1-h-minimal structures on K , and is specific to the subanalytic situation. See [5] for such examples with infinitely many polynomials of bounded degree on the transcendental part of a set definable in a Hensel minimal structure. We leave the discovery of higher dimensional variants of Theorem 4.1.6 to the future.

4.1. We first define sets of rational points of bounded height, as in [6], [7].

4.1.1. **Definition.** For any integer $H > 0$ and any set $X \subset K^n$ with K a field extension of \mathbb{Q}_p , write $X(H)$ for

$$X(H) := \{x \in X \cap \mathbb{Z}^n \mid 0 \leq x_i \leq H \text{ for each } i\}.$$

In the real setting, convexity together with bounds on C^r -norms leads to Taylor approximation results, but in our setting we have to control Taylor approximation in other ways (by lack of convexity and by total disconnectedness). This is captured by the T_r notion from [6] and [7].

4.1.2. **Definition** (T_r maps). Fix a positive integer r . A function $f : U \subseteq K^m \rightarrow K$ is said to be T_r if for each $y \in U$ there is a polynomial $T_{f,y}^{<r}(x)$ of degree less than r and coefficients in K such that for each $x, y \in U$

$$(4.1.1) \quad |f(x) - T_{f,y}^{<r}(x)| \leq |x - y|^r.$$

A map $f : U \subseteq K^m \rightarrow K^n$ is called T_r if each component function is.

We recall [7, Lemma 5.1.3], amending it with a correction for the exponent (See Remark 4.1.4).

4.1.3. Lemma (Counting algebraic hypersurfaces). *For all positive integers d, n, m with $m < n$, consider the integers r, V, e and μ as defined in [7, Section 5.1]. Fix a finite field extension K of degree ν over \mathbb{Q}_p and with q_K many elements in the residue field of K . Let a subset $U \subseteq \mathcal{O}_K^m$ and a T_r map $\psi : U \rightarrow \mathcal{O}_K^n$ be given and put $X = \psi(U)$. Then for each $H \geq 1$, the set $X(H)$ is contained in at most*

$$(4.1.2) \quad q_K^m (\mu!)^{m/e} H^{\nu m V/e}$$

hypersurfaces of degree at most d . Moreover, when d goes to infinity, mV/e goes to 0.

4.1.4. Remark. Note that the exponent ν of H in (4.1.2) is forgotten in the statement of [7, Lemma 5.1.3], but comes up in the proof given there. This lemma is used in Proposition 5.1.4 and Theorem 5.2.2 of [7], both of which can be corrected by letting the implicit constants depend furthermore on ν . The positive characteristic cases of Lemma 5.1.3, Proposition 5.1.4 and Theorem 5.2.2 of [7] are correct as they are in [7].

The following lemma plays a similar role as Lemma 3.2.14 of [6] and Lemma 3.4.3 of [7]. Similarly as in the Yomdin-Gromov and Pila-Wilkie approaches [20] [11] [17], it states that composing with well-chosen powers maps yields good Taylor approximation.

4.1.5. Lemma (Composing with power maps). *Let an integer $r > 0$ and a finite field extension K of \mathbb{Q}_p be given. Let B be an open ball of the form $d(1 + N\mathcal{M}_K)$ for some integer $N > 0$ and some $d \in \mathcal{O}_K$. Let $f : B \rightarrow \mathcal{O}_K$ be a T_1 function such that f and each of the iterated derivatives $f', f'', \dots, f^{(r)}$ of f up to order r has the Jacobian property, and, such that f satisfies Taylor approximation up to level r as in Theorem 3.1.2. Then there is an integer $r' \geq r$ such that the map*

$$x \mapsto f(a + bx^{r'})$$

is T_r on B' for any choice of $a, b \in \mathcal{O}_K$ and any ball $B' \subset \mathcal{O}_K$ which maps into B under the map $x \mapsto a + bx^{r'}$.

The proof of Lemma 4.1.5 is similar to the proofs of Lemma 3.4.3 of [7] and Lemma 3.2.14 of [6] where the analyticity condition is replaced by the Jacobian property for the functions $f, f', f'', \dots, f^{(r)}$. Slightly more details of the proof can be found in [6, 7].

Proof of Lemma 4.1.5. The Jacobian property for f and the $f^{(i)}$ and the fact that f is T_1 imply that $|f^{(i)}(b + bNx)| \leq |x|^{1-i}$ for all i with $0 < i \leq r$ and all x with $b + bNx \in B$. Write $f_{r', B'}$ for a composition of the form $x \in B' \mapsto f(a + bx^{r'})$ for some choice of r', a, b and B' as in the lemma. By the chain rule for differentiation, we find that $|f_{r', B'}^{(i)}(x)| \leq |r'/N^r|$ for all i with $0 < i \leq r$ and all x in B' . Since Taylor approximation up to level r as in Theorem 3.1.2 still holds for $f_{r', B'}$ since it is preserved under compositions (see Lemma 3.2.7 of [6]), we are done for suitable choice of r' . \square

Call an infinite set $C \subset K^n$ purely transcendental if every algebraic curve in K^n has finite intersection with C .

4.1.6. Theorem. *Let K be a finite field extension of \mathbb{Q}_p . Let $C \subset K^n$ be a purely transcendental set which is definable in a 1-h-minimal structure on K and which is of dimension 1 (i.e. there is a linear projection $p : K^n \rightarrow K$ such that $p(C)$ is infinite and such that p has finite fibers on C). Then for each $\varepsilon > 0$ there is a constant c such that for all $H \geq 1$ one has*

$$\#C(H) \leq cH^\varepsilon.$$

Moreover, the constant c can be chosen uniformly in a definable family of such curves in the following sense: Consider a 1-h-minimal structure on K and a definable family $\{C_y\}_{y \in Y}$ of definable sets $C_y \subset K^n$ where y runs over some definable set $Y \subset K^m$. Then for each $\varepsilon > 0$ there is a constant c such that for all $y \in Y$ and all $H \geq 1$ one has, if C_y is of dimension 1 and purely transcendental, then

$$\#C_y(H) \leq cH^\varepsilon.$$

Proof. Let K and C be given as in the beginning of the theorem and fix $\varepsilon > 0$. Write n' for $\binom{n}{2}$. Fix d (and corresponding value for r , V and e) such that $\nu V/e < \varepsilon/n'$, with notation from Lemma 4.1.3 for plane curves (namely the case of that lemma with $m = 1$ and $n = 2$). We may suppose that $\text{acl} = \text{dcl}$ by Proposition 3.2.2, and thus, we can speak more easily about cells. Clearly we may suppose that $C \subset \mathcal{O}_K^n$. By the Cell Decomposition Theorem 3.3.2 with Addendum 4 and possibly after composing with $x \rightarrow a + Mx$ for some $a \in \mathcal{O}_K$ to kill the factor $|1/M|$ from the addendum, we may suppose that we have N maps

$$\varphi_i : P_i \subset \mathcal{O}_K \rightarrow C$$

the union of whose images equals C , and which are 1-Lipschitz. Indeed, by varying over finite choices of a , we can ensure that the union of images equals C . Furthermore, after translation by the center we may suppose that each P_i is a cell with center 0, that each of the component functions of each partial derivative of φ_i up to order r has the Jacobian property, and, that each component functions of φ_i satisfies Taylor approximation up to level r , as in Theorem 3.1.2. By Lemma 4.1.5, we find an integer $r' \geq r$ such that the map $x \mapsto \varphi_i(a + bx^{r'})$ is T_r on P'_i , for any choice of $a, b \in \mathcal{O}_K$ and any cell P'_i which is mapped into P_i under $x \mapsto a + bx^{r'}$. Indeed, on any twisted box B' in P'_i this is exactly given by Lemma 4.1.5, and, obtaining Equation (4.1.1) on P'_i for x and y lying in different twisted boxes of P'_i is more easy. By varying over a, b and P'_i we can ensure that the union of the images of the φ_i still equals C .

Hence, we may suppose that the maps φ_i are already T_r themselves. Write C_i for $\varphi_i(P_i)$. Apply Lemma 4.1.3 to the sets $p_j(C_i)$ in K^2 for each choice of coordinate projection $p_j : K^n \rightarrow K^2$. This yields $N' > 0$ and for each i, j and $H \geq 1$ a collection of at most $N'H^{\varepsilon/n'}$ algebraic curves $S_{ij\ell}$ of degree d whose union contains $p_j(C_i)(H)$. Since any intersection of the form $\bigcap_j p_j^{-1}(S_{ij\ell})$ is algebraic and of dimension at most 1 (see for example the argument on transversal cylinders on page 45 of [6]), we

have for each $H \geq 1$ that the set $C_i(H)$ is contained in no more than $N^{n'} H^\varepsilon$ many algebraic curves (all of bounded complexity in terms of d and n). Each such algebraic curve has finite intersection with C_i by the pure transcendence of C_i . Since the intersections of C_i with the mentioned algebraic curves appear in a definable family of finite definable sets, and since one has uniform upper bounds on the size of finite sets in definable families in 1-h-minimal structures by (3) of Lemma 2.3.1, the theorem follows. \square

4.1.7. *Remark.* A version of Theorem 4.1.6 which works uniformly in all local fields K of large residue field characteristic (namely \mathbb{Q}_p and $\mathbb{F}_p((t))$ and their finite field extensions, for large p) can also be formulated and proved along the same lines. This extends the one-dimensional case (with $m = 1$, and with implied constant as specified in Remark 4.1.4) of Theorem 5.2.2 of [7] to the 1-h-minimal situation (instead of subanalytic).

5. SOME QUESTIONS

We end the paper with some questions for future research.

5.1. Suppose that \mathcal{L}' is an $\bigcup_n \text{RV}_n$ -expansion of \mathcal{L} . Then ℓ -h^{mix}-minimality of $\text{Th}_{\mathcal{L}}(K)$ implies ℓ -h^{mix}-minimality of $\text{Th}_{\mathcal{L}'}(K)$ for $\ell = 0, 1, \omega$ (see Proposition 2.6.5 and Theorem 2.2.7).

5.1.1. *Question.* Does the converse also hold, i.e., does ℓ -h-minimality of $\text{Th}_{\mathcal{L}'}(K)$ imply ℓ -h-minimality of $\text{Th}_{\mathcal{L}}(K)$?

5.1.2. *Remark.* Hensel minimality is not preserved by passing to reducts in general. Indeed, suppose that $\text{Th}_{\mathcal{L}}(K)$ is ω -h-minimal and that K is \aleph_0 -saturated. Fix a ball $B = B_{<\lambda}(a) \subset K$ which is strictly contained in a ball disjoint from $\text{acl}_K(\emptyset)$ (so that B cannot be prepared by a finite, \emptyset -definable $C \subset K$). Then $\text{Th}_{\mathcal{L}(a,\lambda)}(K)$ is ω -h-minimal but the reduct $\text{Th}_{\mathcal{L} \cup \{B\}}(K)$ is not even 0-h-minimal (where by “ B ”, we mean a predicate for that ball). It would be interesting to find a tameness notion which is preserved on reducts.

5.2. Suppose that $\text{Th}(K)$ is ω -h^{ecc}-minimal and that we have a definable coarsening $|\cdot|_c$ of the valuation; write K_c for K considered as a valued field with the coarsened valuation $|\cdot|_c$, and write k_c for the residue field of K_c , and, put the full induced structure on k_c . By Corollary [8, Corollary 4.2.4], resp. by Theorem 2.2.7, one has

- (1) if $\text{Th}(K)$ is ω -h^{ecc}-minimal, then so is $\text{Th}(K_c)$, and, resp., if $\text{Th}(K)$ is 1-h-minimal, then so is $\text{Th}(K_c)$.

5.2.1. *Question.* (2) Does ω -h^{ecc}-minimality of $\text{Th}(K)$ imply ω -h^{ecc}-minimality of $\text{Th}(k_c)$?

- (3) Do ω -h^{ecc}-minimality of $\text{Th}(K_c)$ and ω -h^{ecc}-minimality of $\text{Th}(k_c)$ together imply ω -h^{ecc}-minimality of K ?

And one may ask (2) and (3) also for 1-h-minimality instead of ω -h^{ecc}-minimality. One may also ask whether (1) holds under other variants of Hensel minimality.

5.3. Any C^1 -function $U \subset \mathbb{R}^n \rightarrow \mathbb{R}$ also has a strict derivative (see Definition [8, Definition 3.1.2]). This is not the case in valued fields:

5.3.1. **Example.** Define $f: K^2 \rightarrow K$ by $f(x, y) = x^2$ if $|x|^4 \leq |y|$ and $f(x, y) = x^3$ otherwise. This function is C^1 everywhere, but at 0, the strict derivative does not exist, since $\frac{f(x, x^4) - f(x, 0)}{x^4} = x^{-2} - x^{-1}$, which diverges for $x \rightarrow 0$.

In view of this example, and by our knowledge that strict C^1 is the better notion for rank one valued fields for several reasons (see e.g. [2], and where strict C^1 means that the strict derivative exists everywhere and is continuous), one may try to build a good working notion of definable strict C^1 submanifolds of K^n , assuming a suitable form of Hensel minimality. The following is a first question in this direction.

5.3.2. *Question.* Does the Implicit Function Theorem hold for definable, strict C^1 functions, say, assuming 1-h-minimality (with a natural definition of “strict C^1 ”)?

5.4. Addendum 3 of Cell Decomposition (Theorem 3.3.2) speaks about simultaneous preparation of the domain and range of functions, but only in the one-variable case.

5.4.1. *Question.* Is there a version of Addendum 3 for functions from K^n to K^m which works for n and/or m bigger than 1? What would even be the right formulation?

The following question might be related:

5.4.2. *Question.* Theorem 2.2.7 (3) provides a characterization of 1-h-minimality in terms of functions from K to K . Is there an analogous characterization of ω -h-minimality?

5.5. Theorem 3.3.6 can be seen as a order one Taylor approximations result, for functions in several variables. This viewpoint suggests that we might have the following variant of [8, Theorem 5.6.1] (which is a result on higher order Taylor approximations of functions in several variables):

5.5.1. *Question.* Given a definable function $f: K^n \rightarrow K$ in a 1-h-minimal structure and an integer $r \geq 1$, does there exist a definable map $\chi: K^n \rightarrow \text{RV}_{|N|}^k$ such that [8, Equation (5.6.1)] (or a similar kind of Taylor approximation) holds on each n -dimensional fiber of χ ?

Such a result would be strictly stronger than [8, Theorem 5.6.1], which yields Taylor approximations only on boxes disjoint from a lower-dimensional definable set C . Indeed, given χ , one can easily find a C such that every box disjoint from C is contained in a fiber of χ (namely by 1-preparing χ fiberwise using Proposition 2.3.2). On the other hand, the family of maximal boxes disjoint from C cannot, in general, be parametrized by a tuple from RV. An answer to Question 5.5.1 is important to generalize the diophantine application of Section 4 to arbitrary dimension instead of just curves.

REFERENCES

1. M. Aschenbrenner, A. Chernikov, A. Gehret, and M. Ziegler, *Distality in valued fields and related structures*, (2020), arXiv:2008.09889.
2. W. Bertram, H. Glöckner, and K.-H. Neeb, *Differential calculus over general base fields and rings*, *Expo. Math.* **22** (2004), no. 3, 213–282. MR 2069671
3. G. Binyamini, R. Cluckers, and D. Novikov, *Finiteness results and Wilkie’s conjecture for Pfaffian and Noetherian functions in the non-archimedean case*, (2020), preprint.
4. E. Bombieri and J. Pila, *The number of integral points on arcs and ovals*, *Duke Math. J.* **59** (1989), no. 2, 337–357. MR 1016893
5. V. Cantoral-Farfán, K. H. Nguyen, and F. Vermeulen, *A Pila-Wilkie theorem for Hensel minimal curves*, (2021), preprint.
6. R. Cluckers, G. Comte, and F. Loeser, *Non-archimedean Yomdin-Gromov parametrizations and points of bounded height*, *Forum of Mathematics, Pi* **3** (2015), no. e5, 60 pages, doi:10.1017/fmp.2015.4.
7. R. Cluckers, A. Forey, and F. Loeser, *Uniform Yomdin-Gromov parametrizations and points of bounded height in valued fields*, *Algebra Number Theory* **14** (2020), no. 6, 1423–1456.
8. R. Cluckers, I. Halupczok, and S. Rideau, *Hensel minimality I*, **arXiv:1909.13792** (2019).
9. R. Cluckers and F. Loeser, *b-minimality*, *J. Math. Log.* **7** (2007), no. 2, 195–227.
10. L. van den Dries, *Tame topology and o-minimal structures*, *Lecture note series*, vol. 248, Cambridge University Press, 1998.
11. M. Gromov, *Entropy, homology and semialgebraic geometry*, *Séminaire Bourbaki* **1985/1986** (1987), no. Astérisque 145–146, 225–240.
12. D. Haskell and D. Macpherson, *Cell decompositions of C-minimal structures*, *Ann. Pure Appl. Logic* **66** (1994), no. 2, 113–162.
13. ———, *A version of o-minimality for the p-adics*, *J. Symbolic Logic* **62** (1997), no. 4, 1075–1092.
14. E. Hrushovski and D. Kazhdan, *Integration in valued fields*, *Algebraic geometry and number theory*, *Progr. Math.*, vol. 253, Birkhäuser Boston, Boston, MA, 2006, pp. 261–405.
15. D. Macpherson and C. Steinhorn, *On variants of o-minimality*, *Annals of Pure and Applied Logic* **79** (1996), no. 2, 165–209.
16. J. Pas, *Uniform p-adic cell decomposition and local zeta functions*, *Journal für die reine und angewandte Mathematik* **399** (1989), 137–172.
17. J. Pila and A. J. Wilkie, *The rational points of a definable set*, *Duke Math. J.* **133** (2006), no. 3, 591–616.
18. A. Pillay and C. Steinhorn, *Definable sets in ordered structures. I*, *Trans. Amer. Math. Soc.* **295** (1986), no. 2, 565–592. MR 833697
19. K. Tent and M. Ziegler, *A course in model theory*, *Lecture Notes in Logic*, vol. 40, Association for Symbolic Logic, La Jolla, CA; Cambridge University Press, Cambridge, 2012. MR 2908005
20. Y. Yomdin, *C^k -resolution of semialgebraic mappings. Addendum to: “Volume growth and entropy”*, *Israel J. Math.* **57** (1987), no. 3, 301–317. MR 889980

UNIV. LILLE, CNRS, UMR 8524 - LABORATOIRE PAUL PAINLEVÉ, F-59000 LILLE, FRANCE,
AND, KU LEUVEN, DEPARTMENT OF MATHEMATICS, B-3001 LEUVEN, BELGIUM

Email address: Raf.Cluckers@univ-lille.fr

URL: <http://rcluckers.perso.math.cnrs.fr/>

LEHRSTUHL FÜR ALGEBRA UND ZAHLENTHEORIE, MATHEMATISCHES INSTITUT, UNIVER-
SITÄTSSTR. 1, 40225 DÜSSELDORF, GERMANY

Email address: math@karimmi.de

URL: <http://www.immi.karimmi.de/en/>

CNRS, UNIVERSITÉ PARIS DIDEROT, SORBONNE UNIVERSITÉ, INSTITUT DE MATHÉMA-
TIQUES DE JUSSIEU - PARIS RIVE GAUCHE (UMR 7586), 8 PLACE AURÉLIE NEMOURS, BOITE
COURRIER 7012, 75205 PARIS CEDEX 13, FRANCE

Email address: silvain.rideau@imj-prg.fr

URL: <https://webusers.imj-prg.fr/~silvain.rideau/en/>

KU LEUVEN, DEPARTMENT OF MATHEMATICS, B-3001 LEUVEN, BELGIUM

Email address: floris.vermeulen@kuleuven.be

URL: <https://sites.google.com/view/floris-vermeulen/homepage>