

Lacon-, Shrub- and Parity-Decompositions: Characterizing Transductions of Bounded Expansion Classes

Jan Dreier
 TU Wien
 dreier@ac.tuwien.ac.at

January 28, 2022

Abstract

The concept of bounded expansion provides a robust way to capture sparse graph classes with interesting algorithmic properties. Most notably, every problem definable in first-order logic can be solved in linear time on bounded expansion graph classes. First-order interpretations and transductions of sparse graph classes lead to more general, dense graph classes that seem to inherit many of the nice algorithmic properties of their sparse counterparts.

In this paper, we show that one can encode graphs from a class with structurally bounded expansion via lacon-, shrub- and parity-decompositions from a class with bounded expansion. These decompositions are useful for lifting properties from sparse to structurally sparse graph classes.

1 Introduction

Many hard graph problems become easier if we assume the input to be *well-structured*. Usually, this is enforced by assuming that the input belongs to a certain kind of (infinite) graph class. This can for example be the class of all planar graphs or the class of all graphs with maximal degree at most d for some number d . Many of the established tractable graph classes are *sparse* in the sense that their graphs have relatively few edges compared to their number of possible edges.

Algorithmic meta-theorems state that whole families of problems become more tractable if we restrict the input to certain graph classes. The best known example is Courcelle’s theorem [3], which states that every problem definable in monadic second-order logic can be solved in linear time on bounded treewidth classes. It has been shown in a series of papers that all problems definable in first-order logic can be solved in linear time on bounded degree [31], excluded minor [9], locally bounded treewidth [10] and more general sparse graph classes [8, 18]. These results are obtained by solving the *first-order model-checking problem* on those graph classes. The input to this problem is a first-order sentence φ and a graph G and the task is to decide whether φ is satisfied on G or not. This problem can trivially be solved in time $O(|G|^{|\varphi|})$, which is optimal




Figure 1: Hierarchy of selected sparse graph classes and transductions thereof. The complexity of the first-order model-checking problem encoded using colors. White means that there is an fpt algorithm. Gray means that there is no fpt algorithm for monotone graph classes unless $\text{FPT} = \text{AW}[*]$. Striped means that it is unknown whether there is an fpt algorithm.

under complexity theoretic assumptions [2]. A first-order model-checking algorithm is considered *efficient* on some graph class if it solves the problem in time $f(|\varphi|)|G|^{O(1)}$ for some function f , i.e., in *fpt time*.

Nešetřil and Ossona de Mendez introduce *bounded expansion* and *nowhere dense* graph classes. These are very robust notions that generalize the previously mentioned graph classes (see left side of Figure 1) and have interesting algorithmic properties (e.g., [21, 30, 32, 7, 18, 8]). Most notably, Dvořák, Král and Thomas [8] solve the first-order model-checking problem in linear fpt time on bounded expansion classes, and Grohe, Kreutzer and Siebertz [18] solve this problem in almost linear fpt time on nowhere dense graph classes. For sparse graphs, this is in a sense the best possible result of this type: If a graph class is monotone (i.e., closed under taking subgraphs) and *somewhere dense* (i.e., not nowhere dense) then the first-order model-checking problem is as hard as on all graphs [18].

We therefore have reached a natural barrier in the study of meta-theorems for sparse graphs. But there are other well-structured graph classes that do not fit into the framework of sparsity. One of the current *main goals* in this area is to push the theory to account for such *dense* (or non-monotone) classes as well. In particular, we want to find dense, but *structurally simple* graph classes on which one can solve the first-order

model-checking problem in fpt time.

An excellent tool to capture such graph classes are *first-order interpretations and transductions*. For a given first-order formula $\varphi(x, y)$, the corresponding *first-order interpretation* is a function that maps an input a graph G to the graph $I_\varphi(G)$ with vertex set $V(G)$ and edge set $\{uv \mid u, v \in V(G), u \neq v, G \models \varphi(u, v) \vee \varphi(v, u)\}$. For example with $\varphi(x, y) = \neg \text{edge}(x, y)$, the interpretation complements the input graph. With $\varphi(x, y) = \exists z \text{edge}(x, z) \wedge \text{edge}(z, y)$ it computes the square of a graph. For a given formula $\varphi(x, y)$, the *first-order interpretation of a graph class \mathcal{G}* is the graph class $\{I_\varphi(G) \mid G \in \mathcal{G}\}$. *First-order transductions* are slightly more powerful than interpretations, since they also have the ability to copy, delete and nondeterministically color vertices. As the precise definition of transductions is not important for now, we delay it until Section 2. We say a graph class \mathcal{G} has *structurally property X* if it is the transduction of a graph class with property X (see right side of Figure 1 for examples). For the properties considered in this paper, this is equivalent to saying that \mathcal{G} consists of induced subgraphs of an interpretation of a class with property X . This definition has nice closure properties in the sense that applying a transduction to a structurally property X class yields again a structurally property X class.

There have been recent efforts to lift the meta-theorems for sparse graph classes to transductions of these classes. Gajarský et al. [12] solve the first-order model-checking problem for structurally bounded degree graphs using a two-step approach:

- (1) First they provide a new structural characterization of these graphs. They define *near-uniform graph classes* and show that they are equivalent to structurally bounded degree graph classes.
- (2) Then they use this new characterization to compute for a graph G from a structurally bounded degree class a bounded degree graph G' that transduces to G via some fixed transduction.

It is then trivial to reduce the problem of evaluating a first-order formula φ on G to evaluating a modified formula φ' on the bounded degree graph G' . If the transduction from step 2) is based on the formula $\psi(x, y)$ then φ' is obtained from φ by replacing every edge relation $\text{edge}(x, y)$ with $\psi(x, y)$. Now the existing meta-theorems for sparse graphs immediately give us an fpt model-checking algorithm for structurally bounded degree graphs. A very promising approach to obtain meta-theorems for transductions of more general sparse graph classes is a similar two step process: First find new structural characterizations of the dense graph class and then use them as a tool to compute a sparse preimage, i.e., a graph from a sparse class that transduces to a given dense graph. Most notably, Oum used this approach to solve the MSO model-checking problem on bounded cliquewidth [28].

Given the existing result for structurally bounded degree graph classes [12], and an extension thereof [13], a *significant milestone* would be an fpt model-checking algorithm for *structurally bounded expansion*. Bounded expansion can be characterized in many ways, one of which is via *low treedepth covers* [25]. Analogously, Gajarský et al. show

that structurally bounded expansion can be characterized using so-called *low shrubdepth covers* and that one can solve the first-order model-checking problem if one can compute a low shrubdepth 2-cover [14]. Shrubdepth [4, 15] can be understood as a dense equivalent to treedepth. We have a large and flexible toolset to build algorithms on bounded expansion (grad, low treedepth covers, transitive-fraternal-augmentations, generalized coloring numbers, ...) [27], but low shrubdepth covers are at this point the only such tool for structurally bounded expansion. So far, they have not yet led to an fpt model-checking algorithm for these graph classes. In this paper, we add three new tools to our toolbox for structurally bounded expansion: *lacon-decompositions*, *shrub-decompositions* and *parity-decompositions*. We show that the first-order model-checking problem on structurally bounded expansion can be reduced to computing sparse lacon-, shrub- or parity-decompositions.

We should mention another important approach to obtain dense, structurally simple graph classes. Bonnet, Kim, Thomassé and Watrigant [1] introduce a graph parameter called *twinwidth* that captures many interesting non-monotone graph classes. The authors provide an algorithm that — given an appropriate twinwidth contraction sequence — solves the first-order model-checking problem in linear fpt time [1]. The notion of bounded twinwidth is orthogonal to structurally bounded expansion or structurally nowhere denseness.

1.1 Introducing Lacon-, Shrub- and Parity-Decompositions

A lacon-, shrub- or parity-decomposition of a graph G is a labeled graph G' that encodes G in a possibly much sparser way. These decompositions are not tied to any fixed sparsity measure, instead one can characterize different dense graph classes by applying different sparsity requirements to G' . In this paper, we use them to capture structurally bounded expansion, structurally bounded treewidth and structurally bounded treedepth, i.e., shrubdepth (Theorem 1, 2, 3). Due to their flexibility, it seems quite likely that they can be used to capture transductions of other sparse classes as well. Generally speaking, we show that a graph class has structural property X if and only if each graph in the graph class has a lacon-, shrub- or parity-decomposition with property X .

A decomposition of a graph G consists of the so-called *target vertices* $V(G)$ and additional *hidden vertices*. “Lacon” is an acronym for **l**argest **c**ommon **n**eighbor, since there is an order on the hidden vertices and the label of the largest common neighbor of two vertices (either “0” or “1”) determines whether they are adjacent or not. Formally, this is defined as follows. A linear order π on the vertices of a graph G is represented by an injective function $\pi: V(G) \rightarrow \mathbf{N}$.

Definition 1 (Lacon-decomposition). *A lacon-decomposition is a tuple (L, π) satisfying the following properties.*

- (1) *L is a bipartite graph with sides T, H . We say T are the target vertices and H are the hidden vertices of the decomposition.*
- (2) *Every hidden vertex is labeled with either “0” or “1”.*

- (3) π is a linear order on the vertices of L with $\pi(h) < \pi(t)$ for all $t \in T, h \in H$.
- (4) For all $t, t' \in T$ holds $N(t) \cap N(t') \neq \emptyset$. The largest vertex with respect to π in $N(t) \cap N(t')$ is called the dominant vertex between t and t' .

We say (L, π) is the lacon-decomposition of a graph G if

- (5) $V(G) = T$,
- (6) for all $t \neq t' \in T$ there is an edge between t and t' in G if and only if the dominant vertex of t and t' is labeled with “1”.

This definition leads to an illustrative process that reconstructs a graph G from its lacon-decomposition (L, π) : In the beginning, all edges in G are unspecified. Then we reveal the hidden vertices of L one by one in ascending order by π . If we encounter a hidden vertex h with label “1” we fully connect its neighborhood $N(h)$ in G . On the other hand, if h has label “0” we remove all edges between vertices from $N(h)$ in G . The edges inserted or removed by h “overwrite” the edges that were inserted or removed by a previous hidden vertex. After all hidden vertices have been revealed, the resulting graph is exactly G . See the middle Figure 2 for an example of a lacon-decomposition.

Before we use lacon-decompositions to characterize transductions of different sparse graph classes, let us also introduce shrub-decompositions. Our definition closely follows the wording used by Ganian et al. [15, 4] to define shrubdepth.

Definition 2 (Shrub-decomposition). *Let m and d be nonnegative integers. A shrub-decomposition of m colors and diameter d of a graph G is a pair (F, S) consisting of an undirected graph F and a set $S \subseteq \{1, 2, \dots, m\}^2 \times \{1, 2, \dots, d\}$ (called signature) such that*

- (1) the distance in F between any two vertices is at most d ,
- (2) the set of pendant vertices (i.e., degree-one vertices) of F is exactly the set $V(G)$ of vertices of G ,
- (3) each pendant vertex of F is assigned one of the colors $\{1, 2, \dots, m\}$, and
- (4) for any i, j, l it holds $(i, j, l) \in S$ iff $(j, i, l) \in S$ (symmetry in the colors), and for any two vertices $u, v \in V(G)$ such that u is colored with i and v is colored with j and the distance between u, v in F is l , the edge uv exists in G if and only if $(i, j, l) \in S$.

Item (4) says that the existence of a G -edge between $u, v \in V(G)$ depends only on the colors of u, v and their distance in the decomposition. See the right side of Figure 2 for an example. A shrub-decomposition is a generalization of a *tree-model* used in the definition of shrubdepth. In fact, we can define shrubdepth (i.e., structurally bounded treedepth) using shrub-decompositions.




Figure 2: Top: A graph G . Left: A lacon-decomposition of G . The hidden vertices are listed in ascending order from left to right. Middle: A shrub-decomposition of G with two colors and diameter three. Vertices from G are adjacent if they have distance two or distance three and the same color. Right: A parity-decomposition of G with target-degree two.

Definition 3 (Shrubdepth [4, 15]). *We say a tree-model of m colors and depth d is a shrub-decomposition (F, S) of m colors where F is a rooted tree and the length of every root-to-leaf path in F is exactly d . A graph class \mathcal{G} is said to have bounded shrubdepth if there exist numbers m and d such that every graph in \mathcal{G} has a tree-model of m colors and depth d .*

Instead of requiring F to be a tree, we can require F to come from a bounded expansion class to obtain a characterization of structurally bounded expansion, as we will see soon. At last, we also introduce parity-decompositions, where adjacency between vertices is encoded by the parity of the intersection of their neighborhoods in the decomposition.

Definition 4 (Parity-Decomposition). *Let d be a nonnegative integer. A parity-decomposition of target-degree d is a bipartite graph P with sides T, H , where every vertex in T has degree at most d . We say T are the target vertices and H are the hidden vertices of the decomposition. We say P is the parity-decomposition of a graph G if $V(G) = T$, and for all $t \neq t' \in T$ there is an edge between t and t' in G if and only if $|N(t) \cap N(t')|$ is odd.*

1.2 Generalized Coloring Numbers and Bounded Expansion

Before we present our results, we introduce the so-called *generalized coloring numbers*, as we use them to pose a sparsity requirement on lacon-decompositions. These numbers can be used to characterize bounded expansion, treewidth and treedepth and have numerous algorithmic applications. Let G be an undirected graph and π be an ordering on the vertices of G . We say a vertex u is r -reachable from v with respect to π if $\pi(u) \leq \pi(v)$ and there is a path of length at most r from v to u and for all vertices w on the path

either $w = u$ or $\pi(w) \geq \pi(v)$. A vertex u is *weakly r -reachable* from v with respect to π if there is a path of length at most r from v to u and $\pi(u) \leq \pi(w)$ for all vertices w on that path. If G is a directed graph, we say u is (weakly) r -reachable from v if and only if it is (weakly) r -reachable in the underlying undirected graph.

Let $\text{Reach}_r(G, \pi, v)$ be the set of vertices that are r -reachable from v with respect to π . We similarly define $\text{WReach}_r(G, \pi, v)$ be the set of weakly r -reachable vertices. We set

$$\begin{aligned} \text{col}_r(G, \pi) &= \max_{v \in V(G)} |\text{Reach}_r(G, \pi, v)|, \\ \text{wcol}_r(G, \pi) &= \max_{v \in V(G)} |\text{WReach}_r(G, \pi, v)|. \end{aligned}$$

We define $\Pi(G)$ to be the set of all linear orders on G . Finally, the r -coloring number and weak r -coloring number of a graph G is defined as

$$\begin{aligned} \text{col}_r(G) &= \min_{\pi \in \Pi(G)} \text{col}_r(G, \pi), \\ \text{wcol}_r(G) &= \min_{\pi \in \Pi(G)} \text{wcol}_r(G, \pi). \end{aligned}$$

For small r , the two flavors of generalized coloring numbers are strongly related. It holds that $\text{col}_r(G) \leq \text{wcol}_r(G) \leq \text{col}_r(G)^r$ [22]. For large r , the generalized coloring numbers converge to treewidth ($\text{tw}(G)$) and treedepth ($\text{td}(G)$), respectively [27, 17]

$$\begin{aligned} \text{col}_1(G) &\leq \dots \leq \text{col}_\infty(G) = \text{tw}(G) + 1, \\ \text{wcol}_1(G) &\leq \dots \leq \text{wcol}_\infty(G) = \text{td}(G). \end{aligned}$$

A graph H is an r -shallow minor of a graph G if it is the result of a contraction of mutually disjoint connected subgraphs with radius at most r in G . A graph class \mathcal{G} has *bounded expansion* if there exists a function $f(r)$ such that $\frac{|E(H)|}{|V(H)|} \leq f(r)$ for all $G \in \mathcal{G}$ and r -shallow minors H of G [27]. Zhu first observed that generalized coloring numbers can also be used to characterize bounded expansion [34]. In this paper, we rely heavily on the following characterization by van den Heuvel and Kierstead [33], which is slightly stronger than Zhu's [34] original characterization.

Definition 5 (Bounded Expansion [33]). *A graph class \mathcal{G} has bounded expansion if there exists a function $f(r)$ such that every graph $G \in \mathcal{G}$ has an ordering π of its vertices with $\text{col}_r(G, \pi) \leq f(r)$ for all r .*

As illustrated in Figure 1, examples of bounded expansion graph classes include planar graphs or any graph class of bounded degree.

1.3 Main Result

This paper presents characterizations of structurally bounded expansion based on lacon-, shrub- and parity-decompositions. As a side result, we also obtain characterizations of structurally bounded treewidth and treedepth. For the definition of *first-order transductions* and *structurally bounded expansion* see Section 2. The following theorem contains the main contribution of this paper.

Theorem 1. *Let \mathcal{G} be a graph class. The following statements are equivalent.*

- (1) \mathcal{G} has structurally bounded expansion, i.e., there exists a graph class \mathcal{G}' with bounded expansion and a first-order transduction τ such that $\mathcal{G} \subseteq \tau(\mathcal{G}')$.
- (2) There exists a function $f(r)$ such that every graph in \mathcal{G} has a lacon-decomposition (L, π) with $\text{col}_r(L, \pi) \leq f(r)$ for all r (this implies that L comes from a bounded expansion class).
- (3) There exist a signature S , a number d and a graph class \mathcal{G}' with bounded expansion such that every graph in \mathcal{G} has a shrub-decomposition (F, S) with one color, diameter at most d and $F \in \mathcal{G}'$.
- (4) There exist number d and a graph class \mathcal{G}' with bounded expansion such that every graph in \mathcal{G} has a parity-decomposition $P \in \mathcal{G}'$ with target-degree at most d .

Note that if we are given a graph G from a structurally bounded expansion class \mathcal{G} and compute a lacon-, shrub- or parity-decomposition satisfying the conditions listed in (2), (3) or (4) then we compute a bounded expansion graph that transduces to G (i.e., a preimage), thereby solving the first-order model-checking problem for structurally bounded expansion graph classes. By itself, however, the results of this paper have no algorithmic implications.

Equivalently, structurally bounded expansion classes can also be characterized via *low shrubdepth covers* [14]. This characterization was obtained using quantifier-elimination techniques for bounded expansion classes. It is a big open question whether quantifier-elimination techniques also exist for nowhere dense classes. This question would need to be settled first, before the approach of [14] could lead to new insights to structurally nowhere dense classes. The techniques of this paper (local separators and logical composition theorems) have a better chance to generalize to structurally nowhere dense classes and beyond. In fact, in a follow up paper [5], we build upon this approach to derive shrub-decompositions as well as low shrubdepth covers for structurally nowhere dense graph classes.

Theorem 1 can further be understood as a limit on the expressive power of transductions on bounded expansion classes. For example, it implies that transductions based on boolean combinations of local, purely existential formulas have the same expressive power as general first-order transductions: Assume a graph class \mathcal{G} was obtained from a graph class \mathcal{G}' with bounded expansion via a (possibly very complicated) transduction. We replace \mathcal{G}' with the class of shrub-decompositions of \mathcal{G} , as described in (3). Now \mathcal{G} can be expressed as a very simple transduction of \mathcal{G}' . We merely have to check the color of two vertices and their (bounded) distance. This can be done using a boolean combination of local, existential formulas. Other bounds on the expressive power of transductions on sparse graphs (orthogonal to ours) have been obtained in the literature [14, 26]. As side results, we obtain the following characterizations of structurally bounded treedepth and treewidth. We should note that structurally bounded treedepth is the same as bounded shrubdepth [15].

Theorem 2. *Let \mathcal{G} be a graph class. The following statements are equivalent.*

- (1) \mathcal{G} has structurally bounded treedepth, i.e., there exists a graph class \mathcal{G}' with bounded treedepth and a first-order transduction τ such that $\mathcal{G} \subseteq \tau(\mathcal{G}')$.
- (2) There exists a number d such that every graph in \mathcal{G} has a lacon-decomposition (L, π) with $\text{wcol}_\infty(L, \pi) \leq d$ (this implies L has treedepth at most d).
- (3) There exist a number d and a signature S such that every graph in \mathcal{G} has a shrub-decomposition (F, S) with one color, diameter at most d and treedepth at most d .
- (4) There exist number d and a graph class \mathcal{G}' with bounded treedepth such that every graph in \mathcal{G} has a parity-decomposition $P \in \mathcal{G}'$ with target-degree at most d .

Theorem 3. *Let \mathcal{G} be a graph class. The following statements are equivalent.*

- (1) \mathcal{G} has structurally bounded treewidth, i.e., there exists a graph class \mathcal{G}' with bounded treewidth and a first-order transduction τ such that $\mathcal{G} \subseteq \tau(\mathcal{G}')$.
- (2) There exists a number t such that every graph in \mathcal{G} has a lacon-decomposition (L, π) with $\text{col}_\infty(L, \pi) \leq t - 1$ (this implies L has treewidth at most t).
- (3) There exist a number t and a signature S such that every graph in \mathcal{G} has a shrub-decomposition (F, S) with one color, diameter at most t and treewidth at most t .
- (4) There exist number d and a graph class \mathcal{G}' with bounded treewidth such that every graph in \mathcal{G} has a parity-decomposition $P \in \mathcal{G}'$ with target-degree at most d .

The results of Theorem 1, 2, and 3 are consequences of a more general statement, Theorem 5, which shows that for every transduction of a graph G , we can find an equivalent lacon-decomposition whose generalized coloring numbers are not too far off from the numbers of G .

1.4 Localized Feferman–Vaught Composition Theorem

The Feferman–Vaught theorem [20] states that the validity of first-order formulas on the disjoint union or Cartesian product of two graphs is uniquely determined by the value of first-order formulas on the individual graphs. Makowsky adjusted the theorem for algorithmic use in the context of MSO model-checking [24]. It has numerous applications and in the area of meta-theorems it leads for example to an especially concise proof of Courcelle’s theorem [16].

A highly useful property of first-order logic is *locality*, for example in the form of Hanf’s [19] or Gaifman’s [11] theorem. Intuitively, these theorems state that first-order logic can only express local properties. Locality is a key ingredient in many first-order model-checking algorithms [6, 18, 23, 31].

A central building block in our proofs is a localized variant of the Feferman–Vaught theorem (also proven in [29, Lemma 15]). The essence of this result is the following:

Assume we have a graph G , a first-order formula $\varphi(x, y)$ and two vertices v, w and want to know whether $G \models \varphi(v, w)$. Further assume that we have some kind of “local separator” between v, w , i.e., a tuple of vertices \bar{u} such that all short paths between v and w pass through \bar{u} . We show that knowing whether $G \models \psi(\bar{u}, v)$, $G \models \psi(\bar{u}, w)$ for certain formulas ψ this gives us enough information to compute whether $G \models \varphi(v, w)$. The original Feferman–Vaught theorem claims this only if \bar{u} is an actual separator between v and w , i.e., *all* paths between v and w pass through \bar{u} . To formalize our result, we need to define so-called *q-types*.

Definition 6 (q-type [16]). *Let G be a labeled graph and $\bar{v} = (v_1, \dots, v_k) \in V(G)^k$. The q -type of \bar{v} in G is the set $\text{tp}_q(G, \bar{v})$ of all first-order formulas $\psi(x_1 \dots x_k)$ of rank at most q such that $G \models \psi(v_1 \dots v_k)$.*

We syntactically normalize formulas so that there are only finitely many formulas of fixed quantifier rank and with a fixed set of free variables. Therefore q -types are finite sets. For a tuple $\bar{u} = (u_1, \dots, u_k)$, we write $\{\bar{u}\}$ for the set $\{u_1, \dots, u_k\}$. We follow Grohe’s presentation of the Feferman–Vaught theorem [16].

Proposition 1 ([16, Lemma 2.3]). *Let G, H be labeled graphs and $\bar{u} \in V(G)^k$, $\bar{v} \in V(G)^l$, $\bar{w} \in V(H)^m$, such that $V(G) \cap V(H) = \{\bar{u}\}$. Then for all $q \in \mathbf{N}$, $\text{tp}_q(G \cup H, \bar{u}\bar{v}\bar{w})$ is determined by $\text{tp}_q(G, \bar{u}\bar{v})$ and $\text{tp}_q(H, \bar{u}\bar{w})$.*

In the previous statement, \bar{u} separates \bar{v} and \bar{w} by splitting $G \cup H$ into the subgraphs G and H . We extend this result using a more general notion of separation that goes as follows. The *length* of a path equals its number of edges.

Definition 7. *Let G be a graph, $r \in \mathbf{N}$, and $\bar{u}, \bar{v}_1, \dots, \bar{v}_k$ be tuples of vertices from G . We say \bar{u} r -separates \bar{v}_i and \bar{v}_j if every path of length at most r between a vertex from \bar{v}_i and a vertex from \bar{v}_j contains at least one vertex from \bar{u} . We say \bar{u} r -separates $\bar{v}_1, \dots, \bar{v}_k$ if it r -separates \bar{v}_i and \bar{v}_j for all $i \neq j$.*

Based on this notion of separation, we use the following Feferman–Vaught-inspired result.

Theorem 4. *There exists a function $f(q, l)$ such that for all labeled graphs G , every $q, l \in \mathbf{N}$, and all tuples $\bar{u}, \bar{v}_1, \dots, \bar{v}_k$ of vertices from G such that \bar{u} 4^q -separates $\bar{v}_1, \dots, \bar{v}_k$ and $|\bar{u}| + |\bar{v}_1| + \dots + |\bar{v}_k| \leq l$, the type $\text{tp}_q(G, \bar{u}\bar{v}_1 \dots \bar{v}_k)$ depends only on the types $\text{tp}_{f(q, l)}(G, \bar{u}\bar{v}_1), \dots, \text{tp}_{f(q, l)}(G, \bar{u}\bar{v}_k)$. Furthermore, $\text{tp}_q(G, \bar{u}\bar{v}_1 \dots \bar{v}_k)$ can be computed from $\text{tp}_{f(q, l)}(G, \bar{u}\bar{v}_1), \dots, \text{tp}_{f(q, l)}(G, \bar{u}\bar{v}_k)$.*

1.5 Techniques and Outline

The two cornerstones of our proofs are a localized Feferman–Vaught composition theorem [29], as well as separators derived via generalized coloring numbers. We show the following circular sequence of implications.

- (i) We start with a graph G that is a transduction of a graph G' , where G' has bounded generalized coloring numbers.
- (i) \implies (ii) Then we construct a so-called *directed lacon-decomposition* (L, π) of G whose coloring numbers are bounded as well. This is a generalization of a lacon-decomposition where we allow L to be a directed graph. This step forms the central part of the paper and here we use the localized Feferman–Vaught theorem extensively.
- (ii) \implies (iii) Next, we convert (L, π) into a normal lacon-decomposition without increasing the generalized coloring numbers too much.
- (iii) \implies (iv) We transform the lacon-decomposition into an equivalent shrub-decomposition (F, S) , also with bounded coloring numbers.
- (iii) \implies (v) We transform the lacon-decomposition into an equivalent parity-decomposition P , also with bounded coloring numbers.
- (iv,v) \implies (i) This implies that G is a transduction of F or P , and since the generalized coloring numbers of both are bounded, this brings us back to the start.

Afterwards, we use the fact that generalized coloring numbers can describe bounded expansion, bounded treewidth and bounded treedepth. Therefore, Theorem 1, 2 and 3 follow by posing different bounds on the coloring numbers.

The proof of (i) \implies (ii) is the central technical contribution of this paper. We sketch some of its ideas. We have a graph G' with an ordering π of its vertices and want to obtain a lacon-decomposition of an interpretation $I_\varphi(G')$ of G' . Consider vertices $v, w \in V(G')$ and a tuple \bar{u} consisting of the vertices in $\text{WReach}_r(G', \pi, v) \cap \text{WReach}_r(G', \pi, w)$. It is a basic property of generalized coloring numbers that every path of length at most r between v and w passes through \bar{u} , i.e., \bar{u} *r-separates* v and w . The localized decomposition theorem states that $\varphi(v, w)$ depends only on q -types of $\bar{u}v$ and $\bar{u}w$. For every possible tuple \bar{u} and every combination of q -types, we introduce a hidden vertex that we label with “1” if and only if its two q -types together imply φ to be true. We then connect the hidden vertices with the vertices of G' if their q -types match.

The implication (i) \implies (ii) is proven in Section 3 and (ii) \implies (iii) is proven in Section 4. This is then combined in Section 5 to show (i) \implies (iii). In Section 6 we prove (iii) \implies (iv) and in Section 7 we prove (iii) \implies (v). Then, in Section 8 we combine all these implications to prove our main results Theorem 1, 2 and 3. The localized Feferman–Vaught theorem has been proven before [29, Lemma 15]. We nevertheless finish the paper in Section 9 with a self sustained proof of the theorem.

2 Preliminaries

Any unusual graph notation is limited to specific sections and therefore introduced where it is used. We use first-order logic over labeled and unlabeled graphs. An unlabeled graph

is a relational structure with a binary edge relation. A labeled graph may additionally have vertex labels (or colors), represented by unary relations.

Let $\varphi(x, y)$ be a first-order formula. For a given graph G , the *first-order interpretation* of G under φ , denoted by $I_\varphi(G)$, is the undirected unlabeled graph with vertex set $V(G)$ and edge set $\{uv \mid u, v \in V(G), u \neq v, G \models \varphi(u, v) \vee \varphi(v, u)\}$.

First-order transductions extend interpretations with the ability to delete vertices, as well as copy and color the input graph. We use the same notation as [12]. The most interesting building block for us are *basic transductions*. A basic transduction is a triple $\tau_0 = (\chi, \nu, \varphi)$ of first-order formulas of arity zero, one and two. If $G \not\models \chi$, then $\tau_0(G)$ is undefined. Otherwise $\tau_0(G)$ is the graph with vertex set $\{v \mid v \in V(G), G \models \nu(v)\}$ and edge set $\{uv \mid u, v \in V(G), G \models \varphi(u, v) \vee \varphi(v, u)\}$. There are two more building blocks, which are less important for this work. A *p-parameter expansion* is an operation that maps each graph G to the set of all graphs that can be created by adding p unary predicates (i.e., colors or labels) to G . An *m-copy operation* maps a graph G to a graph G^m with $V(G^m) = V(G) \times \{1, \dots, m\}$ and $E(G^m) = \{(v, i)(w, i) \mid vw \in E(G), 1 \leq i \leq m\}$. Thus G^m consists of m disjoint copies of G . Furthermore, G^m has a binary relation \sim with $(u, i) \sim (v, j)$ iff $u = v$ and unary relations Q_1, \dots, Q_m with $Q_i = \{(v, i) \mid v \in V(G)\}$. Finally, a *transduction* τ is an operation of the form $\tau = \tau_0 \circ \gamma \circ \varepsilon$ where ε is a p -parameter expansion, γ is a m -copy operation and τ_0 is a basic transduction. Notice that the output $\tau(G)$ is a *set of graphs* because of the parameter expansion. For a class \mathcal{G} , we define $\tau(\mathcal{G}) = \bigcup_{G \in \mathcal{G}} \tau(G)$ to be the *transduction of \mathcal{G}* . This now leads us the definition of structurally bounded expansion and related graph classes.

Definition 8 (Structurally bounded expansion [14]). *A graph class \mathcal{G} has structurally bounded expansion if there exists a graph class \mathcal{G}' with bounded expansion and a transduction τ such that $\mathcal{G} \subseteq \tau(\mathcal{G}')$. Generally speaking, \mathcal{G} has structurally property X if \mathcal{G}' has property X .*

3 Interpretations of Bounded Expansion Have Directed Lacon-Decompositions

This section contains the central idea of this paper. We consider a generalization of lacon-decompositions called *directed lacon-decompositions* and construct such a decomposition with bounded generalized coloring numbers. In the following definition we denote the in- and out-neighborhoods of a vertex h by $N^-(h)$ and $N^+(h)$, respectively.

Definition 9 (Directed Lacon-decomposition). *A directed lacon-decomposition is a tuple (L, π) satisfying the following properties.*

- (1) *L is a directed bipartite graph with sides T, H . We say T are the target vertices and H are the hidden vertices of the decomposition.*
- (2) *Every hidden vertex is labeled with either “0” or “1”.*
- (3) *π is a linear order on the vertices of L with $\pi(h) < \pi(t)$ for all $t \in T, h \in H$.*

- (4) For all $t, t' \in T$ holds $(N^-(t) \cap N^+(t')) \cup (N^+(t) \cap N^-(t')) \neq \emptyset$. The largest vertex with respect to π in $(N^-(t) \cap N^+(t')) \cup (N^+(t) \cap N^-(t'))$ is called the dominant vertex between t and t' .

We say (L, π) is the directed lacon-decomposition of an undirected graph G if

- (5) $V(G) = T$,
- (6) for all $t \neq t' \in T$ there is an edge between t and t' in G if and only if the dominant vertex of t and t' is labeled with “1”.

Lemma 1. Let $\varphi(x, y)$ be a first-order formula. There exists a function g such that for every labeled graph G and ordering σ on the vertices of G there exists a directed lacon-decomposition (L, π) of $I_\varphi(G)$ with

- $\text{col}_r(L, \pi) \leq g(|\varphi| + \text{col}_{2,4|\varphi|}(G, \sigma)) \cdot \text{col}_{4|\varphi|r}(G, \sigma)$ for all r ,
- $\text{wcol}_r(L, \pi) \leq g(|\varphi| + \text{col}_{2,4|\varphi|}(G, \sigma)) \cdot \text{wcol}_{4|\varphi|r}(G, \sigma)$ for all r .

In our upcoming proof of Lemma 1 we need the following facts about generalized coloring numbers. As they can be easily derived from their basic definition, we omit a proof.

Proposition 2. Let G be a graph with ordering σ and $r \in \mathbf{N}$.

- (1) For $v \in V(G)$ and $u, w \in \text{WReach}_r(G, \sigma, v)$ with $\sigma(w) < \sigma(u)$ holds $w \in \text{WReach}_{2r}(G, \sigma, u)$.
- (2) Let $v, w \in V(G)$ and $S = \text{WReach}_r(G, \sigma, v) \cap \text{WReach}_r(G, \sigma, w)$. Then every path of length at most r between v and w contains a vertex from S .
- (3) Let v_1, \dots, v_k be vertices with $\sigma(v_1) \leq \sigma(v_k)$ and $\min_{i=2}^k \sigma(v_i) = \sigma(v_k)$ such that for all $1 \leq i < k$ either $v_i \in \text{WReach}_r(G, \sigma, v_{i+1})$ or $v_{i+1} \in \text{WReach}_r(G, \sigma, v_i)$. Then there exists $w \in \text{Reach}_{rk}(G, \sigma, v_k)$ with $v_1 \in \text{WReach}_r(G, \sigma, w)$.
- (4) Let v_1, \dots, v_k be vertices with $\sigma(v_1) = \min_{i=1}^k \sigma(v_i)$ such that for all $1 \leq i < k$ either $v_i \in \text{WReach}_r(G, \sigma, v_{i+1})$ or $v_{i+1} \in \text{WReach}_r(G, \sigma, v_i)$. Then $v_1 \in \text{WReach}_{rk}(G, \sigma, v_k)$.
- (5) $\text{wcol}_r(G, \sigma) \leq \text{col}_r(G, \sigma)^r$.

Proof of Lemma 1. For technical reasons, it is easier to prove the result of this lemma under the additional assumption that the input graph G has an apex vertex. We do so first, and at the end of the proof we generalize the result also to graphs without an apex vertex. Our main proof is outlined as follows: We first construct a directed lacon-decomposition, then prove that the construction is correct and at last bound the coloring numbers.

Constructing a Lacon-Decomposition. Let us fix a graph G (with an apex vertex) and an ordering σ . Let $q = |\varphi|$, $l = \text{wcol}_{2,4^q}(G, \sigma) + 2$ and let $f(q, l)$ be the function from Theorem 4. We define the hidden vertices H of L to be all tuples $(u, \bar{u}, \text{type}_1, \text{type}_2)$ such that

- $u \in V(G)$,
- \bar{u} is the tuple of all vertices in $\text{WReach}_{2,4^q}(G, \sigma, u)$ ordered in ascending order by σ ,
- $\text{type}_1, \text{type}_2$ are $f(q, l)$ -types containing formulas with $|\bar{u}| + 1$ free variables.

Let us fix one such hidden vertex $(u, \bar{u}, \text{type}_1, \text{type}_2)$. Assume vertices $v_1, v_2 \in V(G)$ that are 4^q -separated by \bar{u} and have types $\text{tp}_{f(q,l)}(G, \bar{u}v_1) = \text{type}_1$ and $\text{tp}_{f(q,l)}(G, \bar{u}v_2) = \text{type}_2$. The fact whether $G \models \varphi(v_1, v_2) \vee \varphi(v_2, v_1)$ is determined by the type $\text{tp}_q(G, \bar{u}v_1v_2)$. And according to the localized Feferman–Vaught variant in Theorem 4, $\text{tp}_q(G, \bar{u}v_1v_2)$ is in turn determined by type_1 and type_2 . With this in mind, we iterate over all hidden vertices $(u, \bar{u}, \text{type}_1, \text{type}_2)$ and distinguish two cases:

- *Case 1:* For all $v_1, v_2 \in V(G)$ that are 4^q -separated by \bar{u} with $\text{tp}_{f(q,l)}(G, \bar{u}v_1) = \text{type}_1$ and $\text{tp}_{f(q,l)}(G, \bar{u}v_2) = \text{type}_2$ holds $G \models \varphi(v_1, v_2) \vee \varphi(v_2, v_1)$. In this case we give $(u, \bar{u}, \text{type}_1, \text{type}_2)$ the label “1”.
- *Case 2:* For all $v_1, v_2 \in V(G)$ that are 4^q -separated by \bar{u} with $\text{tp}_{f(q,l)}(G, \bar{u}v_1) = \text{type}_1$ and $\text{tp}_{f(q,l)}(G, \bar{u}v_2) = \text{type}_2$ holds $G \not\models \varphi(v_1, v_2) \vee \varphi(v_2, v_1)$. In this case we give $(u, \bar{u}, \text{type}_1, \text{type}_2)$ the label “0”.

Now, every hidden vertex is labeled with either “1” or “0”. We define the arc set of our directed lacon-decomposition as follows: Let $(u, \bar{u}, \text{type}_1, \text{type}_2)$ be a hidden vertex. For every v_1 with $\text{tp}_{f(q,l)}(G, \bar{u}v_1) = \text{type}_1$ such that $u \in \text{WReach}_{4^q}(G, \sigma, v_1)$ we add an arc from v_1 to $(u, \bar{u}, \text{type}_1, \text{type}_2)$. Similarly, for every v_2 with $\text{tp}_{f(q,l)}(G, \bar{u}v_2) = \text{type}_2$ such that $u \in \text{WReach}_{4^q}(G, \sigma, v_2)$ we add an arc from $(u, \bar{u}, \text{type}_1, \text{type}_2)$ to v_2 .

As the last step of our construction, we fix an ordering π on the vertices of L such that for all hidden vertices $(u, \bar{u}, \text{type}_1, \text{type}_2), (u', \bar{u}', \text{type}'_1, \text{type}'_2) \in H$ with $\sigma(u) < \sigma(u')$ holds $\pi((u, \bar{u}, \text{type}_1, \text{type}_2)) < \pi((u', \bar{u}', \text{type}'_1, \text{type}'_2))$. We further require for every $h \in H$ and every $t \in V(G)$ that $\pi(h) < \pi(t)$. Such an ordering trivially exists.

Correctness of Construction. At first, we need to show that (L, π) is a directed lacon-decomposition as defined in Definition 9. One can easily verify that L is in fact a directed bipartite graph, every hidden vertex is either labeled with “1” or “0” and that π is an ordering on the vertices of L with $\pi(h) < \pi(t)$ for all $h \in H, t \in T = V(G)$. What is left to do, is fix some vertices $v_1, v_2 \in T$ and show that the set $(N^-(v_1) \cap N^+(v_2)) \cup (N^+(v_1) \cap N^-(v_2))$ is non-empty. We define $R_1 = \text{WReach}_{4^q}(G, \sigma, v_1)$ and $R_2 = \text{WReach}_{4^q}(G, \sigma, v_2)$. Since G has an apex vertex, there is a path of length at most two between v_1 and v_2 . We can also generally assume that $4^q \geq 2$ and therefore, by (2) of Proposition 2, $R_1 \cap R_2 \neq \emptyset$. We choose some vertex $u' \in R_1 \cap R_2$ and consider the hidden vertex $h' = (u', \bar{u}', \text{tp}_{f(q,l)}(G, \bar{u}'v_1), \text{tp}_{f(q,l)}(G, \bar{u}'v_2))$ where \bar{u}' is the tuple of

all vertices in $\text{WReach}_{2.4q}(G, \sigma, u')$, ordered in ascending order by σ . We constructed L such that there is an arc from v_1 to h' and an arc from h' to v_2 . We conclude that $(N^-(v_1) \cap N^+(v_2)) \cup (N^+(v_1) \cap N^-(v_2))$ is non-empty and therefore that (L, π) is a directed lacon-decomposition.

Next, we show that (L, π) also is a directed lacon-decomposition of $I_\varphi(G)$. L was constructed such that its target vertices are $T = V(G) = V(I_\varphi(G))$. It remains to show that the dominant vertex of two arbitrary vertices $v_1, v_2 \in T$ is labeled with “1” if and only if $v_1 v_2 \in E(I_\varphi(G))$. Let h be the dominant vertex of v_1 and v_2 . By construction, h is either of the form $h = (u, \bar{u}, \text{tp}_{f(q,l)}(G, \bar{u}v_1), \text{tp}_{f(q,l)}(G, \bar{u}v_2))$ or $h = (u, \bar{u}, \text{tp}_{f(q,l)}(G, \bar{u}v_2), \text{tp}_{f(q,l)}(G, \bar{u}v_1))$. W.l.o.g. we assume it is the former form.

We show that $R_1 \cap R_2 \subseteq \{\bar{u}\}$. To this end, remember our previously defined vertex $u' \in R_1 \cap R_2$ and corresponding hidden vertex h' . If $\sigma(u') > \sigma(u)$ then this would mean that $\pi(h') > \pi(h)$, a contradiction to the fact that we chose h to be the largest vertex in $(N^-(v_1) \cap N^+(v_2)) \cup (N^+(v_1) \cap N^-(v_2))$. Therefore $\sigma(u') < \sigma(u)$. We constructed L such that $u \in \text{WReach}_{4q}(G, \sigma, v_1)$ and we chose $u' \in \text{WReach}_{4q}(G, \sigma, v_1)$. Thus by (1) of Proposition 2, $u' \in \text{WReach}_{2.4q}(G, \sigma, u) = \{\bar{u}\}$. This implies that $R_1 \cap R_2 \subseteq \{\bar{u}\}$.

Thus, (2) of Proposition 2 states that \bar{u} $4q$ -separates v_1 and v_2 . Hence, as discussed earlier, the fact whether $G \models \varphi(v_1, v_2) \vee \varphi(v_2, v_1)$ only depends on $\text{tp}_{f(q,l)}(G, \bar{u}v_1)$ and $\text{tp}_{f(q,l)}(G, \bar{u}v_2)$. We constructed L such that $h = (u, \bar{u}, \text{tp}_{f(q,l)}(G, \bar{u}v_1), \text{tp}_{f(q,l)}(G, \bar{u}v_2))$ is labeled with “1” if and only if $G \models \varphi(v_1, v_2) \vee \varphi(v_2, v_1)$, i.e., $v_1 v_2 \in E(I_\varphi(G))$. This implies that (L, π) is in fact a directed lacon-decomposition of $I_\varphi(G)$.

Bounding the Coloring Numbers. For a hidden vertex $(u, \bar{u}, \text{type}_1, \text{type}_2)$ in L , we say u is its *corresponding vertex in G* . If t is a target vertex, we say the *corresponding vertex in G* is t itself. The corresponding vertex of a vertex $x \in V(L)$ is denoted by $u(x)$.

We start by bounding the number of hidden vertices that have the same corresponding vertex in G . To do so, we need to determine the number of possible types $\text{type}_1, \text{type}_2$ a hidden vertex can have. The size of \bar{u} is bounded by $\text{wcol}_{2.4q}(G, \sigma)$. Thus, the number of $f(q, l)$ -types containing formulas with at most $|\bar{u}| + 1$ free variables is bounded by some function of q and l . This means, there exists a function g' (independent of G and σ) such that

$$\text{for all } v \in V(G) \text{ holds } |\{x \in V(L) \mid u(x) = v\}| \leq g'(q, l). \quad (1)$$

Let h be a hidden vertex and $h' \in \text{Reach}_r(L, \pi, h)$ for some r . Since all target vertices are larger than h with respect to π , we know that h' is a hidden vertex. Remember that L is bipartite. By definition of the generalized coloring numbers, there is a path $h_0 t_0 h_1 t_1 \dots t_{k-1} h_k$ such that $h_0 = h'$, $h_k = h$, $k \leq r/2$, $\pi(h_0) < \pi(h_k)$, and $\pi(h_k) = \min_{i=2}^k \pi(h_i)$. Here, the h_i are hidden vertices and the t_i are target vertices. The order π was constructed such that $\sigma(u(h_i)) > \sigma(u(h_j))$ implies $\pi(h_i) > \pi(h_j)$ for all hidden vertices h_i, h_j . Therefore, $\sigma(u(h_0)) \leq \sigma(u(h_k))$, and $\sigma(u(h_k)) = \min_{i=2}^k \sigma(u(h_i))$. We constructed L such that $u(h_i), u(h_{i+1}) \in \text{WReach}_{4q}(G, \sigma, t_i)$. Thus by (1) of Proposition 2, either $u(h_i) \in \text{WReach}_{2.4q}(G, \sigma, u(h_{i+1}))$ or $u(h_{i+1}) \in \text{WReach}_{2.4q}(G, \sigma, u(h_i))$. By (3) of

Proposition 2, there exists $w \in \text{Reach}_{4qr}(G, \sigma, u(h))$ with $u(h') \in \text{WReach}_{2.4q}(G, \sigma, w)$. In other words, $|\text{Reach}_r(L, \pi, h)|$ is bounded from above by the number of tuples (w, v, h') with $w \in \text{Reach}_{4qr}(G, \sigma, u(h))$, $v \in \text{WReach}_{2.4q}(G, \sigma, w)$ and $u(h') = v$. Using (1), we bound the number of such tuples by

$$|\text{Reach}_r(L, \pi, h)| \leq \text{col}_{4qr}(G, \sigma) \cdot l \cdot g'(q, l). \quad (2)$$

Let us now consider a target vertex t . Observe that

$$|\text{Reach}_r(L, \pi, t)| \leq 1 + \sum_{h \in N(t) \setminus \{t\}} |\text{Reach}_r(L, \pi, h)|. \quad (3)$$

Since for all $h \in N(t)$ holds $u(h) \in \text{WReach}_{4q}(G, \sigma, t)$, we can again use (1) to bound

$$|N(t)| \leq g'(q, l) \cdot l. \quad (4)$$

By (5) of Proposition 2, $l \leq \text{col}_{2.4q}(G, \sigma)^{2.4q} + 2$. Combining equations (2), (3) and (4) yields a function g with

$$\text{col}_r(L, \pi) \leq g(q + \text{col}_{2.4q}(G, \sigma)) \cdot \text{col}_{4qr}(G, \sigma) \text{ for all } r.$$

The statement of this lemma also requires a bound on the weak coloring numbers $\text{wcol}_r(L, \pi)$. This bound can be proven in almost the same way as for $\text{col}_r(L, \pi)$ and therefore we only describe the main difference. Let h be a hidden vertex and $h' \in \text{WReach}_r(L, \pi, h)$. By definition, there is a path $h_0 t_0 h_1 t_1 \dots t_{k-1} h_k$ such that $h_0 = h'$, $h_k = h$, $k \leq r/2$, $\pi(h_0) = \min_{i=1}^k \pi(h_i)$. Then by Proposition 2, Item (4), $u(h') \in \text{WReach}_{4qr}(G, \sigma, u(h))$, i.e., $|\text{WReach}_r(L, \pi, h)|$ is bounded by the number of tuples (v, h') with $v \in \text{WReach}_{4qr}(G, \sigma, u(h))$ and $u(h') = v$. The rest proceeds as for the strong coloring numbers.

No Apex Vertex. It remains to prove this result for the case that the input graph has no apex vertex. We reduce this case to the previously covered case with an apex. Let G be a graph without an apex and σ be an ordering of $V(G)$. We construct a graph G' from G by adding an additional apex vertex and let σ' be the ordering of $V(G')$ that preserves the order of σ but whose minimal element is the new apex vertex. Then $\text{col}_r(G', \sigma') \leq \text{col}_r(G, \sigma) + 1$ and $\text{wcol}_r(G', \sigma') \leq \text{wcol}_r(G, \sigma) + 1$. Assume we have a directed lacon-decomposition (L', π') of $I_\varphi(G')$. We obtain a directed lacon-decomposition (L, π) of $I_\varphi(G)$ by removing the apex vertex from (L', π') . Then $\text{col}_r(L, \pi) \leq \text{col}_r(L', \pi')$ and $\text{wcol}_r(L, \pi) \leq \text{wcol}_r(L', \pi')$. These observations reduce the no-apex case to the apex case. \square

4 Directed and Undirected Lacon-Decompositions Have Same Expressive Power

In this section, we prove the following lemma, stating that directed and normal lacon-decompositions are equally powerful.

Lemma 2. *Assume a graph G has a directed lacon-decomposition (L, π) . Then it also has an (undirected) lacon-decomposition (L', π') with*

- $\text{col}_r(L', \pi') \leq 4^{\text{col}_2(L, \pi)} \cdot \text{col}_r(L, \pi)$ for all r ,
- $\text{wcol}_r(L', \pi') \leq 4^{\text{col}_2(L, \pi)} \cdot \text{wcol}_r(L, \pi)$ for all r .

Proof. Assume we have a directed lacon-decomposition (L, π) of a graph G with target vertices T and hidden vertices H . We construct an undirected lacon-decomposition (L', π') of G with target and hidden vertices T, H' as follows: We start with $H' = \emptyset$. We process the vertices in H one by one in ascending order by π and add in each processing step various new vertices to H' . The ordering π' is thereby defined implicitly as follows (in ascending order): First come the vertices of H' in the order of insertion and then come the vertices of T in the same order as in π .

We will make sure that after a vertex $h \in H$ has been processed the following invariant holds for all pairs of vertices $v_1 \neq v_2 \in T$. If v_1, v_2 have a dominant vertex d in L such that $\pi(d) \leq \pi(h)$ then we guarantee that v_1, v_2 have a dominant vertex d' in L' and that d' is labeled with “1” if and only if d is. After every vertex in H has been processed, this guarantees that (L', π') is an undirected lacon-decomposition of G .

While processing a vertex $h \in H$, certain new vertices are inserted into H' . The newly inserted vertices are in a sense either a copy of h or copies of vertices that have already been inserted into H' . These copies may later again be copied, and so forth. This leads to exponential growth. To keep track of these copies and to show that while being exponential, this growth is nevertheless bounded, every newly inserted vertex h' will have a *memory*, denoted by $m(h')$. This memory consists of a sequence of vertices from H . If h' is derived from h , then we set $m(h') = h$. However, if h' is derived both from h and another vertex h'' that has previously been inserted into H' , then we set $m(h') = m(h'') + h$, i.e., $m(h')$ is obtained by appending h to the memory of h'' . Let us now get into the details of the construction and prove our invariant.

Construction of Lacon-Decomposition. We describe the processing steps of a vertex $h \in H$. If $N^-(h) \cup N^+(h) = \emptyset$ we do nothing. Otherwise, we add a vertex h' to H' that has the same label (“0” or “1”) as h . We further add edges to L' such that $N(h') = N^-(h) \cup N^+(h)$ (here $N(h')$ refers to the neighborhood in L' , while $N^-(h)$, $N^+(h)$ refer to the in- and out-neighborhoods in L). Since h' is derived only from h , we set $m(h') = h$.

We observe a problem. Let $v_1 \neq v_2 \in N^-(h) \setminus N^+(h)$. Currently, h' is the dominant vertex for v_1 and v_2 in L' , even though h is not dominant for these vertices in L . Thus h' connects too many target vertices and we have to “undo” the effect of h' on the edges within the set $N^-(h) \setminus N^+(h)$. The same holds for the set $N^+(h) \setminus N^-(h)$. We do so as follows. We iterate over all vertices $l \in \text{Reach}_2(L', \pi', h')$ in order of insertion into H' . If $N(l) \cap N^+(h) \setminus N^-(h) \neq \emptyset$ we add a vertex l' with $N(l') = N(l) \cap N^+(h) \setminus N^-(h)$. Similarly, if $N(l) \cap N^-(h) \setminus N^+(h) \neq \emptyset$ we add a vertex l'' with $N(l'') = N(l) \cap N^-(h) \setminus N^+(h)$. These two vertices get the same label as l and since they were derived from

both l and h , we set $m(l') = m(l'') = m(l) + h$. These new vertices l' and l'' undo the undesired effects of inserting h' , as we will prove soon. This completes the processing round of h .

By induction, we know that the invariant was satisfied in the last round for all v_1, v_2 . We show that it also holds after h is processed. Let $v_1 \neq v_2$ be two vertices with dominant vertex d in L such that $\pi(d) \leq \pi(h)$. We distinguish five cases.

- $v_1 \notin N^-(h) \cup N^+(h)$ or $v_2 \notin N^-(h) \cup N^+(h)$. In this case, neither h in (L, π) nor any of the newly inserted vertices in (L', π') is dominant for v_1, v_2 . Since the invariant for v_1, v_2 held in the previous round, it also holds in this round.
- $v_1 \in N^-(h)$ and $v_2 \in N^+(h)$. In this case, h and h' are dominant for v_1, v_2 in (L, π) and (L', π') , respectively. The vertex h' is labeled with “1” if and only if h is. Thus, the invariant is fulfilled.
- $v_1 \in N^+(h)$ and $v_2 \in N^-(h)$. As above.
- $v_1, v_2 \in N^+(h) \setminus N^-(h)$. Then $\pi(d) < \pi(h)$. Thus, in the previous round, there was a vertex that was dominant for v_1, v_2 in (L', π') that has the same label as d . Since $v_1, v_2 \in N(h')$, this dominant vertex is contained in $\text{Reach}_2(L', \pi', h')$. In the construction, we iterate over all vertices $l \in \text{Reach}_2(L', \pi', h')$ in order of insertion into H' (i.e., ascending order by π') and insert vertices l' that satisfy $v_1, v_2 \in N(l')$ iff $v_1, v_2 \in N(l)$. The last vertex l with $v_1, v_2 \in N(l)$ that we encounter during this procedure was the dominant vertex of the previous round. The corresponding vertex l' has the same label as l and is the dominant vertex now.
- $v_1, v_2 \in N^-(h) \setminus N^+(h)$. As above, but with l'' instead of l' .

Bounding the Coloring Numbers. For a vertex $h' \in H'$ with $m(h') = h_1 \dots h_k$, we say that h_k is *the corresponding vertex of h' in L* . If t is a target vertex, we say the *corresponding vertex in L* is t itself. The corresponding vertex of a vertex $x \in V(L')$ is denoted by $u(x)$.

We fix a vertex $h \in H$ and ask: How many $h' \in H'$ can there be with $u(h') = h$? We pick a vertex $h' \in H'$ with $u(h') = h$. Then it has a memory $m(h') = h_1 \dots h_k$ with $h_k = h$. The following observations follow from the construction of (L', π') .

- $N(h') \neq \emptyset$
- $N(h') \subseteq N^-(h_i) \cup N^+(h_i)$ for all $1 \leq i \leq k$
- $\pi(h_1) < \dots < \pi(h_k)$

These three observations together imply that $h_1, \dots, h_k \in \text{Reach}_2(L, \pi, h)$. Thus $m(h')$ consists of a subset of $\text{Reach}_2(L, \pi, h)$, written in ascending order by π . We can therefore bound the number of possible memories $m(h')$ of h' by $2^{\text{col}_2(L, \pi)}$.

Furthermore, for each vertex $l \in H'$ with memory $l_1 \dots l_{k-1}$ and every vertex $l_k \in H$ there are at most two vertices (denoted by l' and l'' in the above construction) with

the memory $l_1 \dots l_k$. This means for a fixed memory of length k , there are at most 2^k vertices that have this particular memory. Combining the previous observations, we can conclude that for a fixed vertex $h \in H$ there are at most $2^{\text{col}_2(L, \pi)} \cdot 2^{\text{col}_2(L, \pi)} = 4^{\text{col}_2(L, \pi)}$ vertices $h' \in H'$ with $u(h') = h$.

The fact that $N(h') \subseteq N^-(u(h')) \cup N^+(u(h'))$ for all $h' \in H'$ (see second item in enumeration above) implies the following for all $x, y \in V(L')$: If x, y are adjacent in L' then $u(x), u(y)$ are adjacent in L . Also, the order π' was constructed such that $\pi(u(x)) > \pi(u(y))$ implies $\pi'(x) > \pi'(y)$ for all $x, y \in V(L')$. The last two points together mean

- if $y \in \text{Reach}_r(L', \pi', x)$ then $u(y) \in \text{Reach}_r(L, \pi, u(x))$ for all r ,
- if $y \in \text{WReach}_r(L', \pi', x)$ then $u(y) \in \text{WReach}_r(L, \pi, u(x))$ for all r .

Combining this with the observation that for a fixed $u \in V(L)$ there are at most $4^{\text{col}_2(L, \pi)}$ vertices $y \in V(L')$ with $u(y) = u$ gives us the bounds on $\text{col}_r(L', \pi')$ and $\text{wcol}_r(L', \pi')$ required by this lemma. \square

5 Transductions of Bounded Expansion Have Lacon-Decompositions

Now we combine the previous results into a more concise statement. We show that for every transduction of a graph G , we can find a lacon-decomposition whose coloring numbers are not too far off from the coloring numbers of G .

Theorem 5. *Let τ be a transduction. There exist a constant c and a function f such that for every graph G , ordering σ on the vertices of G , and $D \in \tau(G)$ there exists a lacon-decomposition (L, π) of D with*

- $\text{col}_r(L, \pi) \leq f(\text{col}_c(G, \sigma)) \cdot \text{col}_{cr}(G, \sigma)$ for all r ,
- $\text{wcol}_r(L, \pi) \leq f(\text{col}_c(G, \sigma)) \cdot \text{wcol}_{cr}(G, \sigma)$ for all r .

All the key ideas needed to prove this result have already been established in Lemma 1 and Lemma 2. In this section, we merely combine them. First, we need to following technical but unexciting lemma.

Lemma 3. *Let τ be a transduction. There exists a constant c and a basic transduction τ_0 such that for every graph G , ordering π on the vertices of G , and $D \in \tau(G)$ there exists a labeled graph G' with $\tau_0(G') = D$ and ordering π' on the vertices of G' such that*

- $\text{col}_r(G', \pi') \leq c \cdot \text{col}_r(G, \pi)$ for all r ,
- $\text{wcol}_r(G', \pi') \leq c \cdot \text{wcol}_r(G, \pi)$ for all r .

Proof. We break τ down into $\tau = \tau_0 \circ \gamma \circ \varepsilon$. Let $G' \in (\gamma \circ \varepsilon)(G)$ such that $\tau_0(G') = D$. Note that G' is the result of applying the copy operation ε and a coloring from γ to G . We extend π into an ordering π' on G' by placing the copied vertices next to the corresponding original vertices. If ε copies the graph c times then $\text{col}_r(G', \pi') \leq c \cdot \text{col}_r(G, \pi)$ and $\text{wcol}_r(G', \pi') \leq c \cdot \text{wcol}_r(G, \pi)$. \square

Now we prove Theorem 5 by chaining Lemma 3, 1 and 2 in this order.

Proof of Theorem 5. Assume we have a graph G , an ordering σ on the vertices of G , and $D \in \tau(G)$. Lemma 3 gives us a basic transduction τ_0 , a graph G' with $\tau_0(G') = D$ and an ordering σ' such that

- $\text{col}_r(G', \sigma') \leq c' \cdot \text{col}_r(G, \sigma)$ for all r ,
- $\text{wcol}_r(G', \sigma') \leq c' \cdot \text{wcol}_r(G, \sigma)$ for all r .

for some constant c' depending only on τ . Assume τ_0 to be of the form (χ, ν, φ) . By Lemma 1, there exists a function g and a directed lacon-decomposition (L', π') of $I_\varphi(G')$ with

- $\text{col}_r(L', \pi') \leq g(|\varphi| + \text{col}_{2,4|\varphi|}(G', \sigma')) \cdot \text{col}_{4|\varphi|_r}(G', \sigma')$ for all r ,
- $\text{wcol}_r(L', \pi') \leq g(|\varphi| + \text{col}_{2,4|\varphi|}(G', \sigma')) \cdot \text{wcol}_{4|\varphi|_r}(G', \sigma')$ for all r .

We know $G' \models \chi$, since otherwise $\tau_0(G')$ would be undefined. Since ν describes the vertex set of the transduction, we update L' by removing all target vertices t with $L' \not\models \nu(t)$. (L', π') is now a directed lacon-decomposition of $\tau_0(G') = D$. Since we only deleted vertices, the coloring numbers of (L', π') did not increase, thus the above two bounds remain true. At last, we use Lemma 2 to construct an undirected lacon-decomposition (L, π) of D with

- $\text{col}_r(L, \pi) \leq 4^{\text{col}_2(L', \pi')} \cdot \text{col}_r(L', \pi')$ for all r ,
- $\text{wcol}_r(L, \pi) \leq 4^{\text{col}_2(L', \pi')} \cdot \text{wcol}_r(L', \pi')$ for all r .

Combining the previous 6 bounds then proves the result. \square

6 Converting Lacon- to Shrub-Decompositions

We now convert a lacon-decomposition into an equivalent shrub-decomposition while maintaining a bound on the generalized coloring numbers.

Lemma 4. *Let G be a graph with a lacon-decomposition (L, π) . Then there exists a shrub-decomposition (F, S) of G with one color, diameter at most $4\text{col}_1(L, \pi) + 4$ and*

- $\text{col}_r(F) \leq d + \text{col}_r(L, \pi)$ for all r ,
- $\text{wcol}_r(F) \leq d + \text{wcol}_r(L, \pi)$ for all r .




Figure 3: Left: Part of a lacon-decomposition. Right: Corresponding shrub-decomposition, including distances between round and square vertices.

Proof. Let (L, π) be a lacon-decomposition of G . For every target vertex t and $1 \leq i < |N(t)|$ we define $\nu_i(t)$ to be the i th hidden neighbor in L , counted in descending order by π . We construct a graph F from L by performing the following two operations for every vertex $t \in V(G)$ (see Figure 3).

- Add a vertex t' and make t' the only neighbor of t .
- For each $1 \leq i < |N(t)|$ add a path of length $2i+1$ from t' to $\nu_i(t)$ if $\nu_i(t)$ is labeled with one, and of length $2i$ if $\nu_i(t)$ is labeled with zero.

We furthermore iteratively remove pendant vertices in $V(F) \setminus V(G)$ from F until the pendant vertices are exactly $V(G)$. This last operation does not change the distance between any two vertices from $V(G)$ in F , since no path can pass through a pendant vertex.

It is easy to see that the distance between any two vertices $t_1, t_2 \in V(G)$ is even, and thus can be written as $2d$ for $d \in \mathbb{N}$. Furthermore, the shortest path between t_1 and t_2 passes through their largest common neighbor in L . This leads to the following crucial observation: The half-distance d is even iff the largest common neighbor of t_1 and t_2 is labeled with one. Thus, two vertices are adjacent in G iff their distance in F is a multiple of four.

Since our construction does not rely on any colors, we give every target vertex the same dummy color 1. We then define the signature of the shrub decomposition as

$$S = \{(1, 1, 4d) \mid 1 \leq d \leq \text{col}_1(L, \pi) + 1\}.$$

The diameter of F is at most $4\text{col}_1(L, \pi) + 4$. The bound on the coloring numbers of F can be proven as follows. Let H be the hidden vertices of L . Every connected component of $F[V(F) \setminus H]$ consists of a vertex $t \in V(G)$, a vertex t' and a bounded number of short paths attached to t' . Each component is bounded in size by a function of $\text{col}_2(L, \pi)$. We extend the ordering π of L into an ordering π' of F by making the vertices in $V(F) \setminus V(L)$ larger than all vertices in $V(L)$. Now for every $h \in H$

holds $\text{Reach}_r(F, \pi', h) \subseteq \text{Reach}_r(L, \pi, h)$. For all other vertices $v \in V(F) \setminus H$ holds $\text{Reach}_r(F, \pi', v) \subseteq \text{Reach}_r(L, \pi, t) \cup X$, where X is the connected component of v in $F[V(F) \setminus H]$ and $t \in V(G)$ is the unique target vertex contained in X . An equivalent bound can be obtained for the weak reachability. \square

7 Converting Lacon- to Parity-Decompositions

The following proof is substantially similar to the one of Lemma 2.

Lemma 5. *Assume a graph G has a lacon-decomposition (L, π) . Then it also has a parity-decomposition P with target-degree at most $2^{\text{col}_2(L, \pi)} \cdot \text{col}_1(L, \pi)$ and*

- $\text{col}_r(P) \leq 2^{\text{col}_2(L, \pi)} \cdot \text{col}_r(L, \pi)$ for all r ,
- $\text{wcol}_r(P) \leq 2^{\text{col}_2(L, \pi)} \cdot \text{wcol}_r(L, \pi)$ for all r .

Proof. Assume we have a lacon-decomposition (L, π) of a graph G with target vertices T and hidden vertices H . We construct a parity-decomposition P of G with target and hidden vertices T, H' as follows: We start with $H' = \emptyset$. We process the vertices in H one by one in ascending order by π and add in each processing step various new vertices to H' . This defines an ordering π' of $V(P)$ as follows (in ascending order): First come the vertices of H' in the order of insertion and then come the vertices of T in the same order as in π . Afterwards, we will use π to bound the coloring numbers of P .

We will make sure that after a vertex $h \in H$ has been processed the following invariant holds for all pairs of vertices $v_1 \neq v_2 \in T$. If v_1, v_2 have a dominant vertex d in L such that $\pi(d) \leq \pi(h)$ then we guarantee that $|N(v_1) \cap N(v_2)|$ is odd in P if and only if d labeled with “1”. After every vertex in H has been processed, this guarantees that P is a parity-decomposition of G .

While processing a vertex $h \in H$, certain new vertices are inserted into H' . The newly inserted vertices are in a sense either a copy of h or copies of vertices that have already been inserted into H' . These copies may later again be copied, and so forth. This leads to exponential growth. To keep track of these copies and to show that while being exponential, this growth is nevertheless bounded, every newly inserted vertex h' will have a *memory*, denoted by $m(h')$. This memory consists of a sequence of vertices from H . If h' is derived from h , then we set $m(h') = h$. However, if h' is derived both from h and another vertex h'' that has previously been inserted into H' , then we set $m(h') = m(h'') + h$, i.e., $m(h')$ is obtained by appending h to the memory of h'' . Let us now get into the details of the construction and prove our invariant.

Construction of Parity-Decomposition. We describe the processing steps of a vertex $h \in H$. If $N^L(h) = \emptyset$ we do nothing. Otherwise iterate over all $l \in H'$ with $N^P(l) \cap N^L(h) \neq \emptyset$ in order of insertion to H' , and add a vertex l' to P with $N^P(l') = N^P(l) \cap N^L(h)$ and $m(l') = m(l) + h$. Also, if h is labeled with “1”, we add an additional vertex h' to P with $N^P(h') = N^L(h)$ and $m(h') = h$. By induction, we know that the

invariant was satisfied in the last round for all v_1, v_2 . We show that it also holds after h is processed. Let $v_1 \neq v_2$ be two vertices with dominant vertex d in L such that $\pi(d) \leq \pi(h)$. We distinguish two cases.

- $v_1 \notin N^L(h)$ or $v_2 \notin N^L(h)$. In this case, h is not dominant for v_1, v_2 in (L, π) . Also none of the newly inserted vertices are contained in $|N^P(v_1) \cap N^P(v_2)|$. Since the invariant for v_1, v_2 held in the previous round, it also holds in this round.
- $v_1, v_2 \in N^L(h)$. Now, h is dominant for v_1, v_2 in (L, π) . If h is labeled with “0” then $|N^P(v_1) \cap N^P(v_2)|$ is even. This is because for every element $l \in N^P(v_1) \cap N^P(v_2)$ from the previous round, a “copy” l' with $v_1, v_2 \in N^P(l')$ has been inserted. On the other hand, if h is labeled with “1”, the additional vertex h' with $N^P(h') = N^L(h)$ makes $|N^P(v_1) \cap N^P(v_2)|$ odd.

Bounding the Coloring Numbers. For a vertex $h' \in H'$ with $m(h') = h_1 \dots h_k$, we say that h_k is *the corresponding vertex of h' in L* . If t is a target vertex, we say the *corresponding vertex in L* is t itself. The corresponding vertex of a vertex $x \in V(P)$ is denoted by $u(x)$.

We fix a vertex $h \in H$ and ask: How many $h' \in H'$ can there be with $u(h') = h$? We pick a vertex $h' \in H'$ with $u(h') = h$. Then it has a memory $m(h') = h_1 \dots h_k$ with $h_k = h$. The following observations follow from the construction of P .

- $N^P(h') \neq \emptyset$
- $N^P(h') \subseteq N^L(h_i)$ for all $1 \leq i \leq k$
- $\pi(h_1) < \dots < \pi(h_k)$

These three observations together imply that $h_1, \dots, h_k \in \text{Reach}_2(L, \pi, h)$. Thus $m(h')$ consists of a non-empty subset of $\text{Reach}_2(L, \pi, h)$, written in ascending order by π . We can therefore bound the number of possible memories $m(h')$ of h' by $2^{\text{col}_2(L, \pi)} - 1$. Also, every vertex $h' \in H'$ is uniquely identified by its memory $m(h')$ and every vertex $t \in T$ has $u(t) = t$. We can therefore conclude for every $h \in H$ that there are at most $2^{\text{col}_2(L, \pi)}$ vertices $h' \in V(P)$ with $u(h') = h$.

The fact that $N^P(h') \subseteq N^L(u(h'))$ for all $h' \in H'$ (see second item in enumeration above) implies the following for all $x, y \in V(P)$: If x, y are adjacent in P then $u(x), u(y)$ are adjacent in L . Also, the order π' was constructed such that $\pi(u(x)) > \pi(u(y))$ implies $\pi'(x) > \pi'(y)$ for all $x, y \in V(P)$. The last two points together mean

- if $y \in \text{Reach}_r(P, \pi', x)$ then $u(y) \in \text{Reach}_r(L, \pi, u(x))$ for all r ,
- if $y \in \text{WReach}_r(P, \pi', x)$ then $u(y) \in \text{WReach}_r(L, \pi, u(x))$ for all r .

Combining this with the observation that for a fixed $u \in V(L)$ there are at most $2^{\text{col}_2(L, \pi)}$ vertices $y \in V(P)$ with $u(y) = u$ gives us the bounds on $\text{col}_r(P, \pi')$ and $\text{wcol}_r(P, \pi')$ required by this lemma. Since every target vertex in L has degree at most $\text{col}_1(L, \pi)$, P has target-degree at most $2^{\text{col}_2(L, \pi)} \cdot \text{col}_1(L, \pi)$. \square

8 Proof of Theorem 1, 2 and 3

We finally prove the main result of this paper by combining all results of the previous sections.

Proof of Theorem 1. We prove (1) \implies (2) \implies (3) \implies (1) and then (2) \implies (4) \implies (1).

- (1) \implies (2). Pick any graph $D \in \mathcal{G}$. By (1) there exists a graph $G \in \mathcal{G}'$ with $D \in \tau(G)$. Since \mathcal{G}' has bounded expansion, Definition 5 states that there exists an ordering σ with $\text{col}_r(G, \sigma) \leq g(r)$ for all r (where the function $g(r)$ depends only on \mathcal{G}'). Now, (2) directly follows from Theorem 5.
- (2) \implies (3). This follows from Lemma 4 and Definition 5.
- (3) \implies (1). In first-order logic we can compute the (bounded) distance between two vertices, check their color and implement the lookup table S . Thus, we can easily construct a transduction τ with $\tau(\mathcal{G}') \subseteq \mathcal{G}$.
- (2) \implies (4). This follows from Lemma 5 and Definition 5.
- (4) \implies (1). Since the target-degree is bounded, first-order logic can check the parity of the intersection of two neighborhoods.

□

We proceed in a similar way to prove our characterizations of structurally bounded treedepth and treewidth.

Proof of Theorem 2. The implications (1) \implies (2) \implies (3) \implies (1) and then (2) \implies (4) \implies (1) can be proven analogously to the proof of Theorem 1. We only list the first implication.

- (1) \implies (2). As mentioned in the introduction, the weak coloring numbers converge to treedepth, i.e., $\text{wcol}_1(G) \leq \dots \leq \text{wcol}_\infty(G) = \text{td}(G)$ [17]. Assume every graph in \mathcal{G}' has treedepth at most d' . Thus, for a graph $D \in \mathcal{G}$ there exists a graph $G \in \mathcal{G}'$ with $D \in \tau(G)$ and $\text{wcol}_\infty(G) \leq d'$. Theorem 5 then gives us a laconic decomposition (L, π) of D with $\text{wcol}_\infty(L, \pi) \leq f(\text{col}_c(G)) \cdot \text{wcol}_\infty(G) \leq f(d') \cdot d'$.

□

Proof of Theorem 3. The proof proceeds as the one of Theorems 1 and 2, except this time using $\text{col}_1(G) \leq \dots \leq \text{col}_\infty(G) = \text{tw}(G) + 1$ [17].

□

9 Proof of Localized Feferman–Vaught Composition Theorem

We give a self sustained proof of Theorem 4, using central ideas from the proof of Gaifman’s theorem [16]. An alternative proof exists in [29, Lemma 15]. We repeat the relevant definitions and then prove the result.

Definition 6 (q-type [16]). *Let G be a labeled graph and $\bar{v} = (v_1, \dots, v_k) \in V(G)^k$, for some nonnegative integer k . The first-order q-type of \bar{v} in G is the set $\text{tp}_q(G, \bar{v})$ of all first-order formulas $\psi(x_1 \dots x_k)$ of rank at most q such that $G \models \psi(v_1 \dots v_k)$.*

Definition 7. *Let G be a graph, $r \in \mathbf{N}$, and $\bar{u}, \bar{v}_1, \dots, \bar{v}_k$ be tuples of vertices from G . We say \bar{u} r -separates \bar{v}_i and \bar{v}_j if every path of length at most r between a vertex from \bar{v}_i and a vertex from \bar{v}_j contains at least one vertex from \bar{u} . We say \bar{u} r -separates $\bar{v}_1, \dots, \bar{v}_k$ if it r -separates \bar{v}_i and \bar{v}_j for all $i \neq j$.*

Theorem 4. *There exists a function $f(q, l)$ such that for all labeled graphs G , every $q, l \in \mathbf{N}$, and all tuples $\bar{u}, \bar{v}_1, \dots, \bar{v}_k$ of vertices from G such that \bar{u} 4^q -separates $\bar{v}_1, \dots, \bar{v}_k$ and $|\bar{u}| + |\bar{v}_1| + \dots + |\bar{v}_k| \leq l$, the type $\text{tp}_q(G, \bar{u}\bar{v}_1 \dots \bar{v}_k)$ depends only on the types $\text{tp}_{f(q, l)}(G, \bar{u}\bar{v}_1), \dots, \text{tp}_{f(q, l)}(G, \bar{u}\bar{v}_k)$. Furthermore, $\text{tp}_q(G, \bar{u}\bar{v}_1 \dots \bar{v}_k)$ can be computed from $\text{tp}_{f(q, l)}(G, \bar{u}\bar{v}_1), \dots, \text{tp}_{f(q, l)}(G, \bar{u}\bar{v}_k)$.*

Proof. Our result follows from proving the following claim via structural induction. Let $\varphi(\bar{x}, \bar{y}_1, \dots, \bar{y}_k)$ be a first-order formula with quantifier rank q . Then one can compute a boolean combination $\Phi(\bar{x}, \bar{y}_1, \dots, \bar{y}_k)$ of first-order formulas of the form $\xi(\bar{x}, \bar{y}_i)$ such that for all graphs G and all tuples $\bar{v}_1, \dots, \bar{v}_k$ that are 4^q -separated by a tuple \bar{u} in G holds

$$G \models \varphi(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \iff G \models \Phi(\bar{u}, \bar{v}_1, \dots, \bar{v}_k).$$

We call such a boolean combination Φ a *separated expression*. The claim holds for atomic formulas because \bar{u} 4^0 -separates $\bar{v}_1, \dots, \bar{v}_k$, i.e., there are no edges between $\{\bar{v}_i\} \setminus \{\bar{u}\}$ and $\{\bar{v}_j\} \setminus \{\bar{u}\}$ in G for $i \neq j$. It also holds for boolean combinations and negations. It remains to prove our claim for the case that $\varphi(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) = \exists x \psi$ for some formula ψ .

For $t \in \mathbf{N}$ and $v \in V(G)$, we define $N_t^{\bar{u}}(v)$ to be the set of all vertices in G that are reachable from v via a path of length at most t that contains no vertex from \bar{u} . For a tuple \bar{v} , we define $N_t^{\bar{u}}(\bar{v}) = \bigcup_{v \in \{\bar{v}\}} N_t^{\bar{u}}(v)$. With slight abuse of notation, we write $z \in N_t^{\bar{x}}(\bar{y})$ as a short-hand for a first-order formula $\nu_t(\bar{x}, \bar{y}, z)$ such that for all graphs G , vertex-tuples \bar{u}, \bar{v} and vertices w holds $G \models \nu_t(\bar{u}, \bar{v}, w)$ iff $w \in N_t^{\bar{u}}(\bar{v})$. The formula $\nu_t(\bar{x}, \bar{y}, z)$ can be constructed using $t - 1$ existential quantifiers.

Let $r = 4^{q-1}$. We rewrite φ by distinguishing two cases: Either the existentially quantified variable x is contained in the neighborhood $N_r^{\bar{x}}(\bar{y}_i)$ for some i or for none. This gives us

$$\varphi(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) \equiv \varphi^*(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) \vee \bigvee_{i=1}^k \varphi_i(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) \quad (5)$$

with

$$\begin{aligned}\varphi_i(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) &= \exists x x \in N_r^{\bar{x}}(\bar{y}_i) \wedge \psi(\bar{x}, x, \bar{y}_1, \dots, \bar{y}_k), \\ \varphi^*(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) &= \exists x x \notin \bigcup_{i=1}^k N_r^{\bar{x}}(\bar{y}_i) \wedge \psi(\bar{x}, x, \bar{y}_1, \dots, \bar{y}_k).\end{aligned}$$

We will proceed by finding separated expressions for φ^* and for each φ_i . Since $4^q = 4r$, we consider a graph G and tuples $\bar{v}_1, \dots, \bar{v}_k$ that are $4r$ -separated by a tuple \bar{u} in G . We start with φ_i . This formula asks whether there exists an $x \in N_r^{\bar{x}}(\bar{y}_i)$ satisfying ψ . For every $u \in N_r^{\bar{u}}(\bar{v}_i)$ holds that \bar{u} r -separates the tuples $\bar{v}_i u$ and \bar{v}_j for $i \neq j$, since otherwise there would be a short path from \bar{v}_i via u to \bar{v}_j that contains no vertex from \bar{u} . By the induction hypothesis, there exists a separated expression $\Psi_i(\bar{x}, \bar{y}_1, \dots, \bar{y}_i x, \dots, \bar{y}_k)$ such that for all $u \in N_r^{\bar{u}}(\bar{v}_i)$

$$G \models \psi(\bar{u}, u, \bar{v}_1, \dots, \bar{v}_k) \iff G \models \Psi_i(\bar{u}, \bar{v}_1, \dots, \bar{v}_i u, \dots, \bar{v}_k).$$

Thus $G \models \varphi_i(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \iff G \models \exists x x \in N_r^{\bar{x}}(\bar{v}_i) \wedge \Psi_i(\bar{u}, \bar{v}_1, \dots, \bar{v}_i x, \dots, \bar{v}_k)$. If we assume Ψ_i to be of the form

$$\Psi_i = \bigvee_{l=1}^m \left(\xi_{li}(\bar{x}, \bar{y}_i x) \wedge \bigwedge_{\substack{j=1 \\ j \neq i}}^k \xi_{lj}(\bar{x}, \bar{y}_j) \right)$$

then $G \models \varphi_i(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \iff G \models \Phi_i(\bar{u}, \bar{v}_1, \dots, \bar{v}_k)$ with the separated expression

$$\Phi_i(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) = \bigvee_{l=1}^m \left(\exists x x \in N_r^{\bar{x}}(\bar{y}_i) \wedge \xi_{li}(\bar{x}, \bar{y}_i x) \wedge \bigwedge_{\substack{j=1 \\ j \neq i}}^k \xi_{lj}(\bar{x}, \bar{y}_j) \right).$$

Next, we want to obtain a separated expression for φ^* . Let therefore $u \in V(G)$ with $u \notin N_r^{\bar{u}}(\bar{v}_i)$ for all i . Then the tuples $u, \bar{v}_1, \dots, \bar{v}_k$ are r -separated by \bar{u} . By the induction hypothesis, there exists a separated expression $\Psi^*(\bar{x}, x, \bar{y}_1, \dots, \bar{y}_k)$ such that

$$G \models \psi(\bar{u}, u, \bar{v}_1, \dots, \bar{v}_k) \iff G \models \Psi^*(\bar{u}, u, \bar{v}_1, \dots, \bar{v}_k).$$

We can assume Ψ^* to be of the form

$$\Psi^* = \bigvee_{l=1}^m \left(\xi_l(\bar{x}, x) \wedge \bigwedge_{j=1}^k \xi_{lj}(\bar{x}, \bar{y}_j) \right).$$

Then $G \models \varphi^*(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \iff G \models \Phi^*(\bar{u}, \bar{v}_1, \dots, \bar{v}_k)$ using the separated expression

$$\Phi^*(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) = \bigvee_{l=1}^m \left(\exists x x \notin \bigcup_{i=1}^k N_r^{\bar{x}}(\bar{v}_i) \wedge \xi_l(\bar{x}, x) \wedge \bigwedge_{j=1}^k \xi_{lj}(\bar{x}, \bar{y}_j) \right).$$

Note that Φ^* is not yet a separated expression. Nevertheless, substituting Φ_i and Φ^* into equation (5) yields

$$G \models \varphi(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \iff G \models \bigvee_{i=1}^k \Phi_i(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \vee \bigvee_{l=1}^m \left(\exists x \, x \notin \bigcup_{i=1}^k N_r^{\bar{u}}(\bar{v}_i) \wedge \xi_l(\bar{u}, x) \bigwedge_{j=1}^k \xi_{lj}(\bar{u}, \bar{v}_j) \right).$$

The remaining problematic subformulas are those of the form

$$\gamma(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) = \exists x \, x \notin \bigcup_{i=1}^k N_r^{\bar{x}}(\bar{y}_i) \wedge \xi(\bar{x}, x).$$

To complete the proof, it is sufficient to find a separated expression equivalent to γ . We therefore define the separated expression

$$\Gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) = \bigvee_{i=1}^k \exists x \, x \in N_{3r}^{\bar{u}}(\bar{v}_i) \wedge x \notin N_r^{\bar{u}}(\bar{v}_i) \wedge \xi(\bar{u}, x)$$

and make a case distinction based on it.

- *Case 1:* $G \models \Gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k)$. Since $\bar{v}_1, \dots, \bar{v}_k$ are $4r$ -separated by \bar{u} , we have $N_{3r}^{\bar{u}}(\bar{v}_i) \cap N_r^{\bar{u}}(\bar{v}_j) = \emptyset$ for $i \neq j$. This means $G \models \gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k)$.
- *Case 2:* $G \not\models \Gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k)$. We define the \bar{u} -distance between two vertices to be the length of the shortest path between them that contains no vertex from \bar{u} . For $V' \subseteq V(G)$, we define an $(2r, \bar{u})$ -scattered subset of V' to be a set $S \subseteq V'$ such that all vertices in S pairwise have \bar{u} -distance greater than $2r$ in G . The size of the largest $(2r, \bar{u})$ -scattered subset of V' in G is denoted by $s(V')$. Let further $R(\bar{u}) = \{u \mid u \in V(G), G \models \xi(\bar{u}, u)\}$. Using this new notation, we observe that $G \models \gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k)$ if and only if $R(\bar{u}) \setminus \bigcup_{i=1}^k N_r^{\bar{u}}(\bar{v}_i) \neq \emptyset$. Since we assume $G \not\models \Gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k)$, this means that the sets $R(\bar{u}) \setminus \bigcup_{i=1}^k N_r^{\bar{u}}(\bar{v}_i)$, $R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_1), \dots, R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_k)$ are $2r$ -separated by \bar{u} . Therefore

$$s(R(\bar{u})) = s\left(R(\bar{u}) \setminus \bigcup_{i=1}^k N_r^{\bar{u}}(\bar{v}_i)\right) + \sum_{i=1}^k s\left(R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_i)\right).$$

Thus, $G \models \gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k)$ iff $R(\bar{u}) \setminus \bigcup_{i=1}^k N_r^{\bar{u}}(\bar{v}_i) \neq \emptyset$ iff $s\left(R(\bar{u}) \setminus \bigcup_{i=1}^k N_r^{\bar{u}}(\bar{v}_i)\right) \neq 0$ iff $\sum_{i=1}^k s\left(R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_i)\right) < s(R(\bar{u}))$.

The previous case distinction implies that

$$G \models \gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \iff G \models \Gamma(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \text{ or } \sum_{i=1}^k s\left(R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_i)\right) < s(R(\bar{u})).$$

It now remains to construct a separated expression $\Delta(\bar{x}, \bar{y}_1, \dots, \bar{y}_k)$ such that $G \models \Delta(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \iff \sum_{i=1}^k s(R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_i)) < s(R(\bar{u}))$. For arbitrary vertices a, b, c such that $a, b \in N_r^{\bar{u}}(c)$, we see that a, b have \bar{u} -distance at most $2r$. Since $s(R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_i))$ is the size of a subset of $N_r^{\bar{u}}(\bar{v}_i)$ where all vertices pairwise have \bar{u} -distance greater than $2r$, we have $s(R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_i)) \leq |\bar{v}_i|$.

For $1 \leq i \leq k$ and $0 \leq h \leq |\bar{v}_i|$ we define a first-order formula $\delta_{ih}(\bar{x}, \bar{y}_i)$ which is true iff $s(R(\bar{x}) \cap N_r^{\bar{x}}(\bar{y}_i)) \leq h$:

$$\delta_{ih}(\bar{x}, \bar{y}_i) = \neg \exists s_1 \dots \exists s_{h+1} \bigwedge_{j=1}^{h+1} \left(\xi(\bar{x}, s_j) \wedge s_j \in N_r^{\bar{x}}(\bar{y}_i) \wedge \bigwedge_{l=j+1}^{h+1} s_j \notin N_{2r}^{\bar{x}}(s_l) \right).$$

We further define for $0 \leq h \leq \sum_{i=1}^k |\bar{v}_i|$ a first-order formula $\delta_h(\bar{x})$ which is true iff $s(R(\bar{x})) > h$. This formula can be constructed in a similar way as the formulas $\delta_{ih}(\bar{x}, \bar{y}_i)$ above. We use these formulas to construct the separated expression

$$\Delta(\bar{x}, \bar{y}_1, \dots, \bar{y}_k) = \bigvee_{h_1=0}^{|\bar{v}_1|} \dots \bigvee_{h_k=0}^{|\bar{v}_k|} \delta_{1h_1}(\bar{x}, \bar{y}_1) \wedge \dots \wedge \delta_{kh_k}(\bar{x}, \bar{y}_k) \wedge \delta_{\sum_{i=1}^k h_i}(\bar{x})$$

with $G \models \Delta(\bar{u}, \bar{v}_1, \dots, \bar{v}_k) \iff \sum_{i=1}^k s(R(\bar{u}) \cap N_r^{\bar{u}}(\bar{v}_i)) < s(R(\bar{u}))$. This completes our decomposition of φ into separated expressions. \square

References

- [1] Édouard Bonnet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant. Twin-width I: tractable FO model checking. In *2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 601–612, 2020. [doi:10.1109/FOCS46700.2020.00062](https://doi.org/10.1109/FOCS46700.2020.00062).
- [2] Jianer Chen, Xiuzhen Huang, Iyad A Kanj, and Ge Xia. Strong computational lower bounds via parameterized complexity. *Journal of Computer and System Sciences*, 72(8):1346–1367, 2006.
- [3] Bruno Courcelle. The monadic second-order logic of graphs I. Recognizable sets of finite graphs. *Inf. Comput.*, 85(1):12–75, 1990. [doi:10.1016/0890-5401\(90\)90043-H](https://doi.org/10.1016/0890-5401(90)90043-H).
- [4] Patrice Ossona de Mendez, Jan Obdržálek, Jaroslav Nešetřil, Petr Hliněný, and Robert Ganian. Shrub-depth: Capturing height of dense graphs. *Logical Methods in Computer Science*, 15, 2019.
- [5] Jan Dreier, Jakub Gajarský, Sandra Kiefer, Michał Pilipczuk, and Szymon Toruńczyk. Treelike decompositions for transductions of sparse graphs. 2022. [arXiv:2201.11082](https://arxiv.org/abs/2201.11082).

- [6] Jan Dreier, Philipp Kuinke, and Peter Rossmanith. First-order model-checking in random graphs and complex networks. In *28th Annual European Symposium on Algorithms (ESA 2020)*, volume 173 of *LIPICs*. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- [7] Jan Dreier and Peter Rossmanith. Approximate evaluation of first-order counting queries. In *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1720–1739. SIAM, 2021.
- [8] Zdenek Dvořák, Daniel Král, and Robin Thomas. Deciding First-Order Properties for Sparse Graphs. In *Proceedings of the 51st Conference on Foundations of Computer Science*, pages 133–142, 2010.
- [9] Jörg Flum and Martin Grohe. Fixed-Parameter Tractability, Definability, and Model-Checking. *SIAM Journal on Computing*, 31(1):113–145, 2001.
- [10] Markus Frick and Martin Grohe. Deciding first-order properties of locally tree-decomposable structures. *J. ACM*, 48(6):1184–1206, 2001. [doi:10.1145/504794.504798](https://doi.org/10.1145/504794.504798).
- [11] Haim Gaifman. On local and non-local properties. In *Studies in Logic and the Foundations of Mathematics*, volume 107, pages 105–135. Elsevier, 1982.
- [12] Jakub Gajarský, P Hliněný, J Obdržálek, Daniel Lokshtanov, and MS Ramanujan. A New Perspective on FO Model Checking of Dense Graph Classes. In *Proceedings of the 31st Symposium on Logic in Computer Science*, pages 176–184, 2016.
- [13] Jakub Gajarský and Daniel Král'. Recovering Sparse Graphs. In Igor Potapov, Paul Spirakis, and James Worrell, editors, *43rd International Symposium on Mathematical Foundations of Computer Science (MFCS 2018)*, volume 117 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 29:1–29:15, Dagstuhl, Germany, 2018. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik. [doi:10.4230/LIPIcs.MFCS.2018.29](https://doi.org/10.4230/LIPIcs.MFCS.2018.29).
- [14] Jakub Gajarský, Stephan Kreutzer, Jaroslav Nešetřil, Patrice Ossona De Mendez, Michał Pilipczuk, Sebastian Siebertz, and Szymon Toruńczyk. First-order interpretations of bounded expansion classes. *ACM Trans. Comput. Logic*, 21(4), July 2020. [doi:10.1145/3382093](https://doi.org/10.1145/3382093).
- [15] Robert Ganian, Petr Hliněný, Jaroslav Nešetřil, Jan Obdržálek, Patrice Ossona de Mendez, and Reshma Ramadurai. When trees grow low: Shrubs and fast mso1. In Branislav Rován, Vladimiro Sassone, and Peter Widmayer, editors, *Mathematical Foundations of Computer Science 2012*, pages 419–430, Berlin, Heidelberg, 2012. Springer Berlin Heidelberg.
- [16] Martin Grohe. Logic, graphs, and algorithms. *Logic and Automata*, 2:357–422, 2008.

- [17] Martin Grohe, Stephan Kreutzer, Roman Rabinovich, Sebastian Siebertz, and Konstantinos S. Stavropoulos. Coloring and covering nowhere dense graphs. *SIAM J. Discret. Math.*, 32(4):2467–2481, 2018. doi:[10.1137/18M1168753](https://doi.org/10.1137/18M1168753).
- [18] Martin Grohe, Stephan Kreutzer, and Sebastian Siebertz. Deciding first-order properties of nowhere dense graphs. *J. ACM*, 64(3):17:1–17:32, 2017. doi:[10.1145/3051095](https://doi.org/10.1145/3051095).
- [19] William Hanf. Model-theoretic methods in the study of elementary logic. In J. W. Addison, editor, *Journal of Symbolic Logic*, pages 132–145. Amsterdam: North-Holland Pub. Co., 1965.
- [20] Carol Karp. The first order properties of products of algebraic systems. *fundamenta mathematicae. Journal of Symbolic Logic*, 32(2):276–276, 1967. doi:[10.2307/2271704](https://doi.org/10.2307/2271704).
- [21] Wojciech Kazana and Luc Segoufin. Enumeration of first-order queries on classes of structures with bounded expansion. In *Proceedings of the 32nd ACM SIGMOD-SIGACT-SIGART Symposium on Principles of Database Systems, PODS 2013, New York, NY, USA - June 22 - 27, 2013*, pages 297–308. ACM, 2013. doi:[10.1145/2463664.2463667](https://doi.org/10.1145/2463664.2463667).
- [22] Henry A. Kierstead and Daqing Yang. Orderings on graphs and game coloring number. *Order*, 20(3):255–264, 2003. doi:[10.1023/B:ORDE.0000026489.93166.cb](https://doi.org/10.1023/B:ORDE.0000026489.93166.cb).
- [23] Dietrich Kuske and Nicole Schweikardt. First-order logic with counting. In *32nd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2017, Reykjavik, Iceland, June 20-23, 2017*, pages 1–12. IEEE Computer Society, 2017. doi:[10.1109/LICS.2017.8005133](https://doi.org/10.1109/LICS.2017.8005133).
- [24] Johann A Makowsky. Algorithmic uses of the Feferman–Vaught theorem. *Annals of Pure and Applied Logic*, 126(1-3):159–213, 2004.
- [25] Jaroslav Nešetřil and Patrice Ossona De Mendez. On low tree-depth decompositions. *Graphs and combinatorics*, 31(6):1941–1963, 2015.
- [26] Jaroslav Nešetřil, Patrice Ossona de Mendez, and Sebastian Siebertz. Towards an arboretum of monadically stable classes of graphs. *arXiv preprint arXiv:2010.02607*, 2020.
- [27] Jaroslav Nešetřil and Patrice Ossona de Mendez. *Sparsity - Graphs, Structures, and Algorithms*, volume 28 of *Algorithms and combinatorics*. Springer, 2012. doi:[10.1007/978-3-642-27875-4](https://doi.org/10.1007/978-3-642-27875-4).
- [28] Sang-Il Oum. Approximating rank-width and clique-width quickly. *ACM Transactions on Algorithms (TALG)*, 5(1):1–20, 2008. doi:[10.1145/1435375.1435385](https://doi.org/10.1145/1435375.1435385).

- [29] Michał Pilipczuk, Sebastian Siebertz, and Szymon Toruńczyk. On the number of types in sparse graphs. In *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science*, pages 799–808, 2018.
- [30] Nicole Schweikardt, Luc Segoufin, and Alexandre Vigny. Enumeration for FO queries over nowhere dense graphs. In *Proceedings of the 37th ACM SIGMOD-SIGACT-SIGAI Symposium on Principles of Database Systems, Houston, TX, USA, June 10-15, 2018*, pages 151–163. ACM, 2018. doi:[10.1145/3196959.3196971](https://doi.org/10.1145/3196959.3196971).
- [31] Detlef Seese. Linear time computable problems and first-order descriptions. *Math. Struct. Comput. Sci.*, 6(6):505–526, 1996.
- [32] Szymon Toruńczyk. Aggregate queries on sparse databases. In *Proceedings of the 39th ACM SIGMOD-SIGACT-SIGAI Symposium on Principles of Database Systems, PODS 2020, Portland, OR, USA, June 14-19, 2020*, pages 427–443. ACM, 2020. doi:[10.1145/3375395.3387660](https://doi.org/10.1145/3375395.3387660).
- [33] Jan van den Heuvel and H.A. Kierstead. Uniform orderings for generalized coloring numbers. *European Journal of Combinatorics*, 91:103214, 2021. Colorings and structural graph theory in context (a tribute to Xuding Zhu). doi:<https://doi.org/10.1016/j.ejc.2020.103214>.
- [34] Xuding Zhu. Colouring graphs with bounded generalized colouring number. *Discrete Mathematics*, 309(18):5562–5568, 2009.