

Universal Horn Sentences and the Joint Embedding Property ^{*}

Manuel Bodirsky¹, Jakub Rydval², and André Schrottenloher³

¹ Institute of Algebra, TU Dresden, Germany
manuel.bodirsky@tu-dresden.de

² Institute of Theoretical Computer Science, TU Dresden, Germany
jakub.rydval@tu-dresden.de

³ Cryptology Group, CWI, Amsterdam, The Netherlands
andre.schrottenloher@m4x.org

Abstract. The finite models of a universal sentence Φ in a finite relational signature are the age of a structure if and only if Φ has the *joint embedding property*. We prove that the computational problem whether a given universal sentence Φ has the joint embedding property is undecidable, even if Φ is additionally Horn and the signature of Φ only contains relation symbols of arity at most two.

Keywords: joint embedding property · universal · Horn

1 Introduction

The preservation theorem of McKinsey states that a universal sentence Φ is equivalent to a universal Horn sentence if and only if the class of models of Φ is closed under direct products [Hod93]. Universal Horn sentences are particularly relevant in theoretical computer science; we mention two contexts where they appear.

- The *constraint satisfaction problem* of a structure \mathfrak{B} with a finite relational signature τ is the problem of deciding whether a given finite τ -structure \mathfrak{A} has a homomorphism to \mathfrak{B} . This problem is in NP if the age of \mathfrak{B} can be described by a universal sentence, and it is in P if the age of \mathfrak{B} can be described by a universal Horn sentence.
- Baader and Rydval [BR21] identified which relational structures whose age can be described by a universal sentence yield tractable extensions of the description logic \mathcal{EL} when employed as a concrete domain if equalities are allowed in atomic formulas. This is the case if and only if the age of the structure can be described by a universal Horn sentence.

^{*} Supported by DFG GRK 1763 (QuantLA). Manuel Bodirsky has received funding from the European Research Council through the ERC Consolidator Grant 681988 (CSP-Infinity). André Schrottenloher is supported by ERC-ADG-ALSTRONGCRYPTO (project 740972).

We show that the question whether a given universal sentence is equivalent to a universal Horn sentence can be decided effectively. This motivates the question whether one can also decide whether the class of finite models of a given universal Horn sentence has the JEP. The theories used by Braunfeld to prove the undecidability of JEP for given universal sentences are not universal Horn. We prove that JEP remains undecidable even if the given universal sentence is additionally Horn, and if the signature is binary. Our proof is based on the undecidability of universality of context-free languages and shorter than the argument of Braunfeld.

2 Preliminaries

The set $\{1, \dots, n\}$ is denoted by $[n]$. We use the bar notation for tuples; for a tuple \bar{a} indexed by a set I , the value of \bar{a} at the position $i \in I$ is denoted by $\bar{a}[i]$. For a function $f: A \rightarrow B$ and a tuple $\bar{a} \in A^k$, we set $f(\bar{a}) := (f(\bar{a}[1]), \dots, f(\bar{a}[k]))$. We always use capital Fraktur letters $\mathfrak{A}, \mathfrak{B}, \mathfrak{C}$, etc to denote structures, and the corresponding capital Roman letters A, B, C , etc for their domains.

Let τ be a finite relational signature. An *atomic τ -formula*, or τ -*atom* for short, is a formula of the form $R(x_1, \dots, x_n)$ for $R \in \tau$ and (not necessarily distinct) variables x_1, \dots, x_n . Note that in this text equality atoms, i.e., expressions of the form $x = y$, are not permitted. A class \mathcal{C} of relational τ -structures has the *joint embedding property (JEP)* if: for every pair $\mathfrak{B}_1, \mathfrak{B}_2 \in \mathcal{C}$, there exists $\mathfrak{C} \in \mathcal{C}$ and embeddings $f_i: \mathfrak{B}_i \hookrightarrow \mathfrak{C}$ ($i \in [2]$). It is closed under forming *disjoint unions* if $\mathfrak{C} \in \mathcal{C}$ and $f_i: \mathfrak{B}_i \hookrightarrow \mathfrak{C}$ can be chosen so that $f_1(B_1) \cup f_2(B_2) = C$, $f_1(B_1) \cap f_2(B_2) = \emptyset$, and $R^{\mathfrak{C}} = f_1(R^{\mathfrak{B}_1}) \cup f_2(R^{\mathfrak{B}_2})$ for every $R \in \tau$.

If \mathfrak{A} is a τ -structure and Φ is a τ -sentence, we write $\mathfrak{A} \models \Phi$ if \mathfrak{A} satisfies Φ . If Φ and Ψ are τ -sentences, we write $\Phi \models \Psi$ if every τ -structure that satisfies Φ also satisfies Ψ . For universal sentences, this is equivalent to the statement that every *finite* τ -structure that satisfies Φ also satisfies Ψ , by a standard application of the compactness theorem of first-order logic. If Φ is a τ -sentence, we denote the class of all finite models of Φ by $[[\Phi]]^{<\omega}$. Clearly, every universal formula can be written as $\forall x_1, \dots, x_n. \psi$ where ψ is quantifier-free and in conjunctive normal form. Conjunctions of ψ are also called *clauses*; clauses are disjunctions of atomic formulas and negated atomic formulas. A clause ϕ is a *weakening* of a clause ψ if every disjunct of ψ is also a disjunct of ϕ . We sometimes omit universal quantifiers in universal sentences (all first-order variables are then implicitly universally quantified). It is well-known that a class \mathcal{C} of finite τ -structures can be described by forbidding isomorphic copies of fixed finitely many finite τ -structures if and only if $\mathcal{C} = [[\Phi]]^{<\omega}$ for some universal τ -sentence Φ . With this observation, it is easy to see that Braunfeld [Bra19] proved undecidability for the following question. His proof is based on a reduction from the tiling problem.

INPUT: A universal sentence Φ in a finite relational signature.

QUESTION: Does $[[\Phi]]^{<\omega}$ have the JEP?

A universal sentence Φ is called *Horn* if its quantifier-free part is a conjunction of *Horn clauses*, i.e., disjunctions of positive or negative atomic formulas

each having at most one *positive* disjunct. Every Horn clause can be written equivalently as an implication

$$(\phi_1 \wedge \cdots \wedge \phi_n) \Rightarrow \phi_0$$

where ϕ_1, \dots, ϕ_n are atomic formulas (possibly $n = 0$), and ϕ_0 is either an atomic formula or \perp which stands for the empty disjunction. We refer to $\phi_1 \wedge \cdots \wedge \phi_n$ as the *premise*, and to ϕ_0 as the *conclusion* of the Horn clause. A Horn clause is called a *tautology* if the conclusion equals one of the conjuncts of the premise.

Definition 1. Let Φ be a universal Horn sentence and ψ a Horn clause, both in a fixed relational signature τ . An SLD-derivation of ψ from Φ of length k is a finite sequence of Horn clauses $\psi^0, \dots, \psi^k = \psi$ such that ψ^0 is a conjunct in Φ and each ψ^i ($1 \leq i \leq k$) is a (binary) resolvent of ψ^{i-1} and a conjunct ϕ^i from Φ , i.e., for some atomic formula ψ_j^{i-1} , we have

$$\frac{\overbrace{\psi_1^{i-1} \wedge \cdots \wedge \psi_j^{i-1} \wedge \cdots \wedge \psi_{n^{i-1}}^{i-1} \Rightarrow \psi_0^{i-1}}^{\psi^{i-1}} \quad \overbrace{\phi_1^i \wedge \cdots \wedge \phi_{m^i}^i \Rightarrow \psi_j^{i-1}}^{\phi^i}}{\underbrace{\psi_1^{i-1} \wedge \cdots \wedge \psi_{j-1}^{i-1} \wedge \phi_1^i \wedge \cdots \wedge \phi_{m^i}^i \wedge \psi_{j+1}^{i-1} \wedge \cdots \wedge \psi_{n^{i-1}}^{i-1} \Rightarrow \psi_0^{i-1}}_{\psi^i}}$$

There exists an SLD-deduction of ψ from Φ , written as $\Phi \vdash \psi$, if ψ is a tautology or a weakening of a Horn clause that has an SLD-derivation from Φ up to renaming variables.

The following theorem presents a fundamental property of universal Horn sentences.

Theorem 1 (Theorem 7.10 in [NCdW97]). Let Φ be a universal Horn sentence and ψ a Horn clause, both in a fixed signature τ . Then

$$\Phi \models \psi \quad \text{iff} \quad \Phi \vdash \psi.$$

3 Universal Horn Sentences and the JEP

This section deals with two separate topics. First, we discuss the semantical difference between universal sentences and universal Horn sentences which is described by the convexity condition. Second, we present some important observations about the JEP in the context of universal Horn sentences which we later use in the proof of our main result. Let τ be a relational signature.

A universal τ -sentence Φ is called *convex* if for all atomic τ -formulas $\phi_1, \dots, \phi_n, \psi_1, \dots, \psi_k$, if

$$\Phi \models \bigwedge_{i \leq n} \phi_i \Rightarrow \bigvee_{j \leq k} \psi_j$$

then there exists $j \leq k$ such that

$$\Phi \models \bigwedge_{i \leq n} \phi_i \Rightarrow \psi_j.$$

We say that Φ is *preserved in products* if for every non-empty family $(\mathfrak{A}_i)_{i \in I}$ of models of Φ , the direct product $\prod_I \mathfrak{A}_i$ is also a model of Φ . The following is well known; e.g., the direction (3) \Rightarrow (1) is Corollary 9.1.7 in [Hod93]. Since we also need (4) we provide a proof for the convenience of the reader.

Theorem 2. *Let Φ be a universal sentence with relational signature τ . Then the following are equivalent.*

1. Φ is convex;
2. Φ is equivalent to a universal Horn sentence;
3. Φ is preserved in products;
4. Φ is preserved in binary products of finite structures.

Proof. (1) \Rightarrow (2): We may assume that Φ is in prenex normal form and that its quantifier-free part ϕ is in conjunctive normal form. Every conjunct in ϕ is equivalent to an implication of the form $\bigwedge_{i \leq n} \phi_i \Rightarrow \bigvee_{j \leq k} \psi_j$. Since Φ is convex, we can replace the conjunct by $\bigwedge_{i \leq n} \phi_i \Rightarrow \psi_j$ for some $j \leq k$. In this way, Φ can be rewritten into an equivalent universal Horn sentence.

(2) \Rightarrow (3): Corollary 9.1.6 in [Hod93].

(3) \Rightarrow (4): This direction is trivial.

(4) \Rightarrow (1): Suppose that Φ has m variables and is not convex, i.e., $\Phi \models \bigwedge_{i \leq n} \phi_i \Rightarrow \bigvee_{j \leq k} \psi_j$ but, for every $j \leq k$, there exists a model \mathfrak{A}_j of Φ such that $\mathfrak{A}_j \models (\bigwedge_{i \leq n} \phi_i \wedge \neg \psi_j)(\bar{t}^j)$ for some tuple $\bar{t}^j \in A_j^m$. We may assume that each \mathfrak{A}_j is finite; otherwise we replace it with its substructure on the coordinates of \bar{t}^j while preserving the desired properties. For $\bar{s}^j := ((t_1^1, \dots, t_1^j), \dots, (t_m^1, \dots, t_m^j))$ we have

$$\prod_{i \leq j} \mathfrak{A}_i \models (\bigwedge_{i \leq n} \phi_i \wedge \bigwedge_{i \leq j} \neg \psi_i)(\bar{s}^j).$$

It follows by induction on $j \leq k$ that if Φ is preserved in binary products of finite structures, then $\prod_{i \leq j} \mathfrak{A}_i \models \Phi$. We then obtain a contradiction for $j = k$. \square

It is natural to consider the following question as a decision problem.

INPUT: A universal sentence Φ in a finite relational signature.

QUESTION: Is Φ equivalent to a universal Horn sentence?

In the context of the results of [BR21], one might also be interested in the variant where the input is restricted to those Φ for which $[[\Phi]]^{<\omega}$ has the JEP.

Proposition 1. *Deciding whether a given universal sentence Φ is equivalent to a universal Horn sentence is coNP-complete. The coNP-hardness holds even if the signature is limited to unary relation symbols and $[[\Phi]]^{<\omega}$ has the JEP.*

Proof. We first prove containment in coNP. For a given universal sentence Φ , let $\phi(x_1, \dots, x_n)$ be the quantifier-free part of Φ . If Φ is not equivalent to a universal Horn sentence, then, by Theorem 2, Φ has two finite models $\mathfrak{A}, \mathfrak{B}$ such that $\mathfrak{A} \times \mathfrak{B}$ is not a model of Φ . This means that there exists $\bar{t} \in (A \times B)^n$ such that $\mathfrak{A} \times \mathfrak{B} \not\models \phi(\bar{t})$. But then, by the definition of product of structures, there exist substructures \mathfrak{A}' of \mathfrak{A} and \mathfrak{B}' of \mathfrak{B} of size n with $\bar{t} \in (A' \times B')^n$ and $\mathfrak{A}' \times \mathfrak{B}' \not\models \phi(\bar{t})$. Since models of Φ are preserved under taking substructures, we have $\mathfrak{A}' \models \Phi$ and $\mathfrak{B}' \models \Phi$. Conversely, if there exist two models of Φ of size n whose product is not a model of Φ , then clearly Φ is not equivalent to a universal Horn sentence by Theorem 2. This means that there is a sound and complete algorithm for the complement of the original problem which guesses two models of Φ of size n and then confirms in time polynomial in n that their product is not a model of Φ .

The coNP-hardness can be shown by a reduction from the propositional unsatisfiability problem. From a given propositional formula ϕ we obtain a universal sentence Φ in the unary signature $\tau := \{C_1, \dots, C_n, L, R\}$ as follows. Let X_1, \dots, X_n be the propositional variables which appear in ϕ . We represent them with unary relation symbols C_1, \dots, C_n . We obtain the formula $\phi'(x)$ from ϕ by replacing each occurrence of X_i with the literal $C_i(x)$ where x is a single first-order variable. Now we define Φ as the τ -sentence

$$\forall x(\phi'(x) \Rightarrow (L(x) \vee R(x))).$$

If ϕ is unsatisfiable, then no τ -structure can satisfy ϕ' , which means that every τ -structure satisfies Φ , i.e., Φ is equivalent to an arbitrary tautology. Now suppose that ϕ is satisfiable. Let $f: \{X_1, \dots, X_n\} \rightarrow \{true, false\}$ be a satisfying assignment. We define τ -structures $\mathfrak{A}_1, \mathfrak{A}_2$ as follows:

- $A_1 = A_2 := \{x\}$;
- for every $i \in [n]$, $C_i^{\mathfrak{A}_1} = C_i^{\mathfrak{A}_2} := \{x\}$ if $f(X_i) = true$ and \emptyset otherwise;
- $L^{\mathfrak{A}_1} = R^{\mathfrak{A}_2} := \{x\}$ and $R^{\mathfrak{A}_1} = L^{\mathfrak{A}_2} := \emptyset$.

Then it is easy to see that $\mathfrak{A}_i \models \Phi$ for $i \in \{1, 2\}$ but $\mathfrak{A}_1 \times \mathfrak{A}_2 \not\models \Phi$. By Theorem 2, Φ is not equivalent to any universal Horn sentence. Clearly, $[\Phi]^{<\omega}$ has the JEP because $[\Phi]^{<\omega}$ is even closed under forming disjoint unions. \square

A Horn clause $\phi \Rightarrow \psi$ is called *connected* if the graph with vertex set $\{x_1, \dots, x_n\}$ where x_i and x_j form an edge if they appear jointly in a conjunct of ϕ , forms a connected graph on all variables occurring in $\phi \Rightarrow \psi$ in the usual graph-theoretic sense.

Proposition 2. *Let Φ be a conjunction of connected Horn clauses. Then $[\Phi]^{<\omega}$ has the JEP.*

Proof. The class $[\Phi]^{<\omega}$ is even closed under the formation of disjoint unions. \square

Definition 2 (Subsumption). *Let Φ be a universal Horn sentence over the relational signature τ , and let $\phi(\bar{x})$ and $\psi(\bar{x}, \bar{y})$ be conjunctions of atomic τ -formulas. We say that $\phi(\bar{x})$ subsumes $\psi(\bar{x}, \bar{y})$ with respect to Φ and write $\psi(\bar{x}, \bar{y}) \sqsubseteq_{\Phi} \phi(\bar{x})$.*

$\phi(\bar{x})$ if for every χ that is \perp or an atomic τ -formula with free variables among \bar{x}

$$\Phi \models \forall \bar{x}, \bar{y} (\psi(\bar{x}, \bar{y}) \Rightarrow \chi(\bar{x})) \quad \text{implies} \quad \Phi \models \forall \bar{x} (\phi(\bar{x}) \Rightarrow \chi(\bar{x})).$$

Note that subsumption can be tested using SLD-deduction (Theorem 1). In our proofs later it will be useful to take a proof-theoretic perspective on the amalgamation property.

Lemma 1. *Let Φ be a universal Horn sentence over the relational signature τ . Then the following are equivalent:*

1. $\llbracket \Phi \rrbracket^{<\omega}$ has the joint embedding property.
2. Suppose that $\phi_1(\bar{x}_1)$ and $\phi_2(\bar{x}_2)$ are conjunctions of atomic formulas such that $\phi_1(\bar{x}_1) \wedge \Phi$ is satisfiable for both $i \in [2]$. Then

$$\phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2) \sqsubseteq_{\Phi} \phi_1(\bar{x}_1).$$

Proof. (1) \Rightarrow (2): Let $\phi_1(\bar{x}_1)$ and $\phi_2(\bar{x}_2)$ be as in the first part of (2). Let $\Psi_1(\bar{x}_1)$ and $\Psi_2(\bar{x}_2)$ be the conjunctions of all atomic formulas implied by $\Phi \wedge \phi_1(\bar{x}_1)$ and $\Phi \wedge \phi_2(\bar{x}_2)$, respectively. By our assumption, $\Phi \wedge \Psi_1(\bar{x}_1)$ and $\Phi \wedge \Psi_2(\bar{x}_2)$ are both satisfiable. Define \mathfrak{B}_1 and \mathfrak{B}_2 as the structures whose domains consist of the variables $\{\bar{x}_{1[1]}, \dots\}$, and $\{\bar{x}_{2[1]}, \dots\}$, respectively, and where \bar{z} is a tuple of a relation for $R \in \tau$ if the conjunct $R(\bar{z})$ is contained in Ψ_1 or Ψ_2 , respectively. Note that, by construction, \mathfrak{B}_1 and \mathfrak{B}_2 satisfy every Horn clause in Φ . Since Φ is universal Horn, this implies that $\mathfrak{B}_1, \mathfrak{B}_2 \in \llbracket \Phi \rrbracket^{<\omega}$. Since $\llbracket \Phi \rrbracket^{<\omega}$ has the joint embedding property, there exists $\mathfrak{C} \in \llbracket \Phi \rrbracket^{<\omega}$ together with embeddings $f_i: \mathfrak{B}_i \hookrightarrow \mathfrak{C}$ for $i \in \{1, 2\}$. By the construction of \mathfrak{B}_1 and \mathfrak{B}_2 , it follows that $\Phi \not\models \forall \bar{x}_1, \bar{x}_2 (\phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2) \Rightarrow \perp)$. Let $\chi(\bar{x}_1)$ be an atomic τ -formula such that $\Phi \models \forall \bar{x}_1, \bar{x}_2 (\phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2) \Rightarrow \chi(\bar{x}_1))$. By the construction of \mathfrak{B}_1 and \mathfrak{B}_2 , and because f_1 and f_2 are homomorphisms, there exist a tuple \bar{z} over B_1 such that $\mathfrak{C} \models \chi(f_1(\bar{z}))$. Since f_1 is an embedding, we must also have $\mathfrak{B}_1 \models \chi(\bar{z})$. Thus, by the construction of \mathfrak{B}_1 and \mathfrak{B}_2 , it follows that $\Phi \models \forall \bar{x}_1 (\phi_1(\bar{x}_1) \Rightarrow \chi(\bar{x}_1))$.

(2) \Rightarrow (1): Let $\mathfrak{B}_1, \mathfrak{B}_2 \in \llbracket \Phi \rrbracket^{<\omega}$ be arbitrary. We construct a structure $\mathfrak{C} \in \llbracket \Phi \rrbracket^{<\omega}$ with $f_i: \mathfrak{B}_i \hookrightarrow \mathfrak{C}$ as follows. Without loss of generality we may assume that $B_1 \cap B_2 = \emptyset$. Let \bar{x} be a tuple of variables representing the elements of $B_1 \cap B_2$ in some order, and let $\phi_1(\bar{x}_1), \phi_2(\bar{x}_2)$ be the conjunctions of all atomic τ -formulas which hold in $\mathfrak{B}_1, \mathfrak{B}_2$, respectively. By construction, $\phi_i(\bar{x}_i) \wedge \Phi$ is satisfiable for both $i \in [2]$. Let $\Psi(\bar{x}_1, \bar{x}_2)$ be the conjunction of all atomic formulas implied by $\Phi \wedge \phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2)$. We claim that $\Phi \wedge \Psi$ is satisfiable: otherwise, $\Phi \models \forall \bar{x}_1, \bar{x}_2 (\phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2) \Rightarrow \perp)$, and then (2) implies that $\Phi \models \forall \bar{x}_1 (\phi_1(\bar{x}_1) \Rightarrow \perp)$, which is impossible since $\mathfrak{B}_1 \models \Phi$. Define \mathfrak{C} as the structure with domain $\{\bar{x}_{1[1]}, \dots, \bar{x}_{2[1]}, \dots\}$ and such a tuple $\bar{z} \in R^{\mathfrak{C}}$ if and only if Ψ contains the conjunct $R(\bar{z})$. For $i \in [2]$, let f_i be the identity map. We claim that f_i is an embedding from \mathfrak{B}_i to \mathfrak{C} . It is clear from the construction of \mathfrak{C} that f_i is a homomorphism. Suppose for contradiction that there exists $R \in \tau$ and a tuple \bar{z} over B_i such that $\bar{z} \notin R_i^{\mathfrak{B}_i}$ while $f_i(\bar{z}) \in R^{\mathfrak{C}}$. For the sake of notation, we assume that $i = 1$; the case that $i = 2$ can be shown analogously. Note that the

construction of \mathfrak{C} implies that $\Phi \models \forall \bar{x}_1, \bar{x}_2, (\phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2) \Rightarrow R(\bar{z}))$. Then (2) implies that $\Phi \models \forall \bar{x}_1 (\phi_1(\bar{x}_1) \Rightarrow R(\bar{z}))$, a contradiction to $\mathfrak{B}_1 \in \llbracket \Phi \rrbracket^{<\omega}$. Thus, f_i is an embedding from \mathfrak{B}_i to \mathfrak{C} . This concludes the proof of the joint embedding property. \square

We mention in passing that (1) and (2) are equivalent even if equality is permitted in Φ .

4 Undecidability of the JEP

In this section we prove that the problem of deciding whether the class of all finite models of a given universal Horn sentence Φ has the joint embedding property is undecidable. Our proof is based on a reduction from the problem of deciding the universality of a given context-free grammar.

A context-free grammar is a 4-tuple $G = (N, \Sigma, P, S)$ where

- N is a finite set of *non-terminal symbols*,
- Σ is a finite set of *terminal symbols*,
- P is a finite set of *production rules* of the form $A \rightarrow w$ where $A \in N$ and $w \in (N \cup \Sigma)^*$,
- $S \in N$ is the *start symbol*.

For $u, v \in (N \cup \Sigma)^*$ we write $u \rightarrow_G v$ if there are $x, y \in (N \cup \Sigma)^*$ and $(A \rightarrow w) \in P$ such that $u = xAy$ and $v = xwy$. The transitive closure of \rightarrow_G is denoted by \rightarrow_G^* . The *language of G* is $L(G) := \{w \in \Sigma^* \mid S \rightarrow_G^* w\}$. Note that with this definition the *empty word*, i.e., the word ϵ of length 0, can never be an element of $L(G)$; some authors use a modified definition that also allows rules that derive ϵ , but for our purposes the difference is not essential.

The idea of the reduction is to compute from a given context-free grammar G a universal Horn sentence which consists of two parts, Φ_1 and Φ_2 : the sentence Φ_2 does not depend on G and entails many Horn clauses witnessing failure of the JEP via Lemma 1; the sentence Φ_1 can be computed efficiently from G and is such that $\llbracket \Phi_1 \rrbracket^{<\omega}$ is closed under the formation of disjoint unions and prevents all the failures of the JEP of $\llbracket \Phi_1 \wedge \Phi_2 \rrbracket^{<\omega}$ if and only if G is universal, i.e., $L(G) = \Sigma^*$.

Theorem 3. *For a given universal Horn sentence Φ the question whether $\llbracket \Phi \rrbracket^{<\omega}$ has the JEP is undecidable even if the signature is limited to at most binary relation symbols.*

Proof. The universality problem for context-free grammars is known to be undecidable [Har78] (Lemma 8.4.2, page 259). In this proof, we only consider context-free grammars without production rules of the form $A \rightarrow b$ such that w contains the start symbol S . This is without loss of generality as every context-free grammar can be transformed into such form in polynomial time [Har78]. We also assume that $(A \rightarrow A) \in P$ for every $A \in N$. Note that this assumption does not influence $L(G)$ at all.

Encoding context-free grammars into ages of relational structures. Let τ_1 be the signature that contains the unary symbols I and T , the binary symbol E , and the binary relation symbol R_a for every element $a \in N \cup \Sigma$. Let Φ_1 be the universal Horn sentence that contains for every $(A \rightarrow a_1 \dots a_n) \in P$ with $A \neq S$ the Horn clause

$$\bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1}) \Rightarrow R_A(y, x_{i+1}), \quad (1)$$

and, for every $(S \rightarrow a_1 \dots a_n) \in P$, the Horn clause

$$I(x_1) \wedge T(x_{n+1}) \wedge \bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1}) \Rightarrow \perp. \quad (2)$$

Note that each conjunct of Φ_1 is connected, which means that $\llbracket \Phi_1 \rrbracket^{<\omega}$ has the JEP by Proposition 2. The following correspondence can be shown via a straightforward induction.

Claim 1. *For every $w = a_1 \dots a_n \in (N \cup \Sigma)^*$ and every $A \in N \setminus \{S\}$*

$$A \rightarrow_G^* w \quad \text{if and only if} \quad \Phi_1 \models \forall x_1, \dots, x_{n+1} \left(\bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1}) \Rightarrow R_A(x_1, x_{n+1}) \right). \quad (3)$$

For $A = S$, we have

$$S \rightarrow_G^* w \quad \text{if and only if} \quad \Phi_1 \models \forall x_1, \dots, x_{n+1} \left(I(x_1) \wedge T(x_{n+1}) \wedge \bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1}) \Rightarrow \perp \right). \quad (4)$$

Proof. “ \Rightarrow ”: Suppose that $A \rightarrow_G^* a_1 \dots a_n$ for $A \in N$. Then there is a path in \rightarrow_G from A to $a_1 \dots a_n$ of length $\lambda \geq 1$. We prove the statements by induction on λ .

In the *induction base* $\lambda = 1$ we have $(A \rightarrow a) \in P$ in which case (3) is a conjunct of Φ_1 .

In the *induction step* $\lambda \rightarrow \lambda + 1$, we assume that the claim holds for all paths of length $\leq \lambda$, and that there exists a path of length $\lambda + 1$ from A to $a_1 \dots a_n$, i.e., there exists $(A \rightarrow a_1 \dots a_{k-1} B a_{\ell+1} \dots a_n) \in P$ and a path of length λ from B to $a_k \dots a_\ell$. By the induction hypothesis, after renaming of variables we have that

$$\Phi_1 \models \forall x_k, \dots, x_{\ell+1} \left(\bigwedge_{i \in \{k, \dots, \ell\}} R_{a_i}(x_i, x_{i+1}) \Rightarrow R_B(x_k, x_{\ell+1}) \right). \quad (5)$$

By the construction of Φ_1 , if $A \neq S$, after renaming of variables we also have that

$$\begin{aligned} \Phi_1 \models \forall x_1, \dots, x_k, x_{\ell+1}, \dots, x_{n+1} \left(\bigwedge_{i \in [k-1]} R_{a_i}(x_i, x_{i+1}) \right. \\ \left. \wedge R_B(x_k, x_{\ell+1}) \wedge \bigwedge_{i \in \{\ell+1, \dots, n\}} R_{a_i}(x_i, x_{i+1}) \Rightarrow R_A(x_1, x_{n+1}) \right). \quad (6) \end{aligned}$$

We can now apply an SLD derivation step to (6) with (5) to obtain (3). The argument for $A = S$ is analogous.

“ \Leftarrow ”: Suppose that $\Phi_1 \models (3)$. By Theorem 1, there is an SLD-deduction of (3) from Φ_1 . If (3) is a tautology, then $n = 1$ and $a_1 = A$ in which case the statement is true because $(A \rightarrow A) \in P$ for every $A \in N$. Otherwise, (3) is a weakening of a Horn clause ψ that has an SLD-derivation from Φ_1 , and we prove the claim by induction on the length λ of a shortest possible SLD-derivation for ψ .

In the *base case* $\lambda = 0$, ψ must be a conjunct of Φ_1 . By the construction of Φ_1 , we get that $(A \rightarrow a_1 \dots a_n) \in P$ and thus $A \rightarrow_G^* a_1 \dots a_n$.

In the *induction step* $\lambda \rightarrow \lambda + 1$, we assume that the claim holds if ψ has an SLD-derivation of length $\leq \lambda$. Suppose that ψ requires an SLD-derivation of length $\lambda + 1$. By the construction of Φ_1 , there must exist $(B, a_k \dots a_\ell) \in P$ such that Φ_1 contains a conjunct of the form (1) that is used in the last step in a shortest possible SLD-derivation of ψ . Moreover, there exists an SLD-deduction of

$$\begin{aligned} & \left(\bigwedge_{i \in [k-1]} R_{a_i}(x_i, x_{i+1}) \right) \wedge R_B(x_k, x_{\ell+1}) \\ & \wedge \bigwedge_{i \in [n] \setminus [\ell]} R_{a_i}(x_i, x_{i+1}) \Rightarrow R_A(x_1, x_{n+1}) \end{aligned} \quad (7)$$

from Φ_1 of length $\leq \lambda$. By the induction hypothesis, (7) is equivalent to $A \rightarrow_G^* a_1 \dots a_{k-1} B a_{\ell+1} \dots a_{n-1}$. Therefore, $A \rightarrow_G^* a_1, \dots, a_n$. The argument for $A = S$ is analogous. \square

Creating candidates for failure of the JEP. Let τ_2 be the signature which contains all symbols from τ_1 except for the ones coming from N and additionally the unary symbol U and the binary symbol Q . The sentence Φ_2 consists of the following Horn clauses for every $a \in \Sigma$:

$$U(y) \wedge I(x_1) \Rightarrow Q(y, x_1) \quad (8)$$

$$U(y) \wedge Q(y, x_i) \wedge R_a(x_i, x_{i+1}) \Rightarrow Q(y, x_{i+1}) \quad (9)$$

$$U(y) \wedge Q(y, x_n) \wedge R_a(x_n, x_{n+1}) \wedge T(x_{n+1}) \Rightarrow \perp \quad (10)$$

The proof of the following claim is straightforward and left to the reader.

Claim 2. *Let $\phi(\bar{x})$ be a conjunction of $(\tau_2 \setminus \{Q\})$ -atoms. Then*

$$\Phi_2 \models \forall \bar{x} (\phi(\bar{x}) \Rightarrow \perp)$$

if and only if there exist $n \in \mathbb{N}_{>0}$ and $a_1, \dots, a_n \in \Sigma$ such that ϕ has a subformula of the following form:

$$U(y) \wedge I(x_1) \wedge T(x_{n+1}) \wedge \bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1}) \quad (11)$$

Now we are ready for the main part of the proof. We set $\Phi := \Phi_1 \wedge \Phi_2$.

Claim 3. $[[\Phi]]^{<\omega}$ has the JEP if and only if $L(G) = \Sigma^*$.

Proof. “ \Rightarrow ”: Suppose that $[[\Phi]]^{<\omega}$ has the JEP. Let $a_1 \dots a_n \in \Sigma^*$. Consider the formulas $\phi_1(y)$ and $\phi_2(x_1, \dots, x_{n+1})$ given by

$$\begin{aligned} \phi_1 &:= U(y) \\ \text{and } \phi_2 &:= I(x_1) \wedge T(x_{n+1}) \wedge \bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1}). \end{aligned}$$

By Claim 2, we have

$$\Phi_2 \models \forall x_1, \dots, x_{n+1}, y (\phi_1 \wedge \phi_2 \Rightarrow \perp). \quad (12)$$

Since $[[\Phi]]^{<\omega}$ has the JEP, by Lemma 1(2), $\Phi \wedge \phi_1$ or $\Phi \wedge \phi_2$ is not satisfiable. Note that every Horn clause in Φ_1 contains an R_a -atom for some $a \in N \cup \Sigma$ in its premise, and ϕ_1 contains none. Also note that every Horn clause in Φ_2 contains an I -atom or an R_a -atom for some $a \in N \cup \Sigma$ in its premise, and ϕ_1 contains none. Thus, by Theorem 1, $\Phi \wedge \phi_1$ is satisfiable. It follows that

$$\Phi \models \forall x_1, \dots, x_{n+1} (\phi_2 \Rightarrow \perp).$$

Since every Horn clause in Φ_2 contains a U -atom in its premise and ϕ_2 contains none, by Theorem 1, we have that

$$\Phi_1 \models \forall x_1, \dots, x_{n+1} (\phi_2 \Rightarrow \perp).$$

By Claim 1, $a_1 \dots a_n \in L(G)$ and we are done.

“ \Leftarrow ”: We prove the contrapositive and assume that $[[\Phi]]^{<\omega}$ does not have the JEP. Then there exists a counterexample to Lemma 1(2), i.e., there exists a Horn clause ψ of the following form

$$\phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2) \Rightarrow \chi$$

where χ is either \perp or an atomic τ -formula with free variables among \bar{x}_1 such that

$$\Phi \models \forall \bar{x}_1, \bar{x}_2 (\phi_1 \wedge \phi_2 \Rightarrow \chi) \quad (13)$$

$$\text{and } \Phi \not\models \forall \bar{x}_1 (\phi_1 \Rightarrow \chi) \quad (14)$$

and, for both $i \in \{1, 2\}$,

$$\Phi \not\models \forall \bar{x}_i (\phi_i \Rightarrow \perp). \quad (15)$$

We choose ψ minimal with respect to the number of its atomic subformulas.

Our proof strategy is as follows. First we show that ψ encodes a single word $w \in \Sigma^*$ in the sense of Claim 2. Then we show that the word w may not be contained in $L(G)$, because otherwise a part of the counterexample would encode w in the sense of Claim 1 which would lead to a contradiction.

For a conjunction θ of τ -atoms, we denote by θ' the conjunction obtained by removing all $(\tau_2 \setminus \tau_1)$ -atoms from ψ .

Observation 1. $\Phi_2 \models \psi$ and ψ only contains symbols from τ_2 .

Proof (of Observation 1). By Theorem 1 and (13) we have $\Phi \vdash \psi$. Since $Q \notin \tau_1$ and $R_A \notin \tau_2$ for every $A \in N$, by the construction of Φ_1 and Φ_2 we have either $\Phi_1 \vdash \psi$ or $\Phi_2 \vdash \psi$. Suppose for contradiction that $\Phi_1 \vdash \psi$. Note that $\chi(\bar{x}_1)$ cannot be a subformula of $\phi_1(\bar{x}_1)$, by (14). Also note that $\chi(\bar{x}_1)$ cannot be a subformula of $\phi_2(\bar{x}_2)$ as they do not share any variables. Hence, $\chi(\bar{x}_1)$ is not a subformula of $\phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2)$. Thus, it follows from $\Phi_1 \vdash \psi$ and the fact that Φ_1 only contains τ_1 -atoms that

$$\Phi_1 \models \forall \bar{x}_1, \bar{x}_2 (\phi'_1 \wedge \phi'_2 \Rightarrow \chi). \quad (16)$$

By the construction of Φ_1 , (16) and Theorem 1 imply that either χ is an R_A -atom for some $A \in N$, or χ is \perp . In both cases, (15), (16), and (14) witness that $\llbracket \Phi_1 \rrbracket^{<\omega}$ does not have JEP through an application of Lemma 1. But this is in contradiction to Proposition 2. Thus, $\Phi_1 \vdash \psi$ does not hold, and $\Phi_2 \vdash \psi$ holds instead. In this case, $\phi_1 \wedge \phi_2$ can only contain symbols from τ_2 , since we could otherwise remove all $(\tau_1 \setminus \tau_2)$ -atoms and get a contradiction to the minimality of ψ . Similarly as above, by the construction of Φ_2 , Theorem 1 and $\Phi_2 \vdash \psi$ imply that either χ is a Q -atom, or χ equals \perp . Thus, ψ only contains τ_2 -atoms. \square

Observation 2. *The formula $\phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2)$ does not contain any Q -atom and χ equals \perp .*

Proof (of Observation 2). Recall that, by Observation 1 and Theorem 1, ψ has an SLD-deduction from Φ_2 . By (13) and (14), ψ cannot be a tautology. By the minimality of ψ , we may assume that there exists an SLD-derivation of ψ from Φ_2 . Consider any SLD-derivation ψ^0, \dots, ψ^k of ψ from Φ_2 . Note that, by the construction of Φ_2 , for every $i \in [k]$, if there exists a variable y in ψ^{i-1} such that

- i. every Q -atom contains y in its first argument,
- ii. every U -atom contains y in its only argument,

then ψ^i also satisfies (i) and (ii) for the same variable y . Since every possible choice of ψ^0 from Φ_2 initially satisfies these two conditions, it follows via induction that (i) and (ii) must hold for $\psi = \psi^k$ for some y . Also note that (8) is the only Horn clause in Φ_2 that is not connected, but the lack of connectivity is only because the variable y satisfying (i) and (ii) is isolated from the remaining variables.

We claim that ψ^0 is of the form (10). Otherwise, ψ^0 is of the form (9) or (8), in which case the conclusion of ψ is a Q -atom. Since there exists a variable y that satisfies (i) for ψ and, by (13), the conclusion of ψ may only contain variables from \bar{x}_1 , every Q -atom that can be derived from $\Phi_2 \wedge \phi_1(\bar{x}_1) \wedge \phi_2(\bar{x}_2)$ may only contain variables from \bar{x}_1 and can already be derived from $\Phi_2 \wedge \phi_1(\bar{x}_1)$. The claim implies that χ equals \perp .

It follows by induction that the premise of $\psi = \psi^k$ is connected except perhaps for the variable y which might be isolated from the remaining variables. Since y satisfies (i) for ψ , if ψ contains any Q -atom in the premise, then ψ is connected. But then we get a contradiction to (14). Thus, ψ does not contain any Q -atoms at all. \square

As a consequence of Observation 1 and Observation 2 we have that $\bar{\Phi}_2 \models \psi$ where ψ is of the form $\phi_1 \wedge \phi_2 \Rightarrow \perp$ and ϕ_1, ϕ_2 are conjunctions of atomic $\tau_2 \setminus \{Q\}$ -formulas. Therefore, Claim 2 implies that there exist $a_1, \dots, a_n \in \Sigma$ such that $\phi_1 \wedge \phi_2$ is of the form

$$U(y) \wedge I(x_1) \wedge T(x_{n+1}) \wedge \bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1}) \quad (17)$$

where the variables need not all be distinct. More specifically,

- $\phi_1(\bar{x}_1)$ equals $U(y)$ and
- $\phi_2(\bar{x}_2)$ equals $I(x_1) \wedge T(x_{n+1}) \wedge \bigwedge_{i \in [n]} R_{a_i}(x_i, x_{i+1})$.

Note that, if $L(G) = \Sigma^*$, then Claim 1 implies that

$$\bar{\Phi}_1 \models \forall x_1, \dots, x_{n+1} (\phi_2 \Rightarrow \perp). \quad (18)$$

If some variables among x_1, \dots, x_n are identified in (17), then we still have (18) even if we perform the same identification of variables. But then we get a contradiction to (15). Thus, $L(G) \neq \Sigma^*$, which concludes the proof of Claim 3. \square

We have thus found a reduction from the undecidable universality problem for G to the decidability problem of the JEP for $[[\bar{\Phi}]]^{<\omega}$; note that $\bar{\Phi}$ is universal Horn and can be computed from G in polynomial time. \square

5 Open Problems

An important open question about universal sentences over relational signatures is the decidability of the *amalgamation property* (AP), which is a strong form of the joint embedding property, and which is of fundamental importance in constraint satisfaction [Bod21]. Unlike the JEP, the AP is decidable if all relation symbols in the signature are at most binary (see, e.g., [KL87, BKS20]). For general relational signatures, we ask the following question.

Question 1. Is it decidable whether the class of finite models of a given universal Horn sentence $\bar{\Phi}$ has the amalgamation property?

Note that if the class of finite models of $\bar{\Phi}$ does have amalgamation, then we can compute an amalgam syntactically by computing implied atomic formulas just as in the proof of Lemma 1.

References

- BKS20. M. Bodirsky, S. Knäuer, and F. Starke. ASNP: A tame fragment of existential second-order logic. In Marcella Anselmo, Gianluca Della Vedova, Florin Manea, and Arno Pauly, editors, *Beyond the Horizon of Computability - 16th Conference on Computability in Europe, CiE 2020, Fisciano, Italy, June 29 - July 3, 2020, Proceedings*, volume 12098 of *Lecture Notes in Computer Science*, pages 149–162. Springer, 2020.

- Bod21. M. Bodirsky. *Complexity of Infinite-Domain Constraint Satisfaction*. Lecture Notes in Logic. Cambridge University Press, 2021.
- BR21. F. Baader and J. Rydval. An algebraic view on p-admissible concrete domains for lightweight description logics, 2021. To appear in JELIA 2021, available at <https://tu-dresden.de/inf/lat/reports#BaRy-LTCS-20-10>.
- Bra19. S. Braunfeld. The undecidability of joint embedding and joint homomorphism for hereditary graph classes. *Discrete Mathematics & Theoretical Computer Science*, 21, 2019. Preprint available at arXiv:1903.11932.
- Har78. M. A. Harrison. *Introduction to Formal Language Theory*. Addison-Wesley Longman Publishing Co., Inc., USA, 1st edition, 1978.
- Hod93. W. Hodges. *Model theory*. Cambridge University Press, 1993.
- KL87. J. Knight and A.H. Lachlan. Shrinking, stretching and codes for homogeneous structures. In *Classification Theory (Chicago, IL, 1985)*, Lecture Notes in Math., 1292, pages 192–228. Springer, Berlin-New York, 1987.
- NCdW97. S.-H. Nienhuys-Cheng and R. de Wolf. *Foundations of Inductive Logic Programming*, volume 1228 of *Lecture Notes in Computer Science*. Springer, 1997.