

Analytical investigation of pre-inflationary effects in the primordial power spectrum: From General Relativity to hybrid Loop Quantum Cosmology

Beatriz Elizaga Navascués,^{a,1} Guillermo A. Mena Marugán^b

^aJSPS International Research Fellow, Department of Physics, Waseda University,
3-4-1 Okubo, Shinjuku-ku, 169-8555 Tokyo, Japan

^bInstituto de Estructura de la Materia, IEM-CSIC,
Serrano 121, 28006 Madrid, Spain

E-mail: w.iac20060@kurenai.waseda.jp, mena@iem.cfmac.csic.es

Abstract. This work is an analytical study of the main differences between classical inflationary effects of a fast-roll regime and imprints of hybrid Loop Quantum Cosmology (LQC) on the primordial power spectrum. Effective LQC solutions of phenomenological interest typically contain a classical period of kinetic dominance prior to slow-roll inflation, with consequences on the power spectrum that here we try and tell apart from those of phenomena occurring in the vicinity of a bounce in hybrid LQC. For this comparison, we use the same criterion for the choice of a vacuum state of the primordial perturbations, both in the context of relativistic cosmology and on the LQC cosmological background. As a first approximation to the problem, our study ignores the influence of a quasi-de Sitter evolution (rather than de Sitter) in the slow-roll inflationary regime, and of a typically short transition between the kinetically dominated era and the slow-roll period. Our results show that, while the two studied types of effects lead to a drastic power suppression in the infrared sector of the spectrum, the mode scale at which this happens is much larger for LQC than for classical relativistic cosmology, the corresponding scales being related to the spacetime curvature at the bounce and at the onset of inflation, respectively.

Keywords: Loop quantum cosmology, vacuum state, cosmological perturbations, primordial power spectrum, short-lived inflation

¹Corresponding author.

Contents

1	Introduction	1
2	Perturbations in de Sitter inflation matching a scalar field pre-inflationary phase	4
2.1	The approximate cosmological background	5
2.2	Dynamics of gauge invariants: Continuous solutions	6
2.3	Primordial power spectrum: Non-oscillating behavior	8
3	Approximate mass for the perturbations in hybrid LQC	9
3.1	Approximate background solutions	10
3.2	Pöschl-Teller approximation	12
3.3	Approximate dynamics of gauge invariants: Continuous solutions	13
4	Short-lived inflation vs. loop quantum bounce effects	14
4.1	Choice of vacuum state	14
4.2	Power spectrum for short-lived inflation in GR	16
4.3	Power spectrum for hybrid LQC	19
5	Discussion	23
A	Fixing a vacuum state with the asymptotic expansion of h_k	25

1 Introduction

The Cosmic Microwave Background (CMB) is a useful observational window to the earliest stages of the Universe. Its angular power spectrum of temperature (and polarization) anisotropies can be explained remarkably well starting from a primordial slow-roll inflationary epoch, using standard cosmological perturbation theory in General Relativity (GR) [1, 2]. Despite this success, both WMAP [3] and, more recently, the Planck mission [4, 5] have reported an anomalous lack of power for the lowest multipoles of the observable spectrum, namely for the correlations between points in the sky separated by the largest angles. Even though this sector of the power spectrum is strongly affected by cosmic variance, this anomalous behavior may open the possibility for new physics, beyond the otherwise successful inflationary paradigm.

During last years, a considerable number of works have proposed a primordial origin for this anomalous lack of power in the CMB spectrum, without giving up on a relativistic description of the early Universe (according to GR) nor on its spatial flatness [6–18]. Instead, these investigations often relax the hypothesis that no information on the pre-inflationary Universe survives the slow-roll epoch (this point of view has been reinforced recently e.g. through the study of the path integral formulation of quantum cosmology [19]). Such investigations often consider scenarios in which the slow-roll period is just long enough to satisfy the observational constraints on the necessary e-folds [20–22]. In many cases, this situation is attained by considering inflationary models with multiple, different stages. Among these, perhaps the simplest ones are those that, driven by a single scalar field, contain what has

been called a phase of fast-roll inflation (see e.g. refs. [6, 9, 10, 18]). They describe a cosmological evolution with a (pre-inflationary) epoch in which the kinetic energy of the scalar field dominates over its potential energy, followed by a short period of slow-roll inflation. With appropriate choices for the vacuum state of the cosmological perturbations, the primordial power spectrum resulting from these models displays a strong suppression, which can be interpreted as if there existed an effective infrared cutoff.

Another possibility to address the presence of anomalies in the CMB arises naturally in the context of quantum cosmology. As one approaches the Big-Bang singularity and the spacetime curvature grows unboundedly large, our conventional knowledge of physics indicates that the classical relativistic description eventually breaks down, and we expect that it should be replaced with an underlying quantum theory. The modifications coming from quantum geometry effects in the pre-inflationary evolution of the Universe could naturally involve phenomena leading e.g. to power suppression in the CMB. In this framework, an interesting scenario is provided by Loop Quantum Cosmology (LQC) [23–25]. Apart from developing a non-perturbative quantization of cosmological spacetimes, LQC provides a mechanism for the resolution of the Big-Bang singularity into a bounce of quantum origin [26, 27]. Certain Gaussian states of the theory effectively describe cosmological spacetimes that, evolving as in standard GR cosmology for low densities, depart from the classical trajectories when the energy density approaches the Planck scale and then avoid the singularity [25, 28, 29].

In the presence of an inflaton field, there is a class of especially appealing solutions in the aforementioned effective LQC regime. These are solutions with slow-roll inflation for which the amount of expansion from the bounce to the end of inflation is such that the perturbations with largest observable wavelengths (and only them) felt the quantum curvature effects near the bounce. Remarkably, these solutions in effective LQC display, shortly after the bounce, a classical period that mimicks fast-roll inflation [30–32]. Namely, a pre-inflationary regime of kinetic dominance that eventually leads to a period of slow-roll inflation with just enough e-folds. Focusing on this type of solutions, a number of studies have predicted power suppression in the primordial spectrum that alleviates the CMB anomalies [31–34]. In order to guarantee the potential falsifiability of these models, it then becomes very important to be able to distinguish whether such power suppression is due to genuine LQC effects on the evolution of the perturbations or to fast-roll effects (or to a combination of both). Unfortunately, all the studies of power suppression in LQC have so far relied on numerical computations, which fail to tell apart in detail the actual processes (classical or quantum) underlying their predictions on the modifications of the spectrum.

The aim of this work is to develop an analytical understanding of how the different types of pre-inflationary effects that coexist in the LQC evolution of scalar and tensor perturbations become reflected in the primordial power spectrum. More specifically, in this article we will concentrate our discussion exclusively on a concrete proposal for the description of cosmological perturbations in LQC: the hybrid approach [35–38]. This is one of the proposals from which suppression for large wavelengths of the primordial power spectrum has been reported. An important reason to focus our analysis on hybrid LQC is that it provides a manageable and well-motivated framework to try and disentangle the ultimate origin of the different phenomena that affect the power spectra. Nevertheless, we believe that the techniques and intuition gained with this analysis can be extended to other theoretical frameworks, facilitating a better understanding of the observational consequences of pure relativistic effects and trans-Planckian physics in the very early Universe (even beyond the context of LQC). The dynamical equations for the perturbations in hybrid LQC are derived from a quantization of

the full cosmological system that combines loop quantization techniques for the spatially homogeneous degrees of freedom (the cosmological background) with a Fock quantization for the perturbations. These dynamical equations have the same hyperbolic structure as in leading-order cosmological perturbation theory for GR, but display a time-dependent mass term that differs from the classical one in the vicinity of the bounce [39]. We will analytically study these equations for background solutions of effective LQC in situations of phenomenological interest, on the one hand, and for their counterpart given by fast-roll inflationary solutions in GR cosmology, on the other hand. Then, using the same choice of vacuum state in both cases (a choice that is motivated by physical considerations), we will compare the resulting power spectra and study the differences, which can be rooted in the existence of two distinct scales in the considered situations, namely, the spacetime curvature at the bounce, with strong LQC effects, and the curvature at the onset of slow-roll inflation, which is ruled by GR cosmology.

In order to carry out this analytical study, we will need to approximate in a convenient form the time-dependent mass that appears in the dynamical equations of the perturbations. For that purpose, we will model both the GR cosmological solution with fast-roll inflation, and the part of the LQC background with a similar classical behavior, as a cosmological spacetime minimally coupled to a massless scalar field that instantaneously transitions to a de Sitter spacetime. In this respect, as a first approximation to the problem, we will ignore any effect produced during the passage from kinetic dominance to slow-roll inflation. Furthermore, we will also neglect the departures from an exact de Sitter expansion during slow-roll inflation, which depend on the specific inflaton potential that one is considering. As a consequence, our study will not provide any reliable detail about slow-roll observables such as the spectral index or the tensor-to-scalar ratio. On the other hand, in the vicinity of the LQC bounce, we will approximate the time-dependent mass as certain Pöschl-Teller potential, in a way that improves similar approximations made in the literature of (hybrid) LQC [40] in what respects the time interval of applicability. Although limiting the extent and accuracy of our physical investigation, the simplicity of these approximations will actually allow us to deduce and clearly explain the main differences between the primordial power spectra obtained with pure GR cosmology and with effective hybrid LQC.

The structure of the rest of this paper is the following. In section 2 we approximate and discuss analytically the propagation of cosmological perturbations on a single-field cosmology with fast-roll inflation within GR. Then, we define the primordial power spectrum and comment on some of its properties. The general strategy that we follow in this section is the same as in ref. [6], but maintaining the generality in the choice of vacuum state for the perturbations. In section 3 we revisit the properties of the effective LQC solutions of phenomenological interest, and make a Pöschl-Teller approximation in the vicinity of the bounce to approximate the dynamics of the cosmological perturbations in the hybrid approach. We then explain in which sense this approximation improves similar ones in the literature [40]. In section 4 we set a common criterion for the choice of vacuum state, with original motivation coming from the hybrid quantization process itself and the demand of non-oscillatory properties on the power spectrum. We use this criterion to select a state for the cosmological perturbations, both in the GR cosmological model and in hybrid LQC. We also analytically compute and compare the resulting power spectra. Finally, in section 5 we discuss our results and their prospective applications. Throughout the paper we work in Planck units, setting $G = c = \hbar = 1$.

2 Perturbations in de Sitter inflation matching a scalar field pre-inflationary phase

Let us set up the theoretical model with which we want to approximate a classical, single-field driven, inflationary universe preceded by a classical epoch of decelerated expansion, in the presence of inhomogeneous perturbations. This cosmological evolution takes place in an interval $(\eta_0, \eta_{\text{end}}]$ of the conformal time η , where η_{end} is the instant when slow-roll inflation ends. For the moment, let η_0 be an unspecified choice of initial time. Ignoring any perturbative backreaction, the equations that rule the evolution of the considered background cosmology in conformal time are

$$\left(\frac{a'}{a}\right)^2 = \frac{8\pi}{3}a^2\rho, \quad \frac{a''}{a} = \frac{4\pi}{3}a^2\rho - 4\pi a^2P, \quad (2.1)$$

where a is the cosmological scale factor, the prime denotes the derivative with respect to the conformal time, and ρ and P are the respective energy density and pressure of the homogeneous inflaton field ϕ with potential $V(\phi)$, namely

$$\rho = \frac{1}{2}\left(\frac{\phi'}{a}\right)^2 + V(\phi), \quad P = \rho - 2V(\phi). \quad (2.2)$$

For concreteness, in this work we focus the discussion on the case of a quadratic potential $V(\phi) = m^2\phi^2/2$, where m is the inflaton mass. However, since the details of this potential during inflation will be completely ignored within our approximations, we do not expect our results to be extremely sensitive to this choice.

In the context of dominant-order perturbation theory, the equations of motion for the gauge-invariant perturbations on the considered background are

$$v_{\vec{k}}^{(r)''} + \left[k^2 + s_{\text{GR}}^{(r)}\right]v_{\vec{k}}^{(r)} = 0, \quad k = |\vec{k}|, \quad (2.3)$$

where $v_{\vec{k}}^{(r)}$ generically denotes either the Fourier coefficient of the mode with wavevector \vec{k} of the Mukhanov-Sasaki field [41–43] (field for which we employ the notation $r = s$) or the respective coefficient of the tensor perturbations (case for which $r = t$). The function $s_{\text{GR}}^{(r)}$ plays the role of a time-dependent mass for the respective Mukhanov-Sasaki and tensor perturbations. Explicitly, it is well-known that [44]

$$s_{\text{GR}}^{(s)} = -\frac{z''}{z}, \quad s_{\text{GR}}^{(t)} = -\frac{a''}{a}, \quad (2.4)$$

where $z = a^2\phi'/a'$. Even for times when $\phi' = 0$, the Mukhanov-Sasaki mass is non-singular and can be expressed in terms of the tensor one in a general relation of the form [39]

$$s_{\text{GR}}^{(s)} = s_{\text{GR}}^{(t)} + \mathcal{U}_{\text{GR}}, \quad (2.5)$$

where \mathcal{U}_{GR} is a linear combination (with time-dependent coefficients) of the potential $V(\phi)$, its square, and its first and second derivatives with respect to ϕ . We do not need to show the explicit expression of \mathcal{U}_{GR} here, but only to notice that it is a very small term when $V(\phi)$ and its two first derivatives are negligible compared to the kinetic component of the energy density ρ .

2.1 The approximate cosmological background

Among the solutions of eq. (2.1) for the cosmological background in the period $(\eta_0, \eta_{\text{end}}]$, we restrict our attention to those with a subinterval (η_0, η_i) where the kinetic contribution to the energy density greatly dominates over the potential one. In this way, we focus the discussion on solutions that contain a period of fast-roll inflation. These are the only solutions from which one can expect non-trivial imprints of pre-inflationary classical physics in the largest observable angular scales of the CMB. On top of this behavior of early kinetic dominance, observability of these pre-inflationary effects requires that the slow-roll inflationary phase does not last long enough so as to shift the affected scales to regions of the spectrum that are not yet visible to us nowadays. Such inflationary processes often go by the name of just-enough or short-lived inflation [20, 21]. Consequently, in this work we refer to the effects of the kinetically dominated period on the evolution of the primordial perturbations as short-lived inflation effects.

Deep in the pre-inflationary period of kinetic dominance, the contribution of the inflaton potential $V(\phi)$ to the energy density ρ should be negligible compared to the kinetic term. Therefore, we directly approximate the evolution of the cosmological background in the period (η_0, η_i) by restricting eq. (2.1) to the case $V(\phi) = 0$. The resulting dynamics is analytically solvable, with expanding solutions given by

$$a(\eta) = a_0 \sqrt{1 + 2a_0 H_0 (\eta - \eta_0)}, \quad \rho(\eta) = \rho_0 \left[\frac{a_0}{a(\eta)} \right]^6, \quad (2.6)$$

where $H = \sqrt{8\pi\rho/3}$ is the Hubble parameter, with $H_0 = H(\eta_0)$, $a_0 = a(\eta_0)$, and $\rho_0 = \rho(\eta_0)$. For vanishing inflaton potential, one has $\mathcal{U}_{\text{GR}} = 0$, and then the evolution equations for the Mukhanov-Sasaki and tensor perturbations take both the form

$$v_k^{(r)''} + (k^2 + s_{\text{kin}}) v_k^{(r)} = 0, \quad s_{\text{kin}} = \frac{1}{4} \left(\eta - \eta_0 + \frac{1}{2H_0 a_0} \right)^{-2}. \quad (2.7)$$

According to our hypotheses, the kinetic epoch ends when $\eta = \eta_i$. From this instant on, the potential contribution to the inflaton energy density becomes comparable to the kinetic one, and eventually dominates over it. It is after this transition when the slow-roll inflationary period can start, period in which the cosmological background experiences a quasi-de Sitter expansion. For the purposes of this work, we will simplify the physical evolution and completely neglect any of the transition effects from the kinetically dominated period to the slow-roll inflationary one, as well as any change in the inflaton potential that would produce departs in the expansion from an exact de Sitter behavior during slow-roll. These simplifications will allow an analytical investigation of the perturbations dynamics.

More specifically, we approximate the evolution of our cosmological background in the period $[\eta_i, \eta_{\text{end}}]$ by Friedmann equations in vacuo with a cosmological constant. The expanding solutions are

$$a(\eta) = [a_i^{-1} - H_\Lambda (\eta - \eta_i)]^{-1}, \quad (2.8)$$

where $a_i = a(\eta_i)$ and H_Λ is the constant Hubble parameter in de Sitter. We can single out a unique solution if, at the transition time η_i , we impose continuity of the scale factor and its first derivative with a solution of the preceding kinetically dominated phase. This fixes a_i and H_Λ as

$$a_i = a_0 \sqrt{1 + 2a_0 H_0 (\eta_i - \eta_0)}, \quad H_\Lambda = \sqrt{\frac{8\pi}{3}} \rho_0 \left[\frac{a_0}{a_i} \right]^3. \quad (2.9)$$

On the other hand, on a de Sitter background, the time-dependent mass is given by $-a''/a$ for both the Mukhanov-Sasaki and tensor perturbations. Thus, their dynamical equations in this period are¹

$$v_k^{(r)''} + (k^2 + s_{\text{dS}}) v_k^{(r)} = 0, \quad s_{\text{dS}} = -2H_\Lambda^2 [a_i^{-1} - H_\Lambda(\eta - \eta_i)]^{-2}. \quad (2.10)$$

So, in total, we approximate the evolution of cosmological perturbations on a single-field inflationary universe with a kinetically dominated pre-inflationary interval by means of the mode equations

$$v_k^{(r)''} + (k^2 + \tilde{s}_{\text{GR}}) v_k^{(r)} = 0, \quad r = s, t, \quad (2.11)$$

where the approximate time-dependent mass \tilde{s}_{GR} is piecewise-defined as

$$\tilde{s}_{\text{GR}} = \begin{cases} s_{\text{kin}}, & \eta \in (\eta_0, \eta_i), \\ s_{\text{dS}}, & \eta \in [\eta_i, \eta_{\text{end}}], \end{cases} \quad (2.12)$$

with s_{kin} and s_{dS} respectively given in eqs. (2.7) and (2.10).

2.2 Dynamics of gauge invariants: Continuous solutions

The evolution eqs. (2.11) that we have proposed for the Mukhanov-Sasaki and tensor perturbations present a discontinuity at $\eta = \eta_i$, since we clearly have that $s_{\text{kin}}(\eta_i) \neq s_{\text{dS}}(\eta_i)$. This loss of continuity poses an obstruction to the existence and uniqueness of the solutions in all of $(\eta_0, \eta_{\text{end}}]$. Nonetheless, we can isolatedly study the considered equation restricted to the separate intervals (η_0, η_i) and $[\eta_i, \eta_{\text{end}}]$ where there is continuity. Then, we can characterize all of the functions that, in addition to being solutions in each of these two intervals, are also continuous with a continuous derivative at $\eta = \eta_i$. These solutions are uniquely determined by their initial conditions. They make promising candidates to approximate the exact dynamics of the perturbations in our considered classical background with inflaton, given the smoothness of the time-dependent mass in the exact model. This is motivated by our expectation that a very localized change on the mass \tilde{s}_{GR} around η_i to turn it into a continuous function would not modify too much the solution of the dynamical equation for the perturbative modes. At most, we would expect that this discontinuity (and others of similar nature that we will encounter) might introduce an artificially fast rate of variation of the considered solution around it. This effect might propagate forward and translate into some spuriously generated oscillations, possibly on top of other oscillations associated instead with the choice of vacuum state. If this were the case, we would have to identify those spurious oscillations and consistently remove them, as we will explain later on our discussion.²

In the interval (η_0, η_i) the dynamical equation for the Mukhanov-Sasaki and tensor perturbations is given in eq. (2.7). One can readily check that the general solution μ_k of this equation is

$$\mu_k = C_k \sqrt{\frac{\pi y}{4}} H_0^{(1)}(ky) + D_k \sqrt{\frac{\pi y}{4}} H_0^{(2)}(ky), \quad y = \eta - \eta_0 + \frac{1}{2H_0 a_0}, \quad (2.13)$$

¹Strictly speaking, with our instantaneous transition from the kinetic epoch, the derivative a'' would not exist at η_i . Nevertheless, we can define it as the corresponding limit from the right of its expression for $\eta > \eta_i$.

²For more information on the possible origins of the unwanted behavior that we are referring to, and how to correct it, see the end of section 2, the final paragraph of subsection 4.1, and the remarks in the third paragraph of the concluding section.

where C_k and D_k are complex integration constants, while $H_\nu^{(1)}$ and $H_\nu^{(2)}$, with $\nu \in \mathbb{N}$, respectively denote the Hankel functions of the first and second kind [45]. Notice that the solutions depend only on the norm of the wavevector, $k = |\vec{k}|$. This explains our notation for the subindex of μ_k . Each set of solutions μ_k for all \vec{k} can be understood as the set of positive-frequency solutions associated with a vacuum state for the scalar and tensor perturbations in the kinetically dominated period if they are normalized as

$$\mu_k \mu_k^{*'} - \mu_k' \mu_k^* = i, \quad (2.14)$$

where the symbol $*$ denotes complex conjugation. This normalization condition is equivalent to $|D_k|^2 - |C_k|^2 = 1$. We emphasize that, in the considered time interval, each choice of constants C_k and D_k satisfying this requirement univocally determines a normalized solution μ_k .

On the other hand, for our approximate evolution equations (2.10) in the inflationary period $[\eta_i, \eta_{\text{end}}]$, the general solution μ_k is well known [2]:

$$\begin{aligned} \mu_k = & A_k \frac{e^{ik(\eta - \eta_i - a_i^{-1} H_\Lambda^{-1})}}{\sqrt{2k}} \left[1 + \frac{i}{k(\eta - \eta_i - a_i^{-1} H_\Lambda^{-1})} \right] \\ & + B_k \frac{e^{-ik(\eta - \eta_i - a_i^{-1} H_\Lambda^{-1})}}{\sqrt{2k}} \left[1 - \frac{i}{k(\eta - \eta_i - a_i^{-1} H_\Lambda^{-1})} \right], \end{aligned} \quad (2.15)$$

where A_k and B_k are complex integration constants. This solution is normalized according to eq. (2.14) if $|B_k|^2 - |A_k|^2 = 1$. Once again, we notice that any normalized choice of integration constants, for all \vec{k} , specifies a set of positive-frequency solutions for the cosmological perturbations in the de Sitter regime.

Using the preceding information, we can uniquely characterize any physically sensible family of mode solutions for the Mukhanov-Sasaki and tensor perturbations that is defined in the whole time interval $(\eta_0, \eta_{\text{end}}]$ as follows. Starting with the general expressions (2.13) and (2.15) for the (normalized) solutions in, respectively, (η_0, η_i) and $[\eta_i, \eta_{\text{end}}]$, if we impose continuity up to the first derivative at $\eta = \eta_i$, we obtain after a direct computation that

$$\begin{aligned} A_k = & \frac{e^{ik/(a_i H_\Lambda)}}{4} \sqrt{\frac{k\pi}{a_i H_\Lambda}} \left\{ C_k \left[H_0^{(1)} \left(\frac{k}{2a_i H_\Lambda} \right) - \left(\frac{a_i H_\Lambda}{k} - i \right) H_1^{(1)} \left(\frac{k}{2a_i H_\Lambda} \right) \right] \right. \\ & \left. + D_k \left[H_0^{(2)} \left(\frac{k}{2a_i H_\Lambda} \right) - \left(\frac{a_i H_\Lambda}{k} - i \right) H_1^{(2)} \left(\frac{k}{2a_i H_\Lambda} \right) \right] \right\}, \end{aligned} \quad (2.16)$$

$$\begin{aligned} B_k = & \frac{e^{-ik/(a_i H_\Lambda)}}{4} \sqrt{\frac{k\pi}{a_i H_\Lambda}} \left\{ C_k \left[H_0^{(1)} \left(\frac{k}{2a_i H_\Lambda} \right) - \left(\frac{a_i H_\Lambda}{k} + i \right) H_1^{(1)} \left(\frac{k}{2a_i H_\Lambda} \right) \right] \right. \\ & \left. + D_k \left[H_0^{(2)} \left(\frac{k}{2a_i H_\Lambda} \right) - \left(\frac{a_i H_\Lambda}{k} + i \right) H_1^{(2)} \left(\frac{k}{2a_i H_\Lambda} \right) \right] \right\}. \end{aligned} \quad (2.17)$$

This relation between the integration constants in each of the two subintervals is invertible. Furthermore, if C_k and D_k are normalized as indicated above, then the corresponding coefficients A_k and B_k are also normalized, and vice-versa. It follows that any mode solution of our dynamical eqs. (2.11) defined in $(\eta_0, \eta_{\text{end}}]$ that is continuous and has a continuous derivative is fixed once one provides a choice of (normalized) integration constants C_k and D_k (or alternatively their counterpart A_k and B_k). We henceforth restrict our attention to solutions of this type, and regard any choice of integration constants of this kind as a choice of vacuum state or, equivalently, of initial conditions for the primordial perturbations.

2.3 Primordial power spectrum: Non-oscillating behavior

In single-field inflationary cosmology, the standard definition of the primordial power spectrum for scalar and tensor perturbations is respectively³ [44]

$$\mathcal{P}_{\mathcal{R}}(k) = \frac{k^3}{2\pi^2} \frac{|\mu_k(\eta_{\text{end}})|^2}{z^2(\eta_{\text{end}})}, \quad \mathcal{P}_{\mathcal{T}}(k) = \frac{32k^3}{\pi} \frac{|\mu_k(\eta_{\text{end}})|^2}{a^2(\eta_{\text{end}})}, \quad (2.18)$$

where μ_k are solutions of eq. (2.3) with the normalization (2.14). On the other hand, we recall that observationally admissible inflationary epochs must last at least around 60 e-folds [46]. Taking this fact into account, in a situation where one considers an exact de Sitter inflation, it is reasonable to approximate the primordial power spectra (at dominant order) as

$$\mathcal{P}(k) = \frac{k^3}{2\pi^2} \lim_{a \rightarrow \infty} \left(\frac{|\mu_k|^2}{a^2} \right), \quad (2.19)$$

up to a constant factor in the case of tensor perturbations that is irrelevant for the purposes of this work. Employing eq. (2.8) for the scale factor in de Sitter, and eq. (2.15) for the general solution of the Mukhanov-Sasaki and tensor perturbations in this regime, one concludes that

$$\mathcal{P}(k) = \frac{H_{\Lambda}^2}{4\pi^2} |B_k - A_k|^2. \quad (2.20)$$

This is the formula that we will use to estimate the power spectra in our approximate description of primordial fluctuations over a single-field inflationary cosmology with a non-trivial kinetic pre-inflationary epoch.

It is clear that the specific form of the primordial power spectrum (2.20) depends critically on the choice of (normalized) integration constants A_k and B_k (or, equivalently, of C_k and D_k) that fix the mode solutions for the perturbations. In other words, the behavior of the spectrum is strongly affected by the choice of vacuum state for the perturbations. For example, one can proceed as in standard inflationary models and choose the Bunch-Davies state [47], which is the preferred one in a genuine cosmological de Sitter spacetime. Such vacuum state is characterized by $|B_k| = 1$ and $A_k = 0$, and yields the well-known scale-invariant power spectrum [2]. This choice can be thought of as natural if the perturbations with observationally relevant Fourier scales k were not significantly affected by the pre-inflationary dynamics, as it would happen if slow-roll inflation lasted long enough so as to wash away any pre-inflationary effect. However, if this is not the case, the Bunch-Davies vacuum loses its privileged status and the question of which is the natural state for the cosmological fluctuations becomes especially important in order to extract predictions.

In order to get a preliminary hint of the qualitative behavior that can result for the power spectra from eq. (2.20), it is convenient to re-express this formula as

$$\mathcal{P}(k) = \frac{H_{\Lambda}^2}{4\pi^2} \left[|B_k|^2 + |A_k|^2 - 2|A_k||B_k| \cos(\theta_k^A - \theta_k^B) \right] \quad (2.21)$$

where θ_k^A and θ_k^B are the respective phases of A_k and B_k . This expression is well suited to understand why, in certain cases, the primordial power spectrum is going to display an oscillatory behavior in the scale k , and how to avoid these oscillations. Let us consider all

³The time of evaluation η_{end} can also be set a few e-folds after all of the Fourier scales k that are observable nowadays have exited the Hubble horizon, since their (rescaled) evolution freezes then.

choices of integration constants A_k and B_k such that their complex norms are non-oscillating functions of k . Then, it is clear that any k -dependent relative phase between these constants results into a primordial power spectrum that oscillates in k , unless $|A_k|$ is completely negligible compared to $|B_k|$ (we recall that the normalization condition implies that $|B_k| > |A_k|$). It is also worth noticing that, for many choices of this type, the resulting spectrum can display a mixed behavior, in which there are oscillations in those intervals of k where $|A_k|$ and $|B_k|$ are of comparable order, and no oscillations for all k such that $|A_k| \ll |B_k|$ or the difference of phase between A_k and B_k hardly varies.

In section 4 we will argue in favor of a state for the primordial perturbations leading to a power spectrum without those oscillations in k . One of the reasons for this preference is that those oscillations typically involve an enhancement of power, which can be regarded as artificial. For future use, it is worth explaining a straightforward method for eliminating the oscillations in the spectrum for integration constants A_k and B_k with non-oscillatory norms and a k -dependent dephasing. Actually, the resulting power spectrum captures the main information that gets amplified in the presence of the oscillations, as we will see. One simply has to consider a redefinition of the integration constants of the form

$$A_k \rightarrow \tilde{A}_k = |A_k|, \quad B_k \rightarrow \tilde{B}_k = |B_k|. \quad (2.22)$$

In fact, this is a Bogoliubov transformation, in the sense that it leads to new normalized integration constants. The primordial power spectrum for these new tilded constants is

$$\tilde{\mathcal{P}}(k) = \frac{H_\Lambda^2}{4\pi^2} (|B_k| - |A_k|)^2. \quad (2.23)$$

Clearly, this spectrum does not display fast oscillations. Instead, as a function of k , it follows the trajectory of the minima of the oscillations of the original spectrum. In this sense, it indeed maintains the physically relevant information of the original spectrum once the amplifying oscillations have been removed.

3 Approximate mass for the perturbations in hybrid LQC

Going one step beyond the classical inflationary cosmologies previously discussed, in this section we consider their counterparts in the framework of effective LQC. Let us first introduce the model in hybrid LQC that we will later approximate in order to investigate the evolution of the cosmological perturbations on a background that corresponds to a solution of effective (homogeneous and isotropic) LQC. Without backreaction and in the presence of an inflaton ϕ , these background equations are given in conformal time by [26, 29, 31]

$$\left(\frac{a'}{a}\right)^2 = \frac{8\pi}{3} a^2 \rho \left(1 - \frac{\rho}{\rho_c}\right), \quad \frac{a''}{a} = \frac{4\pi}{3} a^2 \rho \left(1 + 2\frac{\rho}{\rho_c}\right) - 4\pi a^2 P \left(1 - 2\frac{\rho}{\rho_c}\right). \quad (3.1)$$

The energy density ρ and pressure P of the inflaton are defined in eq. (2.2), and $\rho_c \simeq 0.4092$ is the maximum value that the energy density can take in effective LQC. The bounce occurs at the moment when this value is reached. These equations are a way of modelling the trajectories of the peaks of certain Gaussian states in homogeneous and isotropic LQC. When the energy density ρ falls to be much smaller than ρ_c , they are simply the classical Friedmann equations considered in section 2.

In the hybrid approach to the quantization of cosmological perturbations in LQC, the dynamical equations for the Mukhanov-Sasaki and tensor modes in the considered effective regime adopt the form [37, 38]

$$v_{\vec{k}}^{(r)''} + [k^2 + s_h^{(r)}] v_{\vec{k}}^{(r)} = 0, \quad r = t, s, \quad (3.2)$$

where [39]

$$s_h^{(t)} = -\frac{4\pi}{3} a^2 (\rho - 3P), \quad s_h^{(s)} = s_h^{(t)} + \mathcal{U}_h, \quad (3.3)$$

and \mathcal{U}_h is a linear combination (with time-dependent coefficients) of the potential $V(\phi)$, its square, and its first and second derivatives with respect to ϕ . In practice, the time-dependent masses $s_h^{(r)}$ ($r = t, s$) become indistinguishable from the classical masses (2.4) as soon as the background solution enters the low-energy regimes with $\rho \ll \rho_c$. However, the difference between these masses is considerable in the regions surrounding the loop quantum bounce.

3.1 Approximate background solutions

Equations (3.1) in effective LQC lead to solutions with a bounce and that display a classical behavior at low matter energy densities. The specific evolution of the solution after the bounce depends on the initial conditions chosen there. Here, we are particularly interested in solutions with phenomenological relevance. Namely, we will focus our attention on effective LQC solutions such that their perturbations with the largest wavelengths that would be observable today in the CMB may have been affected by loop quantum effects in the neighbourhood of the bounce. Roughly speaking, this means that the lowest values of k corresponding to the observable power spectrum are such that k^2 is close to or slightly smaller than $s_h^{(r)}$ in the surroundings of the bounce of the considered background.

This type of effective LQC solutions have been widely studied over the last decade. Remarkably, they all display the following qualitative behavior. The inflaton energy density is dominated by its kinetic contribution at the bounce and, shortly after, a classical period of decelerated expansion follows, leading finally to a slow-roll inflationary period that is short-lived [30–32, 48]. In the case of a quadratic potential for the inflaton, these solutions correspond to choices of inflaton mass of the order $m \sim 10^{-6}$ and an initial value of ϕ at the bounce of order one (in Planck units). On the one hand, these values for m are derived by requiring a number of e-folds of slow-roll inflation that can be compatible with the minimum allowed by observations. Remarkably, this restriction on m alone guarantees that the scales k that are significantly greater than the spacetime curvature at the bounce lie in the observationally (quasi-)scale-invariant part of the power spectrum. On the other hand, the restriction on the initial value of ϕ at the bounce follows from demanding that inflation does not last long enough so as to shift the smallest observable scales k to be much greater than the spacetime curvature at the bounce.⁴ For this type of effective LQC solutions, the inflaton energy density at the onset of inflation ranges in the interval $[10^{-9}, 10^{-12}]$ [48].

Restricting our discussion to these solutions of phenomenological interest, it seems reasonable that, during a large part of the pre-inflationary epoch, one can approximate the

⁴The restriction on the initial datum for ϕ was first obtained numerically after fixing a specific vacuum state for the perturbations [30]. Nonetheless, we consider reasonable to expect that this numerical restriction does not change much if it is reformulated independently of the choice of vacuum, by asking that the smallest observational scales k are close to the curvature at the bounce [32, 34, 48].

dynamics by setting $V(\phi) = 0$ in eq. (3.1). The evolution can then be solved analytically in the proper time t , obtaining [49]

$$\dot{\phi}(t) = \pm\sqrt{2\rho_c}a^{-3}(t), \quad a(t) = (1 + 24\pi\rho_c t^2)^{1/6}. \quad (3.4)$$

Here, we have arbitrarily set $t = 0$ at the bounce, and we have set the global scale of distances such that $a(t = 0) = 1$, without loss of generality.⁵ We notice that the proper time t is related to the conformal one by

$$\eta - \eta_B = {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{3}{2}; -24\pi\rho_c t^2\right) t, \quad (3.5)$$

where η_B is the value of η at the bounce and ${}_2F_1$ is the Gauss hypergeometric function [45]. In this situation with negligible inflaton potential, one can check that the dynamical equations for both the scalar and tensor perturbations have the form

$$v_k^{(r)''} + (k^2 + s_h^{\text{kin}}) v_k^{(r)} = 0, \quad s_h^{\text{kin}} = \frac{8\pi\rho_c}{3} (1 + 24\pi\rho_c t^2)^{-2/3}. \quad (3.6)$$

As we have already commented, the effective LQC trajectories rapidly adopt a classical behavior after the bounce occurs. Focusing on solutions of the considered type, it is then reasonable to expect that the analysis and approximations developed in section 2 can be applied now as well, starting at a certain instant η_0 in the kinetically dominated regime. In this way, we will have at our disposal a completely analytical description, at least approximate, of our effective LQC background.

More specifically, the initial conditions for eq. (2.6) at that time η_0 (in the kinetic epoch) that can approximate well the effective LQC evolution of our background towards the future are

$$a_0 = (1 + 24\pi\rho_c t_0^2)^{1/6}, \quad \rho_0 = \rho_c a_0^{-6}, \quad (3.7)$$

where $t_0 = t(\eta_0)$ is calculated using eq. (3.5). The accuracy of this approximation depends on the choice of time η_0 for the matching of initial conditions using the LQC data. In particular, since our interest here is the study of the propagation of cosmological perturbations, an acceptable choice of η_0 in the kinetically dominated region is one for which the difference between the hybrid LQC mass s_h^{kin} and the classical mass s_{kin} becomes negligible. This latter mass is obtained from eq. (2.7), evaluated at the initial conditions (3.7) inherited from effective LQC. According to our comments, it is then convenient to introduce the relative difference

$$2 \frac{|s_{\text{kin}} - s_h^{\text{kin}}|}{(s_{\text{kin}} + s_h^{\text{kin}})} \quad (3.8)$$

which measures the relative error made when one approximates the hybrid LQC mass by its classical counterpart in the kinetic epoch. This difference can be computed analytically as a function of time, for different choices of η_0 . The results show that, for $t_0 \gtrsim 0.4$, the error committed grows at most up to a 3% immediately after the initial time t_0 (instant at which both masses coincide), and becomes negligible at large $t > t_0$. However, the maximum of this error rapidly increases if one chooses smaller values of t_0 .

⁵In standard cosmology, one usually sets the present value of the scale factor equal to one. Choosing instead the value at the bounce shifts proportionally all the Fourier scales k in the observable window with respect to those reported by e.g. the Planck mission.

3.2 Pöschl-Teller approximation

Even with the approximated analytical description of the effective LQC background at our disposal, we are not able to solve exactly the mode equation (3.6) for the Mukhanov-Sasaki and tensor perturbations. This situation calls for an additional approximation to the time-dependent mass s_h^{kin} near the bounce, that allows us to find explicit solutions for the cosmological perturbations.

In this respect, refs. [40, 49] suggested that the behavior of the mass s_h^{kin} in the vicinity of the bounce closely resembles that of a Pöschl-Teller potential of the form

$$s_{\text{PT}} = \frac{U_0}{\cosh^2[\alpha(\eta - \eta_B)]}. \quad (3.9)$$

The works that have used this expression to approximate the mass in hybrid LQC have so far chosen the values $U_0 = 8\pi\rho_c/3$ and $\alpha = \sqrt{16\pi\rho_c}$ for the respective peak and width of the potential [40]. These values are obtained by demanding that the functions s_{PT} and s_h^{kin} coincide at the bounce, up to the second time derivative (included). However, if one compares this choice of Pöschl-Teller mass with s_h^{kin} , one realizes that the relative error rapidly grows beyond a 20% for $t \gtrsim 0.25$ (see the dashed lines in figure 1). Then, according to our discussion above, the interval of time for which this Pöschl-Teller mass can be considered a fairly good approximation is too narrow to admit admissible matching points with the classical kinetically dominated background.

With the aim of modelling a more realistic transition to the classical regime, we can improve the Pöschl-Teller approximation of the hybrid LQC mass near the bounce by changing its width to values other than $\alpha = \sqrt{16\pi\rho_c}$. More concretely, we demand that s_{PT} coincide with s_h^{kin} at the bounce and with s_{kin} at the time η_0 where we connect the effective LQC evolution with the classical kinetic epoch. This requirement fixes the peak and width of the Pöschl-Teller potential in the following way:

$$U_0 = \frac{8\pi\rho_c}{3}, \quad \alpha = \frac{\text{arcosh}(a_0^2)}{(\eta_0 - \eta_B)}, \quad (3.10)$$

as one can check using eqs. (2.7), (3.6), and (3.7). Therefore, we can try and approximate the evolution of the primordial cosmological perturbations on the effective LQC background with short-lived inflation as

$$v_k^{(r)''} + (k^2 + \tilde{s}_h) v_k^{(r)} = 0, \quad r = s, t, \quad (3.11)$$

where the time-dependent mass \tilde{s}_h is

$$\tilde{s}_h = \begin{cases} U_0 \cosh^{-2}[\alpha(\eta - \eta_B)], & \eta \in [\eta_B, \eta_0], \\ \tilde{s}_{\text{GR}} & \eta \in (\eta_0, \eta_{\text{end}}], \end{cases} \quad (3.12)$$

and α and U_0 are given in eq. (3.10). Recall that \tilde{s}_{GR} is our approximation to the classical mass [see eq. (2.12)].

Let us notice that the choice of parameters for the Pöschl-Teller potential made in eq. (3.10) depends on the instant η_0 where we place the transition to a classical universe. In order to elucidate which values of η_0 are acceptable, it is convenient to define the relative difference

$$\text{Err} = 2 \frac{|\tilde{s}_h - s_h^{\text{kin}}|}{(\tilde{s}_h + s_h^{\text{kin}})}, \quad (3.13)$$




Figure 1. The relative difference Err as a function of the proper time t , for several choices of $t_0 \geq 0.4$. Dashed lines: The corresponding function Err defined using the Pöschl-Teller potential with $\alpha = \sqrt{16\pi\rho_c}$, for $t_0 \simeq 0.19$ (left panel) and $t_0 \simeq 0.27$ (right panel).

which measures the relative error made when one approximates the hybrid LQC mass in the kinetic epoch as we have suggested. Figure 1 shows this error as a function of time, for several choices of $t_0 \geq 0.4$. For completeness, this figure also displays the error that we would have made if we had defined \tilde{s}_h using the Pöschl-Teller potential with $\alpha = \sqrt{16\pi\rho_c}$. This proves that our proposed parameters (3.10) provide an overall improvement of the existing approximations of the hybrid LQC mass employing a Pöschl-Teller term [40], without compromising its classical behavior. Furthermore, inspecting figure 1, one can argue that the choices of transition time t_0 that optimize our approximation lie around $t_0 \simeq 0.4$.

3.3 Approximate dynamics of gauge invariants: Continuous solutions

The dynamical eq. (3.11), introduced to approximate the evolution of the perturbations in hybrid LQC for effective backgrounds with short-lived inflation, can be solved analytically in the two time intervals $[\eta_B, \eta_0]$ and $(\eta_0, \eta_{\text{end}}]$. For this last interval, which corresponds to the classical period, the general solution with continuity up to the first derivative has been characterized in section 2. On the other hand, in the interval $[\eta_B, \eta_0]$ that describes the vicinity of the bounce, the perturbations evolve according to

$$v_k^{(r)''} + (k^2 + s_{\text{PT}}) v_k^{(r)} = 0, \quad r = s, t, \quad (3.14)$$

where s_{PT} is the Pöschl-Teller mass (3.9) with parameters (3.10). One can check that its general solution μ_k is

$$\begin{aligned} \mu_k = & M_k [x(1-x)]^{-ik/2\alpha} {}_2F_1 \left(b_1^k, b_2^k; b_3^k; x \right) \\ & + N_k x^{ik/2\alpha} (1-x)^{-ik/2\alpha} {}_2F_1 \left(b_1^k - b_3^k + 1, b_2^k - b_3^k + 1; 2 - b_3^k; x \right), \end{aligned} \quad (3.15)$$

where M_k and N_k are integration constants, and we have defined the time variable

$$x = \left[1 + e^{-2\alpha(\eta - \eta_B)} \right]^{-1}. \quad (3.16)$$

In addition, the parameters of the hypergeometric function are

$$b_3^k = 1 - \frac{ik}{\alpha}, \quad (3.17)$$

$$b_1^k = \frac{1}{2} \left(1 + \sqrt{1 + \frac{32\pi\rho_c}{3\alpha^2}} \right) - \frac{ik}{\alpha}, \quad b_2^k = \frac{1}{2} \left(1 - \sqrt{1 + \frac{32\pi\rho_c}{3\alpha^2}} \right) - \frac{ik}{\alpha}. \quad (3.18)$$

This solution can be normalized according to eq. (2.14) with a suitable restriction on the constants M_k and N_k , which we need not show here.

At the instant η_0 , the approximate mass function \tilde{s}_h is continuous by construction. Then, each choice of integration constants M_k and N_k , namely of a solution in the surrounding of the bounce, selects a unique solution (2.13) in the kinetic epoch. The constants C_k and D_k associated with this solution are obtained by demanding its continuity and that of its derivative at η_0 . They are given by

$$C_k = \left[H_0^{(1)} \left(\frac{k}{2H_0 a_0} \right) \right]^{-1} \left[\sqrt{\frac{8H_0 a_0}{\pi}} \mu_k(\eta_0) - D_k H_0^{(2)} \left(\frac{k}{2H_0 a_0} \right) \right], \quad (3.19)$$

$$D_k = i \sqrt{\frac{\pi}{8H_0 a_0}} \left[k H_1^{(1)} \left(\frac{k}{2H_0 a_0} \right) \mu_k(\eta_0) - H_0^{(1)} \left(\frac{k}{2H_0 a_0} \right) H_0 a_0 \mu_k(\eta_0) + H_0^{(1)} \left(\frac{k}{2H_0 a_0} \right) \mu_k'(\eta_0) \right]. \quad (3.20)$$

As we explained in section 2, any choice of these constants fixes in turn a solution in the de Sitter period provided that it is continuous and has a continuous derivative at η_i , by means of eqs. (2.16) and (2.17).

In summary, if we restrict our attention to solutions of our approximate equation (3.11) that are C^1 at the matching points, each choice of integration constants M_k and N_k in the vicinity of the bounce leads to a unique solution throughout the whole pre-inflationary and inflationary periods, solution that can be calculated explicitly. Alternatively, the same happens if one chooses the pair of constants C_k and D_k in the kinetic epoch, or A_k and B_k in the de Sitter period. The vacuum state associated with any of such choices for all \vec{k} leads to a primordial power spectrum (2.20), with properties that we can investigate analytically.

4 Short-lived inflation vs. loop quantum bounce effects

In what follows, we will use the approximations introduced in sections 2 and 3 for an analytical study of the differences arising in the primordial power spectra from the evolution of the perturbations on a purely classical universe, on the one hand, and on a hybrid LQC background, on the other hand, both with a period of short-lived inflation. For a robust comparison, the first and most important step that we need to take is to establish a common criterion for the selection of a vacuum state of the perturbations, on any of the considered backgrounds.

4.1 Choice of vacuum state

As we discussed in section 2, in any situation where a pre-inflationary epoch may have affected the perturbations with scales k that are observable nowadays, the arguments leading to the Bunch-Davies state lose their strength. Instead, one should develop new criteria that distinguish a preferred quantum state for the perturbations, taking into account the pre-inflationary dynamics. In the lack of enough symmetries in the background spacetime that can serve to naturally select such a state, several proposals have emerged over the last decade

in the context of LQC [30, 32, 33, 40, 49–52]. A notorious example is the consideration of states constructed with the imposition of (low-order) adiabatic initial conditions [53, 54]. These conditions are usually set at the bounce, or at some instant in the distant past from it. Other more recent proposals involve the minimization of the fields uncertainty relations in the vicinity of the bounce [32, 51], or of the smeared time variation of the mode solutions throughout a good portion of the evolution from the bounce to the onset of inflation [33].

All of these proposals to select a vacuum state heavily depend on the choice of time to set the initial conditions, and/or require numerical techniques to determine them. Furthermore, the numerical evolution of the associated mode solutions typically leads to primordial power spectra that are highly oscillating with respect to the scale k [30–32, 48, 52]. In order to extract predictions from the theory, these oscillations are often averaged out. In recent years, concerns have been raised about whether these oscillations are physically acceptable, or they rather should be interpreted as an indication that the corresponding state of the perturbations is not a natural vacuum in the theory [33, 48, 55]. For example, the Bunch-Davies vacuum does not present such oscillations, not even when quasi-de Sitter effects of the inflationary background destroy its scale-invariance. This state is undoubtedly a preferred one when the attention is restricted to slow-roll inflation, and it provides accurate tested predictions for most observable scales k . Moreover, oscillations in the power spectrum can be thought of as the remnants of a past evolution of the mode solutions with fast oscillations in time. Such oscillatory dynamics can easily erase relevant information about the effects of the earliest epochs of the pre-inflationary Universe on the perturbations. Finally, oscillations in the primordial spectrum have been seen to often pump a somewhat artificial amplification of power, which in some cases can even be in slight tension with the CMB observations [30, 48].

Let us recall that the aim of this work is to develop manageable tools to investigate the differences between classical (short-lived inflationary) effects and LQC effects on the primordial power spectrum. From this perspective, it would be especially desirable to characterize the vacuum state *analytically* by means of a criterion such that it is motivated by fundamental arguments and can potentially lead to a power spectrum with non-oscillating properties. In this respect, a proposal has been recently put forward in hybrid LQC which seems promising to address both requirements [55–57]:

- i) It can be directly related [57] to the vacuum state associated with a set of annihilation and creation operators that possess appealing physical properties: their evolution in hybrid quantum cosmology is dictated by a diagonal Hamiltonian, in which the self-interacting parts have been eliminated order by order in the local ultraviolet regime $k \rightarrow \infty$ [56]. It is worth noticing that the procedure leading to this vacuum state is very different from an instantaneous diagonalization of the Hamiltonian.
- ii) The norm of the mode solutions for the Mukhanov-Sasaki and tensor perturbations corresponding to this vacuum state in the effective LQC regimes minimize the scale and time-dependent oscillations in the ultraviolet regime $k \rightarrow \infty$ [55].

The aforementioned state is characterized by the mode solutions⁶

$$\mu_k = \sqrt{-\frac{1}{2\text{Im}(h_k)}} e^{i \int_{\eta_0}^{\eta} d\tilde{\eta} \text{Im}(h_k)(\tilde{\eta})} \quad (4.1)$$

⁶The lower end of the integration interval in eq. (4.1) is set equal to η_0 for convenience. The solutions for other choices only differ in global constant phases, irrelevant for the power spectrum.

for all \vec{k} , where h_k is a solution of the differential equation

$$h'_k = k^2 + s_h^{(r)} + h_k^2 \quad (4.2)$$

that displays the following asymptotic expansion

$$kh_k^{-1} \sim i \left[1 - \frac{1}{2k^2} \sum_{n=0}^{\infty} \left(\frac{-i}{2k} \right)^n \gamma_n \right]. \quad (4.3)$$

Here, γ_n are k -independent functions of time that are fixed by the relations

$$\gamma_0 = s_h^{(r)}, \quad \gamma_{n+1} = -\gamma'_n + 4s_h^{(r)} \left[\gamma_{n-1} + \sum_{m=0}^{n-3} \gamma_m \gamma_{n-(m+3)} \right] - \sum_{m=0}^{n-1} \gamma_m \gamma_{n-(m+1)}. \quad (4.4)$$

Although this proposal of vacuum state was first motivated and introduced in the context of hybrid LQC, one can straightforwardly generalize the formulas that define its mode solutions to other time-dependent masses different from $s_h^{(r)}$. In particular, the proposal selects the Bunch-Davies vacuum if the mass of the primordial perturbations becomes the mass associated with a de Sitter spacetime in GR. However, in the case of the time-dependent masses \tilde{s}_{GR} and \tilde{s}_h , with which we respectively approximate the dynamics of the perturbations on classical and LQC backgrounds with short-lived inflation, the above formulas for the selection of a vacuum cannot be directly applied. Actually, we recall that these masses have a discontinuity at the beginning of inflation, at η_i . In addition, \tilde{s}_h is not smooth at η_0 . In spite of these obstructions, the formulas are nonetheless applicable in each of the pre-inflationary periods (η_0, η_i) and $[\eta_B, \eta_0]$.⁷ Thus, we will use them as a first step to characterize, with the same criteria, the respective preferred sets of mode solutions on the classical and the effective LQC backgrounds. With the resulting primordial spectra at our disposal, we will then employ the method explained at the end of section 2 to suitably redefine the vacuum in order to eliminate all the spurious oscillations that amplify the power.

4.2 Power spectrum for short-lived inflation in GR

Let us now consider the evolution of primordial perturbations in a classical relativistic cosmology with a short-lived single-field inflationary period. More specifically, we want to study the primordial power spectrum for the Mukhanov-Sasaki and tensor perturbations that propagate on a background satisfying the Friedmann equation in the interval $(\eta_0, \eta_{\text{end}}]$, where η_0 is the instant for the Big-Bang singularity. With this purpose, we approximate the dynamics of the gauge invariant perturbations, describing it by means of eq. (2.11), that models the evolution on a kinetically dominated background that instantaneously transitions to a de Sitter phase.

If we decided to fix the mode solutions in the de Sitter phase $[\eta_i, \eta_{\text{end}}]$ by using eqs. (4.1) and (4.3), we would simply arrive at the Bunch-Davies vacuum and a scale-invariant power spectrum. This choice of state would not encode any information from the pre-inflationary period, regardless of how long inflation lasts. This seems unphysical in those cases in which there exist observable Fourier scales k of the order of the time-dependent mass in the pre-inflationary epoch. Therefore, for background solutions with short-lived inflation, this choice of a vacuum is not natural anymore.

⁷To use eq. (4.4) at the right end of the interval $[\eta_B, \eta_0]$, we define all of the derivatives of the mass \tilde{s}_h as the corresponding limits from the left.

Another possibility, motivated by our criterion for the selection of vacuum state, is to employ formulas (4.1) and (4.3) instead in the kinetic epoch (η_0, η_i) to try and fix the mode solutions there. As shown in the Appendix, this procedure leads to solutions of the form (2.13) with integration constants given by $C_k = 0$ and $D_k = 1$. These solutions have been previously selected to study the possible effects of short-lived inflation on the power spectrum, motivated by certain adiabatic considerations [6]. Actually, they can be obtained by demanding that the mode solutions correspond to the Poincaré vacuum in the limit $\eta \rightarrow \infty$. However, we should remark that the motivation and range of applicability of the formulas used here to select those solutions are much more general.

Let us recall that the continuity of the solutions and of their first derivatives at η_i leads to a unique primordial power spectrum [of the form (2.20)]. For the considered choice of vacuum state, the constants A_k and B_k are those corresponding to the values $C_k = 0$ and $D_k = 1$ in eqs. (2.16) and (2.17). If we define the Fourier scale $k_I = a_i H_\Lambda$, associated with the spacetime curvature effects at the onset of inflation, the resulting power spectrum is scale-invariant when $k \gg k_I$, oscillates when $k \gtrsim 2k_I$, and presents a steep suppression for smaller k (see the dashed line in the right panel of figure 2). We can understand this behavior by studying the different k -regions relative to k_I separately.

In the region $k \gg k_I$, one can check that, for $C_k = 0$ and $D_k = 1$,

$$A_k = \mathcal{O}(\sqrt{k_I/k}), \quad B_k = e^{-3ik/(2k_I)+i\pi/4} + \mathcal{O}(\sqrt{k_I/k}). \quad (4.5)$$

We recall that the symbol \mathcal{O} stands for terms of the order of its argument or subdominant. This result follows from the asymptotic behavior of the Hankel functions $H_\nu^{(j)}$ [45],

$$H_\nu^{(j)}(x) = \sqrt{\frac{2}{\pi x}} \exp[-i(-1)^j(x - \nu\pi/2 - \pi/4)] + \mathcal{O}(x^{-1}), \quad x \in \mathbb{R}, \quad x \rightarrow \infty, \quad (4.6)$$

and it explains the scale-invariance of the power spectrum in the ultraviolet region (relative to k_I) at dominant order.

In the region $k \ll k_I$, we have that, for $C_k = 0$ and $D_k = 1$,

$$A_k = \sqrt{\frac{k\pi}{16k_I}} \left[\left(1 + i\frac{k}{k_I}\right) H_0^{(2)}\left(\frac{k}{2k_I}\right) - \left(\frac{k_I}{k} + \frac{k}{k_I}\right) H_1^{(2)}\left(\frac{k}{2k_I}\right) \right] + A_k \mathcal{O}(k^2/k_I^2), \quad (4.7)$$

$$B_k = \sqrt{\frac{k\pi}{16k_I}} \left[\left(1 - i\frac{k}{k_I}\right) H_0^{(2)}\left(\frac{k}{2k_I}\right) - \left(\frac{k_I}{k} + \frac{k}{k_I}\right) H_1^{(2)}\left(\frac{k}{2k_I}\right) \right] + B_k \mathcal{O}(k^2/k_I^2). \quad (4.8)$$

On the other hand, the following relations at dominant order (denoted with the symbol \sim) are valid for $|z| \ll 1$ [45]:

$$-iH_0^{(1)}(z) \sim iH_0^{(2)}(z) \sim \frac{2}{\pi} \ln z, \quad -iH_1^{(1)}(z) \sim iH_1^{(2)}(z) \sim -\frac{2}{\pi z}. \quad (4.9)$$

Therefore, we get the dominant behavior

$$|B_k - A_k| \sim \sqrt{\frac{k^3}{\pi k_I^3}} \left| \ln\left(\frac{k}{2k_I}\right) \right|, \quad (4.10)$$

which rapidly tends to zero as k/k_I does. This explains the sharp suppression in the infrared region (relative to k_I).




Figure 2. *Left panel:* Functions $|A_k|$ and $|B_k|$ of k away from the ultraviolet regime $k \gg k_I$. *Right panel:* Non-oscillating power spectrum $4\pi^2\mathcal{P}_{\text{kin}}(k)/H_\Lambda^2$ (continuous line), compared with its oscillating version corresponding to the choice $C_k = 0$ and $D_k = 1$ (dashed line).

Finally, in the intermediate region $k \gtrsim 2k_I$ one can check in eqs. (2.16) and (2.17) that, for $C_k = 0$ and $D_k = 1$, the relative phase between A_k and B_k contains the dominant term $2k/k_I$. This ratio is of order one (or slightly bigger) in the considered region. It immediately follows that, unless $|A_k|$ and $|B_k|$ are oscillatory functions of k , the power spectrum (2.20) oscillates for $k \gtrsim 2k_I$ owing to the commented phase. Indeed, in the left panel of figure 2 we plot $|A_k|$ and $|B_k|$ away from the ultraviolet region $k \gg k_I$ and we see that they do not oscillate.

Looking at the dashed line in the right panel of figure 2, we also see that the oscillations in the intermediate region $k \gtrsim 2k_I$ result in a mean amplification of the power in that sector. In this context, we refer to our discussion at the end of section 2 and, accordingly, we proceed to eliminate this oscillatory behavior by considering instead the integration constants

$$A_k \rightarrow A_k^{\text{kin}} = |A_k|, \quad B_k \rightarrow B_k^{\text{kin}} = |B_k|. \quad (4.11)$$

With them, we obtain the non-oscillating power spectrum

$$\mathcal{P}_{\text{kin}}(k) = \frac{H_\Lambda^2}{4\pi^2} \left(B_k^{\text{kin}} - A_k^{\text{kin}} \right)^2, \quad (4.12)$$

that captures the non-amplified information about the oscillatory region of the original spectrum for $k \gtrsim 2k_I$, while respecting the infrared and ultraviolet behaviors. We show this non-oscillating spectrum in the right panel of figure 2, comparing it to its original version for the vacuum determined by the choice $C_k = 0$ and $D_k = 1$.

In summary, in a cosmological model with short-lived inflation ruled by the Friedmann equation of GR, we have singled out a set of mode solutions starting with a general criterion applied in the kinetically dominated regime, and then eliminating the oscillations that amplify the resulting power spectrum. Adopting this procedure, the effects on the primordial power spectrum of a purely classical pre-inflationary period in a short-lived inflationary cosmology are as follows. Given the scale k_I , equal to the inverse of the radius of the comoving Hubble horizon at the onset of inflation, one distinguishes two sectors of Fourier scales k with very different properties. On the one hand, for $k \gg k_I$ the power spectrum is nearly scale-invariant, something that can be interpreted as the consequence that the perturbations are in the Bunch-Davies state at the end of inflation for those scales. On the other hand, as k

becomes comparable to k_I and gets smaller, the power spectrum gets highly suppressed. This indicates that, for such scales, the perturbations are in a(n excited) state different than the Bunch-Davies solution.

Remarkably, this depart from the Bunch-Davies state only for Fourier scales smaller than k_I happens even when *all the scales* end up crossing the Hubble horizon as one approaches the Big-Bang singularity in the kinetically dominated phase. It is often argued that this crossing can excite the corresponding perturbations with respect to their Bunch-Davies state. Nonetheless, our results show that there exists in fact a natural mechanism in the kinetic epoch that can de-excite all modes with $k \gg k_I$. This means that, strictly speaking, the existence of an upper bound in the cosmological curvature (as e.g. in the case of effective LQC [32, 34]) is not completely necessary to explain a power suppression affecting only the infrared sector, as indicated by observations. Therefore, one needs additional features and details of the spectrum to distinguish effects in our primordial Universe that can be attributed exclusively to LQC.

4.3 Power spectrum for hybrid LQC

Let us finally analyze the primordial power spectrum of the cosmological perturbations in effective scenarios of hybrid LQC with short-lived inflation, using the same criterion for the choice of vacuum state as in the GR case. As we have motivated in section 3, we approximate the corresponding evolution of the Mukhanov-Sasaki and tensor perturbations by means of eq. (3.11). It contains a period $[\eta_B, \eta_0]$ in the vicinity of the bounce where the mass behaves as a Pöschl-Teller potential, and then a classical kinetically dominated regime (η_0, η_i) , followed by a de Sitter inflation during the interval $[\eta_i, \eta_{\text{end}}]$. Recalling the properties of the LQC backgrounds of interest, our approximation requires that the energy density at the instant η_i be in the range $[10^{-9}, 10^{-12}]$. Taking, e.g., a value of the order of 10^{-9} , i.e. the largest possible one, this requirement implies that $k_I = a_i H_\Lambda$ must be of the order of 10^{-3} .

Formulas (4.1) and (4.3), associated with our preferred criterion for the choice of vacuum state, can be separately applied in any of the intervals $[\eta_B, \eta_0]$, (η_0, η_i) , or $[\eta_i, \eta_{\text{end}}]$. The resulting set of mode solutions, that are continuous and have a continuous derivative at the matching points, changes with the interval chosen among the three commented options. As mentioned above, if we use formulas (4.1) and (4.3) in the interval $[\eta_i, \eta_{\text{end}}]$, we fix the Bunch-Davies vacuum, which is not the most natural choice if the pre-inflationary dynamics has affected the perturbations with observable scales k . On the other hand, if we instead focus our analysis on the classical kinetic period (η_0, η_i) , we obtain the state and power spectrum studied in the previous subsection. This spectrum only encodes information from the effects that the pre-inflationary relativistic evolution of the background can have imprinted on the perturbations. In the context of effective LQC, quantum geometry phenomena can importantly affect the evolution of the perturbations near the bounce. One then reasonably expects that there should exist a different, yet physically preferred, vacuum state with a power spectrum that captures such LQC effects. This leads us to consider the last remaining possibility, namely, that we apply the considered formulas (4.1) and (4.3) in the interval $[\eta_B, \eta_0]$ in order to fix the mode solutions for the perturbations.

As we show in the Appendix, the particular solution μ_k of eq. (3.11) obtained with eqs. (4.1) and (4.3) in $[\eta_B, \eta_0]$ corresponds to

$$h_k = -i\alpha\tilde{k} - 2\alpha x(1-x) \frac{cd}{1+i\tilde{k}} \frac{{}_2F_1(c+1, d+1; 2+i\tilde{k}; x)}{{}_2F_1(c, d; 1+i\tilde{k}; x)}, \quad (4.13)$$

where $\tilde{k} = k/\alpha$ and we have introduced the following k -independent constants:

$$c = \frac{1}{2} \left(1 + \sqrt{1 + \frac{32\pi\rho_c}{3\alpha^2}} \right), \quad d = \frac{1}{2} \left(1 - \sqrt{1 + \frac{32\pi\rho_c}{3\alpha^2}} \right). \quad (4.14)$$

We recall that α and x are defined in eqs. (3.10) and (3.16). It is possible to show that the resulting normalized solution μ_k is that given by the integration constants $M_k = 1/\sqrt{2k}$ and $N_k = 0$ (up to a constant phase) in eq. (3.15). Actually, this solution has been previously considered in the study of the primordial power spectrum arising from hybrid LQC, motivated by the imposition of adiabatic conditions in the far past region of the Pöschl-Teller term [40, 49]. These previous analyses adopted much simpler approximations for the kinetic epoch than those presented here. In addition, we emphasize that our criterion for the selection of vacuum state is supported by fundamental arguments applicable in more general scenarios than the presence of a time-dependent mass of Pöschl-Teller form, and do not rest on particular asymptotic regimes of this mass (or of the associated cosmological background).

Starting with the normalized solutions μ_k in the interval $[\eta_B, \eta_0]$ for each \tilde{k} , obtained by substituting in eq. (4.1) the function h_k of eq. (4.13), we fix a set of mode solutions for the perturbations on the whole of our approximated LQC cosmology after the bounce, provided that we impose continuity up to the first time derivative. The resulting power spectrum is given by eq. (2.20), where the integration constants A_k and B_k are those specified in eqs. (2.16) and (2.17) in terms of C_k and D_k , which are in turn fixed by means of eqs. (3.19) and (3.20). In particular, we notice that

$$\mu_k(\eta_0) = \sqrt{-\frac{1}{2\text{Im}(h_k)(\eta_0)}}, \quad \mu'_k(\eta_0) = -h_k^*(\eta_0)\mu_k(\eta_0), \quad (4.15)$$

From our expressions, one can realize that the behavior of the resulting spectrum depends on the considered region of k relative to the scale $k_I \sim 10^{-3}$ at the onset of inflation, on the one hand, and to the quantities $a_0 H_0$ and α , on the other hand. For transition times between the LQC and the classical regimes such that our approximations to the mass are reasonably good (i.e., around $t_0 = 0.4$), we have that the parameters $a_0 H_0$ and α are not far from one and three in Planck units, respectively. Remarkably, these values are of the same order as the time-dependent mass s_h^{kin} at the bounce, in hybrid LQC. In view of this, we define (e.g.) the scale $k_{\text{LQC}} = \alpha$ associated with the potentially observable LQC effects on the evolution of the perturbations near the bounce.

Let us first consider the sector of Fourier scales with $k \gg 1$. This covers the ultraviolet region, relative to both k_I and k_{LQC} . Taking into account the asymptotic behavior of the hypergeometric function ${}_2F_1(a_1, a_2; a_3, z)$ for large a_3 [45], we have

$$\text{Im}(h_k) = -k + \mathcal{O}(k^{-1}), \quad \text{Re}(h_k) = \mathcal{O}(k^{-2}), \quad (4.16)$$

and therefore

$$\mu_k(\eta_0) = \sqrt{\frac{1}{2k}} + \mathcal{O}(k^{-5/2}), \quad \mu'_k(\eta_0) = -i\sqrt{\frac{k}{2}} + \mathcal{O}(k^{-3/2}). \quad (4.17)$$

Inserting this in eqs. (3.19) and (3.20) (which give the integration constants C_k and D_k) and using the asymptotic behavior of the Hankel functions, we get

$$C_k = \mathcal{O}(k^{-1/2}), \quad D_k = e^{ik/(2H_0 a_0) - i\pi/4} + \mathcal{O}(k^{-1}), \quad (4.18)$$




Figure 3. The functions in both panels are computed with a transition time between the LQC and classical epochs equal to $t_0 = 0.4$. *Left panel:* Functions $|C_k|$ and $|D_k|$ of k away from the ultraviolet region $k \gg 1$. *Right panel:* Functions $|A_k|$ and $|B_k|$ of k away from the ultraviolet region $k \gg 1$.

from which we obtain the following ultraviolet behavior for A_k and B_k :

$$|A_k| = \mathcal{O}(k^{-1/2}), \quad |B_k| = 1 + \mathcal{O}(k^{-1/2}). \quad (4.19)$$

The resulting power spectrum is therefore scale-invariant for $k \gg 1$, in agreement with the standard result derived by using a Bunch-Davies state.

We next consider the region with $k \lesssim k_{\text{LQC}}$. Recalling the orders of magnitude that the scales k_{LQC} and k_I take in our system, in this region $3 \gtrsim k \gg k_I$. It then follows that

$$A_k = e^{3ik/(2k_I) - i\pi/4} C_k \left[1 + \mathcal{O}(\sqrt{k_I/k}) \right] + D_k \mathcal{O}(\sqrt{k_I/k}), \quad (4.20)$$

$$B_k = e^{-3ik/(2k_I) + i\pi/4} D_k \left[1 + \mathcal{O}(\sqrt{k_I/k}) \right] + C_k \mathcal{O}(\sqrt{k_I/k}). \quad (4.21)$$

To improve this asymptotic formula, we need more information about the relative order of magnitude of the integration constants C_k and D_k . In the considered region, a priori, we do not expect this relative order to depend on k/k_I , because the formulas that determine those constants do not involve the scale k_I . We confirm our expectations in the left panel of figure 3, where we plot the values of $|C_k|$ and $|D_k|$ given by their analytical expressions. Therefore, in the considered region $k \lesssim k_{\text{LQC}}$ and at dominant order with respect to k/k_I , we obtain the following behavior for the integration constants A_k and B_k :

$$A_k \sim e^{3ik/(2k_I) - i\pi/4} C_k, \quad B_k \sim e^{-3ik/(2k_I) + i\pi/4} D_k. \quad (4.22)$$

Notice the relative phase between them, that necessarily contains $3k/k_I$. Consequently, we expect the appearance of rapid oscillations in the resulting power spectrum around $k_{\text{LQC}} \gg k_I$. Furthermore, we would also find slower, superposed oscillations if a k -dependent relative phase happened to exist between the constants C_k and D_k .

The oscillations in the power spectrum suggest that we should still improve our choice of vacuum state, in the same spirit that was explained at the end of section 2. Indeed, our choice so far determines constants A_k and B_k that have non-oscillatory norms. Apart from the previous discussion, valid for k around k_{LQC} , we show this in the right panel of figure 3 for all scales that do not belong to the ultraviolet region $k \gg 1$. In view of this, we follow *exactly the same procedure* as in the case of the classical GR cosmology regime studied above, and redefine the integration constants A_k and B_k in such a way that the oscillations that amplify




Figure 4. *Left panel:* Non-oscillating power spectrum $4\pi^2\mathcal{P}_{\text{LQC}}(k)/H_\Lambda^2$ in hybrid LQC (continuous line), compared with its analog $4\pi^2\mathcal{P}_{\text{kin}}(k)/H_\Lambda^2$ in the GR context (dashed line). *Right panel:* Subportion of the previous panel, to show the sensitivity on the choice of transition time t_0 between the LQC and classical epochs.

the resulting power spectrum are eliminated, while respecting its scale-invariant ultraviolet behavior. Explicitly, we introduce the following Bogoliubov transformation:

$$A_k \rightarrow A_k^{\text{LQC}} = |A_k|, \quad B_k \rightarrow B_k^{\text{LQC}} = |B_k|, \quad (4.23)$$

and consider the associated power spectrum

$$\mathcal{P}_{\text{LQC}}(k) = \frac{H_\Lambda^2}{4\pi^2} \left(B_k^{\text{LQC}} - A_k^{\text{LQC}} \right)^2. \quad (4.24)$$

We plot this non-oscillating spectrum \mathcal{P}_{LQC} in figure 4. There, we also compare it with the spectrum \mathcal{P}_{kin} that we have previously obtained in the case of a purely classical background, particularizing it to $k_I = 2.8 \times 10^{-3}$. Recalling that $k_I = a_i H_\Lambda$, this choice defines a classical cosmology where the inflation starts, roughly speaking, with the largest energy density that is possible among the effective LQC solutions of interest. In addition, in the case of \mathcal{P}_{LQC} , we show the result for three different choices of the transition time t_0 between the LQC and classical epochs. The behavior of the resulting spectra is only slightly sensitive to this choice, for scales k around k_{LQC} . More importantly, we see that for $k < 2k_{\text{LQC}}$, the primordial power \mathcal{P}_{LQC} gets strongly suppressed, in sharp contrast with the scale-invariant behavior of \mathcal{P}_{kin} , that is only lost when $k < 2k_I$.

The plots in figure 4 are restricted to the interval $k \in [10^{-4}, 10^2]$. For the considered effective LQC solutions with phenomenological application, this interval covers (with some margin in the infrared region and except for the most ultraviolet part in some cases) the window of wavenumbers that are directly observable nowadays [30, 32, 33]. Roughly speaking, this window corresponds to $10^{-4} \text{Mpc}^{-1} \lesssim k/a_{\text{today}} \lesssim 10^{-1} \text{Mpc}^{-1}$, where a_{today} is the current value of the scale factor [4]. Actually, the lower end of the observable interval of k depends on the specific LQC solution that one selects. In particular, for those solutions which accumulate more e-folds from the bounce till the present, this end can be relatively large and range in $[10^{-1}, 1]$ [30, 32], hence excluding the scale of inflation k_I from the observable interval.

To our degree of approximation, the region where we have found power suppression, starting at distinct values of k , is the most important difference between the power spectra obtained in effective LQC and in purely classical backgrounds with short-lived inflation. It depends strongly on the difference between the scales k_{LQC} and k_I , which are related to the

spacetime curvature around the vicinity of the bounce where LQC corrections are important and to the onset of inflation in relativistic cosmology, respectively. In fact, the order of magnitude of both of these scales is fixed for background solutions in effective LQC such that the largest sector of observable angular scales in the CMB may have been affected by quantum geometry effects. The suppression for $k < k_{\text{LQC}}$ can be interpreted as a consequence of the vacuum excitation of these scales with respect to the Bunch-Davies state, owing to the presence of the bounce. Furthermore, the later period of decelerated expansion does not de-excite these scales.

Finally, let us remark that the power spectra \mathcal{P}_{LQC} and \mathcal{P}_{kin} have been obtained using the same procedure for the choice of vacuum state. That is, starting with eqs. (4.1) and (4.3), applied in the earliest period of the cosmological evolution, we have fixed the respective vacua by selecting integration constants $(A_k^{\text{LQC}}, B_k^{\text{LQC}})$ and $(A_k^{\text{kin}}, B_k^{\text{kin}})$ that avoid, in a specific way, the oscillations that amplify the resulting power spectrum. In this sense, the differences between both power spectra shown in our analysis reflect genuine traces of the loop quantum bounce, distinguishing them from classical pre-inflationary effects.

5 Discussion

This work is the first analytic comparison between the main effects on the primordial power spectrum arising from a single-field inflationary cosmology with a fast-roll period within GR, on the one hand, and from the cosmological evolution of effective LQC backgrounds of phenomenological interest, on the other hand. In this last case, we have focused our full attention on the dynamical equations of the Mukhanov-Sasaki and tensor perturbations that result from the hybrid loop quantization approach. These equations have the same structure as in GR, but with a different and very specific time-dependent mass. Both of the considered cosmological backgrounds (i.e., the classical relativistic and the effective LQC backgrounds) for the propagation of the perturbations coincide a few Planck times after the instant that would correspond in effective LQC to the bounce. They display a classical kinetically dominated era that transitions into a slow-roll inflation that is short-lived.

In order to study the problem analytically, as a first approximation, we have ignored any transition effects between the kinetic and slow-roll regimes, and described the classical cosmological evolution as a Friedmann solution with a massless scalar field that is instantaneously matched with a de Sitter spacetime. Furthermore, in the LQC case we have approximated the time-dependent mass function for the dynamics of the perturbations in the vicinity of the bounce by some Pöschl-Teller potential, selected to improve similar approximations that have been previously proposed in the literature [40]. We have then derived general analytic formulas to characterize any choice of vacuum state for the perturbations and its resulting primordial power spectrum, for both of our considered classical and LQC cosmologies.

The form and properties of this power spectrum depend critically on the specific choice of vacuum state. It is then evident that a robust comparison between the effects of the different cosmological evolutions on the spectrum requires a common criterion for the choice of such a state. Inspired by recent studies in hybrid LQC [56, 57], and by the physical interest of attaining a non-(rapidly-)oscillating power spectrum [33, 55], we have proposed a criterion that is applicable for both of our types of backgrounds (classical and in effective LQC). The starting point in both cases is to choose the state that results from imposing an asymptotic, time-dependent (non-instantaneous) diagonalization of the Hamiltonian during the earliest period of cosmological evolution in the respective approximate models. This procedure can

be physically motivated from hybrid LQC, and its application only finds one obstruction for our background models, namely the discontinuities that our approximations introduce in the matching between the different epochs. These discontinuities might introduce some rapidly changing frequencies in the vacuum solutions, that have to adapt themselves fast to the discontinuities of the evolution. Therefore, one should expect that the resulting power spectrum can present some undesired properties, such as the kind of unwanted oscillations that we try to avoid. Nonetheless, it is possible to redefine the considered choice of vacuum by inspecting the resulting primordial spectrum and eliminating the oscillations that amplify the power (in the sense explained in section 2), which happen to be easily identifiable in this situation. The power spectrum obtained in this way captures the essential properties of the originally chosen vacuum that do not get artificially altered by those oscillations.

The main differences that our analysis finds between both power spectra (namely, that arising from a purely classical GR background with fast-roll effects and that from effective LQC) are notorious. They depend on the difference between two natural scales of the cosmological models, k_I and k_{LQC} , that are respectively related to the spacetime curvature at the onset of inflation and at the bounce. With our choice of vacuum state for the perturbations, the power spectrum of the studied classical backgrounds does not oscillate in the Fourier scale k , and displays a drastic infrared suppression for $k < k_I$, while displaying scale-invariance for larger k . This scale-invariant behavior in the ultraviolet combined with a strong suppression in the infrared has already been reported in past investigations, e.g. in refs. [6, 10, 18, 20–22], however often using choices of vacuum state that lead to an artificially oscillating power for $k \gtrsim k_I$. On the other hand, our non-oscillating spectrum corresponding to the effective LQC cosmologies (in the sector of phenomenological interest) has strongly suppressed power for $k < k_{\text{LQC}}$ and is scale-invariant for larger k . Since all those cosmologies must involve a (physically significant) classical kinetic epoch prior to slow-roll inflation, it follows that $k_{\text{LQC}} \gg k_I$ for any of them. Therefore, we conclude that the main effects of the loop quantum bounce on the primordial perturbations translate into a huge infrared suppression in their spectrum (viewable in practice as if there existed a cutoff) that covers a much wider region of scales than it would if the perturbations were solely affected by a kinetically dominated pre-inflationary dynamics within GR. In this sense, it is worth commenting that, for the set of effective LQC solutions that we have considered (as well as for their analogues within GR), the scale of inflation k_I lies either in the very lower end of the window of directly observable scales k , or just out of it for those backgrounds with the largest number of e-folds of expansion. One would then expect that the net suppression on the angular power spectrum of temperature anisotropies should turn out to be much greater if one takes into account all of the LQC effects rather than only the classical fast-roll ones.

Although yet very approximate, it is enlightening to compare our analytic results with previous ones attained in the literature of hybrid LQC by solving numerically the exact equations for the cosmological background and perturbations [31, 33]. There, the predicted power spectrum is also non-oscillating and displays a clear infrared suppression. However, this strong power suppression starts at a Fourier scale that is of the typical order of $k_I \sim 10^{-3}$, rather than that of k_{LQC} . In view of our conclusions, this behavior seems to indicate that the choice of vacuum state adopted in those works is only capable of capturing the classical fast-roll effects on the evolution of the perturbations, but not the LQC effects associated with a bounce in the high-curvature regime. Other remarkable investigations in LQC [32, 34] have also predicted an infrared suppression in the power spectrum, obtained after averaging its fast superimposed oscillations. In this case, the suppression does start from a Fourier scale that

can be related to the spacetime curvature at the bounce. However, the theoretical framework underlying these works is rather different from the hybrid approach. It is usually called the dressed-metric formalism [58–60], and the corresponding time-dependent mass in effective LQC is not the same. For the effective LQC solutions that we have considered, perhaps the most remarkable difference is the negativity of this mass at the bounce, compared to the positivity of $s_h^{(r)}$. It would be interesting to investigate if one can apply similar approximations and choice of vacuum state to analytically study the dressed-metric case, comparing the resulting power spectrum with the hybrid one.

Finally, it is worth commenting on the fact that, because of the simplicity of our approximations, we have been forced to ignore in this work all relevant inflationary aspects related with quantities such as the spectral index and the tensor-to-scalar ratio, which reflect the departure of the slow-roll period from an exact de Sitter evolution. This mainly depends on the specific form of the inflaton potential. We believe that the analytical techniques presented in this work can be further developed to incorporate such inflationary aspects, taking into account the typical values of the slow-roll parameters in the effective LQC solutions of interest. This kind of studies would then provide a more realistic power spectrum for potential comparison with the CMB observations, while facilitating a clear understanding of the reasons underlying its behavior.

A Fixing a vacuum state with the asymptotic expansion of h_k

Let us first show how we can fix the vacuum state associated with the choice of normalized mode solution (2.13) with $C_k = 0$ and $D_k = 1$ in the kinetically dominated interval (η_0, η_i) , by applying eqs. (4.1) and (4.3). In this period, the function h_k satisfies by construction

$$h'_k = k^2 + \frac{1}{4y^2} + h_k^2, \quad y = \eta - \eta_0 + \frac{1}{2H_0 a_0}. \quad (\text{A.1})$$

If we introduce the change of function

$$h_k = -k \left[\frac{1}{2T_k} + \frac{d}{dT_k} (\log \sigma) \right], \quad T_k = ky, \quad (\text{A.2})$$

we are led to the Bessel equation

$$T_k^2 \frac{d^2 \sigma}{dT_k^2} + T_k \frac{d\sigma}{dT_k} + T_k^2 \sigma = 0. \quad (\text{A.3})$$

The general solution of this equation can be expressed as a linear combination of the Hankel functions of zeroth order $H_0^{(1)}(ky)$ and $H_0^{(2)}(ky)$. On the other hand, one can check that the asymptotic restriction imposed by eqs. (4.3) and (4.4) on h_k implies

$$h_k = -ik [1 + \mathcal{O}(k^{-2})], \quad (\text{A.4})$$

where the subdominant terms $\mathcal{O}(k^{-2})$ must form a series in inverse powers of $T_k = ky$. Comparing this condition on h_k with eq. (A.2), and integrating the resulting equality in the asymptotic regime $k \rightarrow \infty$, we have that σ must admit an asymptotic expansion of the form

$$\sigma = e^{iT_k} (T_k)^{-1/2} \sum_{n=0}^{\infty} (T_k)^{-n} \sigma_n \quad (\text{A.5})$$

when $T_k \rightarrow \infty$, where σ_n are certain constants. Recalling the asymptotic behavior of the Hankel functions for large arguments, given in eq. (4.6), we conclude that the only solution of eq. (A.3) with the desired asymptotic behavior is

$$\sigma = \sigma_0 \sqrt{\frac{\pi}{2}} e^{i\pi/4} H_0^{(1)}(ky), \quad (\text{A.6})$$

for arbitrary constant σ_0 . Therefore, the asymptotic eqs. (4.3) and (4.4) allow us to fix h_k as

$$h_k = k \frac{H_1^{(1)}(ky)}{H_0^{(1)}(ky)} - \frac{1}{2y}. \quad (\text{A.7})$$

This selects a specific normalized solution μ_k of the dynamical eq. (2.7) for the perturbations via eq. (4.1). Explicitly, one can check that the squared norm of this solution is

$$|\mu_k|^2 = \frac{\pi y}{4} |H_0^{(2)}(ky)|^2. \quad (\text{A.8})$$

In the derivation of this formula, we have used the identity [45]

$$H_1^{(1)}(ky)H_0^{(2)}(ky) - H_0^{(1)}(ky)H_1^{(2)}(ky) = -\frac{4i}{\pi ky}. \quad (\text{A.9})$$

Finally, the phase of the mode solution can be determined in eq. (4.1) noting that

$$\text{Im}(h_k) = -\arg [H_0^{(1)}(ky)]' = \arg [H_0^{(2)}(ky)]'. \quad (\text{A.10})$$

Thus, our asymptotic expansion for h_k indeed fixes the vacuum state associated with the mode solutions (2.13) with $C_k = 0$ and $D_k = 1$, up to a constant irrelevant phase.

Let us now show that the asymptotic eqs. (4.3) and (4.4) fix as well the vacuum state associated with our normalized mode solutions in the interval $[\eta_B, \eta_0]$ near the bounce. We recall that these mode solutions are obtained by inserting in eq. (4.1) the function h_k given in eq. (4.13). In this case, h_k satisfies by construction that

$$h_k' = k^2 + \frac{U_0}{\cosh^2 \alpha(\eta - \eta_B)} + h_k^2, \quad (\text{A.11})$$

where U_0 and α were defined in eq. (3.10). If we introduce the change of function

$$h_k = ik(1 - 2x) - 2\alpha x(1 - x) \frac{d}{dx} (\log u_k), \quad (\text{A.12})$$

with x given in eq. (3.16), then we obtain the hypergeometric equation

$$x(1 - x) \frac{d^2 u_k}{dx^2} + [b_3^k - (b_1^k + b_2^k + 1)x] \frac{du_k}{dx} - b_1^k b_2^k u_k = 0. \quad (\text{A.13})$$

The parameters b_1^k , b_2^k , and b_3^k are defined in general as

$$b_3^k = 1 - \frac{ik}{\alpha} \quad (\text{A.14})$$

$$b_1^k = \frac{1}{2} \left(1 + \sqrt{1 + \frac{4U_0}{\alpha^2}} \right) - \frac{ik}{\alpha}, \quad b_2^k = \frac{1}{2} \left(1 - \sqrt{1 + \frac{4U_0}{\alpha^2}} \right) - \frac{ik}{\alpha}. \quad (\text{A.15})$$

Clearly, they reproduce those given in eqs. (3.17) and (3.18) for $U_0 = 8\pi\rho_c/3$. The general solution of eq. (A.13) is a linear combination of the two hypergeometric functions [45]

$${}_2F_1 \left(b_1^k, b_2^k; b_3^k; x \right), \quad x^{ik/\alpha} {}_2F_1 \left(b_1^k - b_3^k + 1, b_2^k - b_3^k + 1; 2 - b_3^k; x \right). \quad (\text{A.16})$$

In our case, we have

$$b_1^k - b_3^k + 1 = \frac{1}{2} \left(1 + \sqrt{1 + \frac{4U_0}{\alpha^2}} \right), \quad b_2^k - b_3^k + 1 = \frac{1}{2} \left(1 - \sqrt{1 + \frac{4U_0}{\alpha^2}} \right). \quad (\text{A.17})$$

Again, for $U_0 = 8\pi\rho_c/3$ these formulas reproduce the respective values of the constants c and d in eq. (4.14).

Taking into account that, in terms of x ,

$$\frac{U_0}{\cosh^2 \alpha(\eta - \eta_B)} = 4U_0 x(1 - x), \quad (\text{A.18})$$

it is easy to check that the asymptotic behavior required by eqs. (4.3) and (4.4) for h_k implies

$$h_k = -ik \left[1 + \frac{2U_0}{k^2} x(1 - x) + \mathcal{O}(k^{-3}) \right]. \quad (\text{A.19})$$

Furthermore, the subdominant terms $\mathcal{O}(k^{-3})$ must depend polynomially on x . If we introduce this behavior in eq. (A.12) and integrate the result in the asymptotic regime of large k , we get that u_k must admit an asymptotic expansion of the form

$$u_k = x^{ik/\alpha} \sum_{n=0}^{\infty} v_n^k x^n, \quad (\text{A.20})$$

where the complex constants v_n^k are linear combinations of inverse powers of k . At this stage, we recall that $x < 1$ by the very construction of this variable, and that the hypergeometric functions that appear in the solutions of eq. (A.13) admit representations as power series of x that are convergent precisely when $x < 1$ [45]. In view of this, we impose that the behavior (A.20) holds for all k . The only solution of eq. (A.13) that can depend on x as in eq. (A.20) is

$$u_k = v_0^k x^{ik/\alpha} {}_2F_1 \left(b_1^k - b_3^k + 1, b_2^k - b_3^k + 1; 2 - b_3^k; x \right). \quad (\text{A.21})$$

Introducing this solution in eq. (A.12), we finally obtain

$$h_k = -ik - 2\alpha x(1 - x) \frac{(b_1^k - b_3^k + 1)(b_2^k - b_3^k + 1)}{2 - b_3^k} \frac{{}_2F_1 \left(b_1^k - b_3^k + 2, b_2^k - b_3^k + 2; 3 - b_3^k; x \right)}{{}_2F_1 \left(b_1^k - b_3^k + 1, b_2^k - b_3^k + 1; 2 - b_3^k; x \right)}, \quad (\text{A.22})$$

which indeed coincides with eq. (4.13) when $U_0 = 8\pi\rho_c/3$.

Acknowledgments

This work was partially supported by Project. No. MICINN FIS2017-86497-C2-2-P from Spain (with extension Project. No. MICINN PID2020-118159GB-C41 under evaluation). B.E.N. acknowledges financial support from the Standard program of JSPS Postdoctoral Fellowships for Research in Japan.

References

- [1] A.R. Liddle and D.H. Lyth, *Cosmological Inflation and Large-Scale Structure*, Cambridge University Press (2000).
- [2] V. Mukhanov, *Physical Foundations of Cosmology*, Cambridge University Press (2005).
- [3] C.L. Bennett et al., *Nine-year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Final maps and results*, *ApJS* **208** (2013) 20.
- [4] N. Aghanim et al., *Planck 2018 results. VI. Cosmological parameters*, *A&A* **641** (2020) A6.
- [5] Y. Akrami et al., *Planck 2018 results. X. Constraints on inflation*, *A&A* **641** (2020) A10.
- [6] C.R. Contaldi, M. Peloso, L. Kofman, and A.D. Linde, *Suppressing the lower multipoles in the CMB anisotropies*, *JCAP* **0307** (2003) 002.
- [7] J.M. Cline, P. Crotty, and J. Lesgourgues, *Does the small CMB quadrupole moment suggest new physics?*, *JCAP* **0309** (2003) 010.
- [8] I. Wang and K. Ng, *Effects of a preinflation radiation-dominated epoch to CMB anisotropy*, *Phys. Rev. D* **77** (2008) 083501.
- [9] C. Destri, H.J. de Vega, and N.G. Sanchez, *CMB quadrupole depression produced by early fast-roll inflation: Monte Carlo Markov chains analysis of WMAP and SDSS data*, *Phys. Rev. D* **78** (2008) 023013.
- [10] C. Destri, H.J. de Vega, and N.G. Sanchez, *Preinflationary and inflationary fast-roll eras and their signatures in the low CMB multipoles*, *Phys. Rev. D* **81** (2010) 063520.
- [11] D. Boyanovsky, C. Destri, H.J. de Vega, and N.G. Sanchez, *The effective theory of inflation in the standard model of the Universe and the CMB+LSS data analysis*, *Int. J. Mod. Phys. A* **24** (2009) 3669.
- [12] R.K. Jain, P. Chingangbam, J. Gong, L. Sriramkumar, and T. Souradeep, *Punctuated inflation and the low CMB multipoles*, *JCAP* **01** (2009) 009.
- [13] F. Scardigli, C. Gruber, and P. Chen, *Black hole remnants in the early universe*, *Phys. Rev. D* **83** (2011) 063507.
- [14] M.H. Namjoo, H. Firouzjahi, and M. Sasaki, *Multiple inflationary stages with varying equation of state*, *JCAP* **12** (2012) 018.
- [15] E. Dudas, N. Kitazawa, S.P. Patil, and A. Sagnotti, *CMB imprints of a pre-inflationary climbing phase*, *JCAP* **05** (2012) 012.
- [16] S. Kouwn, O. Kwon, and P. Oh, *Large scale suppression of scalar power on a spatial condensation*, *Phys. Rev. D* **91** (2015) 063521.
- [17] P. Chen and Y. Lin, *What initial condition of inflation would suppress the large-scale CMB spectrum?*, *Phys. Rev. D* **93** (2016) 023503.
- [18] L.T. Hergt, W.J. Handley, M.P. Hobson, and A.N. Lasenby, *Constraining the kinetically dominated Universe*, *Phys. Rev. D* **100** (2019) 023501.

- [19] A. Di Tucci, J. Feldbrugge, J.-L. Lehners, and N. Turok, *Quantum Incompleteness of Inflation*, *Phys. Rev. D* **100** (2019) 063517.
- [20] E. Ramirez and D.J. Schwarz, *Predictions of just-enough inflation*, *Phys. Rev. D* **85** (2012) 103516.
- [21] E. Ramirez, *Low power on large scales in just-enough inflation models*, *Phys. Rev. D* **85** (2012) 103517.
- [22] A. Scacco and A. Albrecht, *Transients in finite inflation*, *Phys. Rev. D* **92** (2015) 083506.
- [23] M. Bojowald, *Loop Quantum Cosmology*, *Living Rev. Rel.* **11** (2008) 4.
- [24] A. Ashtekar, M. Bojowald, and J. Lewandowski, *Mathematical structure of loop quantum cosmology*, *Adv. Theor. Math. Phys.* **7** (2003) 233.
- [25] A. Ashtekar and P. Singh, *Loop quantum cosmology: A status report*, *Class. Quant. Grav.* **28** (2011) 213001.
- [26] A. Ashtekar, T. Pawłowski, and P. Singh, *Quantum nature of the big bang*, *Phys. Rev. Lett.* **96** (2006) 141301.
- [27] M. Martín-Benito, G.A. Mena Marugán, and J. Olmedo *Further improvements in the understanding of isotropic loop quantum cosmology*, *Phys. Rev. D* **80** (2009) 104015.
- [28] V. Taveras, *Corrections to the Friedmann equations from loop quantum gravity for a universe with a free scalar field*, *Phys. Rev. D* **78** (2008) 064072.
- [29] A. Bhardwaj, E.J. Copeland, and J. Louko, *Inflation in loop quantum cosmology*, *Phys. Rev. D* **99** (2019) 063520.
- [30] I. Agullo and N.A. Morris, *Detailed analysis of the predictions of loop quantum cosmology for the primordial power spectra*, *Phys. Rev. D* **92** (2015) 124040.
- [31] L. Castelló Gomar, G.A. Mena Marugán, D. Martín de Blas, and J. Olmedo, *Hybrid loop quantum cosmology and predictions for the cosmic microwave background*, *Phys. Rev. D* **96** (2017) 103528.
- [32] A. Ashtekar and B. Gupta, *Quantum gravity in the sky: Interplay between fundamental theory and observations*, *Class. Quant. Grav.* **34** (2017) 014002.
- [33] D. Martín de Blas and J. Olmedo, *Primordial power spectra for scalar perturbations in loop quantum cosmology*, *JCAP* **06** (2016) 029.
- [34] A. Ashtekar, B. Gupta, D. Jeong, and V. Sreenath, *Alleviating the tension in CMB using Planck-scale physics*, *Phys. Rev. Lett.* **125** (2020) 051302.
- [35] M. Fernández-Méndez, G.A. Mena Marugán, and J. Olmedo, *Hybrid quantization of an inflationary model: The flat case*, *Phys. Rev. D* **88** (2013) 044013.
- [36] L. Castelló Gomar, M. Fernández-Méndez, G.A. Mena Marugán, and J. Olmedo, *Cosmological perturbations in hybrid loop quantum cosmology: Mukhanov–Sasaki variables*, *Phys. Rev. D* **90** (2014) 064015.
- [37] L. Castelló Gomar, M. Martín-Benito, and G.A. Mena Marugán, *Gauge-invariant perturbations in hybrid quantum cosmology*, *JCAP* **06** (2015) 045.
- [38] F. Benítez Martínez and J. Olmedo, *Primordial tensor modes of the early universe*, *Phys. Rev. D* **93** (2016) 124008.
- [39] B. Elizaga Navascués, D. Martín de Blas, and G.A. Mena Marugán, *Time-dependent mass of cosmological perturbations in the hybrid and dressed metric approaches to loop quantum cosmology*, *Phys. Rev. D* **97** (2018) 043523.
- [40] Q. Wu, T. Zhu, and A. Wang, *Nonadiabatic evolution of primordial perturbations and non-Gaussianity in hybrid approach of loop quantum cosmology*, *Phys. Rev. D* **98** (2018) 103528.

- [41] V. Mukhanov, *Quantum theory of gauge-invariant cosmological perturbations*, *Zh. Eksp. Teor. Fiz.* **94**(1988) 1 [*Sov. Phys. JETP* **67** (1988) 1297].
- [42] M. Sasaki, *Gauge invariant scalar perturbations in the new inflationary universe*, *Prog. Theor. Phys.* **70** (1983) 394.
- [43] H. Kodama and M. Sasaki, *Cosmological perturbation theory*, *Prog. Theor. Phys. Suppl.* **78** (1984) 1.
- [44] D. Langlois, *Inflation and cosmological perturbations*, *Lect. Notes Phys.* **800** (2010) 1.
- [45] M. Abramovitz and I.A. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Natl. Bur. Stand. Appl. Math. Ser. No. 55, revised 9th ed. (1970).
- [46] A.R. Liddle and S.M. Leach, *How long before the end of inflation were observable perturbations produced?*, *Phys. Rev. D* **68** (2003) 103503.
- [47] T.S. Bunch and P. Davies, *Quantum field theory in de Sitter space: Renormalization by point splitting*, *Proc. R. Soc. Lond. A* **360** (1978) 117.
- [48] B. Elizaga Navascués, D. Martín de Blas, and G.A. Mena Marugán, *The vacuum state of primordial fluctuations in hybrid loop quantum cosmology*, *Universe* **4** (2018) 98.
- [49] T. Zhu, A. Wang, G. Cleaver, K. Kirsten, and Q. Sheng *Pre-inflationary universe in loop quantum cosmology*, *Phys. Rev. D* **96** (2017) 083520.
- [50] I. Agullo, W. Nelson, and A. Ashtekar *Preferred instantaneous vacuum for linear scalar fields in cosmological space-times*, *Phys. Rev. D* **91** (2015) 064051.
- [51] A. Ashtekar and B. Gupt, *Initial conditions for cosmological perturbations*, *Class. Quant. Grav.* **34** (2017) 035004.
- [52] M. Martín-Benito, R.B. Neves, and J. Olmedo, *States of Low Energy in bouncing inflationary scenarios in Loop Quantum Cosmology*, arXiv:2104.03035 (2021).
- [53] L. Parker, *Quantized fields and particle creation in expanding universes I*, *Phys. Rev.* **183** (1969) 1057.
- [54] C. Lüders and J.E. Roberts, *Local quasiequivalence and adiabatic vacuum states*, *Commun. Math. Phys.* **134** (1990) 29.
- [55] B. Elizaga Navascués, G.A. Mena Marugán, and S. Prado, *Non-oscillating power spectra in loop quantum cosmology*, *Class. Quant. Grav.* **38** (2021) 035001.
- [56] B. Elizaga Navascués, G.A. Mena Marugán, and T. Thiemann, *Hamiltonian diagonalization in hybrid quantum cosmology*, *Class. Quant. Grav.* **36** (2019) 18.
- [57] B. Elizaga Navascués and G.A. Mena Marugán, *Constraining quantization ambiguities from effective descriptions of cosmological perturbations in hybrid Loop Quantum Cosmology*, arXiv:2102.00124 (2021).
- [58] I. Agullo, A. Ashtekar, and W. Nelson, *A quantum gravity extension of the inflationary scenario*, *Phys. Rev. Lett.* **109** (2012) 251301.
- [59] I. Agullo, A. Ashtekar, and W. Nelson, *Extension of the quantum theory of cosmological perturbations to the Planck era*, *Phys. Rev. D* **87** (2013) 043507.
- [60] I. Agullo, A. Ashtekar, and W. Nelson, *The pre-inflationary dynamics of loop quantum cosmology: Confronting quantum gravity with observations*, *Class. Quant. Grav.* **30** (2013) 085014.