

# First evidence that non-metricity $f(Q)$ gravity can challenge $\Lambda$ CDM

Fotios K. Anagnostopoulos,<sup>1</sup> Spyros Basilakos,<sup>2,3</sup> and Emmanuel N. Saridakis<sup>2,4,5</sup>

<sup>1</sup>*Department of Physics, National & Kapodistrian University of Athens, Zografou Campus GR 157 73, Athens, Greece*

<sup>2</sup>*National Observatory of Athens, Lofos Nymfon, 11852 Athens, Greece*

<sup>3</sup>*Academy of Athens, Research Center for Astronomy and Applied Mathematics, Soranou Efessiou 4, 11527, Athens, Greece*

<sup>4</sup>*CAS Key Laboratory for Researches in Galaxies and Cosmology, Department of Astronomy, University of Science and Technology of China, Hefei, Anhui 230026, P.R. China*

<sup>5</sup>*School of Astronomy, School of Physical Sciences, University of Science and Technology of China, Hefei 230026, P.R. China*

We propose a novel model in the framework of  $f(Q)$  gravity, which is a gravitational modification class arising from the incorporation of non-metricity. The model has General Relativity as a particular limit, it has the same number of free parameters to those of  $\Lambda$ CDM, however at a cosmological framework it gives rise to a scenario that does not have  $\Lambda$ CDM as a limit. Nevertheless, confrontation with observations at both background and perturbation levels, namely with Supernovae type Ia (SNIa), Baryonic Acoustic Oscillations (BAO), cosmic chronometers (CC), and Redshift Space Distortion (RSD) data, reveals that the scenario, according to AIC, BIC and DIC information criteria, is at some cases statistically preferred comparing to  $\Lambda$ CDM cosmology. Finally, the model does not exhibit early dark energy features, and thus it immediately passes BBN constraints, while the variation of the effective Newton's constant lies well inside the observational bounds.

## Introduction

Although General Relativity (GR) is the well-established theory for the description of the gravitational interaction, there are two main motivations that justify the large amount of research devoted to its modification and extension. The first arises from cosmological grounds, since modified gravity is very efficient in describing the universe's two phases of accelerated expansion [1], and moreover it can alleviate the two possible tensions of  $\Lambda$ CDM cosmology, namely the  $H_0$  [2] and the  $\sigma_8$  [3] ones. The second, and chronologically older, motivation is purely theoretical, and aims towards the improvement of the renormalizability of General Relativity with the further goal to finally reach to a quantum gravitational theory [4]. Hence, the goal is to construct gravitational theories that possess General Relativity as a particular limit, but which in general include extra degree(s) of freedom that are able to fulfill the above requirements.

In this Letter we propose a specific model in the framework of the recently constructed  $f(Q)$  modified gravity, which as we show is very efficient in fitting the cosmological data. In this class of modification, one starts from the so-called symmetric teleparallel theories, which is an equivalent description of gravity using the *non-metricity* scalar  $Q$  [5], and extends it to an arbitrary function  $f(Q)$ .  $f(Q)$  gravity leads to interesting applications [6–14], and trivially passes the constraints arising from gravitational wave observations [15]. By confronting our new model with data from Supernovae type Ia (SNIa), Baryonic Acoustic Oscillations (BAO), Hubble parameter cosmic chronometers (CC) and Redshift Space Distortions (RSD)  $f\sigma_8$  observations, we deduce that the scenario at hand is, in some cases, statistically better than  $\Lambda$ CDM, although it does not include it as a particular limit.

## A novel model in $f(Q)$ gravity

The action of non-metricity-based modified gravity is [5]

$$S = -\frac{1}{16\pi G} \int d^4x \sqrt{-g} f(Q), \quad (1)$$

where the nonmetricity scalar

$$Q = -\frac{1}{4} Q_{\alpha\beta\gamma} Q^{\alpha\beta\gamma} + \frac{1}{2} Q_{\alpha\beta\gamma} Q^{\gamma\beta\alpha} + \frac{1}{4} Q_{\alpha} Q^{\alpha} - \frac{1}{2} Q_{\alpha} \tilde{Q}^{\alpha}, \quad (2)$$

with  $Q_{\alpha} \equiv Q_{\alpha}{}^{\mu}{}_{\mu}$  and  $\tilde{Q}^{\alpha} \equiv Q_{\mu}{}^{\mu\alpha}$ , arises from contractions of the non-metricity tensor  $Q_{\alpha\mu\nu} \equiv \nabla_{\alpha} g_{\mu\nu}$ . From these it is apparent that Symmetric Teleparallel Equivalent of General Relativity (and thus General Relativity) is obtained in the case of  $f(Q) = Q$ .

Varying the total action  $S + S_m$ , with  $S_m$  the matter sector action, one obtains the field equations as [6, 8]:

$$\begin{aligned} & \frac{2}{\sqrt{-g}} \nabla_{\alpha} \left\{ \sqrt{-g} g_{\beta\nu} f_Q \left[ -\frac{1}{2} L^{\alpha\mu\beta} + \frac{1}{4} g^{\mu\beta} (Q^{\alpha} - \tilde{Q}^{\alpha}) \right. \right. \\ & \quad \left. \left. - \frac{1}{8} (g^{\alpha\mu} Q^{\beta} + g^{\alpha\beta} Q^{\mu}) \right] \right\} \\ & + f_Q \left[ -\frac{1}{2} L^{\mu\alpha\beta} - \frac{1}{8} (g^{\mu\alpha} Q^{\beta} + g^{\mu\beta} Q^{\alpha}) \right. \\ & \quad \left. + \frac{1}{4} g^{\alpha\beta} (Q^{\mu} - \tilde{Q}^{\mu}) \right] Q_{\nu\alpha\beta} + \frac{1}{2} \delta_{\nu}^{\mu} f = T^{\mu}_{\nu}, \quad (3) \end{aligned}$$

with  $L^{\alpha}_{\mu\nu} = \frac{1}{2} Q^{\alpha}_{\mu\nu} - Q_{(\mu}{}^{\alpha}{}_{\nu)}$  the disformation tensor,  $T_{\mu\nu}$  the energy-momentum tensor, and  $f_Q \equiv \partial f / \partial Q$ . In order to apply it in a cosmological framework we impose a flat Friedmann-Robertson-Walker (FRW) metric of the form  $ds^2 = -dt^2 + a^2(t) \delta_{ij} dx^i dx^j$ , in which case equa-

tions (3) give rise to the two Friedmann equations [8]

$$6f_Q H^2 - \frac{1}{2}f = 8\pi G(\rho_m + \rho_r), \quad (4)$$

$$(12H^2 f_{QQ} + f_Q)\dot{H} = -4\pi G(\rho_m + p_m + \rho_r + p_r), \quad (5)$$

where  $H \equiv \dot{a}/a$  is the the Hubble function and with  $\rho_m$ ,  $\rho_r$  and  $p_m$ ,  $p_r$  the energy densities and pressures of the matter and radiation perfect fluids. Additionally, note that the nonmetricity scalar  $Q$  in an FRW background becomes  $Q = 6H^2$ . Finally, the equations close by the consideration of the matter and radiation conservation equations

$$\begin{aligned} \dot{\rho}_m + 3H(\rho_m + p_m) &= 0 \\ \dot{\rho}_r + 3H(\rho_r + p_r) &= 0. \end{aligned} \quad (6)$$

Proceeding to the perturbation level, elaborating the full perturbation equations [8] one can extract the evolution equation for the matter overdensity,  $\delta \equiv \delta\rho_m/\rho_m$ , at sub-horizon scales in terms of the scale factor as

$$\delta''_m + \left(\frac{H'}{H} + \frac{3}{a}\right)\delta'_m = \frac{3\Omega_{m0}}{2H^2 a^5} \frac{G_{\text{eff}}}{G} \delta_m, \quad (7)$$

with primes denoting derivatives with respect to the scale factor, and where we have introduced the density parameters through  $\Omega_i \equiv \frac{8\pi G\rho_i}{3H^2}$ , with the subscript ‘‘0’’ denoting the value at present time. Finally,  $G_{\text{eff}}$  is the effective Newton’s constant in  $f(Q)$  gravity, which is given as  $G_{\text{eff}} \equiv G/f_Q$  [8].

In this work we propose the following model

$$f(Q) = Qe^{\lambda\frac{Q_0}{Q}}, \quad (8)$$

where  $\lambda$  is the sole model free parameter, and  $Q_0 = 6H_0^2$  with  $H_0$  the current value of the Hubble parameter. For  $\lambda = 0$  GR is recovered but not  $\Lambda$ CDM since the cosmological constant is absent, thus this model alleviates the cosmological constant problem. Note that in certain periods of cosmic history, as the term  $Q_0/Q$  decreases, our model effectively reduces to the polynomial case  $f(Q) = Q^n$ . Thus, in a sense, it could be thought as an encapsulation of many  $f(Q)$  models, where at a given cosmic time a term with particular  $n$  becomes dominant. Using (8) and (4), and considering  $w_m \equiv p_m/\rho_m = 0$  and  $w_r \equiv p_r/\rho_r = 1/3$ , the corresponding normalized Hubble parameter  $E^2 \equiv H^2/H_0^2$  is written as

$$(E^2 - 2\lambda)e^{\lambda/E^2} = \Omega_{m0}a^{-3} + \Omega_{r0}a^{-4}. \quad (9)$$

Applying the above equation at present, the parameter  $\lambda$  can be expressed as

$$\lambda = 0.5 + \mathcal{W}_0 \left( \frac{\Omega_{m0} + \Omega_{r0}}{2e^{1/2}} \right), \quad (10)$$

with  $\mathcal{W}_0$  the principal branch of the Lambert function. Hence, the scenario at hand has exactly the same number

of free parameters with  $\Lambda$ CDM, and as we will see this is a key reason for its statistical preference over the latter. Lastly, the effective Newton’s constant becomes

$$G_{\text{eff}} = \frac{G}{e^{\lambda\frac{Q_0}{Q}} \left(1 - \lambda\frac{Q_0}{Q}\right)}. \quad (11)$$

### Data, Methodology and Results

We employ Bayesian analysis and the likelihood function  $\mathcal{L}_{\text{tot}} \sim \exp(-\chi_{\text{tot}}^2/2)$ , where  $\chi_{\text{tot}}^2$  is obtained from different sums of  $\chi_{\text{SNia}}^2, \chi_{\text{BAOs}}^2, \chi_{\text{CC}}^2, \chi_{\text{RSD}}^2$ , to be specified below. The quantities  $\chi_{\text{SNia}}^2, \chi_{\text{RSD}}^2$  and  $\chi_{\text{CC}}^2$  are defined and described in [16]. Since RSD data were extracted by imposing  $\Lambda$ CDM as a reference model, we use the simple correction factor described in [17]. Additionally, we utilize the BAOs dataset of [18], without the last three points, that they do not have fiducial cosmology correction (i.e the term  $r_d/r_{\text{fid}}$ ). We sample the posterior distribution of the parameters applying the MCMC method as implemented within the open-source Python package emcee [19], for the following cases:

1. SNIa+CC datasets, with free parameters  $\Omega_{m0}, h, \mathcal{M}$ .
2. SNIa+CC+BAOs, with free parameters  $\Omega_{m0}, h, r_d, \mathcal{M}$ .
3. SNIa+CC+RSD, with free parameters  $\Omega_{m0}, h, \sigma_8, \mathcal{M}$ .

Note that  $\mathcal{M}$  is the intrinsic free parameter of Pantheon dataset (see [16] and references therein) and we neglect  $\Omega_{r0}$ . We apply the aforementioned setup for our  $f(Q)$  model (8), along with  $\Lambda$ CDM to allow for a direct comparison. In all cases 1000 walkers and 2500 states are employed along with flat priors on the parameters. Although the particular  $f(Q)$  model considered here has the same number of free parameters with  $\Lambda$ CDM, the functional form of the Hubble rate is different and thus the relative fitting quality could only be compared in the context of information criteria differences.

We present the results on the parameters in Table I. Additionally, in Fig. 1 we show the corresponding contour plots for the extracted model parameters for all considered datasets. Finally, in Fig. 2 we depict the contour plots for both the  $f(Q)$  and  $\Lambda$ CDM models, for the case of Pantheon+CC+BAOs datasets. Concerning the parameter values, in all cases  $\Omega_{m0}$  is close to the Planck value, while the deviations from the concordance model is evident. Moreover, as a consistency check, we compare the sound horizon at baryon drag epoch,  $r_d$ , with the model-independent one from [20], and we observe  $1\sigma$  compatibility. Note that concerning the  $H_0$  and  $\sigma_8$  tensions, in order to be able to extract a safe results we should

Model	$\Omega_{m0}$	$h$	$r_d$	$\sigma_8$	$\mathcal{M}$	$\chi^2_{\min}$	$\chi_{\min}/\text{dof}$
SNIa/CC							
$Qe^{\lambda \frac{Q_0}{Q}}$	$0.349^{+0.022}_{-0.021}$	$0.6917 \pm 0.011$	—	—	$-19.384 \pm 0.027$	39.318	0.578
$\Lambda\text{CDM}$	$0.296^{+0.022}_{-0.021}$	$0.6992^{+0.0117}_{-0.0115}$	—	—	$-19.355 \pm 0.027$	39.297	0.578
SNIa/CC/BAOs							
$Qe^{\lambda \frac{Q_0}{Q}}$	$0.356 \pm 0.020$	$0.6882^{+0.0105}_{-0.0103}$	$146.646^{+1.550}_{-1.541}$	—	$-19.392 \pm 0.025$	41.615	0.555
$\Lambda\text{CDM}$	$0.306 \pm 0.021$	$0.6941^{+0.0108}_{-0.0116}$	$144.892^{+1.558}_{-1.550}$	—	$-19.366 \pm 0.025$	42.737	0.570
SNIa/CC/RSD							
$Qe^{\lambda \frac{Q_0}{Q}}$	$0.339^{+0.021}_{-0.020}$	$0.6971 \pm 0.011$	—	$0.703^{+0.029}_{-0.029}$	$-19.372 \pm 0.026$	54.526	0.613
$\Lambda\text{CDM}$	$0.290^{+0.021}_{-0.020}$	$0.7024^{+0.0113}_{-0.0110}$	—	$0.772^{+0.034}_{-0.034}$	$-19.348 \pm 0.026$	52.300	0.588

TABLE I. Observational constraints and the corresponding  $\chi^2_{\min}$  for the new  $f(Q)$  gravity model (8), where “dof” stands for degrees of freedom (defined as the number of the used data points minus the number of fitted parameters). In order to allow direct comparison the concordance  $\Lambda\text{CDM}$  model is also included.




FIG. 1. The  $1\sigma$  and  $2\sigma$  iso-likelihood contours for the  $f(Q)$  model (8), for the 2D subsets of the parameter space  $(\Omega_{m0}, h, \mathcal{M})$ , using graphic package *getdist* [21]. We have used joint analysis of various datasets (see text).

analyze the full CMB spectrum and compare the corresponding parameter values with the current ones. Lastly, we mention that for the case of Pantheon+CC+BAOs dataset the parameter  $\lambda$  is found as  $\lambda = 0.598 \pm 0.005$  at  $1\sigma$  confidence level.

Proceeding forward, in Table II we compare the fitting quality of the present  $f(Q)$  model with the corresponding one of  $\Lambda\text{CDM}$  cosmology, through the use of the Akaike Information Criterion (AIC), the Bayesian Information Criterion (BIC), and the Deviance Information Criterion




FIG. 2. The  $1\sigma$  and  $2\sigma$  iso-likelihood contours for the  $f(Q)$  model (8), as well as for the  $\Lambda\text{CDM}$  scenario, for the 2D subsets of the parameter space  $(\Omega_{m0}, h, \mathcal{M}, r_d)$ , using graphic package *getdist* [21]. We have used the joint analysis of SNIa+CC+BAOs datasets (see text).

(DIC), see [16] and references therein. For SNIa + CC datasets, we find that  $\Delta\text{IC}$  is almost 0, and thus the two models are statistically compatible. For SNIa + BAOs + CC, our  $f(Q)$  model is slightly more preferred by the data. In contrast, for SNIa + BAOs + RSD the  $f(Q)$  model is deemed inferior by the data, however still statistically indistinguishable from  $\Lambda\text{CDM}$ . We mention here (without considering the possibility that the BAOs data may include numerical relics) that the fact that these

BAOs employ a free parameter to correct for the imposed cosmology, while the  $f\sigma_8$  data do not, could be an indication for circularity problem with the RSD data. Hence, although prominent, the present  $f(Q)$  model, since it has no  $\Lambda$ CDM limit, gets “punished” by those datasets that incorporate  $\Lambda$ CDM as a reference model. Finally, we mention an additional reason that our model is statistically preferable comparing to  $\Lambda$ CDM one, namely while it includes many models as subcases (i.e polynomial) at particular times within the cosmic history, it maintains the same number of free parameters with  $\Lambda$ CDM cosmology.

Model	AIC	$\Delta$ AIC	BIC	$\Delta$ BIC	DIC	$\Delta$ DIC
SNIa/CC						
$Qe^{\lambda \frac{Q_0}{Q}}$	45.588	0.068	52.916	0.829	45.329	0.048
$\Lambda$ CDM	45.656	0	52.086	0	45.281	0
SNIa/CC/BAOs						
$Qe^{\lambda \frac{Q_0}{Q}}$	50.156	0	59.092	0	49.623	0
$\Lambda$ CDM	51.278	1.122	60.215	1.123	50.722	1.098
SNIa/CC/RSD						
$Qe^{\lambda \frac{Q_0}{Q}}$	62.981	2.227	72.657	2.227	62.494	2.220
$\Lambda$ CDM	60.754	0	70.430	0	60.274	0

TABLE II. The information criteria AIC, BIC and DIC for the examined cosmological models, alongside the corresponding differences  $\Delta$ IC  $\equiv$  IC  $-$  IC<sub>min</sub>.

### Conclusions

We proposed a novel  $f(Q)$  model and we confronted it against observational data (SNIa, BAOs, CC and RSD). For CC + SNIa datasets the two models are statistically compatible, however for CC + SNIa + BAOs datasets the  $f(Q)$  model is statistically preferred comparing to  $\Lambda$ CDM one. On the other hand, in the case of RSD + CC + SNIa data, the  $\Lambda$ CDM paradigm is slightly preferred by the data, although the two models remain statistically equivalent.

In the large redshift limit (i.e at large  $E^2(z) \equiv H^2(z)/H_0^2$ ) the proposed  $f(Q)$  tends to  $Q$  and thus the scenario at hand tends to GR, hence it trivially passes the early universe constraints and in particular the BBN ones. Additionally, knowing the observational bounds of  $E^2(z)$  throughout the evolution, and using (10), from (11) we deduce that throughout the evolution  $|\frac{G_{\text{eff}}}{G} - 1|$  remains smaller than 0.1 and therefore it satisfies the observational constraints [22].

In summary, this result could be used as motivation for further study of the present model, as well as  $f(Q)$

gravity in general, as it constitutes one of the first alternatives to the concordance model that apart from being preferred by the data (at least by some datasets), it additionally possesses a Lagrangian description. Further studies on this model, using the full CMB and LSS spectra, weak lensing data and other datasets, could enlighten our findings and verify whether the present  $f(Q)$  model outperforms the concordance one or not.

- 
- [1] S. Capozziello and M. De Laurentis, Phys. Rept. **509**, 167-321 (2011).
  - [2] E. Di Valentino, L. A. Anchordoqui, O. Akarsu, Y. Ali-Haimoud, L. Amendola, N. Arendse, M. Asgari, M. Ballardini, S. Basilakos and E. Battistelli, *et al.* [[arXiv:2008.11284](https://arxiv.org/abs/2008.11284)] [[astro-ph.CO](https://arxiv.org/abs/2008.11284)].
  - [3] E. Di Valentino, L. A. Anchordoqui, O. Akarsu, Y. Ali-Haimoud, L. Amendola, N. Arendse, M. Asgari, M. Ballardini, S. Basilakos and E. Battistelli, *et al.* [[arXiv:2008.11285](https://arxiv.org/abs/2008.11285)] [[astro-ph.CO](https://arxiv.org/abs/2008.11285)].
  - [4] K. S. Stelle, Phys. Rev. D **16**, 953 (1977).
  - [5] J. Beltrán Jiménez, L. Heisenberg and T. Koivisto, Phys. Rev. D **98**, no.4, 044048 (2018).
  - [6] K. F. Dialektopoulos, T. S. Koivisto and S. Capozziello, Eur. Phys. J. C **79**, no.7, 606 (2019).
  - [7] B. J. Barros, T. Barreiro, T. Koivisto and N. J. Nunes, Phys. Dark Univ. **30**, 100616 (2020).
  - [8] J. Beltrán Jiménez, L. Heisenberg, T. S. Koivisto and S. Pekar, Phys. Rev. D **101**, no.10, 103507 (2020).
  - [9] F. Bajardi, D. Vernieri and S. Capozziello, Eur. Phys. J. Plus **135**, no.11, 912 (2020).
  - [10] I. Ayuso, R. Lazkoz and V. Salzano, Phys. Rev. D **103**, no.6, 063505 (2021).
  - [11] K. Flathmann and M. Hohmann, Phys. Rev. D **103**, no.4, 044030 (2021).
  - [12] N. Frusciante, Phys. Rev. D **103**, no.4, 044021 (2021).
  - [13] W. Khyllep, A. Paliathanasis and J. Dutta, [[arXiv:2103.08372](https://arxiv.org/abs/2103.08372)] [[gr-qc](https://arxiv.org/abs/2103.08372)].
  - [14] F. D’Ambrosio, M. Garg and L. Heisenberg, Phys. Lett. B **811**, 135970 (2020) [[arXiv:2004.00888](https://arxiv.org/abs/2004.00888)] [[gr-qc](https://arxiv.org/abs/2004.00888)].
  - [15] I. Soudi, G. Farrugia, V. Gakis, J. Levi Said and E. N. Saridakis, Phys. Rev. D **100**, no.4, 044008 (2019).
  - [16] F. K. Anagnostopoulos, S. Basilakos and E. N. Saridakis, Phys. Rev. D **100**, no.8, 083517 (2019).
  - [17] E. Macaulay, I. K. Wehus and H. K. Eriksen, Phys. Rev. Lett. **111**, no.16, 161301 (2013).
  - [18] J. Ryan, Y. Chen and B. Ratra, Mon. Not. Roy. Astron. Soc. **488**, no.3, 3844-3856 (2019).
  - [19] D. Foreman-Mackey, D. W. Hogg, D. Lang and J. Goodman, Publ. Astron. Soc. Pac. **125**, 306-312 (2013).
  - [20] B. S. Haridasu, V. V. Luković, M. Moresco and N. Vittorio, JCAP **10**, 015 (2018).
  - [21] Lewis, A. GetDist, *Monte Carlo sample analyzer*, Astrophysics Source Code Librar.
  - [22] S. Nesseris, G. Pantazis and L. Perivolaropoulos, Phys. Rev. D **96**, no.2, 023542 (2017) [[arXiv:1703.10538](https://arxiv.org/abs/1703.10538)] [[astro-ph.CO](https://arxiv.org/abs/1703.10538)].