

HOLOGRAPHIC ENTANGLEMENT NEGATIVITY IN FLAT SPACE GENERALIZED MINIMAL MASSIVE GRAVITY

M. R. Setare ¹, M. Koohgard ²

Department of Science,

Campus of Bijar, University of Kurdistan, Bijar, Iran

Abstract

In this paper we study the application of holographic entanglement negativity proposal for bipartite states in the 2d Galilean conformal field theory ($GCFT_2$) dual to bulk asymptotically flat spacetimes in the context of generalized minimal massive gravity (GMMG) model. $GCFT_2$ is considered on the boundary side of the duality and the bulk gravity is described by GMMG that is asymptotically symmetric under the Galilean conformal transformations. In this paper, the replica technique, based on the two-point and the four-point twist correlators, is utilized and the entanglement entropy and the entanglement negativity are obtained in the bipartite configurations of the system in the boundary. This paper generalizes similar studies of $Flat_3/GCFT_2$ holography in Einstein gravity and topologically massive gravity (TMG).

Keywords: Replica technique, GMMG, Entanglement negativity, Entanglement entropy, Galilean conformal field theory

¹E-mail: rezakord@ipm.ir

²E-mail: m.koohgard@modares.ac.ir

1 Introduction

Recently, the non-relativistic version of the duality between gravity theory and boundary quantum field theory has been attracted a lot of attention. In this paper, we would consider a version of nonrelativistic conformal group, named Galilean conformal (GC) group, which arises as parametric contraction of conformal group. Galilean conformal algebra (GCA) was explored in two dimensions by two copies of Virasoro algebra in non-relativistic limit [1]. One option for bulk description of 2d GCA, with asymmetric central charges would be generalized minimal massive gravity (GMMG) in three dimensions [2]. GMMG model providing a new example of a theory that avoids the bulk-boundary clash and therefore, as Minimal Massive Gravity (MMG) [3], the theory possesses both, positive energy excitations around the maximally AdS_3 vacuum as well as a positive central charge in the dual CFT. Such clash is present in the previously constructed gravity theories with local degrees of freedom in 2+1-dimensions, namely Topologically Massive Gravity [4, 5] and the cosmological extension of New Massive Gravity [6]. MMG supplements the field equations of Topologically Massive Gravity with a symmetric, ranked two-tensor containing up to second derivatives of the metric. GMMG model is a general version of MMG in which by adding the CS deformation term (stands for Chern-Simons term for the one-form Levi-Civita affine connection which produces a propagating massive graviton), and an extra term to pure Einstein gravity with a negative cosmological constant. Since these theories avoid the bulk-boundary clash, they define excellent arenas to explore the structure of asymptotically AdS solutions, asymptotic symmetries, their algebra and other holographically inspired questions. We study some aspects of a class of non-AdS holography where the $3d$ bulk gravity is given by GMMG. We investigate a derivation of the holographic entanglement negativity using $flat_3/GCFT_2$ in the context of GMMG.

The study of the quantum entanglement of many-body systems has increased the knowledge of these systems (see for a review [7]). The quantum entanglement has an important role to study of a wide variety of the fields in physics from the condensed matter systems to quantum gravity. One of the most notable measures of quantum entanglement is the entanglement entropy [8]. The entanglement entropy can be defined by the von Neumann entropy of the corresponding reduced density matrix of a subsystem of a bipartite quantum system. When the whole system is in a state $|\Psi\rangle$ with the density matrix $\rho = |\Psi\rangle\langle\Psi|$, the reduced density matrix of A is $\rho_A = Tr_B \rho$. The observer A can observe only the subset of a complete set of commuting observables. The other observer B can observe the remainder of the system. Using the von Neumann entropy, the entanglement entropy is $S_A = -Tr_A \rho_A \log \rho_A$. This approach can not give a proper measure about the entanglement between two non-complementary subsections in a mixed state [9].

Some logarithmic measures of the entanglement in the bipartite states has been introduced in [10, 11, 12, 13], but these measures have some problems to be evaluated analytically. Another measure of the entanglement has been introduced in [14] (see also [15]) that is a computable one. This measure is the *entanglement negativity* that can be obtained from

any mixed state. A classical approach to compute the negativity in QFT and many-body systems has been introduced in [9, 16]. This method provides a good measure for entanglement between two parts of a larger system in a pure state and has been used a lot to study the ground-states of CFTs [17, 18, 19, 20]. The authors in [21] extended the works of [22, 23] on the entanglement entropy of GCFTs to compute the entanglement negativity of the $GCFT_2$. The replica technique has been used to compute the entanglement entropy of the bipartite configurations of $GCFT_2$ in [21]. It can be seen that the universal features of the entanglement negativity in CFT_2 can be reproduced in $GCFT_2$ in [21]. In [24], a covariant construction in calculation of the entanglement negativity has been introduced in the context of bipartite states in the duality $GCFT_2$ /TMG where this is in a class of flat-holography. It can be seen that the universal part of the entanglement negativity is same as the universal part of the negativity calculated in [21].

In this paper, we extend the measurement of the quantum entanglement to another duality in the flat holography that the bipartite system can be considered in the boundary side of the duality and the GMMG is the gravity theory in the bulk. To compute the entanglement entropy and the entanglement negativity utilizing the replica technique, it is crucial to find out the two-point and four-point twist correlators for the bipartite configuration of the subsystem. We find a viable form of the correlators in the $GCFT_2$ /GMMG that can be obtained by comparing the result of the replica technique for the entanglement entropy with the result in our previous work in [25]. The equality of the universal features of the entanglement negativity in this work matches with the universal parts of the negativity in [21].

This paper is organized as follows. In section 2 the general features of the GCA is presented. We give the essential tools that is needed to utilize the replica technique in this context. We present the argument to find out the two-point twist correlator in the GMMG by asymptotically flat geometry, in section 3. The first case we consider is related to a single interval configurations. These configurations are considered in section 4 on the zero and finite temperature and on the finite-sized systems. The bipartite configurations with two adjacent intervals and with two disjoint intervals are considered in sections 5 and 6, respectively. We conclude in section 7 with a summary of our results.

2 Replica approach in $GCFT_2$

The replica techniques have been used to compute the entanglement entropy and the entanglement negativity in some special cases of the flat holography. Also this technique has been used to compute the entanglement negativity in a CFT_2 [9, 16]. Another application of the replica technique can be found in the computation of the entanglement entropy in Galilean conformal field theories ($GCFT_2$) in [22, 23]. The two dimensional Galilean conformal field theories [26, 27, 1, 28] are a class of conformal field theories with Galilean invariance. The $GCFT_2$ involves the Galilean conformal algebra (GCA) that may be obtained via usual

conformal algebra in two dimensions as follows

$$t \rightarrow t, \quad x \rightarrow \epsilon x, \quad (2.1)$$

with $\epsilon \rightarrow 0$. The generators of GCA_2 in the plane coordinates are given as follows

$$L_n = t^{n+1}\partial_t + (n+1)t^n x \partial_x, \quad M_n = t^{n+1}\partial_x, \quad (2.2)$$

where these generators satisfy the following Lie algebra with a central extension

$$\begin{aligned} [L_n, L_m] &= (m-n)L_{n+m} + \frac{c_L}{12}(n^3-n)\delta_{n+m}, \\ [L_n, M_m] &= (m-n)M_{n+m} + \frac{c_M}{12}(n^3-n)\delta_{n+m}, \\ [M_n, M_m] &= 0, \end{aligned} \quad (2.3)$$

where c_L and c_M are the central charges of the algebra GCA_2 . To compute the entanglement entropy, the replica manifold can be considered as an n -sheeted surface. This surface involves n copies of the $GCFT_2$ plane glued together. The reduced density matrix ρ_A^n corresponding to the interval A can be considered as follows [9]

$$Tr \rho_A^n = \langle \Phi_n(\partial_1 A) \Phi_{-n}(\partial_2 A) \rangle, \quad (2.4)$$

where the twist fields Φ_n are primary fields of the $GCFT_2$ with scaling dimensions as follows

$$\Delta_n = \frac{c_L}{24}(n - \frac{1}{n}), \quad \chi_n = \frac{c_M}{24}(n - \frac{1}{n}). \quad (2.5)$$

Now, the entanglement entropy for the bipartite state corresponding to the interval A can be computed by taking the replica limit as follows [9]

$$S_A = - \lim_{n \rightarrow 1} \frac{\partial}{\partial n} Tr \rho_A^n. \quad (2.6)$$

To define the entanglement negativity as [14], a tripartite system is considered that consists of the subsystems A_1 , A_2 and B where $A = A_1 \cup A_2$ and $B = A^c$ represent the rest of the system. The entanglement negativity for the bipartite mixed state $A = A_1 \cup A_2$ can be defined as follows

$$\mathcal{E} = \log Tr \|\rho_A^{T_2}\|, \quad (2.7)$$

where $\rho_A^{T_2}$ is the partial transpose of the reduced density matrix for the system A with respect to the subsystem A_2 that can be defined as follows

$$\langle e_i^{(1)} e_j^{(2)} | \rho_A^{T_2} | e_k^{(1)} e_l^{(2)} \rangle = \langle e_i^{(1)} e_l^{(2)} | \rho_A | e_k^{(1)} e_j^{(2)} \rangle, \quad (2.8)$$

where $|e_i^{(1)}\rangle$ and $|e_j^{(2)}\rangle$ are the bases for the Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 . The replica technique can be used to compute the entanglement negativity \mathcal{E} for the bipartite mixed state A as follows [21]

$$\mathcal{E} = \lim_{n \rightarrow 1} \log Tr (\rho_A^{T_2})^n, \quad (2.9)$$

where we have used the replica limit $n \rightarrow 1$. The quantity $Tr (\rho_A^{T_2})^n$ can be expressed as follows

$$Tr (\rho_A^{T_2})^n = \langle \Phi_n(x_1, t_1) \Phi_{-n}(x_2, t_2) \Phi_{-n}(x_3, t_3) \Phi_n(x_4, t_4) \rangle. \quad (2.10)$$

3 Two-point twist correlator in GMMG

The holographic entanglement entropy for a single interval in the vacuum state of a $GCFT_2$ can be done by the replica technique. To this end, the geometry of the bulk should be considered. So far, this technique has been utilized in the Einstein gravity and in the flat space topological massive gravity (TMG) [24]. At first we review the results of [24] briefly, then we apply the replica approach in the generalized minimal massive gravity (GMMG) [2] in this section. In this context, the flat space geometry in Eddington-Finkelstein coordinates is given by the following metric

$$ds^2 = dr^2 - du^2 + r^2 d\phi^2. \quad (3.1)$$

An interval $A = [(u_1, \phi_1), (u_2, \phi_2)]$ on the boundary plane is located at the infinity of the flat spacetime. The bulk extremal curve joining the endpoints of the interval A is given by [29]

$$L_{tot}^{extr} = \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right|, \quad (3.2)$$

where

$$u_{12} = u_1 - u_2, \quad \phi_{12} = \phi_1 - \phi_2. \quad (3.3)$$

For the Einstein gravity in the flat space we have the following central charges for dual $GCFT_2$

$$c_L = 0, \quad c_M = \frac{3}{G}. \quad (3.4)$$

Substituting these central charges into (2.5), the field scaling dimensions can be found as follows

$$\Delta_n = 0, \quad \chi_n = \frac{1}{8G} \left(n - \frac{1}{n} \right). \quad (3.5)$$

As described in [29], the twist fields inserted at the endpoints of the interval can be considered as a particle with mass $m_n = \chi_n$ that propagates in the bulk. The two point correlator for the twisted fields can be considered as follows [29]

$$\langle \Phi_n(\partial_1 A) \Phi_{-n}(\partial_2 A) \rangle = \exp(-\chi_n S_{on-shell}), \quad (3.6)$$

where

$$S_{on-shell} = \sqrt{\eta_{\mu\nu} \dot{X}^\mu \dot{X}^\nu} = L_{tot}^{extr}, \quad (3.7)$$

where $S_{on-shell}$ is the on-shell action of the particle that propagates along an trajectory X^μ homologous to the interval. Substituting (3.7), (3.5) and (3.2) into (3.6), one can find the two-point correlator as follows

$$\langle \Phi_n(\partial_1 A) \Phi_{-n}(\partial_2 A) \rangle = \exp \left(- \frac{1}{8G} \left(n - \frac{1}{n} \right) \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right| \right), \quad (3.8)$$

Using the replica limit (2.6), and Eqs.(2.4), (3.8) the entanglement entropy can be found as follows

$$S_A = \frac{1}{4G} L_{tot}^{extr} = \frac{1}{4G} \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right|. \quad (3.9)$$

By considering the TMG in flat space [4, 5], a non-vanishing Δ introduces a spin for the massive particle. In this context, the bulk picture can be modified by introducing a gravitational Chern-Simons (CS) term [24, 29]. The action of the TMG has the following form

$$\begin{aligned} \mathcal{S}_{TMG} &= \mathcal{S}_{EH} + \frac{1}{\mu} \mathcal{S}_{CS} \\ &= \frac{1}{16\pi G} \int d^3x \left(R + \frac{2}{l^2} + \frac{1}{2\mu} \epsilon^{\alpha\beta\gamma} (\Gamma_{\alpha\sigma}^\rho \partial_\beta \Gamma_{\gamma\rho}^\sigma + \frac{2}{3} \Gamma_{\alpha\sigma}^\rho \Gamma_{\beta\eta}^\sigma \Gamma_{\gamma\rho}^\eta) \right), \end{aligned} \quad (3.10)$$

where the first term is the usual Einstein-Hilbert term, the second term is the cosmological constant term and the last one is a gravitational CS term. The algebra of the asymptotic Killing vectors of the TMG in AdS_3 is isomorphic to two copies of the Virasoro algebra with the following right and left central charges [31]:

$$c_{TMG}^+ = \frac{3l}{2G} \left(1 + \frac{1}{\mu l} \right), \quad c_{TMG}^- = \frac{3l}{2G} \left(1 - \frac{1}{\mu l} \right). \quad (3.11)$$

By taking Inonue-Wigner contraction [32, 33], the asymptotic symmetry group leads to the Galilean conformal algebra (GCA) with the following central charges:

$$\begin{aligned} c_L &= c_{TMG}^+ - c_{TMG}^- = \frac{3}{\mu G}, \\ c_M &= \frac{(c_{TMG}^+ + c_{TMG}^-)}{l} = \frac{3}{G}. \end{aligned} \quad (3.12)$$

In [24], the authors computed the entanglement entropy of the flat-TMG. As easily can be seen from (3.10), the CS-term is unaffected by the flat limit $l \rightarrow \infty$ and the propagating particle along the bulk trajectory is a massive spinning one with mass $m = \chi_n$ and spin $s = \Delta_n$. For such a particle propagating along the trajectory homologous to the interval, the two-point twist correlator has the following form:

$$\langle \Phi_n(\partial_1 A) \Phi_{-n}(\partial_2 A) \rangle = \exp(-\chi_n S_{on-shell}^{EH} - \Delta_n S_{on-shell}^{CS}), \quad (3.13)$$

where $S_{on-shell}^{CS}$ has the following form in a pure Minkowski spacetime [24]

$$S_{on-shell}^{CS} \equiv \Delta \eta_{12} = 2 \log \left(\frac{2}{\epsilon} \sin \frac{\phi_{12}}{2} \right). \quad (3.14)$$

Using the replica limit (2.6), (2.4),(2.5),(3.13), the entanglement entropy for a single interval in a pure Minkowski spacetime can be obtained as follows

$$\begin{aligned}
S_A &= -\lim_{n \rightarrow 1} \frac{\partial}{\partial n} \langle \Phi_n(\partial_1 A) \Phi_{-n}(\partial_2 A) \rangle \\
&= \lim_{n \rightarrow 1} [\partial_n \chi_n S_{on-shell}^{EH} + \partial_n \Delta_n S_{on-shell}^{CS}] \\
&= \frac{c_M}{12} S_{on-shell}^{EH} + \frac{c_L}{12} S_{on-shell}^{CS}.
\end{aligned} \tag{3.15}$$

Using (3.7) and substituting (3.2), (3.14) into (3.15), the following result can be found

$$S_A = \frac{c_M}{12} \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right| + \frac{c_L}{6} \log \left(\frac{2}{\epsilon} \sin \frac{\phi_{12}}{2} \right). \tag{3.16}$$

After a short review on the Einstein gravity and on the flat-TMG, we find the two-point twist correlator in the flat-GMMG. The Lagrangian of the GMMG model has the following form [2]

$$L_{GMMG} = L_{TMG} - \frac{1}{m^2} (f \cdot R + \frac{1}{2} e \cdot f \times f) + \frac{\alpha}{2} e \cdot h \times h \tag{3.17}$$

where L_{TMG} is the Lagrangian of the TMG and R is the Ricci scalar. f is an auxiliary one-form field, and m is a mass parameter term. e is a dreibein and h is the auxiliary field. The CS-term in the L_{TMG} has the following form in the first order formalism

$$CS - term : \frac{1}{2\mu} (\omega \cdot d\omega + \frac{1}{3} \omega \cdot \omega \times \omega) \tag{3.18}$$

where ω is dualised spin-connection. Our guess is that the only change in the CS-term in the two-point twist correlator (3.13) compared to the TMG is a change in the scaling dimension of the fields. The effects of other terms of the GMMG Lagrangian on the two-point twist correlator (3.13) can be addressed into the Einstein-Hilbert term. We can find the changes just in the scaling dimension χ_n compared to the flat-TMG. To provide a proof, we consider the two-point twist correlator in the flat-GMMG similar to the flat-TMG correlator (3.13). The only difference is in the scaling dimensions χ_n and Δ_n . The algebra of the asymptotic conserved charges of asymptotically AdS_3 spacetimes in the context of GMMG is isomorphic to two copies of the Virasoro algebra with the following right and left central charges [30]

$$c_{GMMG}^+ = \frac{3l}{2G} \left(-\sigma - \frac{\alpha H}{\mu} - \frac{F}{m^2} + \frac{1}{\mu l} \right), \quad c_{GMMG}^- = \frac{3l}{2G} \left(-\sigma - \frac{\alpha H}{\mu} - \frac{F}{m^2} - \frac{1}{\mu l} \right), \tag{3.19}$$

where H and F are constant. By taking Inonue-Wigner contraction, we can find the asymptotic symmetry group as a GCA with the following central charges

$$\begin{aligned}
c_L &= c_{GMMG}^+ - c_{GMMG}^- = \frac{3}{\mu G}, \\
c_M &= \frac{(c_{GMMG}^+ + c_{GMMG}^-)}{l} = \frac{3}{G} \left(-\sigma - \frac{\alpha H}{\mu} - \frac{F}{m^2} \right).
\end{aligned} \tag{3.20}$$

Substituting c_L and c_M (3.20) into (3.16), we find the entanglement entropy for the single interval A in the pure Minkowski spacetime as follows

$$S_A = \frac{1}{4G} \left(-\sigma - \frac{\alpha H}{\mu} - \frac{F}{m^2} \right) \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right| + \frac{1}{2\mu G} \log \left(\frac{2}{\epsilon} \sin \frac{\phi_{12}}{2} \right). \quad (3.21)$$

Previously, we have calculated the entanglement entropy for the GMMG in flat holography in another paper [25]. In the flat holography, we have considered the BMS_3 symmetry group [34, 35, 36] as an asymptotically symmetry group of GMMG in flat space. Since the BMS_3 field theories and the $GCFT_2$ theories have the same symmetry algebra, we expect the entanglement entropy to be the same in both cases. We arrive at this conclusion, and the entropy (3.21) obtained from (3.13) with the help of the replica approach is the same as the universal part of the result in [25]. The proof presented in this section shows us that we can use (3.13) with (3.20) in the all correlators that we need in all related computations to the entanglement entropy and the entanglement negativity.

4 Entanglement negativity and entanglement entropy for a single interval

In this section, we compute the entanglement negativity for various state configurations in a $GCFT_2$. These configurations consist of a single interval $A = [(x_1, t_1), (x_2, t_2)]$. To derive the entanglement negativity we need the four-point twist correlator (2.10) that is considered for a tripartite system. In the bipartite limit $B \rightarrow \emptyset$, the four-point correlator (2.10) can be turned into a two-point correlator as follows

$$Tr(\rho_A^{T_2})^n = \langle \Phi_n^2(x_1, t_1) \Phi_{-n}^2(x_2, t_2) \rangle. \quad (4.1)$$

For even n , there is following relation between the n -sheeted surface correlator and the $n/2$ -sheeted surface correlator [16]

$$\langle \Phi_n^2(x_1, t_1) \Phi_{-n}^2(x_2, t_2) \rangle = (\langle \Phi_{n/2}(x_1, t_1) \Phi_{-n/2}(x_2, t_2) \rangle)^2 \quad (4.2)$$

Substituting (3.13) into the left-hand side of (4.2), we find the two-point correlator (4.1) as follows

$$\langle \Phi_n^2(\partial_1 A) \Phi_{-n}^2(\partial_2 A) \rangle = \exp(-\chi_{n/2} S_{on-shell}^{EH} - \Delta_{n/2} \Delta \eta_{12}), \quad (4.3)$$

where

$$\begin{aligned} \chi_{n/2} &= \frac{c_M}{24} \left(\frac{n}{2} - \frac{2}{n} \right) = \frac{1}{8G} \left(-\sigma - \frac{\alpha H}{\mu} - \frac{F}{m^2} \right) \left(\frac{n}{2} - \frac{2}{n} \right) \\ \Delta_{n/2} &= \frac{c_L}{24} \left(\frac{n}{2} - \frac{2}{n} \right) = \frac{1}{8G\mu} \left(\frac{n}{2} - \frac{2}{n} \right), \end{aligned} \quad (4.4)$$

where we have used c_L and c_M central charges (3.20) into the right-hand sides of the equations in (4.4).

4.1 Single interval at zero temperature

We consider a holographic $GCFT_2$ at zero temperature at the boundary of the space. A single interval is considered in the vacuum state of the field theory. The geometry of the bulk in context of the GMMG is a pure Minkowski spacetime. We use (3.14) for the CS-contribution into the two-point correlator. The bulk extremal curve (3.9) can be utilized into (4.3). Using the replica limit (2.9) and $\chi_{n/2}$ and $\Delta_{n/2}$ (4.4), we find the entanglement negativity for a single interval at zero temperature as follows

$$\begin{aligned}\mathcal{E} &= \frac{c_L}{4} \log\left(\frac{2}{\epsilon} \sin \frac{\phi_{12}}{2}\right) + \frac{c_M}{8} \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right| \\ &= \frac{3}{4G\mu} \log\left(\frac{2}{\epsilon} \sin \frac{\phi_{12}}{2}\right) - \frac{3}{8G} \left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2}\right) \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right|\end{aligned}\quad (4.5)$$

where we have used (4.4) into the 2nd line of the entanglement negativity (4.5). Using the following transformations between the planar coordinates (x, t) and the cylinder coordinates (u, ϕ)

$$t = e^{i\phi}, \quad x = iue^{i\phi}, \quad (4.6)$$

we can find the following relations

$$\begin{aligned}t_{12} &= 2 \sin \frac{\phi_{12}}{2} \\ \frac{x_{12}}{t_{12}} &= \frac{1}{2} \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right|.\end{aligned}\quad (4.7)$$

Substituting relations (4.7) into (4.5), we find the entanglement negativity in the plane coordinates as follows

$$\begin{aligned}\mathcal{E} &= \frac{c_L}{4} \log\left(\frac{t_{12}}{\epsilon}\right) + \frac{c_M}{4} \frac{x_{12}}{t_{12}} \\ &= \frac{3}{4G\mu} \log\left(\frac{t_{12}}{\epsilon}\right) - \frac{3}{2G} \left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2}\right) \frac{x_{12}}{t_{12}}\end{aligned}\quad (4.8)$$

Given that the central charges of the GMMG are different from the charges of the TMG, the result for the flat-TMG in [24] can be extended to the GMMG in (4.8). The result (4.8) corresponds to the universal part of the result for the single interval configuration at zero temperature in [21].

To find the holographic entanglement entropy for a single interval at zero temperature, we can repeat the steps to find (3.15). The two-point twist correlator is the same as (3.13), just the scaling dimensions differ in GMMG in compare to the TMG. Substituting the central charges c_L and c_M (4.4) into the holographic entanglement entropy (3.15), we find

the following result for a zero-temperature single interval configuration

$$\begin{aligned}
S_A &= \frac{c_M}{12} S_{on-shell}^{EH} + \frac{c_L}{12} S_{on-shell}^{CS} \\
&= \frac{c_M}{12} \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right| + \frac{c_L}{6} \log \left(\frac{2}{\epsilon} \sin \frac{\phi_{12}}{2} \right) \\
&= -\frac{1}{4G} \left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2} \right) \left| \frac{u_{12}}{\tan \frac{\phi_{12}}{2}} \right| + \frac{1}{2\mu G} \log \left(\frac{2}{\epsilon} \sin \frac{\phi_{12}}{2} \right)
\end{aligned} \tag{4.9}$$

where we have used (3.2) and (3.14) in 2nd line. By comparing the holographic entanglement negativity (4.5) and the holographic entanglement entropy (4.9), we find the following relation between these results

$$\mathcal{E} = \frac{3}{2} S_A. \tag{4.10}$$

This result is consistent with the universal part of the result in [21].

4.2 Single interval at a finite-sized system

In this subsection we find the holographic entanglement negativity for the bipartite state of a single interval in a $GCFT_2$ with finite-sized system with periodic boundary conditions. The bulk geometry is a global Minkowski spacetime in the null infinity of the GMMG. Instead of the metric (3.1), we have the following one [37]

$$ds^2 = dr^2 - \left(\frac{2\pi}{l_\phi} \right)^2 du^2 + r^2 d\phi^2. \tag{4.11}$$

where the boundary periodic condition is as follows [37]

$$(u, \phi) \sim (u, \phi + l_\phi). \tag{4.12}$$

To find the entanglement negativity and the entanglement entropy, we need $\Delta\eta_{12}$ and L_{tot}^{extr} that we can use our previous result in [25]. For a field theory with following identification

$$(u, \phi) \sim (u + i\beta_u, \phi - i\beta_\phi) \tag{4.13}$$

we have the entanglement entropy for a single interval as follows

$$S_A = \frac{c_L}{6} \log \left(\frac{\beta_\phi}{\pi\epsilon_\phi} \sinh \frac{\pi l_\phi}{\beta_\phi} \right) + \frac{c_M}{6} \frac{1}{\beta_\phi} \left(\pi(l_u + \frac{\beta_u}{\beta_\phi} l_\phi) \coth \frac{\pi l_\phi}{\beta_\phi} - \beta_u \right) - \frac{c_M}{6} \frac{\epsilon_u}{\epsilon_\phi}. \tag{4.14}$$

By comparing (4.12) and (4.13), we find the following identifications to turn the entanglement entropy (4.14) for a finite-sized system,

$$\begin{aligned}
\beta_u &= 0, & \beta_\phi &= iL_\phi, \\
l_\phi &= \phi_{12} \\
l_u &= u_{12}
\end{aligned} \tag{4.15}$$

Substituting (4.15) into (4.14), we find the entanglement entropy for a single interval in a $GCFT_2$ with finite-sized system as follows

$$S_A = \frac{c_L}{6} \log \left(\frac{L_\phi}{\pi\epsilon} \sin \frac{\pi\phi_{12}}{L_\phi} \right) + \frac{c_M}{6} \frac{\pi u_{12}}{L_\phi} \cot \frac{\pi\phi_{12}}{L_\phi}. \quad (4.16)$$

By comparing (4.16) and (4.15), we find the following results for the extremal curve L_{tot}^{extr} and $\Delta\eta_{12}$ as follows

$$L_{tot}^{extr} = \frac{2\pi u_{12}}{L_\phi} \cot \frac{\pi\phi_{12}}{L_\phi} \quad (4.17)$$

and

$$\Delta\eta_{12} = 2 \log \left(\frac{L_\phi}{\pi\epsilon} \sin \frac{\pi\phi_{12}}{L_\phi} \right) \quad (4.18)$$

The bulk extremal curve (4.17) and the spinning contribution (4.18) can be utilized into the two-point correlator (4.3). Utilizing the replica technique (2.9), we find the entanglement negativity as follows

$$\begin{aligned} \mathcal{E} &= \frac{c_M}{8} L_{tot}^{extr} + \frac{c_L}{8} \Delta\eta_{12} \\ &= \frac{c_M}{4} \frac{\pi u_{12}}{L_\phi} \cot \frac{\pi\phi_{12}}{L_\phi} + \frac{c_L}{4} \log \left(\frac{L_\phi}{\pi\epsilon} \sin \frac{\pi\phi_{12}}{L_\phi} \right) \end{aligned} \quad (4.19)$$

This result is the same as the result in flat-TMG in [24], where the only difference is in the central charges between GMMG and TMG. We can utilize the central charges (3.20) into (4.19) that the following relation for the entanglement negativity can be found

$$\mathcal{E} = -\frac{3}{4G} \left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2} \right) \frac{\pi u_{12}}{L_\phi} \cot \frac{\pi\phi_{12}}{L_\phi} + \frac{3}{4G\mu} \log \left(\frac{L_\phi}{\pi\epsilon} \sin \frac{\pi\phi_{12}}{L_\phi} \right) \quad (4.20)$$

Substituting the central charges (3.20) into the entanglement entropy (4.16), we find the following result

$$S_A = -\frac{1}{2G} \left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2} \right) \frac{\pi u_{12}}{L_\phi} \cot \frac{\pi\phi_{12}}{L_\phi} + \frac{1}{2G\mu} \log \left(\frac{L_\phi}{\pi\epsilon} \sin \frac{\pi\phi_{12}}{L_\phi} \right). \quad (4.21)$$

By comparing (4.20) and (4.21), the following relation between the entanglement entropy and the entanglement negativity for a single interval in a finite-sized system can be found

$$\mathcal{E} = \frac{3}{2} S_A. \quad (4.22)$$

This result is the same as the universal part of the result of [21] that has been obtained by another approach there.

4.3 Single interval at a finite temperature

For a single interval A at a finite temperature, the $GCFT_2$ can be considered as a thermal field theory on a cylinder with complex circumference β . A naive two-point twist correlator as (4.3) to compute the entanglement negativity leads to an incorrect result [21]. For this reason, two other finite intervals B_1 and B_2 adjacent to the interval A are necessary to be considered. The bulk spacetime is described by GMMG in FSC geometries. To find the entanglement negativity, we compute the following four-point twist correlator instead of the two-point correlator (4.3)

$$\begin{aligned}
\langle \Phi_n(x_1, t_1) \Phi_{-n}^2(x_2, t_2) \Phi_n^2(x_3, t_3) \Phi_{-n}(x_4, t_4) \rangle &= \frac{t_{13}^{\Delta_n + \Delta_n^{(2)}} t_{24}^{\Delta_n + \Delta_n^{(2)}}}{t_{12}^{\Delta_n + \Delta_n^{(2)}} t_{23}^{\Delta_n + \Delta_n^{(2)}} t_{34}^{\Delta_n + \Delta_n^{(2)}} t_{14}^{2\Delta_n}} \\
&\times \exp \left[\frac{x_{13}}{t_{13}} (\chi_n + \chi_n^{(2)}) + \frac{x_{24}}{t_{24}} (\chi_n + \chi_n^{(2)}) \right. \\
&- \frac{x_{12}}{t_{12}} (\chi_n + \chi_n^{(2)}) - \frac{x_{23}}{t_{23}} (2\chi_n) \\
&\left. - \frac{x_{34}}{t_{34}} (\chi_n + \chi_n^{(2)}) - \frac{x_{14}}{t_{14}} (2\chi_n) \right] \mathcal{G}(t, \frac{x}{t})
\end{aligned} \tag{4.23}$$

where we have just write the universal part of the four-point correlator. To write the right-hand side of (4.23), we have used the general form of the four-point twist field correlator in [21]. Substituting the following identities

$$\left(\frac{t_{13} t_{24}}{t_{12} t_{34}} \right)^{\Delta_n} \equiv t^{-\Delta_n} \tag{4.24}$$

and

$$\chi_n \left(\frac{x_{12}}{t_{12}} + \frac{x_{34}}{t_{34}} - \frac{x_{13}}{t_{13}} - \frac{x_{24}}{t_{24}} \right) \equiv \chi_n \frac{x}{t} \tag{4.25}$$

and inserting these identities into the $\mathcal{G}(t, \frac{x}{t})$, we can find the four-point correlator as follows

$$\begin{aligned}
\langle \Phi_n(x_1, t_1) \Phi_{-n}^2(x_2, t_2) \Phi_n^2(x_3, t_3) \Phi_{-n}(x_4, t_4) \rangle &= \exp \left[-\chi_n L_{14}^{extr} - \chi_{n/2} (2L_{23}^{extr} + L_{12}^{extr} + L_{34}^{extr}) \right. \\
&- L_{13}^{extr} - L_{24}^{extr} - \Delta_n \Delta \eta_{14} - \Delta_{n/2} (2\Delta \eta_{23} \\
&\left. + \Delta \eta_{12} + \Delta \eta_{34} - \Delta \eta_{13} - \Delta \eta_{14}) \right]
\end{aligned} \tag{4.26}$$

where we have used the following relations

$$L_{ij}^{extr} = 2 \frac{x_{ij}}{t_{ij}} \tag{4.27}$$

and

$$\Delta \eta_{ij} = 2 \log \frac{t_{ij}}{\epsilon}. \tag{4.28}$$

Substituting (4.26) into the replica technique (2.9), we can find the entanglement negativity as follows

$$\mathcal{E} = \frac{c_M}{16}(2L_A + L_{B_1} + L_{B_2} - L_{AUB_1} - L_{AUB_2}) \quad (4.29)$$

$$+ \frac{c_L}{16}(2\Delta\eta_A + \Delta\eta_{B_1} + \Delta\eta_{B_2} - \Delta\eta_{AUB_1} - \Delta\eta_{AUB_2}) \quad (4.30)$$

where we have used the following definitions [24]

$$L_{12}^{extr} = L_{B_1}, \quad L_{23}^{extr} = L_A, \quad L_{34}^{extr} = L_{B_2}, \quad (4.31)$$

$$L_{13}^{extr} = L_{AUB_1}, \quad L_{24}^{extr} = L_{AUB_2}, \quad L_{14}^{extr} = L_{AUB}. \quad (4.32)$$

Substituting the central charges (3.20) into (4.29), we find the following result for the entanglement negativity

$$\mathcal{E} = \frac{3}{16G}(2\mathcal{L}_A + \mathcal{L}_{B_1} + \mathcal{L}_{B_2} - \mathcal{L}_{AUB_1} - \mathcal{L}_{AUB_2}) \quad (4.33)$$

where

$$\mathcal{L}_X = -(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2})L_X + \frac{1}{\mu}\Delta\eta_X. \quad (4.34)$$

To find the exact form of the universal part of the entanglement negativity, one can write the right-hand side of (4.23) as follows

$$\begin{aligned} \langle \Phi_n(x_1, t_1) \Phi_{-n}^2(x_2, t_2) \Phi_n^2(x_3, t_3) \Phi_{-n}(x_4, t_4) \rangle &= t^{-\Delta_n} t^{-\Delta_n^{(2)}} t_{14}^{-2\Delta_n} t_{23}^{-2\Delta^{(2)}} \\ &\times \exp \left[-\chi_n^{(2)} \frac{x}{t} - \chi_n \frac{x}{t} \right. \\ &\left. - 2\chi_n^{(2)} \frac{x_{23}}{t_{23}} - 2\chi_n \frac{x_{14}}{t_{14}} \right]. \end{aligned} \quad (4.35)$$

In the bipartite limit ($L \rightarrow \infty$), we have the following relations for x and $\frac{x}{t}$ as follows [21]

$$\lim_{L \rightarrow \infty} t = \exp\left(-\frac{2\pi}{\beta}\phi_{23}\right), \quad \lim_{L \rightarrow \infty} \frac{x}{t} = -\frac{2\pi}{\beta}u_{23}. \quad (4.36)$$

Utilizing the entanglement entropy (4.14) that we have computed in our previous work [25] for a finite temperature field theory, we can find the following result

$$t_{ij} = \frac{\beta}{\pi} \sinh\left(\frac{\pi}{\beta}\phi_{ij}\right) \quad (4.37)$$

$$\frac{x_{ij}}{t_{ij}} = \frac{\pi}{\beta} u_{ij} \coth\left(\frac{\pi}{\beta}\phi_{ij}\right) \quad (4.38)$$

where we have used the following identifications into (4.14)

$$\begin{aligned} \beta_u &= 0, \quad \beta_\phi = \beta, \\ l_\phi &= \phi_{12}, \\ l_u &= u_{12}. \end{aligned} \quad (4.39)$$

Substituting (4.37) and (4.36) into (4.35) and using the replica limit (2.9), we find the entanglement negativity for a finite temperature $GCF T_2$ as follows

$$\begin{aligned}\mathcal{E} &= \frac{c_L}{4} \log\left(\frac{\beta}{\pi\epsilon} \sinh\left(\frac{\pi}{\beta}\phi_{23}\right)\right) - \frac{c_L}{4} \frac{\pi}{\beta} \phi_{23} \\ &+ \frac{c_M}{4} \frac{\pi}{\beta} u_{23} \coth\left(\frac{\pi}{\beta}\phi_{23}\right) - \frac{c_M}{4} \frac{\pi}{\beta} u_{23}\end{aligned}\quad (4.40)$$

This is same as the result in [24] where the only difference is in the central charges (3.20). The result (4.40) is compatible with the universal part of the result in [21]. To compute the entanglement entropy using the replica technique (3.15), we use (4.37) and (4.38) to find L_{FSC}^{extr} and $\Delta\eta_{ij}$

$$L_{FSC}^{extr} = 2\frac{\pi}{\beta} u_{23} \coth\left(\frac{\pi}{\beta}\phi_{23}\right) \quad (4.41)$$

and

$$S_{FSC}^{CS} = 2 \log\left(\frac{\beta}{\pi\epsilon} \sinh\left(\frac{\pi}{\beta}\phi_{23}\right)\right). \quad (4.42)$$

Substituting (4.41) and (4.42) into two-point correlator (3.13) and utilizing the replica technique (3.15), we find the holographic entanglement entropy as follows

$$\begin{aligned}S_A &= \frac{c_M}{6} \frac{\pi}{\beta} u_{23} \coth\left(\frac{\pi}{\beta}\phi_{23}\right) + \frac{c_L}{6} \log\left(\frac{\beta}{\pi\epsilon} \sinh\left(\frac{\pi}{\beta}\phi_{23}\right)\right) \\ &= -\frac{1}{2G}\left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2}\right) \frac{\pi}{\beta} u_{23} \coth\left(\frac{\pi}{\beta}\phi_{23}\right) \\ &+ \frac{1}{2G\mu} \log\left(\frac{\beta}{\pi\epsilon} \sinh\left(\frac{\pi}{\beta}\phi_{23}\right)\right).\end{aligned}\quad (4.43)$$

By comparing (4.40) and (4.43), one can find the following result between the holographic entanglement entropy and the holographic entanglement negativity

$$\mathcal{E} = \frac{3}{2}(S_A - S_{th}) \quad (4.44)$$

where S_{th} is the thermal entropy and has the following form

$$S_{th} = \frac{c_L}{4} \frac{\pi}{\beta} \phi_{23} + \frac{c_M}{4} \frac{\pi}{\beta} u_{23}. \quad (4.45)$$

where can be calculated by the Cardy formula [38, 39]. The Cardy formula for the thermal entropy can be considered as follows

$$S_{th} = -\frac{\pi^2}{3}\left(c_L \frac{b}{a} + c_M \frac{\bar{a}b - a\bar{b}}{a^2}\right), \quad (4.46)$$

where there is the following identification between the thermal circle and the spatial circle

$$(\tilde{u}, \tilde{\phi}) \sim (\tilde{u} + i\bar{a}, \tilde{\phi} - ia) \sim (\tilde{u} + 2\pi\bar{b}, \tilde{\phi} - 2\pi b), \quad (4.47)$$

where (a, \bar{a}) parameterizes the thermal circle and (b, \bar{b}) parameterizes the spatial circle. The formula (4.46) is the general form of the thermal entropy. We can use this formula in the configuration of a single interval at finite temperature $GCFT_2$, utilizing the following identification

$$(u, \phi) \sim (u, \phi - i\beta) \sim (u + u_{23}, \phi + \phi_{23}). \quad (4.48)$$

By comparing (4.47) and (4.48), we find the special parameters of the circles. Substituting the parameters into (4.46), the result (4.45) for the thermal entropy can be established. The result (4.44) in the FSC background of the GMMG and for the finite temperature $GCFT_2$ is the same as the result in [21] in the $GCFT_2$ with a bipartite system of a single interval at finite temperature.

5 Entanglement negativity and entanglement entropy for adjacent intervals

In this section, we consider the $GCFT_2$ on the boundary with two adjacent intervals A_1 and A_2 . To compute the entanglement negativity, the GMMG can be considered as the bulk gravity theory with asymptotically flat space geometry. As we have two adjacent intervals, the four-point twist correlator (2.10) can be used to find the entanglement negativity. Since these intervals A_1 and A_2 are not far apart, the correlator (2.10) can be turned into the following form

$$Tr(\rho_A^{T_2})^n = \langle \Phi_n(x_1, t_1) \Phi_{-n}^2(x_2, t_2) \Phi_n(x_3, t_3) \rangle, \quad (5.1)$$

where we have utilized the following limit into (2.10)

$$(x_2, t_2) \rightarrow (x_3, t_3). \quad (5.2)$$

By applying the limit (5.2) into the four-point correlator, we can find the three-point correlator (5.1) as follows

$$\begin{aligned} \langle \Phi_n(x_1, t_1) \Phi_{-n}^2(x_2, t_2) \Phi_n(x_3, t_3) \rangle &= \exp \left[-\chi_n L_{13}^{extr} - \chi_{n/2} (L_{12}^{extr} + L_{23}^{extr} - L_{13}^{extr}) \right. \\ &\quad \left. - \Delta_n \Delta \eta_{13} - \Delta_{n/2} (\Delta \eta_{12} + \Delta \eta_{23} - \Delta \eta_{13}) \right] \end{aligned} \quad (5.3)$$

Utilizing the central charges c_L and c_M (3.20) into (5.3) and by substituting (5.3) into the replica limit (2.9), the holographic entanglement negativity can be found as follows

$$\mathcal{E} = \frac{3}{16G} \left[-\left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2}\right) (L_{12}^{extr} + L_{23}^{extr} - L_{13}^{extr}) + \frac{1}{\mu} (\Delta \eta_{12} + \Delta \eta_{23} - \Delta \eta_{13}) \right] \quad (5.4)$$

This equation can be expressed as follows

$$\mathcal{E} = \frac{3}{16G} \left[\mathcal{L}_{A_1} + \mathcal{L}_{A_2} - \mathcal{L}_{A_1 \cup A_2} \right] \quad (5.5)$$

where we have defined \mathcal{L}_X as follows

$$\mathcal{L}_X = -\left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2}\right)L_X + \frac{1}{\mu}\Delta\eta_X. \quad (5.6)$$

This result is the same as the entanglement negativity of two adjacent intervals in [24], where the only difference is the scaling dimensions of the twist fields.

5.1 Adjacent intervals at zero temperature

To consider the $GCFT_2$ at zero temperature, the two adjacent intervals should be in the vacuum state of the theory on the boundary. We have computed the extremal curve L_{tot}^{extr} and the spinning contribution $\Delta\eta_{ij}$ in the FSC bulk geometry in (4.27) and (4.28), respectively. Substituting L_{tot}^{extr} and $\Delta\eta_{ij}$ into (5.4), we find the holographic entanglement negativity as follows

$$\mathcal{E} = -\frac{3}{8G}\left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2}\right)\left(\frac{x_{12}}{t_{12}} + \frac{x_{23}}{t_{23}} - \frac{x_{13}}{t_{13}}\right) + \frac{3}{8G}\log\frac{t_{12}t_{23}}{\epsilon t_{13}}. \quad (5.7)$$

We can write (5.7) with the central charges (3.20) as follows

$$\mathcal{E} = \frac{c_M}{8}\left(\frac{x_{12}}{t_{12}} + \frac{x_{23}}{t_{23}} - \frac{x_{13}}{t_{13}}\right) + \frac{c_L}{8}\log\frac{t_{12}t_{23}}{\epsilon(t_{12} + t_{23})}. \quad (5.8)$$

The holographic entanglement negativity (5.8) is the same as the result for flat-TMG in [24], but with different central charges. To compute the holographic entanglement entropy, we utilize the general form of the entropy in (3.15) as follows

$$S_A = \frac{c_M}{12}L_{ij}^{extr} + \frac{c_L}{12}\Delta\eta_{ij}, \quad (5.9)$$

Utilizing (4.27) and (4.28) into (5.9), we find the entanglement entropy of a single interval $[(x_i, t_i), (x_j, t_j)]$ in FSC bulk geometry as follows

$$S_{ij} = \frac{c_M}{6}\frac{x_{ij}}{t_{ij}} + \frac{c_L}{6}\log\frac{t_{ij}}{\epsilon}. \quad (5.10)$$

By comparing the entanglement negativity (5.8) and the entanglement entropy (5.10), we can easily find the following relation between them

$$\mathcal{E} = \frac{3}{4}(S_{12} + S_{23} - S_{13}), \quad (5.11)$$

where S_{12} , S_{23} and S_{13} can be found by (5.10). The relation between the entanglement entropy and the entanglement negativity in (5.11) is same as the result between the two types of the entropies in [21].

5.2 Adjacent intervals at a finite-sized system

We find the holographic entanglement negativity for a configuration of two adjacent intervals in a $GCFT_2$ with a finite size. The $GCFT_2$ is the boundary side theory of the duality that can be considered on an infinite cylinder with L_ϕ circumference [24]. The gravity model in the bulk is the GMMG model in FSC geometry. We utilize (4.17) and (4.18) for extremal curve L_{ij}^{extr} and the spinning contribution $\Delta\eta_{ij}$, respectively. Comparing (4.27) and (4.28) with (4.17) and (4.18), we find the following result

$$\frac{x_{ij}}{t_{ij}} = \frac{\pi u_{ij}}{L_\phi} \cot \frac{\pi\phi_{ij}}{L_\phi} \quad (5.12)$$

and

$$t_{ij} = \frac{L_\phi}{\pi} \sin \frac{\pi\phi_{ij}}{L_\phi}. \quad (5.13)$$

Substituting (5.12) and (5.13) into the entanglement negativity relation (5.8)

$$\begin{aligned} \mathcal{E} &= \frac{c_L}{8} \log \left(\frac{L_\phi \sin \frac{\pi\phi_{12}}{L_\phi} \sin \frac{\pi\phi_{23}}{L_\phi}}{\pi\epsilon \sin \frac{\pi\phi_{13}}{L_\phi}} \right) \\ &+ \frac{c_M}{8} \left(\frac{\pi u_{12}}{L_\phi} \cot \frac{\pi\phi_{12}}{L_\phi} + \frac{\pi u_{23}}{L_\phi} \cot \frac{\pi\phi_{23}}{L_\phi} - \frac{\pi u_{13}}{L_\phi} \cot \frac{\pi\phi_{13}}{L_\phi} \right) \end{aligned} \quad (5.14)$$

This is same as the result in [24]. The result (5.14) for asymptotically flat GMMG is different with the TMG result in flat TMG. The only difference is due to the different central charges for these massive gravity models. Substituting (5.13) and (5.12) into (5.10), we can reproduce (5.11) for the finite-sized system. This result is consistent with the result in [21].

5.3 Adjacent intervals at a finite temperature

In this subsection, we focus on the configuration of two adjacent intervals in a finite temperature $GCFT_2$. As the previous section, we have the theory on an infinite cylinder but with complex circumference β . The bulk geometry at null infinity is FSC geometry that is a solution of GMMG model. To compute the following holographic entanglement negativity, we utilize relations (4.37) and (4.38) into (5.8)

$$\begin{aligned} \mathcal{E} &= \frac{c_L}{8} \log \left(\frac{\beta \sinh \frac{\pi\phi_{12}}{\beta} \sinh \frac{\pi\phi_{23}}{\beta}}{\pi\epsilon \sinh \frac{\pi\phi_{13}}{\beta}} \right) \\ &+ \frac{c_M}{8} \left(\frac{\pi u_{12}}{\beta} \coth \frac{\pi\phi_{12}}{\beta} + \frac{\pi u_{23}}{\beta} \coth \frac{\pi\phi_{23}}{\beta} - \frac{\pi u_{13}}{\beta} \coth \frac{\pi\phi_{13}}{\beta} \right). \end{aligned} \quad (5.15)$$

Except for the difference in the central charges, there is no difference between the result (5.15) in the GMMG and the entanglement negativity of two adjacent intervals in a

finite temperature case in [24]. Substituting relations (4.41) and (4.42) into the holographic entanglement entropy (5.10) for a single interval, we find the following result

$$S_{ij} = \frac{c_M \pi}{6 \beta} u_{ij} \coth\left(\frac{\pi}{\beta} \phi_{ij}\right) + \frac{c_L}{6} \log\left(\frac{\beta}{\pi \epsilon} \sinh\left(\frac{\pi}{\beta} \phi_{ij}\right)\right). \quad (5.16)$$

By comparing (5.15) and (5.16), the result (5.11) can be reproduced, that is the same as the result in [21]. We have had this result in all three cases of the two adjacent intervals configurations.

6 Entanglement negativity and entanglement entropy for disjoint intervals

In this section, we compute the holographic entanglement negativity and the holographic entanglement entropy for the state of two disjoint intervals in a $GCFT_2$. In the bulk side of the duality, we have the GMMG in pure Minkowski spacetime. As we have two disjoint intervals, the four-point twist correlator (2.10) can be used to find the entanglement negativity. The four point twist correlator for two disjoint intervals $A_1 = [(x_1, t_1), (x_2, t_2)]$ and $A_2 = [(x_3, t_3), (x_4, t_4)]$ can be considered as follows [24]

$$\begin{aligned} \langle \Phi_n(x_1, t_1) \Phi_{-n}(x_2, t_2) \Phi_n(x_3, t_3) \Phi_{-n}(x_4, t_4) \rangle &= \exp \left[\frac{c_M}{16} (L_{13}^{extr} + L_{24}^{extr} - L_{14}^{extr} - L_{23}^{extr}) \right. \\ &\quad \left. + \frac{c_L}{16} (\Delta\eta_{13} + \Delta\eta_{24} - \Delta\eta_{14} - \Delta\eta_{23}) \right], \quad (6.1) \end{aligned}$$

Substituting (6.1) into the replica limit (2.9), we find the entanglement negativity as follows

$$\mathcal{E} = \frac{3}{16G} (\mathcal{L}_{13}^{extr} + \mathcal{L}_{24}^{extr} - \mathcal{L}_{14}^{extr} - \mathcal{L}_{23}^{extr}), \quad (6.2)$$

where

$$\mathcal{L}_X^{extr} = -\left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2}\right) L_X^{extr} + \frac{1}{\mu} \Delta\eta_X. \quad (6.3)$$

We can use this result to compute the entanglement negativity in different configurations involving two disjoint intervals.

6.1 Disjoint intervals at zero temperature

To consider the two disjoint intervals configurations, the $GCFT_2$ should be in the ground state that is dual to the GMMG in pure Minkowski spacetime. By comparing (4.5) and (4.7), we use the extremal curve and the spinning contribution in the pure Minkowski spacetime as follows

$$L_{ij}^{extr} = 2 \frac{x_{ij}}{t_{ij}}, \quad \Delta\eta_{ij} = 2 \log\left(\frac{t_{ij}}{\epsilon}\right). \quad (6.4)$$

Substituting L_{ij}^{extr} and $\Delta\eta_{ij}$ (6.4) into the entanglement negativity (6.2), we find the entanglement negativity for two disjoint intervals at zero temperature as follows

$$\mathcal{E} = -\frac{3}{8G}\left(\sigma + \frac{\alpha H}{\mu} + \frac{F}{m^2}\right)\left(\frac{x_{13}}{t_{13}} + \frac{x_{24}}{t_{24}} - \frac{x_{14}}{t_{14}} - \frac{x_{23}}{t_{23}}\right) + \frac{3}{8G\mu} \log\left(\frac{t_{13}t_{24}}{t_{14}t_{23}}\right) \quad (6.5)$$

where we can write this result as following

$$\mathcal{E} = \frac{c_M}{8}\left(\frac{x_{13}}{t_{13}} + \frac{x_{24}}{t_{24}} - \frac{x_{14}}{t_{14}} - \frac{x_{23}}{t_{23}}\right) + \frac{c_L}{8} \log\left(\frac{t_{13}t_{24}}{t_{14}t_{23}}\right). \quad (6.6)$$

This result is same as the result for two disjoint intervals configurations at zero temperature holographic $GCFT_2$ in [24].

6.2 Disjoint intervals at a finite-sized system

In this subsection, the two disjoint intervals are considered in a finite-sized system. The boundary theory is on a cylinder with circumference L_ϕ . The gravity in the bulk side of the theory is described by FSC geometry in the GMMG. Substituting (5.12) and (5.13) into (4.27) and (4.28) respectively, we find the extremal curve and the spinning contribution as follows

$$\begin{aligned} L_{ij}^{extr} &= \frac{2\pi}{L_\phi} u_{ij} \cot\left(\frac{\pi}{l_\phi} \phi_{ij}\right), \\ \Delta\eta_{ij} &= 2 \log\left(\frac{L_\phi}{\pi} \sin\frac{\pi}{l_\phi} \phi_{ij}\right). \end{aligned} \quad (6.7)$$

Substituting (6.7) into (6.2), we find the entanglement negativity as follows

$$\begin{aligned} \mathcal{E} &= \frac{c_L}{8} \log\left(\frac{L_\phi \sin\frac{\pi\phi_{13}}{L_\phi} \sin\frac{\pi\phi_{24}}{L_\phi}}{\pi \sin\frac{\pi\phi_{14}}{L_\phi} \sin\frac{\pi\phi_{23}}{L_\phi}}\right) \\ &+ \frac{c_M}{8} \left(\frac{\pi u_{13}}{L_\phi} \cot\frac{\pi\phi_{13}}{L_\phi} + \frac{\pi u_{24}}{L_\phi} \cot\frac{\pi\phi_{24}}{L_\phi} - \frac{\pi u_{14}}{L_\phi} \cot\frac{\pi\phi_{14}}{L_\phi} - \frac{\pi u_{23}}{L_\phi} \cot\frac{\pi\phi_{23}}{L_\phi}\right). \end{aligned} \quad (6.8)$$

6.3 Disjoint intervals at a finite temperature

In this subsection, we have a configuration state of two disjoint intervals in a finite temperature $GCFT_2$. The boundary is living on a cylinder with complex circumference β . The bulk side of the duality is described by the FSC geometry in the GMMG. To find the holographic entanglement entropy, we utilize the extremal curve (4.41) and the spinning contributions

(4.42) into (6.2) as follows

$$\begin{aligned}
\mathcal{E} &= \frac{c_L}{8} \log \left(\frac{\beta \sinh \frac{\pi\phi_{13}}{\beta} \sinh \frac{\pi\phi_{24}}{\beta}}{\pi \sinh \frac{\pi\phi_{14}}{\beta} \sinh \frac{\pi\phi_{23}}{\beta}} \right) \\
&+ \frac{c_M}{8} \left(\frac{\pi u_{13}}{\beta} \coth \frac{\pi\phi_{13}}{\beta} + \frac{\pi u_{24}}{\beta} \coth \frac{\pi\phi_{24}}{\beta} - \frac{\pi u_{14}}{\beta} \coth \frac{\pi\phi_{14}}{\beta} \right. \\
&\left. - \frac{\pi u_{23}}{\beta} \coth \frac{\pi\phi_{23}}{\beta} \right). \tag{6.9}
\end{aligned}$$

7 Conclusion

In this paper we give an evidence for non-relativistic holography between the gravity theory and the quantum field theory in $2 + 1$ dimension. The asymptotic symmetry group of the spacetime is considered to be the Galilean conformal (GC) group that is a non-relativistic version of the conformal group. The theory in two-dimensional boundary is a Galilean conformal field theory ($GCFT_2$) [22, 23] and the bulk description would be the generalized minimal massive gravity (GMMG)[2]. GMMG model is an extension of MMG model [3] by adding the CS-term and an extra term to pure Einstein gravity with a negative cosmological constant. The asymptotic symmetry of the asymptotically flat spacetime solutions of GMMG model Galilean conformal algebra (GCA) in two dimensions [30]. In this context we obtain the holographic entanglement negativity for bipartite states in $GCFT_2$ dual to bulk asymptotically flat space times.

The holography provides a good construction to find out a viable measurement for the quantum entanglement [7] of the bipartite state of the $GCFT_2$. The entanglement entropy is not a proper measure about the entanglement in a mixed states [9], but the entanglement negativity can be used as a measure of the entanglement in the bipartite state configuration of the $GCFT_2$ s [9]. We use the replica technique to compute the entanglement negativity and the entanglement entropy. To this end the manifold can be considered as an n -sheeted surface. The relation between the reduced density matrix corresponding to the interval A and the entanglement of the interval introduced in (2.6). By defining the partial transpose of the reduced density matrix in (2.8), the replica limit (2.9) can be used to compute the entanglement entropy.

The two-point twist correlator (3.13) has an important role in computation of the entanglement entropy. We give an argument that the form (3.13) of the correlator can be used in the GMMG model. The argument works because we can reproduce our previous result of the holographic entanglement entropy in [25]. The correlator (3.13) has an S_{EH} contribution which originates from the pure Einstein gravity and its extra terms added to it in the GMMG. Another contribution in (3.13) is a spinning one that is related to the CS-term in the GMMG. The importance of the two-point twist correlator (3.13) is not only in the holographic entanglement entropy, but also plays an important role in the computations related to the entanglement negativity.

We computed the bipartite state configurations in the cases a single interval, two-adjacent interval and two-disjoint interval. In each case, we considered the field theory on the boundary side of the duality in ground state, finite-sized system and finite-temperature states. One can find the results of the single interval in three cases in (4.8), (4.20) and (4.40). The result of the entanglement negativity related to two-adjacent intervals can be found in (5.4), (5.14) and (5.15). The result related to the different cases of the two-disjoint intervals can be found in (6.2), (6.8) and (6.9). As we expected, the only difference between the GMMG results and the TMG result is in their central charges that these charges are the coefficients of the Einstein-Hilbert contribution and the spinning contribution in the entanglement negativity.

Another important conclusion that we have obtained is the relationship between the holographic entanglement negativity and the entanglement entropy in each cases of the bipartite states. The general form of these relationships in the $GCFT_2$ have been introduced in [21]. We extend the results to the GMMG model. In the single interval cases, the entanglement negativity is three-thirds of the entanglement entropy in zero-temperature and in the finite-sized state. The only difference is in the finite temperature state that the contribution of the thermal entropy is also included that can be seen in (4.44). The relationship between two types of the entanglement measures in the two-adjacent cases has a same form in each cases of the $GCFT_2$ situations that can be bound in (5.11). According to the results of the present paper we can say with more sure that the dual theory of the asymptotically flat spacetime solutions of GMMG is a $GCFT_2$.

References

- [1] A. Bagchi, R. Gopakumar, I. Mandal, and A. Miwa, *GCA in 2d*, *JHEP* **08** (2010) 004.
- [2] M. R. Setare, *On the generalized minimal massive gravity*, *Nucl. Phys. B* **898** (2015) 259–275.
- [3] E. Bergshoeff, O. Hohm, W. Merbis, A. J. Routh and P. K. Townsend, *Class. Quant. Grav.* **31**, 145008 (2014).
- [4] S. Deser, R. Jackiw and S. Templeton, *Three-Dimensional Massive Gauge Theories*, *Phys. Rev. Lett.* **48** (1982) 975–978.
- [5] S. Deser, R. Jackiw and S. Templeton, *Topologically Massive Gauge Theories*, *Annals Phys.* **140** (1982) 372–411.
- [6] E. A. Bergshoeff, O. Hohm and P. K. Townsend, *Phys. Rev. Lett.* **102**, 201301 (2009)
- [7] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, *Rev. Mod. Phys.* **80**, 517 (2008); J. Eisert, M. Cramer, and M. B. Plenio, *ibid.* **82**, 277 (2010); *Entanglement entropy in extended systems*, P. Calabrese, J. Cardy, and B. Doyon, *J. Phys. A* **42**, 500301 (2009).

- [8] C. H. Bennett, H. J. Bernstein, S. Popescu, and B. Schumacher, *Concentrating Partial Entanglement by Local Operations*, *Phys. Rev. A* **53** (1996) 2046.
- [9] P. Calabrese, J. Cardy, and E. Tonni, *Entanglement negativity in quantum field theory*, *Phys. Rev. Lett.* **109** (2012) 130502.
- [10] M. B. Plenio and S. Virmani, *An introduction to entanglement measures*, *Quant. Inf. Comput.* **7**, 1 (2007).
- [11] J. Eisert and M. B. Plenio, *A Comparison of entanglement measures*, *J. Mod. Opt.* **46**, 145 (1999).
- [12] A. Peres, *Separability criterion for density matrices*, *Phys. Rev. Lett.* **77**, 1413 (1996); M. Horodecki, P. Horodecki and R. Horodecki, *Mixed state entanglement and distillation: Is there a ‘bound’ entanglement in nature?*, *Phys. Rev. Lett.* **80**, 5239 (1998);
- [13] K. Zyczkowski, P. Horodecki, A. Sanpera and M. Lewenstein, *On the volume of the set of mixed entangled states*, *Phys. Rev. A* **58**, 883 (1998); L. M. Duan, G. Giedke, J. I. Cirac and P. Zoeller, *Inseparability criterion for continuous variables systems*, *Phys. Rev. Lett.* **84**, 2722 (2000); R. Simon, *Peres-Horodecki separability criterion for continuous variables systems*, *Phys. Rev. Lett.* **84**, 2726 (2000).
- [14] G. Vidal and R. F. Werner, *Computable measure of entanglement*, *Phys. Rev. A* **65** (2002) 032314.
- [15] G. Vidal, *Entanglement monotones*, *J. Mod. Opt.* **47** (2000) 355
- [16] P. Calabrese, J. Cardy, and E. Tonni, *Entanglement negativity in extended systems: A field theoretical approach*, *J. Stat. Mech.* **1302** (2013) P02008.
- [17] V. Alba, *Entanglement negativity and conformal field theory: a Monte Carlo study*, *J. Stat. Mech.* P05013 (2013).
- [18] P. Calabrese, L. Tagliacozzo and E. Tonni, *Entanglement negativity in the critical Ising chain*, *J. Stat. Mech.* P05002 (2013).
- [19] M. Rangamani and M. Rota, *Comments on Entanglement Negativity in Holographic Field Theories*, arXiv:1406.6989.
- [20] M. Kulaxizi, A. Parnachev, and G. Policastro, *Conformal Blocks and Negativity at Large Central Charge*, arXiv:1407.0324
- [21] V. Malvimat, H. Himanshu, B. Paul and G. Sengupta, *Entanglement Negativity in Galilean Conformal Field Theories*, *Phys.Rev.D* **100** (2019) 2, 026001.
- [22] A. Bagchi, R. Basu, D. Grumiller and M. Riegler, *Entanglement entropy in Galilean conformal field theories and flat holography*, *Phys. Rev. Lett.* **114** (2015) 111602.

- [23] R. Basu and M. Riegler, *Wilson lines and holographic entanglement entropy in Galilean conformal field theories*, *Physical Review D* **93** (2016).
- [24] D. Basu, A. Chaandra, A. Parihar and G. Sengupta, *Entanglement Negativity in Flat Holography*, *arXiv: 2102.05685* .
- [25] M. R. Setare and M. Koohgard, *Holographic Entanglement Entropy in flat limit of the Generalized Minimal Massive Gravity model*, *arXiv: 2101.12013*.
- [26] J. Isberg, U. Lindstrom, B. Sundborg, and G. Theodoridis, *Classical and quantized tensionless strings*, *Nucl. Phys. B* **411** (1994) 122–156.
- [27] A. Bagchi and R. Gopakumar, *Galilean Conformal Algebras and AdS/CFT*, *JHEP* **07** (2009) 037.
- [28] A. Bagchi and I. Mandal, *On Representations and Correlation Functions of Galilean Conformal Algebras*, *Phys. Lett. B* **675** (2009) 393–397.
- [29] E. Hijano and C. Rabideau, *Holographic entanglement and Poincaré blocks in three-dimensional flat space*, *JHEP* **05** (2018) 068.
- [30] M. R. Setare and H. Adami, *Enhanced asymptotic BMS3 algebra of the flat spacetime solutions of generalized minimal massive gravity*, *Nucl. Phys. B* **926** (2018) 70-82.
- [31] J. D. Brown and M. Henneaux, *Central Charges in the Canonical Realization of Asymptotic Symmetries: An Example from Three-Dimensional Gravity*, *Commun.Math.Phys.* **104** (1986) 207-226.
- [32] E. Inonu and E. P. Wigner, *Representations of the Galilei Group*, *Nuovo Cimento IX*, **705**, (1952).
- [33] A. Bagchi and R. Gopakumar, *Galilean Conformal Algebras and AdS/CFT*, *JHEP* **07** (2009) 037.
- [34] H. Bondi, M.G.J. van der Burg, A.W.K. Metzner, *Gravitational waves in general relativity, VII. Waves from axi-symmetric isolated system*, *Proc. R. Soc. Lond. Ser. A* **269** (Aug., 1962) 21 - 52.
- [35] R. K. Sachs, *Gravitational waves in general relativity VIII. Waves in asymptotically flat space-time*, *Proc. R. Soc. Lond. Ser. A* **270** (Oct., 1962) 103 - 126 .
- [36] R. K. Sachs, *Asymptotic Symmetries in Gravitational Theory*, *Phys. Rev.* **128** (1962) 2851.
- [37] H. Jiang, W. Song and Q. Wen, *Entanglement Entropy in Flat Holography*, *JHEP* **07** (2017) 142.

- [38] J.L. Cardy, *Operator Content of Two-Dimensional Conformally Invariant Theories*, *Nucl. Phys. B* **270** (1986) 186.
- [39] T. Hartman, C.A. Keller and B. Stoica, *Universal Spectrum of 2d Conformal Field Theory in the Large c Limit*, *JHEP* **09** (2014) 118.