

# Action principle selection of regular black holes

Breno L. Giacchini,<sup>\*</sup> Tibério de Paula Netto,<sup>†</sup> and Leonardo Modesto<sup>‡</sup>

*Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China*

**Abstract.** We elaborate on the role of higher-derivative curvature invariants as a quantum selection mechanism of regular spacetimes in the framework of the Lorentzian path integral approach to quantum gravity. We show that for a large class of black hole metrics prominently regular there are higher-derivative curvature invariants associated with a singular term in the action. Therefore, according to the finite action principle applied to a general higher-derivative gravity model, not only singular spacetimes but also some of the regular ones seem to not contribute to the path integral.

## I. INTRODUCTION

It is widely acknowledged that quantum effects would be a way of resolving the spacetime singularities present in Einstein's theory. We hereby consider the appealing possibility that a finite action principle for gravity, consistent with good quantum properties like renormalizability and/or finiteness, may be at the core of the resolution to the singularity issue.

In one of the perturbative approaches to quantum gravity (QG) it is common to consider higher-derivative actions, as they can result in a (super)renormalizable theory [1, 2]. From the classical point of view, at least in the linear approximation, higher derivatives can regularize the modified Newtonian potential [3–5] and the linearized curvature invariants associated with a Dirac delta source [5–7]. However, this is not enough to claim that singularities cannot appear in the nonlinear regime nor rule out singular solutions that couple to other energy configurations (see, for instance, [8–12]).

In the work [13] it was remarked that higher derivatives could also have a fruitful role in the Lorentzian path integral approach to QG, in what concerns the cosmological singularity. The central assumption here is that metrics which make the action to diverge produce a destructive interference in the quantum gravity functional integral

$$\int Dg_{\mu\nu} e^{iS[g_{\mu\nu}]}.$$
 (1)

Therefore, the integration over all possible geometries selects with higher weight the spacetime configurations for which the action is finite. (Earlier considerations on the finite action principle were formulated in [14]; see also [15–18] for a more recent discussion and further applications.) We remind that in the path integral (1) all kinds of metrics are included, not only those which are solutions of the classical equations of motion.

A similar reasoning was applied, in Ref. [16], to the black hole singularities. The authors argued that viable gravitational actions are the ones which diverge for

singular metrics. A simple example is provided by the Schwarzschild metric, which is singular but Ricci-flat. Thus, the Einstein–Hilbert action evaluated at this solution is null, while the general fourth-derivative gravity action (see Eq. (5) below) diverges. Following [16], this observation would support fourth-derivative gravity against Einstein gravity because, in the former, the singular Schwarzschild metric would be filtered out.

In the present work we elaborate more on the role played by higher-derivative invariants in the gravitational action as a selection mechanism of regular spacetimes. Here we go beyond the analysis of [16], and study also the regularity of the terms *involving derivatives of curvature tensors*. Our central argument is that these scalars impose further constraints on the space of regular metrics which contribute to the path integral. It is instructive to begin by recalling the relevance of such terms already in the perturbative framework, where they are closely related to the problems of renormalizability and unitarity in QG.

In fact, starting from the quantum version of general relativity, the power-counting analysis shows that there are diagrams with logarithmic superficial divergence that satisfy the relation

$$d = 2L + 2,$$
 (2)

where  $d$  is the number of derivatives acting on the external graviton lines and  $L$  denotes the number of loops in the diagram. Hence, from (2) and the covariance of the counterterms, one can expect the presence, at the one-loop level, of the counterterms [19–21]

$$R_{\mu\nu\alpha\beta}^2, \quad R_{\mu\nu}^2, \quad R^2, \quad \square R,$$
 (3)

( $\square = g^{\mu\nu}\nabla_\mu\nabla_\nu$  denotes the d'Alembertian) while at the two-loop level one has

$$R_{\mu\nu\alpha\beta} R^{\alpha\beta}{}_{\cdot\rho\sigma} R^{\rho\sigma\mu\nu}, \quad R_{\mu\nu} R^{\mu\lambda\alpha\beta} R^\nu{}_{\lambda\alpha\beta}, \quad \dots, \quad R^3, \\ (\nabla_\lambda R_{\mu\nu\alpha\beta})^2, \quad \dots, \quad R_{\mu\nu} \square R^{\mu\nu}, \quad R \square R, \quad \square^2 R,$$
 (4)

and so on. Since the first structure in (4) does not vanish on the mass shell [22, 23], the theory is not renormalizable. New types of counterterms must be introduced at every order in the loop expansion and, for each of them, it is necessary to fix a renormalization condition—which involves performing a measurement. Thus, in the

<sup>\*</sup> breno@sustech.edu.cn

<sup>†</sup> tiberio@sustech.edu.cn

<sup>‡</sup> lmodesto@sustech.edu.cn

usual sense, the non-renormalizability of Einstein gravity means that the theory has no predictive power.

However, general relativity is well suited for the effective theory approach [24], because the graviton is a massless field and the next degrees of freedom due to the higher derivatives have huge masses (see, e.g., [25]). Indeed, performing an expansion in the energy  $\mathcal{E}$ , the higher-derivative terms are suppressed by powers of  $\mathcal{E}/M_{\text{P}}$ , where  $M_{\text{P}} \sim 10^{19}$  GeV is the Planck mass. At high energies all the relevant operators should be considered in the gravitational action accordingly. In particular, the power counting (2) shows that at  $L$ -loops order it is necessary to introduce into the action terms with up to  $2L + 2$  derivatives of the metric. For a recent discussion about a gauge- and parametrization-independent formulation of the renormalization group in effective quantum gravity, see [26].

The renormalization of QG models depends, of course, on the choice of the initial classical action. Regarded as a fundamental (i.e., not effective) quantum field theory, the Einstein–Hilbert action  $S_{\text{EH}}$  (including the cosmological constant) augmented by fourth-derivate invariants leads to a quantum theory which is proven to be renormalizable [1], namely

$$S = S_{\text{EH}} + \int d^4x \sqrt{-g} \{a_1 C^2 + a_2 E + a_3 \square R + a_4 R^2\}, \quad (5)$$

where  $C^2 = R^2_{\mu\nu\alpha\beta} - 2R^2_{\mu\nu} + 1/3 R^2$  is the square of the Weyl tensor and  $E = R^2_{\mu\nu\alpha\beta} - 4R^2_{\mu\nu} + R^2$  denotes the integrand of the topological Gauss–Bonnet term in four dimensions. The theory (5) also represents the minimal set of terms required in the gravitational action for having a consistent quantum field theory of matter fields in curved spacetime. In fact, even if the metric is not quantized, but treated as an external classical field, the higher-derivative terms in (5) are generated with logarithmic divergent coefficients in vacuum diagrams (see, e.g., [27–29]).

It deserves to be noted that although the Gauss–Bonnet and the  $\square R$  terms do not affect the classical dynamics of gravity, their renormalization is responsible for important quantum effects. For example, a significant part of the conformal (trace) anomaly is due to these structures' renormalization (see, e.g., [27, 28, 30]). Also, the integration of the  $\square R$ -term in the conformal anomaly can generate a finite local term  $\int d^4x \sqrt{-g} R^2$  [31–33], which can have fruitful applications in the construction of cosmological inflationary models [34–39]. Therefore, such types of operators cannot be forgotten and must be included into the classical action entering the path integral (1).

One of the drawbacks of higher-derivative models are the ghost-like particles that usually exist in the spectrum if the Feynman quantization prescription is applied [1]. However, recent developments on Lee–Wick [2, 40–45] and nonlocal theories [46–49] have brought new insights on how to handle (or avoid) the ghosts and restore uni-

tarity at any perturbative order<sup>1</sup>. In both classes of models, the classical action contains *more* than four derivatives of the metric. In the former case, besides the terms in (5) we also include operators that are polynomial in the d'Alembertian,

$$\sum_{n=1}^N \int d^4x \sqrt{-g} \left\{ \omega_n^C C_{\mu\nu\alpha\beta} \square^n C^{\mu\nu\alpha\beta} + \omega_n^R R \square^n R \right\}, \quad (6)$$

where the coefficients  $\omega_n^{C,R}$  are chosen in such a way that all the ghosts correspond to complex conjugate pairs of poles in the propagator. The unitarity of the  $S$ -matrix can be preserved if the complex poles are quantized *à la* Lee–Wick [43, 50–52].

The presence of derivatives higher than four also renders this theory super-renormalizable. This can be readily seen from the power-counting formula which, instead of (2), now yields [2]

$$d = 4 - 2N(L - 1) \quad (7)$$

(here  $N$  is the highest power of the d'Alembertian, see Eq. (6), and we assume  $\omega_N^C, \omega_N^R \neq 0$ ). Since (7) decreases with the number of loops, for  $N = 1$  the divergences exist up to  $L = 3$ , while for  $N \geq 3$  only one-loop divergences remain. For the same reason, the theory with  $N \geq 3$  has one-loop exact beta functions (regardless of  $N$ , these beta functions do not depend on the gauge and the parametrization of the quantum variables [2], as shown in [2, 53, 54]). We stress that for a given  $N$ , if there are one-loop diverging diagrams, the counterterms always have zero, two, or four derivatives of the metric. Therefore, the coefficients  $\omega_n^{C,R}$  do not get renormalized, being fixed by the choice of the initial bare classical action.

The terms quadratic in curvature in (6) represent the minimal set of operators required for having super-renormalizability. Since they contribute both to the propagator and the vertices, we call them *kinetic terms*.

In addition to the kinetic terms, there is also the possibility of including a non-minimal structure characterized by  $O(\mathcal{R}^3)$ -terms, which, therefore, only contributes to the interaction vertices (here and in the following  $\mathcal{R}$  denotes a generic curvature tensor). If such terms contain up to  $2N + 4$  derivatives of the metric tensor, the power-counting formula (7) does not change and the super-renormalizability of the model is not spoiled. For example, one can include the following operators,

$$R^2 \square^{N-2} R^2 \quad \text{and} \quad R_{\mu\nu} R^{\mu\nu} \square^{N-2} R_{\alpha\beta} R^{\alpha\beta}. \quad (8)$$

These terms can be used to kill the loop divergences by a judicious choice of their front coefficients, and the theory turns out to be finite at quantum level [2, 41, 55, 56].

<sup>1</sup> As the technical details regarding unitarity in these models lie beyond the scope of this work, we refer the interested reader to the original references cited above for further considerations.

It is worthwhile to mention that in principle one could also include in the action the following Gauss–Bonnet-like structures,

$$\text{GB}_n = R_{\mu\nu\alpha\beta}\square^n R^{\mu\nu\alpha\beta} - 4R_{\mu\nu}\square^n R^{\mu\nu} + R\square^n R, \quad (9)$$

which are not topological for  $n \neq 0$ . However, the terms in (9) can be converted to  $O(\mathcal{R}^3)$  invariants by means of the Bianchi identities (see, e.g., [2, 57]). Therefore, they do not contribute to the flat spacetime propagator. Because of this, one may change the kinetic operators which are written in (6) in the Weyl-tensor basis to another basis that uses  $R_{\mu\nu\alpha\beta}\square^n R^{\mu\nu\alpha\beta}$  or  $R_{\mu\nu}\square^n R^{\mu\nu}$  without modifying the flat-space propagator [2]. The procedure is similar to what is usually done in the fourth-derivative gravity (5).

In the discussion above we emphasized the importance, in different QG models, of curvature invariants which can be simply polynomials in the curvature tensors or polynomials in covariant derivatives of the curvatures<sup>2</sup>. As already mentioned, our goal in the present work is to consider the effect of such terms in the path integral approach to QG as a mechanism for selecting regular spacetime configurations. To this end, we here study the regularity of the curvature-derivative scalars evaluated at regular black hole metrics<sup>3</sup>. We mainly focus on invariants of the type  $\mathcal{R}\square^n\mathcal{R}$ , for these structures contain the highest number of derivatives acting on a single metric component and, thus, may produce the strongest singularity. For the sake of simplicity, we only consider static spherically symmetric metrics, defined by the line element

$$ds^2 = -A(r)dt^2 + B(r)dr^2 + r^2d\Omega^2, \quad (10)$$

where  $A(r)$  and  $B(r)$  are two generic functions and  $d\Omega^2 = d\theta^2 + \sin^2\theta d\varphi^2$  is the metric of the unit two-sphere. We also adopt units such that  $c = \hbar = G = 1$ .

The rest of the paper is organized as follows. In Sections II and III we evaluate and study the curvature invariants and curvature-derivative invariants for a large class of spherically symmetric spacetimes. It turns out that for some regular black holes the curvature scalars are bounded, but some curvature-derivative scalars diverge. In Section IV we use the results of the previous sections to evaluate the contribution of the curvature-derivative invariants to the action. It is the main result of this paper that also some regular black holes do not contribute to the Lorentzian path integral. Finally, in Section V we briefly comment on non-analytic regular metrics; our conclusions are summarized in Section VI.

## II. EXAMPLES OF SINGULAR CURVATURE INVARIANTS FOR REGULAR BLACK HOLES

Many “regular black hole” metrics have been presented in the literature [62–85] as exact or approximate solutions of the equations of motion in a certain classical gravity theory, as an effective treatment of quantum corrections, or motivated by other theoretical grounds. In what follows we are going to discuss some of these metrics, but before moving on it is necessary to fix the convention of what we call a “regular” black hole. Definitions of regular black holes can rely on the regularity of the metric components in a given coordinate chart, on the regularity of the associated Christoffel symbols, the regularity of a set of curvature invariants, or in a limiting curvature condition (see, for instance, [62, 63] and references therein). In the case of the definition based on the finiteness of the curvature invariants, it is usually assumed that the curvature-squared scalars  $\mathcal{R}^2$  are nonsingular. In the present work we will adopt this same definition. More precisely, since we only consider spacetime configurations which may yield diverging scalars at  $r = 0$ , we shall say that a metric in the form (10) is regular if all the invariants  $R_{\mu\nu\alpha\beta}^2$ ,  $R_{\mu\nu}^2$ ,  $C^2$  and  $R$  are finite at  $r = 0$ .

In this spirit, throughout this paper (except for Sec. V) we only consider metric components which are analytic functions around  $r = 0$ , namely

$$A(r) = \sum_{\ell=0}^{\infty} a_{\ell}r^{\ell}, \quad B(r) = \sum_{\ell=0}^{\infty} b_{\ell}r^{\ell}. \quad (11)$$

One can verify that the conditions for regularity then read (see, e.g., [62] or the Sec. III below)

$$a_0 \neq 0, \quad b_0 = 1, \quad a_1 = b_1 = 0. \quad (12)$$

The regularity of curvature-derivative invariants in the Newtonian limit has been studied in detail in Ref. [7]. The main result can be summarized as follows: *assuming that (12) holds and given a positive integer  $n$ , if the Taylor series of both the potentials  $A(r)$  and  $B(r)$  do not contain any odd power of  $r$  up to (including)  $r^{2n+3}$ , then all the curvature-derivative invariants with at most  $2n$  covariant derivatives are regular.*

For example, if the first non-zero odd coefficient occurs at  $O(r^5)$ , all invariants of the type  $\mathcal{R}\square\mathcal{R}$  are finite, while the first singularity appears in the set of the operators of the type  $\mathcal{R}\square^2\mathcal{R}$ . This means, in particular, that if the metric potentials are even functions of  $r$ , all polynomials constructed from the curvature tensors and their derivatives are finite at  $r = 0$ . *Here we conjecture that the same is true in the nonlinear regime as well.*

A thorough investigation of this problem is beyond the scope of the paper, as for our purpose it suffices to identify diverging curvature-derivative scalars—more exactly, scalars that yield a diverging action, see discussion in Sec. IV. However, it is instructive to present some explicit examples supporting the conjecture. In the Table I

<sup>2</sup> Sometimes we shall refer to these scalars as “curvature-derivative” invariants.

<sup>3</sup> Polynomial curvature-derivative invariants also have an interesting application in identifying the position of black hole horizons [58–61].

we list some regular black hole metrics that are even analytic functions. Using WOLFRAM MATHEMATICA [86] and the tensor algebra package xACT [87, 88], we have evaluated a set of curvature-derivative invariants containing up to twenty-two derivatives of a single metric component: all the invariants calculated are finite at  $r = 0$ .

Metric functions: $A(r) = 1/B(r)$	Ref.
$A(r) = 1 - \frac{2mr^2}{(r^2 + \alpha^2)^{3/2}}$	[64]
$A(r) = 1 - \frac{2mr^2}{(r^2 + q^2)^{3/2}} + \frac{q^2 r^2}{(r^2 + q^2)^2}$	[65]
$A(r) = 1 - \frac{4m}{\pi r} \left( \tan^{-1} \frac{r}{r_0} - \frac{rr_0}{r^2 + r_0^2} \right)$	[66]
$A(r) = 1 + \frac{4M}{\sqrt{\pi r}} \gamma \left( \frac{3}{2}, \frac{r^2}{4\theta} \right)$	[67]

TABLE I. Examples of regular metrics defined by even functions for which curvature polynomials and curvature-derivative polynomial invariants are singularity-free.

Let us mention that other examples of spherically symmetric metrics defined by even analytic functions have been obtained in the fourth-derivative gravity (namely, the so-called (0,0) family of solutions [8–10]) and in the framework of nonlocal higher-derivative gravity [46–49]. See, for instance, the solutions in [72–76]; for more general results, including logarithmic quantum corrections, the reader can consult Ref. [7].

In what follows we present some examples of regular black hole metrics for which some curvature invariants are singular.

### A. Renormalization group improved black hole

In Ref. [77] it was considered the improvement of the Schwarzschild metric due to the running of the Newton constant, obtained from the flow equation for the average effective action truncated at the Einstein–Hilbert order. The resulting metric is defined by (10) with the functions

$$A(r) = \frac{1}{B(r)} = 1 - \frac{2Mr^2}{r^3 + \omega(r + \gamma M)}, \quad (13)$$

where  $M$  is the mass of the black hole measured by a distant observer, and  $\omega$  and  $\gamma$  are parameters related to the running. Expanding the function  $A(r)$  around  $r = 0$  we obtain

$$A(r) = 1 - \frac{2}{\gamma\omega} \left[ r^2 - \frac{r^3}{\gamma M} + \frac{r^4}{(\gamma M)^2} \right] + O(r^5). \quad (14)$$

It is straightforward to verify that  $R$  and the quadratic-curvature scalars such as  $R_{\mu\nu}^2$  and  $R_{\mu\nu\alpha\beta}^2$  are finite [77], as expected for a metric with a de Sitter core. For example,

$$R(r) = \frac{24}{\gamma\omega} - \frac{40}{\gamma^2 M \omega} r + \frac{60}{\gamma^3 M^2 \omega} r^2 + O(r^3). \quad (15)$$

However, since the first odd power in the series (14) occurs at  $r^3$ , as anticipated, the invariants with two deriva-

tives of the curvature tensors may already diverge. Indeed,

$$\square R \underset{r \rightarrow 0}{\sim} -\frac{80}{\gamma^2 M \omega r}, \quad (16)$$

while those containing higher derivatives can diverge to a higher power. For instance,

$$R_{\mu\nu} \square^2 R^{\mu\nu} \underset{r \rightarrow 0}{\sim} \frac{256}{\gamma^4 M^2 \omega^2 r^2}, \quad (17)$$

$$R_{\mu\nu} \square^3 R^{\mu\nu} \underset{r \rightarrow 0}{\sim} \frac{256 (27\gamma^2 M^2 + 32\gamma M^2 + 27\omega)}{\gamma^6 M^4 \omega^3 r^2}, \quad (18)$$

$$R_{\mu\nu} \square^4 R^{\mu\nu} \underset{r \rightarrow 0}{\sim} \frac{36864}{\gamma^6 M^3 \omega^3 r^3}, \quad (19)$$

$$R_{\mu\nu} \square^5 R^{\mu\nu} \underset{r \rightarrow 0}{\sim} \frac{221184}{\gamma^6 M^3 \omega^3 r^5}. \quad (20)$$

The same singularity pattern presented above occurs for the invariants  $R_{\mu\nu\alpha\beta} \square^n R^{\mu\nu\alpha\beta}$  and  $C_{\mu\nu\alpha\beta} \square^n C^{\mu\nu\alpha\beta}$ , while all  $\square^n R$  and  $R \square^n R$  we have evaluated diverge like  $1/r$ .

### B. Hayward spacetime

The Hayward black hole [78] is defined by the function

$$A(r) = \frac{1}{B(r)} = 1 - \frac{2Mr^2}{r^3 + 2L^3}, \quad (21)$$

where  $L > 0$  is a parameter related to the de Sitter center. The behavior of the metric (21) for small  $r$  differs from the (13). For (21) the leading odd-order term in the series expansion occurs at the order  $r^5$ :

$$A(r) = 1 - M \left[ \frac{r^2}{L^3} - \frac{r^5}{2L^6} + \frac{r^8}{4L^9} + O(r^{11}) \right]. \quad (22)$$

It is not difficult to verify that the associated Ricci scalar,

$$R = -\frac{24L^3 M (r^3 - 4L^3)}{(r^3 + 2L^3)^3} = \frac{12M}{L^3} + O(r^3), \quad (23)$$

is regular, as well as the curvature-squared scalars. In Ref. [16] the calculation of some regular scalars containing up to eight derivatives of the metric was presented, and it was shown, e.g., that also the six-derivative scalar  $(\nabla_\rho R_{\mu\nu\alpha\beta})^2$  is nonsingular for the Hayward spacetime. Nonetheless, another scalar containing six derivatives of the metric diverges,

$$\square^2 R \underset{r \rightarrow 0}{\sim} -\frac{504M}{L^6 r}. \quad (24)$$

If one is worried that (24) is a total derivative, there exist other singular invariants with eight and more derivatives of the metric that diverge too. For example,

$$R \square^2 R \underset{r \rightarrow 0}{\sim} -\frac{6048M^2}{L^9 r}. \quad (25)$$

Invariants with more derivatives can diverge even more strongly, e.g.,

$$(\nabla_\alpha \nabla_\beta \nabla_\gamma \nabla_\delta \nabla_\varepsilon R_{\mu\nu\rho\sigma})^2 \underset{r \rightarrow 0}{\sim} \frac{304560M^2}{L^{12}r^4}, \quad (26)$$

$$R_{\mu\nu\alpha\beta} \square^7 R^{\mu\nu\alpha\beta} \underset{r \rightarrow 0}{\sim} -\frac{6065280}{L^{12}M^3r^3}. \quad (27)$$

### C. Dymnikova spacetime

A similar situation occurs for the Dymnikova metric [79], defined by (10) with the function

$$A(r) = \frac{1}{B(r)} = 1 - \frac{2M(1 - e^{-\frac{r^3}{L^3}})}{r}, \quad (28)$$

where  $L \equiv 2Mr_0^2$  and  $r_0$  is a constant related to the de Sitter core. The power series of  $A(r)$  around  $r = 0$  reads:

$$A(r) = 1 - 2M \left[ \frac{r^2}{L^3} - \frac{r^5}{2L^6} + \frac{r^8}{6L^9} + O(r^{11}) \right]. \quad (29)$$

Since the series (22) and (29) coincide to the order  $r^6$  (up to a factor 2 associated with  $M$ ), it follows that the small- $r$  behavior of the scalars  $\square^2 R$  and  $R \square^2 R$  for the Dymnikova spacetime are given by the Eqs. (24) and (25) up to the replacement  $M \mapsto 2M$ . Higher-derivative invariants have diverging behavior similar to the ones of the Hayward metric, but with different coefficients. Therefore, although  $R$  and the quadratic scalars are bounded [79], there are singular scalars with six and more derivatives of the metric.

### D. Modified Hayward spacetimes

Now we consider an example of metric for which  $A(r) \neq 1/B(r)$ . In Ref. [80] it was proposed a modification of the Hayward metric that includes the leading infrared (IR) one-loop quantum corrections to the Newtonian potential [89, 90]<sup>4</sup>

$$\phi(r) = -\frac{M}{r} \left( 1 + \beta \frac{1}{r^2} \right). \quad (30)$$

The metric has the form

$$A(r) = \left( 1 - \frac{2M\alpha\beta}{r^3\alpha + M\beta} \right) \frac{1}{B(r)}, \quad (31)$$

$$B(r) = \left( 1 - \frac{2Mr^2}{r^3 + 2L^3} \right)^{-1}. \quad (32)$$

Bearing in mind the Hayward metric (21), the extra factor in  $A(r)$  has the property of encoding a non-trivial time delay between  $r = 0$  and an observer in the infinity because  $A(0) = 1 - 2\alpha \neq 1$ .

Expanding  $A(r)$  around  $r = 0$  we get:

$$A(r) = 1 - 2\alpha - \frac{(1 - 2\alpha)M}{L^3} r^2 + \frac{2\alpha^2}{M\beta} r^3 + O(r^5). \quad (33)$$

Differently from the Hayward metric (21), now the first odd-power term occurs already at order  $r^3$  [c.f. Eq. (22)], whereas for  $B(r)$  it is still at  $O(r^5)$  because this function is unchanged. By direct computation one can verify that, although  $R$  and the curvature-squared scalars are bounded at  $r = 0$ ,

$$\square R \underset{r \rightarrow 0}{\sim} -\frac{48\alpha^2}{(1 - 2\alpha)\beta M r} \quad (34)$$

diverges, and other curvature-derivative invariants have singular behavior alike those discussed in Sec. II A.

Another class of modified Hayward metrics with non-trivial time shift, introduced in [62], has a similar feature. If the series of  $A(r)$  contains the power  $r^3$  then  $\square R$  already diverges; otherwise, it is necessary to apply two d'Alembertians to get a singular scalar, like in the case of the original Hayward metric. This happens because the series of  $B(r)$  is unchanged and does contain the term  $r^5$ .

## III. ON THE EXISTENCE OF SINGULAR CURVATURE INVARIANTS

In the previous section we provided examples of regular black hole metrics with singular curvature-derivative invariants. Since the calculation of a large number of curvature scalars is a cumbersome task, here we present some more general considerations on the existence of singular curvature invariants for a given regular metric in the form (10).

We start assuming that  $A(r)$  and  $B(r)$  are smooth functions that have a Taylor series representation around  $r = 0$ ,

$$A(r) = a_0 + \sum_{\ell=1}^{\infty} a_\ell r^\ell, \quad B(r) = 1 + \sum_{\ell=1}^{\infty} b_\ell r^\ell. \quad (35)$$

Moreover, let us assume that their first odd power are, respectively,  $r^{2n_a+1}$  and  $r^{2n_b+1}$  for some  $n_a, n_b \in \mathbb{N}$ . This means that  $a_{2\ell+1} = 0$  for all  $\ell$  smaller than  $n_a$ ,  $b_{2\ell+1} = 0 \forall \ell < n_b$ , and  $a_{2n_a+1}, b_{2n_b+1} \neq 0$ .

It is useful to introduce a notation for the property of a function  $F(r)$  to have the first odd-order term in the Taylor expansion at a certain order  $r^{2n+1}$ . If a function  $F(r)$  has this property, we shall denote it by writing  $\mathcal{P}[F] = n$ . In terms of this notation, we have  $\mathcal{P}[A] = n_a$  and  $\mathcal{P}[B] = n_b$ .

<sup>4</sup> See also [7, 91] for a recent discussion concerning the universality of this correction in higher-derivative gravity models.

Now, consider the action of the d'Alembert operator on a smooth scalar  $F(r)$ ,

$$\square F = \frac{1}{B} \left[ 2 \frac{F'}{r} + F'' - \frac{(AB' - A'B)F'}{2AB} \right]. \quad (36)$$

The function  $F$  could be, for example, the scalar curvature  $R$ , or  $R_{\mu\nu}^2$ . Expanding  $F(r)$  as a power series around  $r = 0$ ,

$$F(r) = \sum_{\ell=0}^{\infty} f_{\ell} r^{\ell}, \quad (37)$$

it is straightforward to verify that if  $\mathcal{P}[F] \equiv n_f \geq 1$ , then

$$\mathcal{P}[F'/r] = \mathcal{P}[F''] = n_f - 1,$$

whereas the first odd-order power of the last term in (36) is at least  $O(r^{2 \min\{n_f, n_a, n_b\} + 1})$ .

Hereafter, we assume  $n_f < n_a, n_b$ , so that the order of the first overall odd-power term only depends on  $F$ . This particular choice will be justified later, when we substitute this function by the scalar curvature. In fact, under this assumption, the first odd-power term in the series of  $\square F(r)$  is

$$2(n_f + 1)(2n_f + 1) f_{2n_f+1} r^{2n_f-1} \neq 0. \quad (38)$$

Therefore,  $\mathcal{P}[\square F] = n_f - 1$ . From Eq. (38) one can see that if  $n_f = 0$  then  $\square F$  diverges like  $r^{-1}$ ; otherwise it is regular. The iteration of this procedure shows that  $\mathcal{P}[\square^{n_f} F] = 0$ ; hence,  $\square^{n_f+1} F$  has a singularity of the type  $r^{-1}$ .

Let us now turn the attention to the Ricci scalar associated with the metric in (10),

$$R = \frac{(A')^2}{2A^2B} + 2 \frac{B-1}{r^2B} + 2 \frac{AB' - A'B}{rAB^2} + \frac{A'B' - 2A''B}{2AB^2}. \quad (39)$$

The requirement that  $R$  is regular immediately gives<sup>5</sup>

$$a_0 \neq 0 \quad \text{and} \quad a_1 = 2a_0b_1. \quad (40)$$

It is easy to verify, however, that even when (40) holds for  $b_1 \neq 0$  the invariants  $R_{\mu\nu\alpha\beta}^2$  and  $R_{\mu\nu}^2$  are singular. For example, if this is the case,

$$R_{\mu\nu\alpha\beta}^2 \underset{r \rightarrow 0}{\sim} \frac{14b_1^2}{r^2} \quad \text{and} \quad R_{\mu\nu}^2 \underset{r \rightarrow 0}{\sim} \frac{11b_1^2}{2r^2}. \quad (41)$$

In contrast, only if

$$a_1 = b_1 = 0, \quad (42)$$

all the curvature scalars built only with curvature tensors (i.e., without its covariant derivatives) are finite. This is,

of course, in agreement with the known results in the literature (see, e.g., [62]) and with the discussion in the beginning of the Sec. II. Therefore, since our main objective is to study the new singularities produced when covariant derivatives act on regular quantities, in the rest of the paper we assume (42).

A moment's reflection reveals that, in most of the cases, the first odd-order term in the Taylor expansion of  $R(r)$  is at order  $r^{2N-1}$ , where  $N \equiv \min\{n_a, n_b\}$ . In fact, for the first term in the right-hand side of (39), the first odd-order power is at least  $O(r^{2n_a+1})$ ; for the second term it is exactly at  $O(r^{2n_b-1})$ ; for the third term it is  $O(r^{2N-1})$ , and for the last one it is  $O(r^{2n_a-1})$ . The cancellation of the  $O(r^{2N-1})$ -terms can take place only if the functions  $A$  and  $B$  are related in a very particular way.

Indeed, assuming  $N = n_a < n_b$  one can explicitly evaluate the coefficient of the  $O(r^{2N-1})$ -term in the series of  $R$ , which reads,

$$-2(n_a + 1)(2n_a + 1) \frac{a_{2n_a+1}}{a_0} \neq 0, \quad (43)$$

since  $a_{2n_a+1} \neq 0$  by definition [see the discussion after Eq. (35)]. On the other hand, if  $N = n_b < n_a$ , the coefficient of the  $O(r^{2N-1})$ -term is

$$4(n_b + 1)b_{2n_b+1} \neq 0. \quad (44)$$

The remaining scenario is when  $n_a = n_b = N$ . In this case the first odd-power term is

$$-2(N + 1) \left[ (2N + 1) \frac{a_{2N+1}}{a_0} - 2b_{2N+1} \right] r^{2N-1}, \quad (45)$$

which vanishes if and only if

$$a_{2N+1} = \frac{2a_0b_{2N+1}}{2N + 1}. \quad (46)$$

Solely for this particular combination of coefficients the term  $r^{2N-1}$  is absent in the Taylor representation of  $R(r)$ . However, higher odd powers can occur depending on the other coefficients of  $A(r)$  and  $B(r)$ . It is important to remark that the relation (46) cannot be satisfied for metrics with  $A(r) = 1/B(r)$  because, although  $n_a = n_b$ , in this case the coefficients satisfy the relation  $a_{2N+1} = -a_0^2 b_{2N+1}$ , which is not compatible with (46).

Hence, unless (46) is satisfied, the Ricci scalar is such that

$$\mathcal{P}[R(r)] = N - 1 < n_a, n_b. \quad (47)$$

In particular, the hypothesis  $n_f < n_a, n_b$  evoked in the considerations concerning the regularity of  $\square F$  is satisfied by the scalar  $F = R$ . Thus, from the previous results it follows that  $\square R \sim r^{-1}$  for  $N = 1$ , while if  $N \geq 2$  we have  $\mathcal{P}[\square R] = N - 2$ . Furthermore,  $\mathcal{P}[\square^{N-1} R] = 0$  and, thus,  $\square^N R$  has a singularity of the type  $r^{-1}$ .

Some of the examples presented in Sec. II can be derived from this general result. Indeed, for the metrics of the subsections A, B and C we immediately get

<sup>5</sup> The choice  $B(0) = 1$  assumed in (35) is also necessary for the regularity of  $R$ .

$\mathcal{P}[R(r)] = N - 1$ , inasmuch as they have  $B(r) = 1/A(r)$ . For the Hayward and Dymnikova black holes, the first odd-order term in the series of the metric is  $r^5$  (i.e.,  $N = 2$ ), thus  $R$  and  $\square R$  are finite but  $\square^2 R$  is singular. In contrast,  $N = 1$  for the renormalization group improved black hole metric, so  $\square R$  already diverges. Finally, for the modified Hayward metric (31) of subsection D we have  $n_a = 1$  and  $n_b = 2$ ; whence  $N = n_a = 1$  and  $\square R$  is singular at  $r = 0$ .

#### IV. SINGULARITIES OF SUPERFICIAL AND KINETIC CURVATURE INVARIANTS

In the previous section we showed that if the series expansion of the metric (10) around  $r = 0$  contains the first odd-power term at order  $r^{2N+1}$ , given a regular curvature invariant  $F(r)$  such that  $\mathcal{P}[F] = n_f < N$ , the operator  $\square^{n_f+1} F$  might diverge like  $r^{-1}$ . This is the case for the scalar curvature if the relation (46) does not hold. Moreover, if  $N \geq 1$ , all the terms of the type  $R\square^n R$  for  $n < N$  are finite; while  $R\square^N R$  may diverge like  $r^{-1}$  as  $r \rightarrow 0$ .

However, from the point of view of the finite action principle, singularities of the type  $r^{-1}$  and  $r^{-2}$  are harmless in four-dimensional spacetimes because they do not produce a divergent action. In fact, the contribution of a given curvature invariant  $F(r)$  to the gravitational action is through the term

$$S_F(\Omega_{UV}, \Omega_{IR}) = \int_{\Omega_{UV}}^{\Omega_{IR}} dr \sqrt{-g} F(r), \quad (48)$$

where, following the procedure of [16], we introduced infrared and ultraviolet cutoffs and restrict considerations to the integral on the coordinate  $r$ , which is the only relevant one for static spherically symmetric metrics. Since the factor

$$\sqrt{-g} = \sqrt{A(r)B(r)} r^2 = \sqrt{a_0} r^2 + O(r^4) \quad (49)$$

in the measure is  $O(r^2)$  for a regular metric, only if  $F(r)$  is more singular than  $r^{-2}$  the action term  $S_F$  will diverge when we let  $\Omega_{UV} \rightarrow 0$ . Thus, our goal in this section is to investigate whether there are scalars of the type  $\square^n F$ , or  $\mathcal{R}\square^n \mathcal{R}$ , with  $n > N$  that diverge with a higher inverse power of  $r$ .

To this end, we assume that a generic curvature invariant  $I(r)$  has a  $r^{-1}$ -type of singularity, i.e.,

$$I(r) = \frac{c}{r} + F(r), \quad (50)$$

where  $c \neq 0$  is a constant and  $F(r)$  is an analytic function given by the power series (37). Hence, when we apply a d'Alembert operator on it we get, for  $r \neq 0$  [see Eq. (36)]:

$$\begin{aligned} \square I &= \frac{c(AB' - A'B)}{2AB^2 r^2} + \square F \\ &= \frac{c(a_0 b_1 - a_1)}{2a_0 r^2} + O(1/r). \end{aligned} \quad (51)$$

Therefore, only if  $a_0 b_1 \neq a_1$  it happens that  $\square I \sim r^{-2}$ . Otherwise, it can either be regular or diverge as  $r^{-1}$ , depending on the value of the coefficients  $c$ ,  $a_1$ ,  $a_2$ ,  $b_1$  and  $b_2$ , and also on  $F$ . Inasmuch as  $a_1 = b_1 = 0$  for regular metrics, the application of this result iteratively to the scalar  $\square^N R$ —which has a singularity of the type  $r^{-1}$  in the form of (50)—leads to the following conclusion: *given a regular analytic metric,  $\square^n R$  (for any power  $n$ ) can only diverge like  $r^{-1}$  for  $r \rightarrow 0$ .* In the sense of distributions, however, for  $n > N$  there can be singularities involving Dirac deltas and its derivatives.

The regularity of the kinetic scalars of the type  $R\square^n R$  also depends on the behavior of  $R$  near  $r = 0$ . Nevertheless, if  $R$  is regular,  $R\square^n R$  cannot be more diverging than  $\square^n R$ . As an important particular case, if the metric has  $B = 1/A$  and has a de Sitter core, i.e.,  $a_1 = b_1 = 0$  and  $b_2 = -a_2 \neq 0$ , then  $R(0) = -12a_2$  and  $R\square^n R$  always diverges as  $r^{-1}$  if  $\square^n R$  diverges. This explains why we found only  $1/r$ -type of divergences for these invariants in the explicit evaluations considered in Sec. II.

However, the explicit examples of Sec. II have also shown that the kinetic terms which are not quadratic in the Ricci scalar can diverge faster than  $r^{-1}$ . Consider, for instance, the invariant

$$K_n \equiv R_{\mu\nu\alpha\beta} \square^n R^{\mu\nu\alpha\beta}. \quad (52)$$

For simplicity, let us also assume that  $B = 1/A$ . If  $a_3 \neq 0$ , the invariant  $K_1$  diverges like  $r^{-1}$ , but  $K_2$  and  $K_3$  go with  $r^{-2}$ ,  $K_4 \sim r^{-3}$  and  $K_5 \sim r^{-5}$ . If one sets  $a_3 = 0$ , then all these five invariants may diverge only like  $r^{-1}$ ; nonetheless, pushing the calculations further one finds  $K_6 \sim a_4 a_5^2 r^{-2}$  and  $K_7 \sim a_4 a_5^2 r^{-4}$ . If, in addition,  $a_4 = 0$ , then the lowest-order kinetic invariant diverging stronger than  $r^{-1}$  is  $K_7 \sim a_5^3 r^{-3}$ ; this is the case of the Hayward and Dymnikova metrics, see Eq. (27). A similar situation occurs for the kinetic scalars quadratic in the Ricci or Weyl tensors [see, for instance, Eqs. (17)-(20)].

Therefore, we conclude that even for regular analytic spacetimes, curvature-derivative terms can yield a diverging action. Still, as mentioned above, it seems that there exists a relation between the order of the first odd-power term in the Taylor expansion of the metric and the maximal power of the d'Alembertian in a generic regular kinetic scalar needed to produce a divergence stronger than  $r^{-2}$ .

#### V. COMMENT REGARDING NON-ANALYTIC METRICS

So far, our considerations involved only analytic metrics in  $r = 0$ , i.e., those that can be expressed as Taylor series. However, under certain circumstances, a similar analysis can be carried out for regular metrics that are not analytic.

The most interesting scenario concerns the situation when  $A(r)$  and  $B(r)$  are non-analytic smooth functions. Many of these regular metrics have been obtained in the

literature—see, e.g., [81–85]. The main observation here is that if  $A(r)$  and  $B(r)$  tend to a constant value (as  $r \rightarrow 0$ ) faster than any polynomial, all the invariants polynomial in curvatures and their derivatives are regular. For example, consider the metric of Ref. [83],

$$A(r) = \frac{1}{B(r)} = 1 - \frac{2M}{r} [1 - \tanh(r_0/r)], \quad (53)$$

where  $r_0$  is a positive parameter. Since

$$R = \frac{4Mr_0^2 \tanh(r_0/r) \operatorname{sech}^2(r_0/r)}{r^5} \quad (54)$$

and  $\operatorname{sech}(r_0/r)$  is exponentially suppressed by a factor  $2e^{-r_0/r}$  as  $r \rightarrow 0$ , all the covariant derivatives of  $R$  will still vanish faster than any polynomial.

Another possibility of non-analytic metrics are those which are not smooth but still admit a Taylor representation around  $r = 0$ , with a non-polynomial remainder function. Though it is clear that the non-existence of a derivative of a certain order at  $r = 0$  implies that there will be singular curvature-derivative scalars, the minimal number of derivatives in such diverging scalars may be less than the order of differentiability of the functions  $A(r)$  and  $B(r)$ . Take, for example, a metric defined by the functions  $A(r)$  and  $B(r)$  which are  $n$ -times continuously differentiable at  $r = 0$  (with  $n \geq 2$ ). Let us define  $\mathcal{P}[A] = n_a$  and  $\mathcal{P}[B] = n_b$  such that the first odd-power terms in the Taylor polynomial of  $A(r)$  and  $B(r)$  are respectively at order  $r^{2n_a+1}$  and  $r^{2n_b+1}$ . Then, from the discussion of Sec. III, if  $2N \equiv 2 \min\{n_a, n_b\} < n - 1$ , it can happen that  $\square^N R$  diverges as  $r^{-1}$ . Higher-order invariants, however, can have other types of singularities, depending on the remainder function.

For instance, in the context of the sixth-derivative gravity with logarithmic quantum corrections, in Ref. [7] it was obtained a metric defined by the function

$$A(r) = \sum_{\ell=0}^{\infty} a_{\ell} r^{\ell} + \log(\mu r) \sum_{\ell=2}^{\infty} c_{2\ell+1} r^{2\ell+1}, \quad (55)$$

with  $a_1 \equiv 0$  but  $a_3, c_5 \neq 0$  ( $\mu > 0$  is a parameter and  $B(r)$  has the same general structure). Even though the function  $A(r)$  is four-times differentiable in  $r = 0$ ,  $\square R$  already diverges like  $r^{-1}$  due to the presence of the term

$r^3$  in (55), whereas  $\square^2 R \sim \log r/r$  for small  $r$ , as a consequence of non-analyticity.

## VI. DISCUSSION

In this paper we stressed that regular black hole metrics can still produce divergent curvature-derivative scalars, like  $R_{\mu\nu\alpha\beta} \square^n R^{\mu\nu\alpha\beta}$  and similar ones. The inclusion of these higher-derivative operators in a gravitational action may (or may not) cause the action to diverge. Indeed, as argued in [16], the contribution of a curvature invariant  $F(r)$  evaluated at a static spherically symmetric metric in the form (10) is

$$S_F(\Omega_{\text{UV}}, \Omega_{\text{IR}}) = \int_{\Omega_{\text{UV}}}^{\Omega_{\text{IR}}} dr \sqrt{A(r)B(r)} r^2 F(r). \quad (56)$$

Since  $\sqrt{AB}$  is finite for a regular metric, if  $F(r)$  diverges like  $r^{-n}$  with  $n > 2$ , then  $S_F(\Omega_{\text{UV}}, \Omega_{\text{IR}})$  will diverge when we send the UV cutoff to zero. If this is the case, following the finite action principle [13–16], the corresponding spacetime configuration is going to be filtered out of the functional integration over all geometries.

In view of the results presented in Sec. III, if the fourth-derivative gravity is taken as the fundamental QG theory, then for any regular black hole metric (10) the divergence that may exist at  $r = 0$  in the term  $\square R$  (which is required by renormalization) does not yield an infinite action. This happens because for these metrics such term can only diverge like  $1/r$ ; moreover, this is the sole scalar with at most four derivatives that can diverge on a regular metric.

However, as pointed out in Ref. [16], the requirement of finiteness may be a way of distinguishing viable gravitational actions in their capability of selecting regular geometries. In this spirit, a theory containing higher-derivative structures like those in (6) with large  $N$  is favored in comparison to the simple fourth-derivative gravity (5). In fact, the space of regular metrics associated with a finite action is more constrained if the the number of derivatives in the action is larger.

## ACKNOWLEDGMENTS

This work was supported by the Basic Research Program of the Science, Technology and Innovation Commission of Shenzhen Municipality (grant no. JCYJ20180302174206969).

- 
- [1] K. S. Stelle, *Renormalization of Higher Derivative Quantum Gravity*, Phys. Rev. D **16**, 953 (1977).  
 [2] M. Asorey, J. L. López and I. L. Shapiro, *Some remarks on high derivative quantum gravity*, Int. J. Mod. Phys. A

- 12**, 5711 (1997), arXiv:hep-th/9610006.  
 [3] L. Modesto, T. de Paula Netto and I. L. Shapiro, *On Newtonian singularities in higher derivative gravity models*, J. High Energy Phys. **1504**, 098 (2015),

- arXiv:1412.0740.
- [4] B. L. Giacchini, *On the cancellation of Newtonian singularities in higher-derivative gravity*, Phys. Lett. B **766**, 306 (2017), arXiv:1609.05432.
- [5] B. L. Giacchini and T. de Paula Netto, *Effective delta sources and regularity in higher-derivative and ghost-free gravity*, J. Cosmol. Astropart. Phys. **1907**, 013 (2019), arXiv:1809.05907.
- [6] B. L. Giacchini and T. de Paula Netto, *Weak-field limit and regular solutions in polynomial higher-derivative gravities*, Eur. Phys. J. C **79**, 217 (2019), arXiv:1806.05664.
- [7] N. Burzillà, B. L. Giacchini, T. de Paula Netto and L. Modesto, *Higher-order regularity in local and nonlocal quantum gravity*, arXiv:2012.11829.
- [8] K. S. Stelle, *Classical Gravity with Higher Derivatives*, Gen. Rel. Grav. **9**, 353 (1978).
- [9] H. Lü, A. Perkins, C. N. Pope and K. S. Stelle, *Black Holes in Higher-Derivative Gravity*, Phys. Rev. Lett. **114**, 171601 (2015), arXiv:1502.01028; *Spherically Symmetric Solutions in Higher-Derivative Gravity*, Phys. Rev. D **92**, 124019 (2015), arXiv:1508.00010.
- [10] B. Holdom and J. Ren, *Not quite a black hole*, Phys. Rev. D **95**, 084034 (2017), arXiv:1612.04889.
- [11] A. Bonanno and S. Silveravalle, *Characterizing black hole metrics in quadratic gravity*, Phys. Rev. D **99**, 101501 (2019), arXiv:1903.08759.
- [12] B. Holdom, *On the fate of singularities and horizons in higher derivative gravity*, Phys. Rev. D **66**, 084010 (2002), arXiv:hep-th/0206219.
- [13] J. L. Lehners and K. S. Stelle, *Safe beginning for the universe?*, Phys. Rev. D **100**, 083540 (2019), arXiv:1909.01169.
- [14] J. D. Barrow and F. J. Tipler, *Action principles in nature*, Nature **331**, 31 (1988).
- [15] J. D. Barrow, *Finite action principle revisited*, Phys. Rev. D **101**, 023527 (2020), arXiv:1912.12926.
- [16] J. N. Borissova and A. Eichhorn, *Towards black-hole singularity-resolution in the Lorentzian gravitational path integral*, Universe **7**, 48 (2021), arXiv:2012.08570.
- [17] C. Jonas, J. L. Lehners and J. Quintin, *Cosmological consequences of a principle of finite amplitudes*, arXiv:2102.05550.
- [18] J. Chojnacki and J. H. Kwapisz, *Finite Action Principle and Horava-Lifshitz Gravity: early universe, black holes and wormholes*, arXiv:2102.13556.
- [19] G. 't Hooft and M. Veltman, *One loop divergencies in the theory of gravitation*, Ann. Inst. H. Poincaré Phys. Theor. A **20**, 69 (1974).
- [20] S. Deser and P. van Nieuwenhuisen, *One-loop divergences of quantized Einstein-Maxwell fields*, Phys. Rev. D **10**, 401 (1974).
- [21] R. E. Kallosh, O. V. Tarasov and I. V. Tyutin, *One-loop finiteness of quantum gravity off mass shell*, Nucl. Phys. B **137**, 145 (1978).
- [22] M. H. Goroff and A. Sagnotti, *Quantum gravity at two loops*, Phys. Lett. **B160**, 81 (1985); *The ultraviolet behavior of Einstein gravity*, Nucl. Phys. **B266**, 709 (1986).
- [23] A. E. M. van de Ven, *Two loop quantum gravity*, Nucl. Phys. **B378**, 309 (1992).
- [24] J. F. Donoghue, *General relativity as an effective field theory: The leading quantum corrections*, Phys. Rev. D **50**, 3874 (1994), arXiv:gr-qc/9405057.
- [25] A. Accioly, B. L. Giacchini and I. L. Shapiro, *On the gravitational seesaw in higher-derivative gravity*, Eur. Phys. J. C **77**, 540 (2017), arXiv:1604.07348.
- [26] B. L. Giacchini, T. de Paula Netto and I. L. Shapiro, *On the Vilkovisky-DeWitt approach and renormalization group in effective quantum gravity*, J. High Energy Phys. **2020**, 011 (2020), arXiv:2009.04122.
- [27] N. D. Birrell and P. C. W. Davies, *Quantum fields in curved space* (Cambridge University Press, 1982).
- [28] I. L. Buchbinder, S. D. Odintsov and I. L. Shapiro, *Effective Action in Quantum Gravity* (IOPP, Bristol, 1992).
- [29] I. L. Shapiro, *Effective Action of Vacuum: Semiclassical Approach*, Class. Quant. Grav. **25**, 103001 (2008), arXiv:0801.0216.
- [30] M. J. Duff, *Twenty years of the Weyl anomaly*, Class. Quant. Grav. **11**, 1387 (1994), arXiv:hep-th/9308075.
- [31] R. J. Riegert, *A Nonlocal Action for the Trace Anomaly*, Phys. Lett. B **134**, 56 (1984).
- [32] E. S. Fradkin and A. A. Tseytlin, *Conformal Anomaly in Weyl Theory and Anomaly Free Superconformal Theories*, Phys. Lett. B **134**, 187 (1984).
- [33] M. Asorey, E. V. Gorbar and I. L. Shapiro, *Universality and ambiguities of the conformal anomaly*, Class. Quant. Grav. **21** 163 (2003), arXiv:hep-th/0307187.
- [34] A. A. Starobinsky, *A new type of isotropic cosmological models without singularity*, Phys. Lett. B **91**, 99 (1980).
- [35] A. A. Starobinsky, *The Perturbation Spectrum Evolving from a Nonsingular Initially De-Sitter Cosmology and the Microwave Background Anisotropy*, Sov. Astron. Lett. **9**, 302 (1983).
- [36] J. C. Fabris, A. M. Pelinson and I. L. Shapiro, *Anomaly induced effective action for gravity and inflation*, Grav. Cosmol. **6**, 59 (2000), arXiv:gr-qc/9810032.
- [37] J. C. Fabris, A. M. Pelinson and I. L. Shapiro, *On the gravitational waves on the background of anomaly-induced inflation*, Nucl. Phys. B **597**, 539 (2001), arXiv:hep-th/0009197.
- [38] A. M. Pelinson, I. L. Shapiro and F. I. Takakura, *On the stability of the anomaly induced inflation*, Nucl. Phys. B **648**, 417 (2003), arXiv:hep-ph/0208184.
- [39] T. de Paula Netto, A. M. Pelinson, I. L. Shapiro and A. A. Starobinsky, *From stable to unstable anomaly-induced inflation*, Eur. Phys. J. C **76**, 544 (2016), arXiv:1509.08882.
- [40] L. Modesto and I. L. Shapiro, *Superrenormalizable quantum gravity with complex ghosts*, Phys. Lett. B **755**, 279 (2016), arXiv:1512.07600.
- [41] L. Modesto, *Super-renormalizable or finite Lee-Wick quantum gravity*, Nucl. Phys. B **909**, 584 (2016), arXiv:1602.02421.
- [42] D. Anselmi and M. Piva, *A new formulation of Lee-Wick quantum field theory*, J. High Energy Phys. **06**, 066 (2017), arXiv:1703.04584.
- [43] D. Anselmi and M. Piva, *Perturbative unitarity of Lee-Wick quantum field theory*, Phys. Rev. D **96**, 045009 (2017), arXiv:1703.05563.
- [44] D. Anselmi, *Fakeons and Lee-Wick Models*, J. High Energy Phys. **02**, 141 (2018), arXiv:1801.00915.
- [45] D. Anselmi, *On the quantum field theory of the gravitational interactions*, J. High Energy Phys. **06**, 086 (2017), arXiv:1704.07728.
- [46] N. V. Krasnikov, *Nonlocal Gauge Theories*, Theor. Math. Phys. **73**, 1184 (1987) [Teor. Mat. Fiz. **73**, 235 (1987)].

- [47] Yu. V. Kuz'min, *Finite nonlocal gravity*, Sov. J. Nucl. Phys. **50**, 1011 (1989) [Yad. Fiz. **50**, 1630 (1989)].
- [48] E. T. Tomboulis, *Superrenormalizable gauge and gravitational theories*, arXiv:hep-th/9702146.
- [49] L. Modesto, *Super-renormalizable Quantum Gravity*, Phys. Rev. D **86**, 044005 (2012), arXiv:1107.2403.
- [50] T. D. Lee and G. C. Wick, *Negative metric and the unitarity of the S Matrix*, Nucl. Phys. B **9**, 209 (1969).
- [51] T. D. Lee and G. C. Wick, *Finite Theory of Quantum Electrodynamics*, Phys. Rev. D **2**, 1033 (1970).
- [52] R. E. Cutkosky, P. V. Landshoff, D. I. Olive and J. C. Polkinghorne, *A non-analytic S matrix*, Nucl. Phys. B **12**, 281 (1969).
- [53] L. Modesto, L. Rachwał and I. L. Shapiro, *Renormalization group in super-renormalizable quantum gravity*, Eur. Phys. J. C **78**, 555 (2018), arXiv:1704.03988.
- [54] L. Rachwał, L. Modesto, A. Pinzul and I. L. Shapiro, *Renormalization Group in Six-derivative Quantum Gravity*, arXiv:2104.13980.
- [55] L. Modesto and L. Rachwał, *Super-renormalizable and finite gravitational theories*, Nucl. Phys. B **889** 228 (2014), arXiv:1407.8036.
- [56] L. Modesto, M. Piva and L. Rachwał, *Finite quantum gauge theories*, Phys. Rev. D **94**, 025021 (2016), arXiv:1506.06227.
- [57] S. Deser and A. N. Redlich, *String Induced Gravity and Ghost Freedom*, Phys. Lett. B **176**, 350 (1986).
- [58] D. N. Page and A. A. Shoom, *Local Invariants Vanishing on Stationary Horizons: A Diagnostic for Locating Black Holes*, Phys. Rev. Lett. **114**, 141102 (2015), arXiv:1501.03510.
- [59] D. D. McNutt and D. N. Page, *Scalar Polynomial Curvature Invariant Vanishing on the Event Horizon of Any Black Hole Metric Conformal to a Static Spherical Metric*, Phys. Rev. D **95**, 084044 (2017), arXiv:1704.02461.
- [60] A. Coley and D. McNutt, *Identification of black hole horizons using scalar curvature invariants*, Class. Quant. Grav. **35**, 025013 (2018), arXiv:1710.08773.
- [61] D. D. McNutt, M. A. H. MacCallum, D. Gregoris, A. Forget, A. A. Coley, P. C. Chavy-Waddy and D. Brooks, *Cartan Invariants and Event Horizon Detection, Extended Version*, Gen. Rel. Grav. **50**, 37 (2018), arXiv:1709.03362.
- [62] V. P. Frolov, *Notes on nonsingular models of black holes*, Phys. Rev. D **94**, 104056 (2016), arXiv:1609.01758.
- [63] T. Berry, A. Simpson and M. Visser, *General class of "quantum deformed" regular black holes*, arXiv:2102.02471.
- [64] J. M. Bardeen, *Non-singular general-relativistic gravitational collapse*, in Proceedings of International Conference GR5, (Tbilisi University Press, Tbilisi, USSR, 1968), p. 174.
- [65] E. Ayón-Beato and A. García, *Regular black hole in general relativity coupled to nonlinear electrodynamics*, Phys. Rev. Lett. **80**, 5056 (1998), arXiv:gr-qc/9911046.
- [66] I. Dymnikova, *Regular electrically charged vacuum structures with de Sitter centre in nonlinear electrodynamics coupled to general relativity*, Class. Quant. Grav. **21**, 4417 (2004), arXiv:gr-qc/0407072.
- [67] P. Nicolini, A. Smailagic and E. Spallucci, *Noncommutative geometry inspired Schwarzschild black hole*, Phys. Lett. B **632**, 547 (2006), arXiv:gr-qc/0510112, P. Nicolini, *Noncommutative Black Holes, The Final Appeal To Quantum Gravity: A Review*, Int. J. Mod. Phys. A **24**, 1229 (2009), arXiv:0807.1939.
- [68] K. A. Bronnikov and J. C. Fabris, *Regular phantom black holes*, Phys. Rev. Lett. **96**, 251101 (2006), arXiv:gr-qc/0511109.
- [69] A. Simpson and M. Visser, *Black-bounce to traversable wormhole*, J. Cosmol. Astropart. Phys. **02**, 042 (2019), arXiv:1812.07114.
- [70] J. Boos, *Non-singular "Gauss" black hole from non-locality: a simple model with a de Sitter core, mass gap, and no inner horizon*, arXiv:2104.00555.
- [71] Y. Zhang, Y. Zhu, L. Modesto and C. Bambi, *Can static regular black holes form from gravitational collapse?*, Eur. Phys. J. C **75**, 96 (2015), arXiv:1404.4770.
- [72] A. A. Tseytlin, *On singularities of spherically symmetric backgrounds in string theory*, Phys. Lett. B **363**, 223 (1995), arXiv:hep-th/9509050.
- [73] V. P. Frolov, A. Zelnikov and T. de Paula Netto, *Spherical collapse of small masses in the ghost-free gravity*, J. High Energy Phys. **1506**, 107 (2015), arXiv:1504.00412.
- [74] V. P. Frolov, *Mass-gap for black hole formation in higher derivative and ghost free gravity*, Phys. Rev. Lett. **115**, 051102 (2015), arXiv:1505.00492.
- [75] V. P. Frolov and A. Zelnikov, *Head-on collision of ultrarelativistic particles in ghost-free theories of gravity*, Phys. Rev. D **93**, 064048 (2016), arXiv:1509.03336.
- [76] L. Buoninfante, A. S. Koshelev, G. Lambiase, J. Marto and A. Mazumdar, *Conformally-flat, non-singular static metric in infinite derivative gravity*, J. Cosmol. Astropart. Phys. **06**, 014 (2018), arXiv:1804.08195.
- [77] A. Bonanno and M. Reuter, *Renormalization group improved black hole space-times*, Phys. Rev. D **62**, 043008 (2000), arXiv:hep-th/0002196.
- [78] S. A. Hayward, *Formation and evaporation of regular black holes*, Phys. Rev. Lett. **96**, 031103 (2006), arXiv:gr-qc/0506126.
- [79] I. Dymnikova, *Vacuum nonsingular black hole*, Gen. Rel. Grav. **24**, 235 (1992).
- [80] T. De Lorenzo, C. Pacilio, C. Rovelli and S. Speziale, *On the Effective Metric of a Planck Star*, Gen. Rel. Grav. **47**, 41 (2015), arXiv:1412.6015.
- [81] E. Ayón-Beato and A. García, *Nonsingular charged black hole solution for nonlinear source*, Gen. Rel. Grav. **31**, 629 (1999), arXiv:gr-qc/9911084.
- [82] E. Ayón-Beato and A. García, *New regular black hole solution from nonlinear electrodynamics*, Phys. Lett. B **464**, 25 (1999), arXiv:hep-th/9911174.
- [83] K. A. Bronnikov, *Regular magnetic black holes and monopoles from nonlinear electrodynamics*, Phys. Rev. D **63**, 044005 (2001), arXiv:gr-qc/0006014.
- [84] W. Berej, J. Matyjasek, D. Tryniecki and M. Woronowicz, *Regular black holes in quadratic gravity*, Gen. Rel. Grav. **38**, 885 (2006), arXiv:hep-th/0606185.
- [85] L. Balart and E. C. Vagenas, *Regular black holes with a nonlinear electrodynamics source*, Phys. Rev. D **90**, 124045 (2014), arXiv:1408.0306.
- [86] Wolfram Research, Inc., MATHEMATICA, Version 8, <https://www.wolfram.com/mathematica/>.
- [87] J. M. Martín-García, *xAct: Efficient tensor computer algebra for the Wolfram Language*, <http://xact.es>, accessed: 2021-01-27.
- [88] D. Yllanes and J. M. Martín-García, *xCoba: General component tensor computer algebra*, <http://xact.es/xCoba/>, accessed: 2021-01-27.

- [89] M. J. Duff, *Quantum corrections to the Schwarzschild solution*, Phys. Rev. D **9**, 1837 (1974).
- [90] J. F. Donoghue, *Leading quantum correction to the Newtonian potential*, Phys. Rev. Lett. **72**, 2996 (1994), arXiv:gr-qc/9310024; *General relativity as an effective field theory: The leading quantum corrections*, Phys. Rev. D **50**, 3874 (1994), arXiv:gr-qc/9405057.
- [91] N. Burzillà, B. L. Giacchini, T. de Paula Netto and L. Modesto, *Newtonian potential in higher-derivative quantum gravity*, Phys. Rev. D **103**, 064080 (2021), arXiv:2012.06254.