

MEGADOCK-Web-Mito: human mitochondrial protein–protein interaction prediction database

Masahito Ohue¹, Hiroki Watanabe^{2,3}, and Yutaka Akiyama²

¹ Department of Computer Science, School of Computing, Tokyo Institute of Technology, G3-56, 4259 Nagatsutacho, Midori-ku, Yokohama City, Kanagawa 226-8503, Japan

ohue@c.titech.ac.jp

² Department of Computer Science, School of Computing, Tokyo Institute of Technology, W8-76, 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8550, Japan

h.watanabe@bi.c.titech.ac.jp, akiyama@c.titech.ac.jp

³ RWBC-OIL, National Institute of Advanced Industrial Science and Technology, 1-1-1 Umezono, Tsukuba, Ibaraki 305-8560, Japan

Abstract. Mitochondrial diseases are largely caused by dysfunction in mitochondrial proteins. However, annotations of human mitochondrial proteins are scattered across various public databases and individual studies. To facilitate research aimed at elucidating mitochondrial functions, we constructed the MEGADOCK-Web-Mito database as a protein–protein interaction (PPI) prediction data archive, including prediction results for exhaustive protein pairs of 654 mitochondria-related human proteins. MEGADOCK-Web-Mito enables users to search for all PPI prediction results efficiently and comprehensively. In particular, we linked functional annotations to each human mitochondrial protein. The comprehensive and specialized human mitochondrial PPI prediction results and searching function of MEGADOCK-Web-Mito will support further research on mitochondria and mitochondrial diseases.

Keywords: protein–protein interaction (PPI), mitochondrial protein, MEGADOCK, MEGADOCK-Web

1 Introduction

Mitochondria are organelles of eukaryotic cells that serve as the primary energy production sites. Mitochondria are also involved in various other essential cellular processes, including the production of reactive oxygen species, apoptosis, calcium ion storage, and infection defense. In recent years, there has been increased research attention on the range of mitochondrial functions, especially in the context of mitochondrial diseases [1]. Toward this end, it is essential to elucidate the function of mitochondria-associated proteins.

Many proteins in living organisms interact with each other [2], and such protein–protein interactions (PPIs) play a central role in biological phenomena. PPIs themselves, or their dysregulation, have been associated with certain diseases, and drugs have recently been developed to inhibit pathological PPIs [3,4].

Along with progress in molecular technologies, enabling the discovery of approximately 196,111 human proteins [based on Swiss-Prot and TrEMBL entries in Universal Protein Resource (UniProt) [5] as of April 15, 2021], comes the challenge of identifying the large amount of potential PPIs exhaustively in biochemical experiments because of the extensive time and costs involved in such a process. Therefore, computational prediction of PPIs is desirable, which can narrow down the candidate proteins that are most likely to interact with each other prior to experimentation.

To address this challenge, we have developed MEGADOCK [6,7], a structure-based PPI prediction method. Although there are various methods for PPI prediction, the structure-based method has the advantage of discovering novel PPIs that do not depend on known interaction information. Moreover, since the three-dimensional structures of approximately 40% of mitochondria-localized proteins have been resolved and reported, structure-based prediction by MEGADOCK is particularly useful for mitochondria-related proteins. We previously reported development of the MEGADOCK-Web database, which is an archive of PPI predictions made by MEGADOCK [8]. This database implements a search and display function for information on predicted PPIs precomputed by MEGADOCK, and further allows users to browse comprehensive PPI prediction results for approximately 2,000 representative human proteins, including predicted complex structural models.

However, MEGADOCK-Web is a database for general human PPI prediction results, and does not have a specific function specifically related to the mitochondria. For example, since one of the major causes of mitochondrial diseases is mutation or abnormality of genes with sequence information of mitochondria-related proteins, it is important to be able to refer to mitochondrial-specific gene information. Therefore, to elucidate the biological functions of mitochondria and the causes of mitochondrial diseases, it is necessary to have a database specialized in mitochondrial PPIs, including information on genes with mitochondrial protein sequence information.

In this paper, we report the results of comprehensive PPI prediction using MEGADOCK for human proteins localized in the mitochondria, including proteins encoded in both the mitochondrial and nuclear genomes. We further introduce our newly developed MEGADOCK-Web-Mito database, which integrates mitochondria-related PPI prediction results and mitochondria-related protein information.

2 Related Work

2.1 MitoProteome

The MitoProteome is an object-related database developed at the UCSD Supercomputer Center, which contains information on mitochondria-localized proteins that are primarily encoded in the human nuclear genome [9,10]. Each entry in the MitoProteome corresponds to a gene encoding a protein that is localized in




Fig. 1. Page transition diagram of MEGADOCK-Web.

the mitochondria and its basic information, along with annotations of isoforms, splice variants, and functions of the corresponding protein.

2.2 MEGADOCK-Web

MEGADOCK-Web [8] is a PPI prediction results database and browsing system. MEGADOCK-Web contains the results of comprehensive PPI predictions performed by MEGADOCK for representative human proteins.

Fig. 1 shows the structure of the MEGADOCK-Web webpage. The PPI score is a value calculated from the results of PPI prediction calculations by MEGADOCK. Normally, a certain threshold is set for the PPI score, and protein pairs with PPI scores higher than the threshold are predicted to interact.

There are three main functions of MEGADOCK-Web: (1) searching for PPIs predicted by MEGADOCK, (2) visualizing the predicted interaction partners on the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway map [11], and (3) visualizing protein complex models by Molmil viewer [12].

3 Implementation

In this study, we developed MEGADOCK-Web-Mito, a database containing predicted results of mitochondria-related PPIs based on the MEGADOCK-Web framework.

3.1 Data Resources

The proteins targeted by MEGADOCK-Web-Mito were those that satisfied certain criteria by querying the Protein Data Bank (PDB) [13] via UniProt [5] among all proteins registered in the MitoProteome. The details of each database are provided below.

MitoProteome We defined mitochondria-associated proteins as those contained in the MitoProteome [9,10]. The MitoProteome assigns Mito IDs, represented by “MT” and six digits such as “MT000001,” to mitochondria-associated protein sequences, and provides a correspondence between the Mito ID and UniProt AC, which is the general ID of the protein sequence.

PDB PDB [13] is a database of the three-dimensional structures of molecules such as proteins and nucleic acids. Each PDB structure is assigned a four-letter identifier (PDB ID), which may be provided in the form of a complex of multiple molecules such as proteins. A chain ID was assigned to each molecular chain in the PDB ID structure. For example, the protein structure 1A4I contains two chains, chain A and chain B; therefore, the molecular chain structure is presented as 1A4I_A or 1A4I_B.

UniProt UniProt [5] is a database that contains protein sequence data and annotation information related to the functions. UniProt assigns an accession number (UniProt AC) to each protein sequence, and RESTful application programming interfaces (APIs) can be used to retrieve data from UniProt in text, XML, and other formats.

3.2 Data Collection

The results of PPI prediction to be included in MEGADOCK-Web-Mito were obtained using the chain structure as input. Considering that there are multiple possible structures for a given protein, even with an identical sequence, the chain structures of the proteins in the MitoProteome were obtained to satisfy the following conditions:

1. The chain structure included in MEGADOCK-Web-Mito is one of the possible structures of the corresponding mitochondria-associated protein.
2. If there are multiple PDB IDs with the same UniProt AC, MEGADOCK-Web-Mito will include the PDB ID with the highest number of residues resolved. If the number of residues is the same, the structure determined by X-ray crystallography is selected. If there are multiple structures determined by X-ray crystallography, that with the best resolution is selected.

Based on these criteria, a total of 1,675 chain structures were obtained, which were classified into 654 protein types, corresponding to the protein sequence (UniProt AC).

3.3 PPI Prediction

The obtained 1,675 chain structures were used for PPI prediction calculations by running MEGADOCK on the TSUBAME 3.0 supercomputer owned by the Tokyo Institute of Technology. MEGADOCK calculations were performed on $1,675 \times 1,675 = 2,805,625$ chain structure pairs.

Suppose the MEGADOCK computation is performed on one `f_node` with TSUBAME 3.0 [two Intel Xeon E5-2680 V4 processors (14 cores, 2.4 GHz) and four GPUs (NVIDIA TESLA P100 for NVlink-Optimized)], it takes approximately 10 seconds per chain structure pair. An exhaustive PPI prediction calculation using all 1,675 types of chain structures requires approximately 320 node-days. Therefore, we developed a script that uses multiple nodes of TSUBAME 3.0 simultaneously, and achieved exhaustive PPI prediction calculation in only 4.3 days.

3.4 Link to Mito ID and Entrez Gene ID

MEGADOCK-Web contains information on proteins, PPI prediction results, and KEGG pathways. MEGADOCK-Web-Mito contains this same information, in addition to the Mito ID and Entrez Gene ID. Each ID links to MitoProteome and Entrez Gene [14] entries to facilitate access to detailed protein information. In addition, a search function according to Mito ID and Entrez Gene ID was implemented.

3.5 System Architecture

As for MEGADOCK-Web, we developed MEGADOCK-Web-Mito database using Play Framework [15], which is a Java and Scala web application framework that includes components and APIs necessary for web application development. MEGADOCK-Web-Mito adopts a client-server model in the same way as a general web application. Requests from clients are classified into requests to static resources (such as protein PDB files and docking result files) and requests to the web pages. In the former case, Apache directly returns the corresponding resources to clients. In the latter case, Apache redirects requests to the web application running on the local port. We stored information in the relational H2 Database Engine [16] database with MySQL [17] wrapped around Play Framework for server-side implementation. Receiving requests from users invokes the web application, which generates HTML dynamically to refer to the database. Server-side implementation is multi-threaded to respond immediately. JavaScript and jQuery are adopted as client-side implementation tools to control the web pages. The web server was built on-premises and can be accessed at the following URL: <http://www.bi.cs.titech.ac.jp/megadock-web-mito>.


4 Database Features and Applications

4.1 Protein Statistics

The 654 proteins obtained were classified based on their EC number (enzyme number) [18], which are summarized in Table 1.

Table 1. Breakdown of the 654 mitochondria-related proteins in MEGADOCK-Web-Mito.

class	#protein
Oxidoreductases	97
Transferases	116
Hydrolases	94
Lyases	17
Isomerases	12
Ligases	28
Non-enzymes	290

**Fig. 2.** Page transition diagram of MEGADOCK-Web-Mito (pages marked in orange are those that have been modified or extended from MEGADOCK-Web).

4.2 Database Structure

The page structure of MEGADOCK-Web-Mito is shown in Fig. 2, and each page is described in further detail below.

– Top page

The top page of MEGADOCK-Web-Mito has four text fields as search windows, as found in MEGADOCK-Web. As a unique feature of MEGADOCK-Web-Mito, Mito IDs and Entrez Gene IDs have been added to the pull-down menu items placed to the left of each text field, allowing protein searches by Mito ID and Entrez Gene ID, respectively.

– Protein pair selection page and Protein selection page

In MEGADOCK-Web-Mito, the columns for “Mito ID” and “Entrez Gene ID” attributes were added to the table of the protein list of search hits. In addition, Mito ID and Entrez Gene IDs were linked to MitoProteome and Entrez Gene, respectively, to enable convenient access to additional information (see Fig. 3).

– PPI prediction list page

In the PPI prediction list page of MEGADOCK-Web-Mito, we added a row for the “Mito ID” and “Entrez Gene ID” attributes to the table of protein information and PPI scores (see Fig. 4), which can be searched and linked to

the MitoProteome and Entrez Gene ID, respectively. In addition, we added a line that displays a list of KEGG pathways that contain the protein to be searched, enabling the visualization of KEGG pathways for individual proteins.

4.3 Use Case

This section describes a specific example of the use of MEGADOCK-Web-Mito to retrieve the PPI prediction information for the protein with Mito ID MT000972 in the MitoProteome, which is related to Parkinson’s disease. The following is an example of using MEGADOCK-Web-Mito to retrieve PPI prediction information for the protein with the Mito ID ‘MT000972.’

1. A search for the protein with Mito ID MT000972 in MEGADOCK-Web-Mito revealed the following (Fig. 5): the gene symbol is HTRA2, and there are two chain structures: 2PZD_A and 5FHT_A (Fig. 5(a)). Referring to hsa05012 Parkinson disease from KEGG pathway information (Fig. 5(b)), there is a protein named DJ-1 in the vicinity of HTRA2 (Fig. 5(c)).
2. Searching for the protein DJ-1 in MEGADOCK-Web-Mito, we found the associated gene symbol PARK7, with Mito ID MT000850. There are four PDB structures (1PDW_A, 1PE0_A, 2RK3_A, 3BWE_A) for this protein.
3. One of the PPI prediction results for HTRA2 and DJ-1 is shown in Fig. 6, demonstrating the interaction for the structures 5FHT_A and 3BWE_A.

MEGADOCK-Web-Mito

There are 310 matches for the query "mitochondria". Please click "View" to go to your protein page.

UniProt AC	Organism	Protein Names	Gene Names	Mito ID	Entrez Gene ID	PDB ID	Chain	
Q02127	Homo sapiens (Human)	Dihydrorotate dehydrogenase (quinone), mitochondrial	DHODH	MT000182	1723	1D3H	A	View
Q13423	Homo sapiens (Human)	NAD(P) transhydrogenase, mitochondrial	NNT	MT000894	23530	1DJL	A	View
P23368	Homo sapiens (Human)	NAD-dependent malic enzyme, mitochondrial	ME2	MT000352	4200	1EFK	A	View
O00746	Homo sapiens (Human)	Nucleoside diphosphate kinase, mitochondrial	NME4	MT000434	4833	1EHW	A	View
P00480	Homo sapiens (Human)	Ornithine carbamoyltransferase, mitochondrial	OTC	MT000450	5009	1FVO	A	View
P00390	Homo sapiens (Human)	Glutathione reductase, mitochondrial	GSR	MT000271	2936	1GRG	A	View
P30044	Homo sapiens (Human)	Peroxiredoxin-5, mitochondrial	PRDX5	MT000915	25824	1H4O	A	View
Q13825	Homo sapiens (Human)	Methylglutaconyl-CoA hydratase, mitochondrial	AUH	MT000066	549	1HZD	A	View
P49675	Homo sapiens (Human)	Steroidogenic acute regulatory protein, mitochondrial	STAR	MT000581	6770	1IMG	A	View
P26440	Homo sapiens (Human)	Isovaleryl-CoA dehydrogenase, mitochondrial	IVD	MT000324	3712	1IVH	A	View

◀ 1 2 3 ▶

Fig. 3. Example of a protein selection page for MEGADOCK-Web-Mito.

Protein Information

Your Query: PCCA

Protein Names	Propionyl-CoA carboxylase alpha chain; mitochondrial
Gene Names	PCCA
PDB ID	2CQY (Chain A)
UniProt AC	P05165
Mito ID	MT000458
Entrez Gene ID	5095
KEGG Pathway	<ul style="list-style-type: none"> • hsa00640 Propanoate metabolism - Homo sapiens (human) • hsa01100 Metabolic pathways - Homo sapiens (human) • hsa00630 Glyoxylate and dicarboxylate metabolism - Homo sapiens (human) • hsa00280 Valine, leucine and isoleucine degradation - Homo sapiens (human) • hsa01200 Carbon metabolism - Homo sapiens (human)

Predicted PPI Information

Show KEGG pathways in candidates

PPIScore > [Show Pathway](#)

UniProt AC	Organism	Protein Names	Gene Names	Mito ID	Entrez Gene ID	PDB ID	PPI Score	View	Databases
Q06124	Homo sapiens (Human)	Tyrosine-protein phosphatase non-receptor type 11	PTPN11	MT000519	5781	4H1O (Chain A)	11.7992	View	
P04637	Homo sapiens (Human)	Cellular tumor antigen p53	TP53	MT000600	7157	5ECG (Chain A)	11.3691	View	
O75874	Homo sapiens (Human)	Isocitrate dehydrogenase [NADP] cytoplasmic	IDH1	MT000311	3417	3INM (Chain A)	10.5039	View	

Fig. 4. Example PPI prediction list page for MEGADOCK-Web-Mito.

(a) MEGADOCK-Web-Mito

There are 2 matches for the query "MT000972". Please click "View" to go to your protein page.

UniProt AC	Organism	Protein Names	Gene Names	Mito ID	Entrez Gene ID	PDB ID	Chain	
O43464	Homo sapiens (Human)	Serine protease HTRA2; mitochondrial	HTRA2	MT000972	27429	2PZD	A	View
O43464	Homo sapiens (Human)	Serine protease HTRA2; mitochondrial	HTRA2	MT000972	27429	5FHT	A	View

(b) Protein Information

Your Query: HTRA2

Protein Names	Serine protease HTRA2; mitochondrial
Gene Names	HTRA2
PDB ID	2PZD (Chain A)
UniProt AC	O43464
Mito ID	MT000972
Entrez Gene ID	27429
KEGG Pathway	<ul style="list-style-type: none"> • hsa04210 Apoptosis - Homo sapiens (human) • hsa04215 Apoptosis - multiple species - Homo sapiens (human) • hsa05012 Parkinson disease - Homo sapiens (human)




Fig. 5. (a) Protein selection page of the result of searching for “MT000972”. (b) Detailed page of protein with Mito ID MT000972. (c) KEGG pathway map of the Parkinson’s disease pathway extracted from the PPI prediction results of HTRA2, and the predicted interaction partners of HTRA2.

Docking Information
 PPIScore: 8.473203 [Show KEGG Pathway](#)

Protein	Protein Names	Gene Names	PDB ID
Protein1	Serine protease HTRA2; mitochondrial	HTRA2	5FHT (Chain A)
Protein2	Protein/nucleic acid deglycase DU-1	PARK7	3BWE (Chain A)

Docking results from rank 1 to 10
[Download complex in this page](#)
[Download docking result file](#)

Rank	Score	Model	Download
1	4908.32		5FHT_A-3BWE_A.decoy1.pdb
2	4872.64		5FHT_A-3BWE_A.decoy2.pdb

Fig. 6. List of predicted complex structures of 5FHT_A and 3BWE_A.

5 Conclusions

In this study, we developed MEGADOCK-Web-Mito, a PPI prediction database related to the mitochondria, based on the existing MEGADOCK-Web framework. The PPIs of 654 human mitochondrial proteins with a total of 1,675 protein structures were comprehensively predicted, and the database contains all relevant predictions. In addition, we integrated the annotation information of proteins related to human mitochondria to enable an efficient and detailed investigation of the predicted PPI results. Since MEGADOCK-Web-Mito has comprehensive PPI prediction results for mitochondria-related proteins and a

search function specific to human mitochondria, it is expected to contribute to the study of the molecular mechanisms underlying diseases, particularly mitochondrial diseases.

Although all of the proteins in MEGADOCK-Web-Mito are localized in the human mitochondria, it is also necessary to include proteins localized in the mouse and rat mitochondria, which have similar three-dimensional structures to human proteins, which would further improve PPI prediction results. This will be a challenge for future research.

Acknowledgements

This work was partially supported by Japan Society for the Promotion of Science (JSPS) KAKENHI grants (18K18149, 20H04280), Platform Project for Supporting Drug Discovery and Life Science Research [Basis for Supporting Innovative Drug Discovery and Life Science Research (BINDS) grant no. JP20am0101112] from the Japan Agency for Medical Research and Development (AMED). This work was partially conducted as part of the research activities of AIST-Tokyo Tech Real World Big-Data Computation Open Innovation Laboratory (RWBC-OIL). The authors thank Editage (www.editage.com) for English language editing.

References

1. Schapira, A. H. V.: Mitochondrial diseases. *Lancet* 379, 1825–1834 (2012).
2. de Las Rivas, J., Fontanillo, C.: Protein-protein interactions essentials: Key concepts to building and analyzing interactome networks. *PLoS Comput. Biol.* 6, e1000807 (2010).
3. Oltersdorf, T., Elmore, S.W., Shoemaker, A. R., Armstrong, R.C., Augeri, D.J., Belli, B.A., et al.: An inhibitor of Bcl-2 family proteins induces regression of solid tumours. *Nature* 435, 677–681 (2005).
4. Popowicz, G.M., Czarna, A., Wolf, S., Wang, K., Wang, W., Dömling, A., Holak, T.A.: Structures of low molecular weight inhibitors bound to MDMX and MDM2 reveal new approaches for p53-MDMX/MDM2 antagonist drug discovery. *Cell Cycle*. 9, 1104–1111 (2010).
5. The UniProt Consortium. UniProt: The universal protein knowledgebase in 2021. *Nucleic Acids Res.* 49, D480–D489 (2021).
6. Ohue, M., Shimoda, T., Suzuki, S., Matsuzaki, Y., Ishida, T., Akiyama, Y.: MEGADOCK 4.0: an ultra-high-performance protein-protein docking software for heterogeneous supercomputers. *Bioinformatics* 30, 3281–3283 (2014).
7. Ohue, M., Matsuzaki, Y., Uchikoga, N., Ishida, T., Akiyama, Y.: MEGADOCK: An all-to-all protein-protein interaction prediction system using tertiary structure data. *Protein Pept. Lett.* 21, 766–778 (2014). doi:10.2174/09298665113209990050
8. Hayashi, T., Matsuzaki, Y., Yanagisawa, K., Ohue, M., Akiyama, Y.: MEGADOCK-Web: an integrated database of high-throughput structure-based protein-protein interaction predictions. *BMC Bioinform.* 19, 62 (2018).

9. Cotter, D., Guda, P., Fahy, E., Subramaniam, S.: MitoProteome: mitochondrial protein sequence database and annotation system. *Nucleic Acids Res.* 32, D463–D467 (2004).
10. Guda, P., Subramaniam, S., Guda, C.: MitoProteome: human heart mitochondrial protein sequence database. *Methods Mol. Biol.* 357, 375–383 (2007).
11. Kanehisa, M., Furumichi, M., Sato, Y., Ishiguro-Watanabe, M., Tanabe, M.: KEGG: integrating viruses and cellular organisms. *Nucleic Acids Res.* 49, D545–D551 (2021).
12. Bekker, G.J., Nakamura, H., Kinjo, A.R.: Molmil: a molecular viewer for the PDB and beyond. *J. Cheminform.* 8, 42 (2016).
13. Burley, S.K., Bhikadiya, C., Bi, C., Bittrich, S., Chen, L., Crichlow, G.V., Zhuravleva, M.: RCSB Protein Data Bank: powerful new tools for exploring 3D structures of biological macromolecules for basic and applied research and education in fundamental biology, biomedicine, biotechnology, bioengineering, and energy sciences. *Nucleic Acids Res.* 49, D437–D451 (2021).
14. Maglott, D., Ostell, J., Pruitt, K.D., Tatusova, T.: Entrez Gene: gene-centered information at NCBI. *Nucleic Acids Res.* 39, D52–D57 (2010).
15. Play Framework. <https://www.playframework.com/>. Accessed 19 April 2021.
16. H2 Database Engine. <http://www.h2database.com/>. Accessed 19 April 2021.
17. MySQL. <https://www.mysql.com/>. Accessed 19 April 2021.
18. International Union of Biochemistry and Molecular Biology. <https://iubmb.or>. Accessed 19 April 2021.