

# Ladder Symmetries of Black Holes: Implications for Love Numbers and No-Hair Theorems

Lam Hui,<sup>a\*</sup> Austin Joyce,<sup>b†</sup> Riccardo Penco,<sup>c‡</sup>  
Luca Santoni,<sup>a§</sup> and Adam R. Solomon<sup>c¶</sup>

<sup>a</sup>*Center for Theoretical Physics, Department of Physics,  
Columbia University, New York, NY 10027*

<sup>b</sup>*Delta-Institute for Theoretical Physics,  
University of Amsterdam, Amsterdam, 1098 XH, The Netherlands*

<sup>c</sup>*McWilliams Center for Cosmology, Department of Physics,  
Carnegie Mellon University, Pittsburgh, PA 15213*

## Abstract

It is well known that asymptotically flat black holes in general relativity have a vanishing static tidal response and no static hair. We show that both are a result of linearly realized symmetries governing static (spin 0,1,2) perturbations around black holes. The symmetries have a geometric origin: in the scalar case, they arise from the (E)AdS isometries of a dimensionally reduced black hole spacetime. Underlying the symmetries is a ladder structure which can be used to construct the full tower of solutions, and derive their general properties: (1) solutions that decay with radius spontaneously break the symmetries, and must diverge at the horizon; (2) solutions regular at the horizon respect the symmetries, and take the form of a finite polynomial that grows with radius. Property (1) implies no hair; (1) and (2) combined imply that static response coefficients—and in particular Love numbers—vanish. We also discuss the manifestation of these symmetries in the effective point particle description of a black hole, showing explicitly that for scalar probes the worldline couplings associated with a non-trivial tidal response and scalar hair must vanish in order for the symmetries to be preserved.

---

\*lh399@columbia.edu

†a.p.joyce@uva.nl

‡rpenco@andrew.cmu.edu

§luca.santoni@columbia.edu

¶adamsolo@andrew.cmu.edu

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>Ladder in Schwarzschild</b>	<b>6</b>
<b>3</b>	<b>Ladder in Kerr</b>	<b>10</b>
<b>4</b>	<b>Ladder in Spin: From Scalar to Vector and Tensor</b>	<b>13</b>
<b>5</b>	<b>Symmetries in the Infrared</b>	<b>15</b>
<b>6</b>	<b>Discussion</b>	<b>18</b>
<b>A</b>	<b>Geometric Origin of the Ladder Symmetries</b>	<b>20</b>
<b>B</b>	<b>Pöschl–Teller Potential</b>	<b>23</b>
<b>C</b>	<b>Ladder in Regge–Wheeler and Zerilli</b>	<b>24</b>
<b>D</b>	<b>Ladder in Teukolsky</b>	<b>26</b>
<b>E</b>	<b>Supersymmetric Structure</b>	<b>28</b>
<b>F</b>	<b>Hypergeometric Identities</b>	<b>29</b>
<b>G</b>	<b>Scalar-Gauss–Bonnet Theories</b>	<b>30</b>
	<b>References</b>	<b>32</b>

# 1 Introduction

Chandrasekhar famously wrote [1]: “the black holes of nature are the most perfect macroscopic objects there are in the universe: the only elements in their construction are our concepts of space and time.” Indeed, the nonlinear vacuum solutions of the Einstein equation, first discovered a century ago, are elegant in their simplicity, yet deeply mysterious. Students of black holes learn to cajole them to give up their secrets by thought experiments: we throw objects at them and see what comes out; we perturb them and see how they respond. Of the many remarkable properties of black holes, we wish to focus on two: the vanishing of the so called Love numbers, and the absence of hair.

Consider a black hole in a tidal environment, which could take the form of an external gravitational field generated by other objects, but could also be an external scalar or vector field. The situation is analogous to placing a dielectric under the influence of an external electric field. The charges inside the material get displaced, setting up a multipole moment which sources an observable response field. For black holes, though, such a polarizability, or tidal response, appears to be absent. Let us be more precise: under the influence of a weak static tidal field (spin 0, 1, or 2), black holes have vanishing Love numbers [2–18]. The Love numbers quantify the tidal response: suppose the external tidal field goes as  $r^\ell$  at large distance  $r$  ( $\ell$  is the multipole of interest), an object such as a star that gets tidally deformed would source a response field that goes as  $1/r^{\ell+1}$  far away.<sup>1</sup> The coefficient of this  $1/r^{\ell+1}$  tail, normalized by the amplitude of the tidal field, is the Love number, or rather Love numbers (or Love tensors even), for there are in general different types of tidal fields one could turn on.<sup>2</sup>

For a weak static tidal field, the problem reduces to that of studying static linear perturbations around the object of interest, be it a star or a black hole. At a large distance from the object, the perturbation, suitably defined (let’s call it  $\phi$ ), obeys:

$$\nabla^2\phi = 0, \tag{1.1}$$

giving two branches of solutions, growing  $r^\ell$  and decaying  $1/r^{\ell+1}$  at multipole order  $\ell$ . For a star, the growing (tidal) solution is generally accompanied by the decaying tail, signifying deformability. For a black hole, the growing solution is regular at the horizon while the decaying one is not (and therefore discarded). These two facts combined tell us its Love numbers vanish.<sup>3</sup>

The second fact, that the decaying solution diverges at the horizon, ought to remind us of no-hair theorems. No-hair theorems tell us we would inevitably fail if we attempt to endow black holes with extra features, in other words, hair that is static and falls off with distance. Known no-hair theorems are stronger than the version we focus on in this paper: they apply even when some nonlinear interactions are present,

---

<sup>1</sup>The precise exponent depends on the choice of field one focuses on. For instance, for the gravitational field, while the metric fluctuations have  $r^\ell$  and  $1/r^{\ell+1}$  asymptotics at large  $r$ , the Weyl tensor asymptotes to  $r^{\ell-2}$  and  $1/r^{\ell+3}$ .

<sup>2</sup>There is some discussion in the literature on exactly how to define Love numbers [3, 5, 13, 14, 19]. Here, we follow the definition laid out in [8, 16–18, 20]. The Love numbers are identified with the coefficients of operators in the point-particle action involving the square of the tidal field (in the case of spin 2, it is the square of the Weyl tensor, suitably contracted). As such, the definition is explicitly gauge invariant, and solutions to the Teukolsky equation can be related directly to the Love numbers. We rely in particular on the matching computations in [17, 18]. This definition differentiates non-dissipative response (what we are interested in) from the dissipative one, relevant especially for rotating black holes [13–16, 18].

<sup>3</sup>More details about the growing solution are needed, in particular that as a power series in  $r$ , only certain powers appear, as will be explained below.

and/or when the metric is not a solution to the Einstein equation (but crucially must have a horizon) [21–30].<sup>4</sup> Nonetheless, our limited version—for linear, scalar/vector/tensor hair around a Schwarzschild or Kerr black hole—is a manifestation of these theorems, and it connects naturally with the phenomenon of tidal response.

Both characteristics, the lack of hair and tidal response, are well known of course. Our main task is not to derive them, although we will. Rather, we hope to peel back the mystery a little, and ask: why? What is the symmetry underpinning of these phenomena? We will work out the symmetries governing static, linear perturbations of general spin around Schwarzschild or Kerr black holes. Underlying them is a ladder structure that connects solutions at different  $\ell$ 's. The symmetries give rise to conserved quantities which can be used to connect asymptotic behaviors at the horizon and at infinity. Interestingly, the growing branch of solutions linearly realizes the symmetries while the decaying branch does not. This gives us a richer perspective on the choice of one branch over another: the boundary conditions that differentiate between a star and a black hole are seen to pick out states that make or break the ladder symmetries. In the scalar case at least, the symmetries have an identifiable geometric origin: they follow from the isometries of an Euclidean anti-de Sitter space of a suitably rescaled, dimensionally reduced black hole spacetime. Along the way, we will uncover a second ladder structure, one that connects perturbations of different spin.

The symmetry understanding is interesting from an infrared perspective. A far away observer, if they detect a  $\phi \sim 1/r^{\ell+1}$  profile, would conclude that the object of interest has some sort of (multipolar) charge that sources such a profile. That is, they would model it by adding a source term to eq. (1.1), localized on the object and proportional to the charge.<sup>5</sup> A black hole cannot have such a charge, or hair. As we will see, the ladder symmetries have a non-trivial large distance limit, telling us that a charge-less object like the black hole is an object with enhanced symmetries. Likewise, if the far away observer finds a  $1/r^{\ell+1}$  tail when an  $r^\ell$  tidal field is applied, they would conclude the object of interest has a non-trivial static response. That is, they would model it by adding another source term to eq. (1.1), localized at the position of the object and proportional to the tidal field. The proportionality constant is the Love number. In the effective worldline description of the system, these couplings break the infrared manifestations of the symmetries, so that the point in parameter space where these couplings vanish (as is appropriate to model a black hole) is an enhanced symmetry point.

Two papers are particularly relevant to our investigation. One is an interesting recent paper by Charalambous, Dubovsky, and Ivanov [43]. They pointed out a different set of symmetries that are also relevant for the vanishing of the black hole Love numbers. Their symmetry transformation involves time, while ours is purely static. The connection between the two is an interesting question which we hope to explore in the future. Another influential paper is by Compton and Morrison [44]. The ladder structure we find for black holes mirrors the one they discovered for de Sitter space. As we will explain, there is a simple reason: at least for the scalar case, the relevant metric, suitably reduced and rescaled, can be analytically continued to de Sitter. Our strategy for identifying conserved charges follows theirs.

---

<sup>4</sup>We would be remiss not to mention exceptions: they generally involve hair that is not static, or does not fall off at infinity, or is non-minimally coupled to gravity, or has special interactions [31–42].

<sup>5</sup>The approach of modeling distant behavior by adding terms to the point particle action is a standard effective field theory technique (see e.g., [20]).

The discussion is organized as follows. We start with the simplest case in Section 2: a massless scalar in a Schwarzschild background. All the essential features that govern higher spin perturbations in a Kerr background can be found in this case. Section 3 covers a scalar in the Kerr spacetime, and Section 4 extends the argument to vector and tensor perturbations. In Section 5, we discuss the infrared limit of the ladder symmetries. We conclude in Section 6, commenting on the extension to finite frequency and general number of dimensions. The Appendices contain a number of useful results: Appendix A on the geometric origin of the ladder symmetries, B on the Pöschl–Teller potential and its relation to our problem, C and D on the ladder structures in the Regge–Wheeler and Teukolsky equations respectively, E on the supersymmetric structure of the ladder symmetries, F on certain hypergeometric identities helpful for revealing the ladders, and G on black hole hair and tidal response for a scalar coupled to the Gauss–Bonnet curvature invariant.

## 2 Ladder in Schwarzschild

The simplest manifestation of the ladder symmetry structure we wish to describe is in the equation for a (massless) scalar propagating in a fixed Schwarzschild background. The Schwarzschild metric is

$$ds^2 = -\frac{\Delta}{r^2} dt^2 + \frac{r^2}{\Delta} dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (2.1)$$

where  $\Delta \equiv r(r - r_s)$  and  $r_s$  is the Schwarzschild radius. A static, massless scalar  $\phi$  in this background satisfies the equation of motion

$$\partial_r(\Delta\partial_r\phi_\ell) - \ell(\ell + 1)\phi_\ell = 0, \quad (2.2)$$

where we have decomposed  $\phi$  in spherical harmonics, i.e.,  $\phi_\ell$  is the field amplitude associated with a particular  $Y_{\ell m}(\theta, \varphi)$ , with  $\ell$  the angular momentum quantum number (the equation is independent of the magnetic quantum number  $m$ , and therefore we will suppress this index in what follows). It will prove convenient to multiply eq. (2.2) by  $-\Delta$ , rewriting the equation of motion as:

$$H_\ell\phi_\ell = 0, \quad \text{where} \quad H_\ell \equiv -\Delta(\partial_r(\Delta\partial_r) - \ell(\ell + 1)), \quad (2.3)$$

which defines a ‘‘Hamiltonian,’’  $H_\ell$ , where derivatives act on everything to their right. For large  $r$ , it is easy to see that the two linearly independent solutions to eq. (2.2) are  $\phi_\ell \sim r^\ell$  (growing) or  $1/r^{\ell+1}$  (decaying). For  $r \rightarrow r_s$ ,  $\phi_\ell$  goes either as a constant or a logarithm,  $\ln[(r - r_s)/r_s]$ . The tidal response of a black hole is determined by the ratio of the decaying and growing modes at infinity, calculated under the assumption that the scalar remains regular at the horizon. The key task is therefore to connect the growing/decaying branch of solutions to the horizon asymptotics.

Let us define the following operators:

$$\begin{aligned} D_\ell^+ &\equiv -\Delta\partial_r + \frac{\ell + 1}{2}(r_s - 2r), \\ D_\ell^- &\equiv \Delta\partial_r + \frac{\ell}{2}(r_s - 2r). \end{aligned} \quad (2.4)$$

The geometric origin of these operators is discussed in Appendix A. An explicit calculation shows that they satisfy the following relations:<sup>6</sup>

$$\begin{aligned} D_{\ell+1}^- D_\ell^+ - D_{\ell-1}^+ D_\ell^- &= \frac{(2\ell + 1)r_s^2}{4}, \\ H_\ell = D_{\ell+1}^- D_\ell^+ - \frac{(\ell + 1)^2 r_s^2}{4} &= D_{\ell-1}^+ D_\ell^- - \frac{\ell^2 r_s^2}{4}. \end{aligned} \quad (2.6)$$

The  $D_\ell^\pm$  operators are raising and lowering operators in the sense that their algebra with the Hamiltonian

---

<sup>6</sup>Though it is not important to our argument, it is worth noting that if one deforms  $D^\pm$  by introducing a phase that catalogues the  $\ell$  value as:

$$L_\pm \equiv \frac{2}{r_s} e^{\pm i\alpha} \left( \mp \Delta\partial_r + \frac{1}{2}\Delta'(r)i\partial_\alpha \right), \quad L_0 \equiv -i\partial_\alpha, \quad (2.5)$$

then  $L_\pm$  and  $L_0$  satisfy an SL(2) algebra:  $[L_\pm, L_0] = \mp L_\pm$ ,  $[L_+, L_-] = -2L_0$ .

is<sup>7</sup>

$$H_{\ell+1}D_\ell^+ = D_\ell^+H_\ell, \quad H_{\ell-1}D_\ell^- = D_\ell^-H_\ell. \quad (2.7)$$

Using this structure, from a solution at level  $\ell$ , i.e.,  $\phi_\ell$  such that  $H_\ell\phi_\ell = 0$ , we can raise to a level  $\ell + 1$  solution. That is,  $D_\ell^+\phi_\ell$  solves  $H_{\ell+1}(D_\ell^+\phi_\ell) = 0$ . Similarly, we can lower to a level  $\ell - 1$  solution, i.e.,  $D_\ell^-\phi_\ell$  solves  $H_{\ell-1}(D_\ell^-\phi_\ell) = 0$ . Also, both  $D_{\ell+1}^-D_\ell^+\phi_\ell$  and  $D_{\ell-1}^+D_\ell^-\phi_\ell$  reproduce  $\phi_\ell$  (up to a constant normalization), provided  $H_\ell\phi_\ell = 0$ .

From this algebraic structure a whole tower of solutions falls into our lap immediately. First, the shift symmetry of the scalar guarantees that  $\phi_0 = \text{constant}$  is a good  $\ell = 0$  solution; then applying  $D_{\ell-1}^+ \cdots D_1^+ D_0^+ \phi_0$  gives us the corresponding solution at level  $\ell$ . From the form of the raising operator, one can see the level  $\ell$  solution is a polynomial in  $r$  of the form  $\phi_\ell \sim 1 + r + \cdots + r^\ell$ . Such a finite polynomial is manifestly regular at the horizon,  $r = r_s$ . This solution belongs to the growing branch: going as  $r^\ell$  at large  $r$ .

It is not difficult to write down the precise decaying solution for  $\ell = 0$ , and raise it to arbitrary  $\ell$  to fill out the decaying branch.<sup>8</sup> Instead we take a different tack: let us delve deeper into the symmetries behind the ladder structure. Here, we follow the strategy of [44] who pointed out an analogous ladder in de Sitter space.<sup>9</sup>

We can use the algebraic structure to construct conserved quantities for each  $\ell$ . At the  $\ell = 0$  level, it is easy to see:

$$Q_0 \equiv \Delta\partial_r \quad \text{satisfies} \quad [Q_0, H_0] = 0. \quad (2.8)$$

Thus  $Q_0$  generates a symmetry in the  $\ell = 0$  sector. Climbing the ladder, we can construct  $Q_1 \equiv D_0^+Q_0D_1^-$  which satisfies  $[Q_1, H_1] = 0$ , i.e.,  $D_0^+Q_0D_1^-H_1 - H_1D_0^+Q_0D_1^- = D_0^+[Q_0, H_0]D_1^- = 0$ . Proceeding inductively, we have:

$$Q_\ell \equiv D_{\ell-1}^+Q_{\ell-1}D_\ell^- \quad \text{which satisfies} \quad [Q_\ell, H_\ell] = 0. \quad (2.9)$$

It can be shown  $\delta\phi_\ell = Q_\ell\phi_\ell$  is a symmetry of the action, from which a corresponding Noether current can be derived.<sup>10</sup> In the static limit, current conservation takes the form  $\partial_r J_\ell^r(r) = 0$ , i.e., the  $r$  component of the Noether current is a *conserved* quantity, in the sense of being *r-independent*. For this reason, we will refer to it as “charge” in what follows.

It is simpler to identify the conserved quantities by inspection. Equation (2.2) tells us, for  $\ell = 0$ :

$$P_0 \equiv \Delta\partial_r\phi_0 \quad \text{is conserved:} \quad \partial_r P_0 = 0, \quad (2.10)$$

<sup>7</sup>In this sense, the ladder structure that we are discussing is slightly different from the more familiar ladder structure in, for example, the harmonic oscillator. Here the ladder operators map a solution of a given Hamiltonian—indexed by  $\ell$ —to a corresponding eigenstate of a *different* Hamiltonian, indexed by  $\ell \pm 1$ . This is in contrast to the harmonic oscillator where ladder operators map between different states of the same Hamiltonian.

<sup>8</sup>The exact  $\ell = 0$  decaying solution is  $\ln[(r - r_s)/r]$ .

<sup>9</sup>In fact, as we discuss in Appendix A, one can change variables to cast the static equation of motion (2.2) as the equation for a (massive) scalar in a three-dimensional Euclidean anti-de Sitter geometry. From this perspective, the vanishing of the static response is related by analytic continuation to the transparency of odd-dimensional de Sitter spaces.

<sup>10</sup>Strictly speaking, there should be a dimensionful factor relating  $\delta\phi_\ell$  and  $Q_\ell\phi_\ell$ , which we keep implicit.

where  $\phi_0$  is now an arbitrary solution of the  $\ell = 0$  equation. Conserved quantities at higher  $\ell$  can be obtained by climbing the ladder:

$$\partial_r P_\ell = 0, \quad \text{where} \quad P_\ell \equiv \Delta \partial_r (D_1^- D_2^- \cdots D_\ell^- \phi_\ell). \quad (2.11)$$

The philosophy is that, starting from any solution at level  $\ell$ , one can descend to  $\ell = 0$  by a series of lowering operations, and then invoke the conservation of  $P_0$  to see that  $P_\ell$  is conserved. The Noether current associated to the symmetry  $\delta\phi_\ell = Q_\ell\phi_\ell$  turns out to be actually  $J_\ell^r = (P_\ell)^2$ , but clearly this implies that  $P_\ell$  is also conserved.<sup>11</sup> We will sometimes abuse the terminology and refer to  $P_\ell$  itself as the conserved charge.

The utility of the conserved quantities  $P_\ell$  is that they can be used to deduce properties of the solutions to (2.2), without knowing their precise form. In particular this allows us to understand how solutions with particular behaviors in the two asymptotic regions (near the horizon and near  $r = \infty$ ) connect to each other without explicitly solving the differential equation. Consider the decaying branch of solutions, which goes like  $\phi_\ell \sim 1/r^{\ell+1}$  at large  $r$ . Each  $D^-$  operator essentially increases the power in  $r$  by 1 at large  $r$ . Thus  $D_1^- \cdots D_\ell^- (1/r^{\ell+1})$  scales as  $1/r$  at large  $r$ , so that its corresponding  $P_\ell$  is nonzero. Near the horizon, this solution must have the same  $P_\ell$ , since it is conserved as a function of  $r$ . Of the two possible horizon asymptotics, the constant solution yields  $P_\ell = 0$  while the logarithm,  $\ln[(r-r_s)/r_s]$ , yields a non-vanishing  $P_\ell$  as  $r \rightarrow r_s$ . Thus, conservation of  $P_\ell$  tells us the decaying branch of solutions must diverge logarithmically at the horizon.<sup>12</sup>

We already know that the growing branch, defined by  $\phi_\ell = D_{\ell-1}^+ \cdots D_0^+ \phi_0$ , with a constant  $\phi_0$ , is regular at the horizon (by virtue of being a finite polynomial). But we can see this by charge conservation, too. The series of lowering operators in  $P_\ell$  merely reverts  $\phi_\ell$  back to the constant  $\phi_0$  (up to normalization) which implies that  $P_\ell = 0$  for the growing branch. Conservation of  $P_\ell$  thus tells us this solution, when extended to the horizon, must approach a constant rather than the divergent logarithm.<sup>13</sup>

It is worth emphasizing the vanishing of the Love number follows from *two* facts: (1) the purely decaying solution ( $\sim 1/r^{\ell+1}$  at large  $r$ ) is divergent at the horizon, and (2) the solution that is regular at the horizon is a finite polynomial going as  $1 + r + \cdots + r^\ell$ . The second fact by itself is not sufficient: it does not forbid us from adding to it a decaying tail that goes as  $1/r^{\ell+1}$ .<sup>14</sup> Indeed, an object like a star should have this decaying tail (assuming it is coupled to the scalar), signifying that the star develops a multipole moment under the deforming influence of the  $r^\ell$  tidal field. A black hole, by the demand of regularity at the horizon, cannot have this tail—implying that its static response vanishes, meaning zero Love number.

<sup>11</sup>Notice that  $P_\ell$  and  $\delta\phi_\ell$  happen to coincide for  $\ell = 0$ , but are different from each other for non-zero values of  $\ell$ .

<sup>12</sup>The solution that is purely decaying mode at infinity  $\sim 1/r^{\ell+1}$  is an admixture of the constant and logarithmic fall-offs near the horizon, but the important point is that it is not regular at the horizon.

<sup>13</sup>Interestingly, charge conservation implies that the solution relevant for the computation of Love numbers has a single fall-off when expanded in each of the asymptotic regions. This is closely related to the phenomenon of reflection-less transmission in quantum mechanics, which we elaborate on in Appendix B.

<sup>14</sup>In order to infer that Love numbers vanish, in addition to the polynomial solution, we need to know that the other solution diverges at the horizon. This can alternatively be inferred from analysis of the differential equation in the near-horizon regime. However, to make statements about black hole hair we need to be able to connect the solution that goes as  $1/r^{\ell+1}$  at infinity to these near-horizon solutions, which fact (1) achieves.

Fact (1) is also the content of the no-hair theorem. Indeed, Bekenstein showed that a black hole cannot sustain static, scalar hair that decays at infinity [26]. His celebrated proof makes minimal assumptions about the spacetime, and is thus more general, but a crucial ingredient is the existence of a horizon. Our derivation shows that both the no-hair theorem, and the vanishing of static response, follow from the ladder symmetries of a massless scalar in a Schwarzschild background.

A pleasing feature of the story is that one can also motivate the disposal of the decaying solution by *symmetry*, rather than by the requirement of *regularity*. For instance, at the  $\ell = 0$  level, it is simple to check that the symmetry transformation  $\delta\phi_0 \equiv Q_0\phi_0$  vanishes if  $\phi_0$  is a constant (the growing solution), and is non-zero if  $\phi_0$  goes as  $1/r$  at large  $r$  (the decaying solution). The same pattern can be established for the higher  $\ell$ 's by ascending the ladder. That is,  $\delta\phi_\ell \equiv Q_\ell\phi_\ell$  vanishes for the growing branch (thus respecting the symmetry) and is non-zero for the decaying branch (thus spontaneously breaking the symmetry). The symmetry perspective is helpful especially when viewing the black hole from far away—such an observer might not even be aware of the existence of a horizon (let alone regularity on it), but could determine whether the field configuration at large  $r$  is symmetric. This is discussed further in Section 5.

As an aside, it is worth stressing there are two different kinds of symmetries at work. One mixes  $\ell$  levels, which we call a *vertical* symmetry. It acts on the fields as

$$\delta\phi_\ell = D_{\ell-1}^+\phi_{\ell-1}, \quad \delta\phi_{\ell-1} = -D_\ell^-\phi_\ell. \quad (2.12)$$

This is a symmetry of the action, and has a geometric origin (see Appendix A). Its existence is responsible for the ladder structure. The other symmetry—associated with the conservation of  $P_\ell$ —stays within the same level (acts on a given rung of the ladder). We call this a *horizontal* symmetry:  $\delta\phi_\ell = Q_\ell\phi_\ell$ . It owes its existence to the horizontal symmetry at  $\ell = 0$ , which can then be bootstrapped to higher  $\ell$ 's by climbing the ladder.

It is also worth mentioning that we have specialized to  $D = 4$ , but the equation of motion for a scalar in general dimension can be cast in the form (2.2) with  $\ell$  replaced with  $\hat{\ell} \equiv \ell/(D-3)$ . All of the same ladder structure is present in general dimensions, but only integer  $\hat{\ell}$  can be connected to  $\hat{\ell} = 0$  via the ladder, so only these values have conserved quantities. Indeed, precisely these values have vanishing Love numbers in general dimension [8, 17].

We next extend the discussion to a scalar field in the Kerr geometry, before treating the general case.

### 3 Ladder in Kerr

The algebraic ladder structure that we have discussed has a direct analogue in a Kerr background. The Kerr line element in Boyer–Lindquist (BL) coordinates is:

$$ds^2 = -\frac{\Delta}{\rho^2} (dt - a \sin^2 \theta d\varphi)^2 + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \frac{\sin^2 \theta}{\rho^2} (a dt - (r^2 + a^2) d\varphi)^2, \quad (3.1)$$

where we have defined the quantities

$$\rho^2 \equiv r^2 + a^2 \cos^2 \theta, \quad \Delta \equiv r^2 - rr_s + a^2, \quad (3.2)$$

where also  $\Delta = (r - r_+)(r - r_-)$  with  $r_{\pm} \equiv r_s/2 \pm \sqrt{(r_s/2)^2 - a^2}$  being the locations of the inner and outer horizons. It is useful to note the relations:  $r_+ r_- = a^2$ ,  $a^2 + r_+^2 = r_s r_+$  and  $g^{\varphi\varphi} = (1 - \rho^{-2} r_s r) / (\Delta \sin^2 \theta)$ . The equation of motion for a static, massless scalar in a Kerr background is, after decomposing in spherical harmonics  $Y_{\ell m}$ :

$$\partial_r (\Delta \partial_r \phi_\ell) + \frac{a^2 m^2}{\Delta} \phi_\ell - \ell(\ell + 1) \phi_\ell = 0, \quad (3.3)$$

where the magnetic quantum number,  $m$ , dependence of  $\phi_\ell$  is again suppressed. It is useful to make the following definitions:

$$\begin{aligned} z &\equiv r - r_-, & z_k &\equiv r_+ - r_- \\ q &\equiv \frac{am}{z_k}, & \phi_\ell &\equiv e^{-iq \ln[(z - z_k)/z]} \psi_\ell. \end{aligned} \quad (3.4)$$

Note that  $\partial_r e^{-iq \ln[(z - z_k)/z]} = -(iam/\Delta) e^{-iq \ln[(z - z_k)/z]}$ , and  $\Delta = z(z - z_k)$ . After these changes of variables, the scalar equation of motion can thus be recast as:

$$\partial_z (\Delta \partial_z \psi_\ell) - 2iqz_k \partial_z \psi_\ell - \ell(\ell + 1) \psi_\ell = 0. \quad (3.5)$$

As before, it is convenient to multiply this equation by  $-\Delta$  and rewrite:

$$H_\ell \psi_\ell = 0, \quad \text{with} \quad H_\ell \equiv -\Delta (\partial_z (\Delta \partial_z) - 2iqz_k \partial_z - \ell(\ell + 1)). \quad (3.6)$$

At large  $z$ , the growing/decaying solutions go as  $z^\ell$  or  $1/z^{\ell+1}$ , respectively. Close to the horizon ( $z \rightarrow z_k$ ), the two asymptotic solutions are  $\psi_\ell \sim \text{constant}$ , and  $\psi_\ell \sim e^{2iq \ln[(z - z_k)/z_k]}$ .

The equation (3.5) has a similar ladder structure to the Schwarzschild scalar. It can be discerned by defining:

$$\begin{aligned} D_\ell^+ &\equiv -\Delta \partial_z + \frac{(\ell + 1)}{2} (z_k - 2z) + iqz_k, \\ D_\ell^- &\equiv \Delta \partial_z + \frac{\ell}{2} (z_k - 2z) - iqz_k. \end{aligned} \quad (3.7)$$

These operators obey very similar relations to (2.6):

$$\begin{aligned} D_{\ell+1}^- D_\ell^+ - D_{\ell-1}^+ D_\ell^- &= \frac{(2\ell + 1)z_k^2}{4}, \\ H_\ell = D_{\ell+1}^- D_\ell^+ - \left( \frac{(\ell + 1)^2}{4} + q^2 \right) z_k^2 &= D_{\ell-1}^+ D_\ell^- - \left( \frac{\ell^2}{4} + q^2 \right) z_k^2. \end{aligned} \quad (3.8)$$

In addition, exactly the same “commutators” as in eq. (2.7) apply here, from which the interpretation of  $D^\pm$  as raising and lowering operators follows. Note that the ladder structure extends in  $\ell$  but not in  $m$ . It is possible to formally climb to any  $\ell$ , including  $\ell < |m|$ . Equation (3.3) does not by itself forbid this. It is the connection to  $Y_{\ell m}$  that ultimately limits  $|m| \leq \ell$ .

The introduction of  $\psi_\ell$  to replace  $\phi_\ell$  in eq. (3.4) is useful not only for making the equation resemble its Schwarzschild counterpart, but also for thinking about horizon regularity. The logarithmic phase that relates  $\phi_\ell$  and  $\psi_\ell$  is motivated by the azimuthal coordinate that was used originally by Kerr [45]:  $\varphi_{\text{Kerr}} = \varphi - (a/z_k) \ln[(z - z_k)/z]$ . The point is that a scalar quantity like  $(\partial\phi)^2$ , when evaluated in BL coordinates, has a potentially divergent term at the horizon coming from  $g^{\varphi\varphi} \propto 1/\Delta$ . The phase is chosen so this is precisely cancelled by another potentially dangerous term from the radial gradients. Imposing horizon regularity is thus translated into demanding the regularity of  $\psi_\ell$ , or more concretely, demanding that  $\Delta\partial_z\psi_\ell = 0$  as  $z \rightarrow z_k$ . Moreover, according to calculations by [15, 16, 18], the static Love number can be inferred by examining the power series for  $\psi_\ell$ : absence of a  $1/r^{\ell+1}$  term in  $\psi_\ell$  is tantamount to a vanishing Love number.<sup>15</sup>

Returning to the ladder structure: just as in the Schwarzschild case we can define a set of horizontal symmetries that act only at a fixed  $\ell$ . The symmetry transformation is  $\delta\psi_\ell = Q_\ell\psi_\ell$  with  $Q_\ell$  given by

$$\begin{aligned} Q_0 &\equiv \Delta\partial_z, & [Q_0, H_0] &= 0, \\ Q_\ell &\equiv D_{\ell-1}^+ Q_{\ell-1} D_\ell^-, & [Q_\ell, H_\ell] &= 0. \end{aligned} \tag{3.9}$$

The corresponding conserved charge is, as before, most easily obtained by inspection, starting from  $\ell = 0$ :

$$P_0 \equiv (\Delta\partial_z - 2iqz_k)\psi_0 \quad \text{is conserved :} \quad \partial_z P_0 = 0. \tag{3.10}$$

Higher  $\ell$  charges,  $P_\ell$ , are found by climbing the ladder:

$$\partial_z P_\ell = 0, \quad \text{where} \quad P_\ell \equiv (\Delta\partial_z - 2iqz_k) D_1^- \cdots D_\ell^- \psi_\ell, \tag{3.11}$$

for any level- $\ell$  solution  $\psi_\ell$ .

The growing branch of solutions can again be raised from the  $\psi_0 = \text{constant}$  solution:  $\psi_\ell = D_{\ell-1}^+ \cdots D_0^+ \psi_0$  is the corresponding level  $\ell$  solution. It is manifestly a finite polynomial  $\sim 1 + \cdots + z^\ell$  (from the form of  $D^+$ ), and is thus regular at the horizon.

The conserved charge  $P_\ell$  evaluated for the so-defined growing branch is non-zero, unlike in the Schwarzschild case (for example, eq. (3.10) tells us  $P_0 \propto iam$  for a constant, non-vanishing  $\psi_0$ ). A decaying solution that goes as  $1/z^{\ell+1}$  at large  $z$  can be shown to give a non-vanishing  $P_\ell$  as well. Of the two horizon asymptotics, the constant gives a non-vanishing  $P_\ell$  while  $e^{2iq \ln[(z-z_k)/z_k]}$  yields  $P_\ell = 0$ . Thus, charge conservation does not by itself tell us whether the decaying solution contains a  $e^{2iq \ln[(z-z_k)/z_k]}$  contribution when it is extended to the horizon (though it does tell us it must contain the constant contribution).

---

<sup>15</sup>If one were to expand out the logarithmic phase, i.e., look at the power series expansion for  $\phi_\ell$  instead of  $\psi_\ell$ , a  $1/r^{\ell+1}$  term is actually present. But its presence is unrelated to the Love number, defined by the coefficient of the tidal response operator  $\sim (\partial^\ell \phi)^2$  in the point-particle action [8, 16–18, 46]. The coefficient of the  $1/r^{\ell+1}$  tail from expanding out the logarithmic phase is instead capturing the dissipative effects of tidal heating [15, 16, 18].

In order to figure out how the solution that decays at infinity behaves near the horizon, what we need is the Wronskian:

$$W[\psi_\ell^a, \psi_\ell^b] = e^{-2iq \ln \left[ \frac{z-z_k}{z} \right]} (\psi_\ell^a \Delta \partial_z \psi_\ell^b - \psi_\ell^b \Delta \partial_z \psi_\ell^a), \quad (3.12)$$

where  $\psi_\ell^a$  and  $\psi_\ell^b$  are any two level- $\ell$  solutions. It can be shown  $\partial_z W = 0$  (this is easiest to see by relating back to  $\phi_\ell$  and inspecting eq. (3.3)). We can choose  $\psi_\ell^a$  to be the regular, growing solution, and  $\psi_\ell^b$  to be the decaying solution. Evaluating  $W$  at large  $z$  tells us  $W$  is proportional to the normalization of the  $1/z^{\ell+1}$  tail. The decaying solution, when extended to the horizon, in general can have both the constant and the  $e^{2iq \ln[(z-z_k)/z_k]}$  contributions, but the constant cannot contribute to  $W$  (for  $\psi_\ell^a$  also goes to a constant at the horizon) while  $e^{2iq \ln[(z-z_k)/z_k]}$  contributes to a non-vanishing  $W$ . By the constancy of  $W$ , we infer that the decaying solution must be irregular at the horizon. We say *irregular* in the sense that  $\Delta \partial_z \psi_\ell \neq 0$  as  $z \rightarrow z_k$ , for  $\psi_\ell \sim e^{2iq \ln[(z-z_k)/z_k]}$ —recalling that  $(\partial\phi)^2$  (or its spherical average) contains  $\Delta |\partial_z \psi_\ell|^2$ , this tells us  $(\partial\phi)^2$  diverges at the horizon.

Fact (1)—that the purely decaying solution is irregular at the horizon—and fact (2)—that the regular solution is a finite polynomial  $\sim 1 + \dots + z^\ell$ —are thus established, leading to no-hair theorem and vanishing Love number for a scalar around a Kerr black hole. Furthermore, the same symmetry understanding as in the Schwarzschild case applies here: the decaying solution spontaneously breaks the horizontal ladder symmetry, i.e.,  $\delta\psi_\ell \equiv Q_\ell \psi_\ell \neq 0$ , while the growing (regular) solution respects it. Next, we extend the argument to vector and tensor perturbations.

## 4 Ladder in Spin: From Scalar to Vector and Tensor

Finally, we want to treat the most general case of interest, a field of arbitrary (integer) spin  $s \leq 2$  in a Kerr background. Such perturbations are described by the Teukolsky equation, which in the static limit in BL coordinates is [47, 48]:

$$\partial_r \left( \Delta \partial_r \phi_\ell^{(s)} \right) + s(2r - r_s) \partial_r \phi_\ell^{(s)} + \left( \frac{a^2 m^2 + is(2r - r_s)am}{\Delta} - (\ell - s)(\ell + s + 1) \right) \phi_\ell^{(s)} = 0, \quad (4.1)$$

where a (spin-weighted) spherical harmonic transform is assumed, and the  $m$  dependence of the Newman–Penrose variable  $\phi_\ell^{(s)}$  is suppressed. Making the same substitutions as in eq. (3.4), we have:<sup>16</sup>

$$\tilde{H}_{\ell s} \psi_\ell^{(s)} = 0, \quad \text{where} \quad \tilde{H}_{\ell s} \equiv \partial_z (\Delta \partial_z) - [(z_k - 2z)s + 2iqz_k] \partial_z - (\ell - s)(\ell + s + 1). \quad (4.2)$$

The large  $z$  asymptotics are:  $\psi_\ell^{(s)} \sim z^{\ell-s}$  and  $1/z^{\ell+s+1}$ . The near-horizon asymptotics are:  $\psi_\ell^{(s)} \sim \text{constant}$  and  $e^{(2iq-s)\ln[(z-z_k)/z_k]}$ . A ladder structure raising and lowering  $\ell$  can be found (see Appendix D), but there is another ladder more useful to us. We define the operators

$$E^+ \equiv \partial_z, \quad E_s^- \equiv \Delta \partial_z - s(z_k - 2z) - 2iqz_k. \quad (4.3)$$

These are raising and lowering operators in the spin,  $s$ :

$$\begin{aligned} E^+ E_s^- - E_{s+1}^- E^+ &= 2s, \\ \tilde{H}_{\ell s+1} E^+ &= E^+ \tilde{H}_{\ell s}, & \tilde{H}_{\ell s-1} E_s^- &= E_s^- \tilde{H}_{\ell s}. \end{aligned} \quad (4.4)$$

This implies that given a solution  $\psi_\ell^{(s)}$ , one can generate  $\psi_\ell^{(s+1)} = E^+ \psi_\ell^{(s)}$  and  $\psi_\ell^{(s-1)} = E_s^- \psi_\ell^{(s)}$ . We can use this spin ladder structure to construct conserved charges at each  $\ell$  and  $s$  from the known one at  $\ell = 0$ ,  $s = 0$  (eq. (3.10)):

$$\partial_z P_0^{(0)} = 0, \quad \text{where} \quad P_0^{(0)} \equiv (\Delta \partial_z - 2iqz_k) \psi_0^{(0)}. \quad (4.5)$$

For instance, the conserved charge for a general  $\ell$  and  $s = 2$  solution  $\psi_\ell^{(2)}$  is

$$P_\ell^{(2)} \equiv (\Delta \partial_z - 2iqz_k) D_1^- \cdots D_\ell^- E_1^- E_2^- \psi_\ell^{(2)}, \quad (4.6)$$

where the  $D^-$ 's are  $\ell$ -lowering operators for the scalar (3.7). The two  $E^-$  operators serve to lower the spin from 2 to 0, and the  $D^-$  operators then lower  $\ell$  to 0. The resulting  $\psi_0^{(0)}$  can be plugged into eq. (4.5) to show  $\partial_z P_\ell^{(2)} = 0$ .

The growing branch of solutions for  $s = 2$  can be obtained by, once again, starting from the scalar  $\psi_0^{(0)} = \text{constant}$  solution. Then  $\psi_\ell^{(2)} = E^+ E^+ D_{\ell-1}^+ \cdots D_0^+ \psi_0^{(0)}$  is manifestly a finite polynomial of the form  $1 + z + \cdots + z^{\ell-2}$ , judging from the form of  $E^+$  and  $D^+$ . This implies that the purely growing at infinity solutions are regular at the horizon.

The decaying branch of solutions for  $s = 2$  can be obtained from  $E^+ E^+ \psi_\ell^{(0)}$  where  $\psi_\ell^{(0)}$  is the level  $\ell$  scalar decaying solution we have worked out previously. Recall that  $\psi_\ell^{(0)}$  scales as  $1/z^{\ell+1}$  at large  $z$  and

<sup>16</sup>Note that here we have not multiplied by an additional factor of  $-\Delta$  in order to define  $\tilde{H}$ . This is because the spin ladder structure is most simply phrased in terms of  $\tilde{H}$ . In Appendix D we describe the  $\ell$  ladder structure, which uses  $H$  defined analogously to the previous cases.

contains a  $e^{2iq \ln[(z-z_k)/z_k]}$  contribution as  $z \rightarrow z_k$ . Therefore,  $E^+ E^+ \psi_\ell^{(0)}$  goes as  $1/z^{\ell+3}$  at large  $z$  and contains  $e^{(2iq-2) \ln[(z-z_k)/z_k]}$  close to the horizon. This means that the decaying branch is irregular at the horizon in the same sense as before.

We have thus established facts (1) and (2) for tensor perturbations  $s = 2$ , from which no-hair theorem and vanishing Love numbers follow.<sup>17</sup> The same arguments translate straightforwardly to vector perturbations  $s = 1$ . It is important to emphasize the spin-weighted spherical harmonics restrict us to  $\ell \geq s$ . Thus, the absence of hair and Love numbers holds for  $\ell \geq 2$  tensor perturbations, and  $\ell \geq 1$  vector ones.

The symmetry story works as before (recognizing that with the Teukolsky equation, we can only work with the equation of motion as opposed to an action): the growing branch respects the horizontal ladder symmetry while the decaying branch spontaneously breaks it. Details are given in Appendix D.

Here we have focused on the Teukolsky equation to see the symmetries underlying the computation of black hole Love numbers, but it is reasonable to ask whether similar structures can be seen at the level of metric perturbations (as opposed to Weyl scalars). Around a Schwarzschild background this is indeed possible, as we discuss explicitly in Appendix C.

---

<sup>17</sup>To go from the Teukolsky solution  $\psi_\ell^{(2)}$  to the Love numbers (really elements of the Love tensor), a matching computation is needed to connect the fall-offs of  $\psi_\ell^{(2)}$  to the coefficients of the tidal response operators  $\sim (\partial^{\ell-2} C)^2$  in the point-particle action, with  $C$  the electric/magnetic part of the Weyl tensor. See [16, 18]. Similar comments as in footnote 15 apply here.

## 5 Symmetries in the Infrared

Our discussion so far has revolved around symmetries present in the microphysical computation of static responses of black holes. However, to a distant observer, such microscopic details of the black hole geometry are invisible. Instead, they will infer the black hole to be effectively a point particle, whose properties they can probe for example by applying an external electric field, or gravitational tidal field, and measuring the response. In this effective description, the properties of the black hole are encoded in an infinite number of coefficients appearing in the point-particle action; the values of these coefficients are determined by the requirement that the measured responses are correctly reproduced. From this effective field theory (EFT) point of view, the vanishing of black hole static response coefficients is somewhat puzzling, because it would seem to be a fine-tuning [49–51]. It is natural to ask whether the ultraviolet symmetries we have identified can provide a symmetry explanation in the infrared, and in this section we explore this question.

For concreteness, we will specialize to the simplest setting: the worldline EFT of a massive (non-spinning) particle coupled to a scalar field, which is the relevant effective theory describing, for example, the response of a Schwarzschild black hole to an externally applied scalar field profile. The effective theory (up to quadratic order in  $\phi$ ) governing static field configurations takes the form of a bulk action plus the point-particle worldline action:

$$S = -\frac{1}{2} \int d^4x (\partial\phi)^2 + \int d\tau \gamma \left[ \frac{1}{2} \gamma^{-2} \dot{x}^\nu \dot{x}_\nu - \frac{\mu^2}{2} - g\phi + \sum_{\ell=0}^{\infty} \frac{\lambda_\ell}{2\ell!} (\partial_{(a_1} \cdots \partial_{a_\ell)_T} \phi)^2 \right], \quad (5.1)$$

where  $x^\nu$  denotes the spacetime position of the point particle, with  $\mu$  its mass. The variable  $\gamma$  is a worldline vielbein enforcing reparametrization invariance of the effective action and  $(\cdots)_T$  denotes the symmetrized traceless part of the enclosed indices. The indices  $a_1 \cdots a_\ell$  are defined by projecting space-time indices onto the particle rest frame—see e.g. [17] for an explicit definition of such projection.

It is simplest to think of the particle as being at rest at the origin, in which case a term like  $g\phi$  has the effect of adding a source term to the (static)  $\phi$  equation of motion, i.e.,  $\nabla^2\phi = g\delta_D$  where  $\delta_D$  is the 3D Dirac delta function centered at the origin. As such,  $g$  can be thought of as the monopole scalar charge. If one wishes to describe a non-spherically-symmetric object, the possibility of higher multipole hair could be included by adding linear terms of the form  $\partial^\ell\phi$ . The couplings  $\lambda_\ell$  are the response coefficients that encode the particle’s response to a static external field with multipole moment  $\ell$ . Their presence means the static  $\phi$  equation of motion takes the schematic form  $\nabla^2\phi \sim \lambda_\ell \partial^\ell(\partial^\ell\phi \delta_D)$ . Imagine the external tidal field as the input  $\phi$  on the right hand side; solving this equation then gives the linear response, that is, an  $r^\ell$  tidal field sources an  $r^{-(\ell+1)}$  response. Note that in eq. (5.1) the metric is treated as flat, as is appropriate at a large distance  $r$ . Beyond the lowest order, one would also need to take into account couplings with metric perturbations, sourced by the particle itself, whose effect can be included order by order in  $r_s/r$ .

A computation in the ultraviolet, such as those discussed in the previous sections, can be compared against the infrared predictions to fix the EFT coefficients  $g$  and  $\lambda_\ell$ . Indeed, we conclude that they would vanish if the point particle were a black hole. But here, we wish to ask a different question: what symmetries, visible in the infrared, might tell us that the hair or tidal response coefficients should vanish?

A possible candidate is the large distance limit of the ultraviolet symmetries we have worked out—the

ladder symmetries. As we show in Appendix A, the geometric perspective, in which a suitably rescaled and reduced Schwarzschild black hole is a Euclidean anti-de Sitter space in disguise, offers a particularly simple way to think about the whole ladder structure. The associated isometries have a well defined flat space (i.e. infrared) limit, giving the symmetry transformation in (A.16):

$$\delta\phi = r^2 \cos\theta \partial_r \phi + r \partial_\theta (\sin\theta \phi), \quad (5.2)$$

after scaling out an irrelevant overall factor. The angular structure of this transformation mixes different  $\ell$  values in spherical harmonic space, leading to the (large  $r$  limit of)  $D_\ell^\pm$  operators defined in (2.4). The transformation (5.2) may look unfamiliar, but it is actually nothing more than a special conformal transformation, written in spherical coordinates.<sup>18</sup>

We now want to understand how the symmetry (5.2), or equivalently (5.3), constrains the effective action (5.1). The bulk  $(\partial\phi)^2$  term is clearly invariant in the static limit, but none of the worldline-localized couplings are invariant. This is more subtle than it might appear at first—since the transformation (5.3) involves factors of  $\vec{x}$ , one may be tempted to conclude that the variation of an operator localized on the worldline vanishes when computed at the location of the point particle,  $\vec{x} = 0$ . However, this makes a tacit assumption about the behavior of the field at  $\vec{x} = 0$ . This issue can be settled by showing that the would-be Noether currents associated with the symmetry transformation (5.2) in  $\ell$  space are not conserved on-shell when any of the worldline couplings are present.<sup>19</sup> Thus, (5.2) is the sought-after infrared symmetry which forbids Love number (and hair) couplings in the point-particle effective theory.

This appears to be a consistent story for black holes. How about stars, which in general can have a non-trivial hair and tidal response? The symmetry (5.2) would appear to be broken for them. How does that happen? To understand this, it is useful to recall the *horizontal* part of the ladder symmetries, effected by  $Q_\ell$  defined in (2.8) and (2.9), for which a large  $r$  limit also exists.<sup>20</sup> Recall the purely growing mode solution  $\phi_\ell^{(g)} \sim r^\ell + \dots$  preserves the horizontal symmetry  $Q_\ell \phi_\ell^{(g)} = 0$ . On the other hand, any solution that has an admixture of the decaying solution at infinity,  $\phi_\ell^{(d)} \sim 1/r^{\ell+1} + \dots$ , breaks the symmetry spontaneously, in the sense that  $Q_\ell \phi_\ell^{(d)} \neq 0$ . For stars, the boundary conditions force us to include some contribution from this decaying branch, leading to them having a nonzero tidal response. From this viewpoint, we would expect the symmetry (5.2) to be either realized nonlinearly in the EFT, or absent. In either case, it is consistent that the EFT couplings that reproduce tidal responses are not invariant under the linearly realized symmetry.

It is worth stressing that imposing the symmetry (5.2) rules out *all* multipolar hair and response couplings in one fell swoop (as is appropriate for a 4D black hole in general relativity). For a more general object,

---

<sup>18</sup>This can be made manifest by transforming to Cartesian coordinates and writing the transformation (5.2) in a rotationally covariant way, where it takes the more familiar form:

$$\delta\phi = c_i (x^i - \vec{x}^2 \partial^i + 2x^i \vec{x} \cdot \vec{\partial}) \phi, \quad (5.3)$$

which is a special conformal transformation for a field with weight  $1/2$ . This equation reproduces (5.2) for  $c_i = (0, 0, 1)$ . Since we are assuming that the theory is also rotationally invariant, the symmetry (5.3) with arbitrary  $c_i$  is also present.

<sup>19</sup>The non-invariance of the worldline couplings  $\lambda_\ell$  also follows from the fact that they inevitably introduce a scale in the theory, thus breaking the conformal symmetry. This is because these terms capture finite-size effects [52].

<sup>20</sup>Given the relation of  $D_\ell^\pm$  to conformal symmetry, it is tempting to speculate that the  $Q_\ell$  symmetries for the free scalar in flat space are related to known higher-spin symmetries, a subject for further investigation.

it is conceivable the tidal response or hair vanishes for some multipoles but not for others. In such a case, one would need a more flexible formulation of the ladder symmetries. Decomposing  $\phi$  in spherical harmonics, (5.2) implies a specific pattern of  $\delta\phi_\ell$ 's (A.13). One could instead consider a modified version where only a particular neighboring pair, e.g.,  $\delta\phi_\ell$  and  $\delta\phi_{\ell-1}$ , are non-vanishing (2.12)—this would be an example of the vertical symmetries restricted to a particular subset of  $\ell$ s. There are also the horizontal symmetries, one for each  $\ell$ :  $\delta\phi_\ell = Q_\ell\phi_\ell$ . Since these symmetries act on a single multipole, the structure allows—in principle—the breaking of the  $Q_\ell$  symmetries for some  $\ell$ s but not others. Exactly how this may manifest in the worldline EFT is something that we plan to explore in the future.

In this section, we have focused on the simplest situation where the hair and the tidal field is a scalar, indeed a free scalar. We expect the symmetry (5.2) to be broken once interactions are introduced. For instance, in our infrared discussion so far, we have ignored gravitational interactions by working in the flat space limit. What is interesting, perhaps surprising from the infrared point of view, is that the metric perturbations can be systematically included, in fact resummed into the Schwarzschild background, around which the symmetry does not get destroyed, but elevated to (A.12):

$$\delta\phi = \Delta \cos\theta \partial_r \phi + \left(r - \frac{r_s}{2}\right) \partial_\theta(\sin\theta\phi). \quad (5.4)$$

We have thus come full circle to the ultraviolet symmetries that got us started in this EFT discussion.

A number of questions remain to be addressed. First, it would be interesting to work out in detail how the flat limit of our symmetries prevents black holes from having non-zero spin 1 and spin 2 response. In a separate direction, the case of a scalar around a rotating black hole is harder to interpret from a geometric perspective (see Appendix A), but it remains true that the ladder symmetries have a well defined large distance limit. The same can be said for vector or tensor perturbations around a Kerr background. We expect that the ultraviolet symmetries we have identified will have similar consequences in the infrared, although the symmetries in question could only be guessed at by inspecting the Teukolsky equation. A formulation in terms of the original metric perturbations rather than Newman–Penrose variables would be desirable in order to draw a connection with the low-energy EFT of a spinning point-particle [53, 54].

## 6 Discussion

We have seen that static linearized perturbations of massless fields in black hole backgrounds have a hidden structure of ladder-like symmetries and conserved quantities. These conserved quantities allow us to relate the behaviors of solutions near infinity to their behaviors near the black hole horizon *without* explicitly solving the relevant differential equations everywhere in between. This has useful implications for understanding both black hole no-hair theorems and the vanishing of black hole Love numbers from a symmetry perspective.

The question of whether or not a black hole can have hair is one that is naturally posed from very far away. We define hair to be the coefficient of the  $1/r^{\ell+1}$  component of a static solution and the question of whether a black hole can have static hair is equivalent to asking whether or not a solution that has only this behavior exists in a black hole background. We have seen that the conservation of charges associated to what we have called the horizontal ladder symmetries demands that this solution be singular at the horizon of a black hole—indicating that it cannot have hair. The nice feature of this argument is that it is based on symmetry, but it is noticeably less general than other no-hair theorems, which apply nearly irrespective of interactions, and essentially rely only on the existence of a horizon [26, 29, 30, 55]. It is natural to wonder if there is some more refined version of the arguments, that we have given in the simplest free case, to these more general situations.

In contrast to the question of black hole hair, the question of how a black hole responds to a static external field is naturally posed near the black hole horizon. In the vicinity of a black hole, we can deduce that the two linearly independent solutions to the static equations of motion of a field are such that one of them is regular at the horizon and the other diverges. The generic expectation is that, if we were to take the (unique) regular solution and expand it as  $r \rightarrow \infty$ , there would be a linear combination of the two linearly independent fall-offs at infinity (one growing and one decaying). We call the ratio of the decaying solution to the growing solution the static response, or Love number, and it tells us how the black hole responds if we apply an external growing mode background subject to the constraint that the field profile be regular at the horizon. We have seen that for black holes, symmetry forces the solution that is regular at the horizon to *only* have the growing fall-off at infinity, meaning that black hole static response coefficients vanish. For an ordinary object, we impose different boundary conditions at the surface of the object, leading to a component of the subleading, decaying fall-off at infinity, which spontaneously breaks the symmetries.

Identifying these symmetries clarifies some aspects of the effective point particle description of black holes. In the effective theory, the vanishing of hair or static responses correspond to apparently fine-tuned points in parameter space. However, we have seen that (at least in the scalar case) the locus in parameter space where hair and response coefficients vanish is an enhanced symmetry point. Of course, this does not by itself guarantee that there should be a microphysical theory that actually gives rise to this effective theory. However, since we originally identified these symmetries in the ultraviolet, the microphysical model consisting of linearized fields in a black hole background leads precisely to this low-energy description.

The symmetry viewpoint that we are advocating proves to be useful beyond pure general relativity. Once we understand in the microscopic theory what symmetry is responsible for either the vanishing of

Love numbers or for hair not being present, we can ask if additional interactions preserve or break these symmetries and infer their properties, without having to explicitly solve the equations. As an example, it is well-known that adding a coupling of a scalar field to the Gauss–Bonnet curvature invariant of the form  $\phi R_{\text{GB}}^2$  can lead to black hole hair. We can understand this from the symmetry viewpoint: to leading order (i.e. neglecting back-reaction), this coupling breaks the  $\ell = 0$  horizontal ladder symmetry, and so now we are allowed to include operators in the effective theory that give rise to a black hole hair. (See Appendix G for more details.) We expect that it will be similarly possible to understand some broad features of other proposals for gravitational physics [56] from this perspective.

We have focused on the setting of exactly static perturbations in four spacetime dimensions for simplicity, but it is possible to relax these assumptions. In particular, the extension to finite (but small) frequencies is straightforward. The extension to other dimensions is similarly easy: as we have mentioned, the only essential difference is that the quantum number  $\ell$  should be replaced with  $\hat{\ell} = \ell/(D - 3)$ . The ladder structures will still be present, but only integer values of  $\hat{\ell}$  will be connected via the ladder to the  $\ell = 0$  equation of motion, which is necessary to be able to write down conserved charges and infer that Love numbers vanish. This is satisfying because it is known that Love numbers are actually nonzero for certain values of  $\hat{\ell}$  in general dimension, but they do indeed vanish for integer  $\hat{\ell}$  in all dimensions [8, 17], completely consistent with the expectation coming from the ladder symmetries.

It is important to stress that the symmetries we have identified apply only to linearized fields in the background of a black hole, and in that regard they are deeply mysterious—why should they be present at all? It is tantalizing to speculate that they are the manifestation of some structure present in full general relativity, specialized to this simple setting. While we do not know what such a deeper explanation is, a possible hint in this direction is provided by the geometrization of the symmetries. As we discuss in Appendix A, the ladder symmetries that we have used to understand the properties of static solutions admit a geometric description, where the linearized fields propagate in an effective AdS background—with the isometries of this space leading to the ladder structure. A related promising avenue is to further investigate the algebraic structure underlying both the vertical and horizontal ladder symmetries, which is still somewhat mysterious. We expect that understanding these geometric and algebraic structures more fully will lead to additional insights, hopefully into the nonlinear regime.

**Acknowledgements** RP would like to acknowledge the hospitality of the Abdus Salam International Centre for Theoretical Physics (ICTP) during the final stage of this project. LH is supported by the DOE DE-SC0011941 and a Simons Fellowship in Theoretical Physics. The work of AJ is part of the Delta-ITP consortium. The work of RP is supported in part by the National Science Foundation under Grant No. PHY-1915611. LS is supported by Simons Foundation Award No. 555117. ARS is supported by DOE HEP grants DOE DE-FG02-04ER41338 and FG02-06ER41449 and by the McWilliams Center for Cosmology, Carnegie Mellon University.

## A Geometric Origin of the Ladder Symmetries

Some of the mystery surrounding the ladder symmetries introduced in the main text can be eliminated by viewing them as arising geometrically. Here we show how this works for a scalar field in a Schwarzschild background, showing that in the static limit it can be recast as propagating in an effective Euclidean anti-de Sitter space. The generalization to the other cases will be discussed elsewhere.

Consider a static scalar  $\phi$  in a Schwarzschild background.<sup>21</sup> Its action takes the form:

$$S = \frac{1}{2} \int d\theta d\varphi dr \sqrt{g} \phi \square \phi, \quad (\text{A.1})$$

where the 3D metric,  $g_{ij}$ , is given by:

$$ds^2 = dr^2 + \Delta (d\theta^2 + \sin^2 \theta d\varphi^2). \quad (\text{A.2})$$

Let us rescale the metric and the scalar by the following Weyl transformation

$$\tilde{g}_{ij} = \Omega^2 g_{ij}, \quad \tilde{\phi} = \Omega^{-\frac{d-2}{2}} \phi, \quad (\text{A.3})$$

where  $d = 3$  for our case of interest. The action transforms according to the following identity:

$$S = \frac{1}{2} \int d^d x \sqrt{g} \phi \square \phi = \frac{1}{2} \int d^d x \sqrt{\tilde{g}} \left[ \tilde{\phi} \square \tilde{\phi} - \Omega^{\frac{d-2}{2}} \frac{1}{\sqrt{\tilde{g}}} \partial_i \left( \sqrt{\tilde{g}} \tilde{g}^{ij} \partial_j \Omega^{-\frac{d-2}{2}} \right) \tilde{\phi}^2 \right]. \quad (\text{A.4})$$

Choosing  $d = 3$  and  $\Omega = L^2/\Delta$ , where  $L$  is an arbitrary length scale, it can be shown that the action takes the form

$$S = \frac{1}{2} \int d^3 x \sqrt{\tilde{g}} \phi \square \phi = \frac{1}{2} \int d^3 x \sqrt{\tilde{g}} \left( \tilde{\phi} \square \tilde{\phi} + \frac{r_s^2}{4L^4} \tilde{\phi}^2 \right). \quad (\text{A.5})$$

This means that the rescaled scalar  $\tilde{\phi} = (\sqrt{\Delta}/L)\phi$  behaves as a massive scalar in the following background:

$$\tilde{ds}^2 = \left( \frac{L^2}{\Delta} \right)^2 \left( dr^2 + \Delta d\theta^2 + \Delta \sin^2 \theta d\varphi^2 \right). \quad (\text{A.6})$$

Though it is not obvious, this is a constant curvature space, with Ricci scalar

$$\tilde{R}_{\text{EAdS}} = -\frac{3r_s^2}{2L^4}. \quad (\text{A.7})$$

Since the curvature is negative, it is hyperbolic space (Euclidean AdS<sub>3</sub>). It might seem somewhat peculiar that the scalar field, which is *not* conformally coupled,<sup>22</sup> can be mapped via a Weyl transformation to a scalar with a constant mass in EAdS. The underlying reason for this is that the action for a scalar in the full four-dimensional Schwarzschild spacetime is conformally coupled, as we will explain elsewhere.

<sup>21</sup>For simplicity we restrict to completely static field configurations in this appendix. However, all of our manipulations can be extended to finite frequencies whose wavelength is large compared to the other scales in the problem (which is the near-horizon limit of [43, 57]).

<sup>22</sup>One way to check this is to note that the scalar's mass in EAdS is  $m^2 = -\frac{1}{6}\tilde{R}_{\text{EAdS}}$ , which is the  $d = 4$  conformally coupled value, instead of the  $d = 3$  conformally coupled value  $m^2 = -\frac{1}{8}\tilde{R}_{\text{EAdS}}$ .

Regardless of its origin, the fact that (A.1) is related by a Weyl transformation to a scalar theory in EAdS (A.5) implies that the isometries of EAdS (which act as conformal Killing vectors on the original metric) have a natural action on  $\phi$ :

$$\delta\phi = \xi^i \partial_i \phi + \frac{d-2}{2d} \nabla_i \xi^i \phi, \quad (\text{A.8})$$

where  $\xi^i$  are conformal Killing vectors of the metric  $g_{ij}$  and true Killing vectors of the metric  $\tilde{g}_{ij}$ . Conceptually, the first part of this transformation comes from the ordinary coordinate transformation that is a conformal Killing isometry, while the second part comes from the compensating Weyl transformation needed to bring the metric back to its original form.

What are the isometries of  $\tilde{g}_{ij}$ ? They are easier to recognize by defining a new radial coordinate:

$$\begin{aligned} r_* &\equiv \frac{L^2}{r_s} \ln \frac{r - r_s}{r}, & dr_* &\equiv \frac{L^2 dr}{\Delta}, \\ r &= \frac{r_s}{1 - e^{r_* r_s / L^2}}, & \Delta &= \frac{r_s^2}{4 \sinh^2(r_* r_s / (2L^2))}, \end{aligned} \quad (\text{A.9})$$

such that the line element (A.6) is cast into the form of EAdS<sub>3</sub> in global coordinates

$$\widetilde{ds}^2 = dr_*^2 + \frac{4L^4}{r_s^2} \sinh^2\left(\frac{r_* r_s}{2L^2}\right) (d\theta^2 + \sin^2\theta d\varphi^2). \quad (\text{A.10})$$

The space has 6 Killing vectors: 3 rotations and 3 translations (or “boosts”). As an example, consider the translation that mixes  $r_*$  and  $\theta$ , i.e., the Killing vector (in  $\tilde{g}_{ij}$  sense):

$$J_{r_*\theta} = -2L^2 \cos\theta \partial_{r_*} + r_s \coth\left(\frac{r_* r_s}{2L^2}\right) \sin\theta \partial_\theta. \quad (\text{A.11})$$

From this, we can extract the vector  $\xi^i$ . It is a conformal Killing vector of the original metric (A.2) and its action on the scalar is then given by (A.8)

$$\delta\phi = -2\Delta \cos\theta \partial_r \phi + (r_s - 2r) \partial_\theta (\sin\theta \phi). \quad (\text{A.12})$$

Note that the arbitrary scale  $L$  disappears from  $\delta\phi$ , as it should. Decomposing  $\phi$  into spherical harmonics, it can be shown that this symmetry acts on the various  $\ell$  components as [44]

$$\begin{aligned} \delta\phi_\ell &= f(\ell-1) D_{\ell-1}^+ \phi_{\ell-1} - f(\ell) D_{\ell+1}^- \phi_{\ell+1}, \\ \delta\phi_{\ell-1} &= f(\ell-2) D_{\ell-2}^+ \phi_{\ell-2} - f(\ell-1) D_\ell^- \phi_\ell, \\ f(\ell) &\equiv \left( \frac{4(\ell+1-m)(\ell+1+m)}{3+4\ell(\ell+2)} \right)^{\frac{1}{2}}, \end{aligned} \quad (\text{A.13})$$

where the  $D_\ell^\pm$  operators appearing are precisely the ones in (2.4). We highlight the pair  $\delta\phi_\ell$  and  $\delta\phi_{\ell-1}$  (among the whole tower of  $\delta\phi$ 's) to emphasize that the invariance of the total action comes from the cancellation between the two  $f(\ell-1)$  terms. It is useful in fact to zero in on the pair  $\phi_\ell$  and  $\phi_{\ell-1}$  (ignoring  $\phi_{\ell+1}$  and  $\phi_{\ell-2}$  contributions to  $\delta\phi_\ell$  and  $\delta\phi_{\ell-1}$ ). In spherical harmonic space, the action is a sum over  $\ell$  of the radial integral of  $\phi_\ell^* H_\ell \phi_\ell$ , the variation of the  $\ell^{\text{th}}$  term is

$$\delta(\phi_\ell^* H_\ell \phi_\ell) = f(\ell-1) [\phi_\ell^* H_\ell D_{\ell-1}^+ \phi_{\ell-1} + (D_{\ell-1}^+ \phi_{\ell-1})^* H_\ell \phi_\ell], \quad (\text{A.14})$$

and the variation of the  $(\ell - 1)^{\text{th}}$  term,  $\phi_{\ell-1}^* H_{\ell-1} \phi_{\ell-1}$  is

$$\delta(\phi_{\ell-1}^* H_{\ell-1} \phi_{\ell-1}) = -f(\ell - 1) [\phi_{\ell-1}^* H_{\ell-1} D_{\ell-1}^- \phi_{\ell} + (D_{\ell-1}^- \phi_{\ell})^* H_{\ell-1} \phi_{\ell-1}] . \quad (\text{A.15})$$

One can see these terms cancel, upon recognizing that  $H_{\ell} D_{\ell-1}^+ = D_{\ell-1}^+ H_{\ell-1}$  and  $H_{\ell-1} D_{\ell}^- = D_{\ell}^- H_{\ell}$ , as well as  $D_{\ell}^{-*} \phi_{\ell}^* = \phi_{\ell}^* D_{\ell-1}^+$  and  $D_{\ell-1}^+ \phi_{\ell-1}^* = \phi_{\ell-1}^* D_{\ell}^-$  from integration by parts. The precise form of  $f(\ell - 1)$  does not matter, except one feature: that  $f(\ell - 1)$  implicitly depends on  $m$  and vanishes when  $m = \pm\ell$ . This prevents the generation of terms with  $\ell < |m|$ , and allows the whole tower to terminate at  $\ell = 0$ . The pairing up of  $\phi_{\ell}$  and  $\phi_{\ell-1}$  in this way points to a supersymmetric structure, spelled out in Appendix E. We refer to this symmetry that mixes levels as a *vertical* ladder symmetry.

The charge conservation we use in Section 2 arises from the *horizontal* symmetry transformation  $\delta\phi_{\ell} = Q_{\ell}\phi_{\ell}$ , which does *not* mix different  $\ell$ 's. It arises from the simple observation that  $Q_0 \equiv \Delta\partial_r$  is a symmetry at the  $\ell = 0$  level; symmetries at higher  $\ell$ 's are built up from  $Q_0$  by climbing the ladder.

It is worth noting that this whole geometric perspective has a well-defined flat space  $r_s \rightarrow 0$  limit under which:  $\Omega = L^2/r^2$ ,  $r_* = -L^2/r$ ,  $\widetilde{ds}^2 = dr_*^2 + r_*^2(d\theta^2 + \sin^2\theta d\varphi^2)$ , and  $\delta\phi$  becomes<sup>23</sup>

$$\delta\phi = -2r^2 \cos\theta \partial_r \phi - 2r \partial_{\theta}(\sin\theta \phi) , \quad (\text{A.16})$$

which is relevant for our discussion in Section 5.

In the Kerr case, the static scalar effectively lives in the metric  $g_{ij}^K$  given by

$$ds_K^2 = \frac{\rho^2 - rr_s}{\Delta} \left( dr^2 + \Delta d\theta^2 + \frac{\Delta^2 \sin^2\theta}{\rho^2 - rr_s} d\varphi^2 \right) . \quad (\text{A.17})$$

There does not seem to be a rescaling which brings this to (E)AdS form. However, the action is not much more complicated than that for the Schwarzschild case:

$$S_K = \frac{1}{2} \int d^3x \sqrt{g^K} \phi \square_K \phi = \frac{1}{2} \int d^3x (\sqrt{g} \phi \square \phi) - \frac{a^2 \sin\theta}{\Delta} \phi \partial_{\varphi}^2 \phi , \quad (\text{A.18})$$

where  $g_{ij}$  and  $\square$  on the right hand side refers to the metric in eq. (A.2). Suppose we apply the same rescaling as before:  $\widetilde{g}_{ij} = (L^4/\Delta^2)g_{ij}$  and  $\widetilde{\phi} = (\sqrt{\Delta}/L)\phi$ , then

$$S_K = \frac{1}{2} \int d^3x \sqrt{g^K} \phi \square_K \phi = \frac{1}{2} \int d^3x \sqrt{\widetilde{g}} \widetilde{\phi} \left( \widetilde{\square} + \frac{r_s^2}{4L^4} - \frac{a^2}{L^4} \partial_{\varphi}^2 \right) \widetilde{\phi} . \quad (\text{A.19})$$

Separating the  $\partial_{\varphi}^2$  term from the other derivatives might seem strange, but the point is that in spherical harmonic space,  $\partial_{\varphi}^2 = -m^2$ , and so the Kerr geometry merely modifies the mass term for  $\widetilde{\phi}$ . Because of this, a scalar in the Kerr background has the same ladder structure (with modified  $D^{\pm}$ 's) as a scalar in the Schwarzschild background.

The geometric view of the ladder symmetries can be extended beyond spin 0, as well as non-vanishing frequency. This will be discussed in a separate paper.

<sup>23</sup>In fact, in this limit the symmetry (A.16) is nothing but a special conformal transformation (5.3). It would be very interesting to understand in more detail the relation between these flat space conformal symmetries and the conformal Killing symmetries in the Schwarzschild background.

## B Pöschl–Teller Potential

We can understand the simplicity of the static scalar in Schwarzschild by relating it to the well-known class of Pöschl–Teller potentials [58]. Starting from the equation of motion for the scalar in Schwarzschild in the form (2.3)

$$\Delta(\partial_r(\Delta\partial_r) - \ell(\ell + 1))\phi_\ell = 0, \quad (\text{B.1})$$

we make the change of variables  $dz_\star = r_s\Delta^{-1}dr$  which implies

$$z_\star = 2 \operatorname{arctanh}\left(\frac{r_s}{r_s - 2r}\right). \quad (\text{B.2})$$

After this change of variables, the equation (B.1) takes the form of a Schrödinger equation with a potential

$$\left(\partial_{z_\star}^2 - \frac{\ell(\ell + 1)}{4} \operatorname{csch}^2\left(\frac{z_\star}{2}\right)\right)\phi_\ell = 0, \quad (\text{B.3})$$

which is precisely of the Pöschl–Teller form.<sup>24</sup> This class of potentials has a number of very interesting features. Famously, for  $\ell \in \mathbb{Z}$ , scattering in the Pöschl–Teller potential is reflectionless—a wave sent in from one side of the potential barrier passes through the barrier with no reflected amplitude. The vanishing of a black hole’s tidal response is essentially a zero-frequency version of this reflectionless property: the physically relevant solution has only a single fall-off in both asymptotic regions. Interestingly the Pöschl–Teller potential also plays a prominent role in discussions of the transparency of odd-dimensional de Sitter spaces [44, 60], revealing a surprising mathematical connection between black hole physics and de Sitter space.<sup>25</sup>

The transformation to the  $z_\star$  variable also makes manifest some features of the solutions. By inspection it is clear the two  $\ell = 0$  solutions are  $\phi_0 \sim \text{constant}$  and  $\phi_0 \sim z_\star$ , valid everywhere. Transforming back to the  $r$  coordinate, the  $z_\star$  solution is the one that diverges at the horizon (which is mapped to  $z_\star = -\infty$ ).

The Pöschl–Teller potential has a well-studied connection to supersymmetric quantum mechanics (which we elaborate on in Appendix E), where the ladder operators (2.4) are essentially the supercharges. The potential is in particular an example of a shape-invariant potential, since potentials with consecutive integer values of  $\ell$  are superpartners [61].

Here we have shown how the scalar in Schwarzschild can be cast in the Pöschl–Teller form, but the more general Teukolsky equation (4.1) can also be put in a generalized Pöschl–Teller form. This class of potentials has an algebraic structure equivalent to the one that we have discussed [62–64].

---

<sup>24</sup>Note that often the Pöschl–Teller potential is written in terms of  $\operatorname{sech}$  rather than  $\operatorname{csch}$ , but the two forms can be transformed into each other [59]. The  $\operatorname{csch}$  structure appears here because we are choosing  $z_\star$  to cover the region of  $r$  that lies outside the black hole.

<sup>25</sup>In fact, in Appendix A we saw that one can understand the black hole ladder symmetries geometrically in terms of a scalar field propagating in an effective Euclidean AdS geometry, which is simply related by analytic continuation to the de Sitter problem.

## C Ladder in Regge–Wheeler and Zerilli

In the main text we argued for the vanishing of Love numbers using the symmetries of the Teukolsky equation. One might wonder, however, if it is possible to see the phenomenon directly at the level of metric perturbations. Here we show that in Schwarzschild background a similar structure is present in the Regge–Wheeler and Zerilli equations. We start by considering the Regge–Wheeler equation. The action governing the linearized dynamics of odd-parity spin-2 perturbations around Schwarzschild black holes in general relativity is (see [17] for a detailed derivation in general spacetime dimensions)

$$S_{\text{RW}} = \sum_{\ell} \int dt dr_{\star} \left( \frac{1}{2} \dot{\Psi}_{\ell}^2 - \frac{1}{2} \left( \frac{\partial \Psi_{\ell}}{\partial r_{\star}} \right)^2 - \frac{1}{2} V_{\text{RW}}(r) \Psi_{\ell}^2 \right), \quad (\text{C.1})$$

where  $V_{\text{RW}}(r)$  is the Regge–Wheeler potential

$$V_{\text{RW}}(r) = f(r) \left( \frac{\ell(\ell+1)}{r^2} - f'(r) \frac{3}{r} \right), \quad f(r) \equiv 1 - \frac{r_s}{r}, \quad (\text{C.2})$$

and where  $r_{\star}$  is the standard tortoise coordinate, defined as  $dr_{\star} \equiv dr/f(r) = r^2 dr/\Delta$ . Note that  $\Psi_{\ell}$  in (C.1) is not quite the usual Regge–Wheeler variable [65], which is, instead, up to a numerical factor, the time derivative of our  $\Psi_{\ell}$  [17]. In (C.1) we have integrated over the angles to remove the dependence on the spherical harmonics. In the static limit, the Regge–Wheeler equation, obtained from the action (C.1), is

$$\frac{d^2 \Psi_{\ell}}{dr_{\star}^2} - V_{\text{RW}}(r) \Psi_{\ell} = 0. \quad (\text{C.3})$$

One can introduce the following operators

$$D_{\ell}^{+} \equiv -\Delta \partial_r + \frac{\ell^2 + 3}{2(\ell + 1)} r_s - \ell r, \quad (\text{C.4})$$

$$D_{\ell}^{-} \equiv \Delta \partial_r + \frac{\ell^2(r_s - 2r) - 2\ell(r - r_s) + 4r_s}{2\ell}, \quad (\text{C.5})$$

which satisfy

$$D_{\ell+1}^{-} D_{\ell}^{+} - D_{\ell-1}^{+} D_{\ell}^{-} = (2\ell + 1) r_s^2 \left[ \frac{1}{4} - \frac{4}{\ell^2(\ell + 1)^2} \right]. \quad (\text{C.6})$$

It can be shown that the equations of motion (C.3) can be equivalently written as  $r^{-4} H_{\ell} \Psi_{\ell} = 0$ , where

$$H_{\ell} \equiv D_{\ell-1}^{+} D_{\ell}^{-} - \frac{(\ell^2 - 4)^2 r_s^2}{4\ell^2} = D_{\ell+1}^{-} D_{\ell}^{+} - \frac{(\ell^2 + 2\ell - 3)^2 r_s^2}{4(\ell + 1)^2}. \quad (\text{C.7})$$

The algebra of  $D_{\ell}^{\pm}$  with  $H_{\ell}$  is the same as (2.7):  $H_{\ell+1} D_{\ell}^{+} = D_{\ell}^{+} H_{\ell}$  and  $H_{\ell-1} D_{\ell}^{-} = D_{\ell}^{-} H_{\ell}$ , which implies that  $D_{\ell}^{+}$  and  $D_{\ell}^{-}$  are raising and lowering operators in the sense that, if  $\Psi_{\ell}$  solves  $H_{\ell} \Psi_{\ell} = 0$  for some  $\ell$ , then  $H_{\ell+1} (D_{\ell}^{+} \Psi_{\ell}) = 0$  and  $H_{\ell-1} (D_{\ell}^{-} \Psi_{\ell}) = 0$ . One can then repeat the analysis of Section 2. In particular, by direct inspection of the equations of motion for modes with  $\ell = 2$ , we can identify the quantity  $P_2 \equiv r^4 \Delta \partial_r (\frac{\Psi_2}{r^3})$  which is conserved on-shell, i.e.,  $\partial_r P_2 = 0$  if  $H_2 \Psi_2 = 0$ . From the conservation of  $P_2$  at large distances, one can find the large- $r$  behavior of the two independent solutions to the Regge–Wheeler equation (C.3): one is  $\Psi_2^{(1)}(r \rightarrow \infty) \sim r^3$  (which is actually a good solution everywhere and corresponds

to  $P_2 = 0$ ), while the other goes as  $\Psi_2^{(2)}(r \rightarrow \infty) \sim r^{-2}$  (which corresponds to  $P_2 = \text{constant} \neq 0$ ). From the conservation of  $P_2$ , one then infers that only  $\Psi_2^{(1)}$  is regular at the black hole horizon, while  $\Psi_2^{(2)}$  has to diverge logarithmically as  $\ln(1 - r_s/r)$ . One can then generate regular solutions for all  $\ell$ 's by repeated use of the raising operator  $D_\ell^+$ ,  $\Psi_\ell^{\text{reg}} = D_{\ell-1}^+ \cdots D_2^+ \Psi_2^{(1)}$ . As is manifest from the definition (C.4),  $\Psi_\ell^{\text{reg}}$  is a finite polynomial with only positive powers of  $r$ . Then, using the constraint equations to compute the metric fluctuation  $\delta g_{t\phi}$  from the solution  $\Psi_\ell^{\text{reg}}$ , or, alternatively, performing the matching with the point-particle EFT (5.1) [17], one concludes that the odd-type Love numbers vanish for all  $\ell$ 's.

In analogy with the scalar case, conserved quantities at higher  $\ell$  can be obtained by climbing the ladder:

$$\partial_r P_\ell = 0, \quad \text{where} \quad P_\ell \equiv r^4 \Delta \partial_r (r^{-3} D_1^- D_2^- \cdots D_\ell^- \Psi_\ell). \quad (\text{C.8})$$

We stress that  $P_2$ —which was inferred from the structure of the Regge–Wheeler equation for  $\ell = 2$ —could have been equivalently derived from the action (C.1). Indeed, it can be shown that, in the static limit, where one neglects time derivatives acting on  $\Psi$ , the sector with  $\ell = 2$  in (C.1) is invariant, up to total derivative terms, under the transformation  $\delta \Psi_2 = Q_2 \Psi_2$ , where  $Q_2 \equiv r^4 \Delta \partial_r - 3r^3 \Delta$ . Following the Noether procedure, one finds the conserved current  $J_2 = -\frac{1}{2} r^8 \Delta^2 [\partial_r (\frac{\Psi_2}{r^3})]^2 = -\frac{1}{2} P_2^2$ . Thus, the conservation of  $J_2$  ( $\partial_{r_*} J_2 = 0$ ) is equivalent to  $\partial_r P_2 = 0$ . The procedure can be generalized to all  $\ell$ 's. In particular, one can introduce recursively the quantities

$$Q_\ell \equiv D_{\ell-1}^+ Q_{\ell-1} D_\ell^-, \quad (\text{C.9})$$

which can be shown to leave the action (C.1) invariant. The operators (C.9) should be understood as follows: starting from a solution  $\Psi_\ell$  at level  $\ell$ , the chain of lowering operators  $D_3^- \cdots D_\ell^-$  generate a solution at level  $\ell = 2$ ;  $Q_2$  is then used to produce another solution with  $\ell = 2$ ; finally, the raising operators  $D_{\ell-1}^- \cdots D_2^-$  yield a level- $\ell$  solution.

The previous argument has shown that the vanishing of the black hole Love numbers of odd type can be traced back to the existence of symmetries governing the static regime of the Regge–Wheeler equation. Similar considerations can be in principle extended and applied also to the case of the Zerilli equation [66] and the corresponding Love numbers of even type. The logic that we will follow here is, however, slightly different. It is in fact much more convenient to understand the vanishing of the even-type Love numbers by combining the previous result with another property of black hole perturbations in general relativity, which avoids going through the derivation of ladder operators for the Zerilli equation. We shall use instead the existence of a duality symmetry relating perturbations of different parity. This duality, which is the basis of isospectrality of massless spin-2 fields around Schwarzschild black holes in general relativity [67],<sup>26</sup> can be shown to be also responsible for the *equality* of the even and odd Love numbers, at any given  $\ell$  [69]. Combining the equality of the Love numbers, as a result of the duality, with the vanishing of the static response for the Regge–Wheeler equation, which we inferred above from the ladder symmetries, one can then conclude immediately that the Love numbers of even type in general relativity must also vanish.

We conclude by mentioning that similar considerations and results, following from the existence of ladder operators that raise and lower  $\ell$ , straightforwardly apply also to the case of spin-1 perturbations around Schwarzschild black holes [17].

---

<sup>26</sup>For a generalization to the case of partially massless spin-2 fields on Schwarzschild-de Sitter spacetimes, see [68].

## D Ladder in Teukolsky

In the main text, we were able to discern the properties of solutions to the Teukolsky equation by exploiting a ladder structure in spin, which relates solutions of the static spinning Teukolsky equation to its scalar counterpart. However, there is also a more direct route, which is to identify a ladder in  $\ell$  structure in the Teukolsky equation at fixed  $s$ . In this Appendix we exhibit this structure directly.

In order to see this ladder structure, it is convenient to multiply eq. (4.2) by  $-\Delta$ , and rewrite the equation of motion as:

$$H_{\ell s} \psi_\ell^{(s)} = 0, \quad \text{with} \quad H_{\ell s} \equiv -\Delta \tilde{H}_{\ell s}. \quad (\text{D.1})$$

Define the  $\ell$  raising and lowering operators:

$$\begin{aligned} D_{\ell s}^+ &\equiv -\Delta \partial_z + \frac{\ell + s + 1}{2(\ell + 1)} [(\ell + 1)(z_k - 2z) + 2iqz_k], \\ D_{\ell s}^- &\equiv \Delta \partial_z + \frac{\ell - s}{2\ell} [\ell(z_k - 2z) - 2iqz_k]. \end{aligned} \quad (\text{D.2})$$

These operators obey the algebraic relations:

$$\begin{aligned} D_{\ell+1s}^- D_{\ell s}^+ - D_{\ell-1s}^+ D_{\ell s}^- &= \frac{(2\ell + 1)z_k^2}{4} + \frac{q^2 s^2 z_k^2 (2\ell + 1)}{\ell^2 (\ell + 1)^2}, \\ H_{\ell s} &= D_{\ell+1s}^- D_{\ell s}^+ - (\ell + s + 1)(\ell - s + 1) \frac{((\ell + 1)^2 + 4q^2)z_k^2}{4(\ell + 1)^2}, \\ &= D_{\ell-1s}^+ D_{\ell s}^- - (\ell + s)(\ell - s) \frac{(\ell^2 + 4q^2)z_k^2}{4\ell^2}, \end{aligned} \quad (\text{D.3})$$

and as before  $D_{\ell s}^\pm$  act as ladder operators on the Hamiltonian:

$$H_{\ell+1s} D_{\ell s}^+ = D_{\ell s}^+ H_{\ell s}, \quad H_{\ell-1s} D_{\ell s}^- = D_{\ell s}^- H_{\ell s}. \quad (\text{D.4})$$

The appearance of non-local looking terms, factors of  $\ell$  in the denominators of (D.2), should not surprise us. Recall that for  $s = 1, 2$ , the Newman–Penrose variables are first and second derivatives of the gauge fields respectively (thus the Teukolsky equation involves more than two derivatives on the fundamental fields). Ignoring this subtlety, we can still think of the raising and lowering operators as giving us vertical symmetries of the equation of motion: i.e., given  $\psi_\ell^{(s)}$  which solves  $H_{\ell s} \psi_\ell^{(s)} = 0$ ,  $D_{\ell s}^+ \psi_\ell^{(s)}$  satisfies  $H_{\ell+1s} (D_{\ell s}^+ \psi_\ell^{(s)}) = 0$ , and  $D_{\ell s}^- \psi_\ell^{(s)}$  satisfies  $H_{\ell-1s} (D_{\ell s}^- \psi_\ell^{(s)}) = 0$ . The fundamental origin of these symmetries is a bit obscure in this case, because we do not have an action for the Teukolsky equation. Nevertheless, they provide a useful device for manipulating solutions to the equation.

Proceeding in the same formal manner, we can guess the horizontal ladder symmetries. Let us start from the  $\ell = s$  level. It can be shown the corresponding equation of motion

$$\beta^2 H_{ss} \psi_s^{(s)} = 0, \quad \text{with} \quad \beta \equiv \Delta^s e^{-2iq \ln[(z-z_k)/z]}, \quad (\text{D.5})$$

takes the simple form:

$$-\beta \Delta \partial_z (\beta \Delta \partial_z) \psi_s^{(s)} = 0. \quad (\text{D.6})$$

It is easy to see that a symmetry of this equation is

$$\delta\psi_s^{(s)} \equiv Q_s^{(s)}\psi_s^{(s)}, \quad \text{where} \quad Q_s^{(s)} \equiv \beta\Delta\partial_z. \quad (\text{D.7})$$

This transformation is a symmetry in the sense that  $[\beta^2 H_{ss}, Q_s^{(s)}] = 0$ , so that if  $\psi_s^{(s)}$  is a solution, so is  $Q_s^{(s)}\psi_s^{(s)}$ . We can bootstrap to higher  $\ell > s$  by:

$$Q_\ell^{(s)} = D_{\ell-1s}^+ \cdots D_{ss}^+ Q_s^{(s)} D_{s+1s}^- \cdots D_{\ell s}^-. \quad (\text{D.8})$$

Starting from a level  $\ell$  solution  $\psi_\ell^{(s)}$ , the combination  $D_{s+1s}^- \cdots D_{\ell s}^- \psi_\ell^{(s)}$  produces a solution at level  $\ell = s$ . We already know acting on it with  $Q_s^{(s)}$  produces another level  $\ell = s$  solution. Finally raising the result back with  $D_{\ell-1s}^+ \cdots D_{ss}^+$  produces another level  $\ell$  solution. We thus have an equation-of-motion symmetry in the sense that if  $\psi_\ell^{(s)}$  is a solution, so is  $Q_\ell^{(s)}\psi_\ell^{(s)}$ . Note, however, that  $[\beta^2 H_{\ell s}, Q_\ell^{(s)}] \neq 0$ .

Considering  $s = 2$  for instance, the growing solution is  $\psi_2^{(2)} = E^+ E^+ D_1^+ D_0^+ \psi_0^{(0)}$  with  $\psi_0^{(0)} = \text{constant}$ . (Here, the  $D^\pm$ 's are the raising and lowering operators for scalar.) It is not hard to show  $\psi_2^{(2)}$  is a constant too, and thus  $Q_2^{(2)}\psi_2^{(2)} = 0$ . Conversely, the decaying solution is  $\psi_2^{(2)} = E^+ E^+ D_1^+ D_0^+ \psi_0^{(0)}$  with  $\psi_0^{(0)} = e^{2iq \ln[(z-z_k)/z]}$ . It can be checked that  $Q_2^{(2)}\psi_2^{(2)} \neq 0$ . The obvious extensions of these statements to  $\ell > 2$  can be derived by climbing the ladder.

## E Supersymmetric Structure

The  $\ell$ -ladder structure, with the equation of motion operator  $H_\ell$  written as some sort of square, is reminiscent of supersymmetric quantum mechanics. The supersymmetric structure can be made explicit as follows. It is convenient to define  $A_\ell$  and  $A_\ell^\dagger$ :

$$\begin{aligned} A_\ell &\equiv D_\ell^- = \Delta\partial_z + \frac{1}{2}[\ell(z_k - 2z) - 2iqz_k] \\ A_\ell^\dagger &\equiv D_{\ell-1}^+ = -\Delta\partial_z + \frac{1}{2}[\ell(z_k - 2z) + 2iqz_k] . \end{aligned} \quad (\text{E.1})$$

where we have used  $D^\pm$  defined in eq. (3.7). The operator  $A_\ell^\dagger$  is the hermitian conjugate of  $A_\ell$  in the sense that the action can be thought of as  $\int dy \psi_\ell^* H_\ell \psi_\ell$  with  $\partial_y \equiv \Delta\partial_z$ . Noting that

$$A_\ell^\dagger A_\ell = H_\ell + \frac{(\ell^2 + 4q^2)z_k^2}{4} , \quad A_\ell A_\ell^\dagger = H_{\ell-1} + \frac{(\ell^2 + 4q^2)z_k^2}{4} , \quad (\text{E.2})$$

we are motivated to define the operator

$$\mathcal{H}_\ell \equiv \begin{pmatrix} A_\ell^\dagger A_\ell & 0 \\ 0 & A_\ell A_\ell^\dagger \end{pmatrix} , \quad (\text{E.3})$$

such that its eigenvectors  $\Psi_\ell$  satisfying

$$\mathcal{H}_\ell \Psi_\ell = E_\ell \Psi_\ell , \quad E_\ell \equiv \frac{(\ell^2 + 4q^2)z_k^2}{4} , \quad (\text{E.4})$$

consist precisely of the  $\psi_\ell$  and  $\psi_{\ell-1}$  solutions packaged together. In particular, it's simple to check that two possible solutions are:

$$\Psi_\ell = \begin{pmatrix} \psi_\ell \\ 0 \end{pmatrix} , \quad \Psi_\ell = \begin{pmatrix} 0 \\ A_\ell \psi_\ell \end{pmatrix} , \quad (\text{E.5})$$

where  $\psi_\ell$  solves  $H_\ell \psi_\ell = 0$  and  $A_\ell \psi_\ell$  solves  $H_{\ell-1}(A_\ell \psi_\ell) = 0$ . The operator  $\mathcal{H}_\ell$  can be thought of as coming from an anti-commutator of ‘‘supercharges’’

$$\mathcal{H}_\ell = \{\mathcal{Q}_\ell, \mathcal{Q}_\ell^\dagger\} , \quad \text{with} \quad \mathcal{Q}_\ell \equiv \begin{pmatrix} 0 & 0 \\ A_\ell & 0 \end{pmatrix} , \quad (\text{E.6})$$

which satisfy  $[\mathcal{H}_\ell, \mathcal{Q}_\ell] = [\mathcal{H}_\ell, \mathcal{Q}_\ell^\dagger] = 0$ . From this perspective, the Hamiltonian at level  $\ell$  is the supersymmetric partner of the Hamiltonian at level  $\ell + 1$  and so on (indicating that these potentials are shape invariant). We can then understand the simplicity of the tower of potentials from the fact that the  $\ell = 0$  potential is just a constant.<sup>27</sup> We have spelled out the supersymmetric structure for the scalar in a Kerr background. The same can be done for the Teukolsky equation, with conjugation involving  $s \rightarrow -s$ . Incidentally, in the context of metric perturbations around a Schwarzschild background, there is a different sort of supersymmetry that has been discussed in the literature, one that relates parity even and odd modes, obeying the Zerilli and Regge–Wheeler equations. The Teukolsky equation provides an economical description, with the even and odd modes described by the real and imaginary parts of the Newman–Penrose variable, making the symmetry between them manifest.

<sup>27</sup>This supersymmetric structure at  $q = 0$  is just the standard structure of the Pöschl–Teller potential, discussed in Appendix B.

## F Hypergeometric Identities

In uncovering the ladder structures in black hole perturbations, we were aided by the known static solutions which are in the form of the (Gauss) hypergeometric function  ${}_2F_1$ .<sup>28</sup> Neighboring hypergeometric functions, with arguments offset by unity, are known to be related. The identities we find the most helpful in revealing the  $\ell$ -ladder are:

$$\begin{aligned} \frac{d}{dz} {}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] &= \frac{a[(c-a-1) - (b-a-1)(1-z)]}{z(b-a-1)(1-z)} {}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] \\ &\quad - \frac{a(c-b)}{z(b-a-1)(1-z)} {}_2F_1 \left[ \begin{matrix} a+1, & b-1 \\ & c \end{matrix} \middle| z \right], \end{aligned} \quad (\text{F.1})$$

$$\begin{aligned} \frac{d}{dz} {}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] &= \frac{b[(c-b-1) - (a-b-1)(1-z)]}{z(a-b-1)(1-z)} {}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] \\ &\quad - \frac{b(c-a)}{z(a-b-1)(1-z)} {}_2F_1 \left[ \begin{matrix} a-1, & b+1 \\ & c \end{matrix} \middle| z \right]. \end{aligned} \quad (\text{F.2})$$

Note that the growing branch of the static solutions to the Teukolsky equation takes the form of

$$\psi_\ell^{(s)} = {}_2F_1 \left[ \begin{matrix} -\ell + s, & \ell + s + 1 \\ & 1 + s + 2iq \end{matrix} \middle| z \right]. \quad (\text{F.3})$$

The identities (F.1) and (F.2) can be derived from known identities in the literature:

$$\begin{aligned} (c-a-1) {}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] &= (b-a-1)(1-z) {}_2F_1 \left[ \begin{matrix} a+1, & b \\ & c \end{matrix} \middle| z \right] \\ &\quad + (c-b) {}_2F_1 \left[ \begin{matrix} a+1, & b-1 \\ & c \end{matrix} \middle| z \right], \end{aligned} \quad (\text{F.4})$$

$$\begin{aligned} [(a-b)(a-b-1)(1-z) + b(c-b-1)] {}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] &= a(a-b-1)(1-z) {}_2F_1 \left[ \begin{matrix} a+1, & b \\ & c \end{matrix} \middle| z \right] \\ &\quad + b(c-a) {}_2F_1 \left[ \begin{matrix} a-1, & b+1 \\ & c \end{matrix} \middle| z \right], \end{aligned} \quad (\text{F.5})$$

$${}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] = {}_2F_1 \left[ \begin{matrix} a+1, & b \\ & c \end{matrix} \middle| z \right] - \frac{bz}{c} {}_2F_1 \left[ \begin{matrix} a+1, & b+1 \\ & c+1 \end{matrix} \middle| z \right], \quad (\text{F.6})$$

$$\frac{d}{dz} {}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] = \frac{ab}{c} {}_2F_1 \left[ \begin{matrix} a+1, & b+1 \\ & c+1 \end{matrix} \middle| z \right]. \quad (\text{F.7})$$

Equations (F.4) and (F.5), generalizing the Gauss contiguous relations, can be found in 1.27 and 2.1 of [70]. Equation (F.6) can be found on p. 43 of [71]. Equation (F.7) is well known and can be found on p. 41 of [72].

The identities useful for revealing the  $s$ -ladder are eq. (F.7) and:

$$\frac{d}{dz} \left( z^{c-1}(1-z)^{a+b-c} {}_2F_1 \left[ \begin{matrix} a, & b \\ & c \end{matrix} \middle| z \right] \right) = (c-1)z^{c-2}(1-z)^{a+b-c-1} {}_2F_1 \left[ \begin{matrix} a-1, & b-1 \\ & c-1 \end{matrix} \middle| z \right], \quad (\text{F.8})$$

from p. 42 of [72].

---

<sup>28</sup>For a review of some of the elementary properties of hypergeometric functions, see Appendix B of [17].

## G Scalar-Gauss–Bonnet Theories

In this appendix, we consider the case of a scalar field coupled to the Gauss–Bonnet curvature invariant  $R_{\text{GB}}^2 \equiv R^{\mu\nu\rho\sigma}R_{\mu\nu\rho\sigma} - 4R^{\mu\nu}R_{\mu\nu} + R^2$ . Such a coupling makes it possible to evade the no-hair theorem, sourcing a nonzero scalar profile around a black hole with a decaying  $1/r$  tail at large  $r$  [35]. This example is an interesting laboratory that allows us to test the consequences of our ladder symmetries for Love numbers and hair in an explicit microscopic model.

Let us start by adding a linear coupling of the form  $\mathcal{L}_{\text{GB}} \sim \alpha\phi R_{\text{GB}}^2$  to the ultraviolet theory of a scalar field coupled to gravity. For simplicity, we shall treat the coupling  $\alpha$  in  $\mathcal{L}_{\text{GB}}$  perturbatively and look for a solution to the field equations in powers of  $\alpha$ . At linear order in  $\alpha$ , we can ignore the backreaction of the metric (corrections to the metric enter at order  $\alpha^2$ ) and study the scalar’s dynamics on a fixed Schwarzschild geometry. The scalar’s equation of motion is

$$\square\phi = -\alpha R_{\text{GB}}^2. \tag{G.1}$$

After decomposing  $\phi$  in spherical harmonics, the Gauss–Bonnet term in (G.1) plays the role of a source for the  $\ell = 0$  mode, while the equations for the higher multipoles are unchanged at this order in  $\alpha$  with respect to (2.2). In the static limit, the solution for the monopole that is regular at the horizon is

$$\phi_{\ell=0}(r) = \phi_\infty + \alpha \left( \frac{4}{rr_s} + \frac{2}{r^2} + \frac{4r_s}{3r^3} \right) + \mathcal{O}(\alpha^2), \tag{G.2}$$

where  $\phi_\infty$  is a constant, corresponding to the value of  $\phi(r)$  at  $r = \infty$ . The term  $4\alpha/(rr_s)$  is the well-known fall-off that corresponds to the scalar hair generated by the Gauss–Bonnet coupling [35]. On the other hand, it is clear from (G.2) that there is no static response induced by a constant tidal field, which would be proportional to  $\phi_\infty$ .

The above calculations can be understood in terms of (exact and broken) symmetries in the following way. The point is that the coupling to the Gauss–Bonnet operator is responsible for an explicit breaking of the  $Q_0$  symmetry, so that hair is no longer forbidden. At the level of the point-particle EFT (5.1), the breaking of  $Q_0$  implies that the tadpole  $g\phi$  coupling can appear in the infrared description and should therefore be included in the EFT. In particular—combined with the spherical symmetry of the system—it implies that if we expand  $g\phi$  in spherical harmonics only the monopole is allowed by symmetries. And indeed this is precisely what the microscopic calculation shows: the Gauss–Bonnet coupling allows for nonzero monopole hair (but not for higher-multipole-type of hair). Even though the coupling to the Gauss–Bonnet operator breaks  $Q_0$  explicitly, one finds that  $Q_{\ell \geq 1}$  are still symmetries of the theory, up to linear order in  $\alpha$ . This is enough to ensure the vanishing of the Love numbers with  $\ell \geq 1$ .<sup>29</sup> On the other hand, the absence of a static response for  $\ell = 0$  is enforced by the shift symmetry,  $\phi \mapsto \phi + c$ , which is an exact invariance of the

<sup>29</sup>Note that this result follows from knowing the exact microscopic dynamics. Formulating a similar statement directly at the level of the EFT (5.1) seems more subtle. Indeed, the breaking of  $Q_0$  allows in general all the  $\lambda_\ell$ ’s in (5.1). Presumably, understanding how to implement a subset of our symmetries (at selected  $\ell$ ’s) at the level of the worldline action (5.1), as mentioned in Section 5, would allow us to use the remaining symmetries to constrain (at least some of) the other couplings  $\lambda_\ell$ . Nevertheless, the vanishing of the monopole Love number can at least be understood in terms of the shift symmetry  $\phi \mapsto \phi + c$ , which clearly forbids  $\phi^2$  in (5.1), providing in the infrared a symmetry explanation of why  $\lambda_0 = 0$ .

theory in the presence of a linear coupling,  $\mathcal{L}_{\text{GB}} \sim \alpha\phi R_{\text{GB}}^2$ . As a result, all the black hole’s scalar Love numbers vanish.

It is instructive to contrast this result with the case of a scalar field coupled quadratically to the Gauss–Bonnet operator, i.e.,  $\mathcal{L}_{\text{GB}} \sim \frac{1}{2}\alpha\phi^2 R_{\text{GB}}^2$  [39]. In contrast to the previous example, the shift symmetry is now broken explicitly in the ultraviolet theory. Therefore, there is no reason now for the monopole Love number to vanish and the quadratic worldline operator  $\phi^2$  not to be allowed in the point-particle effective description. This is indeed what one finds from an explicit calculation in the microscopic theory and after matching with the EFT. To show this, we again solve the scalar equation of motion, which now reads

$$\square\phi + \alpha\phi R_{\text{GB}}^2 = 0, \tag{G.3}$$

perturbatively in  $\alpha$ . Focusing for simplicity on the monopole  $\ell = 0$ , the solution to (G.3) that is regular at  $r = r_s$  is now

$$\phi_{\ell=0}(r) = \phi_\infty \left( 1 + \alpha \left( \frac{4}{rr_s} + \frac{2}{r^2} + \frac{4r_s}{3r^3} \right) + \mathcal{O}(\alpha^2) \right), \tag{G.4}$$

where we have discarded terms that are order  $\alpha^2$  or higher. The constant  $\phi_\infty$  now multiplies the whole solution—this follows from the fact that, as opposed to (G.1), eq. (G.3) is homogeneous in  $\phi$ —and one can genuinely interpret the  $1/r$  tail as the static response to the monopole tidal field profile  $\phi_\infty$ . According to [39], this theory can even generate hair, which from the EFT perspective is described by a linear  $\phi$  coupling to the worldline. This represents a spontaneous breaking of the discrete  $\phi \rightarrow -\phi$  symmetry that is present in the microscopic theory. This so called scalarization phenomenon seems to occur only if  $\alpha$  is sufficiently large.



Our exploration of the black hole ladders—black hole arpeggios as it were—was partly inspired by Bach’s music. A ladder structure can be discerned for instance in the celebrated prelude of Cello Suite No. 1 in G major.



## References

- [1] S. Chandrasekhar, “The mathematical theory of black holes,” in *Oxford, UK: Clarendon (1992) 646 p.*, *Oxford, UK: Clarendon (1985) 646 P.* 1985.
- [2] W. H. Press, “Time Evolution of a Rotating Black Hole Immersed in a Static Scalar Field,” *ApJ* **175** (July, 1972) 243.
- [3] K. Martel and E. Poisson, “Gravitational perturbations of the Schwarzschild spacetime: A Practical covariant and gauge-invariant formalism,” *Phys. Rev. D* **71** (2005) 104003, [arXiv:gr-qc/0502028](#).
- [4] H. Fang and G. Lovelace, “Tidal coupling of a Schwarzschild black hole and circularly orbiting moon,” *Phys. Rev. D* **72** (2005) 124016, [arXiv:gr-qc/0505156](#).
- [5] T. Damour and O. M. Lecian, “On the gravitational polarizability of black holes,” *Phys. Rev.* **D80** (2009) 044017, [arXiv:0906.3003 \[gr-qc\]](#).
- [6] T. Damour and A. Nagar, “Relativistic tidal properties of neutron stars,” *Phys. Rev. D* **80** (2009) 084035, [arXiv:0906.0096 \[gr-qc\]](#).
- [7] T. Binnington and E. Poisson, “Relativistic theory of tidal Love numbers,” *Phys. Rev.* **D80** (2009) 084018, [arXiv:0906.1366 \[gr-qc\]](#).
- [8] B. Kol and M. Smolkin, “Black hole stereotyping: Induced gravito-static polarization,” *JHEP* **02** (2012) 010, [arXiv:1110.3764 \[hep-th\]](#).
- [9] P. Landry and E. Poisson, “Gravitomagnetic response of an irrotational body to an applied tidal field,” *Phys. Rev. D* **91** no. 10, (2015) 104026, [arXiv:1504.06606 \[gr-qc\]](#).
- [10] P. Landry and E. Poisson, “Tidal deformation of a slowly rotating material body. External metric,” *Phys. Rev. D* **91** (2015) 104018, [arXiv:1503.07366 \[gr-qc\]](#).
- [11] N. Gürlebeck, “No-hair theorem for Black Holes in Astrophysical Environments,” *Phys. Rev. Lett.* **114** no. 15, (2015) 151102, [arXiv:1503.03240 \[gr-qc\]](#).
- [12] E. Poisson, “Gravitomagnetic Love tensor of a slowly rotating body: post-Newtonian theory,” [arXiv:2007.01678 \[gr-qc\]](#).
- [13] A. Le Tiec and M. Casals, “Spinning Black Holes Fall in Love,” [arXiv:2007.00214 \[gr-qc\]](#).
- [14] A. Le Tiec, M. Casals, and E. Franzin, “Tidal Love Numbers of Kerr Black Holes,” *Phys. Rev. D* **103** (2021) 084021, [arXiv:2010.15795 \[gr-qc\]](#).
- [15] H. S. Chia, “Tidal Deformation and Dissipation of Rotating Black Holes,” [arXiv:2010.07300 \[gr-qc\]](#).
- [16] W. D. Goldberger, J. Li, and I. Z. Rothstein, “Non-conservative effects on Spinning Black Holes from World-Line Effective Field Theory,” [arXiv:2012.14869 \[hep-th\]](#).
- [17] L. Hui, A. Joyce, R. Penco, L. Santoni, and A. R. Solomon, “Static response and Love numbers of Schwarzschild black holes,” *JCAP* **04** (2021) 052, [arXiv:2010.00593 \[hep-th\]](#).
- [18] P. Charalambous, S. Dubovsky, and M. M. Ivanov, “On the Vanishing of Love Numbers for Kerr Black Holes,” [arXiv:2102.08917 \[hep-th\]](#).

- [19] S. E. Gralla, “On the Ambiguity in Relativistic Tidal Deformability,” *Class. Quant. Grav.* **35** no. 8, (2018) 085002, [arXiv:1710.11096 \[gr-qc\]](#).
- [20] W. D. Goldberger and I. Z. Rothstein, “An Effective field theory of gravity for extended objects,” *Phys. Rev. D* **73** (2006) 104029, [arXiv:hep-th/0409156](#).
- [21] W. Israel, “Event horizons in static vacuum space-times,” *Phys. Rev.* **164** (1967) 1776–1779.
- [22] B. Carter, “Global structure of the Kerr family of gravitational fields,” *Phys. Rev.* **174** (1968) 1559–1571.
- [23] B. Carter, “Axisymmetric Black Hole Has Only Two Degrees of Freedom,” *Phys. Rev. Lett.* **26** (1971) 331–333.
- [24] R. M. Wald, “Final states of gravitational collapse,” *Phys. Rev. Lett.* **26** (1971) 1653–1655.
- [25] J. B. Hartle, “Long-range neutrino forces exerted by kerr black holes,” *Phys. Rev. D* **3** (1971) 2938–2940.
- [26] J. D. Bekenstein, “Nonexistence of baryon number for static black holes,” *Phys. Rev. D* **5** (1972) 1239–1246.
- [27] E. D. Fackerell and J. R. Ipser, “Weak electromagnetic fields around a rotating black hole,” *Phys. Rev. D* **5** (1972) 2455–2458.
- [28] R. H. Price, “Nonspherical Perturbations of Relativistic Gravitational Collapse. II. Integer-Spin, Zero-Rest-Mass Fields,” *Phys. Rev. D* **5** (1972) 2439–2454.
- [29] J. Bekenstein, “Novel “no-scalar-hair” theorem for black holes,” *Phys. Rev. D* **51** no. 12, (1995) 6608.
- [30] L. Hui and A. Nicolis, “No-Hair Theorem for the Galileon,” *Phys. Rev. Lett.* **110** (2013) 241104, [arXiv:1202.1296 \[hep-th\]](#).
- [31] T. Jacobson, “Primordial black hole evolution in tensor scalar cosmology,” *Phys. Rev. Lett.* **83** (1999) 2699–2702, [arXiv:astro-ph/9905303](#).
- [32] K.-M. Lee, V. P. Nair, and E. J. Weinberg, “Black holes in magnetic monopoles,” *Phys. Rev. D* **45** (1992) 2751–2761, [arXiv:hep-th/9112008](#).
- [33] E. J. Weinberg, “Black holes with hair,” *NATO Sci. Ser. II* **60** (2002) 523–544, [arXiv:gr-qc/0106030](#).
- [34] S. Alexander and N. Yunes, “Chern-Simons Modified General Relativity,” *Phys. Rept.* **480** (2009) 1–55, [arXiv:0907.2562 \[hep-th\]](#).
- [35] T. P. Sotiriou and S.-Y. Zhou, “Black hole hair in generalized scalar-tensor gravity,” *Phys. Rev. Lett.* **112** (2014) 251102, [arXiv:1312.3622 \[gr-qc\]](#).
- [36] E. Babichev and C. Charmousis, “Dressing a black hole with a time-dependent Galileon,” *JHEP* **08** (2014) 106, [arXiv:1312.3204 \[gr-qc\]](#).
- [37] T. P. Sotiriou and S.-Y. Zhou, “Black hole hair in generalized scalar-tensor gravity: An explicit example,” *Phys. Rev. D* **90** (2014) 124063, [arXiv:1408.1698 \[gr-qc\]](#).
- [38] C. A. R. Herdeiro and E. Radu, “Kerr black holes with scalar hair,” *Phys. Rev. Lett.* **112** (2014) 221101, [arXiv:1403.2757 \[gr-qc\]](#).
- [39] H. O. Silva, J. Sakstein, L. Gualtieri, T. P. Sotiriou, and E. Berti, “Spontaneous scalarization of black holes and compact stars from a Gauss-Bonnet coupling,” *Phys. Rev. Lett.* **120** no. 13, (2018) 131104, [arXiv:1711.02080 \[gr-qc\]](#).

- [40] L. K. Wong, A.-C. Davis, and R. Gregory, “Effective field theory for black holes with induced scalar charges,” *Phys. Rev. D* **100** no. 2, (2019) 024010, [arXiv:1903.07080 \[hep-th\]](#).
- [41] K. Clough, P. G. Ferreira, and M. Lagos, “Growth of massive scalar hair around a Schwarzschild black hole,” *Phys. Rev. D* **100** no. 6, (2019) 063014, [arXiv:1904.12783 \[gr-qc\]](#).
- [42] L. Hui, D. Kabat, X. Li, L. Santoni, and S. S. Wong, “Black Hole Hair from Scalar Dark Matter,” *JCAP* **06** (2019) 038, [arXiv:1904.12803 \[gr-qc\]](#).
- [43] P. Charalambous, S. Dubovsky, and M. M. Ivanov, “Hidden Symmetry of Vanishing Love,” [arXiv:2103.01234 \[hep-th\]](#).
- [44] G. Compton and I. A. Morrison, “Hidden symmetries for transparent de Sitter space,” *Class. Quant. Grav.* **37** no. 12, (2020) 125001, [arXiv:2003.08023 \[gr-qc\]](#).
- [45] R. P. Kerr, “Gravitational field of a spinning mass as an example of algebraically special metrics,” *Phys. Rev. Lett.* **11** (1963) 237–238.
- [46] W. D. Goldberger and I. Z. Rothstein, “Dissipative effects in the worldline approach to black hole dynamics,” *Phys. Rev. D* **73** (2006) 104030, [arXiv:hep-th/0511133](#).
- [47] S. Teukolsky, “Rotating black holes - separable wave equations for gravitational and electromagnetic perturbations,” *Phys. Rev. Lett.* **29** (1972) 1114–1118.
- [48] S. A. Teukolsky, “Perturbations of a rotating black hole. 1. Fundamental equations for gravitational electromagnetic and neutrino field perturbations,” *Astrophys. J.* **185** (1973) 635–647.
- [49] I. Z. Rothstein, “Progress in effective field theory approach to the binary inspiral problem,” *Gen. Rel. Grav.* **46** (2014) 1726.
- [50] R. A. Porto, “The Tune of Love and the Nature(ness) of Spacetime,” *Fortsch. Phys.* **64** no. 10, (2016) 723–729, [arXiv:1606.08895 \[gr-qc\]](#).
- [51] R. F. Penna, “Near-horizon Carroll symmetry and black hole Love numbers,” [arXiv:1812.05643 \[hep-th\]](#).
- [52] A. Ross, “Multipole expansion at the level of the action,” *Phys. Rev. D* **85** (2012) 125033, [arXiv:1202.4750 \[gr-qc\]](#).
- [53] R. A. Porto, “Post-Newtonian corrections to the motion of spinning bodies in NRGR,” *Phys.Rev.* **D73** (2006) 104031, [arXiv:gr-qc/0511061 \[gr-qc\]](#).
- [54] L. V. Delacrétaz, S. Endlich, A. Monin, R. Penco, and F. Riva, “(Re-)Inventing the Relativistic Wheel: Gravity, Cosets, and Spinning Objects,” *JHEP* **1411** (2014) 008, [arXiv:1405.7384 \[hep-th\]](#).
- [55] J. D. Bekenstein, “Transcendence of the law of baryon-number conservation in black hole physics,” *Phys. Rev. Lett.* **28** (1972) 452–455.
- [56] V. Cardoso, E. Franzin, A. Maselli, P. Pani, and G. Raposo, “Testing strong-field gravity with tidal Love numbers,” *Phys. Rev. D* **95** no. 8, (2017) 084014, [arXiv:1701.01116 \[gr-qc\]](#). [Addendum: *Phys.Rev.D* 95, 089901 (2017)].
- [57] S. Bertini, S. L. Cacciatori, and D. Klemm, “Conformal structure of the Schwarzschild black hole,” *Phys. Rev. D* **85** (2012) 064018, [arXiv:1106.0999 \[hep-th\]](#).

- [58] G. Pöschl and E. Teller, “Bemerkungen zur quantenmechanik des anharmonischen oszillators,” *Zeitschrift für Physik* **83** no. 3, (1933) 143–151.
- [59] D. Anninos, S. A. Hartnoll, and D. M. Hofman, “Static Patch Solipsism: Conformal Symmetry of the de Sitter Worldline,” *Class. Quant. Grav.* **29** (2012) 075002, [arXiv:1109.4942 \[hep-th\]](#).
- [60] P. Lagogiannis, A. Maloney, and Y. Wang, “Odd-dimensional de Sitter Space is Transparent,” [arXiv:1106.2846 \[hep-th\]](#).
- [61] F. Cooper, A. Khare, and U. Sukhatme, “Supersymmetry and quantum mechanics,” *Phys. Rept.* **251** (1995) 267–385, [arXiv:hep-th/9405029 \[hep-th\]](#).
- [62] A. O. Barut, A. Inomata, and R. Wilson, “Algebraic treatment of second poschl-teller, morse-rosen and eckart equations,” *Journal of Physics A: Mathematical and General* **20** no. 13, (Sep, 1987) 4083–4096.
- [63] J. Wu, Y. Alhassid, and F. Gürsey, “Group theory approach to scattering. IV. Solvable potentials associated with  $SO(2, 2)$ ,” *Annals Phys.* **196** no. 1, (1989) 163–181.
- [64] J. Wu and Y. Alhassid, “The potential group approach and hypergeometric differential equations,” *Journal of Mathematical Physics* **31** no. 3, (Mar., 1990) 557–562.
- [65] T. Regge and J. A. Wheeler, “Stability of a Schwarzschild singularity,” *Phys. Rev.* **108** (1957) 1063–1069.
- [66] F. J. Zerilli, “Gravitational field of a particle falling in a schwarzschild geometry analyzed in tensor harmonics,” *Phys. Rev.* **D2** (1970) 2141–2160.
- [67] S. Chandrasekhar, “On the equations governing the perturbations of the schwarzschild black hole,” *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* **343** no. 1634, (1975) 289–298. <http://www.jstor.org/stable/78870>.
- [68] R. A. Rosen and L. Santoni, “Black hole perturbations of massive and partially massless spin-2 fields in (anti) de Sitter spacetime,” *JHEP* **03** (2021) 139, [arXiv:2010.00595 \[hep-th\]](#).
- [69] L. Hui, A. Joyce, R. Penco, L. Santoni, and A. Solomon, *To appear*.
- [70] M. A. Rakha, A. K. Rathie, and P. Chopra, “On some new contiguous relations for the Gauss hypergeometric function with applications,” *Computers and Mathematics with Applications* **61** (2011) 620–629.
- [71] J. Stalker, “Complex Analysis - Fundamentals of the Classical Theory of Functions,” *Complex Analysis - Fundamentals of the Classical Theory of Functions*, Birkhauser (1998) .
- [72] W. Magnus, F. Oberhettinger, and R. P. Soni, “Formulas and Theorems for the Special Functions of Mathematical Physics,” *Formulas and Theorems for the Special Functions of Mathematical Physics*, Springer-Verlag (1966) .