

Effects of coalescence and isospin symmetry on the freezeout of light nuclei and their anti-particles

M. Waqas^{1,*}, G. X. Peng^{1,2,3 †}, Fu-Hu Liu^{4,5} (fuhuliu@163.com; fuhuliu@sxu.edu.cn),
Z. Wazir⁶ (zwazir@gudgk.edu.pk)

¹ School of Nuclear Science and Technology, University of Chinese Academy of Sciences, Beijing 100049, China,

² Theoretical Physics Center for Science Facilities, Institute of High Energy Physics, Beijing 100049, China,

³ Synergetic Innovation Center for Quantum Effects & Applications, Hunan Normal University, Changsha 410081, China,

⁴ Institute of Theoretical Physics & State Key Laboratory of Quantum Optics and Quantum Optics Devices, Shanxi University, Taiyuan, 030006, China,

⁵ Collaborative Innovation Center of Extreme Optics, Shanxi University, Taiyuan, 030006, China,

⁶ Department of physics, Ghazi University, Dera Ghazi Khan, Pakistan.

Abstract: The transverse momentum spectra of light nuclei (deuteron, triton and helion) produced in various centrality intervals in Gold-Gold (Au-Au), Lead-Lead (Pb-Pb) and proton-Lead (p-Pb) collisions, as well as in inelastic (INEL) proton-proton (pp) collisions are analyzed by the blast wave model with Boltzmann Gibbs statistics. The model results are nearly in agreement with the experimental data measured by STAR and ALICE Collaborations in special transverse momentum ranges. We extracted the bulk properties in terms of kinetic freezeout temperature, transverse flow velocity and freezeout volume. It is observed that deuteron and anti-deuteron freezeout later than triton and helion as well as their anti-particles due to its smaller mass, while helion and triton, and anti-helion and anti-triton freezeout at the same time due to isospin symmetry at higher energies. It is also observed that light nuclei freezeout earlier than their anti-nuclei due to the large coalescence of nucleons for light nuclei compared to their anti-nuclei. The kinetic freezeout temperature, transverse flow velocity and kinetic freezeout volume decrease from central to peripheral collisions. Furthermore, the transverse flow velocity depends on mass of the particle which decreases with increasing the mass of the particle.

Keywords: light nuclei, kinetic freezeout temperature, transverse flow velocity, freezeout volume, coalescence, isospin symmetry.

PACS: 12.40.Ee, 13.85.Hd, 25.75.Ag, 25.75.Dw, 24.10.Pa

1 Introduction

A new form of matter, the so called quark-gluon plasma (QGP) is produced under the high temperatures and energy densities in relativistic heavy ion collisions. This matter is formed in the early stage of collisions that survives for a very short period of time ($\sim 7-10$ fm/c), after which the QGP gets transformed rapidly to a system of hadron gas. Due to multi-partonic interactions throughout the evolution time in the collisions, the information about the initial condition of the system get mostly lost. The final state behavior of such colliding system can be attained from the measurement of the number as well as the identity of the produced particles

along with their energy and momentum spectra. The final state information is very useful to understand the particle production mechanisms and the nature of the matter created in these high energy collisions.

Temperature is one of the most crucial factor in subatomic physics. There are different types of temperatures present in literature [1–5]. Chemical freezeout temperature (T_{ch}), which describes the excitation degree of interacting system at the stage of chemical freezeout. At chemical freezeout the chemical components (relative fraction) of the particles are invariant. The excitation degree of the interacting system at the stage of thermal or kinetic freezeout is described by the kinetic freezeout temperature (T_0). At kinetic freezeout, the trans-

*Corresponding author. Email (M.Waqas): waqas_phy313@yahoo.com; waqas_phy313@ucas.ac.cn

†Corresponding author. Email (G. X. Peng): gxpeng@ucas.ac.cn

verse momentum spectra of the particles are no longer changed. Another type of temperature is the effective (T_{eff}). It is not a real temperature but is related to the particle mass and can be extracted from the transverse momentum spectra by using various distribution laws such as standard (Boltzmann, Bose-Einstein and Fermi-Dirac), Tsallis, and so forth.

The chemical freezeout temperature is usually obtained from the particle ratio [6–8]. Due to chemical equilibrium being meanwhile or earlier than the kinetic equilibrium, the chemical freezeout temperature is equal or higher than the kinetic freezeout temperature. The effective temperature, due to mass and flow velocity are also larger than the kinetic freezeout temperature [9, 10]. Due to more violent interactions in central collisions, both the chemical freezeout temperature and effective temperature are larger in central collisions than in the peripheral collisions. However, the situation for kinetic freezeout temperature is not clear. Several literatures claim larger T_0 in central collisions [11–16] which decrease towards periphery, while several claims larger T_0 in peripheral collisions [17–20] which decrease towards the central collisions. In addition, volume is also very important parameter in high energy collisions. The volume occupied by the ejectiles when the correlative interactions become negligible and the only force they experience is Coulombic force, is said to be kinetic freezeout volume (V). Most of the literatures agreed to the larger V as well as the transverse flow velocity (β_T) in central collisions which decrease towards periphery.

Freezeout scenario is very important in high energy collisions. Different freezeout scenarios are discussed in literature at different stages of the freezeout, but we will focus on kinetic freezeout scenarios in the present work. There are several kinetic freezeout scenarios in literature [15, 16, 21–24] which include single, double, triple and multiple freezeout scenario. In the study of production of light nuclei, it is expected that the freezeout of the particles may also be dependent on the nucleon coalescence and isospin symmetry at higher energies.

The transverse momentum spectra (p_T) of the particles are very important observables because they give very crucial information about the equilibrium dynamics and the anisotropy of the produced system in heavy ion collisions [24]. In the present work, we will analyze the p_T spectra of deuteron (d), anti-deuteron (\bar{d}), triton (t), anti-triton (\bar{t}), helion (3He) and anti-helion (${}^3\bar{He}$) in Gold-Gold (Au-Au), Lead-Lead (Pb-Pb), Proton-Lead (p-Pb) and proton-proton (pp) collisions.

2 The method and formalism

A few methods can be used for the extraction of T_0 and β_T , including but not limited to, (1) the blast-wave model with Boltzmann-Gibbs statistics (BGBW) [26–28], (2) the blast-wave model with Tsallis statistics [29], (3) an alternative method using the Boltzmann distribution [22, 28, 30–35], and (4) the alternative method using the Tsallis distribution [36, 37]. It is noteworthy that T_0 is the intercept in the linear relation $T-m_0$ in alternative method, where m_0 is the rest mass; and β_T is the slope in the linear relation $\langle p_T \rangle - \bar{m}$, where $\langle p_T \rangle$ is the mean transverse momentum and \bar{m} is the mean moving mass (i.e., the mean energy).

Ref. [38] confirms that the above methods are harmonious. Among these methods, the first method is the most direct with fewer parameters, though it has been revised in various ways and applied to other quantities [39–43]. We have used the first method, i.e., BGBW in the present work. Due to their coherence, other models will not be used [38].

BGBW is a phenomenological model which is used for the spectra of hadrons based on flowing local thermal sources with global variables of temperature, volume and transverse flow velocity. According to [26–28], the p_T distribution of BGBW can be written as

$$f(p_T) = \frac{1}{N} \frac{dN}{dp_T} = \frac{1}{N} \frac{gV}{(2\pi)^2} p_T m_T \int_0^R r dr \times I_0 \left[\frac{p_T \sinh(\rho)}{T} \right] K_1 \left[\frac{m_T \cosh(\rho)}{T} \right], \quad (1)$$

where N is the number of particles, g represents the degeneracy factor of the particle (which is different for different particles, based on $g_n = 2S_n + 1$, while S_n is the spin of the particle), V is the freezeout volume, m_T is the transverse mass ($m_T = \sqrt{p_T^2 + m_0^2}$), I_0 and K_1 are the modified Bessel functions, $\rho = \tanh^{-1}[\beta(r)]$, $\beta(r) = \beta_S (r/R)^{n_0}$ is the transverse radial flow of the thermal source at radius $0 \leq r \leq R$ with surface velocity β_S and $n_0 = 1$ [29]. In general, $\beta_T = (2/R^2) \int_0^R r \beta(r) dr = 2\beta_S / (n_0 + 2) = 2\beta_S / 3$.

Eqs. (1) can be used for the fitting of p_T spectra to obtain the parameters T_0 , V and β_T . It should be noted that Eqs. (1) can be only valid in a narrow p_T range i.e: they describe only soft excitation process. However we have to consider the hard scattering process for the spectra in a wide p_T range. In general, the contribu-

tion of hard process can be parameterized to an inverse power law [44–46], i.e., Hagedorn function [47,48]

$$f_0(p_T) = \frac{1}{N} \frac{dN}{dp_T} = Ap_T \left(1 + \frac{p_T}{p_0}\right)^{-n}, \quad (2)$$

which is resulted from the calculus of quantum chromodynamics (QCD) [44, 45, 46], where p_0 and n are the free parameters and A is the normalization constant that depends on p_0 and n .

In order to describe a wide p_T range, the superposition of both the soft and hard process can be used, which is

$$f_0(p_T) = kf_S(p_T) + (1 - k)f_H(p_T), \quad (3)$$

where k shows the contribution fraction of the first component (soft excitation), while $(1 - k)$ represents the contribution fraction of the second component (hard scattering) in Eq. 3, and according to Hagedorn model [48] the usual step function can be also be used for the superposition of soft and hard components.

According to Hagedorn thermal model [48], the two-component BGBW distribution function can also be structured by using the usual step function,

$$f_0(p_T) = \frac{1}{N} \frac{dN}{dp_T} = A_1 \theta(p_1 - p_T) f(p_T) + A_2 \theta(p_T - p_1) f(p_T), \quad (4)$$

where A_1 and A_2 are the constants which give the two components to be equal to each other at $p_T = p_1$.

3 Results and discussion

The transverse momentum (p_T) spectra, $(1/2\pi p_T)d^2N/dp_T dy$, of d , \bar{d} and t produced in Au-Au collisions at 54.4 GeV are analyzed by BGBW statistics in different centrality classes are demonstrated in fig. 1. The symbols represent the experimental data of the STAR Collaboration measured in the mid-rapidity range $|y| < 0.5$ and the solid curve are the results of our fitting by using Eq. (1). The well approximate description of the model result to the experimental data of the STAR Collaboration [49] in the special p_T range can be seen. The event centralities and the values of free parameters are listed in Table 1. The free parameters include kinetic freezeout temperature (T_0), transverse flow velocity (β_T), kinetic freezeout volume (V), normalization constant (N_0), χ^2 and the degree of freedom (dof). Each panel is followed by the results of its corresponding ratio of the data/fit. Fig. 2 demonstrates

the p_T spectra, $(1/N_{ev})d^2N/dp_T dy$ of d , \bar{d} , 3He and ${}^3\bar{He}$ in various centrality classes in Pb-Pb collisions at 5.02 TeV. The symbols stands for the experimental data of the ALICE Collaboration [50] by the Large Hadron Collider (LHC) and the solid curve represent our fitting results by using the BGBW statistics. In fig. 2 some spectra are scaled; such as the spectra of d and \bar{d} in centrality intervals 5–10%, 10–20%, 20–30%, 30–40%, 40–50%, 50–60%, 60–70%, 70–80% and 90–90% are multiplied by 1/2, 1/4, 1/8, 1/16, 1/30, 1/40, 1/40, 1/60 and 1/60, respectively.

Fig. 3 is similar to fig. 2, but it shows the p_T spectra of d , \bar{d} and $({}^3He + {}^3\bar{He})/2$ produced in different centrality intervals in p-Pb collisions at 5.02 TeV. The symbols show the experimental data measured by the ALICE Collaboration in the rapidity region $-1 \leq y \leq 1$ and $-1 < y < 0$ respectively, and the curves are the results of our fitting by using Eq. (1). The well approximate description of the model results to the experimental data of the ALICE Collaboration [51, 52] in the special p_T range can be seen.

In Fig. 4 the p_T spectra of d , \bar{d} , t , \bar{t} , 3He and ${}^3\bar{He}$ in inelastic (INEL) pp collisions at 7 TeV are presented. The symbols represent the experimental data of the ALICE Collaboration by the LHC in the rapidity interval of $|y| < 0.5$ and the results of our fitting is represented by curve. The spectra of \bar{d} , t , \bar{t} , 3He and ${}^3\bar{He}$ are multiplied by 1/2.5, 800, 400, 100 and 50, respectively. One can see the well approximate description of the model results to the experimental data of the ALICE Collaboration [53] in the special p_T range.

Fig. 5 shows the variation trend of parameters with centrality (mass). Panels (a), (b), (c) and (d) show the results for Au-Au collisions at 54.4 GeV, Pb-Pb collisions at 5.02 TeV, p-Pb collisions at 5.02 TeV and pp collisions at 7 TeV respectively. Panels (a), (b) and (c) show the dependence of T_0 on centrality, and panel (d) shows the dependence of T_0 on m_0 . The types of particles are represented by different symbols. In Fig. 5, panels (a), (b) and (c), one can see that d , \bar{d} , t , \bar{t} , 3He and ${}^3\bar{He}$ in all collisions (Au-Au, Pb-Pb and p-Pb) results in larger T_0 in central collisions which decrease towards periphery. The reason behind this is, in central collisions, large number of participants involve in interaction and the collisions are more violent that results in higher degree of excitation of the system and the kinetic freezeout temperature is high. However, the collisions become less violent as the centrality decreases and less number of participants involve in the interactions which results in comparatively low kinetic freezeout tempera-




Fig. 1. The transverse momentum (p_T) spectra of (a) d (b) \bar{d} and (c) t produced in $|y| < 0.5$ in different centrality classes in Au-Au collisions at 54.4 GeV. The symbols are the experimental data of the STAR Collaboration [49] and the curves are our fitting by using BGBW statistics. Each panel is followed by the ratio of the data/fit.

ture. This is in agreement with [11–16], but in disagreement with [17–20].

Fig. 5 includes Panel (a) d , \bar{d} , and t ; Panel (b) d , \bar{d} , t and \bar{t} ; Panel (c) d , \bar{d} and $({}^3\text{He} + {}^3\bar{\text{He}})/2$; and Panel (d) d , \bar{d} , t , \bar{t} , ${}^3\text{He}$ and ${}^3\bar{\text{He}}$. In some panels some particles and their anti-particles are missing due to the unavailability of data. In panel (a) one can see that T_0 is larger for t than both of d and \bar{d} due to its mass. While d and \bar{d} has the same mass but d freezeout earlier than \bar{d} . Similarly in panel (b) d , \bar{d} , ${}^3\text{He}$ and ${}^3\bar{\text{He}}$, the mass of ${}^3\text{He}$ and ${}^3\bar{\text{He}}$ is larger than d and \bar{d} , therefore they freezeout earlier than d and \bar{d} , but d and ${}^3\text{He}$ freezeout

earlier than \bar{d} and ${}^3\bar{\text{He}}$ respectively, while in panel (c) $({}^3\text{He} + {}^3\bar{\text{He}})/2$ has larger T_0 than d and \bar{d} whereas d freezeout earlier than \bar{d} , and in Panel (d) ${}^3\text{He}$ and ${}^3\bar{\text{He}}$, as well as t and \bar{t} freezeout earlier than d and \bar{d} , and the values for ${}^3\text{He}$ and t , and ${}^3\bar{\text{He}}$ and \bar{t} are respectively the same. Basically, the formation of light nuclei occur by the coalescence of nucleons with similar momenta. In the present work, we believe that the coalescence of nucleons is larger for d , t and ${}^3\text{He}$ compared to their anti-particles and therefore T_0 is larger for light nuclei than for their anti-nuclei. Furthermore, we observed that ${}^3\text{He}$, and t , and ${}^3\bar{\text{He}}$ and \bar{t} freezeout at the same




Fig. 2. Transverse momentum spectra of d , \bar{d} , ${}^3\text{He}$ and ${}^3\bar{\text{H}}e$ in $|y| < 0.5$ produced in different centrality intervals in Pb-Pb collisions at 5.02 TeV. The symbols represent the experimental data measured by the ALICE Collaboration [50], while the curves are our fitted results by using BGBW statistics, Eqs. (1). Each panel is followed by the ratio of the data/fit.

time. In our opinion this is due to the isospin symmetry at high energies which occurs in nearly identical masses (e.g. triton and helion) where an up quark is replaced by a down quark. In addition, T_0 in Pb-Pb is larger than in Au-Au and in the later, it is larger than in pp collisions which shows its dependence on the cross-section of interacting system. However T_0 is larger in p-Pb than in Au-Au collisions because the center of mass energy for p-Pb is 5.02 TeV which is very larger than the center of mass energy of Au-Au collision (54.4 GeV), and this may reveal its dependence on energy.

Fig. 6 is similar to Fig. 5, but shows the dependence of β_T on centrality in panels (a), (b) and (c), while panel (d) shows its dependence on m_0 . It can be seen that β_T decrease from central to peripheral collisions due to large number of participants in central collision that experience more violent squeeze and results in a rapid expansion of the system. While this expansion becomes steady from central to periphery due to decreasing the participant nucleons which results in comparatively weak squeeze. Furthermore, β_T is mass dependent. Greater the mass of the particle is, smaller




Fig. 3. Transverse momentum spectra of (a) d , (b) \bar{d} and (c) $({}^3\text{He}+{}^3\bar{\text{He}})/2$ produced in various centrality bins in p-Pb collisions at 5.02 TeV. The symbols represent the experimental data measured by ALICE Collaboration [51, 52], while the curves are our fitted results by using BGBW statistics, Eq. (1). Each panel is followed by the ratio of the data/fit.

the value of β_T . β_T for nuclei and anti-nuclei is the same. Besides, β_T shows dependence on the cross-section of interacting system. Larger the cross-section of interacting system, larger the β_T is. However β_T is slightly larger in p-Pb collisions than in Au-Au collisions due to the effect of very large center of mass energy of p-Pb than Au-Au collisions.

Fig. 7 is similar to Fig. 6, but shows the dependence of V on centrality (mass). It can be seen that V decreases from central to peripheral collisions in panel (a), (b) and (c), as the number of participant nucleons de-

creases from central collisions to periphery depending on the interaction volume. The system with more participants reaches to equilibrium quickly due to large number of secondary collisions by the re-scattering of partons, which decreases towards periphery and the system goes away from equilibrium state. In addition, V for deuteron and anti-deuteron is larger than that of triton and anti-triton as well as from helion and anti-helion. The parameter V for nuclei is larger than their anti-nuclei due to larger coalescence of nucleons for the nuclei than for their anti-nuclei. In case of triton and anti-triton and

Table 1. List of the parameters. (- is used in some places instead of dof. In fact it is not the fit result. if dof<0, the we put - instead of negative values.)

Collisions	Centrality	Particle	T_0 (GeV)	β_T (c)	$V(fm^3)$	q	N_0	χ^2/dof
Fig. 1	0-10%	d	0.118 ± 0.006	0.460 ± 0.009	3500 ± 183	$4.5 \times 10^{-4} \pm 5 \times 10^{-5}$	10/11	
Au-Au	10-20%	-	0.113 ± 0.006	0.452 ± 0.008	3350 ± 176	$1.7 \times 10^{-4} \pm 5 \times 10^{-5}$	4/11	
54.4 GeV	20-40%	-	0.109 ± 0.005	0.440 ± 0.007	3200 ± 150	$4 \times 10^{-5} \pm 5 \times 10^{-6}$	11/11	
	40-60%	-	0.105 ± 0.005	0.427 ± 0.009	3000 ± 207	$9 \times 10^{-6} \pm 5 \times 10^{-7}$	14/11	
	60-80%	-	0.101 ± 0.006	0.388 ± 0.010	2800 ± 210	$1.6 \times 10^{-6} \pm 6 \times 10^{-7}$	17/9	
Au-Au	0-10%	d	0.110 ± 0.005	0.460 ± 0.011	3200 ± 170	$5.9 \times 10^{-5} \pm 4 \times 10^{-6}$	12/10	
54.4 GeV	10-20%	-	0.105 ± 0.006	0.453 ± 0.010	3100 ± 180	$2.45 \times 10^{-5} \pm 5 \times 10^{-6}$	10/10	
	20-40%	-	0.100 ± 0.005	0.440 ± 0.008	2900 ± 180	$7.2 \times 10^{-6} \pm 6 \times 10^{-7}$	22/10	
	40-60%	-	0.097 ± 0.006	0.427 ± 0.011	2760 ± 170	$1.6 \times 10^{-6} \pm 4 \times 10^{-7}$	44/10	
	60-80%	-	0.092 ± 0.006	0.388 ± 0.012	2650 ± 140	$3.7 \times 10^{-7} \pm 5 \times 10^{-8}$	34/8	
Au-Au	0-10%	t	0.130 ± 0.005	0.420 ± 0.010	2700 ± 190	$1.3 \times 10^{-6} \pm 6 \times 10^{-7}$	16/3	
54.4 GeV	10-20%	-	0.126 ± 0.005	0.400 ± 0.012	2500 ± 150	$5.5 \times 10^{-7} \pm 4 \times 10^{-8}$	2/3	
	20-40%	-	0.122 ± 0.006	0.388 ± 0.009	2360 ± 128	$1.5 \times 10^{-7} \pm 3 \times 10^{-8}$	8/3	
	40-60%	-	0.118 ± 0.005	0.363 ± 0.012	2200 ± 146	$3.3 \times 10^{-8} \pm 5 \times 10^{-9}$	16/3	
Fig. 2	0-5%	d	0.156 ± 0.005	0.545 ± 0.009	5100 ± 172	0.077 ± 0.004	1/14	
Pb-Pb	5-10%	-	0.153 ± 0.004	0.540 ± 0.012	4940 ± 160	0.037 ± 0.005	3/14	
5.02 TeV	10-20%	-	0.150 ± 0.005	0.535 ± 0.012	4710 ± 160	0.015 ± 0.004	4/14	
	20-30%	-	0.147 ± 0.005	0.527 ± 0.009	4500 ± 167	0.006 ± 0.0004	5/14	
	30-40%	-	0.145 ± 0.005	0.510 ± 0.009	4300 ± 167	0.0023 ± 0.0004	6/14	
	40-50%	-	0.143 ± 0.005	0.500 ± 0.009	4200 ± 167	$7.5 \times 10^{-4} \pm 3 \times 10^{-5}$	6/14	
	50-60%	-	0.140 ± 0.005	0.480 ± 0.012	4100 ± 200	$2.3 \times 10^{-4} \pm 3 \times 10^{-5}$	21/14	
	60-70%	-	0.137 ± 0.006	0.450 ± 0.010	3910 ± 190	$6.3 \times 10^{-5} \pm 4 \times 10^{-6}$	29/11	
	70-80%	-	0.133 ± 0.005	0.440 ± 0.012	3500 ± 160	$2.3 \times 10^{-5} \pm 3 \times 10^{-6}$	16/9	
	80-90%	-	0.130 ± 0.006	0.420 ± 0.010	3300 ± 150	$3 \times 10^{-6} \pm 5 \times 10^{-7}$	12/6	
Pb-Pb	0-5%	d	0.148 ± 0.005	0.545 ± 0.009	4700 ± 181	0.086 ± 0.005	3/14	
5.02 TeV	5-10%	-	0.145 ± 0.006	0.540 ± 0.012	4500 ± 190	0.04 ± 0.006	5/14	
	10-20%	-	0.142 ± 0.005	0.535 ± 0.011	4300 ± 172	0.0166 ± 0.004	5/14	
	20-30%	-	0.140 ± 0.005	0.527 ± 0.012	4100 ± 150	0.007 ± 0.0004	6/14	
	30-40%	-	0.137 ± 0.006	0.510 ± 0.010	3900 ± 170	0.0026 ± 0.0004	9/14	
	40-50%	-	0.133 ± 0.005	0.500 ± 0.012	3700 ± 180	$8.5 \times 10^{-4} \pm 5 \times 10^{-5}$	12/14	
	50-60%	-	0.130 ± 0.004	0.480 ± 0.010	3500 ± 156	$2.6 \times 10^{-4} \pm 4 \times 10^{-5}$	26/14	
	60-70%	-	0.128 ± 0.005	0.450 ± 0.008	3300 ± 160	$7.2 \times 10^{-5} \pm 6 \times 10^{-6}$	22/11	
	70-80%	-	0.125 ± 0.004	0.440 ± 0.010	3100 ± 180	$2.6 \times 10^{-5} \pm 5 \times 10^{-6}$	16/9	
	80-90%	-	0.122 ± 0.005	0.420 ± 0.012	2900 ± 140	$3.2 \times 10^{-6} \pm 4 \times 10^{-7}$	15/6	
Pb-Pb	0-10%	3He	0.164 ± 0.005	0.510 ± 0.007	4400 ± 160	$1.5 \times 10^{-7} \pm 4 \times 10^{-8}$	3/1	
5.02 TeV	10-40%	-	0.160 ± 0.005	0.500 ± 0.008	4200 ± 140	$9 \times 10^{-8} \pm 4 \times 10^{-9}$	2/1	
	40-100%	-	0.156 ± 0.004	0.470 ± 0.010	4000 ± 180	$4.7 \times 10^{-8} \pm 4 \times 10^{-9}$	10/1	
Pb-Pb	0-10%	3He	0.158 ± 0.004	0.510 ± 0.007	4000 ± 140	$1.4 \times 10^{-7} \pm 4 \times 10^{-8}$	4/1	
5.02 TeV	10-40%	-	0.154 ± 0.005	0.505 ± 0.010	3800 ± 166	$9.5 \times 10^{-8} \pm 4 \times 10^{-8}$	3/1	
	30-100%	-	0.151 ± 0.005	0.475 ± 0.009	3600 ± 155	$5.3 \times 10^{-8} \pm 4 \times 10^{-9}$	4/1	
Fig. 3	0-10%	d	0.148 ± 0.006	0.480 ± 0.008	4720 ± 170	$1.25 \times 10^{-4} \pm 4 \times 10^{-5}$	12/7	
p-Pb	10-20%	-	0.143 ± 0.006	0.470 ± 0.011	4550 ± 160	$4.5 \times 10^{-5} \pm 5 \times 10^{-6}$	18/7	
5.02 TeV	20-40%	-	0.139 ± 0.006	0.440 ± 0.007	4400 ± 186	$1.7 \times 10^{-5} \pm 6 \times 10^{-6}$	13/4	
	40-60%	-	0.135 ± 0.005	0.430 ± 0.008	4200 ± 179	$5 \times 10^{-6} \pm 5 \times 10^{-7}$	21/4	
	60-100%	-	0.131 ± 0.005	0.400 ± 0.010	4130 ± 180	$8 \times 10^{-7} \pm 6 \times 10^{-8}$	35/4	
p-Pb	0-10%	d	0.140 ± 0.005	0.480 ± 0.012	4400 ± 184	$1.31 \times 10^{-4} \pm 6 \times 10^{-5}$	8/7	
5.02 TeV	10-20%	-	0.136 ± 0.006	0.470 ± 0.010	4200 ± 140	$4.4 \times 10^{-5} \pm 6 \times 10^{-6}$	18/5	
	20-40%	-	0.132 ± 0.005	0.440 ± 0.009	4000 ± 160	$1.75 \times 10^{-5} \pm 7 \times 10^{-6}$	8/5	
	40-60%	-	0.128 ± 0.004	0.430 ± 0.008	3800 ± 154	$5.5 \times 10^{-6} \pm 7 \times 10^{-7}$	21/4	
	60-100%	-	0.122 ± 0.004	0.400 ± 0.010	3600 ± 168	$1 \times 10^{-6} \pm 4 \times 10^{-7}$	16/4	
p-Pb	0-10%	$({}^3He + {}^3He)/2$	0.154 ± 0.006	0.470 ± 0.008	4000 ± 191	$5 \times 10^{-8} \pm 4 \times 10^{-9}$	4/-	
5.02 TeV	10-20%	-	0.150 ± 0.005	0.443 ± 0.011	3800 ± 166	$1.4 \times 10^{-8} \pm 5 \times 10^{-9}$	3/-	
	20-40%	-	0.146 ± 0.005	0.413 ± 0.009	3600 ± 165	$5 \times 10^{-9} \pm 3 \times 10^{-10}$	26/-	
	40-100%	-	0.142 ± 0.005	0.410 ± 0.011	3400 ± 150	$7 \times 10^{-10} \pm 3 \times 10^{-11}$	1/-	
Fig.4	-	d	0.090 ± 0.006	0.420 ± 0.008	3000 ± 158	$2 \times 10^{-6} \pm 4 \times 10^{-7}$	141/17	
pp	-	\bar{d}	0.080 ± 0.005	0.420 ± 0.007	2600 ± 145	$2 \times 10^{-6} \pm 4 \times 10^{-7}$	81/16	
7 TeV	-	t	0.105 ± 0.004	0.350 ± 0.009	2000 ± 155	$1.3 \times 10^{-10} \pm 3 \times 10^{-11}$	0.02/-	
	-	\bar{t}	0.095 ± 0.004	0.350 ± 0.008	2000 ± 145	$1.6 \times 10^{-10} \pm 4 \times 10^{-11}$	0.02/-	
	-	3He	0.105 ± 0.004	0.300 ± 0.007	2000 ± 155	$1.3 \times 10^{-10} \pm 3 \times 10^{-11}$	0.6/-	
	-	${}^3\bar{He}$	0.095 ± 0.005	0.350 ± 0.007	2000 ± 145	$6 \times 10^{-11} \pm 4 \times 10^{-12}$	38/-	




Fig. 4. is similar to Fig. 2 and 3, but it shows p_T spectra of d , \bar{d} , t , \bar{t} , ${}^3\text{He}$ and ${}^3\bar{\text{He}}$ in $|y| < 0.5$ produced in INEL pp collisions at 7 TeV. The symbols represent the experimental data measured by the ALICE Collaboration [53], while the curves are our fitted results by using BGBW statistics, Eqs (1). Each panel is followed by the ratio of the data/fit.

helion and anti-helion, V of triton and helion, and anti-triton and anti-helion are respectively the same. Besides, V is larger in Pb-Pb collisions than the rest, and in Au-Au collisions as well as p-Pb collisions it is larger than in pp collisions which shows the dependence of V on the cross-section of interacting system. However V is larger in p-Pb than in Au-Au collisions. We think that this is due to the effect of very higher center energy of p-Pb collisions compared to Au-Au collisions, because higher energy corresponds to longer evolution time which may lead to larger partonic system.

Figs. 8 (a), (b) and (c) show the dependence of N_0 on centrality. One can see that N_0 decrease with decreasing centrality. Furthermore, the parameter N_0 depends on mass of the particle. In panels (a)-(d) the parameter N_0 for deuteron and anti-deuteron are larger than triton, and in (a) it is larger for deuteron than anti-deuteron due to large coalescence of deuteron, while in panels (b)-(d) the parameter N_0 for nuclei and their anti-particles are the almost same. In general, the parameter N_0 is the same for nuclei and their anti-nuclei. The parameter N_0 for triton and helion, and anti-triton




Fig. 5. Panels (a), (b) and (c): Variation of T_0 with centrality and (d): variation of T_0 with m_0 for d , \bar{d} , t , \bar{t} , ${}^3\text{He}$ and ${}^3\bar{\text{He}}$ or $({}^3\text{He} + {}^3\bar{\text{He}})/2$ in Au-Au, Pb-Pb, p-Pb and pp collisions.

and anti-helion is the same due to the isospin symmetry. In deed N_0 is only a normalization factor and the data are not cross-section, but they are proportional to the volumes of sources of producing various particles. Therefore it is significant to study N_0 dependence.

Fig. 9 shows the variation of T_0 with β_T . It is observed that central collisions correspond to larger T_0 and β_T . The correlation between T_0 and β_T is positive. This is in agreement with [16] and in disagreement with [18]. In panel (a) the correlation of T_0 and β_T is larger for triton and that of deuteron is larger than anti-deuteron. In panel (b) the correlation of T_0 and β_T is larger for helions than deuterons and that of deuteron is larger for than anti-deuteron. Similarly in panel (c) the

helions has larger correlation between T_0 and β_T than deuteron and anti-deuteron, and for deuteron it is larger than anti-deuteron. In general, the massive particles has larger correlation between of T_0 and β_T , and the particles has larger correlation than their anti-particles due to less coalescence of anti-particles.

Fig. 10 shows the variation of T_0 with V . It is observed that central collisions correspond to larger T_0 and V . The correlation between T_0 and V is positive. In panel (a) the correlation of T_0 and V is larger for triton, while the correlation of T_0 and V for deuterons is slightly larger than anti-deuteron. In panel (b) the correlation of T_0 and V is larger for helions than deuterons, and that of helions and deuterons is slightly larger for than




Fig. 6. Panels (a), (b), (c) Variation of β_T with centrality and (d) variation of β_T with m_0 d , \bar{d} , t , \bar{t} , ${}^3\text{He}$ and ${}^3\bar{\text{He}}$ or $({}^3\text{He} + {}^3\bar{\text{He}})/2$ in Au-Au, Pb-Pb, p-Pb and pp collisions.

their anti-particles. Similarly in panel (c) the helions has larger correlation between T_0 and V than deuteron and anti-deuteron and deuteron has larger correlation of T_0 and V than anti-deuteron. In short the massive particles has larger correlation between of T_0 and V , and the particles has larger correlation than their anti-particles due to less coalescence of anti-particles.

4 Conclusions

The main observations and conclusions are summarized here.

(a) The transverse momentum spectra of light nuclei and their anti-nuclei produced in Au-Au, Pb-Pb and p-

Pb and produced in inelastic pp collisions in different centrality intervals are analyzed by the BGBW model. The model results show an agreement with the experimental data in the special p_T range measured by the STAR and ALICE Collaborations.

(b) Kinetic freezeout temperature is larger for triton and helions and their anti-particles than deuteron and anti-deuteron due to their mass. While helions and tritons have the same value of Kinetic freezeout temperature due to isospin symmetry, and deuteron, triton and helion freezeout earlier than their anti-particles respectively due to large coalescence of nucleons for the light nuclei than anti-nuclei.

(c) Kinetic freezeout temperature decrease from cen-




Fig.7 Panels (a), (b), (c) Variation of V with centrality and (d) variation of β_T with m_0 for d , \bar{d} , t , \bar{t} , ${}^3\text{He}$ and ${}^3\bar{\text{He}}$ or $({}^3\text{He} + {}^3\bar{\text{He}})/2$ in Au-Au, Pb-Pb, p-Pb and pp collisions.

tral to peripheral collisions due to the decrease of participant nucleons in the peripheral collisions which lead to the decrease in the degree of excitation of the system.

d) Transverse flow velocity increases from peripheral to central collisions due the reason that the collisions become more violent in central collisions which also expands the system rapidly.

e) The kinetic freezeout volume decreases from central to peripheral collisions and the system reaches quickly to equilibrium state due to large number of secondary collisions by the re-scattering of partons in central collisions that decreases towards periphery. In addition, The normalization constant is larger in central collisions than in peripheral collisions.

)

Data availability

The data used to support the findings of this study are included within the article and are cited at relevant places within the text as references.

Compliance with Ethical Standards

The authors declare that they are in compliance with ethical standards regarding the content of this paper.

Acknowledgements

The authors would like to thank support from the




Fig.8 Panels (a), (b), (c) Dependence of N_0 on centrality and (d) Dependence of N_0 on m_0 , for d , \bar{d} , t , \bar{t} , ${}^3\text{He}$ and ${}^3\bar{\text{He}}$ or $({}^3\text{He} + {}^3\bar{\text{He}})/2$ in Au-Au, Pb-Pb, p-Pb and pp collisions.

National Natural Science Foundation of China (Grant Nos. 11875052, 11575190, and 11135011).

References

- [1] Chatterjee. S, Das. S, Kumar. L, Mishra. D, Mohanty. B, Sahoo. R and Sharma. N, “Freeze-Out Parameters in Heavy-Ion Collisions at AGS, SPS, RHIC, and LHC Energies,” *Adv. High Energy Phys.* **2015**, 349013, (2015). doi:10.1155/2015/349013
- [2] Xu. N [STAR], “An Overview of STAR Experimental Results,” *Nucl. Phys. A* **931**, 1-12, (2014). doi:10.1016/j.nuclphysa.2014.10.022 [arXiv:1408.3555 [nucl-ex]].
- [3] Chatterjee. S and Mohanty. B, “Production of Light Nuclei in Heavy Ion Collisions Within Multiple Freeze-out Scenario,” *Phys. Rev. C* **90**, 034908, (2014). doi:10.1103/PhysRevC.90.034908 [arXiv:1405.2632 [nucl-th]].
- [4] Chatterjee. S, Mohanty. B and Singh. R, “Freeze-out hypersurface at energies available at the CERN Large Hadron Collider from particle spectra: Flavor and centrality dependence,” *Phys. Rev. C* **92**, 024917, (2015). doi:10.1103/PhysRevC.92.024917 [arXiv:1411.1718 [nucl-th]].
- [5] asänen S. S. R [ALICE], “ALICE overview,” *EPJ Web Conf.* **126**, 02026, (2016). doi:10.1051/epjconf/201612602026 [arXiv:1603.03320 [nucl-ex]].




Fig.9 Variation of T_0 with β_T .

- [6] Das. S, Mishra. D, Chatterjee. S and Mohanty. B, “Freeze-out conditions in proton-proton collisions at the highest energies available at the BNL Relativistic Heavy Ion Collider and the CERN Large Hadron Collider,” *Phys. Rev. C* **95**, 014912, (2017). doi:10.1103/PhysRevC.95.014912 [arXiv:1605.07748 [nucl-th]].
- [7] Floris. M, “Hadron yields and the phase diagram of strongly interacting matter,” *Nucl. Phys. A* **931**, 103-112, (2014). doi:10.1016/j.nuclphysa.2014.09.002 [arXiv:1408.6403 [nucl-ex]].
- [8] Huovinen. P, “Chemical freeze-out temperature in hydrodynamical description of Au+Au collisions at $s(\text{NN})^{1/2} = 200\text{-GeV}$,” *Eur. Phys. J. A* **37**, 121-128, (2008). doi:10.1140/epja/i2007-10611-3 [arXiv:0710.4379 [nucl-th]].
- [9] Andronic. A, “An overview of the experimental study of quark-gluon matter in high-energy nucleus-nucleus collisions,” *Int. J. Mod. Phys. A* **29**, 1430047, (2014). doi:10.1142/S0217751X14300476 [arXiv:1407.5003 [nucl-ex]].
- [10] De. B, “Non-extensive statistics and understanding particle production and kinetic freeze-out process from p_T -spectra at 2.76 TeV,” *Eur. Phys. J. A* **50**, 138, (2014). doi:10.1140/epja/i2014-14138-2 [arXiv:1409.3079 [nucl-th]].
- [11] Lao. H. L, Liu. F. H and Lacey. R. A, “Extracting kinetic freeze-out temperature and radial flow velocity from an improved Tsallis distribution,” *Eur. Phys. J. A* **53**, 44, (2017). [erratum: *Eur. Phys. J. A* **53** (2017) no.6, 143] doi:10.1140/epja/i2017-12238-1 [arXiv:1611.08391 [nucl-th]].




Fig.10 Variation of T_0 with V .

- [12] Wang. Q and Liu. F. H, "Initial and final state temperatures of antiproton emission sources in high energy collisions," *Int. J. Theor. Phys.* **58**, 4119-4138, (2019). doi:10.1007/s10773-019-04278-2 [arXiv:1909.02390 [hep-ph]].
- [13] Waqas. M, Peng. G. X and Liu. F. H, "An evidence of triple kinetic freezeout scenario observed in all centrality intervals in Cu-Cu, Au-Au and Pb-Pb collisions at high energies," doi:10.1088/1361-6471 [arXiv:2101.07971 [hep-ph]].
- [14] Waqas. M, Liu. F. H and Wazir. Z, "Dependence of temperatures and kinetic freeze-out volume on centrality in Au-Au and Pb-Pb collisions at high energy," *Adv. High Energy Phys.* **2020**, 8198126, (2020). doi:10.1155/2020/8198126, [arXiv:2004.03773 [hep-ph]].
- [15] Waqas. M, Liu. F. H, Fakhraddin. S and Rahim. M. A, "Possible scenarios for single, double, or multiple kinetic freeze-out in high energy collisions," *J. Phys.* **93**, 1329-1343, (2019). doi:10.1007/s12648-019-01396-9 [arXiv:1806.04312 [nucl-th]].
- [16] Waqas. M and Li. B. C, "Kinetic freeze-out temperature and transverse flow velocity in Au-Au collisions at RHIC-BES energies," *Adv.High Energy Phys.* **2020**, 1787183, (2020). [arXiv:1909.11339 [hep-ph]].
- [17] Che. G. R, Gu. J. B, Zhang. W. C and Zheng. H, "Identified particle spectra in Pb-Pb and p-Pb collisions with a modified Tsallis blast-wave model," [arXiv:2010.14880 [nucl-th]].
- [18] Adamczyk. L *et al.* [STAR], "Bulk Properties of the Medium Produced in Relativistic Heavy-Ion Collisions from the Beam Energy Scan Program," *Phys. Rev. C* **96**, 044904, (2017). doi:10.1103/PhysRevC.96.044904

- [arXiv:1701.07065 [nucl-ex]].
- [19] Uddin. S, Bhat. R. A and Bashir. I. u, “Study of Centrality Dependence of Transverse Momentum Spectra of Hadrons and the Freeze-out Parameters at $\sqrt{s(NN)}$ of 62.4 GeV, 130 GeV and 200 GeV,” [arXiv:1412.2663 [hep-ph]].
- [20] Bashir. I. u and Uddin. S, “Centrality Dependence of $K^*(892)^0$ and $\phi(1020)$ Production at LHC,” Commun. Theor. Phys. **68**, 500, (2017). doi:10.1088/0253-6102/68/4/500 [arXiv:1510.05874 [hep-ph]].
- [21] Khuntia. A, Sharma. H, Kumar Tiwari. S, Sahoo. R and Cleymans. J, “Radial flow and differential freeze-out in proton-proton collisions at $\sqrt{s} = 7$ TeV at the LHC,” Eur. Phys. J. A **55**, 3, (2019). doi:10.1140/epja/i2019-12669-6 [arXiv:1808.02383 [hep-ph]].
- [22] Lao. H. L, Wei. H. R, Liu. F. H and Lacey. R. A, “An evidence of mass-dependent differential kinetic freeze-out scenario observed in Pb-Pb collisions at 2.76 TeV,” Eur. Phys. J. A **52**, 203, (2016). doi:10.1140/epja/i2016-16203-2 [arXiv:1601.00045 [nucl-th]].
- [23] Thakur. D, Tripathy. S, Garg. P, Sahoo. R and Cleymans. J, “Indication of Differential Kinetic Freeze-out at RHIC and LHC Energies,” Acta Phys. Polon. Supp. **9**, 329, (2016). doi:10.5506/APhysPolBSupp.9.329 [arXiv:1603.04971 [hep-ph]].
- [24] Shao. M, Yi. Y, Tang. Z, Chen. H, Li. C and Xu. Z, “Examine the species and beam-energy dependence of particle spectra using Tsallis Statistics,” J. Phys. G **37**, 085104, (2010). doi:10.1088/0954-3899/37/8/085104 [arXiv:0912.0993 [nucl-ex]].
- [25] Tawfik. A. N, “Equilibrium statistical-thermal models in high-energy physics,” Int. J. Mod. Phys. A **29**, 1430021, (2014). doi:10.1142/S0217751X1430021X [arXiv:1410.0372 [hep-ph]].
- [26] Abelev. B. I, *et al.* [STAR], “Systematic Measurements of Identified Particle Spectra in pp, d^+ Au and Au+Au Collisions from STAR,” Phys. Rev. C **79**, 034909, (2009). doi:10.1103/PhysRevC.79.034909 [arXiv:0808.2041 [nucl-ex]].
- [27] Schnedermann. E, Sollfrank. J and Heinz. U. W, “Thermal phenomenology of hadrons from 200-A/GeV S+S collisions,” Phys. Rev. C **48**, 2462-2475, (1993). doi:10.1103/PhysRevC.48.2462 [arXiv:nucl-th/9307020 [nucl-th]].
- [28] Abelev. B. I *et al.* [STAR], “Identified particle production, azimuthal anisotropy, and interferometry measurements in Au+Au collisions at $\sqrt{s(NN)}^{1/2} = 9.2$ - GeV,” Phys. Rev. C **81**, 024911, (2010). doi:10.1103/PhysRevC.81.024911 [arXiv:0909.4131 [nucl-ex]].
- [29] Tang. Z, Xu. Y, Ruan. L, Buren G. van, Wang. F and Xu. Z, “Spectra and radial flow at RHIC with Tsallis statistics in a Blast-Wave description,” Phys. Rev. C **79**, 051901, (2009). doi:10.1103/PhysRevC.79.051901 [arXiv:0812.1609 [nucl-ex]].
- [30] Russo. R, “Measurement of D^+ meson production in p-Pb collisions with the ALICE detector,” [arXiv:1511.04380 [nucl-ex]].
- [31] Takeuchi. S, Murase. K, Hirano. T, Huovinen. P and Nara. Y, “Effects of hadronic rescattering on multistrange hadrons in high-energy nuclear collisions,” Phys. Rev. C **92**, 044907, (2015). doi:10.1103/PhysRevC.92.044907 [arXiv:1505.05961 [nucl-th]].
- [32] Heinz. U. W, Lecture Notes for Lectures Presented at the 2nd CERN Latin-American School of High-Energy Physics (1 C14June 2003, San Miguel Regla, Mexico), arXiv:hep-ph/0407360 (2004)
- [33] Heiselberg. H and Levy. A. M, “Elliptic flow and HBT in noncentral nuclear collisions,” Phys. Rev. C **59**, 2716-2727, (1999). doi:10.1103/PhysRevC.59.2716 [arXiv:nucl-th/9812034 [nucl-th]].
- [34] Wei H. R, Liu F. H and Lacey R. A, “Kinetic freeze-out temperature and flow velocity extracted from transverse momentum spectra of final-state light flavor particles produced in collisions at RHIC and LHC,” Eur. Phys. J. A **52**, 102, (2016). doi:10.1140/epja/i2016-16102-6 [arXiv:1601.07045 [hep-ph]].
- [35] Wei H. R, Liu F. H and Lacey R. A, “Disentangling random thermal motion of particles and collective expansion of source from transverse momentum spectra in high energy collisions,” J. Phys. G **43**, 125102, (2016) doi:10.1088/0954-3899/43/12/125102 [arXiv:1509.09083 [nucl-ex]].
- [36] Cleymans. J and Worku. D, “Relativistic Thermodynamics: Transverse Momentum Distributions in High-Energy Physics,” Eur. Phys. J. A **48**, 160, (2012). doi:10.1140/epja/i2012-12160-0 [arXiv:1203.4343 [hep-ph]].
- [37] Zheng. H and Zhu. L, “Comparing the Tsallis Distribution with and without Thermodynamical Description in $p + p$ Collisions,” Adv. High Energy Phys. **2016**, 9632126, (2016). doi:10.1155/2016/9632126 [arXiv:1512.03555 [nucl-th]].
- [38] Lao H. L, Liu F. H, Li B. C and Duan. M. Y, “Kinetic freeze-out temperatures in central and peripheral collisions: Which one is larger?,” Nucl. Sci. Tech. **29**, 82, (2018). doi:10.1007/s41365-018-0425-x [arXiv:1703.04944 [nucl-th]].
- [39] Zhang. S, Ma. Y. G, Chen. J. H and Zhong. C, “Production of Kaon and Λ in Nucleus-Nucleus Collisions at Ultrarelativistic Energy from a Blast-Wave Model,” Adv. High Energy Phys. **2015**, 460590, (2015). doi:10.1155/2015/460590 [arXiv:1411.1500 [nucl-th]].
- [40] Bialas. A, Florkowski. W and Zalewski. K, “Blast-wave model description of the Hanbury-Brown-Twiss radii in pp collisions at LHC energies,” J. Phys. G **42**,

- 045001, (2015). doi:10.1088/0954-3899/42/4/045001 [arXiv:1409.5240 [hep-ph]].
- [41] Sun. X, Masui. H, Poskanzer. A. M and Schmah. A, “Blast Wave Fits to Elliptic Flow Data at $\sqrt{s_{NN}} = 7.7\text{--}2760$ GeV,” *Phys. Rev. C* **91**, 024903, (2015). doi:10.1103/PhysRevC.91.024903 [arXiv:1410.1947 [hep-ph]].
- [42] Florkowski. W, “Blast-wave model description of the HBT radii measured in pp collisions at the LHC energies,” *Acta Phys. Polon. B* **47**, 2241, (2016). doi:10.5506/APhysPolB.47.2241 [arXiv:1603.06418 [nucl-th]].
- [43] Cimerman. J, Tomasik. B, Csanad. M and Lokos. S, “Higher-order anisotropies in the Blast-Wave Model - disentangling flow and density field anisotropies,” *Eur. Phys. J. A* **53**, 161, (2017). doi:10.1140/epja/i2017-12349-7 [arXiv:1702.01735 [nucl-th]].
- [44] Arnison. G *et al.* [UA1], “Transverse Momentum Spectra for Charged Particles at the CERN Proton anti-Proton Collider,” *Phys. Lett. B* **118**, 167-172, (1982). doi:10.1016/0370-2693(82)90623-2
- [45] Odorico. R, “DOES A TRANSVERSE ENERGY TRIGGER ACTUALLY TRIGGER ON LARGE P(T) JETS?,” *Phys. Lett. B* **118**, 151-154, (1982). doi:10.1016/0370-2693(82)90620-7
- [46] Biyajima. M, Mizoguchi. T and Suzuki. N, “Analyses of whole transverse momentum distributions in $p\bar{p}$ and pp collisions by using a modified version of Hagedorn’s formula,” *Int. J. Mod. Phys. A* **32**, 1750057, (2017). doi:10.1142/S0217751X17500579 [arXiv:1604.01264 [hep-ph]].
- [47] Abelev. B. B *et al.* [ALICE], “Production of $\Sigma(1385)^\pm$ and $\Xi(1530)^0$ in proton-proton collisions at $\sqrt{s} = 7$ TeV,” *Eur. Phys. J. C* **75**, 1, (2015). doi:10.1140/epjc/s10052-014-3191-x [arXiv:1406.3206 [nucl-ex]].
- [48] Hagedorn. R *Riv. Nuovo Cimento* **6** 1, (1983)
- [49] Zhang. D [STAR], “Light Nuclei (d, t) Production in Au + Au Collisions at $\sqrt{s_{NN}} = 7.7\text{--}200$ GeV,” *Nucl. Phys. A* **1005**, 121825, (2021). doi:10.1016/j.nuclphysa.2020.121825 [arXiv:2002.10677 [nucl-ex]].
- [50] Puccio. M “Study of the production of nuclei and anti-nuclei at the LHC with the ALICE experiment,” CERN-THESIS-2017-338.
- [51] Acharya. S *et al.* [ALICE], “Multiplicity dependence of light (anti-)nuclei production in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Lett. B* **800**, 135043, (2020). doi:10.1016/j.physletb.2019.135043 [arXiv:1906.03136 [nucl-ex]].
- [52] Acharya. S *et al.* [ALICE], “Production of (anti-) ^3He and (anti-) ^3H in p-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys. Rev. C* **101**, 044906, (2020). doi:10.1103/PhysRevC.101.044906 [arXiv:1910.14401 [nucl-ex]].
- [53] Acharya. S *et al.* [ALICE], “Production of deuterons, tritons, ^3He nuclei and their antinuclei in pp collisions at $\sqrt{s} = 0.9, 2.76$ and 7 TeV,” *Phys. Rev. C* **97**, 024615, (2018). doi:10.1103/PhysRevC.97.024615 [arXiv:1709.08522 [nucl-ex]].