

On a temperature-volume phase diagram for water and three-phase invariant reactions in pure substances

Matthew J. Powell-Palm

Department of Mechanical Engineering, University of California at Berkeley, Berkeley, CA 94720, USA

To whom correspondence should be addressed:

MJPP : mpowellp@berkeley.edu

Phase diagrams are integral to the application and interpretation of materials thermodynamics, and none is more ubiquitous than the common Temperature-Pressure diagram of water and its many icy polymorphs. Inspired by recent advances in isochoric thermodynamics, in this Communication we present a new Temperature-Volume phase diagram for water and five of its solid polymorphs. Fundamental similarities between this T - V diagram and conventional binary temperature-concentration (T - x) diagrams are demonstrated and explained, the notion of isochoric three-phase invariant reactions in pure substances (e.g. a “volumetric eutectic”) is introduced, and experimental implications are discussed.

Phase diagrams, which map material equilibrium, may be constructed to reflect a wide variety of natural thermodynamic variables¹⁻⁵. Amongst the most common diagrams for pure substances and binary substances respectively are the Temperature-Pressure and Temperature-concentration diagrams, which have been deployed widely since the turn of the 20th century. These two diagrams exhibit markedly different geometry, and are characterized by markedly different features.

The Temperature-Pressure diagram for a single-component substance is constructed by projecting the Gibbs free energy surfaces $G(T,P)$ of each phase onto T - P axes. Because temperature and pressure are both intensive variables and thus concave in the G - T - P energy space, T - P phase diagrams produce 2-dimensional single-phase regions, bounded by 1-dimensional boundary lines (such as the liquidus curve) along which two phases may coexist and producing 0-dimensional three-phase coexistence points at the meetings of any three single-phase regions (so-called triple points)³.

The Temperature-Concentration diagram for a binary substance is constructed for a given pressure (typically atmospheric) using the Gibbs free energy surfaces $G(T,x)$ of each phase. Critically however, concentration x is an extensive variable (a density of the particle number N , the extensive conjugate of the intensive chemical potential μ), and thus introduces convexity in the G - T - x energy space, as guaranteed by the fundamental thermodynamic stability criterion $\frac{\partial^2 U}{\partial X^2} \geq 0$, in which U is the internal energy of the substance and X is an arbitrary extensive variable⁶.

As Gibbs demonstrated in his seminal *On the Equilibrium of Heterogeneous Substances*⁷, this convexity introduces the opportunity for a system to reduce its free energy by forming a heterogeneous mixture of two or more co-existing phases across a range

of values along the convex axis (here concentration), defined by the tangent line in G - x space that is common to each of the phases. Following the advent of modern convex geometry theory in the 1960s, it has become standard to map the myriad common tangents that join single-phase regions and define multi-phase regions by taking the convex hull of the free energy surfaces of each phase^{3,8,9}. It is the projection of the lowest face of this convex hull (as opposed to the simple projection of the energy surfaces themselves) onto axes of temperature and concentration that then provides the standard binary T - x phase diagram.

As a consequence of this convexity of the free energy surfaces, binary T - x diagrams possess many features that the T - P diagrams of pure substances do not. Most evidently, they possess both 2-dimensional single-phase regions and 2-dimensional two-phase coexistence regions, alongside 1-dimensional three-phase coexistence lines as opposed to the 0-dimensional three-phase triple points found in T - P diagrams. These varied heterogeneous equilibrium regions are also accompanied by various reactions or transitions that describe the unique passage between them. Most notable of these reactions is the eutectic transition, which describes when a single liquid phase or a two-phase liquid-solid mixture transitions into a two-phase solid-solid mixture. This transition or reaction is often described as a “three-phase invariant”, as the transition to the solid-solid mixture requires traversing the three-phase, liquid-solid-solid coexistence line (i.e. the eutectic line). Other three-phase invariant reactions include the peritectic, monotectic, eutectoid, peritectoid, etc. These reactions are integral features of multicomponent phase diagrams, but are not typically thought to occur in pure systems (as reinforced by the absence of such features on the pure-substance T - P diagram).




Figure 1. Temperature-Volume diagram of water. This diagram was constructed by projecting the lower convex hull of the specific Helmholtz free energy surfaces of each phase onto axes of temperature and specific volume. All free energy data were generated using the SeaFreeze framework. Two-phase co-existence regions are labeled, and horizontal lines mark three-phase coexistence lines, the corresponding pressures of which are also labeled.

Here, we will demonstrate that these reactions are in fact not endemic to multicomponent phase diagrams, but may and will also occur in pure-substance or single-component diagrams formulated using appropriate natural variables. In Figure 1, we have constructed the Temperature-Volume phase diagram for water in the temperature range 190 – 353.5 K, including the equilibrium regions of liquid water, ice-Ih, ice-II, ice-III, ice-V, and ice-VI. This temperature range was chosen based on the phases for which mutually-consistent equations of state were available. To construct this phase diagram, we followed the method of Powell-Palm, Rubinsky & Sun: the 2D convex hull of the Helmholtz free energy surfaces $F(T, V)$ of each phase was calculated for temperatures within the stated range, with an increment of 0.08 K; single-phase points on the lower convex hull, which represent equilibrium states, were projected onto axes of the natural variables of the system (T, V); two-phase coexistence regions and three-phase coexistence lines, which definitionally join the single-phase regions within the convex hull, were identified and

marked accordingly. All Helmholtz free energy data were generated via the SeaFreeze framework¹⁰.

Important morphological aspects of this diagram are noted as follows: Firstly, the reader will notice that this T - V diagram appears identical in structure to conventional binary T - x diagrams, even though this diagram describes a pure substance. This similarity is a consequence of the fundamental similarity between all extensive and intensive variables in Gibbsian thermodynamic geometry—free energy surfaces are convex in extensive variables, leading to heterogeneous or multiphase equilibrium, and concave in intensive variables, leading to homogeneous or single-phase equilibrium^{7,11}. Thus, enforcing volume (extensive) as a natural variable of the system is directly thermodynamically analogous to enforcing concentration (extensive) as a natural variable of the system. In this way, we can understand by analogy not only the two-phase equilibrium regions, the composition of which may be analyzed by the Lever Rule as in a binary phase diagram, but also the three-phase invariant lines. These lines, which occur at the temperatures *and*

pressures of the triple points common in T - P phase diagrams, are directly analogous to three-phase invariant lines in binary diagrams; they mark a line in the extensive axis (here volume) along which one may move at a given temperature and maintain three-phase coexistence. Furthermore, the point at which a single-phase region (such as the liquid water region) touches the three-phase coexistence line provides a direct analogue to the binary eutectic point. At the volume coordinate of this point, one may cool directly from liquid water to an ice-Ih – ice-III two-phase mixture, bypassing the liquid - ice-Ih and liquid - ice-III two-phase coexistence regions. To the author's knowledge, the premise of eutectic-like transitions in pure substances has not been previously explored, and we will thus label the aforementioned eutectic-like point as the "volumetric eutectic". The volumetric eutectic point of liquid water as calculated herein occurs at $9.16 \times 10^{-4} \text{ m}^3/\text{kg}$ and 251.2 K.

Additionally, as stands to reason, other features analogous to binary phase diagrams can be found in Fig. 1, including volumetric eutectoid points (e.g. from ice-III to the ice-II – ice-Ih two-phase mixture [$8.64 \times 10^{-4} \text{ m}^3/\text{kg}$, 238.2 K]); volumetric peritectic points (e.g. from the ice-VI – Liquid Water two-phase mixture to single-phase ice-V [$7.89 \times 10^{-4} \text{ m}^3/\text{kg}$, 273.3K]); and volumetric peritectoid points (e.g. from the ice-V – ice-III two-phase mixture to single-phase ice-II [$8.38 \times 10^{-4} \text{ m}^3/\text{kg}$, 249.4 K]).

This fundamental similarity between T - x and T - V phase diagrams (regardless of the number of components in the system) is generalizable to diagrams featuring any arbitrary pairing of an intensive natural variable (such as T or P) with an extensive natural variable (such as x or V); the geometric features are not inherent products of any given physical behavior possessed by multicomponent systems (or isochoric systems for that matter), but instead to the convexity of free energy curves in the free energy – extensive variable space.

This analysis provides two useful conclusions. Firstly, by enforcing any extensive natural variable (e.g. volume, concentration, strain, magnetic moment,

polarization, etc.), varied states of heterogeneous multi-phase equilibrium may be achieved in even pure or single-component substances, analogous to those encountered in classical binary systems, and these varied multiphase equilibria may be mapped by phase diagrams identical in nature and geometry to the common binary T - x phase diagrams.

The difficulty of enforcing extensive variables experimentally may vary substantially from application to application, but a recent example of this principle put into practice is isochoric cryopreservation¹²⁻¹⁴, in which an aqueous medium containing a sensitive biologic is confined completely in a rigid high-strength metallic container without air infiltration, inducing constant-volume conditions, and the biologic is stored safely at sub-zero temperatures in the liquid portion of the stable two-phase liquid - ice-Ih system that develops.

Secondly, under these same extensive variable conditions, the suite of three-phase invariant reactions typical to binary systems, such as the eutectic transition, may be realized between the condensed phases of pure substances. To the author's knowledge such reactions have never been attempted, but theory would suggest that they are certainly possible.

In sum, the fundamental similarity between the phase diagrams (and accordingly phase equilibria) of binary systems with enforced temperature-concentration conditions and pure systems with enforced temperature-volume conditions has been demonstrated by way of a new T - V diagram for water and its icy polymorphs. Volumetric three-phase invariant features have been identified on this new diagram, suggesting by analogy that such reactions should be possible amongst the phases of pure substances. The author suggests that this geometric analogy may be established identically between any phase diagrams featuring this characteristic pairing of an intensive and extensive natural variable, and that creation of such phase diagrams from existing equations of state may provide a route to all manner of interesting materials applications.

Acknowledgements: The author acknowledges no external funding for this work.

References:

1. Zhu, W. *et al.* Room temperature electrofreezing of water yields a missing dense ice phase in the phase diagram. *Nat. Commun.* (2019) doi:10.1038/s41467-019-09950-z.
2. Sun, W., Kitchaev, D. A., Kramer, D. & Ceder, G. Non-equilibrium crystallization pathways of manganese oxides in aqueous solution. *Nat. Commun.* (2019) doi:10.1038/s41467-019-08494-6.
3. Powell-Palm, M. J., Rubinsky, B. & Sun, W. Freezing water at constant volume and under confinement. *Commun. Phys.* **3**, (2020).
4. Li, Y. L. & Chen, L. Q. Temperature-strain phase diagram for BaTiO₃ thin films. *Appl. Phys. Lett.* **88**, (2006).
5. Das, P., Vinutha, H. A. & Sastry, S. Unified phase diagram of reversible–irreversible, jamming, and yielding transitions in cyclically sheared soft-sphere packings. *Proc. Natl. Acad. Sci. U. S. A.* **117**, (2020).
6. Callen, H. B. & Scott, H. L. Thermodynamics and an Introduction to Thermostatistics, 2nd ed. *Am. J. Phys.* **66**, (1998).
7. Gibbs, J. W. On the equilibrium of heterogeneous substances. *Am. J. Sci.* **s3-16**, (1878).
8. Hildebrandt, D. & Glasser, D. Predicting phase and chemical equilibrium using the convex hull of the Gibbs free energy. *Chem. Eng. J. Biochem. Eng. J.* **54**, (1994).
9. Hegde, V. I., Aykol, M., Kirklin, S. & Wolverton, C. The phase stability network of all inorganic materials. *Sci. Adv.* **6**, (2020).
10. Journaux, B. *et al.* Holistic Approach for Studying Planetary Hydrospheres: Gibbs Representation of Ices Thermodynamics, Elasticity, and the Water Phase Diagram to 2,300 MPa. *J. Geophys. Res. Planets* **125**, (2020).
11. Gibbs, J. W. A method of geometrical representation of the thermodynamic properties of substances by means of surfaces. *Trans. Connect. Acad. Arts Sci.* **2**, 382–404 (1873).
12. Powell-Palm, M. J., Zhang, Y., Aruda, J. & Rubinsky, B. Isochoric conditions enable high subfreezing temperature pancreatic islet preservation without osmotic cryoprotective agents. *Cryobiology* (2019) doi:10.1016/j.cryobiol.2019.01.003.
13. Rubinsky, B., Perez, P. A. & Carlson, M. E. The thermodynamic principles of isochoric cryopreservation. *Cryobiology* (2005) doi:10.1016/j.cryobiol.2004.12.002.
14. Wan, L. *et al.* Preservation of rat hearts in subfreezing temperature isochoric conditions to – 8 °C and 78 MPa. *Biochem. Biophys. Res. Commun.* (2018) doi:10.1016/j.bbrc.2018.01.140.