

What happens once an accelerating observer has detected a Rindler particle?

Angel Garcia-Chung^{1,2*}, Benito A. Juárez-Aubry^{3†}
and Daniel Sudarsky^{3‡}

¹Departamento de Física, Universidad Autónoma Metropolitana - Iztapalapa,
San Rafael Atlixco 186, Ciudad de México 09340, México.

²Tecnológico de Monterrey, Escuela de Ingeniería y Ciencias,
Carr. al Lago de Guadalupe Km. 3.5, Estado de Mexico 52926, Mexico.

³Departamento de Gravitación y Teoría de Campos, Instituto de Ciencias Nucleares,
Universidad Nacional Autónoma de México, A. Postal 70-543, Mexico City 045010,
Mexico

April 26, 2022

Abstract

In a seminal paper, Unruh and Wald found that the detection of a right Rindler particle by a linearly uniformly accelerated detector coupled to a Klein-Gordon field in the Minkowski vacuum leads to the creation of a Minkowski particle from the inertial viewpoint. In this paper, we revisit the framework studied by Unruh and Wald, but now consider in addition what happens once the particle has been measured somewhere in the right Rindler wedge. From an orthodox point of view, the change in the field state induced by the measurement is non-local and occurs both in the left and right Rindler wedges. If one takes semiclassical gravity seriously in this context, this opens the possibility for designing superluminal communication protocols between two spacelike separated observers confined to the right and left Rindler wedges respectively. We discuss the possible ways of avoiding such measurement-induced, faster-than-light signaling protocols.

1 Introduction

The Unruh effect posits that on a linearly uniformly accelerated trajectory the Minkowski vacuum looks like a thermal state at a temperature proportional to the trajectory's proper acceleration [1, 2, 3]. This is the Unruh temperature, given by $T_U = a/(2\pi)$ in natural units. The study of the Unruh effect continues to be of great interest in theoretical physics, see e.g. [4] for a study of the problem as return to equilibrium, [5] for the characterisation of thermalisation time for the Unruh effect, [6] in the context of entanglement harvesting. The intimate relationship between the Unruh effect and black holes in equilibrium, which indeed originally motivated Unruh's work [3], is by now well established (see e.g. [7, Chap. 5]), and can

*alechung@xanum.uam.mx

†benito.juarez@correo.nucleares.unam.mx

‡sudarsky@nucleares.unam.mx

be seen very explicitly in two-dimensional situations [8]. [9, 10] include thorough discussions of the Unruh effect, including applications and structural properties. Very recently a concrete experimental proposal has been put forward in [11] to detect for the first time the analogous Unruh temperature along uniformly accelerated circular motions. This is an analogous circular Unruh effect, see e.g. [12, 13, 14].

Sitting at the heart of the Unruh effect is the fact that the Minkowski vacuum state restricted to, say, the right Rindler wedge of Minkowski spacetime, $|t| < x$, can be formally represented as thermal mixture of so-called *Rindler particles* supported on the right Rindler wedge. These are nothing but particles defined in the Fulling-Rindler quantisation in flat spacetime, for which the notion of positive energy is defined with respect to Lorentz boosts, $b^a = a(x \partial_t^a + t \partial_x^a)$, which generate the natural notion of time evolution for linearly accelerated observers.

Following this observation, in 1984 Unruh and Wald wrote a seminal paper [15] where they clarified what occurs when a linearly uniformly accelerated observer detects a Rindler particle: From the point of view of an inertial observer in Minkowski spacetime, the absorption of a Rindler particle – modelled as a two-level detector excitation – corresponds to the emission of a Minkowski particle. The paper [15] is remarkable in that not only did it illustrate the relativity of the notion of a particle, detection and emission, but clarified the fact that working in terms of quantum fields (and taking particles to have an operational meaning) is fully consistent with the basic ideas underlying the equivalence principle.

Furthermore, [15] has served as the starting ground for further developments. For example, the study of *bremssstrahlung* as seen from the point of view of accelerated observers [16, 17], the analysis of the decay of accelerated protons, and the finding that such behavior approaches that of accelerated neutrons, as the mass scale characterizing that acceleration – i.e., the corresponding Unruh temperature – increases, and disappears exponentially as that quantity grows beyond the value of the proton-neutron mass gap [18, 19].

In any case, there are three central questions that are addressed in [15]. The first one is to analyse the unitary evolution of the joint field-detector system when the field is initially in the Minkowski vacuum state and the two-level detector, initially switched off and prepared in the ground state, follows a linearly uniformly accelerated trajectory in the right Rindler wedge. This is done using perturbation theory in the interaction picture up to first order. (Second order contributions were further studied in [20].) The second question is to see what the updated state of the field is assuming the detector has detected a Rindler particle after some interaction time has elapsed, namely a one-particle state in the Minkowski folium, and to obtain the updated stress-energy tensor. It is found that, since the updated field state is a one-particle state, the energy of the field has increased upon detection. The third question addressed in [15] is whether detecting Rindler particles can be used as a mechanism for extracting an unbounded amount of energy from the field or to send a superluminal signal from the right to the left Rindler wedges. In both cases, the analysis leads to a negative answer.

The motivation of this work is two-fold. First, we wish to revisit the three central questions discussed in [15]. Concerning the first one, we note that the calculation in [15] is performed by exploiting an analogy with the situation of a detector interacting with a field in a thermal state that describes a *proper mixture* [21], i.e., such that the actual state of the system is pure, but there is a degree of ignorance as to what the state of the system actually is, which is encoded in weights accounting for a probability distribution of the possible (pure) states of the system. This results in a mixed state description of a pure state due to ignorance. On the other hand, the most natural description for the Minkowski vacuum from the point of view of an accelerated observer is that of a thermal state as an *improper mixture* [21] (see footnote 2 for more details), as the left Rindler wedge degrees of freedom must be traced out,

yielding a reduced mixed state. In sec. 2 we will carry out the calculation from the improper-mixture viewpoint. While the results coincide mathematically, as they should, we think that this treatment is conceptually clearer.

We then proceed to calculate the updated expectation value of the stress-energy tensor once the detector has clicked. We obtain expressions in both the right and the left Rindler wedge, adding to the result displayed in [15] for the right Rindler wedge, as we show in sec. 2. Concerning the third central question addressed in [15], on the point of energy extraction, we agree with the no-go argument presented by Unruh and Wald: while the energy is *not conserved* for a single measurement, it is on the long run for very many successive measurements, but the discussion on superluminal communication ties in with the second motivation of this paper:

Here we shall raise the point that there is a potential issue after a single measurement has been carried out, if one is to trust the semiclassical approximation of quantum gravity “before” and “after” the measurement has been performed. The point will be that in semiclassical gravity the expectation value of the stress-energy tensor can be used to actually source geometry, see eq. (34) below. Thus, an abrupt change of this quantity should be detectable by an experiment on the gravitational sector. Where and how this abrupt change occurs, i.e., where and how the state of the field can be seen as collapsing upon a measurement of the detector must undoubtedly play a rôle on how to prevent this apparent paradox from occurring, but our current understanding of these questions is fuzzy – hence the use of inverted commas around the words *before* and *after*. Thus, it seems to us that the resolution of this apparent contradiction is connected with the *measurement problem* of quantum theory.

On this point we wish to emphasise that, while Unruh and Wald correctly point out in their discussion in [15, Sec. IV] that the presence of a detector (switched on or otherwise) in the right Rindler wedge has no influence on the left Rindler wedge, their argument is based on the Heisenberg-picture observation that the effects of the detector can only affect the causal future of the coupling region between the detector and the field. In fact, this observation does not even depend on the details of the detector or the field observables, see e.g. [22] for a precise statement in some generality. The limitation of that argument is that it does not take into account the non-unitary measurement of the detector, which is typically described as a projection onto the out-state in the interaction picture. This is a central difference with the position assumed in this paper.

The organisation of this paper is as follows: In Sec. 2 we describe the evolution of the field-detector system using a left and right doubled Fock space representation for the field, and we obtain the stress-energy tensor after a Rindler particle has been measured. In doing so, we do not make any assumptions on the details of the coupling of the detector and the field, in particular we do not assume a long-time limit for the interaction, other than assuming that the coupling is weak, which allows us to conform ourselves with first-order effects in the coupling. We then discuss the non-conservation of energy upon measurements in Sec. 4 in a simplified setting, for the sake of clarity. The possibility of faster than light signalling, its implications and potential avoidances appear in Sec. 5. Discussions and conclusions appear in Sec. 6.

2 What happens once an accelerating observer has detected a Rindler particle

Consider as in [15] a particle detector coupled to a Klein-Gordon field in Minkowski spacetime following a linearly uniformly accelerated trajectory with acceleration a in the right Rindler wedge. In other words, the particle detector follows the integral curve generated by the boost $b^a = a(x\partial_t^a + t\partial_x^a)$. While currently the pointlike Unruh-DeWitt detector [23, 24, 5] is the most prominent detector model used in studies about the relativistic quantum information and QFT

in curved spacetime literature, we shall model our detector as Unruh and Wald have in [15] to stay closer to their original treatment.

The detector is a two-level system with Hilbert space \mathbb{C}^2 spanned by energy eigenstates $|\uparrow\rangle$ and $|\downarrow\rangle$. The detector Hamiltonian is $\hat{H}_D := \Omega \hat{A}^* \hat{A}$, where \hat{A} and \hat{A}^* are raising and lowering operators, and $\Omega > 0$ is the energy of the excited state, i.e., $\hat{H}_D |\uparrow\rangle = \Omega |\uparrow\rangle$ and $\hat{H}_D |\downarrow\rangle = 0$.

The coupled detector-field theory is described by the interaction Hamiltonian

$$\hat{H} = \hat{H}_D \otimes \mathbb{1} + \mathbb{1}_D \otimes \hat{H}_\Phi + \hat{H}_I, \quad (1)$$

where \hat{H}_Φ is the Klein-Gordon Hamiltonian and the interaction Hamiltonian is defined by

$$\hat{H}_I(\tau) = \epsilon(\tau) \int_{\Sigma} e^{2a\xi} d\xi dy dz \left[\psi(\xi, y, z) \hat{A}(\tau) + \bar{\psi}(\xi, y, z) \hat{A}^*(\tau) \right] \otimes \hat{\Phi}(\tau, \xi, y, z), \quad (2)$$

where $\hat{\Phi}$ is the Klein-Gordon field, $\psi \in C_0^\infty(\Sigma)$ defines the profile of the spatial extension of the detector and $\epsilon \in C_0^\infty(\mathbb{R})$ is a switching function that controls the interaction between the detector and the Klein-Gordon field along the linearly uniformly accelerated trajectory of the detector. We shall assume that the interaction between the detector and the field is weaker than any other scale in the problem and that it takes place for sufficiently long times.

The coupling takes place in the right Rindler wedge, where the flat metric can be written in terms of Rindler coordinates

$$t = \frac{1}{a} e^{a\xi} \sinh(a\tau), \quad x = \frac{1}{a} e^{a\xi} \cosh(a\tau). \quad (3)$$

It takes the form

$$ds^2 = -e^{2a\xi} (d\tau^2 - d\xi^2) + dy^2 + dz^2. \quad (4)$$

Furthermore, in the right Rindler wedge, the Klein-Gordon field can be written as

$$\hat{\Phi}(\tau, \xi, y, z) = \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa (v_{I\vec{\kappa}}(\tau, \xi, y, z) \hat{a}_{R\vec{\kappa}} + \overline{v_{I\vec{\kappa}}}(\tau, \xi, y, z) \hat{a}_{R\vec{\kappa}}^*), \quad (5)$$

with $\vec{\kappa} = (\omega, \kappa_y, \kappa_z)$, and where the right Rindler modes can be written in terms of the modified Bessel function of the second kind or MacDonald's function,

$$v_{I\vec{\kappa}}(x) = \sqrt{\frac{\sinh\left(\frac{\pi\omega}{a}\right)}{4\pi^4 a}} K_{i\omega/a} \left[\frac{\sqrt{\kappa_y^2 + \kappa_z^2 + m^2}}{a} e^{a\xi} \right] e^{-i\omega\tau + i(y\kappa_y + z\kappa_z)} \quad (6)$$

and the formal sharp-momentum annihilation and creation operators are $\hat{a}_{R\vec{\kappa}} := \hat{a}(\overline{v_{R\vec{\kappa}}})$ and $\hat{a}_{R\vec{\kappa}}^* := \hat{a}^*(v_{R\vec{\kappa}})$ respectively. The annihilation operator annihilates the right Fulling-Rindler vacuum, Ω_R , while creation operators create Rindler particles.

A fully analogous description of the quantum theory holds in the left Rindler wedge. Introducing the left Rindler coordinates

$$t = \frac{1}{a} e^{a\tilde{\xi}} \sinh(a\tilde{\tau}), \quad -x = \frac{1}{a} e^{a\tilde{\xi}} \cosh(a\tilde{\tau}) \quad (7)$$

the field in the left Rindler wedge takes the analogous form

$$\hat{\Phi}(\tilde{\tau}, \tilde{\xi}, y, z) = \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \left(v_{II\vec{\kappa}}(\tilde{\tau}, \tilde{\xi}, y, z) \hat{a}_{L\vec{\kappa}} + \overline{v_{II\vec{\kappa}}}(\tilde{\tau}, \tilde{\xi}, y, z) \hat{a}_{L\vec{\kappa}}^* \right), \quad (8)$$

where the left Rindler modes have an identical form to the right modes (6) upon the replacement of τ and ξ by $\tilde{\tau}$ and $\tilde{\xi}$. It is very well known that the Minkowski vacuum restricted to the (right

or left) Rindler wedges looks like a thermal mixture of (right or left, resp.) Rindler particles. See appendix A for details.

In any case, in the case at hand we consider that the state of the system is prepared initially (before the switch-on of ϵ) as the tensor product

$$|s_{-\infty}\rangle = |\downarrow\rangle \otimes |\Omega_M\rangle. \quad (9)$$

In the interaction picture, the late-time state of the system (after the switch-off of ϵ) is given by

$$|s_{\infty}\rangle = T\left(e^{-i\int_{\mathbb{R}} d\tau \hat{H}_I}\right) |s_{-\infty}\rangle = |s_{-\infty}\rangle + \left[-i\int_{\mathbb{R}} d\tau \hat{H}_I + O(\epsilon^2)\right] |s_{-\infty}\rangle, \quad (10)$$

under the assumption that the coupling is weak. To first order in perturbation theory, the late-time state of the system is

$$|s_{\infty}\rangle = |\downarrow\rangle \otimes |\Omega_M\rangle - i|\uparrow\rangle \otimes \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_I d\text{vol}(x) \zeta(x) \left(v_{I\vec{\kappa}}(x) \hat{a}_{R\vec{\kappa}} + \overline{v_{I\vec{\kappa}}(x)} \hat{a}_{R\vec{\kappa}}^*\right) |\Omega_M\rangle, \quad (11)$$

where the volume element in the right Rindler wedge is locally $d\text{vol}(x) = e^{2a\xi} d\tau d\xi dy dz$ and

$$\zeta(x) := e^{i\Omega\tau} \varepsilon(\tau) \overline{\psi(\xi, y, z)}. \quad (12)$$

In [15] the assumption has been made that $\varepsilon(\tau)$ is nearly constant, physically representing a long interaction between the detector and the field, such that switching effects are negligible. In this case, the τ integral can be performed directly and one obtains a factor proportional to $\delta(\Omega - \omega)$ that indicates that only the modes $v_{I\vec{\kappa}}$ with frequency highly localised around Ω will contribute to first order in eq. (10). This makes perfect sense, after a long interaction time the only modes that get excited from the vacuum state are those whose energy coincides with the frequency gap of the detector.

Let us however digress at this point and not make this approximation. The reason is that in our case we are interested in post-measurement effects for measurements carried out after a finite time of interaction between the field and the detector. On the assumption that the detector clicks, the updated state for the field becomes

$$|f\rangle = -i\mathcal{N} \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_I d\text{vol}(x) \zeta(x) \left(v_{I\vec{\kappa}}(x) \hat{a}_{R\vec{\kappa}} + \overline{v_{I\vec{\kappa}}(x)} \hat{a}_{R\vec{\kappa}}^*\right) |\Omega_M\rangle, \quad (13)$$

where the normalisation \mathcal{N} is given by

$$\mathcal{N} = \left(\int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_I d\text{vol}(x) \int_I d\text{vol}(x') \overline{\zeta(x)} \zeta(x') \left(\frac{v_{I\vec{\kappa}}(x) \overline{v_{I\vec{\kappa}}(x')}}{1 - e^{-2\pi\omega/a}} + \frac{\overline{v_{I\vec{\kappa}}(x)} v_{I\vec{\kappa}}(x')}{e^{2\pi\omega/a} - 1} \right) \right)^{-1/2} \quad (14)$$

as can be seen in appendix B.

We are interesting in the change in the expectation value of the stress-energy tensor of the field in the updated state, i.e., we are interested in

$$\Delta T_{ab} := \langle f | \hat{T}_{ab}(x) f \rangle - \langle \Omega_M | \hat{T}_{ab} \Omega_M \rangle \quad (15)$$

in the right and left Rindler wedges. (Note that imposing $\langle \Omega_M | \hat{T}_{ab} \Omega_M \rangle = 0$, we have that $\Delta T_{ab} = \langle f | \hat{T}_{ab}(x) f \rangle$.) To this end, if we use a point-splitting prescription for renormalising the stress-energy tensor the object of interest is to obtain the two-point function in the left

and right Rindler wedges. It follows from the calculations in appendices C and D that the two-point function in the updated state takes the form

$$\langle f|\hat{\Phi}(x)\hat{\Phi}(x')f\rangle = \langle \Omega_M|\hat{\Phi}(x)\hat{\Phi}(x')\Omega_M\rangle + \Delta_R(x, x') \text{ in the right Rindler wedge and,} \quad (16)$$

$$\langle f|\hat{\Phi}(x)\hat{\Phi}(x')f\rangle = \langle \Omega_M|\hat{\Phi}(x)\hat{\Phi}(x')\Omega_M\rangle + \Delta_L(x, x') \text{ in the left Rindler wedge.} \quad (17)$$

Here, Δ_R and Δ_L are real, smooth, symmetric bi-functions given by

$$\begin{aligned} \Delta_R(x, x') &= \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \frac{\overline{\zeta(y)}\zeta(y')}{(1 - e^{-2\pi\omega_p/a})(1 - e^{-2\pi\omega_\kappa/a})} \\ &\quad \times \left(v_{I\bar{p}}(y)v_{I\bar{\kappa}}(x)\overline{v_{I\bar{p}}(x')v_{I\bar{\kappa}}(y')} + \overline{v_{I\bar{\kappa}}(y)v_{I\bar{\kappa}}(x)}v_{I\bar{p}}(x')\overline{v_{I\bar{p}}(y')}e^{-2\pi\omega_\kappa} \right. \\ &\quad \left. + \overline{v_{I\bar{p}}(y)v_{I\bar{\kappa}}(x)}v_{I\bar{p}}(x')\overline{v_{I\bar{\kappa}}(y')}e^{-2\pi\omega_p} + \overline{v_{I\bar{\kappa}}(y)v_{I\bar{\kappa}}(x)}\overline{v_{I\bar{p}}(x')v_{I\bar{p}}(y')}e^{-2\pi\omega_\kappa}e^{-2\pi\omega_p} \right) + \text{c.c.}, \end{aligned} \quad (18)$$

$$\begin{aligned} \Delta_L(x, x') &= \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)}\zeta(y') \frac{e^{-\pi\omega_p/a}e^{-\pi\omega_\kappa/a}}{(1 - e^{-2\pi\omega_p/a})(1 - e^{-2\pi\omega_\kappa/a})} \\ &\quad \times \left(v_{I\bar{p}}(y)v_{II\bar{\kappa}}(x)\tilde{v}_{II\bar{p}}(x')\tilde{v}_{II\bar{\kappa}}(y') + v_{I\bar{p}}(y)\tilde{v}_{II\bar{p}}(x)v_{II\bar{\kappa}}(x')\tilde{v}_{II\bar{\kappa}}(y') \right. \\ &\quad \left. + v_{I\bar{p}}(y)\overline{v_{II\bar{\kappa}}(x)\tilde{v}_{II\bar{p}}(x')\tilde{v}_{II\bar{\kappa}}(y')} + \overline{v_{I\bar{p}}(y)\tilde{v}_{II\bar{p}}(x)}v_{II\bar{\kappa}}(x')\tilde{v}_{II\bar{\kappa}}(y') \right) + \text{c.c.}, \end{aligned} \quad (19)$$

where in eq. (19) the tilde on the modes denotes a parity operator in the orthogonal direction to the Rindler wedges, i.e., $\tilde{v}_{II\bar{\kappa}} := v_{I(\omega_\kappa, -\kappa_\perp)}$ and similarly for $\tilde{v}_{II\bar{p}}$.

Let us now compute the expectation value of the stress-energy tensor in the state $|f\rangle$. Note that a difference with the calculation in [15] is that we do not treat the state of the system as an improper thermal mixture in the right Rindler wedge, but rather as a pure state defined in the right and left Rindler wedges, which allows us to obtain the renormalised expectation value of the stress energy tensor for points in the right and left Rindler wedges.

Here we use the terminology introduced by d’Espagnat [21] to clarify that the same mathematical object, a density matrix, can be used to represent two very different physical situations: 1) The case in which one is interested in an ensemble of identical quantum systems, each one of which is in one pure and definite quantum state among a list possible such states $\{|i\rangle\}$ and where the fraction of such states in the ensemble is given by a certain classical distribution function $f(i)$, and 2) the case in which a system of interest S is a subsystem of a larger system $S + E$ with the latter in a given pure quantum state, but with our interest focused just on S which can therefore be characterised in terms of the reduced density matrix obtained after tracing over the degrees of freedom of E . For the first case one reserves the name “proper mixture” and says the density matrix represents such proper mixture, and for the second case one reserves the name “improper mixture”, and equally indicates the density matrix is to be understood as representing the improper mixture. We note that another situation one might want to consider, is one in which one is dealing with a single quantum system S which is in a pure state, which however is not completely known, and for which one has information about the classical probability $p(i)$ of the system being in each one of the quantum states. For such situation one can often use the characterisation provided by case 1) by considering a corresponding imaginary ensemble, in which the fraction is arranged to match the given probability i.e., $f(i) = p(i)$. That situation one also talks by extension about a proper mixture, even though the state of the system is pure, and thus its “properness” (or the fact that we do not express the state as a pure one) is just a result of our ignorance. Finally, as is usual, a density matrix is characterised as thermal if its representation in the energy basis has the standard thermal weights. Thus a thermal density matrix can be proper or improper.

3 The change in the stress-energy tensor

In order to compute the stress-energy tensor in the updated state, $\langle f|\hat{T}_{ab}(x)f\rangle$, in the right/left wedge we apply the point-splitting operator

$$\mathcal{T}_{ab} := g_b^{b'} \nabla_a \nabla_{b'} - \frac{1}{2} g_{ab} g^{cd'} \nabla_c \nabla_{d'} - \frac{1}{2} g_{ab} m^2, \quad (20)$$

where $g_a^{a'}$ is the parallel-transport propagator, to $\Delta_{R/L}(x, x')$ given by eq. (53), and then take the coincidence limit as

$$\langle f|\hat{T}_{ab}(x)f\rangle = \lim_{x' \rightarrow x} \mathcal{T}_{ab} \Delta_{R/L}(x, x'). \quad (21)$$

It is useful to locally express the parallel-transport components in Rindler coordinates by using the formula $g_\mu^{\mu'}(x, x') = e_\mu^I(x) e_I^{\mu'}(x')$ in terms of the soldering form and its inverse. We then have that in the right Rindler wedge the parallel-transport propagator has components $g_\eta^{\eta'}(x, x') = e^{a\xi} e^{-a\xi'}$, $g_\xi^{\xi'}(x, x') = e^{a\xi} e^{-a\xi'}$, $g_y^{y'} = 1$ and $g_z^{z'} = 1$, with all other components vanishing.

We are chiefly concerned with changes in the left Rindler wedge, which is causally disconnected from the detector that clicks. Inserting (19) into eq. (21) we have that in the left Rindler wedge stress-energy tensor in the updated state takes a diagonal form and the form it takes can be read directly from (21). For instance, for the energy density of a massless field we have in the left Rindler wedge

$$\begin{aligned} \langle f|\hat{T}_{\eta\eta}(x)f\rangle &= \Re \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{e^{-\pi\omega_p/a} e^{-\pi\omega_\kappa/a}}{(1 - e^{-2\pi\omega_p/a})(1 - e^{-2\pi\omega_\kappa/a})} \\ &\times \left(v_{I\bar{p}}(y) \partial_\eta v_{II\bar{\kappa}}(x) \partial_\eta \tilde{v}_{II\bar{p}}(x) \tilde{v}_{I\bar{\kappa}}(y') + e^{2a\xi} \sum_{i=1}^3 g^{ii} v_{I\bar{p}}(y) \partial_i v_{II\bar{\kappa}}(x) \partial_i \tilde{v}_{II\bar{p}}(x) \tilde{v}_{I\bar{\kappa}}(y') \right. \\ &+ v_{I\bar{p}}(y) \partial_\eta \tilde{v}_{II\bar{p}}(x) \partial_\eta v_{II\bar{\kappa}}(x) \tilde{v}_{I\bar{\kappa}}(y') + e^{2a\xi} \sum_i^3 g^{ii} v_{I\bar{p}}(y) \partial_i \tilde{v}_{II\bar{p}}(x) \partial_i v_{II\bar{\kappa}}(x) \tilde{v}_{I\bar{\kappa}}(y') \\ &+ v_{I\bar{p}}(y) \partial_\eta \overline{v_{II\bar{\kappa}}(x)} \partial_\eta \tilde{v}_{II\bar{p}}(x) \overline{\tilde{v}_{I\bar{\kappa}}(y')} + e^{2a\xi} \sum_i^3 g^{ii} v_{I\bar{p}}(y) \partial_i \overline{v_{II\bar{\kappa}}(x)} \partial_i \tilde{v}_{II\bar{p}}(x) \overline{\tilde{v}_{I\bar{\kappa}}(y')} \\ &\left. + \overline{v_{I\bar{p}}(y)} \partial_\eta \overline{\tilde{v}_{II\bar{p}}(x)} \partial_\eta v_{II\bar{\kappa}}(x) \tilde{v}_{I\bar{\kappa}}(y') + e^{2a\xi} \sum_i^3 g^{ii} \overline{v_{I\bar{p}}(y)} \partial_i \overline{\tilde{v}_{II\bar{p}}(x)} \partial_i v_{II\bar{\kappa}}(x) \tilde{v}_{I\bar{\kappa}}(y') \right), \quad (22) \end{aligned}$$

where \Re denotes the real part and with $g^{11} = e^{-2a\xi}$ and $g^{22} = g^{33} = 1$. In the massive case, one adds the term

$$\frac{1}{2} m^2 e^{2a\xi} \Delta_L(x, x) \quad (23)$$

to the right-hand side of eq. (22).

Similar expressions can be obtained for the components $\langle f|\hat{T}_{\xi\xi}(x)f\rangle$, $\langle f|\hat{T}_{yy}(x)f\rangle$ and $\langle f|\hat{T}_{zz}(x)f\rangle$ in the left Rindler wedge, and for the four non-vanishing components in the right Rindler wedge.

Instead of spelling out in detail all of the remaining components, we point out that eq. (22) suffices to a central point of the paper, which is that a measurement that occurs in the right Rindler wedge has non-trivial effects on the causally-disconnected left Rindler wedge. It is natural to ask “when” or “where” in spacetime the state collapses after a detector has detected

a Rindler particle. This is far from obvious, but a natural assumption seems to be that the state collapses along a Cauchy surface of spacetime (see [25]) intersecting the “detection event” on the right Rindler wedge and extending into the left Rindler wedge.

We should emphasise that regardless of “how big” this change might be, it is a principled statement that by including state collapses with semiclassical gravity has produced an abrupt change in the a region causally disconnected from where the measurement took place, in this case by means of a detector click.

One can see by direct inspection of (22) and (23) that for high accelerations there exists spacetime regions in the left Rindler wedge where the expectation value of the energy density becomes large. For example, for sufficiently small $\tilde{\xi}$, say $\tilde{\xi} \sim 1/a$, the Rindler modes do not exhibit the large-argument suppression of the MacDonald function, but the integrand factors $e^{-\pi\omega/a}/(1 - e^{-2\pi\omega/a})$ exhibit the behaviour

$$\frac{e^{-\pi\omega/a}}{1 - e^{-2\pi\omega/a}} = \frac{1}{2 \sinh(\pi\omega/a)} = \frac{a}{2\omega} + O(\omega/a). \quad (24)$$

This observation is consistent with what one would expect in the long interaction time limit case presented in [15], as can be seen from eq. (3.29) in that paper in the small $\beta = 2\pi/a$ regime.

4 Energetic considerations

In this section we revisit some of the energetic considerations discussed in [15] but focusing on the various possible *individual outcomes* of the “detection attempts”, rather than on the ensemble averages of detector measurement outcomes, which are the quantities considered in most of the discussion of said reference.

Following [15] and to simplify the discussion we will consider the case of (ensembles of) harmonic oscillators, and entangled pairs of harmonic oscillators instead of quantum fields. There is no loss of conceptual clarity in doing so, but the treatment is mathematically less involved.

Consider a harmonic oscillator with energy eigenstates $\{|n\rangle\}$ with renormalised energies $n\epsilon$ (i.e., we removed the zero point or ground state energy for simplicity of analysis), and a detector with two states $|\downarrow\rangle$ and $|\uparrow\rangle$ with energy levels 0 and ϵ respectively. Let us assume that the initial state of the combined system is:

$$|\Psi\rangle = N \left(|0\rangle \otimes |\downarrow\rangle + \frac{1}{\sqrt{\alpha}} |n\rangle \otimes |\uparrow\rangle \right), \quad (25)$$

with $N^2 = 1/(1 + \alpha^{-1})$. The expectation value of the energy is $\langle E \rangle_\Psi = \frac{n\epsilon}{1+\alpha}$.

If an observer finds the detector in the unexcited state $|\downarrow\rangle$, the value of the energy becomes $\langle E \rangle_{\text{unex}} = 0$, and the probability for this is $P_{\text{unex}} = N^2$. On the other hand if they find the detector in the excited state, the energy becomes $\langle E \rangle_{\text{ex}} = n\epsilon$ and the probability for this is $P_{\text{ex}} = N^2/\alpha$. The average result is of course $\langle E \rangle = \frac{n\epsilon}{1+\alpha}$, which is the same as $\langle E \rangle_\Psi$. We note however that in each specific case (i.e., for each specific outcome of the observation) the actual value of the energy differs from $\langle E \rangle_\Psi$, the expectation value of the initial state of the combined system. This is of course not surprising, given that the state $|\Psi\rangle$ is not an eigenstate of the total energy operator.

As has been argued in [27], the relevant issue regarding energy conservation is not its conservation “on average”, but its conservation on each single individual instance of any experiment. In this sense the possibility of preparing a state such as $|\Psi\rangle$ and to make the observations

described above already set serious doubts about the general validity of anything like a law of “energy conservation” in the quantum setting.

4.1 Proper mixture

Consider now the case of a system in thermal equilibrium at temperature T . We take the system in question once again to be a simple harmonic oscillator. This corresponds, as is traditionally treated on statistical mechanic textbooks, to an ensemble (a canonical ensemble) of identical particles and might be described by the proper mixture:

$$\rho = N \sum_{n=0}^{\infty} e^{-\beta n \epsilon} |n\rangle \langle n|, \quad (26)$$

where $N = 1 - e^{-\beta \epsilon}$ is a normalisation constant ensuring $\text{Tr} \rho = 1$. The mean energy is then $\langle E \rangle_T = \frac{\epsilon e^{-\beta \epsilon}}{1 - e^{-\beta \epsilon}}$. Let us now consider a two level detector (as in the previous discussion) initially in the un-excited state (and vanishing energy) and make it interact with our thermal ensemble (for this purpose we in fact consider an ensemble of identical detectors). The initial situation will thus be described by the density matrix:

$$\rho \otimes |\downarrow\rangle \langle \downarrow| = \left[\left(1 - e^{-\beta \epsilon}\right) \sum_{n=0}^{\infty} e^{-\beta n \epsilon} |n\rangle \langle n| \right] \otimes |\downarrow\rangle \langle \downarrow|. \quad (27)$$

After letting the system interact for a suitably long time we will find a result analogous to that encountered in the equation (3.25) of [15]. That is:

$$\rho_{\text{late}} = N_{\text{late}} \sum_{n=0}^{\infty} e^{-\beta n \epsilon} \left[|n\rangle \otimes |\downarrow\rangle - i\gamma\sqrt{n}|n-1\rangle \otimes |\uparrow\rangle + \dots \right] \left[\langle n| \otimes \langle \downarrow| + i\gamma\sqrt{n}\langle n-1| \otimes \langle \uparrow| + \dots \right] \quad (28)$$

where γ is a small parameter representing the strength and time duration of the interaction and N_{late} is a normalisation constant ensuring $\text{Tr} \rho_{\text{late}} = 1$.

4.2 Improper mixture

Consider now the situation in which we are informed that the detector is excited. To do this we simply compute a partial trace after applying the corresponding projector $|\uparrow\rangle \langle \uparrow|$. The resulting density matrix up, to first order in the expansion is

$$\rho_{\text{post}} = N_{\text{post}} \sum_{n=1}^{\infty} e^{-\beta n \epsilon} n |n-1\rangle \langle n-1|, \quad (29)$$

where $N_{\text{post}} = \left(\frac{e^{-\beta \epsilon}}{(1 - e^{-\beta \epsilon})^2} \right)^{-1}$ is still another normalisation constant ensuring $\text{Tr} \rho_{\text{post}} = 1$.

We note that the absence of the first term ought be considered to be the result of both the measurement of the excitation level of the excited detector and the post selection within the ensemble in which the elements containing the un-excited detector state were removed. In other words, we have that in going from eq. 28 to eq. 29 actually modifies the set of systems composing the ensemble itself.

The next observation is that the resulting ensemble, represented by the density matrix 29 is not longer thermal, which in turn implies that the expectation value of the energy

$$\text{Tr}(\hat{H} \rho_{\text{post}}) = N_{\text{post}} \sum_{m=0}^{\infty} e^{-\beta(m+1)\epsilon} (m+1)m\epsilon = \frac{2\epsilon e^{-\beta \epsilon}}{(1 - e^{-\beta \epsilon})}. \quad (30)$$

differs slightly from that of a truly thermal ensemble ($\langle E \rangle_T = \frac{\epsilon e^{-\beta \epsilon}}{1 - e^{-\beta \epsilon}}$). We note again that even after adding the energy of excitation of the detector ϵ the expectation value of the energy has changed as the result of the measurement (and post-selection).

Finally let us consider the case of a single pair of entangled harmonic oscillators with “thermal weights”. That is the case of a pure state, which upon consideration of the reduced density matrices for each oscillator would result in a thermal density matrix, but of course one of an improper nature. Let us add a detector arranged to interact just with the harmonic oscillator (II) and which is initially prepared in the unexcited state, so the state of the complete system is:

$$|\Psi\rangle_{\text{initial}}^{(T)} = A_{\text{initial}} \sum_{n=0}^{\infty} e^{-\beta n \epsilon} |n\rangle^{(I)} \otimes |n\rangle^{(II)} \otimes |\downarrow\rangle, \quad (31)$$

where $A_{\text{initial}} = \sqrt{1 - e^{-2\beta\epsilon}}$. After letting the system interact for a suitably long time, the state of the system will be :

$$|\Psi\rangle_{\text{late}}^{(T)} = A_{\text{late}} \sum_{n=0}^{\infty} e^{-\beta n \epsilon} |n\rangle^{(I)} \otimes \left[|n\rangle^{(II)} \otimes |\downarrow\rangle - i\gamma\sqrt{n} |n-1\rangle^{(II)} \otimes |\uparrow\rangle + \dots \right]. \quad (32)$$

If we project on the subspace corresponding to the excited detector (ignoring the irrelevant factor i and ensuring that the final state is normalised) we find:

$$|\Psi\rangle_{\text{pos-tsel}}^{(T)} = A_{\text{post-sel}} \sum_{n=0}^{\infty} e^{-\beta n \epsilon} |n\rangle^{(I)} \otimes \left[\sqrt{n} |n-1\rangle^{(II)} + \dots \right], \quad (33)$$

where in this case $A_{\text{post-sel}} \approx \left(\frac{e^{-2\beta\epsilon}}{(1 - e^{-2\beta\epsilon})^2} \right)^{-1/2}$.

Now the expectation values of the energy of the harmonic oscillator I is $\langle E \rangle_I = \frac{\epsilon(1 + e^{-2\beta\epsilon})}{(1 - e^{-2\beta\epsilon})}$, which is different from that of a purely thermal state, while the expectation value of the harmonic oscillator II is $\langle E \rangle_{II} = \frac{2\epsilon e^{-2\beta\epsilon}}{(1 - e^{-2\beta\epsilon})}$. Note that if we add the energy of the excited detector, ϵ , we have that $\langle E \rangle_{II} + \epsilon = \langle E \rangle_I$. On the other hand the expectation energies of both cases have definitely changed as a result of the “projection”.

At this point we might find nothing seriously puzzling because we have been dealing with systems that are not initially in energy eigen-states. The change in the case of the harmonic oscillator II in the above example, is of course a bit puzzling due to the fact that this oscillator has not been made to directly interact with a detector to bring about the “projection”, however this is nothing more than the usual change occurring as a result of quantum entanglement with a second system which has been subjected to a measurement.

It is worth noting that, in dealing with mixed states it is only when the expectation of the energy momentum is extracted from an improper mixture (and thus, indirectly from a pure state) that it make sense to use it in semiclassical gravity. If this is done with a proper mixture what we would obtain is indeed an average value of that quantity over some ensemble (such as in the case of the many realizations involved in stochastic gravity) and then self consistency will be in doubt¹.

The situation considered in Sec. 4 is, however a bit more troublesome as it seems to have the potential for a serious violation of our fundamental ideas about relativity, in particular, the potential to offer a path for superluminal communication.

5 Faster than light signaling?

It is widely agreed that our world contains non-local features. This is reflected for instance in the violations of Bell’s inequalities[28], which have been experimentally confirmed by now in several experiments [29, 30, 31, 32, 33]. While even simpler theoretical settings, such as the

¹In fact we do not know what it means to make averages over collections of space-times and the non-linearity of GR clearly cast serious doubts that say Einstein’s equations would be preserved under any kind of averaging one might want to consider.

GHZ construction [34] (which have not been experimentally realised due to technical difficulties) are expected to provide further and even more transparent evidence for non-locality. See for instance the discussion about the GHZ scheme in [35].

Nevertheless, there are a number of arguments and general widespread conviction that such non-locality cannot be exploited to communicate superluminally. Indeed, the non-locality present in the situation examined by Bell does not allow for superluminal communication. Faster-than-light communication would force us to revise the very foundations of special relativity – and of physics as a whole.

In any case, “textbook” quantum mechanics and its alternatives seeking to resolve the measurement problem do not seem to offer paths that allow for faster-than-light communication.

The change that we have noted in the expectation value of the stress-energy tensor in the left Rindler wedge (region II) due to measurements in the right Rindler wedge (region I) – if detectable – however seems to provide a path for superluminal communication, which we describe in the following *gedankenexperiment*:

Gedankenexperiment: Suppose a linearly uniformly accelerated observer in the right Rindler wedge, say Alice, is equipped with a highly-efficient detector for which it is highly probable to detect a Rindler particle and non-detection is negligible. This enables Alice to change the expectation value of the stress-energy tensor in the left Rindler wedge by turning her detector on or to decide not to do it by not switching on her detector. A causally-disconnected observer, say Bob, in the left Rindler wedge that can probe, either the state of the field or more specifically, the expectation value of stress-energy tensor (for example by probing the gravitational field with a torsion balance), would then be able to infer whether Alice has or has not turned on her detector in the right Rindler wedge. This seems to be a path for achieving superluminal communication between Alice and Bob.

A simple specific protocol that realises the above *Gedankenexperiment* can be thought of as follows.

The simplest specific protocol for superluminal communication is the following. Alice and Bob are given instructions that Alice could send to Bob a signal (or instance that she has decided on something and the answer is “yes”) by turning on her detector. She would *not* turn her detector at all if the answer is “no”. Bob will then monitor the value of the expansion of nearby geodesics (which might be hard technically as he is not moving along one), by say looking at freely falling particles he is continuing releasing. Bob would have to be very careful to ensure that nothing he does generates in his surroundings any energy momentum tensor that could mimic that associated with the change in the state resulting from a detection of a field quantum by Alice’s detector (or to take any energy momentum he generates into account so as to be able to distinguish that from the one associated with the latter). If at any time in his world line he detects such geodesic expansion he would know Alice’s decision is “yes”. The point is that no matter when along his world-line that would happen the communication would be superluminal as Alice and Bob are never in causal contact. This is of course a rather poor signaling protocol because Bob could eventually know if Alice’s decision is “yes” but he would never know if Alice’s decision is “no”.

We can however remedy this by having Alice, say, use two different kind of detectors (with say vastly different energy of excitation, say one of them being tenfold, or having different orientation), which could produce two different changes in the energy momentum tensor at Bob’s location. The detection of one of the two changes would mean the answer is “yes” and the other the answer is “no”. This would correspond to a complete 1-bit signaling.

Again the efficiency and reliability of this protocol depends strongly on the magnitude of Alice’s and Bob absolute accelerations a_A and a_B as well as the change on the state of the quantum field induced by the excitation of Alice’s detector (which the timing we recall she

can not control, as she can only decide whether to turn or not the detector on). This change would depend also on Alice detector's which we can modify by selecting, say, the energy gap, Ω , with which the detector operates.

Ignoring switching effects, reasonably good performance for the detector could be achieved by setting the energy gap near the Unruh temperature (in natural units), since for long-time interactions the transition probability of the detector should behave as a Planckian distribution [5]. This part can be modified and we see no obstacle, in principle, to make this quantity arbitrarily large. At the same time the distance D_A along a hypersurface orthogonal to ξ from Alice's intersection world-line and say the bifurcating surface of the Killing Horizon is determined by a_A but decreases as the later increases it, so for larger D_A the protocol becomes poorer.

Optimizing the protocol performance with respect to a_B presumably will have to balance the fact that the analogous distance from Bob to the bifurcating horizon decreases with increasing a_B and the detailed functioning of the device he uses in detecting the expansion of nearby geodesics (an analysis that does not seem very easy to carry out).

In any event we think that even if the optimal level of efficiency is not very high, that does not detract from the fact that in principle there are no trivial or evident obstacles for he protocol to work at some level of reliability and any nonzero value of that would represent the opening of a door for superluminal communication. Indeed, there exist currently serious proposals to measure the gravitational field with high precision, e.g. [36]. We will however discuss at further down the manuscript what seem to be the most natural possibilities for nature to prevent in principle the operation of such protocol.

In any event it seems that, even if not highly effective, the possibility of Alice signaling to Bob in a superluminal way should be considered problematic in view of the implications that would have for our understanding of the world. In fact, the situation is aggravated by the fact that even if the detector does not detect a particle, there are higher-order effects (in the coupling constant) that induce changes in the stress-energy tensor on the left Rindler wedge [37].²

Thus, it seems imperative to consider the caveats that might provide a path to avoid such a problematic communication protocol. Before we start let us make some observations regarding semiclassical gravity. Semiclassical gravity is a formalism in which the gravitational field is taken as dynamical but treated in classical terms as the metric of a spacetime (M, g_{ab}) in general relativity, while the matter is treated in the language of quantum fields on such spacetime. The metric is taken to satisfy the semiclassical version of Einstein's field equations

$$G_{ab}(x) = 8\pi G \langle T_{ab}(x) \rangle, \quad (34)$$

with $\langle T_{ab}^{\text{ren}} \rangle$, the expectation value of the matter's quantum stress-energy tensor in a suitable quantum state, while the quantum matter obeys the dynamics of QFT. See e.g. [38] for a review.

In incorporating state collapses in the semiclassical gravity context we remark that there is an issue, which we have touched on briefly above. The point is that once something like state reduction is considered (be it in the Copenhagen approach or in any of the spontaneous collapse theories models, e.g. [39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50]), one has the ambiguity of which state should be used in computing the right-hand side of eq. (34) should be employed. In other words, it seems natural that if we are interested in the value of the left-hand side of eq. (34) at a point x , the right-hand side should be computed using a state associated with a Cauchy surface that passes trough x , but of course there are infinite such surfaces and the proposal to consider eq. 34 even as an approximation must be completed

²Although it is not clear if the effect when no particle is absorbed is an artifact of perturbation theory.

with a detailed prescription in this regard in order to at least have a well-defined proposal [51, 52, 53, 54, 55]. We should emphasize however that no matter what Cauchy surface is used, a portion of the Cauchy surface must extend to the left Rindler wedge. Thus, assuming that the collapse occurs along any arbitrary Cauchy surface, the left Rindler wedge must include a “pre-collapse” and a “post-collapse” region with different spacetime geometry – according to semiclassical gravity.

Let us now offer, and briefly discuss what we think is the list of serious set of options to be considered, that could help in avoiding the conclusion of super-luminal signaling:

- (i) *Strong enough departures from semiclassical gravity, at least in region II:*
- (ii) *Existence of effects that are indistinguishable from observations of the change in the stress-energy in region II:*
- (iii) *A fundamental undetectability of the change of the state in region II, and in particular that of expectation value of the stress-energy tensor in said region:*
- (iv) *Some fundamental impediment of the construction of the set up.*

Let us now briefly discuss the options (i)-(iv) considered above.

(i) First, and in order to address the issue, we must clarify what is meant by the words “strong enough”. We take that to indicated that as a result of some unknown aspect of physics (with originating, say, in aspects of quantum gravity), the **validity of eq. (34) with the right-hand side in the updated state, cf. the discussion of sec. 3, would be violated** to such a large degree that, for any design of Bob measuring instrument, the expected result will be modified by a factor of at least the same order of magnitude. **As we have pointed out in sec. 3, it is possible to make the expectation value of the stress-energy tensor at points in region II arbitrarily large in the post-collapse state by increasing the detector gap.** However, it must be acknowledged that if Ω becomes large enough modeling the state collapse in semiclassical gravity along a Cauchy surface could be questioned on the basis that a large violation of the stress-energy tensor along a Cauchy surface is too strong a deviation from the semiclassical regime. Thus, one might argue that although one could trust semiclassical gravity before and after the state collapse, there is no way to model the system “during” the state collapse in semiclassical terms, and a more refined understanding of how to model state collapses in semiclassical gravity could be required. We note however that even if that is the case, such conclusion will not prevent the super-luminal signaling as all we need is for Bob to detect the effect at any time.

On the other hand it seems rather unlikely that arbitrarily large departures of semiclassical gravity will be associated with such simple and rather common situation. The full analysis of the question thus requires consideration of the means by which Bob might detect the corresponding change in the spacetime metric in region II ³. As it seems clear that a modification of the spacetime curvature might, in principle, be measured by the study of the geodesic deviation equation and in particular the expansion of the congruence of geodesics in that region, it appears that the question will have to be connected at least to some fundamental limitation on the validity of the notion of test point particles following geodesics of the underlying metric. There are of course some limitations on that arising from simple quantum considerations about the description of so called free point particles which in fact negate the notion of well defined trajectories.

³We will not study here in detail the detection method to be employed by Bob, but just point out that interesting ideas to measure small gravitational effects have been used in say searches for deviation of the universality of free fall, and also note a recent proposl to look for Plank scale dark matter [72].

Moreover, as discussed in [56], the standard quantum mechanical minimal de-localization of a quantum particle (characterized for instance by its Compton wavelength) implies such particle ought not to be considered as point like, and as is well known, in general, even at the classical level, extended objects fail to follow geodesics. It is unclear at this point if these kind of considerations will be enough to dismiss our example, given the fact that, as noted, the effect could be made as large as one wants and the time available for Bob to make the measurement is arbitrarily large.

(ii) Here we must consider the existence of other effects that might not be effectively distinguished from the changes resulting from the collapse. Those might be intrinsically associated with the gravitational effects of either the measuring devices that Bob would be introducing in order to detect the changes in the spacetime metric in region II. They also might be associated with mere existence of (and the non-vanishing stress-energy tensor) corresponding to Alice and her detectors, whose own gravitational effects we have been neglecting so far, although classically these effects will in principle propagate causally and not affect region II, it is likely that deviations from Minkowski spacetime reflect on the *non-local* state of the field from a semiclassical-gravity viewpoint in such a way that the quantum state of the field cannot be in principle the Minkowski vacuum state to begin with. Moreover, the stress-energy of Alice and her apparatus must be considered in the constraints of the theory, and the spacetime will have a non-vanishing ADM mass, reflected in the asymptotic behavior in all spacetime directions.

Another possibility is to consider that the model used for Alice’s detector, although standard in the literature, is a local model (in Region I) of the detector. Treating Alice’s detector as a quantum field yields a detector model with support in both Regions I and II. In the Heisenberg picture we know that the existence of a field-like detector for Alice will only affect the dynamics in the causal future of the coupling region between Alice’s detector (or probe) and the quantum field considered as a system. This can also be formalised in the algebraic QFT language [22]. However, as emphasised in [22], a local and covariant measurement scheme is only able to describe the *probe-system measurement chain* under the assumption that somewhere, someone knows how to measure something. Thus, it seems that the issue of sending a faster-than-light signal by the “act of measurement” cannot be resolved by simply changing the detector model. Sorkin has pointed out in his “impossible measurements” protocol [57] that the existence of spatially extended detectors, would make faster than light signaling almost unavoidable. See however [58].

(iii) This possibility could result from a variety of reasons. For instance, it might be that the appropriate versions of semiclassical gravity that will hold are such that the expectation value of the stress-energy tensor at any point, is taken on states associated with hypersurfaces which would not incorporate the change in the state resulting from a collapse or a measurement on region I. One such option would require to take the expectation value of the stress-energy tensor appearing at the right-hand side of Einstein’s semiclassical equation in x to be computed using the state corresponding to the something like the hypersurface $\partial J^-(x)$ (the boundary of the causal past of x), and that, at the same time, the detailed theory characterising the measurement (i.e., something like the spontaneous collapse theories considered in [59, 60, 61]), is such, that the state associated with $\partial J^-(x)$ is unaffected. A scheme of that kind would ensure that the measurement of Alice would have no consequences from semiclassical gravity for spacelike-separated events. Such proposal is not without difficulties, one of which is the fact that in general $\partial J^-(x)$ is not a Cauchy surface, and another is that it is not smooth at x . The first difficulty might be resolved if one has some good reason to consider that there is a certain initial state associated with an initial Cauchy hypersurface Σ_{in} , which might be considered as the initially “prepared ” state, or perhaps *the initial state of the universe* or something like that, and then to consider, computing the expectation values of quantities of

interest at x , using the state associated with the surface $(I^+(\Sigma_{in}) \cap \partial J^-(x)) \cup (\Sigma_{in} - J^-(x))$, while the second problem might be dealt with, with an adjustment of the recipe based on the taking of appropriate limits of a succession of smooth Cauchy hypersurfaces that have as limit (in a suitable sense), the hypersurface indicated above. The issue is however quite delicate as illustrated by the discussions in [51], the pursuit of which lies outside the scope of the present work.

An argument in favour of the impossibility of detecting the signal is the following. One can assume that stress-energy conservation must hold *on average*, i.e., for successive measurements, which prevents one from extracting arbitrarily large amounts of energy from the quantum field by detector measurements, as argued in [15]. Thus, an arbitrarily-efficient detector must produce arbitrarily-small changes for the stress-energy tensor when it detects a Rindler particle. The issue now becomes whether there exists an “engineering window” where the efficiency of the detector and the amplitude of the signal can be compensated, such that a faster-than-light signal is measurable by Bob and that such signal can be sent with sufficient certainty by Alice. On this point, it follows from the inequality $\Delta E \Delta T \geq \hbar$ that to detect such a small signal Bob requires a large amount of time. On the other hand, in principle, Bob has an infinite time to measure an arbitrarily small signal, for he can orbit along a Killing vector boost orbit in the left Rindler wedge, and one could argue that such engineering window can always be found. However, any physical signal sent by Alice must decay as it approaches \mathcal{I}^+ . Thus, it is expected that the signal will become weaker as Bob’s proper time elapses, and this might render the signal “effectively undetectable”.

(iv) Finally there could be aspects of the set up that would make it simply unfeasible. One possibility we should consider is the following. In order for Bob to be sure that the change in the energy momentum he observes corresponds indeed to a signal that was sent by Alice, he must be sure that similar signals are not reaching him coming from elsewhere. That is, he must be quite sure that the state of the quantum field prior to Alice’s manipulation of her detector is the Minkowski vacuum, and as he must be ready to receive Alice’s signal at any point in his world line, he must be sure that such characterization of the state of the quantum field must be the appropriate one up to arbitrarily distant regions of “space”. So there must be a preparation of the initial state of the quantum field on an extremely large spatial region occurring well before the whole protocol is even started (and if we want to ensure Bob has in effect arbitrarily long time to make the detection it seems the preparation of the state ought to encompass a full Cauchy hypersurface. However, as already noted, the simultaneous (in some reference frame) measurements of observables associated with such extended spatial regions – say, as a means of state preparation – are known to be rather problematic in their own right as argued by Sorkin[57] (again, see however [58]).

In any case, such arguments must be considered with care, because, in principle, even a very unreliable communication protocol which allows faster than light communications would be quite problematic. We think these ideas deserve much deeper exploration.

6 Final remarks

We have revisited the issue of detection of a Rindler particle by an accelerator detector confined to the right Rindler wedge, focusing attention on the implications of the reduction of the state associated with the actual detection or what is often referred to as the measurement part of the process. We have noted that the concomitant modification of the state of the quantum field, imply changes in the (expectation value of the renormalised) stress-energy tensor in both Rindler wedges.

Concerning the right Rindler wedge, it is interesting that the resulting state and stress-

energy tensor expectation values become non-thermal which is, in a sense, easy to understand as result of the disruption brought by the interaction with the detector. Of course, one expects that in the long run further interaction between the field and detector or among the field modes themselves will bring the system to a new state of thermal equilibrium.

Concerning the left Rindler wedge, however, the change in the stress-energy tensor expectation value is more problematic, as it seems to open a possibility for faster than light communication. We have noted some of what we see as the most natural possibilities to avoid such conclusion, but further studies of these issues are required in order to get to more definite and solid conclusions.

For the moment, we can only stress that the fuzziness in the theoretical characterization of the act of measuring in quantum theory can bring about complications even in apparently innocuous circumstances, such as in the context of the detection of a Rindler particle in Minkowski spacetime. Despite being overlooked in many practical applications in physics, the literature on the measurement problem is quite large, and the positions taken in its face are quite varied. We do not intend to discuss them in detail. It suffices for us to mention that the paths to overcome the measurement problem have been classified by Maudlin [62] as follows: It is internally inconsistent to hold simultaneously the following three propositions about a quantum theory:

- (i) The description of a physical system as provided by the quantum state or wave function is complete.
- (ii) The evolution of the quantum state is always dictated by the Schrödinger equation (or its relativistic generalisations).
- (iii) Individual experiments produce definite (even if often unpredictable) results.

Thus, one must negate at least one of (i)-(iii) above. Negating (i) leads down, in general, the path of the so-called *hidden variable theories*, of which the de Broglie-Bohm theory is the best known example [63]. Negating (ii) implies that the collapse or reduction of the wave function plays a central rôle, as in the Copenhagen textbook interpretation, as well as in so-called *spontaneous collapse* or *dynamical state reduction* theories, such as GRW, CSL, Penrose-Diósi, etc [64, 65, 66, 67, 59, 60, 61]. The negation of (iii) leads to *many-worlds*- or *many-minds*-type interpretations [68, 69].

In any case, we should restate that the superluminal tension relies on the application of two central hypotheses in this work. The first one is that measurements induce the collapse of the wavefunction, and that such collapse occurs on a Cauchy surface of spacetime. The second one is that the problem studied in (nearly) flat spacetime is within the regime of applicability of semiclassical gravity. It is well possible that either one of the two hypotheses fail, or that they cannot be taken to hold together. If this is the case, it would seem that there exist apparently innocuous situations, such as the one here studied, for which a correct theoretical description of the evolution of the system must resort to quantum gravity. This seems to be in agreement with conclusions drawn from [70, 71]. However if that is the case it seems likely that other situations that are often discussed using semiclassical language, for instance Hawking radiation and the backreaction in black holes (even at early times), might require reconsideration as well.

Finally, it is our hope to draw the attention of the community to these delicate issues that afflict our understanding of quantum theory in general and of QFT in particular, especially in gravitational contexts.

Acknowledgements

We gladly thank Prof. George Matsas for his careful reading of a previous version of this work and very useful comments, and Prof. Stephen A. Fulling for very stimulating discus-

sions, as well as Ruth E Kastner for pointing out a few mistakes in the original version of this work. B.A.J.-A. is supported by a CONACYT postdoctoral fellowship. D.S. acknowledges partial financial support from PAPIIT-UNAM, Mexico (Grant No.IG100120); the Foundational Questions Institute (Grant No. FQXi-MGB-1928); the Fetzer Franklin Fund, a donor advised by the Silicon Valley Community Foundation. B.A.J.-A. and D.S. acknowledge the support of CONACYT grant FORDECYT-PRONACES no. 140630. This article has been prepared as a contribution to the Topical Collection “Acceleration and Radiation: Classical and Quantum, Electromagnetic and Gravitational” of the Journal Symmetry, edited by Prof. Stephen A. Fulling.

A The Minkowski vacuum in the Rindler wedges

The initial state is taken to be the Minkowski vacuum, whose Wightman function takes the following form in the right Rindler wedge

$$\langle \Omega_M | \hat{\Phi}(x) \hat{\Phi}(x') \Omega_M \rangle = \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3 \kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3 \kappa' \langle \Omega_M | \left(v_{I\vec{\kappa}}(x) \hat{a}_{R\vec{\kappa}} + \overline{v_{I\vec{\kappa}}(x)} \hat{a}_{R\vec{\kappa}}^* \right) \left(v_{I\vec{\kappa}'}(x') \hat{a}_{R\vec{\kappa}'} + \overline{v_{I\vec{\kappa}'}(x')} \hat{a}_{R\vec{\kappa}'}^* \right) \Omega_M \rangle. \quad (35)$$

Using the relations [9, eq. 2.125-2.127],

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}}^* \hat{a}_{R\vec{\kappa}'} \Omega_M \rangle = \langle \Omega_M | \hat{a}_{L\vec{\kappa}}^* \hat{a}_{L\vec{\kappa}'} \Omega_M \rangle = \left(e^{2\pi\omega/a} - 1 \right)^{-1} \delta^3(\vec{\kappa} - \vec{\kappa}'), \quad (36a)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}} \hat{a}_{R\vec{\kappa}'}^* \Omega_M \rangle = \langle \Omega_M | \hat{a}_{L\vec{\kappa}} \hat{a}_{L\vec{\kappa}'}^* \Omega_M \rangle = \left(1 - e^{-2\pi\omega/a} \right)^{-1} \delta^3(\vec{\kappa} - \vec{\kappa}'), \quad (36b)$$

$$\langle \Omega_M | \hat{a}_{L\vec{\kappa}} \hat{a}_{R\vec{\kappa}'} \Omega_M \rangle = \langle \Omega_M | \hat{a}_{L\vec{\kappa}}^* \hat{a}_{R\vec{\kappa}'}^* \Omega_M \rangle = \left(e^{\pi\omega/a} - e^{-\pi\omega/a} \right)^{-1} \delta^3(\vec{\kappa} - \vec{\kappa}'), \quad (36c)$$

(and zero otherwise) we obtain that

$$\langle \Omega_M | \hat{\Phi}(x) \hat{\Phi}(x') \Omega_M \rangle = \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3 \kappa \left(\frac{v_{I\vec{\kappa}}(x) \overline{v_{I\vec{\kappa}}(x')}}{1 - e^{-2\pi\omega/a}} + \frac{\overline{v_{I\vec{\kappa}}(x)} v_{I\vec{\kappa}}(x')}{e^{2\pi\omega/a} - 1} \right). \quad (37)$$

Likewise, in the left Rindler wedge the Wightman function takes the form

$$\langle \Omega_M | \hat{\Phi}(x) \hat{\Phi}(x') \Omega_M \rangle = \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3 \kappa \left(\frac{v_{II\vec{\kappa}}(x) \overline{v_{II\vec{\kappa}}(x')}}{1 - e^{-2\pi\omega/a}} + \frac{\overline{v_{II\vec{\kappa}}(x)} v_{II\vec{\kappa}}(x')}{e^{2\pi\omega/a} - 1} \right), \quad (38)$$

where $v_{II\vec{\kappa}}$ are left Rindler modes.

B Normalisation of the updated state

We have seen in sec. 2 that if the detector clicks the updated state of the field becomes

$$|f\rangle = -i\mathcal{N} \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3 \kappa \int_I d\text{vol}(x) \zeta(x) \left(v_{I\vec{\kappa}}(x) \hat{a}_{R\vec{\kappa}} + \overline{v_{I\vec{\kappa}}(x)} \hat{a}_{R\vec{\kappa}}^* \right) |\Omega_M\rangle. \quad (39)$$

In this appendix, we see that the normalisation constant is given by eq. (14). We compute

$$\begin{aligned} \mathcal{N}^{-2} \langle f | f \rangle &= \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3 \kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3 \kappa' \int_I d\text{vol}(x) \int_I d\text{vol}(x') \overline{\zeta(x)} \zeta(x') \\ &\quad \times \langle \Omega_M | \left(v_{I\vec{\kappa}}(x) \hat{a}_{R\vec{\kappa}} + \overline{v_{I\vec{\kappa}}(x)} \hat{a}_{R\vec{\kappa}}^* \right) \left(v_{I\vec{\kappa}'}(x') \hat{a}_{R\vec{\kappa}'} + \overline{v_{I\vec{\kappa}'}(x')} \hat{a}_{R\vec{\kappa}'}^* \right) \Omega_M \rangle. \end{aligned} \quad (40)$$

Using the relations [9, eq. 2.125-2.126], we obtain that

$$\mathcal{N}^{-2}\langle f|f\rangle = \int_{\mathbb{R}^+\times\mathbb{R}^2} d^3\kappa \int_I \text{dvol}(x) \int_I \text{dvol}(x') \overline{\zeta(x)} \zeta(x') \left(\frac{v_{I\bar{\kappa}}(x)v_{I\bar{\kappa}}(x')}{1 - e^{-2\pi\omega/a}} + \frac{\overline{v_{I\bar{\kappa}}(x)}v_{I\bar{\kappa}}(x')}{e^{2\pi\omega/a} - 1} \right). \quad (41)$$

C The two-point function in the updated state in the right Rindler wedge

The updated stress-energy tensor can be computed from the two-point function in the updated state. In this appendix, we show that the two-point function in the updated state in the right Rindler wedge is given by

$$\langle f|\hat{\Phi}(x)\hat{\Phi}(x')f\rangle = \langle \Omega_M|\hat{\Phi}(x)\hat{\Phi}(x')\Omega_M\rangle + \Delta_R(x, x'). \quad (42)$$

where $\Delta_R(x, x')$ is given by eq. (53) below.

The two-point function in the right Rindler wedge reads

$$\begin{aligned} \langle f|\hat{\Phi}(x)\hat{\Phi}(x')f\rangle &= \mathcal{N}^2 \int_{\mathbb{R}^+\times\mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+\times\mathbb{R}^2} d^3\kappa' \int_{\mathbb{R}^+\times\mathbb{R}^2} d^3p \int_{\mathbb{R}^+\times\mathbb{R}^2} d^3p' \int_I \text{dvol}(y) \int_I \text{dvol}(y') \overline{\zeta(y)} \zeta(y') \\ &\times \langle \Omega_M| \left(v_{I\bar{p}}(y)\hat{a}_{R\bar{p}} + \overline{v_{I\bar{p}}(y)}\hat{a}_{R\bar{p}}^* \right) \left(v_{I\bar{\kappa}}(x)\hat{a}_{R\bar{\kappa}} + \overline{v_{I\bar{\kappa}}(x)}\hat{a}_{R\bar{\kappa}}^* \right) \\ &\times \left(v_{I\bar{\kappa}'}(x')\hat{a}_{R\bar{\kappa}'} + \overline{v_{I\bar{\kappa}'}(x')}\hat{a}_{R\bar{\kappa}'}^* \right) \left(v_{I\bar{p}'}(y')\hat{a}_{R\bar{p}'} + \overline{v_{I\bar{p}'}(y')}\hat{a}_{R\bar{p}'}^* \right) \Omega_M \rangle. \quad (43) \end{aligned}$$

Using [9, eq. 2.122-2.124] one can obtain the relations

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{R\vec{\kappa}_2} \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (44a)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{R\vec{\kappa}_2} \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle = 0, \quad (44b)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{R\vec{\kappa}_2}^* \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (44c)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{R\vec{\kappa}_2}^* \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle = 0, \quad (44d)$$

$$\begin{aligned} & \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2} \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle \\ &= \frac{1}{1 - e^{-2\pi\omega_1/a}} \frac{1}{1 - e^{-2\pi\omega_2/a}} (\delta(\vec{\kappa}_1 - \vec{\kappa}_3) \delta(\vec{\kappa}_2 - \vec{\kappa}_4) + \delta(\vec{\kappa}_1 - \vec{\kappa}_4) \delta(\vec{\kappa}_2 - \vec{\kappa}_3)), \end{aligned} \quad (44e)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2} \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle = \frac{1}{1 - e^{-2\pi\omega_1/a}} \frac{1}{1 - e^{-2\pi\omega_3/a}} \delta(\vec{\kappa}_1 - \vec{\kappa}_2) \delta(\vec{\kappa}_3 - \vec{\kappa}_4) \quad (44f)$$

$$+ \frac{1}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-2\pi\omega_3/a}}{1 - e^{-2\pi\omega_3/a}} \delta(\vec{\kappa}_1 - \vec{\kappa}_4) \delta(\vec{\kappa}_2 - \vec{\kappa}_3), \quad (44g)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2} \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle \quad (44h)$$

$$= \frac{e^{-2\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{1}{1 - e^{-2\pi\omega_4/a}} (\delta(\vec{\kappa}_1 - \vec{\kappa}_2) \delta(\vec{\kappa}_3 - \vec{\kappa}_4) + \delta(\vec{\kappa}_1 - \vec{\kappa}_3) \delta(\vec{\kappa}_2 - \vec{\kappa}_4)), \quad (44i)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2} \hat{a}_{R\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle \quad (44j)$$

$$= \frac{1}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-2\pi\omega_4/a}}{1 - e^{-2\pi\omega_4/a}} (\delta(\vec{\kappa}_1 - \vec{\kappa}_2) \delta(\vec{\kappa}_3 - \vec{\kappa}_4) + \delta(\vec{\kappa}_1 - \vec{\kappa}_3) \delta(\vec{\kappa}_2 - \vec{\kappa}_4)), \quad (44k)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2} \hat{a}_{R\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = \frac{e^{-2\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-2\pi\omega_3/a}}{1 - e^{-2\pi\omega_3/a}} \delta(\vec{\kappa}_1 - \vec{\kappa}_2) \delta(\vec{\kappa}_3 - \vec{\kappa}_4) \quad (44l)$$

$$+ \frac{e^{-2\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{1}{1 - e^{-2\pi\omega_2/a}} \delta(\vec{\kappa}_1 - \vec{\kappa}_4) \delta(\vec{\kappa}_2 - \vec{\kappa}_3), \quad (44m)$$

$$\begin{aligned} & \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2}^* \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle \\ &= \frac{e^{-2\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-2\pi\omega_2/a}}{1 - e^{-2\pi\omega_2/a}} (\delta(\vec{\kappa}_1 - \vec{\kappa}_3) \delta(\vec{\kappa}_2 - \vec{\kappa}_4) + \delta(\vec{\kappa}_1 - \vec{\kappa}_4) \delta(\vec{\kappa}_2 - \vec{\kappa}_3)), \end{aligned} \quad (44n)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2}^* \hat{a}_{R\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (44o)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2}^* \hat{a}_{R\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle = 0, \quad (44p)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2}^* \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle = 0, \quad (44q)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2}^* \hat{a}_{R\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (44r)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{R\vec{\kappa}_2}^* \hat{a}_{R\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0. \quad (44s)$$

Inserting eq. (44) into (43) we obtain that the two point function can be obtained as a sum of six contributions, each coming from one of the non-trivial expressions in eq. (44), i.e., it takes the form

$$\langle f | \hat{\Phi}(x) \hat{\Phi}(x') f \rangle(x, x') = \sum_{n=1}^6 G_{Rn}(x, x'). \quad (45)$$

Subtracting eq. (37) from this expression we obtain that

$$\Delta_R(x, x') = \langle f | \hat{\Phi}(x) \hat{\Phi}(x') f \rangle - \langle \Omega_M | \hat{\Phi}(x) \hat{\Phi}(x') \Omega_M \rangle = \sum_{n=1}^6 \Delta_{Rn}(x, x') \quad (46)$$

with

$$\Delta_{R1}(x, x') := \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{v_{I\bar{p}}(y) v_{I\bar{\kappa}}(x) \overline{v_{I\bar{p}}(x')} \overline{v_{I\bar{\kappa}}(y')}}{(1 - e^{-2\pi\omega_p/a})(1 - e^{-2\pi\omega_\kappa/a})}, \quad (47)$$

$$\Delta_{R2}(x, x') := \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{v_{I\bar{p}}(y) \overline{v_{I\bar{p}}(x)} v_{I\bar{\kappa}}(x') \overline{v_{I\bar{\kappa}}(y')}}{(1 - e^{-2\pi\omega_p/a})(1 - e^{-2\pi\omega_\kappa/a})}, \quad (48)$$

$$\begin{aligned} \Delta_{R3}(x, x') &:= \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{\overline{v_{I\bar{\kappa}}(y)} v_{I\bar{\kappa}}(x) v_{I\bar{p}}(x') \overline{v_{I\bar{p}}(y')} e^{-2\pi\omega_\kappa}}{(1 - e^{-2\pi\omega_\kappa})(1 - e^{-2\pi\omega_p})} \\ &+ \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{\overline{v_{I\bar{\kappa}}(y)} v_{I\bar{p}}(x) v_{I\bar{\kappa}}(x') \overline{v_{I\bar{p}}(y')} e^{-2\pi\omega_\kappa}}{(1 - e^{-2\pi\omega_\kappa})(1 - e^{-2\pi\omega_p})}, \end{aligned} \quad (49)$$

$$\begin{aligned} \Delta_{R4}(x, x') &:= \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{v_{I\bar{\kappa}}(y) \overline{v_{I\bar{\kappa}}(x)} v_{I\bar{p}}(x') \overline{v_{I\bar{p}}(y')} e^{-2\pi\omega_p}}{(1 - e^{-2\pi\omega_\kappa})(1 - e^{-2\pi\omega_p})} \\ &+ \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{v_{I\bar{\kappa}}(y) \overline{v_{I\bar{p}}(x)} v_{I\bar{\kappa}}(x') \overline{v_{I\bar{p}}(y')} e^{-2\pi\omega_p}}{(1 - e^{-2\pi\omega_\kappa})(1 - e^{-2\pi\omega_p})}, \end{aligned} \quad (50)$$

$$\Delta_{R5}(x, x') := \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{\overline{v_{I\bar{\kappa}}(y)} v_{I\bar{\kappa}}(x) \overline{v_{I\bar{p}}(x')} \overline{v_{I\bar{p}}(y')} e^{-2\pi\omega_\kappa} e^{-2\pi\omega_p}}{(1 - e^{-2\pi\omega_\kappa})(1 - e^{-2\pi\omega_p})}, \quad (51)$$

$$\Delta_{R6}(x, x') := \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \frac{\overline{v_{I\bar{\kappa}}(y)} \overline{v_{I\bar{p}}(x)} v_{I\bar{\kappa}}(x') \overline{v_{I\bar{p}}(y')} e^{-2\pi\omega_\kappa} e^{-2\pi\omega_p}}{(1 - e^{-2\pi\omega_\kappa})(1 - e^{-2\pi\omega_p})}. \quad (52)$$

Collecting, we can write

$$\Delta_R(x, x') = \Delta_{R1}(x, x') + \Delta_{R3}(x, x') + \Delta_{R5}(x, x') + \text{c.c.}, \quad (53)$$

which yields eq. (18).

D The two-point function in the updated state in the left Rindler wedge

The updated stress-energy tensor can be computed from the two-point function in the updated state. In this appendix, we show that the two-point function in the updated state in the right Rindler wedge is given by

$$\langle f | \hat{\Phi}(x) \hat{\Phi}(x') f \rangle = \langle \Omega_M | \hat{\Phi}(x) \hat{\Phi}(x') \Omega_M \rangle + \Delta_L(x, x'), \quad (54)$$

as in eq. (19).

The two-point function in the left Rindler wedge reads

$$\begin{aligned} \langle f | \hat{\Phi}(x) \hat{\Phi}(x') f \rangle &= \mathcal{N}^2 \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3\kappa' \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p \int_{\mathbb{R}^+ \times \mathbb{R}^2} d^3p' \int_I d\text{vol}(y) \int_I d\text{vol}(y') \overline{\zeta(y)} \zeta(y') \\ &\times \langle \Omega_M | \left(v_{I\bar{p}}(y) \hat{a}_{R\bar{p}} + \overline{v_{I\bar{p}}(y)} \hat{a}_{R\bar{p}}^* \right) (v_{II\bar{\kappa}}(x) \hat{a}_{L\bar{\kappa}} + \overline{v_{II\bar{\kappa}}(x)} \hat{a}_{L\bar{\kappa}}^*) \\ &\times (v_{II\bar{\kappa}'}(x') \hat{a}_{L\bar{\kappa}'} + \overline{v_{II\bar{\kappa}'}(x')} \hat{a}_{L\bar{\kappa}'}^*) \left(v_{I\bar{p}'}(y') \hat{a}_{R\bar{p}'} + \overline{v_{I\bar{p}'}(y')} \hat{a}_{R\bar{p}'}^* \right) \Omega_M \rangle. \end{aligned} \quad (55)$$

Using [9, eq. 2.122-2.124] one can obtain the relations

$$\begin{aligned} \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{L\vec{\kappa}_2} \hat{a}_{L\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle &= \frac{e^{-\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-\pi\omega_2/a}}{1 - e^{-2\pi\omega_2/a}} \tilde{\delta}(\vec{\kappa}_1 - \vec{\kappa}_3) \tilde{\delta}(\vec{\kappa}_2 - \vec{\kappa}_4) \\ &+ \frac{e^{-\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-\pi\omega_3/a}}{1 - e^{-2\pi\omega_3/a}} \tilde{\delta}(\vec{\kappa}_1 - \vec{\kappa}_2) \tilde{\delta}(\vec{\kappa}_3 - \vec{\kappa}_4), \\ \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{L\vec{\kappa}_2} \hat{a}_{L\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle &= 0, \end{aligned} \quad (56a)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{L\vec{\kappa}_2} \hat{a}_{L\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (56b)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{L\vec{\kappa}_2}^* \hat{a}_{L\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (56c)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{L\vec{\kappa}_2} \hat{a}_{L\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (56d)$$

$$\begin{aligned} \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{L\vec{\kappa}_2} \hat{a}_{L\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle &= \frac{1}{1 - e^{-2\pi\omega_1/a}} \frac{1}{1 - e^{-2\pi\omega_2/a}} \delta(\vec{\kappa}_1 - \vec{\kappa}_4) \delta(\vec{\kappa}_2 - \vec{\kappa}_3) \\ &+ \frac{e^{-\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-\pi\omega_3/a}}{1 - e^{-2\pi\omega_3/a}} \tilde{\delta}(\vec{\kappa}_1 - \vec{\kappa}_2) \tilde{\delta}(\vec{\kappa}_3 - \vec{\kappa}_4), \end{aligned} \quad (56e)$$

$$\begin{aligned} \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{L\vec{\kappa}_2}^* \hat{a}_{L\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle &= \frac{e^{-\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-\pi\omega_2/a}}{1 - e^{-2\pi\omega_2/a}} \tilde{\delta}(\vec{\kappa}_1 - \vec{\kappa}_3) \tilde{\delta}(\vec{\kappa}_2 - \vec{\kappa}_4) \\ &+ \frac{1}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-2\pi\omega_2/a}}{1 - e^{-2\pi\omega_2/a}} \delta(\vec{\kappa}_1 - \vec{\kappa}_4) \delta(\vec{\kappa}_2 - \vec{\kappa}_3) \end{aligned} \quad (56f)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{L\vec{\kappa}_2} \hat{a}_{L\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4}^* \Omega_M \rangle = 0, \quad (56g)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{L\vec{\kappa}_2}^* \hat{a}_{L\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (56h)$$

$$\begin{aligned} \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{L\vec{\kappa}_2} \hat{a}_{L\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle &= \frac{e^{-2\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{1}{1 - e^{-2\pi\omega_2/a}} \delta(\vec{\kappa}_1 - \vec{\kappa}_4) \delta(\vec{\kappa}_2 - \vec{\kappa}_3) \\ &+ \frac{e^{-\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-\pi\omega_2/a}}{1 - e^{-2\pi\omega_2/a}} \tilde{\delta}(\vec{\kappa}_1 - \vec{\kappa}_3) \tilde{\delta}(\vec{\kappa}_2 - \vec{\kappa}_4) \end{aligned} \quad (56i)$$

$$\begin{aligned} \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{L\vec{\kappa}_2}^* \hat{a}_{L\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle &= \frac{e^{-2\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-2\pi\omega_2/a}}{1 - e^{-2\pi\omega_2/a}} \delta(\vec{\kappa}_1 - \vec{\kappa}_4) \delta(\vec{\kappa}_2 - \vec{\kappa}_3) \\ &+ \frac{e^{-\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-\pi\omega_3/a}}{1 - e^{-2\pi\omega_3/a}} \tilde{\delta}(\vec{\kappa}_1 - \vec{\kappa}_2) \tilde{\delta}(\vec{\kappa}_3 - \vec{\kappa}_4), \end{aligned} \quad (56j)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1} \hat{a}_{L\vec{\kappa}_2}^* \hat{a}_{L\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (56k)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{L\vec{\kappa}_2} \hat{a}_{L\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (56l)$$

$$\langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{L\vec{\kappa}_2}^* \hat{a}_{L\vec{\kappa}_3} \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle = 0, \quad (56m)$$

$$\begin{aligned} \langle \Omega_M | \hat{a}_{R\vec{\kappa}_1}^* \hat{a}_{L\vec{\kappa}_2}^* \hat{a}_{L\vec{\kappa}_3}^* \hat{a}_{R\vec{\kappa}_4} \Omega_M \rangle &= \frac{e^{-\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-\pi\omega_2/a}}{1 - e^{-2\pi\omega_2/a}} \tilde{\delta}(\vec{\kappa}_1 - \vec{\kappa}_3) \tilde{\delta}(\vec{\kappa}_2 - \vec{\kappa}_4) \\ &+ \frac{e^{-\pi\omega_1/a}}{1 - e^{-2\pi\omega_1/a}} \frac{e^{-\pi\omega_3/a}}{1 - e^{-2\pi\omega_3/a}} \tilde{\delta}(\vec{\kappa}_1 - \vec{\kappa}_2) \tilde{\delta}(\vec{\kappa}_3 - \vec{\kappa}_4). \end{aligned} \quad (56n)$$

Following steps analogous to those in appendix (C) we obtain eq. (19).

References

- [1] P. C. W. Davies, “Scalar particle production in Schwarzschild and Rindler metrics”, *J. Phys. A* **8** (1975), 609-616 doi:10.1088/0305-4470/8/4/022
- [2] S. A. Fulling, “Nonuniqueness of canonical field quantization in Riemannian space-time”, *Phys. Rev. D* **7** (1973), 2850-2862. doi:10.1103/PhysRevD.7.2850

- [3] W. G. Unruh, “Notes on black hole evaporation”, *Phys. Rev. D* **14** (1976), 870 doi:10.1103/PhysRevD.14.870
- [4] S. De Bievre and M. Merkli, “The Unruh effect revisited”, *Class. Quant. Grav.* **23** (2006), 6525-6542 doi:10.1088/0264-9381/23/22/026 [arXiv:math-ph/0604023 [math-ph]].
- [5] C. J. Fewster, B. A. Juárez-Aubry and J. Louko, “Waiting for Unruh”, *Class. Quant. Grav.* **33** (2016) no.16, 165003 doi:10.1088/0264-9381/33/16/165003 [arXiv:1605.01316 [gr-qc]].
- [6] G. Salton, R. B. Mann and N. C. Menicucci, “Acceleration-assisted entanglement harvesting and rangefinding”, *New J. Phys.* **17** (2015) no.3, 035001 doi:10.1088/1367-2630/17/3/035001 [arXiv:1408.1395 [quant-ph]].
- [7] R. M. Wald, *Quantum Field Theory in Curved Space-Time and Black Hole Thermodynamics* (University of Chicago Press, 1994).
- [8] B. A. Juárez-Aubry and J. Louko, “Onset and decay of the 1 + 1 Hawking-Unruh effect: what the derivative-coupling detector saw”, *Class. Quant. Grav.* **31** (2014) no.24, 245007 doi:10.1088/0264-9381/31/24/245007 [arXiv:1406.2574 [gr-qc]].
- [9] L. C. B. Crispino, A. Higuchi and G. E. A. Matsas, “The Unruh effect and its applications”, *Rev. Mod. Phys.* **80** (2008), 787-838 doi:10.1103/RevModPhys.80.787 [arXiv:0710.5373 [gr-qc]].
- [10] B. L. Hu, S. Y. Lin and J. Louko, “Relativistic Quantum Information in Detectors-Field Interactions”, *Class. Quant. Grav.* **29** (2012), 224005 doi:10.1088/0264-9381/29/22/224005 [arXiv:1205.1328 [quant-ph]].
- [11] C. Gooding, S. Biermann, S. Erne, J. Louko, W. G. Unruh, J. Schmiedmayer and S. Weinfurtner, “Interferometric Unruh detectors for Bose-Einstein condensates”, *Phys. Rev. Lett.* **125** (2020) no.21, 213603 doi:10.1103/PhysRevLett.125.213603 [arXiv:2007.07160 [gr-qc]].
- [12] S. Biermann, S. Erne, C. Gooding, J. Louko, J. Schmiedmayer, W. G. Unruh and S. Weinfurtner, “Unruh and analogue Unruh temperatures for circular motion in 3+1 and 2+1 dimensions”, *Phys. Rev. D* **102** (2020) no.8, 085006 doi:10.1103/PhysRevD.102.085006 [arXiv:2007.09523 [gr-qc]].
- [13] B. A. Juárez-Aubry and D. Moustos, “Asymptotic states for stationary Unruh-DeWitt detectors”, *Phys. Rev. D* **100** (2019) no.2, 025018 doi:10.1103/PhysRevD.100.025018 [arXiv:1905.13542 [gr-qc]].
- [14] M. Good, B. A. Juárez-Aubry, D. Moustos and M. Temirkhan, “Unruh-like effects: Effective temperatures along stationary worldlines”, *JHEP* **06** (2020), 059 doi:10.1007/JHEP06(2020)059 [arXiv:2004.08225 [gr-qc]].
- [15] W. G. Unruh and R. M. Wald, “What happens when an accelerating observer detects a Rindler particle”, *Phys. Rev. D* **29** (1984), 1047-1056 doi:10.1103/PhysRevD.29.1047
- [16] A. Higuchi, G. E. A. Matsas and D. Sudarsky, “Bremsstrahlung and Fulling-Davies-Unruh thermal bath”, *Phys. Rev. D* **46** (1992), 3450-3457 doi:10.1103/PhysRevD.46.3450
- [17] A. Higuchi, G. E. A. Matsas and D. Sudarsky, “Bremsstrahlung and zero energy Rindler photons”, *Phys. Rev. D* **45** (1992), R3308-R3311 doi:10.1103/PhysRevD.45.R3308

- [18] D. A. T. Vanzella and G. E. A. Matsas, “Decay of accelerated protons and the existence of the Fulling-Davies-Unruh effect”, *Phys. Rev. Lett.* **87** (2001), 151301 doi:10.1103/PhysRevLett.87.151301 [arXiv:gr-qc/0104030 [gr-qc]].
- [19] G. E. A. Matsas and D. A. T. Vanzella, “Decay of protons and neutrons induced by acceleration”, *Phys. Rev. D* **59** (1999), 094004 doi:10.1103/PhysRevD.59.094004 [arXiv:gr-qc/9901008 [gr-qc]].
- [20] J. Audretsch and R. Muller, “Radiation from a uniformly accelerated particle detector: Energy, particles, and the quantum measurement process”, *Phys. Rev. D* **49** (1994), 6566-6575 doi:10.1103/PhysRevD.49.6566
- [21] B. D’Espagnat, *Conceptual Foundations Of Quantum Mechanics* (CRC Press, 1999).
- [22] C. J. Fewster and R. Verch, “Quantum fields and local measurements”, *Commun. Math. Phys.* **378** (2020) no.2, 851-889 doi:10.1007/s00220-020-03800-6 [arXiv:1810.06512 [math-ph]].
- [23] B. S. DeWitt, “Quantum gravity: the new synthesis”, in *General Relativity: an Einstein centenary survey*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1979) 680.
- [24] J. Louko and A. Satz, “Transition rate of the Unruh-DeWitt detector in curved spacetime”, *Class. Quant. Grav.* **25** (2008), 055012 doi:10.1088/0264-9381/25/5/055012 [arXiv:0710.5671 [gr-qc]].
- [25] B. A. Juárez-Aubry, B. S. Kay and D. Sudarsky, “Generally covariant dynamical reduction models and the Hadamard condition”, *Phys. Rev. D* **97** (2018) no.2, 025010 doi:10.1103/PhysRevD.97.025010 [arXiv:1708.09371 [gr-qc]].
- [26] F. Olver, *Asymptotics and special functions* (CRC Press, 1997).
- [27] T. Maudlin, E. Okon and D. Sudarsky, “On the status of conservation laws in physics: Implications for semiclassical gravity”, *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* **69**, 67-81 (2020).
- [28] Bell, J. (1964). On the Einstein-Podolsky-Rosen paradox. *Physics*, 1:195–200. Bell, J. (1990). La nouvelle cuisine. In *Between Science and Technology*. Elsevier Science Publishers.
- [29] A. Aspect, J. Dalibard and G. Roger, “Experimental test of Bell’s inequalities using time varying analyzers”, *Phys. Rev. Lett.* **49** (1982), 1804-1807 doi:10.1103/PhysRevLett.49.1804
- [30] A. Aspect, P. Grangier and G. Roger, “Experimental Tests of Realistic Local Theories via Bell’s Theorem”, *Phys. Rev. Lett.* **47** (1981), 460-6443 doi:10.1103/PhysRevLett.47.460
- [31] M. Giustina *et al.*, “Significant-loophole-free test of Bell’s theorem with entangled photons” *Phys. Rev. Lett.* **115**, (2015) 250401
- [32] J. Handsteiner *et al.* “Cosmic Bell Test: Measurement Settings from Milky Way Stars”, *Phys. Rev. Lett.* **118** (2017) no.6, 060401 doi:10.1103/PhysRevLett.118.060401 [arXiv:1611.06985 [quant-ph]].
- [33] B. Hensen *et al.*. “Loophole-free Bell inequality violation using electron spins separated by 1.3 kilometres”, *Nature*, **526** (2015) 575:682–686.

- [34] D. Greenberger; M. Horne; A. Shimony; A. Zeilinger (1990). "Bell's theorem without inequalities". *Am. J. Phys.* **58** (12): 1131.
- [35] T. Maudlin *Quantum nonlocality and reality* (Blackwell Publishing, 1994).
- [36] D. Carney, S. Ghosh, G. Krnjaic and J. M. Taylor, "Proposal for gravitational direct detection of dark matter", *Phys. Rev. D* **102** (2020) no.7, 072003 doi:10.1103/PhysRevD.102.072003 [arXiv:1903.00492 [hep-ph]].
- [37] J. Audretsch and R. Muller, "Radiation from a uniformly accelerated particle detector: Energy, particles, and the quantum measurement process", *Phys. Rev. D* **49** (1994), 6566-6575 doi:10.1103/PhysRevD.49.6566
- [38] L. H. Ford, Spacetime in Semiclassical Gravity, in *100 Years of Relativity - Space-time Structure: Einstein and Beyond*, edited by A. Ashtekar, (World Scientific, Singapore 2005) [arXiv:gr-qc/0504096]
- [39] P. Pearle, "Towards explaining why events occur", *Int. J. Theor. Phys* **18** (1979) 489
- [40] G. Ghirardi, A. Rimini, T. Weber, "A model for a unified quantum description of macroscopic and microscopic systems," in *Quantum Probability and Applications* edited by A. L. Accardi, (Springer, 1985) pp. 223
- [41] G. C. Ghirardi, A. Rimini and T. Weber, "A Unified Dynamics for microscopic and macroscopic Systems", *Phys. Rev. D* **34** (1986), 470 doi:10.1103/PhysRevD.34.470
- [42] P. M. Pearle, "Combining Stochastic Dynamical State Vector Reduction With Spontaneous Localization", *Phys. Rev. A* **39** (1989), 2277-2289 doi:10.1103/PhysRevA.39.2277
- [43] G. C. Ghirardi, P. M. Pearle and A. Rimini, "Markov Processes in Hilbert Space and Continuous Spontaneous Localization of Systems of Identical Particles", *Phys. Rev. A* **42** (1990), 78-79 doi:10.1103/PhysRevA.42.78
- [44] L. Diosi, "Gravitation and Quantum Mechanical Localization of Macro-Objects", *Phys. Lett. A* **105** (1984) 199-202
- [45] R. Penrose, "On Gravity's Role in Quantum State Reduction", *Gen. Rel. Gravit.* **28** (1996) 581-600
- [46] A. Bassi and G. C. Ghirardi, "Dynamical reduction models", *Phys. Rept.* **379** (2003), 257 doi:10.1016/S0370-1573(03)00103-0 [arXiv:quant-ph/0302164 [quant-ph]].
- [47] D. J. Bedingham, "Relativistic state reduction dynamics", *Found. Phys.* **41** (2011), 686-704 doi:10.1007/s10701-010-9510-7 [arXiv:1003.2774 [quant-ph]].
- [48] D. J. Bedingham, "Relativistic state reduction model", *J. Phys.: Conf. Ser.* **306** (2011) 012034
- [49] R. Tumulka, "A relativistic version of the Ghirardi-Rimini-Weber model", *J. Stat. Phys.* **125**, (2006) 821
- [50] P. Pearle, "Relativistic dynamical collapse model", *Phys. Rev. D* **91** (2015) 105012 [arXiv:1412.6723 [quant-ph]].

- [51] Y. Aharonov and D. Z. Albert, “Is the usual notion of time evolution adequate for quantum-mechanical systems? II. Relativistic considerations”, *Phys. Rev. D* **29** (1984) 228
- [52] D. Z. Albert, “Physics and Narrative” in *After Physics* (Harvard University Press, 2015)
- [53] W. C. Myrvold, “On peaceful coexistence: is the collapse postulate incompatible with relativity?” *Stud. Hist. Philos. Mod. Phys.* **33** (2002) 435
- [54] D. Z. Albert and R. Galchen, “A quantum threat to special relativity” *Sci.Am.* **300** (2009) 32
- [55] W. C. Myrvold, “Quantum Mechanics and Narratability” *Found Phys.* **46** (2016) 759.
- [56] Y. Bonder, C. Chryssomalakos and D. Sudarsky, “Extracting Geometry from Quantum Spacetime: Obstacles down the road”, *Foundations of Physics* **1** (2018); arXiv:1706.08221 [gr-qc].
- [57] R. D. Sorkin, in *Directions in General Relativity: Volume 2: Proceedings of the 1993 International Symposium, Maryland: Papers in Honor of Dieter Brill, Vol. 1* (Cambridge University Press, 1993) p. 293.
- [58] H. Bostelmann, C. J. Fewster and M. H. Ruep, “Impossible measurements require impossible apparatus”, *Phys. Rev. D* **103** (2021) no.2, 025017 doi:10.1103/PhysRevD.103.025017 [arXiv:2003.04660 [quant-ph]].
- [59] D. J. Bedingham, “Relativistic state reduction dynamics”, *Foundations of Physics* **41** (2011) 686.
- [60] D. J. Bedingham, D. Dürr, G. C. Ghirardi, S. Goldstein, R. Tumulka and N. Zanghì, “Matter density and relativistic models of wave function collapse”, *Journal of Statistical Physics* **154** (2014) 623–631.
- [61] R. Tumulka, “A relativistic version of the Ghirardi-Rimini-Weber model”, *Journal of Statistical Physics* **125** (2006) 821.
- [62] T. Maudlin, “Three measurement problems”, *Topoi* **14** (1995).
- [63] D. Bohm, “A suggested interpretation of quantum theory in terms of “hidden” variables”, *Physics Reviews* **85** (1952) 166–193.
- [64] G. C. Ghirardi, A. Rimini and T. Weber, “Unified dynamics for microscopic and macroscopic systems”, *Phys. Rev. D* **34** (1986) 470–491.
- [65] P. Pearle, “Combining stochastic dynamical state vector reduction with spontaneous localization”, *Phys. Rev. A* **39** (1989) 2277–2289.
- [66] R. Penrose, *The emperor’s new mind: Concerning computers, minds, and the laws of physics* (Oxford University Press, 1999).
- [67] A. Bassi, K. Lochan, S. Satin, T. Singh and H. Ulbricht, “Models of wave-function collapse, underlying theories, and experimental tests”, *Reviews of Modern Physics* **85** (2013) 471.
- [68] H. Everett, “Relative state formulation of quantum mechanics”, *Reviews of Modern Physics*, **29** (1957) 3.

- [69] D. Wallace, “The emergent multiverse: Quantum theory according to the everett interpretation” (Oxford University Press, 2014).
- [70] A. Belenchia, R. M. Wald, F. Giacomini, E. Castro-Ruiz, Č. Brukner and M. Aspelmeyer, “Information Content of the Gravitational Field of a Quantum Superposition”, *Int. J. Mod. Phys. D* **28** (2019) no.14, 1943001 doi:10.1142/S0218271819430016 [arXiv:1905.04496 [quant-ph]].
- [71] R. M. Wald, “Quantum superposition of massive bodies”, *Int. J. Mod. Phys. D* **29** (2020) no.11, 2041003 doi:10.1142/S0218271820410035
- [72] Daniel Carney, Sohritri Ghosh, Gordan Krnjaic, Jacob M. Taylor, “Proposal for gravitational direct detection of dark matter” *Phys. Rev. D* **102**, 072003 (2020)