

# Quantum baryon number fluctuations in subsystems of a hot and dense relativistic gas of fermions

Arpan Das<sup>1</sup>, Wojciech Florkowski<sup>2</sup>, Radoslaw Ryblewski<sup>1</sup>, and Rajeev Singh<sup>1</sup>

<sup>1</sup>*Institute of Nuclear Physics Polish Academy of Sciences, PL-31-342 Kraków, Poland and*

<sup>2</sup>*Institute of Theoretical Physics, Jagiellonian University, PL-30-348 Kraków, Poland*

(Dated: August 27, 2021)

Quantum features of the baryon number fluctuations in subsystems of a hot and dense relativistic gas of fermions are analyzed. We find that the fluctuations in small systems are significantly increased compared to their values known from the statistical physics, and diverge in the limit where the system size goes to zero. The numerical results obtained for a broad range of the thermodynamic parameters expected in heavy-ion collisions are presented. They can be helpful to interpret and shed new light on the experimental data.

## I. INTRODUCTION

Statistical fluctuations in many-body systems (microscopic as well as macroscopic ones) play a very important role in physics as they encode the crucial information about possible phase transitions, dissipation, and clustering phenomena [1–6]. An unexplored novel feature of the fluctuations is their increase for small systems in the cases where the quantum effects become important. Such effects have been analyzed quantitatively in our two recent papers [7, 8] where we addressed the fluctuations of the energy density in a hot gas of bosons and fermions. Our results indicate limitations of the concept of the fluid elements used in relativistic hydrodynamics applied in the description of heavy-ion collisions. As the size of a subsystem drops below about 0.5 fm, the energy density fluctuations (for typical values of temperatures and particle masses) become so large that they are comparable with their mean values. In this case, the physical picture of fluid cells with well defined energy density becomes unjustified. An interesting feature of our calculations [7, 8] is that not only fluctuations diverge for the subsystem with size approaching zero, but also quantum statistical fluctuations agree with thermodynamic fluctuations if the subsystems become sufficiently large [7, 8]. In this way, our approach incorporates quantum features into statistical mechanics in a very natural way.

In the present work, following similar ideas to that developed in [7, 8], we discuss fluctuations of the baryon number density in a hot and dense relativistic gas of fermions. Our analysis is relevant for relativistic heavy-ion physics, in particular, in the context of the beam energy scan (BES). Hunt for the conjectured critical endpoint in the QCD phase diagram has triggered vast theoretical and experimental studies of many fluctuation observables. For example, one studies fluctuations of the conserved quantum numbers in QCD, such as the baryon number, electric charge, or strangeness. They all provide an excellent opportunity to study the critical phenomena [9–16].

Following our previous framework [7, 8], we consider the fluctuation of the baryon number in the subsystem  $S_a$  of the thermodynamic system  $S_V$  described by the

grand canonical ensemble characterized by the temperature ( $T$ ) (or its inverse  $\beta = 1/T$ ) and the baryon chemical potential ( $\mu$ ). The volume  $V$  of the larger system  $S_V$  is larger than the characteristic volume of the subsystem  $S_a$ . We derive a compact formula that defines quantum fluctuations of the baryon number operator in subsystems of a hot and dense relativistic gas. Then we apply this formula to get physical insights into situations expected in relativistic heavy-ion collisions. In the calculation we use the metric tensor with the signature  $(+1, -1, -1, -1)$ . Three-vectors are shown in bold font and a dot is used to denote the scalar product of both four and three-vectors, i.e.,  $a^\mu b_\mu = a \cdot b = a^0 b^0 - \mathbf{a} \cdot \mathbf{b}$ .

The structure of the paper is organized as follows: We start with introducing the basic concepts and definitions in Sec. II, where we also derive the expression for baryon number density fluctuation. Subsequently, in Sec. III we discuss the thermodynamic limit of baryon number density fluctuation. The numerical results of the baryon number density fluctuations are presented in Sec. IV. Finally, in Sec. V we interpret and summarize our results.

## II. BASIC CONCEPTS AND DEFINITIONS

We consider a system of spin-1/2 particles described by the quantum Dirac field in thermal equilibrium. The field operator is expressed in the standard way as [17]

$$\psi(t, \mathbf{x}) = \sum_r \int \frac{dK}{\sqrt{2\omega_{\mathbf{k}}}} \left( U_r(\mathbf{k}) a_r(\mathbf{k}) e^{-ik \cdot x} + V_r(\mathbf{k}) b_r^\dagger(\mathbf{k}) e^{ik \cdot x} \right), \quad (1)$$

where we use the notation  $dK \equiv d^3k/(2\pi)^3$ , while  $a_r(\mathbf{k})$  and  $b_r^\dagger(\mathbf{k})$  are the annihilation and creation operators for particles and antiparticles, respectively. The polarization degree of freedom is represented by the index  $r$ . The fermionic operators  $a_r(\mathbf{k})$  and  $b_r^\dagger(\mathbf{k})$  satisfy the canonical anticommutation relations,  $\{a_r(\mathbf{k}), a_s^\dagger(\mathbf{k}')\} = (2\pi)^3 \delta_{rs} \delta^{(3)}(\mathbf{k} - \mathbf{k}')$  and  $\{b_r(\mathbf{k}), b_s^\dagger(\mathbf{k}')\} = (2\pi)^3 \delta_{rs} \delta^{(3)}(\mathbf{k} - \mathbf{k}')$ . All the other operators anticommute with each other. The Dirac spinors  $U_r(\mathbf{k})$  and  $V_r(\mathbf{k})$  have normalization  $\bar{U}_r(\mathbf{k}) U_s(\mathbf{k}) =$

$2m\delta_{rs}$  and  $\bar{V}_r(\mathbf{k})V_s(\mathbf{k}) = -2m\delta_{rs}$ , and the quantity  $\omega_{\mathbf{k}} = \sqrt{\mathbf{k}^2 + m^2}$  is the energy of a particle.

To perform thermal averaging of quantum operators, it is sufficient to know the thermal expectation values of the products of two and four creation and/or annihilation operators (for both particles and antiparticles) [18–20]

$$\begin{aligned} \langle a_r^\dagger(\mathbf{k})a_s(\mathbf{k}') \rangle &= (2\pi)^3 \delta_{rs} \delta^{(3)}(\mathbf{k} - \mathbf{k}') f(\omega_{\mathbf{k}}), \quad (2) \\ \langle a_r^\dagger(\mathbf{k})a_s^\dagger(\mathbf{k}')a_{r'}(\mathbf{p})a_{s'}(\mathbf{p}') \rangle \\ &= (2\pi)^6 \left( \delta_{rs'} \delta_{r's} \delta^{(3)}(\mathbf{k} - \mathbf{p}') \delta^{(3)}(\mathbf{k}' - \mathbf{p}) \right. \\ &\quad \left. - \delta_{rr'} \delta_{ss'} \delta^{(3)}(\mathbf{k} - \mathbf{p}) \delta^{(3)}(\mathbf{k}' - \mathbf{p}') \right) f(\omega_{\mathbf{k}}) f(\omega_{\mathbf{k}'}). \quad (3) \end{aligned}$$

Here  $f(\omega_{\mathbf{k}}) = 1/(\exp(\beta(\omega_{\mathbf{k}} - \mu)) + 1)$  is the Fermi–Dirac distribution function for particles. For antiparticles, the Fermi–Dirac distribution function differs by the sign of the baryon chemical potential  $\mu$ , i.e.  $\bar{f}(\omega_{\mathbf{k}}) = 1/(\exp(\beta(\omega_{\mathbf{k}} + \mu)) + 1)$ . Using the anticommutation relations for  $a_r(\mathbf{k})$ ,  $a_r^\dagger(\mathbf{k})$ ,  $b_r(\mathbf{k})$ , and  $b_r^\dagger(\mathbf{k})$  any other combination of two and four creation and/or annihilation operators can be obtained from Eqs. (2) and (3).

Following Ref. [21], we define the baryon number density operator  $\hat{J}_a^0$ , associated with the conserved baryon current in a subsystem  $S_a$  using a smooth Gaussian profile placed at the origin of the coordinate system, namely

$$\hat{J}_a^0 = \frac{1}{(a\sqrt{\pi})^3} \int d^3\mathbf{x} \hat{J}^0(x) \exp\left(-\frac{\mathbf{x}^2}{a^2}\right), \quad (4)$$

where  $\hat{J}^0 = \psi^\dagger\psi$ . To remove any possible sharp-boundary effects we consider the smooth profile with a length scale  $a$  instead of a cubic box. The thermal expectation value of the normal ordered operator  $: \hat{J}_a^0 :$  is denoted as  $\langle : \hat{J}_a^0 : \rangle$ . Here standard normal ordering procedure has been introduced to remove an infinite vacuum part coming from zero-point contributions. To determine the baryon number fluctuation of the subsystem  $S_a$ , we consider the variance

$$\sigma^2(a, m, T, \mu) = \langle : \hat{J}_a^0 :: \hat{J}_a^0 : \rangle - \langle : \hat{J}_a^0 : \rangle^2 \quad (5)$$

and the normalized standard deviation as

$$\sigma_n(a, m, T, \mu) = \frac{(\langle : \hat{J}_a^0 :: \hat{J}_a^0 : \rangle - \langle : \hat{J}_a^0 : \rangle^2)^{1/2}}{\langle : \hat{J}_a^0 : \rangle}. \quad (6)$$

Using the Fourier mode expansion of the Dirac field operator as given by Eq. (1), the normal ordered operator

$: \hat{J}_a^0 :$  can be obtained as,

$$\begin{aligned} : \hat{J}_a^0 : &:= \sum_{r,s} \int \frac{dK}{\sqrt{2\omega_{\mathbf{k}}}} \frac{dK'}{\sqrt{2\omega_{\mathbf{k}'}}} \times \\ &\left[ a_s^\dagger(\mathbf{k}') a_r(\mathbf{k}) \bar{U}_s(\mathbf{k}') \gamma^0 U_r(\mathbf{k}) e^{-i(\omega_{\mathbf{k}} - \omega_{\mathbf{k}'})t} e^{-\frac{a^2}{4}(\mathbf{k} - \mathbf{k}')^2} \right. \\ &\quad - b_r^\dagger(\mathbf{k}) b_s(\mathbf{k}') \bar{V}_s(\mathbf{k}') \gamma^0 V_r(\mathbf{k}) e^{i(\omega_{\mathbf{k}} - \omega_{\mathbf{k}'})t} e^{-\frac{a^2}{4}(\mathbf{k} - \mathbf{k}')^2} \\ &\quad + a_s^\dagger(\mathbf{k}') b_r^\dagger(\mathbf{k}) \bar{U}_s(\mathbf{k}') \gamma^0 V_r(\mathbf{k}) e^{i(\omega_{\mathbf{k}} + \omega_{\mathbf{k}'})t} e^{-\frac{a^2}{4}(\mathbf{k} + \mathbf{k}')^2} \\ &\quad \left. + b_s(\mathbf{k}') a_r(\mathbf{k}) \bar{V}_s(\mathbf{k}') \gamma^0 U_r(\mathbf{k}) e^{-i(\omega_{\mathbf{k}} + \omega_{\mathbf{k}'})t} e^{-\frac{a^2}{4}(\mathbf{k} + \mathbf{k}')^2} \right]. \quad (7) \end{aligned}$$

Using the thermal averaging of two creation and/or annihilation operators as given by Eq. (2), the thermal expectation value of  $: \hat{J}_a^0 :$  has the form

$$\langle : \hat{J}_a^0 : \rangle = 2 \int dK [f(\omega_{\mathbf{k}}) - \bar{f}(\omega_{\mathbf{k}})]. \quad (8)$$

This expression agrees with the standard kinetic-theory definition, with the factor of 2 accounting for the spin degeneracy.

In the next step, using the thermal averaging of four creation and/or annihilation operators as given by Eq. (3) we obtain

$$\begin{aligned} \sigma^2(a, m, T, \mu) &= \langle : \hat{J}_a^0 :: \hat{J}_a^0 : \rangle - \langle : \hat{J}_a^0 : \rangle^2 \\ &= \int \frac{dK}{\omega_{\mathbf{k}}} \frac{dK'}{\omega_{\mathbf{k}'}} (\omega_{\mathbf{k}} \omega_{\mathbf{k}'} + \mathbf{k} \cdot \mathbf{k}' + m^2) e^{-\frac{a^2}{2}(\mathbf{k} - \mathbf{k}')^2} \times \\ &\quad [f(\omega_{\mathbf{k}})(1 - f(\omega_{\mathbf{k}'})) + \bar{f}(\omega_{\mathbf{k}})(1 - \bar{f}(\omega_{\mathbf{k}'}))] \\ &\quad - \int \frac{dK}{\omega_{\mathbf{k}}} \frac{dK'}{\omega_{\mathbf{k}'}} (\omega_{\mathbf{k}} \omega_{\mathbf{k}'} + \mathbf{k} \cdot \mathbf{k}' - m^2) e^{-\frac{a^2}{2}(\mathbf{k} + \mathbf{k}')^2} \times \\ &\quad [f(\omega_{\mathbf{k}})(1 - \bar{f}(\omega_{\mathbf{k}'})) + \bar{f}(\omega_{\mathbf{k}})(1 - f(\omega_{\mathbf{k}'}))]. \quad (9) \end{aligned}$$

Although normal ordering removes unwanted vacuum contribution to the baryon number density operator, it is not sufficient to remove all the vacuum divergences in all composite operators. Therefore following Refs. [7, 8] we discard a divergent temperature and chemical potential independent vacuum term that originally appears in Eq. (9). Equation (9) is the main result which represents a fluctuation of the baryon number density in the subsystem  $S_a$ . Note that in Eqs. (8) and (9), the spin and particle-antiparticle degrees of freedom are included. If we take into account the degeneracy factors ( $g$ ) due to other internal degrees of freedom then we should do the following replacements:  $\langle : \hat{J}_a^0 : \rangle \rightarrow g \langle : \hat{J}_a^0 : \rangle$  and  $\sigma^2 \rightarrow g\sigma^2$  [7, 8].

### III. THERMODYNAMIC LIMIT

Before we turn to a discussion of our numerical results based on Eq. (9), it is important to analyze the thermodynamic limit of  $\sigma^2(a, m, T, \mu)$ . Since  $S_a$  is a subsystem of the system  $S_V$ , the thermodynamic limit can




FIG. 1 – Variation of normalized fluctuation  $\sigma_n$  in the subsystem  $S_a$  with the scale  $a$  for different values of the temperature  $T$  and fixed particle mass  $m = 1.0$  GeV and baryon chemical potential  $\mu = 0.5$  GeV.




FIG. 2 – Variation of normalized fluctuation  $\sigma_n$  in the subsystem  $S_a$  with the scale  $a$  for different values of the baryon chemical potential  $\mu$  and fixed particle mass  $m = 1.0$  GeV and temperature  $T = 0.15$  GeV.

be achieved by considering the  $a \rightarrow \infty$  limit (still with  $a^3 \ll V$ ). In the thermodynamic limit quantum fluctuation as obtained here should reduce to the expression for the classical statistical fluctuation. To obtain the classical limit of the fluctuations we look into the susceptibilities describing fluctuations in the baryon number obtained for thermal and chemical equilibrium [22]

$$\chi_l^{(B)} = \frac{\partial^l (P/T^4)}{\partial (\mu/T)^l} \Big|_T. \quad (10)$$




FIG. 3 – Variation of normalized fluctuation  $\sigma_n$  in the subsystem  $S_a$  with the scale  $a$  for different values of the particle mass and fixed temperature  $T = 0.15$  GeV and baryon chemical potential  $\mu = 0.5$  GeV.

Here  $P$  denotes the thermodynamic pressure. Susceptibilities can also be related to the cumulants of the distribution of baryons, e.g.,

$$\begin{aligned} \chi_1^{(B)} &= \frac{1}{VT^3} \langle N_B \rangle = \frac{1}{T^3} \frac{\langle N_B \rangle}{V} = \frac{n_B}{T^3}, \\ \chi_2^{(B)} &= \frac{1}{VT^3} \langle (\Delta N_B)^2 \rangle = \frac{1}{VT^3} \langle (N_B - \langle N_B \rangle)^2 \rangle. \end{aligned} \quad (11)$$

Here  $n_B \equiv N_B/V$  is the net baryon number density. Equation (11) also implies that

$$V \langle (n_B - \langle n_B \rangle)^2 \rangle = T^3 \chi_2^{(B)}. \quad (12)$$

Using Eq. (10), the susceptibility  $\chi_2^{(B)}$  can also be obtained by taking the appropriate derivative of the thermodynamic pressure. The thermodynamic pressure at finite temperature and baryon chemical potential can be expressed as [22]

$$\begin{aligned} \frac{P}{T^4} &= \frac{2g}{T^3} \int dK \left[ \ln \left( 1 + \exp \left( -\frac{\omega_{\mathbf{k}} - \mu}{T} \right) \right) \right. \\ &\quad \left. + \ln \left( 1 + \exp \left( -\frac{\omega_{\mathbf{k}} + \mu}{T} \right) \right) \right]. \end{aligned} \quad (13)$$

Using Eqs. (10) and (13) we can easily find

$$\begin{aligned} \chi_1^{(B)} &= \frac{\partial (P/T^4)}{\partial (\mu/T)} \Big|_T \\ &= \frac{2g}{T^3} \int dK \left[ f(\omega_{\mathbf{k}}) - \bar{f}(\omega_{\mathbf{k}}) \right] \\ &= \frac{n_B}{T^3} \end{aligned} \quad (14)$$

and

$$\begin{aligned}\chi_2^{(B)} &= \left. \frac{\partial^2 (P/T^4)}{\partial (\mu/T)^2} \right|_T \\ &= \frac{2g}{T^3} \int dK \left[ f(\omega_{\mathbf{k}})(1 - f(\omega_{\mathbf{k}})) \right. \\ &\quad \left. + \bar{f}(\omega_{\mathbf{k}})(1 - \bar{f}(\omega_{\mathbf{k}})) \right].\end{aligned}\quad (15)$$

Therefore, we can write

$$\begin{aligned}T^3 \chi_2^{(B)} &= V \langle (n_B - \langle n_B \rangle)^2 \rangle, \\ &= 2g \int dK \left[ f(\omega_{\mathbf{k}})(1 - f(\omega_{\mathbf{k}})) \right. \\ &\quad \left. + \bar{f}(\omega_{\mathbf{k}})(1 - \bar{f}(\omega_{\mathbf{k}})) \right].\end{aligned}\quad (16)$$

In the large volume limit Eq. (9) should be consistent with Eq. (16). This can be verified using the Gaussian representation of the three-dimensional Dirac delta function

$$\delta^{(3)}(\mathbf{k} - \mathbf{k}') = \lim_{a \rightarrow \infty} \frac{a^3}{(2\pi)^{3/2}} e^{-\frac{a^2}{2}(\mathbf{k} - \mathbf{k}')^2}. \quad (17)$$

This leads us to the following expression

$$\begin{aligned}&\lim_{a \rightarrow \infty} a^3 (2\pi)^{3/2} \left[ \langle : \hat{J}_a^0 :: \hat{J}_a^0 : \rangle - \langle : \hat{J}_a^0 : \rangle^2 \right] \\ &= \int \frac{d^3 k}{(2\pi)^3} d^3 k' \frac{1}{\omega_{\mathbf{k}}} \frac{1}{\omega_{\mathbf{k}'}} (\omega_{\mathbf{k}} \omega_{\mathbf{k}'} + \mathbf{k} \cdot \mathbf{k}' + m^2) \\ &\quad [f(\omega_{\mathbf{k}})(1 - f(\omega_{\mathbf{k}'}) + \bar{f}(\omega_{\mathbf{k}})(1 - \bar{f}(\omega_{\mathbf{k}'}))] \delta^{(3)}(\mathbf{k} - \mathbf{k}') \\ &- \int \frac{d^3 k}{(2\pi)^3} d^3 k' \frac{1}{\omega_{\mathbf{k}}} \frac{1}{\omega_{\mathbf{k}'}} (\omega_{\mathbf{k}} \omega_{\mathbf{k}'} + \mathbf{k} \cdot \mathbf{k}' - m^2) \\ &\quad [f(\omega_{\mathbf{k}})(1 - \bar{f}(\omega_{\mathbf{k}'}) + \bar{f}(\omega_{\mathbf{k}})(1 - f(\omega_{\mathbf{k}'}))] \delta^{(3)}(\mathbf{k} + \mathbf{k}') \\ &= 2 \int dK \left[ f(\omega_{\mathbf{k}})(1 - f(\omega_{\mathbf{k}})) + \bar{f}(\omega_{\mathbf{k}})(1 - \bar{f}(\omega_{\mathbf{k}})) \right],\end{aligned}\quad (18)$$

hence,

$$\lim_{a \rightarrow \infty} V_a \left[ \langle : \hat{J}_a^0 :: \hat{J}_a^0 : \rangle - \langle : \hat{J}_a^0 : \rangle^2 \right] = T^3 \chi_2^{(B)}, \quad (19)$$

where  $V_a = a^3 (2\pi)^{3/2}$  may be identified as the volume of the ‘‘Gaussian’’ subsystem  $S_a$ .

#### IV. NUMERICAL RESULTS

Our main result describing fluctuations of the baryon number density in a hot and dense gas of baryons is represented by Eq. (9). By straightforward numerical integration, we can obtain the results for any subsystem of size  $a$ , temperature  $T$ , baryon chemical potential  $\mu$ , and particle mass  $m$ . In Figs. (1), (2) and (3) we show the variation of the normalized fluctuation  $\sigma_n$  as defined




FIG. 4 – Variation of normalized fluctuation  $V_a \sigma^2 / (T^3 \chi_2^{(B)})$  for different values of temperature ( $T$ ) but with fixed baryon chemical potential ( $\mu$ ) and particle mass ( $m$ ).




FIG. 5 – Variation of normalized fluctuation  $V_a \sigma^2 / (T^3 \chi_2^{(B)})$  for different values of baryon chemical potential ( $\mu$ ) but with fixed temperature ( $T$ ) and particle mass ( $m$ ).

in Eq. (6) with the subsystem size  $a$  for different values of temperature, baryon chemical potential and mass, respectively. Keeping in mind the relativistic heavy ion collisions in the regime where large baryon number densities are expected, we consider the following range of the parameters:  $0.1 \text{ GeV} \leq T \leq 0.25 \text{ GeV}$ ,  $0.5 \text{ GeV} \leq \mu \leq 1.0 \text{ GeV}$  and  $0.0 \leq m \leq 1.0 \text{ GeV}$ . The internal degeneracy factor is taken to be  $g = 10$ .

Figs. (1), (2) and (3) show that the normalized fluctuation  $\sigma_n$  decreases with the increase in the size of the subsystem  $S_a$ , which is the expected behavior of fluctuations. Furthermore, for small system size, the fluctuation




FIG. 6 – Variation of normalized fluctuation  $V_a \sigma^2 / (T^3 \chi_2^{(B)})$  for different values of particle mass ( $m$ ) but with fixed temperature ( $T$ ) and baryon chemical potential ( $\mu$ ).

is substantially large, which is the manifestation of the quantum mechanical behavior. From these figures, it is clear that the normalized fluctuation  $\sigma_n$  decreases with an increase in temperature and baryon chemical potential, however,  $\sigma_n$  is large if the particle mass is large. We should mention that variation of  $\sigma$  with temperature, baryon chemical potential and mass is opposite to the variation of  $\sigma_n$  (not shown here).

In Figs. (4), (5) and (6) we demonstrate the thermodynamic limit of the volume scaled fluctuation  $V_a \sigma^2$ . From Eq. (19) we conclude that in the thermodynamic limit, i.e., in the  $a \rightarrow \infty$  limit,  $V_a \sigma^2 / (T^3 \chi_2^{(B)})$  should approach unity. This property can be seen clearly in Figs. (4), (5) and (6), where for  $a > 1$  fm quantum fluctuations approaches the classical limit which also depend upon the temperature ( $T$ ), baryon chemical potential ( $\mu$ ) and particle mass ( $m$ ). Furthermore, for small system size the quantum fluctuations can be significant.

## V. CONCLUSION

In this work, we have analyzed quantum baryon-number fluctuations in subsystems of a hot and dense relativistic gas of fermions and found that they diverge for small system sizes. On the other hand, our results agree with the results known from statistical physics for sufficiently large system size  $a$ . In this way, we have delivered a useful formula that accounts for both statistical and quantum features of the fluctuations. The numerical results have been obtained for a broad range of thermodynamic parameters. They may be useful to interpret and shed new light on the heavy-ion experimental data.

## Acknowledgments

This research was supported in part by the Polish National Science Centre Grants No. 2016/23/B/ST2/00717 and No. 2018/30/E/ST2/00432.

- 
- [1] M. Smoluchowski, “Beitrag zur Theorie der Opaleszenz von Gasen im kritischen Zustande,” *Bulletin internationale de l’Académie des sciences de Cracovie* (1911) 493–502.
- [2] S. Jeon and V. Koch, “Charged particle ratio fluctuation as a signal for quark-gluon plasma,” *Phys. Rev. Lett.* **85** (Sep, 2000) 2076–2079. [[Phys. Rev. Lett.](#) **85** (Sep, 2000) 2076–2079].
- [3] K. Huang, *Statistical Mechanics*. Wiley New York, 2nd ed., 1987.
- [4] R. Kubo, “The Fluctuation-Dissipation Theorem,” *Rep. Prog. Phys.* **29** (1966) 255.
- [5] E. Lifshitz and I. Khalatnikov, “Investigations in relativistic cosmology,” *Adv. Phys.* **12** (1963) 185–249.
- [6] A. H. Guth and S.-Y. Pi, “Fluctuations in the new inflationary universe,” *Phys. Rev. Lett.* **49** (Oct, 1982) 1110–1113.
- [7] A. Das, W. Florkowski, R. Ryblewski, and R. Singh, “Quantum fluctuations of energy in subsystems of a hot relativistic gas,” [arXiv:2012.05662 \[hep-ph\]](#).
- [8] A. Das, W. Florkowski, R. Ryblewski, and R. Singh, “Pseudo-gauge dependence of quantum fluctuations of energy in a hot relativistic gas of fermions,” [arXiv:2103.01013 \[nucl-th\]](#).
- [9] J. Berges and K. Rajagopal, “Color superconductivity and chiral symmetry restoration at nonzero baryon density and temperature,” *Nucl. Phys. B* **538** (1999) 215–232, [arXiv:hep-ph/9804233](#).
- [10] A. M. Halasz, A. D. Jackson, R. E. Shrock, M. A. Stephanov, and J. J. M. Verbaarschot, “On the phase diagram of QCD,” *Phys. Rev. D* **58** (1998) 096007, [arXiv:hep-ph/9804290](#).
- [11] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, “Signatures of the tricritical point in QCD,” *Phys. Rev. Lett.* **81** (1998) 4816–4819, [arXiv:hep-ph/9806219](#).
- [12] M. A. Stephanov, K. Rajagopal, and E. V. Shuryak, “Event-by-event fluctuations in heavy ion collisions and the QCD critical point,” *Phys. Rev. D* **60** (1999) 114028, [arXiv:hep-ph/9903292](#).
- [13] Y. Hatta and T. Ikeda, “Universality, the QCD critical / tricritical point and the quark number susceptibility,” *Phys. Rev. D* **67** (2003) 014028, [arXiv:hep-ph/0210284](#).
- [14] M. Stephanov, “Non-Gaussian fluctuations near the QCD critical point,” *Phys. Rev. Lett.* **102** (2009) 032301, [arXiv:0809.3450 \[hep-ph\]](#).
- [15] B. Berdnikov and K. Rajagopal, “Slowing

- out-of-equilibrium near the QCD critical point,” *Phys. Rev. D* **61** (2000) 105017, [arXiv:hep-ph/9912274](#).
- [16] M. Kitazawa, M. Asakawa, and H. Ono, “Non-equilibrium time evolution of higher order cumulants of conserved charges and event-by-event analysis,” *Phys. Lett. B* **728** (2014) 386–392, [arXiv:1307.2978 \[nucl-th\]](#).
- [17] L. Tinti and W. Florkowski, “Particle polarization, spin tensor and the Wigner distribution in relativistic systems,” [arXiv:2007.04029 \[nucl-th\]](#).
- [18] C. Cohen-Tannoudji, B. Diu, F. Laloë, and S. R. Hemley, *Quantum mechanics: Vol. 3: fermions, bosons, photons, correlations and entanglement*. A Wiley-Interscience publication. Wiley, New York, 1977.
- [19] C. Itzykson and J. Zuber, *Quantum Field Theory*. International Series In Pure and Applied Physics. McGraw-Hill, New York, 1980.
- [20] T. Evans and D. A. Steer, “Wick’s theorem at finite temperature,” *Nucl. Phys. B* **474** (1996) 481–496, [arXiv:hep-ph/9601268](#).
- [21] S. Coleman, *Lectures of Sidney Coleman on Quantum Field Theory*. WSP, Hackensack, 12, 2018.
- [22] M. Nahrgang, M. Bluhm, P. Alba, R. Bellwied, and C. Ratti, “Impact of resonance regeneration and decay on the net-proton fluctuations in a hadron resonance gas,” *Eur. Phys. J. C* **75** no. 12, (2015) 573, [arXiv:1402.1238 \[hep-ph\]](#).