

Higgs boson decay to paired B_c : relativistic and one-loop corrections.

I. N. Belov,^{1,2,*} A. V. Berezhnoy,^{1,†} A. E. Dorokhov,^{3,‡}
A. K. Likhoded,^{4,§} A. P. Martynenko,^{5,¶} and F. A. Martynenko^{5,**}

¹*SINP MSU, Moscow, Russia*

²*Physics department of MSU, Moscow, Russia*

³*Joint Institute of Nuclear Research, BLTP, Moscow region, Dubna, Russia*

⁴*NRC "Kurchatov Institute" IHEP, Protvino, Russia*

⁵*Samara University, Samara, Russia*

The exclusive decays of the Higgs boson to $B_c B_c$ and $B_c^* B_c^*$ pairs are studied. The hard parts of the decay amplitudes are estimated within the perturbative Standard Model up to one-loop corrections. The soft fusion of heavy quarks to the quarkonium is described in framework of the relativistic quark model.

1. INTRODUCTION

One of the main goals of the research programs of the CMS and ATLAS experiments at the LHC is to study the properties of the Higgs boson [1, 2]. The decays into two W bosons, two Z bosons and two photons play a key role in the Higgs boson investigation. Among other decays, the decays into two quarkonia, including the decays into the pair of pseudoscalar mesons $B_c B_c$ or vector mesons $B_c^* B_c^*$, may be of particular interest (note, that the Higgs boson decay process $H \rightarrow B_c^* + B_c$ is forbidden due to the conservation law of angular momentum). Such processes allow to study the coupling constants of the Higgs boson with heavy quarks, as well as to test the theory of bound states of heavy quarks

*Electronic address: ilia.belov@cern.ch

†Electronic address: Alexander.Berezhnoy@cern.ch

‡Electronic address: dorokhov@theor.jinr.ru

§Electronic address: Anatolii.Likhoded@ihep.ru

¶Electronic address: a.p.martynenko@samsu.ru

**Electronic address: f.a.martynenko@gmail.com

produced in decays. In this work, we calculate both the relativistic corrections connected with the relative motion of heavy quarks and the QCD one-loop corrections to the Higgs boson decay width. As is well known from the studies of various authors, starting with the production of J/ψ and η_c mesons, the corrections of both these types significantly contribute to the production cross section of a pair of heavy quarkonia.

A bound state ($\bar{b}c$) with open beauty and charm has a special place among the heavy quarkoniums since its decay mechanism differs significantly from the decay mechanism of charmonium or bottomonium. That is why we believe that the process $H \rightarrow B_c + B_c$ will attract the attention of experimenters.

Our approach to the calculation of the observed Higgs boson decay widths leading to the pair production of B_c mesons is based on the methods of relativistic quark model (RQM) and the perturbative Standard Model [3, 4]. This approach allows a systematic account for relativistic effects throughout the construction of relativistic amplitudes of pair production of mesons, relativistic production cross sections, and in the description of bound states of quarks themselves through the use of the corresponding quark interaction potential.

As it is known from the previous study [5], the paired B_c production is essentially affected by one loop QCD corrections. Thus we take them into account within the same method, as was applied in [5] using modern computer methods for calculating the Feynman interaction amplitudes.

One of the first works devoted to the pair production of quarkonia in Higgs boson decays was done in the nonrelativistic approximation in [6]. The production of single quarkonia in the H decay was investigated in [7, 8] with the account of relativistic corrections and one-loop corrections. In the work [9], various channels of the Higgs boson decay into pairs of heavy quarkonia were studied, including $H \rightarrow ZZ$ and $H \rightarrow WW$. The single B_c meson production rate in Higgs boson decays was calculated within the nonrelativistic QCD framework in [10]. The first experimental searches for decays of the Higgs boson into a pair of J/Ψ and Υ mesons were performed in [11].

While the quarkonia with a hidden flavour have been studied experimentally well enough, the experimental data on the bound states of heavy quark and antiquark with different flavours are rather poor [12]. In fact, such states, B_c mesons, are known for the $1S$ and $2S$ states only. Therefore, the study of various mechanisms for the production of B_c mesons is of obvious interest, which is connected with the study of their properties.

2. GENERAL RQM FORMALISM

Four production amplitudes of the B_c meson pair in leading order of the QCD coupling constant α_s are presented in Fig. 1. We investigate the production channel of a pair of B_c mesons connected with the initial production of a pair of heavy quarks b or c in the Higgs boson decay. There are two stages of B_c meson production process. At the first stage, which is described by the perturbative Standard Model, the Higgs boson transforms into a heavy quark-antiquark pair. Then the heavy quark or antiquark emits a virtual gluon which produces another heavy quark-antiquark pair. At the second stage, heavy quarks and antiquarks combine with some probability into bound states.

Four-momenta of heavy quarks and antiquarks can be expressed in terms of relative and total four momenta as follows:

$$p_1 = \eta_1 P + p, \quad p_2 = \eta_2 P - p, \quad (p \cdot P) = 0, \quad \eta_i = \frac{M_{B_c}^2 \pm m_1^2 \mp m_2^2}{2(M_{B_c})^2}, \quad (1)$$

$$q_1 = \rho_1 Q + q, \quad q_2 = \rho_2 Q - q, \quad (q \cdot Q) = 0, \quad \rho_i = \frac{M_{B_c}^2 \pm m_1^2 \mp m_2^2}{2(M_{B_c})^2},$$

where M_{B_c} is the mass of pseudoscalar or vector B_c^+ (B_c^{*+}) meson consisting of \bar{b} -antiquark and c -quark. $m_{1,2}$ are the masses of c and b quarks. $P(Q)$ are the total four-momenta of mesons B_c^+ and B_c^{*-} , relative quark four-momenta $p = L_P(0, \mathbf{p})$ and $q = L_P(0, \mathbf{q})$ are obtained from the rest frame four-momenta $(0, \mathbf{p})$ and $(0, \mathbf{q})$ by the Lorentz transformation to the system moving with the momenta P and Q . The index $i = 1, 2$ corresponds to plus and minus signs in (1). Heavy quarks c, b and antiquarks \bar{c}, \bar{b} in the intermediate state are outside the mass shell: $p_{1,2}^2 = \eta_i^2 P^2 - \mathbf{p}^2 = \eta_i^2 M_{B_c}^2 - \mathbf{p}^2 \neq m_{1,2}^2$, so that $p_1^2 - m_1^2 = p_2^2 - m_2^2$.

Let consider the production amplitude of pseudoscalar and vector B_c mesons. Initially it can be written as a convolution of perturbative production amplitude of free quarks and antiquarks and the quasipotential wave functions of B_c mesons moving with four-momenta P and Q . Using then the transformation law of the bound state wave functions from the rest frame to the moving one with four-momenta P and Q we can present the meson production amplitude in the form [4, 13, 14]:

$$\mathcal{M}(p_-, p_+, P, Q) = -iM(\sqrt{2}G_F)^{1/2} \frac{2\pi}{3} M_{B_c} \int \frac{d\mathbf{p}}{(2\pi)^3} \int \frac{d\mathbf{q}}{(2\pi)^3} \times \quad (2)$$




Figure 1: The pair B_c -meson production amplitudes in Higgs boson decay. B_c denotes the B_c -meson states with spin 0 and 1. Dashed line shows the Higgs boson and wavy line corresponds to the gluon.

$$\times Sp \left\{ \Psi_{B_c}^{\mathcal{P}, \mathcal{V}}(p, P) \Gamma_1^\nu(p, q, P, Q) \Psi_{B_c}^{\mathcal{P}, \mathcal{V}}(q, Q) \gamma_\nu + \Psi_{B_c}^{\mathcal{P}, \mathcal{V}}(-p, P) \Gamma_2^\nu(p, q, P, Q) \Psi_{B_c}^{\mathcal{P}, \mathcal{V}}(-q, Q) \gamma_\nu \right\},$$

where a superscript \mathcal{P} indicates a pseudoscalar B_c meson, a superscript \mathcal{V} indicates a vector B_c meson, G_F is the Fermi constant, $M = m_1 + m_2$. $\Gamma_{1,2}$ are the vertex functions defined below. The permutation of subscripts b and c in the wave functions indicates corresponding permutation in the projection operators (see below Eqs.(3)-(4). The method for producing the amplitudes in the form (2) is described in detail in our previous studies [3, 4, 15]. The transition of free quark-antiquark pair to meson bound states is described in our approach by specific wave functions. Relativistic wave functions of pseudoscalar and vector B_c mesons accounting for the transformation from the rest frame to the moving one with four momenta P , and Q are

$$\begin{aligned} \Psi_{B_c}^{\mathcal{P}}(p, P) = & \frac{\Psi_{B_c}^0(\mathbf{p})}{\sqrt{\frac{\epsilon_1(p)}{m_1} \frac{(\epsilon_1(p)+m_1)}{2m_1} \frac{\epsilon_2(p)}{m_2} \frac{(\epsilon_2(p)+m_2)}{2m_2}}} \left[\frac{\hat{v}_1 - 1}{2} + \hat{v}_1 \frac{\mathbf{p}^2}{2m_2(\epsilon_2(p) + m_2)} - \frac{\hat{p}}{2m_2} \right] \\ & \times \gamma_5(1 + \hat{v}_1) \left[\frac{\hat{v}_1 + 1}{2} + \hat{v}_1 \frac{\mathbf{p}^2}{2m_1(\epsilon_1(p) + m_1)} + \frac{\hat{p}}{2m_1} \right], \end{aligned} \quad (3)$$

$$\Psi_{B_c^*}^\nu(q, Q) = \frac{\Psi_{B_c^*}^0(\mathbf{q})}{\sqrt{\frac{\epsilon_1(q)}{m_1} \frac{(\epsilon_1(q)+m_1)}{2m_1} \frac{\epsilon_2(q)}{m_2} \frac{(\epsilon_2(q)+m_2)}{2m_2}}} \left[\frac{\hat{v}_2 - 1}{2} + \hat{v}_2 \frac{\mathbf{q}^2}{2m_1(\epsilon_1(q) + m_1)} + \frac{\hat{q}}{2m_1} \right] \\ \times \hat{\varepsilon}_\nu(Q, S_z)(1 + \hat{v}_2) \left[\frac{\hat{v}_2 + 1}{2} + \hat{v}_2 \frac{\mathbf{q}^2}{2m_2(\epsilon_2(q) + m_2)} - \frac{\hat{q}}{2m_2} \right], \quad (4)$$

where the symbol hat denotes convolution of four-vector with the Dirac gamma matrices, $v_1 = P/M_{B_c}$, $v_2 = Q/M_{B_c}$; $\varepsilon_\nu(Q, S_z)$ is the polarization vector of the $B_c^{*-}(1^-)$ meson, relativistic quark energies $\epsilon_{1,2}(p) = \sqrt{\mathbf{p}^2 + m_{1,2}^2}$. Relativistic functions (3)-(4) and the vertex production functions $\Gamma_{1,2}$ do not contain the $\delta(\mathbf{p}^2 - \eta_i^2 M_{B_c}^2 + m_{1,2}^2)$ which corresponds to the transition on the mass shell. In (3) and (4) we have complicated factor including the bound state wave function in the rest frame. Therefore instead of the substitutions $M_{B_c} = \epsilon_1(\mathbf{p}) + \epsilon_2(\mathbf{p})$ and $M_{B_c^*} = \epsilon_1(\mathbf{q}) + \epsilon_2(\mathbf{q})$ in the production amplitude we carry out the integration over the quark relative momenta \mathbf{p} and \mathbf{q} . The color part of the meson wave function in the amplitude (2) is taken as $\delta_{ij}/\sqrt{3}$ (color indexes $i, j, k = 1, 2, 3$). Relativistic wave functions in (3) and (4) are equal to the product of wave functions in the rest frame $\Psi_{B_c}^0(\mathbf{p})$ and spin projection operators that are accurate at all orders in $|\mathbf{p}|/m$. An expression of spin projector in different form for $(c\bar{c})$ system was obtained in [16] where spin projectors are written in terms of heavy quark momenta $p_{1,2}$ lying on the mass shell. Our derivation of relations (3) and (4) accounts for the transformation law of the bound state wave functions from the rest frame to the moving one with four momenta P and Q . This transformation law was discussed in the Bethe-Salpeter approach in [17] and in quasipotential method in [18].

We have omitted here intermediate expressions, leading to the equations (2)-(4) because they were discussed in detail in our previous papers. In the Bethe-Salpeter approach the initial production amplitude has as a form of convolution of the truncated amplitude with two Bethe-Salpeter (BS) B_c meson wave functions. The presence of the $\delta(p \cdot P)$ function in this case allows us to make the integration over relative energy p^0 . In the rest frame of a bound state the condition $p^0 = 0$ allows to eliminate the relative energy from the BS wave function. The BS wave function satisfies a two-body bound state equation which is very complicated and has no known solution. A way to deal with this problem is to find a soluble lowest-order equation containing main physical properties of the exact equation and develop a perturbation theory. For this purpose we continue to work in three-dimensional quasipotential approach. In this framework the double B_c meson production amplitude (2) can be written initially as a product of the production vertex function $\Gamma_{1,2}$ projected onto

the positive energy states by means of the Dirac bispinors (free quark wave functions) and a bound state quasipotential wave functions describing B_c mesons in the reference frames moving with four momenta P, Q . Further transformations include the known transformation law of the bound state wave functions to the rest frame [4, 15]. In the spin projectors we have $\mathbf{p}^2 \neq \eta_i^2 M^2 - m_{1,2}^2$ just the same as in the vertex production functions $\Gamma_{1,2}$. We can consider (3)-(4) as a transition form factors for heavy quark-antiquark pair from free state to bound state. When transforming the amplitude \mathcal{M} we introduce the projection operators $\hat{\Pi}^{\mathcal{P},\nu}$ onto the states of $(Q_1\bar{Q}_2)$ in the B_c meson with total spin 0 and 1 as follows:

$$\hat{\Pi}^{\mathcal{P},\nu} = [v_2(0)\bar{u}_1(0)]_{S=0,1} = \gamma_5(\hat{\varepsilon}^*)\frac{1 + \gamma^0}{2\sqrt{2}}. \quad (5)$$

At leading order in α_s the vertex functions $\Gamma_{1,2}^\nu(p, P; q, Q)$ can be written as $(\Gamma_2^\nu(p, P; q, Q)$ can be obtained from $\Gamma_1^\nu(p, P; q, Q)$ by means of the replacement $p_1 \leftrightarrow p_2, q_1 \leftrightarrow q_2, \alpha_b \rightarrow \alpha_c$)

$$\Gamma_1^\nu(p, P; q, Q) = r_1\alpha_b \left[\gamma_\mu \frac{(\hat{r} - \hat{q}_1 + m_1)}{(r - q_1)^2 - m_1^2 + i\epsilon} + \frac{(\hat{p}_1 - \hat{r} + m_1)}{(p_1 - r)^2 - m_1^2 + i\epsilon} \gamma_\mu \right] D^{\mu\nu}(k_1), \quad (6)$$

$$\Gamma_2^\nu(p, P; q, Q) = r_2\alpha_c \left[\gamma_\mu \frac{(\hat{r} - \hat{q}_2 + m_2)}{(r - q_2)^2 - m_2^2 + i\epsilon} + \frac{(\hat{p}_2 - \hat{r} + m_2)}{(p_2 - r)^2 - m_2^2 + i\epsilon} \gamma_\mu \right] D^{\mu\nu}(k_2), \quad (7)$$

where $r^2 = M_H^2 = (P + Q)^2 = 2M_{B_c}^2 + 2PQ$, the gluon four-momenta are $k_1 = p_1 + q_1, k_2 = p_2 + q_2, \alpha_{c,b} = \alpha_s (m_{1,2}^2 M_H^2 / (m_1 + m_2)^2 \Lambda^2)$. Relative momenta p, q of heavy quarks enter in the gluon propagators $D_{\mu\nu}(k_{1,2})$ and quark propagators as well as in relativistic wave functions (3) and (4). Accounting for the small ratio of relative quark momenta p and q to the mass M_H , we use an expansion of inverse denominators of quark and gluon propagators as follows:

$$\frac{1}{(p_1 + q_1)^2} = \frac{1}{\eta_1 \rho_1 M_H^2}, \quad \frac{1}{(p_2 + q_2)^2} = \frac{1}{\eta_2 \rho_2 M_H^2}, \quad (8)$$

$$\frac{1}{(r - q_1)^2 - m_1^2} = \frac{1}{\rho_2 M_H^2}, \quad \frac{1}{(-r - p_1)^2 - m_1^2} = \frac{1}{\eta_2 M_H^2}, \quad (9)$$

$$\frac{1}{(r - p_2)^2 - m_1^2} = \frac{1}{\eta_1 M_H^2}, \quad \frac{1}{(-r - q_2)^2 - m_1^2} = \frac{1}{\rho_1 M_H^2}. \quad (10)$$

Using expansions (8)-(10) and wave functions (3)-(4) in the amplitude (2) we hold the second-order correction for small ratios $|\mathbf{p}|/m_{1,2}, |\mathbf{q}|/m_{1,2}, |\mathbf{p}|/M_H, |\mathbf{q}|/M_H$ relative to the leading order result. As we take relativistic factors in the denominator of the amplitudes

(3) and (4) unchanged, the momentum integrals are convergent. Calculating the trace in obtained expression in the package FORM [19], we find relativistic amplitudes of the B_c meson pairs production in the form:

$$\mathcal{M}_{\mathcal{P}\mathcal{P}} = \frac{32\pi}{3M_H^4}(\sqrt{2}G_F)^{1/2}M_{B_c}M \left[\frac{\alpha_b r_1}{\eta_2^3} F_{1P} + \frac{\alpha_c r_2}{\eta_1^3} F_{2P} \right] |\tilde{\Psi}_P(0)|^2, \quad (11)$$

$$\mathcal{M}_{\mathcal{V}\mathcal{V}} = \frac{32\pi}{3M_H^4}(\sqrt{2}G_F)^{1/2}M_{B_c}M \varepsilon_1^\lambda \varepsilon_2^\sigma \left[\frac{\alpha_b r_1}{\eta_2^3} F_{1V}^{\lambda\sigma} + \frac{\alpha_c r_2}{\eta_1^3} F_{2V}^{\lambda\sigma} \right] |\tilde{\Psi}_V(0)|^2, \quad (12)$$

where ε_ν is the polarization vector of spin 1 B_c meson. The decay widths of the Higgs boson into a pair of pseudoscalar and vector B_c mesons are determined by the following expressions:

$$\Gamma_{PP} = \frac{512\sqrt{2}\pi G_F M^2 |\tilde{\Psi}_P(0)|^4 \sqrt{\frac{r_3^2}{4} - 1}}{9M_H^5 r_3^5} \left[\frac{\alpha_b r_1}{\eta_2^3} F_{1P} + \frac{\alpha_c r_2}{\eta_1^3} F_{2P} \right]^2, \quad (13)$$

$$F_{1P} = -r_1 - \eta_1 + \frac{3}{2}r_3^2 - \frac{1}{2}r_1 r_3^2 - \frac{1}{2}\eta_1 r_3^2 + \omega_{01}(-12r_2 + 2r_2 r_3^2) + \quad (14)$$

$$\omega_{10}(2r_1 + r_1 r_3^2) + \omega_{10}\omega_{01}(6r_2 - 2r_1 - \frac{3}{2}r_3^2 - r_3^2 r_2 - r_3^2 r_1),$$

$$\Gamma_{VV} = \frac{512\sqrt{2}\pi G_F M^2 |\tilde{\Psi}_V(0)|^4 \sqrt{\frac{r_3^2}{4} - 1}}{9M_H^5 r_3^5} \sum_{\lambda,\sigma} |\varepsilon_1^\lambda \varepsilon_2^\sigma \left[\frac{\alpha_b r_1}{\eta_2^3} F_{1V}^{\lambda\sigma} + \frac{\alpha_c r_2}{\eta_1^3} F_{2V}^{\lambda\sigma} \right]|^2, \quad (15)$$

$$F_{1V}^{\alpha\beta} = g_1 v_1^\alpha v_2^\beta + g_2 g^{\alpha\beta}, \quad F_{2V}^{\alpha\beta} = \tilde{g}_1 v_1^\alpha v_2^\beta + \tilde{g}_2 g^{\alpha\beta}, \quad (16)$$

$$g_1 = -1 + \frac{1}{9}\omega_{10}\omega_{01}, \quad g_2 = -r_1 - \eta_1 + \frac{1}{2}r_3^2 - \frac{4}{3}r_2\omega_{01} + 2r_1\omega_{10} + \omega_{10}\omega_{01}\left(\frac{2}{3}r_2 - \frac{2}{9} - \frac{1}{18}r_3^2\right).$$

In the nonrelativistic limit the Higgs boson decay rates acquire the form:

$$\Gamma_{PP}^{nr} = \frac{512\sqrt{2}\pi G_F M^2 |\tilde{\Psi}_P(0)|^4 \sqrt{\frac{r_3^2}{4} - 1}}{9M_H^5 r_3^5} \times \left[\frac{\alpha_b r_1}{\eta_2^3} \left(\frac{3}{2}r_3^2 - 2r_1 - r_1 r_3^2 \right) + \frac{\alpha_c r_2}{\eta_1^3} \left(\frac{3}{2}r_3^2 - 2r_2 - r_2 r_3^2 \right) \right]^2, \quad (17)$$

$$\begin{aligned}
\Gamma_{VV}^{nr} = & \frac{512\sqrt{2}\pi G_F M^2 |\tilde{\Psi}_V(0)|^4 \sqrt{\frac{r_3^2}{4} - 1}}{9M_H^5 r_3^5} \frac{1}{4r_1^6 r_2^6} \times \\
& \times \left[\alpha_b^2 r_1^8 \left(4r_1^2 (r_3^4 - 4r_3^2 + 12) - 4r_1 (r_3^2 + 2) r_3^2 + 3r_3^4 \right) + \right. \\
& + 2\alpha_b \alpha_c r_1^4 r_2^4 \left(4r_1 r_2 (r_3^4 - 4r_3^2 + 12) - 2r_1 (r_3^2 + 2) r_3^2 - 2r_2 (r_3^2 + 2) r_3^2 + 3r_3^4 \right) + \\
& \left. + \alpha_c^2 r_2^8 \left(4r_2^2 (r_3^4 - 4r_3^2 + 12) - 4r_2 (r_3^2 + 2) r_3^2 + 3r_3^4 \right) \right] \quad (18)
\end{aligned}$$

where the parameter $r_3 = \frac{M_H}{M_{B_c}}$.

The functions F_{iP} , F_{iV} entering in (11)-(12) can be written initially as series in specific relativistic factors $C_{ij} = [(m_1 - \epsilon_1(p))/(m_1 + \epsilon_1(p))]^i [(m_2 - \epsilon_2(q))/(m_2 + \epsilon_2(q))]^j$ with $i + j \leq 2$ connected with the relative momenta p and q of heavy quarks. In final form the functions F_{iP} , F_{iV} and the production cross sections contain relativistic parameters $\omega_{nk}^{P,V}$ which are determined by the momentum integrals I_{nk} and calculated in the quark model:

$$I_{nk}^{P,V} = \int_0^\infty q^2 R_{B_c}^{P,V}(q) \sqrt{\frac{(\epsilon_1(q) + m_1)(\epsilon_2(q) + m_2)}{2\epsilon_1(q) \cdot 2\epsilon_2(q)}} \left(\frac{m_1 - \epsilon_1(q)}{m_1 + \epsilon_1(q)} \right)^n \left(\frac{m_2 - \epsilon_2(q)}{m_2 + \epsilon_2(q)} \right)^k dq, \quad (19)$$

$$\omega_{10}^{P,V} = \frac{I_{10}^{P,V}}{I_{00}^{P,V}}, \quad \omega_{01}^{P,V} = \frac{I_{01}^{P,V}}{I_{00}^{P,V}}, \quad \omega_{20}^{P,V} = \frac{I_{20}^{P,V}}{I_{00}^{P,V}}, \quad \omega_{02}^{P,V} = \frac{I_{02}^{P,V}}{I_{00}^{P,V}}, \quad \omega_{11}^{P,V} = \frac{I_{11}^{P,V}}{I_{00}^{P,V}}, \quad (20)$$

$$\tilde{\Psi}_{B_c}^0(0) = \int \sqrt{\frac{(\epsilon_1(p) + m_1)(\epsilon_2(p) + m_2)}{2\epsilon_1(p) \cdot 2\epsilon_2(p)}} \Psi_{B_c}^0(\mathbf{p}) \frac{d\mathbf{p}}{(2\pi)^3}. \quad (21)$$

Another source of relativistic corrections is related with the Hamiltonian of the heavy quark bound states which allows to calculate the bound state wave functions of pseudoscalar and vector B_c mesons. The exact form of the bound state wave function $\Psi_{B_c}^0(\mathbf{q})$ is important to obtain more reliable predictions for the decay widths. In nonrelativistic approximation the pair B_c meson production cross sections contain fourth power of nonrelativistic wave function at the origin. The value of the cross sections is very sensitive to small changes of $\Psi_{B_c}^0$. In nonrelativistic QCD there exists corresponding problem of determining the magnitude of the color-singlet matrix elements [20]. To account for relativistic corrections to the meson wave functions we describe the dynamics of heavy quarks by the QCD generalization of the

standard Breit Hamiltonian in the center-of-mass reference frame [21–27]:

$$H = H_0 + \Delta U_1 + \Delta U_2, \quad H_0 = \sqrt{\mathbf{p}^2 + m_1^2} + \sqrt{\mathbf{p}^2 + m_2^2} - \frac{4\tilde{\alpha}_s}{3r} + (Ar + B), \quad (22)$$

$$\Delta U_1(r) = -\frac{\alpha_s^2}{3\pi r} [2\beta_0 \ln(\mu r) + a_1 + 2\gamma_E \beta_0], \quad a_1 = \frac{31}{3} - \frac{10}{9}n_f, \quad \beta_0 = 11 - \frac{2}{3}n_f, \quad (23)$$

$$\begin{aligned} \Delta U_2(r) = & -\frac{2\alpha_s}{3m_1 m_2 r} \left[\mathbf{p}^2 + \frac{\mathbf{r}(\mathbf{r}\mathbf{p})\mathbf{p}}{r^2} \right] + \frac{2\pi\alpha_s}{3} \left(\frac{1}{m_1^2} + \frac{1}{m_2^2} \right) \delta(\mathbf{r}) + \frac{4\alpha_s}{3r^3} \left(\frac{1}{2m_1^2} + \frac{1}{m_1 m_2} \right) (\mathbf{S}_1 \mathbf{L}) + \\ & + \frac{4\alpha_s}{3r^3} \left(\frac{1}{2m_2^2} + \frac{1}{m_1 m_2} \right) (\mathbf{S}_2 \mathbf{L}) + \frac{32\pi\alpha_s}{9m_1 m_2} (\mathbf{S}_1 \mathbf{S}_2) \delta(\mathbf{r}) + \frac{4\alpha_s}{m_1 m_2 r^3} \left[\frac{(\mathbf{S}_1 \mathbf{r})(\mathbf{S}_2 \mathbf{r})}{r^2} - \frac{1}{3}(\mathbf{S}_1 \mathbf{S}_2) \right] - \\ & - \frac{\alpha_s^2(m_1 + m_2)}{m_1 m_2 r^2} \left[1 - \frac{4m_1 m_2}{9(m_1 + m_2)^2} \right], \end{aligned} \quad (24)$$

where $\mathbf{L} = [\mathbf{r} \times \mathbf{p}]$, \mathbf{S}_1 , \mathbf{S}_2 are spins of heavy quarks, n_f is the number of flavors, $\gamma_E \approx 0.577216$ is the Euler constant. To improve an agreement of theoretical hyperfine splittings in $(\bar{b}c)$ mesons with experimental data and other calculations in quark models we add to the standard Breit potential (24) the spin confining potential obtained in [21, 28–30]:

$$\Delta V_{conf}^{hfs}(r) = f_V \frac{A}{8r} \left\{ \frac{1}{m_1^2} + \frac{1}{m_2^2} + \frac{16}{3m_1 m_2} (\mathbf{S}_1 \mathbf{S}_2) + \frac{4}{3m_1 m_2} [3(\mathbf{S}_1 \mathbf{r})(\mathbf{S}_2 \mathbf{r}) - (\mathbf{S}_1 \mathbf{S}_2)] \right\}, \quad (25)$$

where we take the parameter $f_V = 0.9$. For the dependence of the QCD coupling constant $\tilde{\alpha}_s(\mu^2)$ on the renormalization point μ^2 in the pure Coulomb term in (22) we use the three-loop result [31]

$$\tilde{\alpha}_s(\mu^2) = \frac{4\pi}{\beta_0 L} - \frac{16\pi b_1 \ln L}{(\beta_0 L)^2} + \frac{64\pi}{(\beta_0 L)^3} [b_1^2(\ln^2 L - \ln L - 1) + b_2], \quad L = \ln(\mu^2/\Lambda^2). \quad (26)$$

In other terms of the Hamiltonians (23) and (24) we use the leading order approximation for α_s . The typical momentum transfer scale in a quarkonium is of order of double reduced mass, so we set the renormalization scale $\mu = 2m_1 m_2 / (m_1 + m_2)$ and $\Lambda = 0.168$ GeV, which gives $\alpha_s = 0.265$ for $(\bar{b}c)$ meson. The coefficients b_i are written explicitly in [31]. The parameters of the linear potential $A = 0.18$ GeV² and $B = -0.16$ GeV have established values in quark models.

The numerical values of the relativistic parameters entering the cross sections (13), and (15) are obtained by the numerical solution of the Schrödinger equation [32]. They are

Table I: Numerical values of relativistic parameters (20) and decay widths of Higgs boson (13), (15) with the account of relativistic corrections.

B_c meson	$n^{2S+1}L_J$	M_{B_c} , GeV	$\Psi_{B_c}^0(0)$, GeV $^{3/2}$	$\omega_{10}^{P,V}$	$\omega_{01}^{P,V}$	$\omega_{20}^{P,V}$	$\omega_{02}^{P,V}$	$\omega_{11}^{P,V}$	Γ_{nr} , in GeV Γ_{rel} , in GeV
B_c	1^1S_0	6.275	0.250	-0.0489	-0.0060	0.0049	0.0001	0.0006	$0.56 \cdot 10^{-12}$ $0.29 \cdot 10^{-12}$
B_c^*	1^3S_1	6.317	0.211	-0.0540	-0.0066	0.0053	0.0001	0.0007	$0.56 \cdot 10^{-12}$ $0.15 \cdot 10^{-12}$

collected in Table I in which we present also the results of nonrelativistic and relativistic calculation of Higgs boson decay widths.

3. ONE LOOP CORRECTIONS

Estimating NLO corrections we calculate LO widths using the workflow, which differs from one applied within RQM. Following both workflows we have obtained the same expressions (17) and (18) for LO widths; this served as a cross-check of our calculations. Another check is the explicitly obtained zero for the prohibited process of $B_c^* B_c$ production at both LO and NLO levels.




Figure 2: Typical one loop diagrams for the paired B_c -meson production in the Higgs boson decay.

Up to the NLO accuracy $\mathcal{O}(\alpha_s^3)$ the matrix element squared is expressed as follows:

$$|\mathcal{M}|^2 = |\mathcal{M}_{LO}|^2 + 2\text{Re}(\mathcal{M}_{LO}\mathcal{M}_{NLO}^*). \quad (27)$$

The Higgs boson decay to paired B_c is described by a set of 86 diagrams at next-to-leading order. The typical diagrams are shown in Fig. 2.

The computation strategy is based on the following toolchain in Wolfram Mathematica: `FeynArts` [33] \rightarrow `FeynCalc` [34] (`FeynCalcFormLink` [35], TIDL) \rightarrow `Apart` [36] \rightarrow `FIRE` [37] \rightarrow X-package [38]. The amplitudes generated with the `FeynArts` package are further processed with `FeynCalc` package, which provides algebraic calculations with Dirac and color matrices, including the evaluation of traces. The Passarino-Veltman reduction is carried out using the TIDL library implemented in `FeynCalc`. The `Apart` function does the extra simplification of the integrals. The `FIRE` package provides the complete reduction of the integrals obtained in the previous stages to master integrals, using the IBP reduction strategy mostly based on the Laporta algorithm [39]. The master integrals are then evaluated by substitution of their analytical expressions with the help of X-package.

The conventional dimensional regularization (CDR) scheme with D -dimensional momenta (loop and external) and Dirac matrices was used. γ^5 is known to be poorly defined in D dimensions. However γ^5 matrices are canceled out in the amplitude of $H \rightarrow B_c B_c$ decay and are completely absent in the amplitude of $H \rightarrow B_c^* B_c^*$ decay. Therefore we do not face the problem of γ^5 definition estimating the one loop QCD corrections for the discussed processes.

After the `FIRE` reduction only one-, two- and three-point integrals (\mathbf{A}_0 , \mathbf{B}_0 , \mathbf{C}_0) are left in the amplitudes. Some integrals of types \mathbf{A}_0 and \mathbf{B}_0 contribute to the amplitude with the singular coefficient $\frac{1}{D-4}$. In such cases we should keep the terms of the order of $\mathcal{O}(\varepsilon)$ in the master integral expansion over ε , because these terms might contribute to the finite part of an amplitude (see [5] for details).

The so-called ‘‘On shell’’ scheme is adopted for masses and spinors renormalization and

\overline{MS} scheme is adopted for coupling constant renormalization:

$$Z_m^{OS} = 1 - \frac{\alpha_s}{4\pi} C_F C_\epsilon \left[\frac{3}{\epsilon_{UV}} + 4 \right] + O(\alpha_s^2), \quad (28)$$

$$Z_2^{OS} = 1 - \frac{\alpha_s}{4\pi} C_F C_\epsilon \left[\frac{1}{\epsilon_{UV}} + \frac{2}{\epsilon_{IR}} + 4 \right] + O(\alpha_s^2), \quad (29)$$

$$Z_g^{\overline{MS}} = 1 - \frac{\beta_0}{2} \frac{\alpha_s}{4\pi} \left[\frac{1}{\epsilon_{UV}} - \gamma_E + \ln(4\pi) \right] + O(\alpha_s^2), \quad (30)$$

where $C_\epsilon = \left(\frac{4\pi\mu^2}{m^2} e^{-\gamma_E} \right)^\epsilon$ and γ_E is the Euler constant.

The isolated singularities in the one loop amplitude $\widetilde{\mathcal{M}}_{NLO}$ are further cancelled with singular parts of \mathcal{M}_{CT} so that $\mathcal{M}_{NLO} = \widetilde{\mathcal{M}}_{NLO} + \mathcal{M}_{CT}$ remains a finite expression for the renormalized amplitude, where

$$\mathcal{M}_{CT} = Z_2^2 \mathcal{M}_{LO} \left| \begin{array}{l} m \rightarrow Z_m m \\ g_s \rightarrow Z_g g_s \end{array} \right. . \quad (31)$$

Note that the amplitudes of the pair production of B_c mesons with the emission of one soft gluon vanish if we describe B_c mesons in the color singlet model.




Figure 3: Scale dependence of the decay widths estimated within LO and NLO: the absolute values (left) and the NLO/LO ratio (right). The filled area displays the range $m_1 M_H / (m_1 + m_2) \leq \mu \leq M_H$.

The calculation results are shown in the Fig. 3 where the dependence on the scale choice is presented for the LO and NLO approaches, and in the Fig. 4, where the dependence on the quark mass values is demonstrated. Numerical values of the NLO decay widths at different energy scales are presented in Table II.

As it can be seen in the Fig. 3, the NLO/LO ratio quite slowly depends on the scale

choice both for $B_c B_c$ and $B_c^* B_c^*$. Also it is interesting that the one loop corrections essentially change the ratio of the decay widths between $B_c B_c$ and $B_c^* B_c^*$ pairs: while the LO approach predicts the ratio $\Gamma_{VV}/\Gamma_{PP} \sim 1$, the NLO corrections increase this value to $1.5 \div 1.9$ depending on the scale choice.

As it is demonstrated in the Fig. 4, the predictions can be sensitive to the choice of the quark mass ratio $m_1/(m_1 + m_2)$. However, we do not focus on these details in the current study, as we think that the problem of mass choice deserves a separate consideration (see for example [40, 41], where the problem of choice of the b quark mass was studied in details).




Figure 4: Dependence of the decay widths estimated within LO and NLO on $r = m_1/(m_1 + m_2)$: absolute values (left) and NLO/LO ratio (right). The vertical line corresponds to $r = 1.55/6.43$.




Figure 5: The dependence of the H boson decay width on the value of the wave function of the B_c meson at zero $\psi_0 = \Psi_{B_c}^0(0)$. The vertical lines correspond to values of ψ_0 in nonrelativistic approximation and with the account of relativistic corrections.

Table II: NLO decay widths at different scales. The renormalization scale μ_R and the strong coupling scale Q are chosen equal: $\mu = \mu_R = Q$.

	$\Gamma_{LO}, \text{ GeV}$	$\Gamma_{NLO}, \text{ GeV}$			
		$m_1 M_H / (m_1 + m_2)$	$M_H / 2$	$m_2 M_H / (m_1 + m_2)$	M_H
$B_c B_c$	$0.56 \cdot 10^{-12}$	$1.26 \cdot 10^{-12}$	$1.01 \cdot 10^{-12}$	$0.91 \cdot 10^{-12}$	$0.85 \cdot 10^{-12}$
NLO/LO	—	2.24	2.30	2.33	2.34
$B_c^* B_c^*$	$0.56 \cdot 10^{-12}$	$0.70 \cdot 10^{-12}$	$0.63 \cdot 10^{-12}$	$0.59 \cdot 10^{-12}$	$0.56 \cdot 10^{-12}$
NLO/LO	—	1.25	1.41	1.49	1.54

4. CONCLUSION

In this work, we investigate the process of pair production of B_c mesons in the decay of the Higgs boson. The exact amplitudes of the decay of the H boson into a pair of scalar and vector B_c mesons are constructed, in which the dependence on the relative momenta of heavy quarks (relativistic corrections) is taken into account. Then the approximate decay amplitudes are obtained, in which the second-order relativistic corrections are retained. The widths of the decay of the H boson are also calculated both in the nonrelativistic approximation and with an account of the second-order relativistic corrections. Along with relativistic corrections in decay amplitudes, we also take into account second-order relativistic corrections when calculating the wave functions of B_c mesons in the framework of the relativistic quark model. The obtained analytical expressions for the decay widths are used to carry out numerical estimates. As in the solution of various previous problems on the pair production of bound quark states in [4, 13–15], our calculations in this work show that relativistic effects are very important for finding reliable values of the Higgs boson decay widths. Taking into account all relativistic corrections leads to a decrease in the decay widths of Γ_{PP} and Γ_{VV} by several times. The main factor leading to this decrease is the value of the wave function of quark bound states at zero. Our calculations show that relativistic corrections reduce this value by 30 percent, and since the decay widths (13) and (15) include the fourth power $|\psi_{B_c}^0(0)|$, the decrease in the decay widths themselves turns out to be very significant (see the results in Table I). The change in the H boson decay width depending on $|\psi_{B_c}^0(0)|$ is shown in a separate Fig. 5 due to the importance of this nonperturbative factor. In this work, a purely quark mechanism for the production of a pair of B_c mesons in the decay of the H boson is investigated. There are other production mechanisms, which are determined, for

example, by the initial decay of the H boson into a pair of ZZ, WW and by other couplings. Our estimates of the contributions of such processes to the decay width show that they are two orders of magnitude smaller than the mechanism studied in our work.

The main parameters that give the theoretical error of the results obtained are the quark masses, the coupling constant α_s and the discarded corrections of order $O(\mathbf{p}^4)$, $O(\mathbf{q}^4)$. In the case of the Higgs boson decay amplitudes we use for the strong coupling constant α_c the two-loop approximation from (26) where the renormalization scale $\mu = \frac{m_1}{m_1+m_2}M_H$ and for α_b $\mu = \frac{m_2}{m_1+m_2}M_H$. The correction of order $O(\mathbf{p}^4)$ gives the most significant uncertainty in the theoretical value of the obtained decay widths (13), (15). Our total theoretical error of calculations is about 20 percent.

As known the one loop QCD corrections can essentially contribute to the paired quarkonia production. Therefore these corrections are also taken into account in the current study. We have found that the one loop contribution increases the width values by 1.3 – 2.3 times. In our model the relativistic corrections and one-loop effects act in different directions changing the Higgs boson decay widths obtained in the nonrelativistic approximation. As a result, it turns out that at the $\mu = m_1M_H/(m_1 + m_2)$ energy scale, the total decay widths into the pair of pseudoscalar and vector B_c mesons are the following: $\Gamma_{PP}^{tot} = 1.0 \cdot 10^{-12}$ GeV and $\Gamma_{VV}^{tot} = 0.3 \cdot 10^{-12}$ GeV.

Acknowledgments

We are grateful to A. L. Kataev for useful remarks. The work of I. N. Belov and F. A. Martynenko is supported by the Foundation for the Advancement of Theoretical Physics and Mathematics "BASIS" (grants No. 20-2-2-2-1 and No. 19-1-5-67-1). The work of A. V. Berezhnoy and A. K. Likhoded is partially supported by RFBR (grant No. 20-02-00154 A).

Appendix A: General structure of paired B_c meson production relativistic amplitudes in the leading order in Higgs boson decay

$$\mathcal{M} = \frac{4\pi}{3} M_{B_c} \times \int \frac{d\mathbf{p}}{(2\pi)^3} \int \frac{d\mathbf{q}}{(2\pi)^3} \frac{\Psi_{B_c}^0(\mathbf{p})}{\sqrt{\frac{\epsilon_1(\mathbf{p})}{m_1} \frac{(\epsilon_1(\mathbf{p})+m_1)}{2m_1} \frac{\epsilon_2(\mathbf{p})}{m_2} \frac{(\epsilon_2(\mathbf{p})+m_2)}{2m_2}}} \frac{\Psi_{B_c}^0(\mathbf{q})}{\sqrt{\frac{\epsilon_1(\mathbf{q})}{m_1} \frac{(\epsilon_1(\mathbf{q})+m_1)}{2m_1} \frac{\epsilon_2(\mathbf{q})}{m_2} \frac{(\epsilon_2(\mathbf{q})+m_2)}{2m_2}}} \times \text{Tr}\{\mathcal{T}_{12} + \mathcal{T}_{34}\}, \quad (\text{A1})$$

$$\begin{aligned} \mathcal{T}_{12} = & \Gamma_c \alpha_b \left[\frac{\hat{v}_1 - 1}{2} + \hat{v}_1 \frac{\mathbf{p}^2}{2m_2(\epsilon_2(\mathbf{p}) + m_2)} - \frac{\hat{p}}{2m_2} \right] \Sigma_{P,V}^{(1)}(1 + \hat{v}_1) \times \\ & \left[\frac{\hat{v}_1 + 1}{2} + \hat{v}_1 \frac{\mathbf{p}^2}{2m_1(\epsilon_1(\mathbf{p}) + m_1)} + \frac{\hat{p}}{2m_1} \right] \left[\frac{\hat{p}_1 - \hat{r} + m_1}{(r - p_1)^2 - m_1^2} \gamma_\mu + \gamma_\mu \frac{\hat{r} - \hat{q}_1 + m_1}{(r - q_1)^2 - m_1^2} \right] D^{\mu\nu}(k_2) \\ & \left[\frac{\hat{v}_2 - 1}{2} + \hat{v}_2 \frac{\mathbf{q}^2}{2m_1(\epsilon_1(\mathbf{q}) + m_1)} + \frac{\hat{q}}{2m_1} \right] \Sigma_{P,V}^{(2)}(1 + \hat{v}_2) \left[\frac{\hat{v}_2 + 1}{2} + \hat{v}_2 \frac{\mathbf{q}^2}{2m_2(\epsilon_2(\mathbf{q}) + m_2)} - \frac{\hat{q}}{2m_2} \right] \gamma_\nu, \end{aligned} \quad (\text{A2})$$

$$\begin{aligned} \mathcal{T}_{34} = & \Gamma_b \alpha_c \left[\frac{\hat{v}_1 - 1}{2} + \hat{v}_1 \frac{\mathbf{p}^2}{2m_1(\epsilon_1(\mathbf{p}) + m_1)} + \frac{\hat{p}}{2m_1} \right] \Sigma_{P,V}^{(1)}(1 + \hat{v}_1) \\ & \left[\frac{\hat{v}_1 + 1}{2} + \hat{v}_1 \frac{\mathbf{p}^2}{2m_2(\epsilon_2(\mathbf{p}) + m_2)} - \frac{\hat{p}}{2m_2} \right] \left[\frac{\hat{p}_2 - \hat{r} + m_2}{(r - p_2)^2 - m_2^2} \gamma_\mu + \gamma_\mu \frac{\hat{r} - \hat{q}_2 + m_2}{(r - q_2)^2 - m_2^2} \right] D^{\mu\nu}(k_1) \\ & \left[\frac{\hat{v}_2 - 1}{2} + \hat{v}_2 \frac{\mathbf{q}^2}{2m_2(\epsilon_2(\mathbf{q}) + m_2)} - \frac{\hat{q}}{2m_2} \right] \Sigma_{P,V}^{(2)}(1 + \hat{v}_2) \left[\frac{\hat{v}_2 + 1}{2} + \hat{v}_2 \frac{\mathbf{q}^2}{2m_1(\epsilon_1(\mathbf{q}) + m_1)} + \frac{\hat{q}}{2m_1} \right] \gamma_\nu, \end{aligned} \quad (\text{A3})$$

where $\Sigma_{P,V}^{(1),(2)}$ is equal to γ_5 for pseudoscalar B_c meson and $\hat{\epsilon}_\nu$ for vector B_c meson. $\Gamma_c = m_c(\sqrt{2}G_F)^{1/2}$, $\Gamma_b = m_b(\sqrt{2}G_F)^{1/2}$. The trace calculation in (A1) leads to amplitudes \mathcal{M}_{PP} and \mathcal{M}_{VV} presented in (11)-(12).

-
- [1] G. Aad et al. (ATLAS Collaboration), Phys. Lett. B **716**, 1 (2012).
 - [2] S. Chatrchyan et al. (CMS Collaboration), Phys. Lett. B **716**, 30 (2012).
 - [3] A. A. Karyasov, A. P. Martynenko and F. A. Martynenko, Nucl. Phys. B **911**, 36 (2016).
 - [4] A. V. Berezhnoy, A. P. Martynenko and F. A. Martynenko and O. S. Sukhorukova, Nucl. Phys. A **986**, 34 (2019).
 - [5] A. V. Berezhnoy, A. K. Likhoded, A. I. Onishchenko and S. V. Poslavsky, Nucl. Phys. B **915**,

- 224 (2017).
- [6] W. J. Keung, Phys. Rev. D **27**, 2762 (1983).
 - [7] M. A. Shifman and M. I. Vysotsky, Nucl. Phys. B **186**, 475 (1981).
 - [8] G. T. Bodwin, H. S. Chung, J.-H. Ee, J. Lee and F. Petriello, Phys. Rev. D **90**, 113010 (2014).
 - [9] V. Kartvelishvili, A. V. Luchinsky, and A. A. Novoselov, Phys. Rev. D **79**, 114015 (2009).
 - [10] J. Jiang and C.-F. Qiao, Phys. Rev. D **93**, 054031 (2016).
 - [11] A. M. Sirunyan et al. [the CMS Collaboration], Phys. Lett. B **797**, 134811 (2019).
 - [12] A. V. Berezhnoy, I. N. Belov, A. K. Likhoded and, A. V. Luhinsky, Mod. Phys. Lett. A **34**, 40 (2019).
 - [13] A. E. Dorokhov, R. N. Faustov, A. P. Martynenko, and F. A. Martynenko, Phys. Rev. D **102**, 016027 (2020).
 - [14] E. N. Elekina and A. P. Martynenko, Phys. Rev. D **81**, 054006 (2010).
 - [15] A. P. Martynenko and A. M. Trunin, Phys. Rev. D **86**, 094003 (2012).
 - [16] G. T. Bodwin and A. Petrelli, Phys. Rev. D **66**, 094011 (2002).
 - [17] S. J. Brodsky and J. R. Primack, Ann. Phys. **52**, 315 (1969).
 - [18] R. N. Faustov, Ann. Phys. **78**, 176 (1973).
 - [19] J. Kuipers, T. Ueda, J. A. M. Vermaseren, and J. Vollinga, Comput. Phys. Commun. **184**, 1453 (2013).
 - [20] G.T. Bodwin, E. Braaten and G.P. Lepage, Phys. Rev. D **51**, 1125 (1995).
 - [21] S. N. Gupta, S. F. Radford and W. W. Repko, Phys. Rev. D **26**, 3305 (1982).
 - [22] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Rev. Mod. Phys. **77**, 1423 (2005).
 - [23] S. Capstick and N. Isgur, Phys. Rev. D **34**, 2809 (1986).
 - [24] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
 - [25] S. S. Gershtein, V. V. Kiselev, A. K. Likhoded, and A. V. Tkabladze, Phys. Usp. **38**, 1 (1995).
 - [26] S. Godfrey, Phys. Rev. D **70**, 054017 (2004).
 - [27] D. Ebert, R.N. Faustov and V.O. Galkin, Phys. Rev. D **67**, 014027 (2003).
 - [28] S. F. Radford and W. W. Repko, Phys. Rev. D **75**, 074031 (2007).
 - [29] S. N. Gupta, Phys. Rev. D **35**, 1736 (1987).
 - [30] S. N. Gupta, J. M. Johnson, W. W. Repko and C. J. Suchyta, Phys. Rev. D **49**, 1551 (1994).
 - [31] K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, Phys. Rev. Lett. **79**, 2184 (1997).
 - [32] W. Lucha and F. F. Schöberl, Int. J. Mod. Phys. C **10**, 607 (1999).

- [33] T. Hahn, *Comput. Phys. Commun.* **140**, 418 (2001).
- [34] V. Shtabovenko, R. Mertig and F. Orellana, *Comput. Phys. Commun.* **256**, 107478 (2020).
- [35] F. Feng and R. Mertig, [arXiv:1212.3522 [hep-ph]].
- [36] F. Feng, *Comput. Phys. Commun.* **183**, 2158 (2012).
- [37] A. V. Smirnov, *JHEP* **10**, 107 (2008).
- [38] H. H. Patel, *Comput. Phys. Commun.* **218**, 66 (2017).
- [39] S. Laporta, *Int. J. Mod. Phys. A* **15**, 5087 (2000).
- [40] A. L. Kataev and V. T. Kim, *Mod. Phys. Lett. A* **9**, 1309 (1994).
- [41] A. L. Kataev and V. T. Kim, *PoS ACAT08* 004 (2008); [arXiv:0902.1442 [hep-ph]].