

DATA FOR SHIMURA VARIETIES INTERSECTING THE TORELLI LOCUS

WANLIN LI, ELENA MANTOVAN, AND RACHEL PRIES

ABSTRACT. For infinitely many Hurwitz spaces parametrizing cyclic covers of the projective line, we provide a method to determine the integral PEL datum of the Shimura variety that contains the image of the Hurwitz space under the Torelli morphism.

Keywords: abelian variety, endomorphism, complex multiplication, principal polarization, cyclotomic field, class group, curve, cyclic cover, Jacobian, moduli space, Shimura variety, PEL type, lattice, Hermitian form.

MSC20 classifications: primary 11G15, 11G18, 11G30, 14G35, 14K10; secondary 11G10, 11R18, 14H10, 14H40, 14K22.

1. INTRODUCTION

1.1. Overview. In [17], Shimura studied unitary groups associated with Hermitian spaces over algebraic number fields and their maximal lattices. In [16], he developed this theory to study isomorphism classes of polarized abelian varieties and Riemann forms. Using this, in [18], Shimura determined the lattice and Hermitian matrix associated with each of six families of cyclic covers of the projective line \mathbb{P}^1 . The lattice and Hermitian matrix determine *the integral PEL datum* of the family, as defined in Sections 1.2 and 2.5.

An important question to ask is whether Shimura’s method applies to other families of curves or to their associated Shimura varieties. As stated in [18, Section 6], it is essential to have a point in the Shimura variety that represents the Jacobian of a curve, in order to have a self-dual lattice. It is natural to start by studying PEL type Shimura varieties that are associated with families of Jacobians of degree m cyclic covers of \mathbb{P}^1 . In this context, it is easier to identify the lattice when the class number of $\mathbb{Q}(\zeta_m)$ is one. One of Shimura’s key results is that, under mild conditions, there is a unique isomorphism class of Hermitian form for a given lattice and given signature type at the infinite places, [18, Appendix, Proposition 8].

In this paper, for m an odd prime such that $\mathbb{Q}(\zeta_m)$ has class number one, we provide a method to determine the lattice and Hermitian matrix, and thus the integral PEL datum, for all positive-dimensional families of degree m cyclic covers of \mathbb{P}^1 , see Theorem 5.5. Our method also works for infinitely many families when m is not prime.

1.2. Shimura Data. Our results generalize what is known about Shimura data of the moduli spaces of principally polarized abelian varieties to the context of certain unitary Shimura varieties. In that light, the foundation of this paper is Riemann’s theorem, which states that a complex torus is an abelian variety if and only if it is polarizable. Thus the

We would like to thank the American Institute of Mathematics for their support through the Square program. Li was partially supported by the Simons Collaboration on Arithmetic Geometry, Number Theory and Computation. Pries was partially supported by NSF grant DMS-19-01819.

We would like to thank Yunqing Tang for her support and advice on this paper. We also thank Eran Assaf, Bjorn Poonen, and John Voight for helpful conversations about class groups.

category of abelian varieties over \mathbb{C} is equivalent to the category of pairs (V, Λ) , where V is a non-trivial \mathbb{C} -vector space of finite dimension and Λ is a lattice in V that admits a Riemann form which is integral on Λ , see Section 2.2.

For $g \geq 1$, let \mathcal{A}_g denote the moduli space of principally polarized abelian varieties of dimension g . Let $\Lambda = \mathbb{Z}^{2g}$ be the standard lattice in $V = \mathbb{Q}^{2g}$, together with the standard symplectic form $\Psi : V \times V \rightarrow \mathbb{Q}$, which is integral on Λ . The points of the Siegel upper half space of dimension g are complex structures J on $V_{\mathbb{R}}$ such that $\Psi_J = \Psi_{\mathbb{R}}(\cdot, J\cdot)$ is a Riemann form. These points parametrize principally polarized complex abelian varieties A of dimension g , equipped with a trivialization $\Lambda \simeq H^1(A, \mathbb{Z})$. By [13, Example 11.12], the Shimura datum for \mathcal{A}_g is given by the symplectic \mathbb{Q} -vector space (V, Ψ) .

We review Deligne's formulation of a Shimura datum in Definition 2.3. Let G denote the reductive algebraic group $G = \mathrm{GSp}(V, \Psi)$. In this formulation, the Shimura datum for \mathcal{A}_g is given by the pair (G, X) , where X is the $G(\mathbb{R})$ -conjugacy class of homomorphisms $h : \mathbb{S} \rightarrow G_{\mathbb{R}}$, where $\mathbb{S} = \mathrm{Res}_{\mathbb{C}/\mathbb{R}} G_m$ is the Deligne torus.

In this paper, we generalize the result about the Shimura datum of \mathcal{A}_g , by replacing \mathcal{A}_g by certain unitary Shimura varieties which we describe in the next section.

1.3. Results. Consider a family of cyclic covers of the projective line \mathbb{P}^1 , indexed by the monodromy datum $\gamma = (m, N, a)$, where m is the degree, N is the number of branch points, and a is the inertia type, see Section 2.1. Let Z_γ be the image of this family in \mathcal{A}_g under the Torelli morphism. In [4], Deligne and Mostow describe the smallest PEL type Shimura subvariety S_γ of \mathcal{A}_g containing Z_γ . Our goal is to compute the Shimura datum of S_γ or, more precisely, the *integral PEL datum* of S_γ as defined in Definition 2.6.

We say that a point P of Z_γ is a *distinguished point* (see Definition 5.1) if the Jacobian J_P of the curve C_P represented by P is a product of principally polarized abelian varieties each having complex multiplication. For every monodromy datum γ such that m is an odd prime, we prove in Proposition 5.2 that Z_γ has a distinguished point P such that all the abelian varieties in J_P have complex multiplication by $\mathbb{Z}[\zeta_m]$. Here J_P is the Jacobian of a curve in the boundary of the Hurwitz family; this curve is an admissible cyclic cover of a tree of projective lines, which can be explicitly computed from γ .

This provides a strategy to achieve the goal above. Suppose m is an odd prime such that $\mathbb{Q}(\zeta_m)$ has class number 1. In other words, $m = 3, 5, 7, 11, 13, 17, 19$. In Theorem 5.5, for every monodromy datum γ for degree m covers, we provide a method that determines the integral PEL datum of S_γ ; specifically, this method analyzes J_P to

- (1) express V as a vector space over $F = \mathbb{Q}(\zeta_m)$; find a \mathcal{O}_F -lattice $\Lambda \subset V$; and
- (2) explicitly find the Hermitian form $\langle \cdot, \cdot \rangle$ on V , taking integral values on Λ .

Our technique to find the Shimura datum of S_γ also applies to infinitely many monodromy data γ when m is a composite number such that $\mathbb{Q}(\zeta_m)$ has class number 1.

1.4. Comparison of methods. Our approach to this problem is somewhat different from Shimura's and may be more accessible to people with background in the areas of algebraic number theory and moduli spaces of curves (Hurwitz spaces).

One advantage of our approach is that it provides a straight-forward method to determine the lattice and the Hermitian form explicitly. We provide many examples of this in Section 6 (for m prime) and Section 7 (for m composite). The reason is that the decomposition of the Jacobian provides a basis for the lattice as a free $\mathbb{Z}[\zeta_m]$ -module and the

Hermitian form is diagonal with respect to that basis. With Shimura's approach, it is necessary to find a Witt decomposition of the signature and lattice; this is straight-forward for Shimura's six examples, five of which are of the form $y^m = f(x)$ for m an odd prime and $f(x)$ a separable polynomial, but could cause subtleties for more complicated families.

One drawback to our approach is that it only works when the CM-types of the abelian varieties produced in the method are simple. The simple property guarantees that the principal polarization on each of the abelian varieties is unique up to isomorphism; this guarantees that the Hermitian form we compute is correct. For $m = 3, 5, 11, 17$, we show that this simple condition is automatic. For $m = 7$, we compute a new distinguished point to avoid the cases where a non-simple CM-type is produced, see Section 6.4. For $m = 13, 19$, we rely on Shimura's result about uniqueness of Hermitian forms.

For composite m , there are several complications with both methods.

In this paper, we also pay careful attention to the subtleties caused by the choice of primitive m th root of unity. See Remark 6.6.

1.5. Outline. Section 2 contains information about families of cyclic covers of \mathbb{P}^1 and information about Shimura varieties and Shimura data.

Section 3 contains some results in algebraic number theory about the narrow class group and units of independent signs that we need for this paper and later papers. In Sections 3.4 - 3.6, we restrict to the case of cyclotomic fields and provide information about the narrow class number, the different, and simple types.

In Section 4, we prove a result about uniqueness of principal polarizations on abelian varieties with complex multiplication, see Propositions 4.4 and 4.5.

Section 5 contains the main result of the paper, Theorem 5.5, which determines the integral PEL data of the unitary Shimura varieties S_γ for all families S_γ when m is an odd prime such that $\mathbb{Q}(\zeta_m)$ has class number 1.

Section 6 contains examples of Theorem 5.5: for all γ when $m = 3$; and for all γ with $N = 4$ branch points when $m = 5$ and $m = 7$. In Section 6.4, we apply the technique of Theorem 5.5 to determine the integral PEL datum using a different kind of distinguished point that represents the Jacobian of a curve with extra automorphisms.

In Section 7, we provide some examples when $m = 4, 6, 10$ to illustrate that the technique of Theorem 5.5 sometimes works when m is composite.

Remark 1.1. In [15], Moonen proved that there are exactly 20 special families of cyclic covers of \mathbb{P}^1 , up to equivalence. The family with monodromy datum γ is *special* if Z_γ is open and dense in S_γ . The six families in Shimura's paper are $M[6]$, $M[10]$, $M[8]$, $M[11]$, $M[16]$, and $M[17]$, where $M[n]$ denotes the n th row in [15, Table 1]. As an application of Theorem 5.5, we determine the integral PEL datum for twelve of Moonen's special families including the six families from [18]. These provide nice examples but we emphasize that the special property is not necessary for either method.

2. HURWITZ FAMILIES AND SHIMURA VARIETIES

2.1. Families of cyclic covers of the projective line. As in [12, Section 2.2], we consider the universal family of μ_m -covers $\psi : \mathcal{C} \rightarrow \mathbb{P}^1$, branched at N points, with *inertia type* $a = (a(1), \dots, a(N))$; the data $\gamma = (m, N, a)$ is called a *monodromy datum*. Over $\mathbb{Q}(\zeta_m)$, such a cover ψ has an equation of the form $y^m = \prod_{i=1}^N (x - t(i))^{a(i)}$, together with a choice

of automorphism of order m given by $(x, y) \mapsto (x, \zeta_m y)$. By [12, Equation 2.2], the genus of the fibers of \mathcal{C} is

$$(2.1) \quad g = g_\gamma = 1 + \frac{(N-2)m - \sum_{i=1}^N \gcd(a(i), m)}{2}.$$

The *signature type* of the monodromy datum γ is a function $\mathfrak{f} : \text{Hom}(\mathbb{Q}[\mu_m], \mathbb{C}) \rightarrow \mathbb{Z}_{\geq 0}$ which we denote by $\mathfrak{f} = (\mathfrak{f}(\tau_1), \dots, \mathfrak{f}(\tau_{m-1}))$, where τ_n and $\mathfrak{f}(\tau_n)$ are defined as follows. For $0 \leq n < m$, let $\tau_n : \mathbb{Q}[\mu_m] \rightarrow \mathbb{C}$ be given by $\tau_n(\zeta_m) = e^{2\pi i n/m}$. We identify

$$\mathbb{Z}/m\mathbb{Z} = \text{Hom}(\mathbb{Q}[\mu_m], \mathbb{C}) \text{ by } n \mapsto \tau_n.$$

We define $\mathfrak{f}(\tau_n)$ to be the dimension of the eigenspace of $H^0(\mathcal{C}(\mathbb{C}), \Omega^1)$ on which the chosen automorphism of order m acts by multiplication by ζ_m^n .

There is a formula for $\mathfrak{f}(\tau_n)$ as follows. For any $x \in \mathbb{Q}$, let $\langle x \rangle$ denote the fractional part of x . By [15, Lemma 2.7, §3.2] (or [4]),

$$(2.2) \quad \mathfrak{f}(\tau_n) = \begin{cases} -1 + \sum_{i=1}^N \langle \frac{-na(i)}{m} \rangle & \text{if } n \not\equiv 0 \pmod{m} \\ 0 & \text{if } n \equiv 0 \pmod{m}. \end{cases}$$

Next, we describe how the inertia type a and signature \mathfrak{f} change under the action of $\text{Aut}(\mu_m) \simeq (\mathbb{Z}/m\mathbb{Z})^*$. Let $\sigma_i \in \text{Aut}(\mu_m)$ denote the automorphism such that $\sigma_i(\zeta_m) = \zeta_m^i$.

Lemma 2.1. *The automorphism $\sigma_i \in \text{Aut}(\mu_m)$ takes the inertia type $a = (a_1, a_2, \dots, a_N)$ to a' and the signature \mathfrak{f} to \mathfrak{f}' as follows:*

$$a' = i^{-1} \cdot a = (i^{-1} \cdot a_1, i^{-1} \cdot a_2, \dots, i^{-1} \cdot a_N) \text{ and } \mathfrak{f}'(\tau_n) = \mathfrak{f}(\tau_{ni^{-1}}).$$

Proof. The j th entry a_j of the inertia type signifies that the canonical generator of inertia above the j th branch point is the a_j th power of the generator ζ_m of μ_m . The canonical generator of inertia is the $i^{-1}a_j$ th power of the new generator ζ_m^i of μ_m , so the j th entry of the inertia type a' is $i^{-1}a_j$.

The formula $\mathfrak{f}'(\tau_n) = \mathfrak{f}(\tau_{ni^{-1}})$ can be deduced from the formula for a' and (2.2). Alternatively, it can be seen directly since the automorphism σ_i permutes the eigenspaces for the action of the chosen automorphism on $H^0(\mathcal{C}(\mathbb{C}), \Omega^1)$, taking the eigenspace indexed by n to the one indexed by ni^{-1} . \square

Let $\gamma = (m, N, a)$ be a monodromy datum. Consider the Hurwitz family of μ_m -covers of \mathbb{P}^1 with monodromy datum γ . As in [12, Definition 2.1], let $Z^0 = Z_\gamma^0$ denote the image of this family in \mathcal{A}_g . Let $Z = Z_\gamma$ denote the closure of Z_γ^0 in \mathcal{A}_g . A point P in Z_γ^0 represents the Jacobian of a smooth curve C_P , for which there is a μ_m -cover $\psi_P : C_P \rightarrow \mathbb{P}^1$ with monodromy datum γ . A point P in $Z_\gamma - Z_\gamma^0$ represents the Jacobian of a singular curve C_P of compact type, for which there is an admissible μ_m -cover $\psi_P : C_P \rightarrow T$ with monodromy datum γ , where T is a tree of projective lines.

2.2. Complex abelian varieties. Let $g \geq 1$ and let $V = \mathbb{R}^{2g}$. Suppose V has a complex structure and $\Lambda \subset V$ is a lattice of rank $2g$. A Riemann form on a pair (V, Λ) is an alternating pairing $\Psi : \Lambda \times \Lambda \rightarrow \mathbb{Z}$ such that the pairing $\Psi_{\mathbb{R}}(\cdot, \sqrt{-1}\cdot) : V \times V \rightarrow \mathbb{R}$ is symmetric and positive definite. Every complex abelian variety of dimension g is isomorphic to V/Λ for some pair (V, Λ) that admits a Riemann form.

By Riemann's Theorem, the category of polarized abelian varieties of dimension g over \mathbb{C} is equivalent to the category of pairs (V, Λ) , where V is a non-trivial \mathbb{C} -vector space of dimension g and Λ is a lattice in V together with a Riemann form.

Remark 2.2. Let $\mathbb{S} = \text{Res}_{\mathbb{C}/\mathbb{R}}\mathbb{G}_m$ be the Deligne torus. A Hodge structure on V is a homomorphism $h : \mathbb{S} \rightarrow GL_V$. For Hodge structures of type $\{(-1, 0), (0, -1)\}$, the \mathbb{R} -linear map $h(i)$, called the Weil operator, defines a complex structure J on V . By the work of Deligne, a complex structure on V gives equivalent data as a Hodge structure on V of type $\{(-1, 0), (0, -1)\}$.

2.3. Definition of Shimura data. Recall that $\mathbb{S} = \text{Res}_{\mathbb{C}/\mathbb{R}}\mathbb{G}_m$ is the Deligne torus.

Definition 2.3. [13, Definition 11.11]. A *Shimura datum* is a pair (G, X) consisting of a reductive algebraic group G over \mathbb{Q} and a $G(\mathbb{R})$ -conjugacy class of homomorphisms $\mathbb{S} \rightarrow G_{\mathbb{R}}$ satisfying certain conditions [5, 2.1.1(1-3)].

Let \mathbb{A} (resp \mathbb{A}_f) denote the ring of adèles (resp. finite adèles) of \mathbb{Q} . To any Shimura datum (G, X) , there is an associated compatible system of smooth complex manifolds $G(\mathbb{Q}) \backslash (X \times G(\mathbb{A}_f)/K)$, for K any (sufficiently small) open compact subgroup of $G(\mathbb{A}_f)$. These are the *Shimura varieties* associated with the datum (G, X) .

Let (G, X) is a Shimura datum and $H \subset G$ be an algebraic subgroup over \mathbb{Q} . Define

$$(2.3) \quad Y_H = \{h \in X \mid h : \mathbb{S} \rightarrow H_{\mathbb{R}} \subseteq G_{\mathbb{R}}\}.$$

The group $H(\mathbb{R})$ acts by conjugation on the set Y_H and Y_H decomposes as a disjoint union of finitely many $H(\mathbb{R})$ -orbits $Y \subseteq Y_H$. We say that (H, Y) is a *sub Shimura datum* of (G, X) .

If (H, Y) is a sub Shimura datum of a Shimura datum (G, X) , then the Shimura varieties associated with (H, Y) are naturally subvarieties of those associated with (G, X) .

Example 2.4. For $g \geq 1$, let $V = \mathbb{Q}^{2g}$ and consider the lattice $\Lambda = \mathbb{Z}^{2g}$. Let Ψ be the standard symplectic form on V . Let $G_g = \text{GSp}(V, \Psi)$ be the algebraic group of symplectic similitudes over \mathbb{Q} . Let X_g be the $G_g(\mathbb{R})$ -conjugacy class of homomorphisms $h : \mathbb{S} \rightarrow G_{g\mathbb{R}}$. Then (G_g, X_g) is a Shimura datum.

Furthermore, a homomorphism $h : \mathbb{S} \rightarrow GL_{V_{\mathbb{R}}}$ which factors through $G_{g\mathbb{R}} \subset GL_{V_{\mathbb{R}}}$ corresponds to a complex structure J on $V_{\mathbb{R}}$ such that the pairing $\psi_J(\cdot, \cdot) = \psi(\cdot, J\cdot)$ is a Riemann form. Thus, X_g is identified with the space of all complex structures J on $V_{\mathbb{R}}$ such that $(x, y) \mapsto \psi(x, Jy)$ is positive definite, see [13, Example 11.12]. It follows that (G_g, X_g) is the Shimura datum for \mathcal{A}_g , the moduli space of principally polarized abelian varieties of dimension g . That is, there is an identification between $\mathcal{A}_g(\mathbb{C})$ and $G_g(\mathbb{Q}) \backslash (X_g \times G_g(\mathbb{A}_f)/K_0)$, for K_0 the stabilizer of Λ in $G_g(\mathbb{A}_f)$, i.e., $K_0 = \text{GSp}(\Lambda, \Psi)(\hat{\mathbb{Z}})$ for $\hat{\mathbb{Z}} \subset \mathbb{A}_f$ the profinite completion of \mathbb{Z} .

Definition 2.5. A Shimura datum is of *Hodge type* if it is a sub Shimura datum of (G_g, X_g) , for some $g \geq 1$.

Shimura varieties of Hodge type are realized as subvarieties of \mathcal{A}_g .

2.4. Shimura data of PEL type. We consider a subclass of Shimura data of Hodge type.

As in [7, Section 4], a *Shimura datum of PEL type* is a Shimura datum arising from a *PEL datum* $(B, *, V, \langle, \rangle, h)$ defined as follows. Let B be a semisimple finite dimensional

\mathbb{Q} -algebra, with a positive involution $*$. Let V be a finitely generated B -module, together with a non-degenerate skew-symmetric $*$ -Hermitian form $\langle \cdot, \cdot \rangle$ on V . Let $h : \mathbb{C} \rightarrow \text{End}_{B \otimes \mathbb{R}}(V_{\mathbb{R}})$ be an \mathbb{R} -algebra homomorphism satisfying $\langle h(z)v, w \rangle = \langle v, h(\bar{z})w \rangle$, for all $v, w \in V_{\mathbb{R}}$ and $z \in \mathbb{C}$. Assume that the symmetric bilinear form $\langle \cdot, h(i)\cdot \rangle$ on V is positive definite.

Consider the algebraic group H over \mathbb{Q} whose points over a \mathbb{Q} -algebra R are given by

$$(2.4) \quad H(R) = \{x \in \text{End}_{B \otimes R}(V_R) \mid xx^* \in R^\times\}.$$

Then the restriction of h to \mathbb{C}^\times defines a homomorphism of real algebraic groups $\mathbb{S} \rightarrow H_{\mathbb{R}}$. Define Y to be the $H(\mathbb{R})$ -conjugacy class of h . The pair (H, Y) is a Shimura datum; it is the Shimura datum associated to the PEL datum $(B, *, V, \langle \cdot, \cdot \rangle, h)$.

By definition, $H = GL_B(V) \cap GSp(V, \langle \cdot, \cdot \rangle)$. Thus (H, Y) can be realized as a sub Shimura datum of (G_g, X_g) for $g = \dim_{\mathbb{Q}}(V)$; hence (H, Y) is a Shimura datum of Hodge type.

To any PEL datum $(B, *, V, \langle \cdot, \cdot \rangle, h)$, there is an associated PEL moduli space $\text{Sh}(H, Y) = \text{Sh}(B, *, V, \langle \cdot, \cdot \rangle, h)$, which is a subvariety of \mathcal{A}_g , see [7, Section 5]. It is defined over the reflex field of (H, Y) which is a finite extension of \mathbb{Q} . By construction, $\text{Sh}(H, Y)$ only depends on $(H_{\mathbb{A}}, Y)$. When the Hasse principle fails, $\text{Sh}(H, Y)(\mathbb{C})$ is the disjoint union of finitely many Shimura varieties, associated with all \mathbb{Q} -forms of $H_{\mathbb{A}}$, see [7, Sections 7-8].

2.5. Integral PEL data. Let $(B, *, V, \langle \cdot, \cdot \rangle, h)$ be a PEL datum, and (H, Y) the associated Shimura datum.

Definition 2.6. An *integral PEL datum* of the PEL datum $(B, *, V, \langle \cdot, \cdot \rangle, h)$ consists of a tuple $(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$ where: \mathcal{O}_B is an order of B that is $*$ -stable; and Λ is a lattice of V that is \mathcal{O}_B -stable; such that the $*$ -Hermitian form $\langle \cdot, \cdot \rangle$ is integral on Λ .

Sometimes an *integral PEL datum* is called an *integral model* of $(B, *, V, \langle \cdot, \cdot \rangle, h)$.

A rational prime p is *good* for an integral PEL datum if \mathcal{O}_B is maximal at p and Λ is self-dual at p . All but finitely many primes are good, [8, Definition 1.4.1.1]. If p is a good prime for an integral model of $(B, *, V, \langle \cdot, \cdot \rangle, h)$, then the PEL type modular variety $\text{Sh}(H, Y)$ has good reduction at p .

To an integral PEL datum $(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$, there is an associated (canonical) integral model $\text{Sh}(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$ of $\text{Sh}(H, Y)$, defined over the ring of integers of the reflex field of (H, Y) localized at good primes, [8, Theorem 1.4.1.11]. If p is a good prime for the integral PEL datum, then $\text{Sh}(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$ has good reduction at p .¹

2.6. Shimura varieties of Deligne and Mostow. Let $\gamma = (m, N, a)$ be a monodromy datum, and let $Z = Z_\gamma$ be as in Section 2.1. In [4], Deligne and Mostow observe that there is a smallest subvariety of PEL type S_γ containing Z_γ . The Shimura datum $(H, Y) = (H_\gamma, Y_\gamma)$ of S_γ is associated with a PEL datum $(B, *, V, \langle \cdot, \cdot \rangle, h)$ as in Section 2.4, satisfying the following conditions:

- $B = \mathbb{Q}[\mu_m]$ is the group algebra of μ_m over \mathbb{Q} ;
- $*$ is the involution on $\mathbb{Q}[\mu_m]$ induced by the inverse map on μ_m ;
- $\dim_{\mathbb{Q}}(V) = 2g_\gamma$.

Our goal is to compute the integral PEL datum of S_γ .

In [4], Deligne and Mostow compute the signature type $\mathfrak{f} = \mathfrak{f}_\gamma$ associated with the monodromy datum γ , see (2.2). A signature type $\mathfrak{f} : (\mathbb{Z}/m\mathbb{Z})^* \rightarrow [0, g] \cap \mathbb{Z}$ uniquely

¹More precisely, for any level K which is neat and hyperspecial at p , the scheme has good reduction at p .

determines up to isomorphism: the structure of V as B -module; the complex structure on $V_{\mathbb{R}}$; and the group $H_{\mathbb{R}}$.

Hence, our goal is to explicitly compute the skew Hermitian form $\langle \cdot, \cdot \rangle$ on V . More precisely, let $\mathcal{O}_B = \mathbb{Z}[\mu_m]$, which is a $*$ -stable order of B . We compute a \mathcal{O}_B -stable lattice Λ of V and a skew Hermitian form $\langle \cdot, \cdot \rangle$ on V which is integral on Λ . Furthermore, for m prime, the lattice Λ is self-dual away from m . That is, we compute an integral PEL datum $(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$ for S_{γ} , for which all primes except m are good.

2.7. Strategy. If Z is an irreducible algebraic subvariety of \mathcal{A}_g , then there is a unique smallest subvariety of Hodge type S containing Z . The subgroup $H \subset G$ in the Shimura datum of S is not uniquely determined, but it can be taken to be the generic Mumford–Tate group on Z ; in particular, it can be computed at a point of Z , see [14, Sections 2.8 and 2.9, also 3.2]. For a monodromy datum $\gamma = (m, N, a)$, it thus suffices to compute the Shimura datum at a distinguished point P of Z_{γ} (Definition 5.1); when m is an odd prime, we prove that such a point P exists in Proposition 5.2. Write the first Betti cohomology of the curve C_P as $V = H^1(C_P(\mathbb{C}), \mathbb{Q})$.

Assume m is prime. We write $F = \mathbb{Q}(\zeta_m)$ for the m -th cyclotomic field, and $\mathcal{O}_F = \mathbb{Z}[\zeta_m]$ for its ring of integers. From the signature type $\mathfrak{f} = \mathfrak{f}_{\gamma}$, we deduce that $V \simeq F^r$ as $\mathbb{Q}[\mu_m]$ -modules, where the action of $\mathbb{Q}[\mu_m]$ on F^r is induced by the natural homomorphism $\mathbb{Q}[\mu_m] \rightarrow F = \mathbb{Q}(\zeta_m)$, and $r = 2g/(m-1)$, for $g = g_{\gamma}$ as in (2.1). Also, $r = \mathfrak{f}(\tau) + \mathfrak{f}(\tau^*)$ for any $\tau : F \rightarrow \mathbb{C}$ by (2.2); the complex structure on $V_{\mathbb{R}}$ is given as $V_{\mathbb{R}} \simeq (F \otimes_{\mathbb{Q}} \mathbb{R})^r \simeq \bigoplus_{\tau} \mathbb{C}^{\mathfrak{f}(\tau)}$. The standard lattice $\Lambda \simeq \mathcal{O}_F^r$ defines a lattice $\Lambda \subset V$.

In Theorem 5.5, we construct an explicit Hermitian form ψ on V , with signature $\mathfrak{f} = \mathfrak{f}_{\gamma}$, which is integral on Λ . When F has class number 1, for $m = 3, 5, 7, 11, 17$, we give a new proof that there is a unique possibility for the lattice and Hermitian form for S_{γ} , see Remark 5.6. Thus we have computed the integral PEL datum $(\mathbb{Z}[\mu_m], *, \Lambda, \psi, h)$ defining the Shimura datum (H_{γ}, Y_{γ}) of S_{γ} .

2.8. Complex multiplication. Let L be a CM-field and let L_0 be the maximal totally real subfield of L . Let $n = [L_0 : \mathbb{Q}]$; hence, $[L : \mathbb{Q}] = 2n$. A CM-type of L is an ordered set Φ of distinct embeddings $\phi_i : L \hookrightarrow \mathbb{C}$, for $1 \leq i \leq n$, no two of which are complex conjugate. A CM-type of L is called simple if it is not induced from the CM-type of a proper CM-subfield of L .

Let A be a complex torus such that $L \subset \text{End}(A) \otimes \mathbb{Q}$. We say that A is of CM-type (L, Φ) if $\dim(A) = n$ and the complex representation of L on $\text{Lie}(A)$ is isomorphic to $\sum_{\phi \in \Phi} \phi$. If, in addition, $\text{End}(A) \simeq \mathcal{O}_L$, we say A has type (\mathcal{O}_L, Φ) .

By [11, Lemma 3.1], when $N = 3$, the Jacobian of a μ_m -cover $\psi : \mathcal{C} \rightarrow \mathbb{P}^1$, branched at 3 points, with inertia type $a = (a_1, a_2, a_3)$ and signature type \mathfrak{f} , has complex multiplication by $\prod_d \mathbb{Q}(\zeta_d)$, where d satisfies $1 < d \mid m$, and $d \nmid a(i)$ for $1 \leq i \leq 3$. Its CM-type Φ satisfies

$$(2.5) \quad \phi \in \Phi \text{ if and only if } \mathfrak{f}(\phi) > 0.$$

The maximal order in $\prod_d \mathbb{Q}(\zeta_d)$ is $\prod_d \mathbb{Z}[\zeta_d]$. The image of the group ring $\mathbb{Z}[\mu_m]$ in $\prod_d \mathbb{Q}(\zeta_d)$ has finite index inside the maximal order. This index is 1 if and only if there is a unique integer d satisfying $1 < d \mid m$, and $d \nmid a(i)$ for $1 \leq i \leq 3$ (for example, if m is prime).

The next result is immediate from Lemma 2.1 and (2.5).

Lemma 2.7. *If $N = 3$, then $\sigma_i \in \text{Aut}(\mu_m)$ takes the CM-type Φ to $\Phi' = i \cdot \Phi = \{\tau_{ni} \mid \tau_n \in \Phi\}$.*

For any CM-type (L, Φ) , there exists a unique 0-dimensional PEL type moduli space $\text{Sh} = \text{Sh}(L, \Phi)$ whose points represent abelian varieties with complex multiplication of type (\mathcal{O}_L, Φ) . (When $L = \mathbb{Q}(\zeta_m)$, the points of Sh represent the Jacobians of curves for which there is a μ_m -cover of \mathbb{P}^1 branched at $N = 3$ points with CM-type Φ .) The signature type \mathfrak{f} of Sh is given as:

$$(2.6) \quad \text{for each } \phi : L \rightarrow \mathbb{C} : \mathfrak{f}(\phi) = 1 \text{ if } \phi \in \Phi \text{ and } \mathfrak{f}(\phi) = 0 \text{ if } \phi \notin \Phi.$$

For any signature type \mathfrak{f} on L taking values in $\{0, 1\}$, there is a unique CM-type Φ of L compatible with \mathfrak{f} .

3. SOME BACKGROUND FROM ALGEBRAIC NUMBER THEORY

3.1. Class group and narrow class group. For a number field L : let \mathcal{O}_L denote its ring of integers; let \mathcal{U}_L denote the units in \mathcal{O}_L ; and let $D_{L/\mathbb{Q}}$ denote the different of L over \mathbb{Q} .

Let cl_L be the class group of L . Recall that $\text{cl}_L = I_L/P_L$, where I_L is the group of non-zero fractional ideals and P_L is the subgroup of non-zero principal fractional ideals. Let $h_L = |\text{cl}_L|$ be the class number of L .

An element $\alpha \in L$ is totally positive if it is positive under every real embedding of L . Let $\mathcal{U}_L^+ \subset \mathcal{U}_L$ be the subgroup of totally positive units. Let $P_L^+ \subset P_L$ be the subgroup of principal ideals generated by a totally positive element. The narrow class group of L is $\text{cl}_L^+ = I_L/P_L^+$ and the narrow class number is $h_L^+ = |\text{cl}_L^+|$. There is a surjection

$$\nu_L : \text{cl}_L^+ \rightarrow \text{cl}_L.$$

3.2. Units of independent signs. Let L be a CM-field and L_0 its maximal totally real subfield. Let $n = [L_0 : \mathbb{Q}]$; hence, $[L : \mathbb{Q}] = 2n$. By [20, Theorem 4.10], h_{L_0} divides h_L .

We fix an ordering τ_1, \dots, τ_n of the n real embeddings $L_0 \hookrightarrow \mathbb{R}$. If L_0/\mathbb{Q} is Galois, we identify these with the elements of $\text{Gal}(L_0/\mathbb{Q})$. Consider the group homomorphism

$$(3.1) \quad \rho_{L_0} : \mathcal{U}_{L_0} \rightarrow \{\pm 1\}^n, \quad \rho_{L_0}(u) = (\tau_i(u)/|\tau_i(u)|)_{1 \leq i \leq n} \text{ for } u \in \mathcal{U}_{L_0}.$$

We say that L_0 has *units of independent signs* if, for each real embedding τ , there is a unit which is negative under τ but positive under all other real embeddings, see [2, Definition 12.1]. This is equivalent to saying that ρ_{L_0} is surjective or that $\nu_{L_0} : \text{cl}_{L_0}^+ \rightarrow \text{cl}_{L_0}$ is an isomorphism, see [2, Lemma 11.2].

We say that L_0 has *units of almost independent signs* if every unit in \mathcal{U}_{L_0} is negative under an even number of real embeddings and, for every pair of real embeddings, there is a unit which is negative under exactly the two embeddings in that pair, see [2, Definition 12.13]. This condition is equivalent to $|\ker(\nu_{L_0})| = 2$ or $|\text{coker}(\rho_{L_0})| = 2$, see [2, Lemma 11.2].

3.3. Norms and the Hasse unit index. Consider the norm map $N : \mathcal{U}_L \rightarrow \mathcal{U}_{L_0}$ given by $N(y) = N_{L/L_0}(y)$ for $y \in \mathcal{U}_L$. By Dirichlet's Unit Theorem, $[\mathcal{U}_{L_0} : \mathcal{U}_{L_0}^2] = 2^n$.

Lemma 3.1. *Suppose L_0 is a totally real field, and L/L_0 is a CM-extension. Then*

$$(3.2) \quad \mathcal{U}_{L_0}^2 \subseteq N(\mathcal{U}_L) \subseteq \mathcal{U}_{L_0}^+ \subseteq \mathcal{U}_{L_0}.$$

Proof. By definition, $\mathcal{U}_{L_0}^+ \subseteq \mathcal{U}_{L_0}$. All elements in $N(\mathcal{U}_L)$ are totally positive units because L is a CM-field, quadratic over its totally real subfield L_0 . So $N(\mathcal{U}_L) \subseteq \mathcal{U}_{L_0}^+$. Also, $\mathcal{U}_{L_0}^2 = N(\mathcal{U}_{L_0})$, hence $\mathcal{U}_{L_0}^2 \subset N(\mathcal{U}_L)$. \square

Let μ_L be the group of roots of unity of L .

Definition 3.2. The *Hasse unit index* of a CM-extension L/L_0 is $Q(L) = [\mathcal{U}_L : \mu_L \mathcal{U}_{L_0}]$.

By [20, Theorem 4.12], $Q(L) = 1$ or $Q(L) = 2$. Since $\text{Ker}(N) = \mu_L$, it follows that $\mathcal{U}_L = \mu_L \mathcal{U}_{L_0}$ if and only if $N(\mathcal{U}_L) = N(\mathcal{U}_{L_0})$. Also $N(\mathcal{U}_{L_0}) = \mathcal{U}_{L_0}^2$. Thus

$$(3.3) \quad Q(L) = [N(\mathcal{U}_L) : \mathcal{U}_{L_0}^2].$$

Let t be the number of finite primes ramified in L/L_0 .

Lemma 3.3. *Suppose L has odd class number. Let L_0 be a totally real field, and L/L_0 a CM-extension. Then $Q(L) = 1$ if and only if $t = 1$; and $Q(L) = 2$ if and only if $t = 0$.*

Proof. The material in [2, Chapter 13] is stated in terms of the type of the CM-extension L/L_0 . By a theorem of Kummer, see [2, Theorem 13.4], L/L_0 has type I (resp. II) if and only if $Q(L) = 1$ (resp. $Q(L) = 2$). The result is then immediate from [2, page 73]. \square

We summarize the result from this section that we need in this paper and later papers.

Lemma 3.4. *Suppose L is a CM-field with maximal totally real subfield L_0 .*

- (1) *Then L_0 has units of independent signs if and only if $Q(L)[\mathcal{U}_{L_0}^+ : N(\mathcal{U}_L)] = 1$.*
- (2) *Also L_0 has units of almost independent signs if and only if $Q(L)[\mathcal{U}_{L_0}^+ : N(\mathcal{U}_L)] = 2$.*

Proof. (1) By [2, Lemma 12.2], L_0 has units of independent signs if and only if every element of $\mathcal{U}_{L_0}^+$ is a square. The result then follows from Lemma 3.1.

- (2) By definition, L_0 has units of almost independent signs if and only if $[\mathcal{U}_{L_0} : \mathcal{U}_{L_0}^+] = 2^{n-1}$. Since $[\mathcal{U}_{L_0} : \mathcal{U}_{L_0}^2] = 2^n$, this is equivalent to $[\mathcal{U}_{L_0}^+ : \mathcal{U}_{L_0}^2] = 2$. \square

3.4. Cyclotomic fields. Let m be a positive integer and let $\zeta_m = e^{2\pi i/m} \in \mathbb{C}$. Let $\mu_m \subset \mathbb{C}$ be the group of m -th roots of unity. Let $\mathbb{Q}[\mu_m]$ be the group algebra of μ_m over \mathbb{Q} .

The cyclotomic field $F = \mathbb{Q}(\zeta_m)$ is a CM-field over \mathbb{Q} with maximal totally real subfield $F_0 = \mathbb{Q}(\zeta_m + \zeta_m^{-1})$. The degree of F_0 over \mathbb{Q} is $n = \phi(m)/2$.

Notation 3.5. The choice of ζ_m fixes an embedding $\sigma_1 : F \hookrightarrow \mathbb{C}$. For $1 \leq i < m$, with $\text{gcd}(i, m) = 1$, let σ_i be the embedding $F \hookrightarrow \mathbb{C}$ (or automorphism in $\text{Gal}(F/\mathbb{Q})$) determined by $\sigma_i(\zeta_m) = \zeta_m^i$. For $x \in F$, let $\bar{x} = \sigma_{m-1}(x)$ denote its complex conjugate.

If $m = p^r$ is a prime power, then F/F_0 is ramified at the unique prime of F_0 above p and at the n infinite primes of F_0 and is unramified at all other primes, [20, Proposition 2.15].

Lemma 3.6. *Let $F = \mathbb{Q}(\zeta_m)$ and $F_0 = \mathbb{Q}(\zeta_m + \zeta_m^{-1})$. Suppose F has class number 1.*

- (1) *If m is a prime power (or twice a prime power), then F_0 has narrow class number 1 and thus has units of independent signs. The complete list of these m is:*

$$m = 1, 2, 4, 8, 16, 32;$$

$$m = 3, 5, 7, 9, 11, 13, 17, 19, 25, 27; \text{ or twice a value of } m \text{ on this row.}$$

- (2) *If m is not a prime power (or twice a prime power), then F_0 has narrow class number 2 and thus has units of almost independent signs. The complete list of these m is:*

$$m = 12, 16, 20, 24, 28, 36, 40, 44, 48, 60, 84;$$

$$m = 15, 21, 33, 35, 45; \text{ or twice a value of } m \text{ on this row.}$$

Proof. The complete list of m such that $h_F = 1$ is well-known; it is the union of the lists in parts (1) and (2). For these m , since $h_F = 1$, also $h_{F_0} = 1$.

- (1) By a result originally due to Hasse, see [2, Corollary 3.9], since h_F and h_{F_0} are odd, then F_0 has units of independent signs if and only if F/F_0 is ramified at exactly one finite prime. This happens if and only if $m = p^r$ or $m = 2 \cdot p^r$, for some prime p .
- (2) Since h_F and h_{F_0} are odd, then F_0 has units of almost independent signs if and only if F/F_0 is not ramified at any finite prime, see [2, Corollary 13.10]. This happens if m is not a prime power (or twice a prime power).

□

3.5. A generator for the different of cyclotomic fields.

Lemma 3.7. *Let $F = \mathbb{Q}(\zeta_m)$. The element β_0 below generates $D_{F/\mathbb{Q}}$ and $\beta_0 = -\bar{\beta}_0$.*

- (1) *If m is an odd prime, then*

$$\beta_0 = m / (\zeta_m^{(m+1)/2} - \zeta_m^{(m-1)/2}).$$

- (2) *If $m = 2^k$ with $k \geq 2$, then $\beta_0 = -2^{k-1}i$.*
- (3) *If $m = 3^k$, then $\beta_0 = -3^{k-1}\sqrt{3}i$.*
- (4) *If $m = pq$ with p, q distinct odd primes, then*

$$\beta_0 = m \frac{\zeta_m^{(m+1)/2} - \zeta_m^{(m-1)/2}}{(\zeta_m^{q(p+1)/2} - \zeta_m^{q(p-1)/2})(\zeta_m^{p(q+1)/2} - \zeta_m^{p(q-1)/2})}.$$

Proof. In each case, β_0 is on the imaginary axis so condition (2) is satisfied. The different $D_{F/\mathbb{Q}}$ is generated by $\langle c'_m(\zeta_m) \rangle$, see [3, Theorem 4.3].

Let $c_m(x)$ denote the m th cyclotomic polynomial. To show that β_0 generates $D_{F/\mathbb{Q}}$, it suffices to show that it is an associate of $\langle c'_m(\zeta_m) \rangle$ in \mathcal{O}_F .

- (1) When m is an odd prime, then $c_m(x) = (x^m - 1)/(x - 1)$. Then

$$c'_m(x) = ((x - 1)mx^{m-1} - (x^m - 1))/(x - 1)^2.$$

So $c'_m(\zeta_m) = m\zeta_m^{-1}/(\zeta_m - 1)$. So β_0 from part (1) is an associate of $c'_m(\zeta_m)$ in \mathcal{O}_F .

- (2) If $m = 2^k$, then $c_m(x) = x^{2^{k-1}} + 1$. So $c'_m(\zeta_m) = 2^{k-1}\zeta_m^{2^{k-1}-1} = -2^{k-1}/\zeta_m$. So β_0 from part (2) is an associate of $c'_m(\zeta_m)$ in \mathcal{O}_F .
- (3) If $m = 3^k$, then $c_m(x) = x^{2 \cdot 3^{k-1}} + x^{3^{k-1}} + 1$. So $c'_m(\zeta_m) = 3^{k-1}\zeta_m^{3^{k-1}-1}\sqrt{-3}$. So β_0 from part (3) is an associate of $c'_m(\zeta_m)$ in \mathcal{O}_F .
- (4) In this case, $c_m(x) = (x^m - 1)(x - 1)/((x^p - 1)(x^q - 1))$. The element β_0 from part (4) is an associate of $c'_m(\zeta_m) = m\zeta_m^{-1}(\zeta_m - 1)/(\zeta_m^p - 1)(\zeta_m^q - 1)$.

□

3.6. Simple types. Let $F = \mathbb{Q}(\zeta_m)$ and recall that $\text{Gal}(F/\mathbb{Q}) \simeq (\mathbb{Z}/m\mathbb{Z})^*$. A CM-type Φ of F is simple, if it is not induced from any CM-field $F' \subset F$.

Let $H \subset \text{Gal}(F/\mathbb{Q})$ and let F^H be the fixed field of F under H . If F^H is a CM-field, then Φ is induced from F^H if and only if it is a union of cosets of H .

Lemma 3.8. *Suppose $m = 4$ or m is a Fermat prime or twice a Fermat prime. Then Φ is simple.*

We will use Lemma 3.8 for $m = 4$ and $m = 3, 5, 17$ and $m = 6, 10$.

Proof. For these values of m , the field F contains no proper non-trivial CM-fields. \square

Lemma 3.9. *Let $m > 3$ be prime. Let $\gamma = (m, 3, a)$ be a monodromy datum with inertia type $a = (1, a_2, a_3)$. Let Φ be the CM-type of F corresponding to a . Then Φ is simple if and only if $a \neq (1, x, x^2)$ for some $x \in (\mathbb{Z}/m\mathbb{Z})^*$ with order 3.*

Proof. This follows from [6, Theorem 2]. See also [9, Theorem 6.2]. \square

For example, when $m = 7$ and $a = (1, a_2, a_3)$, then Φ is simple unless $a = (1, 2, 4)$.

By a direct computation, one can check that Φ is always simple when $m = 25$ or $m = 27$.

4. PRINCIPAL POLARIZATIONS OF CM ABELIAN VARIETIES

Suppose L is a CM-field and L_0 is its maximal totally real subfield. In this section, we study principal polarizations on abelian varieties having complex multiplication by L .

Section 4.1 contains background about complex tori with complex multiplication. In Section 4.2, we study principal polarizations on CM-abelian varieties, following work of Van Wamelen. In Section 4.3, we prove a result about existence and uniqueness of principal polarizations for abelian varieties with simple CM-type when L has class number 1. In Section 4.4, we specialize to the case that L is a cyclotomic field with class number 1.

4.1. Complex tori. We review some complex multiplication theory, following Lang in [9, Chapter 1]. Let A be a complex torus such that $L \subset \text{End}(A) \otimes \mathbb{Q}$. We say that A is of type (L, Φ) if the complex representation of $\text{End}(A) \otimes \mathbb{Q}$ is isomorphic to $\sum_{\phi \in \Phi} \phi$. If, in addition, $\text{End}(A) \simeq \mathcal{O}_L$, we say A has type (\mathcal{O}_L, Φ) .

Theorem 4.1. [9, Chapter 1: Theorems 3.3, 3.5, 4.1, 4.2]

- (1) *If \mathfrak{a} is a lattice in L and Φ is a CM-type of L , then $\mathbb{C}^n / \Phi(\mathfrak{a})$ is a complex torus of type (L, Φ) .*
- (2) *If A is a complex torus of type (L, Φ) , then there exists a lattice \mathfrak{a} of L such that $A \simeq \mathbb{C}^n / \Phi(\mathfrak{a})$.*
- (3) *If Φ is a simple type and \mathfrak{a} is a fractional ideal of L , then $\text{End}(\mathbb{C}^n / \Phi(\mathfrak{a})) \simeq \mathcal{O}_L$.*
- (4) *If Φ is a simple type and $\mathfrak{a}, \mathfrak{b}$ are fractional ideals of L , then $\mathbb{C}^n / \Phi(\mathfrak{a}) \simeq \mathbb{C}^n / \Phi(\mathfrak{b})$ if and only if \mathfrak{a} and \mathfrak{b} are in the same ideal class.*

In particular, if (L, Φ) is a simple CM-type, then the set of isomorphism classes of complex tori of type (\mathcal{O}_L, Φ) is in bijection with the class group of L .

Furthermore, by [9, Chapter 1, Theorem 4.5], every (admissible, non-degenerate) Riemann form on $\mathbb{C}^n / \Phi(\mathfrak{a})$ is given by

$$(4.1) \quad \mathbb{E}(\Phi(x), \Phi(y)) = \text{tr}_{L/\mathbb{Q}}(\xi x \bar{y}), \text{ for } x, y \in L,$$

for some ξ such that $L = L_0(\xi)$, $\xi^2 \in L_0$ is totally negative, and $\text{Im}(\phi(\xi)) > 0$ for $\phi \in \Phi$.

4.2. Principal polarizations on CM-abelian varieties. In [19, page 310], Van Wamelen developed an algorithm to produce isomorphism classes of principally polarized abelian varieties of type (\mathcal{O}_L, Φ) based on the following result.

Theorem 4.2. (Van Wamelen) [19] *Let (L, Φ) be a CM-type.*

- (1) *(Theorem 4)² Writing $L = L_0(\sqrt{-c})$ for some $c \in \mathcal{O}_{L_0}$, then there exist a fractional ideal $\mathfrak{a} \subset L$ and an element $b \in \mathcal{O}_{L_0}$ such that $D_{L/\mathbb{Q}} \cdot \mathfrak{a}\bar{\mathfrak{a}} = \langle b\sqrt{-c} \rangle$.*

²We made a small adjustment to the notation to be consistent with the other parts of this theorem.

- (2) (Theorem 3) Let $\xi \in L$ be such that $L = L_0(\xi)$, $\xi^2 \in L_0$, and $D_{L/\mathbb{Q}} \cdot \mathfrak{a}\bar{\mathfrak{a}} = \langle \xi^{-1} \rangle$, for some fractional ideal \mathfrak{a} of L ; (for example, with the notation of part (1), $\xi = (b\sqrt{-c})^{-1}$). Define a Riemann form $\mathbb{E} : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}$ by

$$\mathbb{E}(z, w) = \sum_{i=1}^n \phi_i(\xi)(\bar{z}_i w_i - z_i \bar{w}_i),$$

for $z, w \in \mathbb{C}^n$. If $\text{Im}(\phi(\xi)) > 0$ for $\phi \in \Phi$, then \mathbb{E} defines a principal polarization on $\mathbb{C}^n / \Phi(\mathfrak{a})$ of type (\mathcal{O}_L, Φ) . Furthermore, if (L, Φ) is a simple CM-type, then all principal polarizations on $\mathbb{C}^n / \Phi(\mathfrak{a})$ of type (\mathcal{O}_L, Φ) are given by such a ξ .³

- (3) (Corollary 1) Two principal polarizations of the same CM-type on $\mathbb{C}^n / \Phi(\mathfrak{a})$ arising from ξ_1 and ξ_2 give isomorphic principally polarized abelian varieties if and only if there exists a unit $u \in \mathcal{O}_L^*$ such that $\xi_1 = u\bar{u}\xi_2$.

Corollary 4.3. Let Φ be a CM-type of L . An element $\xi = \beta^{-1}$ for some $\beta \in \mathcal{O}_L$ defines a principal polarization on $A_\Phi = \mathbb{C}^n / \Phi(\mathcal{O}_L)$ of CM-type (\mathcal{O}_L, Φ) if and only if

- (1) β generates the different $D_{L/\mathbb{Q}}$;
- (2) $\beta = -\bar{\beta}$;
- (3) $\text{Im}(\phi(\beta)) < 0$, for each $\phi \in \Phi$.

Two elements β, β' satisfying the above conditions yield isomorphic principally polarized abelian varieties if and only if there exists a unit $u \in \mathcal{U}_L$ such that $\beta = u\bar{u}\beta'$. If the CM-type Φ is simple, then all principal polarizations of A_Φ of CM-type (\mathcal{O}_L, Φ) arise this way.

Proof. The result follows from Theorem 4.2, replacing ξ, ξ_1, ξ_2 with $\beta^{-1}, \beta^{-1}, \beta'^{-1}$. \square

In Lemma 3.7, we determined an element $\beta_0 \in \mathcal{O}_F$ satisfying conditions (1) and (2) in Corollary 4.3 when $F = \mathbb{Q}(\zeta_m)$ for many values of m .

4.3. Existence and uniqueness of principal polarizations. In this section, we study principal polarizations on CM-abelian varieties. Under conditions on the class group and unit group of the field, we show such principal polarizations exist and are uniquely determined. Recall that L is a CM-field with maximal totally real subfield L_0 and $n = [L_0 : \mathbb{Q}]$.

Proposition 4.4. Suppose L_0 has units of independent signs and (L, Φ) is a CM-type.

- (1) Then there exists a principally polarized CM-abelian variety of type (\mathcal{O}_L, Φ) .
- (2) If (L, Φ) is simple, then any CM-abelian variety of type (\mathcal{O}_L, Φ) has at most one principal polarization, up to isomorphism.
- (3) If (L, Φ) is simple and L has class number 1, then there exists a unique principally polarized CM-abelian variety of type (\mathcal{O}_L, Φ) , up to isomorphism.

Proof. By Theorem 4.2(1), there exists a fractional ideal \mathfrak{a} in L and an element $\beta_0 \in \mathcal{O}_L$ satisfying the conditions: (1) β_0 generates $D_{L/\mathbb{Q}}\mathfrak{a}\bar{\mathfrak{a}}$ and (2) $\beta_0 = -\bar{\beta}_0$. For $\beta' \in \mathcal{O}_L$, then β' also satisfies conditions (1) and (2) if and only if $\beta' = u_0\beta_0$ for a totally real unit $u_0 \in \mathcal{U}_{L_0}$.

By Theorem 4.2(2), $\beta \in \mathcal{O}_L$ defines a principal polarization of type (\mathcal{O}_L, Φ) on $\mathbb{C}^n / \Phi(\mathfrak{a})$ if and only if it satisfies conditions (1) and (2), and also

- (3) $\text{Im}(\phi(\beta)) < 0$, for each $\phi \in \Phi$.

³Note that type (L, Φ) and type (\mathcal{O}_L, Φ) are equivalent in the last two statements.

Hence, to finish the first claim, it suffices to check that there exists $u_0 \in \mathcal{U}_{L_0}$ such that $\beta = u_0\beta_0$ satisfies condition (3), for any CM-type Φ of L . If L_0 has units of independent signs, then ρ_{L_0} is surjective; this shows that the unit u_0 described above exists.

By Theorem 4.2(2), if (L, Φ) is simple, all principal polarizations of type Φ on $\mathbb{C}^n/\Phi(\mathfrak{a})$ arise from some $\beta \in \mathcal{O}_L$ satisfying conditions (1)–(3). By Theorem 4.2(3), $\beta, \beta' \in \mathcal{O}_L$ satisfying conditions (1)–(3) define isomorphic principally polarized abelian varieties if and only if $\beta' = N(u)\beta$ for some unit $u \in \mathcal{U}_L$.

By definition, given a fractional ideal \mathfrak{a} of L , if $\beta \in \mathcal{O}_L$ satisfying conditions (1)–(3) exists, then $\beta' \in \mathcal{O}_L$ also satisfies conditions (1)–(3) if and only if $\beta' = u^+\beta$ for some totally positive unit $u^+ \in \mathcal{U}_{L_0}^+$.

Hence, to finish the second claim, it suffices to check that for any $u^+ \in \mathcal{U}_{L_0}^+$ there is $u \in \mathcal{U}_L$ such that $u^+ = N(u)$. Since L_0 has units of independent signs, this follows from Lemma 3.4. Finally, if L has class number 1, then by Theorem 4.2 and [19, Theorem 5], any CM-abelian variety of type (\mathcal{O}_L, Φ) is isomorphic to $A_\Phi = \mathbb{C}^n/\Phi(\mathcal{O}_L)$. \square

Proposition 4.5. *Suppose L_0 has units of almost independent signs and $Q(L) = 2$, where $Q(L)$ is defined in Definition 3.2. Suppose (L, Φ) is a simple CM-type.*

- (1) *The number of isomorphism classes of principal polarizations on a CM-abelian variety of type (\mathcal{O}_L, Φ) is at most one.*
- (2) *Suppose in addition that L has class number 1. If there exists a principally polarized CM-abelian variety of type (\mathcal{O}_L, Φ) , then it is unique up to isomorphism.*

Proof. The arguments in the proof of Proposition 4.4 still apply, after observing that $\mathcal{U}_{L_0}^+ = N(\mathcal{U}_L)$ when L_0 has units of almost independent signs and $Q(L) = 2$ by Lemma 3.4. \square

4.4. CM-types for cyclotomic fields. We consider the case when L is the cyclotomic field $F = \mathbb{Q}(\zeta_m)$. By Section 3.2, the next result is a special case of Propositions 4.4 and 4.5.

Corollary 4.6. *Let m such that $F = \mathbb{Q}(\zeta_m)$ has class number 1. Let Φ be a CM-type of F .*

- (1) *If F_0 has narrow class number 1, then there exists a principally polarized CM-abelian variety of type (\mathcal{O}_F, Φ) . Furthermore, if (F, Φ) is simple, then it is unique up to isomorphism.*
- (2) *Suppose F_0 has narrow class number 2 and Φ is simple. If there exists a principally polarized CM-abelian variety of type (\mathcal{O}_F, Φ) , then it is unique up to isomorphism.*

In the rest of this section, we explicitly describe the data of the principal polarization.

Corollary 4.7. *Let $m, F = \mathbb{Q}(\zeta_m)$, and β_0 be as in Lemma 3.7. Let Φ be a CM-type of F . Consider conditions (1)–(3) in Corollary 4.3.*

- (1) *If $\Phi' = i\Phi$ and β satisfies conditions (1)–(3) for Φ , then $\beta' = \sigma_{i-1}(\beta)$ satisfies conditions (1)–(3) for Φ' .*
- (2) *Suppose F_0 has units of independent signs. Then, for any CM-type Φ of F , there exists a unit $u_0 \in \mathcal{U}_{F_0}$ such that $\beta = u_0\beta_0$ satisfies conditions (1)–(3) for Φ . Furthermore, β is the unique element satisfying conditions (1)–(3) for Φ , up to multiplication by an element of $N_{F/F_0}(\mathcal{U}_F)$.*

The condition that F_0 has units of independent signs is satisfied if the narrow class number of F_0 is odd [2, bottom of page 58].

Proof. (1) This follows from Lemma 2.1 and Lemma 2.7.

- (2) By Lemma 3.7, β_0 satisfies conditions (1) and (2) of Corollary 4.3. Since F_0 has units of independent signs, there exists $u_0 \in \mathcal{U}_{F_0}$ such that $\beta = u_0\beta_0$ satisfies condition (3). Since $u_0 \in \mathcal{U}_{F_0}$ is a real unit, β also satisfies conditions (1) and (2) of Corollary 4.3. Note that β is unique up to multiplication by a totally positive unit of F_0 . By Lemma 3.4, $\mathcal{U}_{F_0}^+ = N(\mathcal{U}_F)$, proving the uniqueness statement. \square

See Example 6.3 (resp. 6.7) for an example of Corollary 4.7 when $m = 5$ (resp. $m = 7$). As another example, when $m = 8$, then $\beta_0 = -4i$ from Lemma 3.7: if $\mathfrak{f} = (0, 1, 0, 1)$, set $u_0 = -1$ and $\beta = 4i$; if $\mathfrak{f} = (1, 1, 0, 0)$, set $u_0 = \sqrt{2} - 1$ and $\beta = 4(1 - \sqrt{2})i$.

5. SHIMURA DATA

Consider a monodromy datum $\gamma = (m, N, a)$ as in Section 2.1. Recall from Section 2.1 that Z_γ is the closure in \mathcal{A}_g of the image of the Hurwitz space of μ_m -cyclic covers of \mathbb{P}^1 with monodromy datum γ . A point P in Z_γ represents the Jacobian of a curve C_P (either smooth or of compact type), for which there is an admissible μ_m -cover $\psi_P : C_P \rightarrow T$ with monodromy datum γ , where T is a tree of projective lines. Recall that S_γ is the connected component of $\text{Sh}(\gamma)$ containing Z_γ . In this section, under certain assumptions on m , we provide a method to determine the Shimura datum of S_γ .

5.1. Distinguished points. In this section, we prove that there exists a *distinguished point* P in the family Z_γ such that we can compute the Shimura datum of the Jacobian of C_P .

Definition 5.1. A point P in Z_γ is a *distinguished point* if the Jacobian of C_P is a principally polarized abelian variety with complex multiplication by a maximal order in a CM-field, or the direct sum of such together with the product polarization.

Proposition 5.2. *Let m be an odd prime. Let $\gamma = (m, N, a)$ be a monodromy datum. Then Z_γ has a distinguished point P . More specifically, for $r = N - 2$:*

- (1) *In the family of μ_m -covers with monodromy datum γ , there is a point which represents an admissible μ_m -cover $\psi : C_P \rightarrow T$, where T is a tree of r projective lines and C_P is a curve of compact type, with r irreducible components $\mathcal{C}_1, \dots, \mathcal{C}_r$, each of which is a curve of genus $(m - 1)/2$ admitting a μ_m -cover of \mathbb{P}^1 branched at 3 points.*
- (2) *The Jacobian of C_P is of the form $A \simeq \bigoplus_{j=1}^r A_j$, where each A_j is a principally polarized abelian variety of dimension $(m - 1)/2$ having complex multiplication by $\mathcal{O}_F = \mathbb{Z}[\zeta_m]$, together with the product polarization.*

Proof. The fact that Z_γ has a distinguished point P is immediate from part (2), which we will show follows from part (1).

- (1) Let $a = (a(1), \dots, a(N))$ be the inertia type. If $N \geq 4$ and m is an odd prime, then a has the following property: there is a pair (i, j) with $1 \leq i < j \leq N$, such that $a(i) + a(j) \not\equiv 0 \pmod{m}$. Without loss of generality, we can suppose $i = 1$ and $j = 2$.

When $N \geq 4$, then Z_γ^0 is affine. Consider a family of covers, where the first branch point b_1 approaches the second branch point b_2 . When $b_1 = b_2$, the curve becomes singular and its normalization is a μ_m -cover of the join of two projective lines. Because of the condition $a(i) + a(j) \not\equiv 0 \pmod{m}$, this is a degeneration of

compact type as described in [12, Remark 5.2]. It is the admissible join of two μ_m -covers with these monodromy data:

$$\begin{aligned}\gamma_1 &= (m, 3, (a(1), a(2), -(a(1) + a(2)))); \\ \gamma_2 &= (m, N - 1, (a(1) + a(2), a(3), \dots, a(N))).\end{aligned}$$

By induction on N , we see that the universal family degenerates completely to an admissible μ_m -cover $\psi : \mathcal{C} \rightarrow T$, where T is a tree of r projective lines and the restriction of ψ above each component of T is branched at 3 points. By (2.1), each irreducible component of \mathcal{C} has genus $(m - 1)/2$.

- (2) Choose a labeling C_1, \dots, C_r of the irreducible components of \mathcal{C} . Let $A = \text{Jac}(\mathcal{C})$ and $A_j = \text{Jac}(C_j)$. Then A_j is a principally polarized abelian variety of dimension $(m - 1)/2$. By [1, Section 9.2, Equation 8], $A \simeq \bigoplus_{j=1}^r A_j$, and the principal polarization on A decomposes as the product polarization.

Consider the CM-field $F = \mathbb{Q}(\zeta_m)$. Then $\mathcal{O}_F = \mathbb{Z}[\zeta_m] \subset \text{End}(A_j)$ since μ_m acts on C_j for $1 \leq j \leq r$. Since $\deg(F/\mathbb{Q}) = 2 \cdot \dim(A_j)$, the abelian variety A_j has complex multiplication by \mathcal{O}_F for $1 \leq j \leq r$.

□

5.2. Shimura datum for Hurwitz spaces. Let $P \in Z_\gamma$ be a distinguished point as described in Proposition 5.2. When $F = \mathbb{Q}(\zeta_m)$ has class number 1, we determine the Shimura datum for S_γ . Then strategy is to explicitly compute the principally polarized abelian variety $\text{Jac}(C_P)$ represented by P and then the F -vector space $V = H^1(C_P(\mathbb{C}), \mathbb{Q})$, the lattice $\Lambda = H^1(C_P(\mathbb{C}), \mathbb{Z})$, and the Hermitian matrix.

Definition 5.3. Let $P \in Z_\gamma$ be a distinguished point as described in Proposition 5.2. Suppose $\mathbb{Q}(\zeta_m + \zeta_m^{-1})$ has units of independent signs. (This is guaranteed by Lemma 3.6 when m is an odd prime such that $F = \mathbb{Q}(\zeta_m)$ has class number 1.)

For $1 \leq j \leq r$, define β_j and $\xi_j = \beta_j^{-1}$ as follows. The inertia type of the μ_m -cover $C_j \rightarrow \mathbb{P}^1$ determines the signature \mathfrak{f}_j by (2.2), which determines the CM-type Φ_j as in (2.6). Let β_0 be as in Lemma 3.7. By Corollary 4.7, there exists $u_j \in \mathcal{U}_{F_0}$ such that $\beta_j := u_j \beta_0$ satisfies conditions (1)–(3) of Corollary 4.3 for Φ_j . Furthermore, β_j is the unique element satisfying conditions (1)–(3) for Φ_j , up to multiplication by an element of $N_{F/F_0}(\mathcal{U}_F)$.

Remark 5.4. We note that the CM-type (F, Φ_j) of A_j may not be simple. For example, for the Moonen special family $M[17]$ with $m = 7$, then (F, Φ_2) is not simple, see Section 6.3.

Theorem 5.5. Let m be an odd prime. Suppose $F = \mathbb{Q}(\zeta_m)$ has class number 1. Let $\gamma = (m, N, a)$ be a monodromy datum with $N \geq 4$.

Let $P \in Z_\gamma$ be a distinguished point as described in Proposition 5.2. Let $r = N - 2$. For $1 \leq j \leq r$: let A_j be the abelian variety with CM-type (\mathcal{O}_F, Φ_j) from Proposition 5.2; let $\xi_j = \beta_j^{-1}$ be as in Definition 5.3. Then the integral PEL datum for S_γ is given as:

- the F -vector space $V = F^r$, together with the standard \mathcal{O}_F -lattice $\Lambda = (\mathcal{O}_F)^r \subseteq V$;
- the Hermitian form $H_B = \langle \cdot, \cdot \rangle$ on V , which takes integral values on Λ , defined by

$$\langle x, y \rangle = \text{tr}_{F/\mathbb{Q}}(xB\bar{y}^T) \text{ for } B = \text{diag}[\xi_1, \dots, \xi_r] \in \text{GL}_r(F) = \text{GL}(V).$$

It is straight-forward to compute B from γ ; we provide many examples in Section 6.

Proof. By Lemma 3.6, if m is prime and $F = \mathbb{Q}(\zeta_m)$ has class number 1, then $F_0 = \mathbb{Q}(\zeta_m + \zeta_m^{-1})$ has narrow class number 1 and has units of independent signs. By hypothesis, for $1 \leq j \leq r$, the abelian variety A_j has dimension $g = (m-1)/2$; it has complex multiplication by $\mathcal{O}_F = \mathbb{Z}[\zeta_m]$ and has CM-type (\mathcal{O}_F, Φ_j) . Furthermore, β_j defines a principal polarization on A_j ; the corresponding Hermitian form is given by $\mathbb{E}(\Phi(x), \Phi(y)) = \text{tr}_{F/\mathbb{Q}}(\zeta_j x \bar{y})$ for $x, y \in L$ by (4.1).

Since $F = \mathbb{Q}(\zeta_m)$ has class number 1, there is a unique complex torus of CM-type (\mathcal{O}_F, Φ_j) up to isomorphism. Namely, $A_j \simeq \mathbb{C}^g / \Phi(\mathcal{O}_F)$. The product polarization on $\text{Jac}(C_P) \simeq \bigoplus_{j=1}^r A_j$ defines an integral Hermitian space of the prescribed signature on $V = H^1(C_P(\mathbb{C}), \mathbb{Q})$. By [18, Appendix, Proposition 8], there is a unique Hermitian form of given signature on $\text{Jac}(C_P)$. Thus the integral PEL datum for $\text{Jac}(C_P)$ is as given in the statement. The result follows since the integral PEL datum can be determined at a point of Z_γ , see [14, Sections 2.8 and 2.9, also 3.2]. \square

Remark 5.6. The values of m for which Theorem 5.5 applies are $m = 3, 5, 7, 11, 13, 17, 19$.

For $m = 3, 5, 11, 17$, it is not necessary to refer to [18, Appendix, Proposition 8]; The reason is that (F, Φ_j) is automatically simple for $1 \leq j \leq r$ by Lemmas 3.8 and 3.9. Thus by Corollary 4.6(1), there is a unique principally polarized CM-abelian variety of type (\mathcal{O}_F, Φ_j) up to isomorphism. In Section 6.4, we give an alternative approach for $m = 7$ that relies on Corollary 4.6(2) and does not refer to [18, Appendix, Proposition 8].

The strategy of Theorem 5.5 sometimes applies when m is composite; see Section 7.

6. APPLICATIONS FOR m PRIME

Theorem 5.5 gives a method to determine the integral PEL datum for S_γ for all monodromy data $\gamma = (m, N, a)$ when $m = 3, 5, 7, 11, 13, 17, 19$. In this section, we give examples of this. Specifically:

- in Section 6.1, when $m = 3$, we explicitly determine the Hermitian form for all γ ;
- in Sections 6.2-6.3, when $m = 5, 7$, we give explicit examples for all equivalence classes of γ when $N = 4$.
- in Section 6.4, we follow an alternative approach for $m = 7$ which avoids the possible occurrence of non-simple CM-types at the distinguished point.

In particular, we determine the integral PEL datum for the 6 Moonen special families for which the degree m is an odd prime; these are denoted by $M[n]$ as in [15, Table 1].

In each section below, let $F = \mathbb{Q}(\zeta_m)$. For $i = 1, 2$, let Φ_i be the CM-type of F determined by the signature \mathfrak{f}_i , as in (2.6). Given the F -vector space $V = F^r$, together with the standard \mathcal{O}_F -lattice $\Lambda = (\mathcal{O}_F)^r \subseteq V$, let $H_B = \langle \cdot, \cdot \rangle$ denote the Hermitian form on V defined by

$$\langle x, y \rangle = \text{tr}_{F/\mathbb{Q}}(xB\bar{y}^T) \text{ for } B = \text{diag}[\zeta_1, \dots, \zeta_r] \in \text{GL}_r(F) = \text{GL}(V).$$

6.1. Integral PEL datum for all families when $m = 3$. We find the integral PEL data for all families with $m = 3$.

Example 6.1. When $m = 3$, then $\beta_0 = 3/(\zeta_3^2 - \zeta_3) = \sqrt{-3}$ by Lemma 3.7.

For $\mathfrak{f}_1 = (1, 0)$, set $u_1 = -1$ and $b_1 = u_1\beta_0$ so that $b_1 = -\sqrt{-3}$. Then $\text{Im}(\sigma_1(b_1)) < 0$.

For $\mathfrak{f}_2 = (0, 1)$, set $u_2 = 1$ and $b_2 = u_2\beta_0$ so that $b_2 = \sqrt{-3}$. Then $\text{Im}(\sigma_2(b_2)) < 0$.

The CM-type Φ_i is simple by Lemma 3.8 and b_i satisfies Lemma 4.7 with respect to Φ_i .

For $m = 3$, there are simple formulas relating the signature \mathfrak{f} and the inertia type a . If a has d_1 entries of 1 and d_2 entries of 2, then $\mathfrak{f} = (f_1, f_2)$ with $f_1 = (2d_1 + d_2 - 3)/3$ and $f_2 = (d_1 + 2d_2 - 3)/3$. Then $d_1 = 2f_1 - f_2 + 1$ and $d_2 = 2f_2 - f_1 + 1$. One can check that $0 \leq \max(f_1, f_2) \leq 2\min(f_1, f_2) + 1$.

Corollary 6.2. *Let $m = 3$ and $N \geq 4$. Let $\gamma = (3, N, a)$ be a monodromy datum with signature (f_1, f_2) . Let $g = f_1 + f_2$. Let S_γ be the component of the Shimura variety containing Z_γ .*

Let $F = \mathbb{Q}(\zeta_3)$ and $\xi = -1/\sqrt{-3}$. Then the integral PEL datum of S_γ has lattice $(\mathcal{O}_F)^g$ with Hermitian form H_B where $B \in \mathrm{GL}_g(\mathcal{O}_F)$ is diagonal with f_1 entries of ξ and f_2 entries of $-\xi$.

Proof. Consider the admissible μ_3 -cover $\varphi : \mathcal{C}_P \rightarrow T$ represented by the distinguished point from Proposition 5.2. Here T is a tree of $r = f_1 + f_2$ projective lines. Above f_1 (resp. f_2) components of T , the restriction of φ is a μ_3 -cover branched at 3 points with signature $\mathfrak{f}_1 = (1, 0)$ (resp. $\mathfrak{f}_2 = (0, 1)$). The result then follows from Theorem 5.5. \square

In particular, Corollary 6.2 includes the three special families $M[3]$, $M[6]$, $M[10]$.

M	(m, N, a)	B
$M[3]$	$(3, 4, (1, 1, 2, 2))$	$\mathrm{diag}[\xi, -\xi]$
$M[6]$	$(3, 5, (1, 1, 1, 1, 2))$	$\mathrm{diag}[\xi, \xi, -\xi]$
$M[10]$	$(3, 6, (1, 1, 1, 1, 1, 1))$	$\mathrm{diag}[\xi, \xi, \xi, -\xi]$

The cases $M[6] : \xi[1, 1, -1]$ and $M[10] : \xi[1, 1, 1, -1]$ match the table on [18, page 1].

6.2. Integral PEL datum when $m = 5$. We illustrate Theorem 5.5 for several well-chosen examples when $m = 5$, including all families branched at $N = 4$ points up to equivalence and one family branched at $N = 5$ points. This includes the two special Moonen families when $m = 5$, namely $M[11]$ and $M[16]$. A similar result can be obtained for any monodromy datum γ with $N > 4$ by an inductive process.

Example 6.3. The following table summarizes the data for CM-types when $m = 5$.

a	\mathfrak{f}	Φ	β
$(4, 3, 3)$	$(0, 1, 0, 1)$	$\{2, 4\}$	$\beta_1 = 5/(\zeta_5^3 - \zeta_5^2)$
$(3, 1, 1)$	$(1, 1, 0, 0)$	$\{1, 2\}$	$\beta_2 = 5/(\zeta_5 - \zeta_5^4)$
$(1, 2, 2)$	$(1, 0, 1, 0)$	$\{1, 3\}$	$\beta_3 = -\beta_1$
$(2, 4, 4)$	$(0, 0, 1, 1)$	$\{3, 4\}$	$\beta_4 = -\beta_2$

In the i th line of the table, the CM-type Φ_i is simple by Lemma 3.8 and β_i satisfies Corollary 4.7 with respect to Φ_i . The automorphism σ_3 permutes the rows via the cycle $(1, 2, 3, 4)$ and its inverse σ_2 permutes the 4th column via $\beta_1 \rightarrow \beta_2 \rightarrow -\beta_1 \rightarrow -\beta_2 \rightarrow \beta_1$.

Proof. When $m = 5$, then $\beta_0 = 5/(\zeta_5^3 - \zeta_5^2)$ by Lemma 3.7. For $\mathfrak{f}_1 = (0, 1, 0, 1)$, set $u_1 = 1$ and $\beta_1 = u_1\beta_0$. We compute that $\mathrm{Im}(\sigma_j(\beta_1)) < 0$ for $j = 2, 4$. For $\mathfrak{f}_2 = (1, 1, 0, 0)$, set $u_2 = (\zeta_5^3 - \zeta_5^4)/(\zeta_5 - \zeta_5^4)$ and $\beta_2 = u_2\beta_0$. We compute that $\mathrm{Im}(\sigma_j(\beta_2)) < 0$ for $j = 1, 2$. The signature $\mathfrak{f}_3 = (1, 0, 1, 0)$ (resp. $\mathfrak{f}_4 = (0, 0, 1, 1)$) is the complex conjugate of \mathfrak{f}_1 (resp. \mathfrak{f}_2), which negates the value of β . \square

Corollary 6.4. *Let $m = 5$ and $F = \mathbb{Q}(\zeta_5)$. Every family of μ_5 -covers of \mathbb{P}^1 with $N = 4$ is equivalent to either (i), (ii) or $M[11]$ in the table below. Recall $\beta_1, \beta_2 \in \mathcal{O}_F$ from Example 6.3 and*

let $\xi_i = \beta_i^{-1}$. For the monodromy data $\gamma = (5, N, a)$ and $r = N - 2$ as below, the integral PEL datum of S_γ has lattice $(\mathcal{O}_F)^r$ with Hermitian form H_B where $B \in \mathrm{GL}_r(\mathcal{O}_F)$ is as below.

M	$(5, N, a)$	B
(i)	$(5, 4, (1, 1, 4, 4))$	$\mathrm{diag}[\tilde{\zeta}_2, -\tilde{\zeta}_2]$
(ii)	$(5, 4, (1, 2, 3, 4))$	$\mathrm{diag}[-\tilde{\xi}_1, \tilde{\xi}_1]$
$M[11]$	$(5, 4, (1, 3, 3, 3))$	$\mathrm{diag}[\tilde{\xi}_1, \tilde{\xi}_2]$
$M[16]$	$(5, 5, (2, 2, 2, 2, 2))$	$\mathrm{diag}[-\tilde{\xi}_1, -\tilde{\xi}_2, -\tilde{\xi}_1]$

Proof. Suppose $m = 5$ and $N = 4$ and let a be the inertia type of γ . If three of the values of a are the same, the family is equivalent to the one with $a = (1, 3, 3, 3)$, which is $M[11]$. If two of the values of a are the same, the family is equivalent to (i). If all values of a are distinct, the family is equivalent to (ii).

By Theorem 5.5, it suffices to find the CM-type (F, Φ_i) for the abelian varieties A_i in the decomposition of the Jacobian of \mathcal{C}_p . We refer to [12, Remark 5.2, Lemma 6.4] for information about the admissible degeneration, given in short-hand by: (i) $(1, 1, 3) + (2, 4, 4)$; (ii) $(1, 2, 2) + (3, 3, 4)$; $M[11]$ $(1, 3, 1) + (4, 3, 3)$; and $M[16]$ $(2, 2, 1) + (4, 2, 4) + (1, 2, 2)$. Using the table in Example 6.3, we find the entries of the diagonal of B . \square

Remark 6.5. The family (ii), with monodromy datum $\gamma = (5, 4, (1, 2, 3, 4))$, has a second degeneration of the form $(1, 3, 1) + (4, 2, 4)$, whose Hermitian form has matrix $B' = \mathrm{diag}[\tilde{\zeta}_2, -\tilde{\zeta}_2]$. In this case, there are down-to-earth ways to explain why the Hermitian forms determined by B' and $B = \mathrm{diag}[-\tilde{\xi}_1, \tilde{\xi}_1]$ are isomorphic.

First, by Lemma 2.1, the automorphism σ_2 takes the inertia types in the first degeneration to those in the second by multiplying the entries by 3. By Corollary 4.7(1), the action on the entries of B is via $\sigma_{2^{-1}} = \sigma_3$ and

$$\sigma_3(B) = \mathrm{diag}[\sigma_3(-\tilde{\xi}_1), \sigma_3(\tilde{\xi}_1)] = \mathrm{diag}[\tilde{\zeta}_2, -\tilde{\zeta}_2] = B'.$$

Second, for family (ii), S_γ has signature type $(1, 1, 1, 1)$; hence its reflex field is \mathbb{Q} (which is smaller than $F_0 \subset \mathbb{R}$). The matrices B and B' are conjugate under the action of $\sigma_3 \in \mathrm{Gal}(F_0/\mathbb{Q})$ and correspond to the two choices of a \mathbb{Q} -linear embedding $F_0 \hookrightarrow \mathbb{R}$.

Remark 6.6. To compare with Shimura's work, write $w = \zeta_5 + \zeta_5^4$. Then $w^2 + w - 1 = 0$. So $w = (-1 + \sqrt{5})/2$. Then $\tilde{\zeta}_2/\tilde{\xi}_1 = -w - 1 = -(1 + \sqrt{5})/2$ so $\tilde{\xi}_1/\tilde{\zeta}_2 = (1 - \sqrt{5})/2$.

Consider the family $\gamma' = (5, 4, (1, 1, 1, 1, 1))$. A careful look at [18, Section 5] shows that Shimura replaced ζ_5 by ζ_5^3 in his computation for this family. This has the effect of switching to the family $M[16]$ with $\gamma = (5, 4, (2, 2, 2, 2, 2))$; indeed, Shimura computes that the signature is $(2, 0, 3, 1)$. By line 4 of the table in Corollary 6.4, the family γ has $B = -\tilde{\xi}_1[1, 1, \tilde{\zeta}_2/\tilde{\xi}_1] = -\tilde{\xi}_1[1, 1, -(1 + \sqrt{5})/2]$. This does not exactly match what is written in line 5 of the table on [18, page 1], namely $[1, 1, \tilde{\xi}_1/\tilde{\zeta}_2] = [1, 1, (1 - \sqrt{5})/2]$, but it has the same sign signature and thus yields an isomorphic Hermitian form.

Consider the family $\gamma' = (5, 4, (2, 1, 1, 1, 1))$. The details for this family are not included in [18] but it appears that Shimura replaced ζ_5 by ζ_5^3 in his computation for this family also. This has the effect of switching to the family $M[11]$ with $\gamma = (5, 4, (1, 3, 3, 3))$. By line 3 of the table in Corollary 6.4, the family γ has $B = \tilde{\zeta}_2[1, \tilde{\xi}_1/\tilde{\zeta}_2] = \tilde{\zeta}_2[1, (1 - \sqrt{5})/2]$. This matches what is written in line 4 of the table on [18, page 1].

6.3. Integral PEL datum when $m = 7$. We illustrate Theorem 5.5 for several well-chosen examples when $m = 7$, including all families branched at $N = 4$ points up to equivalence. This includes the special Moonen family M [17]. A similar result can be obtained for any monodromy datum γ with $N > 4$ by an inductive process.

Example 6.7. The following table summarizes the cases when $m = 7$.

a	\mathfrak{f}	Φ	β
(1, 1, 5)	(1, 1, 1, 0, 0, 0)	{1, 2, 3}	$\beta_1 = 7/(\zeta_7 - \zeta_7^6)$
(3, 3, 1)	(1, 0, 1, 0, 1, 0)	{1, 3, 5}	$\beta_2 = 7/(\zeta_7^3 - \zeta_7^4)$
(2, 2, 3)	(1, 0, 0, 1, 1, 0)	{1, 4, 5}	$\beta_3 = 7/(\zeta_7^2 - \zeta_7^5)$
(6, 6, 2)	(0, 0, 0, 1, 1, 1)	{4, 5, 6}	$\beta_4 = -\beta_1$
(4, 4, 6)	(0, 1, 0, 1, 0, 1)	{2, 4, 6}	$\beta_5 = -\beta_2$
(5, 5, 4)	(0, 1, 1, 0, 0, 1)	{2, 3, 6}	$\beta_6 = -\beta_3$
(1, 2, 4)	(1, 1, 0, 1, 0, 0)	{1, 2, 4}	$\beta = -\frac{7(\zeta_7^3 - \zeta_7^4)}{(\zeta_7 - \zeta_7^6)(\zeta_7^2 - \zeta_7^5)}$
(3, 1, 5)	(0, 0, 1, 0, 1, 1)	{3, 5, 6}	$\beta' = -\beta$

Lemma 6.8. *In the i th line of the table in Example 6.7, for $1 \leq i \leq 6$, the CM-type Φ_i is simple, and the element β_i satisfies Corollary 4.7 with respect to Φ_i . The generator σ_3 of the Galois group $\text{Gal}(\mathbb{Q}(\zeta_7)/\mathbb{Q})$ permutes these via $\sigma_3^{i-1}(\beta_1) = \beta_i$, for $i = 1, \dots, 6$.*

In the last two lines of the table, the element β (resp. β') satisfies Corollary 4.7 with respect to the CM-type $\Phi = \{1, 2, 4\}$ (resp. $\Phi' = \{3, 5, 6\}$). The CM types Φ, Φ' are not simple.

Proof. When $m = 7$, then $\beta_0 = 7/(\zeta_7^4 - \zeta_7^3)$ by Lemma 3.7.

For $a_1 = (1, 1, 5)$ and $\mathfrak{f}_1 = (1, 1, 1, 0, 0, 0)$, set $u_1 = (\zeta_7^4 - \zeta_7^3)/(\zeta_7 - \zeta_7^6)$ and $\beta_1 = u_1\beta_0$. We compute that $\text{Im}(\sigma_j(\beta_1)) < 0$ for $j = 1, 2, 3$.

Let $\sigma = \sigma_5$, which is a generator of $\text{Gal}(\mathbb{Q}(\zeta_7)/\mathbb{Q})$ and has inverse σ_3 . Then, by Lemmas 2.1 and 2.7, for $1 \leq i \leq 6$, the action of σ_5^{i-1} changes the inertia type to $a_i := (3^{i-1}) \cdot a_1$ and the CM-type to $\Phi_i := 5^{i-1} \cdot \Phi$, which also determines the signature \mathfrak{f}_i . Also, the element $\beta_i = \sigma_5^{i-1}(\beta_1)$ satisfies Corollary 4.7 with respect to Φ_i .

For $a = (1, 2, 4)$ and $\mathfrak{f} = (1, 1, 0, 1, 0, 0)$, set

$$u = -\frac{(\zeta_7^4 - \zeta_7^3)^2}{(\zeta_7 - \zeta_7^6)(\zeta_7^2 - \zeta_7^5)}, \text{ and } \beta = u\beta_0 = -\frac{7(\zeta_7^3 - \zeta_7^4)}{(\zeta_7 - \zeta_7^6)(\zeta_7^2 - \zeta_7^5)}.$$

Then $\text{Im}(\sigma_j(\beta_2)) < 0$ for $j = 1, 2, 4$. The last line follows from Lemma 2.1 by applying complex conjugation to the previous line.

By Lemma 3.9, Φ_i is simple unless it is $\{1, 2, 4\}$ or $\{3, 5, 6\}$. □

For $m = 7$ and $N = 4$, every family is equivalent to one in the next result.

Corollary 6.9. *Let $m = 7$ and $F = \mathbb{Q}(\zeta_7)$. Recall $\beta_1, \beta_2, \beta_3, \beta \in \mathcal{O}_F$ from Example 6.7. Let $\xi_i = \beta_i^{-1}$ for $i = 1, 2, 3$ and $\xi = \beta^{-1}$. For the monodromy data $\gamma = (7, 4, a)$ as below, the integral*

PEL datum of S_γ has lattice $(\mathcal{O}_F)^2$ with Hermitian form H_B where $B \in \mathrm{GL}_2(\mathcal{O}_F)$ is as below.

M	a	B
(i)	(1, 1, 2, 3)	$\mathrm{diag}[\zeta_1, \zeta_3]$
(ii)	(1, 1, 6, 6)	$\mathrm{diag}[\zeta_1, -\zeta_1]$
(iii)	(1, 2, 5, 6)	$\mathrm{diag}[\zeta_1, -\zeta_1]$
$M[17]$	(2, 4, 4, 4)	$\mathrm{diag}[-\zeta_2, \zeta]$

Proof. We refer to [12, Remark 5.2, Lemma 6.4] for information about the admissible degeneration of each family, given in short-hand by: (i) $(1, 1, 5) + (2, 2, 3)$; (ii) $(1, 1, 5) + (2, 6, 6)$; (iii) $(1, 5, 1) + (6, 2, 6)$; and $M[17]$ $(4, 4, 6) + (1, 4, 2)$. For each family, consider the CM-types (F, Φ_1) and (F, Φ_2) for the abelian varieties A_i in the decomposition of the Jacobian at the distinguished point arising from the degeneration. We compute the associated integral PEL data by Theorem 5.5, using the table in Example 6.7. \square

Remark 6.10. In cases (i), (ii), and (iii), both (F, Φ_1) and (F, Φ_2) are simple. For $M[17]$, the CM-type (F, Φ_2) is not simple, but this is not a concern by [18, Appendix, Proposition 8] or Section 6.4. For $M[17]$, we note that $B = \zeta[1, -\zeta_2/\zeta]$ where $-\zeta_2/\zeta = \zeta_7 + \zeta_7^6 - 1$. In line 6 of the table on [18, page 1], for the related family $\gamma = (7, 4, (4, 1, 1, 1))$, Shimura writes the Hermitian form as H_B with $B = \zeta[1, -\sin(3\pi/7)/\sin(2\pi/7)]$.

6.4. Another approach for $M[17]$. In this section, we compute the integral PEL datum for the family $M[17]$ using another approach. This approach utilizes a different kind of distinguished point in the family, namely one that represents a curve with extra automorphisms. In this section, we see that the Jacobian of this curve has complex multiplication by a larger field and that its CM-type is simple.

In fact, we proved that every positive-dimensional family of μ_7 -covers of \mathbb{P}^1 has a distinguished point representing a product of principally polarized abelian varieties, each of which has complex multiplication with a CM-type that is simple. In this way, one can avoid the use of [18, Appendix, Proposition 8] when $m = 7$. We omit the details of this.

We start by finding another distinguished point in the family Z_γ , under certain restrictive conditions on γ . Similarly to Notation 3.5, for $0 \leq n \leq 3m$ with $\mathrm{gcd}(n, 3m) = 1$, let σ_n be the embedding $\mathbb{Q}(\zeta_{3m}) \hookrightarrow \mathbb{C}$ determined by $\sigma_n(\zeta_{3m}) = \zeta_{3m}^n$.

Proposition 6.11. *Let $\gamma = (m, 4, a)$ where $m > 3$ is prime and $a = (1, 1, 1, m - 3)$. Then Z_γ has a distinguished point Q . More specifically: in the family of μ_m -covers with monodromy datum γ , there is a point which represents a μ_m -cover $\psi : C_Q \rightarrow \mathbb{P}^1$, where C_Q is a curve of genus $m - 1$ having an automorphism of order 3.*

Furthermore, the Jacobian of C_Q has complex multiplication by $(\mathbb{Z}[\zeta_{3m}], \Phi_Q)$, where, for any $0 < n < 3m$ with $(n, 3m) = 1$, the embedding σ_n is in Φ_Q if and only if

$$\begin{aligned} n &\in [0, 2m/3] \cup [m, 5m/3] \cup [2m, 8m/3] \text{ if } n \equiv 1 \pmod{3}, \text{ or} \\ n &\in [0, m/3] \cup [m, 4m/3] \cup [2m, 7m/3] \text{ if } n \equiv 2 \pmod{3}. \end{aligned}$$

It is possible to generalize Proposition 6.11, by replacing 3 by an odd prime ℓ relatively prime to m and letting $N = \ell + 1$ and $a = (1, \dots, 1, m - \ell)$. We omit this generalization.

Proof of Proposition 6.11. Let C_Q be the smooth projective curve with equation $y^m = x^3 - 1$. It admits a μ_m -cover to \mathbb{P}^1 branched at $1, \zeta_3, \zeta_3^2, \infty$ with inertia type $a = (1, 1, 1, m - 3)$. The genus of C_Q is $m - 1$ by (2.1). Also C_Q has an automorphism $(x, y) \mapsto (\zeta_3 x, y)$ of order 3.

The Jacobian $J_Q = \text{Jac}(C_Q)$ is a principally polarized abelian variety having dimension $m - 1$. Let $F = \mathbb{Q}(\zeta_m)$, and $L = \mathbb{Q}(\zeta_{3m})$. The field L is a CM-field of degree $2 \cdot \dim(J_Q)$. Then, the inclusion $\mathcal{O}_F \subset \text{End}(J_Q)$ extends to an inclusion $\mathcal{O}_L = \mathcal{O}_F[\zeta_3] \subset \text{End}(J_Q)$. Thus J_Q has complex multiplication by $\mathcal{O}_L = \mathbb{Z}[\zeta_{3m}]$.

Consider the morphism $\phi : C_Q \rightarrow \mathbb{P}^1$, taking $(x, y) \mapsto x^3$. Then ϕ is a μ_{3m} -cover of \mathbb{P}^1 , branched at $0, 1, \infty$. We compute the CM-type of J_Q by finding first the inertia type and then the signature type of ϕ . Let b_0 (resp. b_1 , resp. b_∞) denote the element of $\mathbb{Z}/3m\mathbb{Z}$ that determines the canonical generator of inertia of ϕ above 0 , (resp. 1 , resp. ∞).

First, note that $b_1 = 3$. This is because ϕ is branched at the 3 points $x = 1, \zeta_3, \zeta_3^2$ that lie above $x^3 = 1$; the canonical generator of inertia of ϕ at the points of C_Q above these is, by definition, the automorphism identified with $\zeta_7^1 = \zeta_{21}^3$.

Second, without loss of generality, $b_0 = m$. To see this, note there are m points of C_Q above $x^3 = 0$, so $\gcd(b_0, 3m) = m$; this implies that $b_0 = 2m$ or $b_0 = m$; possibly after replacing the order 3 automorphism with its inverse, we can suppose that $b_0 = m$. Third, $b_\infty = 3m - b_0 - b_1 = 2m - 3$, so $(b_0, b_1, b_\infty) = (m, 3, 2m - 3)$.

Next, we compute the signature f of ϕ from its inertia type. By (2.2), if $0 < n < 3m$, and $(n, 3m) = 1$, then

$$f(\sigma_n) = -1 + \langle -n/3 \rangle + \langle -n/m \rangle + \langle (-2n)/3 + (n/m) \rangle.$$

If $n \equiv 1 \pmod{3}$, then

$$f(\sigma_n) = -1 + (2/3) + \langle -n/m \rangle + \langle (-2/3) + (n/m) \rangle;$$

if $n \equiv 2 \pmod{3}$, then

$$f(\sigma_n) = -1 + (1/3) + \langle -n/m \rangle + \langle (-1/3) + (n/m) \rangle.$$

We deduce that $f(\sigma_n) = 1$ if $n \equiv 1 \pmod{3}$ and $n \in [0, 2m/3] \cup [m, 5m/3] \cup [2m, 8m/3]$ or if $n \equiv 2 \pmod{3}$ and $n \in [0, m/3] \cup [m, 4m/3] \cup [2m, 7m/3]$; otherwise $f(\sigma_n) = 0$.

Finally, by (2.5), the CM-type Φ_Q of J_Q is as given in the statement. \square

For $m = 7$, we verify that the CM-type Φ_Q of the Jacobian of C_Q is simple.

Lemma 6.12. *When $m = 7$, then the CM-type Φ_Q in Proposition 6.11 is simple.*

Proof. Let C_Q be the smooth projective curve with equation $y^7 = x^3 - 1$. From Proposition 6.11, its CM-type is $\Phi = \{\tau_n \mid n = 1, 2, 4, 8, 10, 16\}$. We check that Φ is simple by showing that it is not induced from any proper CM-field. Let $\sigma_i : \mathbb{Q}(\zeta_{3m}) \rightarrow \mathbb{C}$ be the embedding given by $\sigma_i(\zeta_{3m}) = \zeta_{3m}^i$, for $1 \leq i \leq 3m$ with $\gcd(i, 3m) = 1$. Every subgroup of $\text{Gal}(\mathbb{Q}(\zeta_{3m})/\mathbb{Q})$ not containing complex conjugation contains either σ_8 (order 2), σ_{13} (order 2) or σ_4 (order 3). It suffices to show that the subgroup generated by each of these has a coset not contained in Φ but having non-trivial intersection with Φ , so that Φ is not a union of cosets. For $\langle \sigma_8 \rangle$, this is true because of the coset $\{\sigma_4, \sigma_{11}\}$. For $\langle \sigma_{13} \rangle$, this is true because of the coset $\{\sigma_1, \sigma_{13}\}$. For $\langle \sigma_4 \rangle$, this is true because of the coset $\{\sigma_2, \sigma_8, \sigma_{11}\}$. \square

The following lemma will be helpful.

Lemma 6.13. *In the field $L = \mathbb{Q}(\zeta_{21})$, write $\zeta = \zeta_{21}$. Let $\beta_3 = 7/(\zeta^6 - \zeta^{15})$. Let*

$$(6.1) \quad \alpha = (\zeta^7 - \zeta^{14})(\zeta^2 - \zeta^{19}) = (\zeta^9 + \zeta^{12}) - (\zeta^5 + \zeta^{16}).$$

With respect to the CM-type $\Phi = \{1, 2, 4, 8, 10, 16\}$ of L , the following element z satisfies conditions (1)–(3) in Corollary 4.3:

$$(6.2) \quad z = \beta_3 \alpha = 21(\zeta^2 - \zeta^{19}) / ((\zeta^{14} - \zeta^7)(\zeta^6 - \zeta^{15})).$$

Proof. We verify the first claim by computation using that $\zeta^{14} - \zeta^7 = -\sqrt{-3}$. We verify the second claim from Lemma 3.7, part (4). Note that $z = -\sigma_4(\beta_0)$ for β_0 as in *loc. cit.* \square

We now compute the integral PEL datum of $M[17]$ using Proposition 6.11.

Proposition 6.14. *Let $m = 7$ and $F = \mathbb{Q}(\zeta_7)$. Let $\zeta_1 = \beta_1^{-1}$ where $\beta_1 \in \mathcal{O}_F$ is defined in Example 6.7. Consider the units $u_1 = \zeta_7^2 + \zeta_7^5$ and $v = (1 + \zeta_7^3 + \zeta_7^4)^{-1}$ of $\mathbb{Z}[\zeta_7]$. For the family $M[17]$, with monodromy datum $\gamma = (7, 4, (2, 4, 4, 4))$, the integral PEL datum of S_γ has lattice $\Lambda = (\mathcal{O}_F)^2$ with Hermitian form H_B where*

$$B = \zeta_1 v \begin{bmatrix} u_1 & -\zeta_7^2 \\ -\zeta_7^5 & u_1 \end{bmatrix}.$$

Proof. Consider the family $Z = Z_\gamma$ with monodromy datum $\gamma = (7, 4, (1, 1, 1, 4))$. Because $(2, 4, 4, 4) = 4 \cdot (4, 1, 1, 1)$ and $4 \equiv 2^{-1} \pmod{7}$, the action of σ_2 takes Z to the family $M[17]$.

Let

$$(6.3) \quad A = \frac{\zeta_7^2 - \zeta_7^5}{7(1 + \zeta_7 + \zeta_7^6)} \begin{bmatrix} \zeta_7^3 + \zeta_7^4 & -\zeta_7^4 \\ -\zeta_7^3 & \zeta_7^3 + \zeta_7^4 \end{bmatrix}.$$

One can check that A is integral away from 7. We claim that the integral PEL datum of Z is given by the lattice $\Lambda = (\mathcal{O}_F)^2$ and the Hermitian form for the matrix A . This is sufficient to prove the proposition because, by the last statement of Corollary 4.3, the integral PEL datum of $M[17]$ is then given by the lattice $\Lambda = (\mathcal{O}_F)^2$ and the Hermitian form for the matrix $B = \sigma_2^{-1}(A)$.

We turn to computing the integral PEL datum of Z . Let $L = \mathbb{Q}(\zeta_{21})$. Consider the CM-type $\Phi = \{1, 2, 4, 8, 10, 16\}$ for L . Consider the curve C_Q of genus 6 given by the equation $y^7 = x^3 - 1$ and the associated μ_7 -cover $C_Q \rightarrow \mathbb{P}^1$. By Proposition 6.11: this cover is represented by a point Q of Z ; the Jacobian $\text{Jac}(C_Q)$ is a principally polarized abelian variety with complex multiplication by (\mathcal{O}_L, Φ_Q) ; and this CM-type is simple by Lemma 6.12. In other words, Q is a simple distinguished point of Z .

Note that L has class number 1 and L_0 has narrow class number 2. By Corollary 4.6, there exists a unique principally polarized CM-abelian variety of type (\mathcal{O}_L, Φ) up to isomorphism. By Lemma 6.13, the element z from (6.2) satisfies conditions (1)–(3) in Corollary 4.3 with respect to Φ . Thus $\text{Jac}(C_Q)$ is isomorphic to the torus $\mathbb{C}^6 / \Phi(\mathcal{O}_L)$, with principal polarization given by $\zeta = z^{-1}$. Furthermore, the integral PEL datum of Z is given by the lattice $\Lambda = \mathcal{O}_L \subset V = L$ and the Hermitian form $\langle x, y \rangle = \text{tr}_{L/Q}(x\zeta y^T)$. Note that $\mathcal{O}_L = \mathbb{Z}[\zeta_{21}] = \mathcal{O}_F[\zeta_3]$. With respect to the ordered basis $1, \zeta_3$ for \mathcal{O}_L over \mathcal{O}_F , the Hermitian form is given by a matrix in $\text{GL}_2(\mathcal{O}_F)$. By Lemma 6.15, this matrix is A . \square

Lemma 6.15. *With respect to the ordered basis $1, \zeta_3$ for \mathcal{O}_L over \mathcal{O}_F , the Hermitian form $\langle x, y \rangle = \text{tr}_{L/Q}(x\zeta y^T)$ is given by the matrix $A \in \text{GL}_2(\mathcal{O}_F)$, where A is given in (6.3).*

Proof. Let $x, y \in \Lambda$. By Lemma 6.13, since $\beta_3 = 7/(\zeta_7^2 - \zeta_7^5)$ is in \mathcal{O}_{F_0} , we can write:

$$\langle x, y \rangle = \text{tr}_{L/Q}(xz\bar{y}) = \text{tr}_{F/Q}(\beta_3^{-1} \text{tr}_{L/F}(x\alpha^{-1}\bar{y}))$$

where α is as in Equation (6.1).

Let $\tau = \sigma_8 \in \text{Gal}(L/\mathbb{Q})$. Then τ is the generator of $\text{Gal}(L/F)$. Then

$$\text{tr}_{L/\mathbb{Q}}(x\alpha^{-1}\bar{y}) = \text{tr}_{F/\mathbb{Q}}(x\alpha^{-1}\bar{y} + \tau(x)\tau(\alpha^{-1})\tau(\bar{y})).$$

Write $x = x_1 + x_2\zeta_3$ and $y = y_1 + y_2\zeta_3$, for $x_1, x_2, y_1, y_2 \in \mathcal{O}_F$. We compute

$$\text{tr}_{L/\mathbb{Q}}(x\alpha^{-1}\bar{y}) = \text{tr}_{F/\mathbb{Q}}((x_1 + x_2\zeta_3)\alpha^{-1}(\bar{y}_1 + \bar{y}_2\zeta_3^2) + (x_1 + x_2\zeta_3^2)\tau(\alpha^{-1})(\bar{y}_1 + \bar{y}_2\zeta_3)).$$

Write $a_{1,1} = \alpha^{-1} + \tau(\alpha^{-1})$, $a_{1,2} = \zeta_3^2\alpha^{-1} + \zeta_3\tau(\alpha^{-1})$, and $a_{2,1} = \zeta_3\alpha^{-1} + \zeta_3^2\tau(\alpha^{-1})$. Then

$$\text{tr}_{L/\mathbb{Q}}(x\alpha^{-1}\bar{y}) = \text{tr}_{F/\mathbb{Q}}(x_1\bar{y}_1a_{1,1} + x_1\bar{y}_2a_{1,2} + x_2\bar{y}_1a_{2,1} + x_2\bar{y}_2a_{1,1}).$$

We compute that $a_{1,1} = \alpha^{-1} + \tau(\alpha^{-1}) = (\zeta_7^4 + \zeta_7^3)/(1 + \zeta_7 + \zeta_7^6)$, $a_{1,2} = -\zeta_7^4/(1 + \zeta_7 + \zeta_7^6)$, and $a_{2,1} = -\zeta_7^3/(1 + \zeta_7 + \zeta_7^6)$. Thus, $\langle x, y \rangle = \text{tr}_{F/\mathbb{Q}}((x_1, x_2)A(\bar{y}_1, \bar{y}_2)^T)$. \square

7. APPLICATIONS FOR COMPOSITE m

We illustrate how to extend Theorem 5.5 by computing the integral PEL datum in some cases when m is composite. In particular, we determine the integral PEL datum for 6 of the Moonen special families where $m = 4, 6, 10$.

Remark 7.1. Our techniques do not apply well for symplectic Shimura varieties, so we exclude: the modular curve $M[1]$ with $m = 2$; the Picard surface $M[2]$ with $m = 2$; the family $M[7]$ with $m = 4$; and $M[12]$ with $m = 6$. We exclude $M[15]$ with $m = 8$ and $M[20]$ with $m = 12$, since the CM-types for bi-quadratic CM-fields are never simple. Our techniques are not sufficient to handle $M[13]$ with $m = 6$ or $M[19]$ with $m = 9$.

7.1. Difficulties when m is composite. The key to Theorem 5.5 is the existence of a distinguished point P in Z_γ satisfying the properties in Proposition 5.2. When m is composite, this is not always possible for the following reasons.

Remark 7.2. One issue when m is composite is that the curve C_P might not have compact type. For example, the Moonen family $M[12]$, with $\gamma = (6, 4, (1, 1, 1, 3))$, has no admissible degenerations of compact type. The reason is that the two covers with inertia types $a_1 = (1, 1, 4)$ and $a_2 = (2, 1, 3)$ would need to be joined at two points, leading to a cycle in the dual graph of C_P .⁴ The same is true for $M[7]$, with $\gamma = (4, 4, (1, 1, 1, 1))$.

Remark 7.3. Another issue when m is composite is that the integral group algebra $\mathbb{Z}[\mu_m]$ has non-trivial index in $\prod_{1 \leq d|m} \mathbb{Z}[\zeta_d]$. Hence, the Jacobian of a μ_m -cover of \mathbb{P}^1 branched at 3 points might not have complex multiplication by a maximal order. See Section 7.5.

7.2. Integral PEL data for two families with $m = 4$. We find the integral PEL datum for two of the three special families with $m = 4$: $M[4]$ and $M[8]$; we exclude $M[7]$ as it is not unitary. When $m = 4$, the hypotheses of Proposition 5.2 are not satisfied but we can find a distinguished point in the family by direct computation.

Example 7.4. If $m = 4$, then $\beta_0 = -2i$ from Lemma 3.7. Set $u_1 = -1$, so $\beta_1 = -\beta_0 = 2i$. If $f_1 = (0, 1)$, then β_1 satisfies the 3 conditions of Lemma 4.3 and all CM-types of $F = \mathbb{Q}(i)$ are simple.

⁴There is a typo in this case in [12, Lemma 6.4].

Corollary 7.5. *Let $m = 4$ and $F = \mathbb{Q}(i)$. Let $\xi = \beta_1^{-1} = 1/2i$. For the family M with monodromy datum γ below, the integral PEL datum for S_γ has lattice $(\mathcal{O}_F)^r$ with Hermitian form H_B where r and $B \in \mathrm{GL}_r(\mathcal{O}_F)$ are:*

M	$\gamma = (m, N, a)$	r	$B \in \mathrm{GL}_r(\mathcal{O}_F)$
$M[4]$	$(4, 4, (1, 2, 2, 3))$	2	$\mathrm{diag}[\xi, -\xi]$
$M[8]$	$(4, 5, (1, 1, 2, 2, 2))$	3	$\mathrm{diag}[\xi, -\xi, \xi]$

Proof. When $m = 4$, then $a = (3, 2, 3)$ has signature $\mathfrak{f} = (0, 1)$ and $\sigma_2(\beta_1) < 0$. Also $a = (1, 2, 1)$ has signature $\mathfrak{f} = (1, 0)$ and $\sigma_1(-\beta_1) < 0$. By [12, Lemma 6.4], there is a distinguished point in the family $M[4]$ and the family $M[8]$. The family $M[4]$ has an admissible degeneration, expressed in short hand as $(1, 2, 1) + (3, 2, 3)$. The family $M[8]$ has an admissible degeneration, expressed in short hand as $(1, 2, 1) + (3, 2, 3) + (1, 2, 1)$. The result then follows from Theorem 5.5. \square

The case $M[8] : \zeta[1, -1, 1]$ matches line 3 of the table on [18, page 1]

7.3. Remark when $m = 9$. The techniques in this paper are not sufficient to find the integral PEL datum for the Moonen special family $M[19]$ when $m = 9$. This family has inertia type $a = (3, 5, 5, 5)$ and its only admissible degeneration can be expressed in short hand as $(3, 5, 1) + (8, 5, 5)$. For $a_2 = (8, 5, 5)$, the Jacobian has complex multiplication by $\mathbb{Q}[\zeta_3] \times \mathbb{Q}[\zeta_9]$. As the action of $\mathbb{Z}[\mu_9]$ does not extend to the maximal order $\mathbb{Z}[\zeta_3] \times \mathbb{Z}[\zeta_9]$, we do not know how to compute its lattice.

We expect that our technique is sufficient to determine the integral PEL datum when $m = 9$, $N = 4$ and $a = (3, 1, 8, 6)$. We leave the details to the reader to check.

7.4. Generalities for twice a prime.

Notation 7.6. Let $m = 2m'$ where m' is an odd prime. Let $F = \mathbb{Q}(\zeta_m)$ and $F' = \mathbb{Q}(\zeta_{m'})$. We identify $F = F'$ and $\mathbb{Z}[\zeta_m] = \mathbb{Z}[\zeta_{m'}]$, by $\zeta_m = -\zeta_{m'}^{(m'+1)/2}$. There is a bijection between CM types Φ for F and CM types Φ' for F' where $\sigma_j \in \Phi$ for an odd integer $1 \leq j \leq m$ if and only if $\sigma_{j'} \in \Phi'$, where j' is the reduction of j modulo m' .

If β satisfies Corollary 4.7 with respect to the CM type $(\mathbb{Z}[\zeta_{m'}], \Phi')$ then it also satisfies Corollary 4.7 with respect to the CM type $(\mathbb{Z}[\zeta_m], \Phi)$. This can be verified in general but we only need a few cases from Example 6.1 when $m' = 3$ and Example 6.3 when $m' = 5$.

Example 7.7.

m	β	$(\mathbb{Z}[\zeta_{m'}], \Phi')$	$(\mathbb{Z}[\zeta_m], \Phi)$
6	$b_2 = \sqrt{-3}$	$(\mathbb{Z}[\zeta_3], \{2\})$	$(\mathbb{Z}[\zeta_6], \{5\})$
10	$\beta_1 = 5/(\zeta_5^3 - \zeta_5^2)$	$(\mathbb{Z}[\zeta_5], \{2, 4\})$	$(\mathbb{Z}[\zeta_{10}], \{7, 9\})$
10	$\beta_2 = 5/(\zeta_5 - \zeta_5^4)$	$(\mathbb{Z}[\zeta_5], \{1, 2\})$	$(\mathbb{Z}[\zeta_{10}], \{1, 7\})$

7.5. Integral PEL data for 4 families with $m = 6, 10$. We find the integral PEL datum for three special families with $m = 6$, namely $M[5]$, $M[9]$ and $M[14]$, and for the special family with $m = 10$, namely $M[18]$. For $m = 6$, we exclude $M[12]$ as it is not unitary and $M[13]$ (see Remark 7.10). For m composite, the hypotheses of Proposition 5.2 are not satisfied. In each case, we find a distinguished point P in the family for which the issues in Remarks 7.2 and 7.3 do not occur.

Corollary 7.8. *Recall Notation 7.6. For the special family $M[18]$ with monodromy datum $\gamma = (10, 4, (3, 5, 6, 6))$, the integral PEL datum of S_γ has lattice $\Lambda = (\mathcal{O}_{F'}) \oplus (\mathcal{O}_F)^2$ with Hermitian form H_B where $B \in \mathrm{GL}_1(\mathcal{O}_{F'}) \times \mathrm{GL}_2(\mathcal{O}_F)$ is $\mathrm{diag}[\zeta_2] \oplus \mathrm{diag}[\zeta_1, \zeta_2]$ where $\zeta_i = \beta_i^{-1}$.*

Proof. Let $m = 10$ and $m' = 5$. Let $M = M[18]$ with $\gamma = (10, 4, (3, 5, 6, 6))$ and signature type $\mathfrak{f} = (1, 1, 0, 1, 0, 0, 2, 0, 1)$. The proof is similar to that of Theorem 5.5. We produce a simple distinguished point P in the family similar to that in Proposition 5.2(1); it represents an admissible μ_m -cover $\psi : C_P \rightarrow T$, where T is a tree of projective lines and C_P is a curve of compact type such that each of its irreducible components is a curve admitting a μ_m -cover of \mathbb{P}^1 branched at 3 points. We verify by direct computation that each irreducible component of C_P has complex multiplication by either $\mathbb{Z}[\zeta_m]$ or $\mathbb{Z}[\zeta_{m'}]$.

Consider the μ_5 -cover $\psi_2 : C_2 \rightarrow \mathbb{P}^1$ branched at three points, with inertia type $a_2 = (4, 3, 3)$, and signature $\mathfrak{f}_2 = (0, 1, 0, 1)$. Then $A_2 = \mathrm{Jac}(C_2)$ has complex multiplication by $(\mathbb{Z}[\zeta_5], \{2, 4\})$. Consider the induced curve $\tilde{C}_2 = \mathrm{Ind}_5^{10}(C_2)$, which is the disconnected curve consisting of two copies of C_2 , and the induced μ_{10} -cover $\Psi_2 : \tilde{C}_2 \rightarrow \mathbb{P}^1$. Then Ψ_2 is branched at three points and, somewhat imprecisely, we can say that it has inertia type $(8, 6, 6) = \mathrm{Ind}_5^{10}(4, 3, 3)$. Above the first branch point, there are two points η_2 and η'_2 on \tilde{C}_2 , and they are labeled by the two cosets of $\mu_5 \subset \mu_{10}$. Let $\mathcal{A}_2 = \mathrm{Jac}(\tilde{C}_2) \simeq A_2^2$.

As explained in [10, Section 3.1], the signature type of \mathcal{A}_2 is

$$\mathfrak{f}_2 = (0, 1, 0, 1, 0, 0, 1, 0, 1) = (0, 1, 0, 1, 0, 0, 0, 0, 0) + (0, 0, 0, 0, 0, 0, 1, 0, 1).$$

The action of $\mathbb{Z}[\mu_{10}]$ on \mathcal{A}_2 is the diagonal action of $\mathbb{Z}[\zeta_5] \times \mathbb{Z}[\zeta_{10}]$ on A^2 . The first (resp. second) copy of A_2 has CM-type $(\mathbb{Z}[\zeta_5], \{2, 4\})$ (resp. $(\mathbb{Z}[\zeta_{10}], \{7, 9\})$); for this CM-type, the element β_1 (resp. β_1) satisfies Corollary 4.7, as seen in Example 7.7.

Consider the μ_{10} -cover $\psi_1 : C_1 \rightarrow \mathbb{P}^1$ branched at three points, with inertia type $a_1 = (3, 5, 2)$. Above the 3rd branch point, there are two points η_1 and η'_1 on C_1 and they are labeled by the two cosets of $\mu_5 \subset \mu_{10}$. Let $A_1 = \mathrm{Jac}(C_1)$; it has signature type $\mathfrak{f}_1 = (1, 0, 0, 0, 0, 0, 1, 0, 0)$. By [11, Lemma 3.1], A_1 has complex multiplication by $(\mathbb{Z}[\zeta_{10}], \{1, 7\})$. Since $\{1, 7\} = \{3 \cdot 7, 3 \cdot 9\} \pmod{10}$, the element β that satisfies Corollary 4.7 is $\sigma_7(\beta_1) = \beta_2$. All CM types of $\mathbb{Q}(\zeta_5) = \mathbb{Q}(\zeta_{10})$ are simple.

Let \mathcal{C} denote the singular curve, whose components are C_1 and \tilde{C}_2 , formed by identifying $\eta_1 \eta'_1$ with η_2 and η'_2 , so that the cosets are matched correctly, in two ordinary double points. The curve \mathcal{C} admits an admissible μ_{10} -cover Ψ to a chain of two projective lines. By construction, \mathcal{C} has an action by $\mathcal{O}_{F'} \oplus \mathcal{O}_F^2$, with CM-type given by $\mathrm{diag}[\zeta_2] \oplus \mathrm{diag}[\zeta_1, \zeta_2]$.

The last thing to check is that Ψ is represented by a point $P \in Z_\gamma$. Since Ψ is admissible, it can be deformed to a μ_{10} -cover of \mathbb{P}^1 with inertia type $(3, 5, 6, 6)$. This has signature type $\mathfrak{f} = (1, 1, 0, 1, 0, 0, 2, 0, 1)$. Hence, Ψ is represented by a point $P \in Z_\gamma$ and by the preceding paragraph P is a simple distinguished point. \square

Corollary 7.9. *Recall Notation 7.6 and let $\xi = 1/\sqrt{-3}$. For the special families M with monodromy datum $\gamma = (6, N, a)$ as below, the integral PEL datum of S_γ has lattice $\Lambda = (\mathcal{O}_{F'})^{r'} \oplus (\mathcal{O}_F)^r$ with Hermitian form H_B where r', r , and $B \in \mathrm{GL}_{r'}(\mathcal{O}_{F'}) \times \mathrm{GL}_r(\mathcal{O}_F)$ are:*

M	$\gamma = (m, N, a)$	(r', r)	$B \in \mathrm{GL}_{r'}(\mathcal{O}_{F'}) \times \mathrm{GL}_r(\mathcal{O}_F)$
$M[5]$	$(6, 4, (2, 3, 3, 4))$	$(0, 2)$	$\mathrm{diag}[\xi, -\xi]$
$M[9]$	$(6, 4, (1, 3, 4, 4))$	$(1, 2)$	$\mathrm{diag}[-\xi] \oplus \mathrm{diag}[\xi, -\xi]$
$M[14]$	$(6, 5, (2, 2, 2, 3, 3))$	$(1, 3)$	$\mathrm{diag}[\xi] \oplus \mathrm{diag}[\xi, -\xi, \xi]$

Proof. The proof is very similar to that of Corollary 7.8 so we provide only a short sketch.

- (1) Let $M = M[5]$ with $\gamma = (6, 4, (2, 3, 3, 4))$ and $\mathfrak{f} = (1, 0, 0, 0, 1)$. Then C_P is the join of two μ_6 -covers with inertia types $a_1 = (1, 2, 3)$ and $a_2 = (3, 4, 5)$. These have signatures $\mathfrak{f}_1 = (1, 0, 0, 0, 0)$ and $\mathfrak{f}_2 = (0, 0, 0, 0, 1)$. By [11, Lemma 3.1], A_1 has CM by \mathcal{O}_F of type $\Phi_1 = \{1\}$, and A_2 has CM by \mathcal{O}_F of type $\Phi_2 = \{5\}$. Let $b_1 = -b_2 = -\sqrt{-3}$. For $i = 1, 2$, the CM-type Φ_i is simple, and the element $b_i \in \mathcal{O}_F$ satisfies Corollary 4.7 with respect to Φ_i .
- (2) Let $M = M[9]$ with $\gamma = (6, 4, (1, 3, 4, 4))$ and $\mathfrak{f} = (1, 1, 0, 0, 1)$. Then C_P is the join of two covers with inertia types $a_1 = (1, 2, 3)$ and $a_2 = \text{Ind}_3^6(2, 2, 2)$. These have signatures $\mathfrak{f}_1 = (1, 0, 0, 0, 0)$ and $\mathfrak{f}_2 = (0, 1, 0, 0, 1)$. By [11, Lemma 3.1], $A_1 = \text{Jac}(C_1)$ has CM-type $(\mathcal{O}_F, \{1\})$, which has $\beta = b_1 = -b_2$. Also $\mathcal{A}_2 \simeq A^2$ and the action of $\mathbb{Z}[\mu_6]$ on \mathcal{A}_2 is given by the diagonal action of $\mathbb{Z}[\zeta_3] \times \mathbb{Z}[\zeta_6]$ on A^2 . The first copy of A has CM-type $(\mathbb{Z}[\zeta_3], \{2\})$ which has $\beta = b_2$ and the second copy of A has CM-type $(\mathbb{Z}[\zeta_6], \{5\})$ which has $\beta = b_2$.
- (3) Let $M = M[14]$ with $\gamma = (6, 5, (2, 2, 2, 3, 3))$ and $\mathfrak{f} = (2, 0, 0, 1, 1)$. Then C_P is the join of three covers with inertia types $a_1 = (1, 2, 3)$, $a_2 = (3, 4, 5)$ and $a_3 = \text{Ind}_3^6(1, 1, 1)$. These have signatures $\mathfrak{f}_1 = (1, 0, 0, 0, 0)$, $\mathfrak{f}_2 = (0, 0, 0, 0, 1)$, and $\mathfrak{f}_3 = (1, 0, 0, 1, 0)$. By [11, Lemma 3.1], A_1 has CM-type $(\mathcal{O}_F, \{1\})$ which has $\beta = b_1$ and A_2 has CM-type $(\mathcal{O}_F, \{5\})$ which has $\beta = b_2$. Also $\mathcal{A}_3 \simeq A^2$ and the action of $\mathbb{Z}[\mu_6]$ on \mathcal{A}_3 is given by the diagonal action of $\mathbb{Z}[\zeta_3] \times \mathbb{Z}[\zeta_6]$ on A^2 . The first copy of A has CM-type $(\mathbb{Z}[\zeta_3], \{1\})$ which has $\beta = b_1$ and the second copy of A has CM-type $(\mathbb{Z}[\zeta_6], \{4\})$ which has $\beta = b_1$.

□

We checked Corollaries 7.8 and 7.9 independently, using the fact that the Moonen special families for $m = 6, 10$ are subspaces of those for $m' = 3, 5$, up to a Galois twist.

Remark 7.10. Let $M = M[13]$ be the Moonen special family with $\gamma = (6, 4, (1, 1, 2, 2))$ and $\mathfrak{f} = (2, 1, 0, 1, 0)$. This has an admissible degeneration to the join of two covers with inertia types $a_1 = (1, 1, 4)$ and $a_2 = \text{Ind}_3^6(1, 1, 1)$. The first one has signature $\mathfrak{f}_1 = (1, 1, 0, 0, 0)$ and its Jacobian A_1 has CM by an order of finite index in $\mathcal{O}_{F'} \times \mathcal{O}_F$. The index is non-trivial since $\langle x^3 - 1, x^3 + 1 \rangle = \langle 2 \rangle$ in $\mathbb{Z}[x]/\langle x^6 - 1 \rangle$, so our technique does not apply. The other degeneration to $a_1 = (1, 2, 3)$ and $a_2 = (3, 1, 2)$ does not have compact type.

REFERENCES

1. Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 21, Springer-Verlag, Berlin, 1990. MR 1045822 (91i:14034)
2. P. E. Conner and J. Hurrelbrink, *Class number parity*, Series in Pure Mathematics, vol. 8, World Scientific Publishing Co., Singapore, 1988. MR 963648
3. Keith Conrad, *The different ideal*, <https://kconrad.math.uconn.edu/blurbs/gradnumthy/different.pdf>.
4. P. Deligne and G. D. Mostow, *Monodromy of hypergeometric functions and nonlattice integral monodromy*, Inst. Hautes Études Sci. Publ. Math. (1986), no. 63, 5–89. MR 849651
5. Pierre Deligne, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*, Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 247–289. MR 546620

6. Neal Koblitz and David Rohrlich, *Simple factors in the Jacobian of a Fermat curve*, Canadian J. Math. **30** (1978), no. 6, 1183–1205. MR 511556
7. Robert E. Kottwitz, *Points on some Shimura varieties over finite fields*, J. Amer. Math. Soc. **5** (1992), no. 2, 373–444. MR 1124982
8. Kai-Wen Lan, *Arithmetic compactifications of PEL-type Shimura varieties*, London Mathematical Society Monographs Series, vol. 36, Princeton University Press, Princeton, NJ, 2013. MR 3186092
9. Serge Lang, *Complex multiplication*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 255, Springer-Verlag, New York, 1983. MR 713612
10. Wanlin Li, Elena Mantovan, Rachel Pries, and Yunqing Tang, *Newton polygon stratification of the Torelli locus in PEL-type Shimura varieties*, to appear in International Math Research Notices, available on arXiv:1811.00604.
11. ———, *Newton polygons of cyclic covers of the projective line branched at three points*, Research Directions in Number Theory: Women in Numbers IV, AWM series, pages 115–132, available on arXiv:1805.04598.
12. ———, *Newton polygons arising from special families of cyclic covers of the projective line*, Res. Number Theory **5** (2019), no. 1, Art. 12, 31. MR 3897613
13. J. S. Milne, *Shimura varieties and moduli*, Handbook of moduli. Vol. II, Adv. Lect. Math. (ALM), vol. 25, Int. Press, Somerville, MA, 2013, pp. 467–548. MR 3184183
14. Ben Moonen, *Linearity properties of Shimura varieties. I*, J. Algebraic Geom. **7** (1998), no. 3, 539–567. MR 1618140
15. ———, *Special subvarieties arising from families of cyclic covers of the projective line*, Doc. Math. **15** (2010), 793–819. MR 2735989
16. Goro Shimura, *On analytic families of polarized abelian varieties and automorphic functions*, Ann. of Math. (2) **78** (1963), 149–192. MR 156001
17. ———, *Arithmetic of unitary groups*, Ann. of Math. (2) **79** (1964), 369–409. MR 158882
18. ———, *On purely transcendental fields automorphic functions of several variable*, Osaka Math. J. **1** (1964), no. 1, 1–14. MR 176113
19. Paul van Wamelen, *Examples of genus two CM curves defined over the rationals*, Math. Comp. **68** (1999), no. 225, 307–320. MR 1609658
20. Lawrence C. Washington, *Introduction to cyclotomic fields*, second ed., Graduate Texts in Mathematics, vol. 83, Springer-Verlag, New York, 1997. MR 1421575

CENTRE DE RECHERCHES MATHÉMATIQUES, UNIVERSITÉ DE MONTRÉAL, 2920 CHEMIN DE LA TOUR,
MONTRÉAL (QUÉBEC) H3T 1J4, CANADA

Email address: liwanlin@crm.umontreal.ca

DEPARTMENT OF MATHEMATICS, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CA 91125,
USA

Email address: mantovan@caltech.edu

DEPARTMENT OF MATHEMATICS, COLORADO STATE UNIVERSITY, FORT COLLINS, CO 80523, USA

Email address: pries@math.colostate.edu