

**Berry connection from many-body wave functions and
superconductivity: Calculations by the particle number
conserving Bogoliubov-de Gennes equations**

Hiroyasu Koizumi

*Center for computational sciences, University of Tsukuba, Tsukuba, Ibaraki, Japan**

Alto Ishikawa

*Graduate School of Pure and Applied Sciences,
University of Tsukuba, Tsukuba, Ibaraki, Japan*

(Dated: February 18, 2022)

Abstract

A fundamentally revised version of superconductivity theory has been put forward by the present authors since the standard theory of superconductivity based on the BCS theory cannot explain superconductivity in cuprates discovered in 1986, and reexaminations on several experimental results on the conventional superconductors indicate the necessity for a fundamental revision.

The revision is made on the origin of the superconducting phase variable, which is attributed to a Berry connection arising from many-body wave functions. With this revision, the theory can be cast into a particle number conserving formalism.

We have developed a method to calculate superconducting states with the Berry connection using the particle number conserving version of the Bogoliubov-de Gennes equations. An example calculation is made for a model originally built for cuprate superconductors.

I. INTRODUCTION

Since the present work deals with a fundamental revision of the standard theory of superconductivity, we start with historical evolution of superconductivity theory.

After studying the gauge theories in quantum mechanics [1], London developed a phenomenological theory of superconductivity from the view point of the wave mechanics. He put forward the equation

$$\hbar\nabla\chi^{\text{super}} = m\mathbf{v} + \frac{q}{c}\mathbf{A}^{\text{em}} \quad (1)$$

which explains the Meissner effect [2], where \hbar is Planck's constant divided by 2π , χ^{super} is the superpotential, q and m are the charge and mass of the particle, c is the speed of light in vacuum, and \mathbf{A}^{em} is the electromagnetic vector potential, respectively.

London argued that the wave function for the superconducting state is rigid in the sense that when a magnetic field is applied, and the wave function is modified only by the change of the phase factor

$$\Psi_0 \rightarrow \Psi = \Psi_0 e^{i\sum_j \chi^{\text{super}}(\mathbf{r}_j)} \quad (2)$$

where Ψ_0 is the wave function when magnetic field is absent, and Ψ is that when it is present [2].

* koizumi.hiroyasu.fn@u.tsukuba.ac.jp

The supercurrent density \mathbf{j}_s can be obtained using the rigid wave function in Eq. (2) as

$$\mathbf{j}_s = qn_s\mathbf{v} = -\frac{q^2n_s}{mc} \left(\mathbf{A}^{\text{em}} - \frac{\hbar c}{q} \nabla \chi^{\text{super}} \right) \quad (3)$$

where n_s is the number density of the charge carriers. This explains the diamagnetic current that flows when a magnetic field is applied.

When Eq. (3) is combined with one of Maxwell's equation

$$\nabla \times \mathbf{B}^{\text{em}} = \frac{4\pi}{c} \mathbf{j}_s \quad (4)$$

the equation that explains the exclusion of magnetic field from inside superconductors (Meissner effect),

$$\nabla^2 \mathbf{B}^{\text{em}} = -\frac{1}{\lambda_L^2} \mathbf{B}^{\text{em}} \quad (5)$$

is obtained, where the London penetration depth

$$\lambda_L = \sqrt{\frac{mc^2}{4\pi q^2 n_s}} \quad (6)$$

was introduced. This is the length scale where a magnetic field penetrates into a superconductor.

The superpotential gives rise to fluxoid,

$$\Phi_{\text{fluxoid}} = \frac{c\hbar}{q} \oint \nabla \chi^{\text{super}} \cdot d\mathbf{r} \quad (7)$$

In a ring-shaped superconductor where the magnetic field is excluded from the bulk, the flux through the ring is given by

$$\oint_C \mathbf{A}^{\text{em}} \cdot d\mathbf{r} = \Phi_{\text{fluxoid}} \quad (8)$$

where C is a loop along the ring inside the superconductor where $\mathbf{j}_s = 0$.

Experiments confirmed the presence of the fluxoid given by

$$\Phi_{\text{fluxoid}} = \frac{hc}{2e} n \quad (9)$$

where n is an integer [3].

Note that

$$\mathbf{A}^{\text{eff}} = \mathbf{A}^{\text{em}} - \frac{\hbar c}{q} \nabla \chi^{\text{super}} \quad (10)$$

must be gauge invariant since \mathbf{j}_s in Eq. (3) is gauge invariant. Thus, the angular variable χ^{super} that makes \mathbf{A}^{eff} gauge invariant exists in superconductors.

The theory based on the wave mechanics formalism did not succeed in explaining superconductivity. The first successful microscopic theory for superconductivity, the BCS theory, was built by the formalism using creation and annihilation operators [4].

Let $c_{\mathbf{k}\sigma}^\dagger$ be the creation operator for the electron in the metal with wave vector \mathbf{k} and spin σ , the BCS theory uses the ground state composed of different particle number states

$$|\text{BCS}(\theta)\rangle = \prod_{\mathbf{k}} \left(u_{\mathbf{k}} + v_{\mathbf{k}} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger e^{i\theta} \right) |\text{vac}\rangle \quad (11)$$

where $u_{\mathbf{k}}$ and $v_{\mathbf{k}}$ are real parameters that satisfy $u_{\mathbf{k}}^2 + v_{\mathbf{k}}^2 = 1$, and $|\text{vac}\rangle$ is the vacuum that satisfy

$$c_{\mathbf{k}\sigma} |\text{vac}\rangle = 0 \quad (12)$$

The ground state $|\text{BCS}(\theta)\rangle$ is a linear combination of different particle number states, thus, breaks the conservation of the particle number. It has a degeneracy with respect to the choice of θ , and breaking the global $U(1)$ gauge invariance, the invariance of the physical state by the change $c_{\mathbf{k}\sigma} \rightarrow e^{-\frac{i}{2}\theta} c_{\mathbf{k}\sigma}$, $c_{\mathbf{k}\sigma}^\dagger \rightarrow e^{\frac{i}{2}\theta} c_{\mathbf{k}\sigma}^\dagger$, where θ is a constant.

Using the particle number non-conserving formalism, the BCS theory provides a way to calculate the superconducting transition temperature as the energy gap formation temperature by the electron-pairing.

A salient feature of the BCS theory is the presence of the Bogoliubov excitation with an energy gap formed by the electron-pairing [5]. This excitation is most clearly seen if we use the Bogoliubov operators given by

$$\begin{aligned} \gamma_{\mathbf{k}\uparrow} &= u_{\mathbf{k}} e^{-\frac{i}{2}\theta} c_{\mathbf{k}\uparrow} - v_{\mathbf{k}} e^{\frac{i}{2}\theta} c_{-\mathbf{k}\downarrow}^\dagger \\ \gamma_{-\mathbf{k}\downarrow} &= u_{\mathbf{k}} e^{-\frac{i}{2}\theta} c_{-\mathbf{k}\downarrow} + v_{\mathbf{k}} e^{\frac{i}{2}\theta} c_{\mathbf{k}\uparrow}^\dagger \end{aligned} \quad (13)$$

where they satisfy

$$\gamma_{\mathbf{k}\uparrow} |\text{BCS}(\theta)\rangle = 0, \quad \gamma_{-\mathbf{k}\downarrow} |\text{BCS}(\theta)\rangle = 0 \quad (14)$$

This indicates that the superconducting state is the ‘‘vacuum of the Bogoliubov quasiparticles’’, which replaces Eq. (12).

Using the Bogoliubov operators, the BCS Hamiltonian can be cast into the following,

$$H_{\text{BCS}} = \sum_{\mathbf{k}, \sigma} E_{\mathbf{k}} \gamma_{\mathbf{k}\sigma}^{\dagger} \gamma_{\mathbf{k}\sigma} + E_{\text{const}} \quad (15)$$

where

$$E_{\mathbf{k}} = \sqrt{(\mathcal{E}(\mathbf{k}))^2 + \Delta_{\mathbf{k}}^2} \quad (16)$$

with $\mathcal{E}(\mathbf{k})$ being the energy of Bloch electrons measured from the Fermi energy; $\Delta_{\mathbf{k}}$ is the energy gap given by

$$\Delta_{\mathbf{k}} = 2E_{\mathbf{k}}u_k v_k, \quad u_k^2 = \frac{1}{2} \left(1 + \frac{\mathcal{E}(\mathbf{k})}{E_{\mathbf{k}}} \right), \quad v_k^2 = \frac{1}{2} \left(1 - \frac{\mathcal{E}(\mathbf{k})}{E_{\mathbf{k}}} \right) \quad (17)$$

$E_{\mathbf{k}}$ explains the energy gap spectrum observed in superconductors.

In the BCS theory, the induced current by a magnetic field is calculated as a linear response to \mathbf{A}^{em} , which gives the formula very similar to Eq. (3) without the term with $\nabla \chi^{\text{super}}$. The obtained current is not gauge invariant, and the gauge invariance of the induced current became a big issue. After seminal works by Anderson [6, 7], a final resolution was provided by Nambu[8], culminating to the idea ‘‘spontaneous gauge symmetry breaking’’. Nambu showed the the Nambu-Goldstone mode appears due to the degeneracy of the ground state with respect to the choice of θ , and this mode retrieves the gauge invariance.

Independent of the BCS theory, a phenomenological theory was developed by Ginzburg and Landau [9] based in the London theory of superconductivity. The gauge invariant supercurrent is most easily calculated by this Ginzburg-Landau theory [9]. In this theory a free energy for the superconducting state is given as a functional of an order parameter

$$\Psi_{\text{GL}} = n_s^{1/2} e^{i\chi^{\text{super}}} \quad (18)$$

The free energy is composed of a material part

$$F_{\text{mat}} = F_{\text{normal}} + \int d^3r \frac{1}{2m} \left| \left(\frac{\hbar}{i} \nabla - \frac{q}{c} \mathbf{A}^{\text{em}} \right) \Psi_{\text{GL}} \right|^2 + \int d^3r \left(\alpha |\Psi_{\text{GL}}|^2 + \frac{\beta}{2} |\Psi_{\text{GL}}|^4 \right) \quad (19)$$

where $\alpha < 0$, $\beta > 0$ are parameters and F_{normal} is the free energy of the normal phase, and a magnetic field part

$$F_{\text{mag}} = \int d^3r \frac{1}{8\pi} (\mathbf{B}^{\text{em}})^2 \quad (20)$$

In the Ginzburg-Landau theory, the supercurrent is calculated as

$$\mathbf{j}_s = -c \frac{\delta F_{\text{mat}}}{\delta \mathbf{A}^{\text{em}}(\mathbf{r})} = -\frac{q^2 n_s}{mc} \left(\mathbf{A}^{\text{em}} - \frac{\hbar c}{q} \nabla \chi^{\text{super}} \right) \quad (21)$$

in accordance with Eq. (3). The gauge invariance is retrieved by χ^{super} [10].

The connection between the BCS and Ginzburg-Landau theories was provided by Gor'kov [11]. The Gor'kov's derivation yields,

$$m = 2m^*, \quad q = -2e \quad (22)$$

where m^* is the effective mass of electron in the metal.

Although the Gor'kov derivation has been considered as a valid one, it disagrees with the experimental value $m = 2m_e$ from the London moment measurements where m_e is the free electron mass [12, 13].

A notable point of the Ginzburg-Landau theory is that it introduces the coherence length

$$\xi_{\text{GL}} = \sqrt{\frac{\hbar^2}{2m|\alpha|}} \quad (23)$$

The BCS theory also introduces a similar quantity, the BCS coherence length,

$$\xi_{\text{BCS}} = \frac{\hbar v_{\text{Fermi}}}{\pi \Delta} \quad (24)$$

Currently, it is interpreted that it is the size of the electron-pair, where Δ is the energy gap and v_{Fermi} is the velocity of the electron at the Fermi energy [4].

Based on the Ginzburg-Landau theory, Abrikosov showed that there are two types of superconductors depending on the ratio of λ_L and ξ_{GL} ,

$$\text{type I} \quad \frac{\lambda_L}{\xi_{\text{GL}}} < \frac{1}{\sqrt{2}}; \quad \text{type II} \quad \frac{\lambda_L}{\xi_{\text{GL}}} > \frac{1}{\sqrt{2}}, \quad (25)$$

In type II superconductors, applied magnetic field can penetrate the sample as vortices, and ξ_{GL} is the core size of such vortices [14].

Remarkable effects caused by χ^{super} were predicted by Josephson [15]. They are called “dc Josephson effect” and “ac Josephson effect”, and occur in the junction composed of two superconductors separated by a thin barrier (or insulator region) depicted in Fig. 1.

The current through the junction is given by

$$J = J_c \sin \phi \quad (26)$$




FIG. 1. Josephson junction S_L -I- S_R (superconductor-insulator-superconductor junction) considered by Josephson. μ_R and μ_L are chemical potentials of S_L and S_R , respectively. When voltage V exists the relation $eV = \mu_R - \mu_L$ holds.

where J_c is a parameter for the junction, and ϕ is given using the gauge invariant \mathbf{A}^{eff} as

$$\phi = -\frac{q}{\hbar c} \int_L^R \left(\mathbf{A}^{\text{em}} - \frac{\hbar c}{q} \nabla \chi^{\text{super}} \right) + \phi_0 \quad (27)$$

where ϕ_0 is a constant and the integration is taken along the line connecting the two superconductors.

The ac Josephson effect occurs when there is a voltage across the junction. When the voltage is V , time variation of ϕ given by

$$\dot{\phi} = \frac{2eV}{\hbar} \quad (28)$$

occurs.

This effect is most clearly seen if a radiation field with frequency f exists; the resonance condition $\dot{\phi} = 2\pi f n$, where n is an integer, gives

$$V = \frac{\hbar f}{2e} n \quad (29)$$

Then, steps are observed in the I - V curve (current-voltage curve) [16].

Josephson obtained Eq. (28) based on the BCS theory by assuming $q = -2e$ [15]. His derivation had been long considered to be right one. However, it has been argued that it misses a contribution; Eq. (28) can be obtained by $q = -e$ [17–21].

Before the discovery of high temperature superconductivity in 1986 [22], the BCS theory and its extensions have been very successful and was believed that superconductivity was a well-understood solved problem. However, the cuprate superconductivity is markedly different from the BCS superconductivity in a number of ways; for example, the magnetism is harmful in the BCS superconductivity since it causes the breaking of electron-pairing, however magnetism coexists with superconductivity in the cuprate [23]; the superconducting

transition temperature is not the energy gap formation temperature, but corresponds to the stabilizing temperature for loop currents of ξ_{GL} size in optimally doped sample of the cuprates [24]; the charge carriers in the room temperature become small polarons at low temperatures [25].

Since the cuprate superconductivity is so different from the BCS superconductivity, a theory much departs from the standard one is expected to be needed [17]. The superconducting state transition temperature indicates the importance of ξ_{GL} -sized loop currents, and it has been argued that ξ_{GL} -sized spin-vortices and loop currents appear around the small polarons in the bulk, explaining the magnetic excitation spectrum observed by inelastic neutron scattering [26].

Efforts toward the elucidation of the cuprate superconductivity lead some researchers to reexamine superconductivity from very fundamental levels. Those efforts revealed that there are some experimental facts disagreeing with the standard theory even in the conventional superconductors [27]. We would like to point out four major problems below:

1) Supercurrent in the standard theory contradicts the reversible superconducting-normal phase transition in a magnetic field observed in type I superconductors.

Magnetic field is excluded from the bulk of superconductors, thereby the supercurrent is generated in the surface region of width λ_L in Eq. (6) as the screening current for the magnetic field.

However, the Meissner effect is not just the exclusion of the magnetic field. It also indicates that the superconducting state is a thermodynamically stable phase in the $T - H$ plane, where T is the temperature and H is the external magnetic field. Thus, the surface current stops without generating irreversible Joule heat during the superconducting to currentless normal metallic phase transition. Note that the thermodynamically stable phase of the normal metal in a magnetic field is currentless.

Actually, after the discovery of the Meissner effect, the reversibility of the superconducting-normal state transition in the presence of a magnetic field was confirmed [28–32]; and the state of the art calorimetry indicates that 99.99% of the supercurrent stops without current carriers undergoes irreversible collisions (see Appendix B of Ref. [33]). Hirsch argued that the standard theory of superconductivity cannot explain how this transition is possible since the supercurrent generated by the flow of electron pairs inevitably produces the Joule heat

due to the existence of a significant number of broken pairs that flow with dissipation during the transition [33–35].

It is also note worthy that the use of the linear response theory to calculate the supercurrent may be problematic since it is tied to the fluctuation-dissipation relation [36], and the current calculated by it usually causes dissipation. This point has drawn renewed attention in relation to the absence of a dissipative quantum phase transition in Josephson junctions [37–39].

One may say that the Ginzburg-Landau theory is derivable from the BCS theory [11], the supercurrent generation in the latter theory can be used as the evidence that the BCS theory explains the Meissner effect. However, the Gor’kov’s derivation from BCS to Ginzburg-Landau theories disagrees in the mass in the London moment as explained below.

2) Experiments indicate that the mass in the London moment is the free electron mass, however, the standard theory predicts it to be an effective mass.

The London moment is the magnetic moment generated by a rotating superconductor in a magnetic field [2, 40]. Inside the superconductor magnetic field, the “*London field*”, is produced by the supercurrent flowing in the surface region.

Let us consider a superconducting sphere rotating about its symmetry axis with constant angular velocity $\boldsymbol{\omega}$. Then, the velocity at the position \mathbf{r} measured from the center of the sphere is given by

$$\mathbf{v} = \boldsymbol{\omega} \times \mathbf{r} \tag{30}$$

Electrons inside the superconductor move with this velocity to shield the background positive charge. Substituting this in Eq. (1) and using the relation $\mathbf{B}^{\text{em}} = \nabla \times \mathbf{A}^{\text{em}}$, we have

$$\mathbf{B}^{\text{em}} = -\frac{2mc}{q}\boldsymbol{\omega} \tag{31}$$

This is the London field, where m is mass, q is charge, and c is the speed of light.

The London moment and field have been measured many times using different materials, ranging from the conventional superconductor [41–45] to the high T_c cuprates [46, 47] and heavy fermion superconductors [48]. The results always indicate that the mass m is the free electron mass m_e if $q = -e$ is used, not the effective mass m^* predicted by the standard theory including the Gorkov’s derivation.




FIG. 2. Josephson junction in a real experimental situation, connected to leads, denoted as Lead_L and Lead_R . The ac Josephson effect is observed as the quantized voltage given in Eq. (29) under the application a microwave of frequency f and a dc current through the junction.

3) In the standard theory, the breakdown of the global $U(1)$ gauge invariance or the non-conservation of the particle number is essential.

It is sensible to consider that the particle number is conserved in an isolated superconductor. However, the standard theory requires to use the particle number non-conserving formalism as manifested in the use of Eq. (11) [49, 50]. It has been also claimed that the use of θ as a physically meaningful parameter is against the superselection rule for charge [51]. Thus, it has been argued that current formalism is an approximation of the true formalism in which the particle number is conserved [50, 52].

4) The derivation of the ac Josephson effect takes into account only the half of the contributions.

We shall revisit the ac Josephson effect: let us calculate $\dot{\phi}$ using Eq. (27),

$$\dot{\phi} = -\frac{q}{\hbar c} \int_R^L d\mathbf{r} \cdot \left(\partial_t \mathbf{A}^{\text{em}} - \frac{\hbar c}{q} \nabla \partial_t \chi^{\text{super}} \right) = \frac{q}{\hbar} \int_R^L d\mathbf{r} \cdot \mathbf{E}^{\text{em}} - \frac{q}{\hbar} \left(-\varphi^{\text{em}} - \frac{\hbar}{q} \partial_t \chi^{\text{super}} \right) \Big|_R^L \quad (32)$$

where the relation

$$\mathbf{E}^{\text{em}} = -\frac{1}{c} \partial_t \mathbf{A}^{\text{em}} - \nabla \varphi^{\text{em}} \quad (33)$$

is used.

There are two contributions for $\dot{\phi}$. The first one is

$$\frac{q}{\hbar} \int_R^L d\mathbf{r} \cdot \mathbf{E}^{\text{em}} = -\frac{qV}{\hbar} \quad (34)$$

where V is the voltage across the junction.

The second one is

$$-\frac{q}{\hbar} \left(-\varphi^{\text{em}} - \frac{\hbar}{q} \partial_t \chi^{\text{super}} \right) \Big|_R^L = \frac{q}{\hbar} \frac{\mu_L - \mu_R}{e} \quad (35)$$

Here, the time-component partner of Eq. (10),

$$\varphi^{\text{eff}} = \varphi^{\text{em}} + \frac{\hbar}{q} \partial_t^{\text{super}} \chi \quad (36)$$

is identified as

$$\varphi^{\text{eff}} = \frac{\mu}{e} \quad (37)$$

where μ is the chemical potential.

The balance between the voltage and the chemical potential difference yields,

$$eV = \mu_R - \mu_L \quad (38)$$

Thus, the sum of the two contributions yields

$$\dot{\phi} = -\frac{2qV}{\hbar} \quad (39)$$

In order to obtain the result in Eq. (28), q needs to be $q = -e$, this contradicts the value used by Josephson $q = -2e$.

The above discrepancy comes from the fact that only one of the two contributions in Eq. (32) is included in the Josephson's derivation [15, 53].

The first term with \mathbf{E}^{em} in Eq. (32) arises when charged particles go through the electric field in the insulator region. The second term with φ^{em} and χ^{super} arises due to the fact that the chemical potentials are maintained by the contact with the leads; in order to keep the chemical potential constant, electrons enter from one of the leads and exit to the other lead as electrons tunnel through "I" region in Fig. 2. Since experiments are performed in the presence of a dc current in addition to a microwave field, it is sensible to consider that the two contributions both exist.

If $q = -2e$ is used as in the standard theory, the Josephson relation becomes $\dot{\phi} = \frac{4eV}{\hbar}$. In this case, half-integers will appear in Eq. (29). Indeed, such effects have been observed [54].

As Eq. (3) indicates, the most important ingredient for supercurrent generation is χ^{super} . If an angular variable χ with period 2π that satisfy

$$\nabla \chi^{\text{super}} = -\frac{1}{2} \nabla \chi \quad (40)$$

exists, the experimental values of fluxoid are obtained by $q = -e$. $q = -2e$ contributions may exist since the electron pairing occurs in the system; however, the view that the supercurrent is the flow of paired-electrons will not be valid according to the problem given above in the item 1). It is also noteworthy to recognize that careful treatments of the spatial and temporal fluctuations of \mathbf{E}^{em} are needed to understand Josephson effect related phenomena, correctly [21, 55].

The new theory is the one that can be formulated in a particle number conserving manner [13, 18, 19, 21, 56, 57]. We will present a method to calculate properties of superconducting states, such as diamagnetic response current and persistent current, using the particle number conserving formalism.

The organization of the present work is as follows: In Section II particle number conserving version of the BCS theory is explained. This formalism becomes possible due to the presence of non-trivial Berry connection. In Section III model calculations are performed using the particle number conserving Bogoliubov-de Gennes equations derived from the particle number conserving version of the BCS theory. A model constructed for the cuprate superconductivity is used to demonstrate calculations. We conclude this work with some remarks in Section IV.

Four appendices are included: Berry connection from many-body wave functions and the modification of Maxwell's equations are explained in Appendix A; appearance of \mathbf{A}^{fic} from spin-twisting itinerant motion of electrons is explained in Appendix B; reversible superconducting-normal metal phase transition in a magnetic field is discussed in Appendix C; and collective mode χ and associated number changing operators are derived in Appendix D.

II. PARTICLE NUMBER CONSERVING VERSION OF THE BCS THEORY USING THE SUPERCONDUCTING PHASE FROM THE BERRY CONNECTION

In the original BCS theory, electrons in the normal metallic state are assumed to be well-described by the free electrons with the effective mass m^* . Then, the electron field operators are given by

$$\hat{\Psi}_\sigma(\mathbf{r}) = \frac{1}{\sqrt{\mathcal{V}}} \sum_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} c_{\mathbf{k}\sigma} \quad (41)$$

where \mathcal{V} is the volume of the system.

Using the Bogoliubov operators [58, 59] they are expressed as

$$\begin{aligned}\hat{\Psi}_\uparrow(\mathbf{r}) &= \frac{1}{\sqrt{\mathcal{V}}} \sum_{\mathbf{k}} e^{\frac{i}{2}\theta} e^{i\mathbf{k}\cdot\mathbf{r}} \left(\gamma_{\mathbf{k}\uparrow} u_k - \gamma_{-\mathbf{k}\downarrow}^\dagger v_k \right) \\ \hat{\Psi}_\downarrow(\mathbf{r}) &= \frac{1}{\sqrt{\mathcal{V}}} \sum_{\mathbf{k}} e^{\frac{i}{2}\theta} e^{i\mathbf{k}\cdot\mathbf{r}} \left(\gamma_{\mathbf{k}\downarrow} u_k + \gamma_{-\mathbf{k}\uparrow}^\dagger v_k \right)\end{aligned}\quad (42)$$

Now we depart from the standard theory by replacing $e^{\pm\frac{i}{2}\theta}$ in Eq. (42) by the number changing operators $e^{\pm\frac{i}{2}\hat{\chi}}$. Here, we do not specify the coordinate in $\hat{\chi}$ (we will introduce the coordinate dependence, later). Then, the Bogoliubov transformation in Eq. (13) becomes

$$\begin{aligned}\gamma_{\mathbf{k}\uparrow} &= u_k e^{\frac{i}{2}\hat{\chi}} c_{\mathbf{k}\uparrow} - v_k e^{-\frac{i}{2}\hat{\chi}} c_{-\mathbf{k}\downarrow}^\dagger \\ \gamma_{-\mathbf{k}\downarrow} &= u_k e^{\frac{i}{2}\hat{\chi}} c_{-\mathbf{k}\downarrow} + v_k e^{-\frac{i}{2}\hat{\chi}} c_{\mathbf{k}\uparrow}^\dagger\end{aligned}\quad (43)$$

Note that the above Bogoliubov operators conserve particle numbers. Terms like $e^{\frac{i}{2}\hat{\chi}} c_{\mathbf{k}\sigma}$ can be interpreted that an electron in the (\mathbf{k}, σ) single-electron mode is removed and an electron is added to the collective mode described by χ ; those like $e^{-\frac{i}{2}\hat{\chi}} c_{\mathbf{k}\sigma}^\dagger$ create an electron in the (\mathbf{k}, σ) single-electron mode and subtract an electron from the collective mode described by χ . Thus, the Bogoliubov operators cause the fluctuation of the number of electrons in the collective mode by transferring electrons between single-particle modes and collective mode. As a consequence, the ground state becomes a linear combination of states with different number of electrons in the collective mode [56]. This state replaces the state with fluctuating total number of electrons (given by Eq. (11)) in the standard theory.

By including $\frac{1}{\sqrt{\mathcal{V}}} e^{i\mathbf{k}\cdot\mathbf{r}}$ in u_k and v_k , we define

$$u_{\mathbf{k}}(\mathbf{r}) = \frac{1}{\sqrt{\mathcal{V}}} e^{i\mathbf{k}\cdot\mathbf{r}} u_k, \quad v_{\mathbf{k}}(\mathbf{r}) = \frac{1}{\sqrt{\mathcal{V}}} e^{i\mathbf{k}\cdot\mathbf{r}} v_k \quad (44)$$

Then, the field operators become

$$\begin{aligned}\hat{\Psi}_\uparrow(\mathbf{r}) &= \sum_{\mathbf{k}} e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} \left(\gamma_{\mathbf{k}\uparrow} u_{\mathbf{k}}(\mathbf{r}) - \gamma_{\mathbf{k}\downarrow}^\dagger v_{\mathbf{k}}^*(\mathbf{r}) \right) \\ \hat{\Psi}_\downarrow(\mathbf{r}) &= \sum_{\mathbf{k}} e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} \left(\gamma_{\mathbf{k}\downarrow} u_{\mathbf{k}}(\mathbf{r}) + \gamma_{\mathbf{k}\uparrow}^\dagger v_{\mathbf{k}}^*(\mathbf{r}) \right)\end{aligned}\quad (45)$$

where the spatial dependence is included in $\hat{\chi}$.

Now we allow the coordinate dependent functions other than plane waves. We use the label n in place of the wave number \mathbf{k} for them. As a result, the field operators become

$$\begin{aligned}\hat{\Psi}_\uparrow(\mathbf{r}) &= \sum_n e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} \left(\gamma_{n\uparrow} u_n(\mathbf{r}) - \gamma_{n\downarrow}^\dagger v_n^*(\mathbf{r}) \right) \\ \hat{\Psi}_\downarrow(\mathbf{r}) &= \sum_n e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} \left(\gamma_{n\downarrow} u_n(\mathbf{r}) + \gamma_{n\uparrow}^\dagger v_n^*(\mathbf{r}) \right)\end{aligned}\quad (46)$$

The particle number conserving Bogoliubov operator $\gamma_{n\sigma}$ satisfies

$$\gamma_{n\sigma}|\text{Gnd}(N)\rangle = 0 \quad (47)$$

where $|\text{Gnd}(N)\rangle$ is the ground state with the total number of particles N .

The ground state also satisfies

$$e^{\pm\frac{i}{2}\hat{\chi}(\mathbf{r})}|\text{Gnd}(N)\rangle = e^{\pm\frac{i}{2}\chi(\mathbf{r})}|\text{Gnd}(N \pm 1)\rangle \quad (48)$$

The above phase factor gives

$$\langle\text{Gnd}(N)|e^{\frac{i}{2}\hat{\chi}(\mathbf{r}_f)}e^{-\frac{i}{2}\hat{\chi}(\mathbf{r}_i)}|\text{Gnd}(N)\rangle = e^{\frac{i}{2}\int_{\mathbf{r}_i}^{\mathbf{r}_f}\nabla\chi\cdot d\mathbf{r}} \quad (49)$$

indicating that the phase $\frac{1}{2}\int_{\mathbf{r}_i}^{\mathbf{r}_f}\nabla\chi\cdot d\mathbf{r}$ is acquired when the particle travels from \mathbf{r}_i to \mathbf{r}_f , in accordance with the presence of \mathbf{A}^{fic} in Eq. (A20).

Let us consider the electronic Hamiltonian expressed by the field operators

$$H = \sum_{\sigma} \int d^3r \hat{\Psi}_{\sigma}^{\dagger}(\mathbf{r})h(\mathbf{r})\hat{\Psi}_{\sigma}(\mathbf{r}) - \frac{1}{2} \sum_{\sigma,\sigma'} \int d^3r d^3r' V_{\text{eff}}(\mathbf{r}, \mathbf{r}') \hat{\Psi}_{\sigma}^{\dagger}(\mathbf{r})\hat{\Psi}_{\sigma'}^{\dagger}(\mathbf{r}')\hat{\Psi}_{\sigma'}(\mathbf{r}')\hat{\Psi}_{\sigma}(\mathbf{r}) \quad (50)$$

where $h(\mathbf{r})$ is the single-particle Hamiltonian given by

$$h(\mathbf{r}) = \frac{1}{2m_e} \left(\frac{\hbar}{i}\nabla + \frac{e}{c}\mathbf{A}^{\text{em}} \right)^2 + U(\mathbf{r}) - \mu \quad (51)$$

and $U(\mathbf{r})$ is the single particle potential energy, μ is the chemical potential, and $-V_{\text{eff}}$ is the effective interaction between electrons. In the BCS theory, $-V_{\text{eff}}$ describes the attractive interaction between electrons via the virtual phonon exchange.

We perform the following mean field approximation

$$\begin{aligned} H^{\text{MF}} &= \sum_{\sigma} \int d^3r \hat{\Psi}_{\sigma}^{\dagger}(\mathbf{r})h(\mathbf{r})\hat{\Psi}_{\sigma}(\mathbf{r}) + \int d^3r d^3r' \left[\Delta(\mathbf{r}, \mathbf{r}') \hat{\Psi}_{\uparrow}^{\dagger}(\mathbf{r})\hat{\Psi}_{\downarrow}^{\dagger}(\mathbf{r}') e^{-\frac{i}{2}(\hat{\chi}(\mathbf{r})+\hat{\chi}(\mathbf{r}'))} + \text{H.c.} \right] \\ &+ \int d^3r d^3r' \frac{|\Delta(\mathbf{r}, \mathbf{r}')|^2}{V_{\text{eff}}(\mathbf{r}, \mathbf{r}')} \end{aligned} \quad (52)$$

where the gap function $\Delta(\mathbf{r}, \mathbf{r}')$ is defined as

$$\Delta(\mathbf{r}, \mathbf{r}') = V_{\text{eff}}(\mathbf{r}, \mathbf{r}') \langle e^{\frac{i}{2}(\hat{\chi}(\mathbf{r})+\hat{\chi}(\mathbf{r}'))} \hat{\Psi}_{\uparrow}(\mathbf{r})\hat{\Psi}_{\downarrow}(\mathbf{r}') \rangle \quad (53)$$

Due to the factor $e^{\frac{i}{2}(\hat{\chi}(\mathbf{r})+\hat{\chi}(\mathbf{r}'))}$ the expectation value can be calculated using the particle number fixed state.

Using commutation relations for $\hat{\Psi}_\sigma^\dagger(\mathbf{r})$ and $\hat{\Psi}_\sigma(\mathbf{r})$,

$$\begin{aligned}\{\hat{\Psi}_\sigma(\mathbf{r}), \hat{\Psi}_{\sigma'}^\dagger(\mathbf{r}')\} &= \delta_{\sigma\sigma'}\delta(\mathbf{r} - \mathbf{r}') \\ \{\hat{\Psi}_\sigma(\mathbf{r}), \hat{\Psi}_{\sigma'}(\mathbf{r}')\} &= 0 \\ \{\hat{\Psi}_\sigma^\dagger(\mathbf{r}), \hat{\Psi}_{\sigma'}^\dagger(\mathbf{r}')\} &= 0\end{aligned}\quad (54)$$

the following relations are obtained

$$\begin{aligned}[\hat{\Psi}_\uparrow(\mathbf{r}), H_{\text{MF}}] &= h(\mathbf{r})\hat{\Psi}_\uparrow(\mathbf{r}) + \int d^3r' \Delta(\mathbf{r}, \mathbf{r}')\hat{\Psi}_\downarrow^\dagger(\mathbf{r}')e^{-\frac{i}{2}(\hat{\chi}(\mathbf{r})+\hat{\chi}(\mathbf{r}'))} \\ [\hat{\Psi}_\downarrow(\mathbf{r}), H_{\text{MF}}] &= h(\mathbf{r})\hat{\Psi}_\downarrow(\mathbf{r}) - \int d^3r' \Delta(\mathbf{r}, \mathbf{r}')\hat{\Psi}_\uparrow^\dagger(\mathbf{r}')e^{-\frac{i}{2}(\hat{\chi}(\mathbf{r})+\hat{\chi}(\mathbf{r}'))}\end{aligned}\quad (55)$$

The particle number conserving Bogoliubov operators $\gamma_{n\sigma}$ and $\gamma_{n\sigma}^\dagger$ obey fermion commutation relations. They are chosen to satisfy

$$[H_{\text{MF}}, \gamma_{n\sigma}] = -\epsilon_n\gamma_{n\sigma}, \quad [H_{\text{MF}}, \gamma_{n\sigma}^\dagger] = \epsilon_n\gamma_{n\sigma}^\dagger \quad (56)$$

yielding the diagonalized H_{MF} ,

$$H_{\text{MF}} = E_g + \sum_{n,\sigma} \epsilon_n \gamma_{n\sigma}^\dagger \gamma_{n\sigma} \quad (57)$$

where the excitation energies satisfy $\epsilon_n > 0$, and E_g is the ground state energy.

From Eqs. (55), (56), and (57), the following system of equations are obtained

$$\begin{aligned}\epsilon_n e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} u_n(\mathbf{r}) &= h(\mathbf{r})e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} u_n(\mathbf{r}) + \int d^3r' \Delta(\mathbf{r}, \mathbf{r}')e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} v_n(\mathbf{r}') \\ \epsilon_n e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} v_n^*(\mathbf{r}) &= -h(\mathbf{r})e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} v_n^*(\mathbf{r}) + \int d^3r' \Delta(\mathbf{r}, \mathbf{r}')e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} u_n^*(\mathbf{r}')\end{aligned}\quad (58)$$

Using the relation in Eq. (48), the above are cast into the following,

$$\begin{aligned}\epsilon_n u_n(\mathbf{r}) &= \bar{h}(\mathbf{r})u_n(\mathbf{r}) + \int d^3r' \Delta(\mathbf{r}, \mathbf{r}')v_n(\mathbf{r}') \\ \epsilon_n v_n(\mathbf{r}) &= -\bar{h}^*(\mathbf{r})v_n(\mathbf{r}) + \int d^3r' \Delta^*(\mathbf{r}, \mathbf{r}')u_n(\mathbf{r}')\end{aligned}\quad (59)$$

where

$$\bar{h}(\mathbf{r}) = \frac{1}{2m_e} \left(\frac{\hbar}{i} \nabla + \frac{e}{c} \mathbf{A}^{\text{em}} - \frac{\hbar}{2} \nabla \chi \right)^2 + U(\mathbf{r}) - \mu \quad (60)$$

and

$$\Delta(\mathbf{r}, \mathbf{r}') = V_{\text{eff}}(\mathbf{r}, \mathbf{r}') \sum_n [u_n(\mathbf{r})v_n^*(\mathbf{r}')(1 - f(\epsilon_n)) - u_n(\mathbf{r}')v_n^*(\mathbf{r})f(\epsilon_n)] \quad (61)$$

$f(\epsilon_n)$ is the Fermi function given by

$$f(\epsilon_n) = \frac{1}{e^{\frac{\epsilon_n}{k_B T}} + 1} \quad (62)$$

where k_B is the Boltzmann constant.

The above system of equations are Bogoliubov-de Gennes equations [58] using the particle number conserving Bogoliubov operators [56].

Note that the gauge potential in the single particle Hamiltonian $\bar{h}(\mathbf{r})$ is the effective one given by

$$\mathbf{A}^{\text{eff}} = \mathbf{A}^{\text{em}} - \frac{\hbar c}{2e} \nabla \chi \quad (63)$$

In the BCS superconductor, the energy gain by the electron-pair formation exceeds the energy loss by performing the spin-twisting itinerant motion that is necessary to have the number changing operators; thus, even if the spin-orbit interaction is negligibly small, the spin-twisting itinerant motion will occur. This will be the reason for the simultaneous occurrence of superconductivity and energy gap formation.

By solving the system of equations composed of Eqs. (59), (60), and (61), with the condition $\mathbf{A}^{\text{eff}} = 0$, we obtain the currentless solutions for u_n, v_n , which we denote as \acute{u}_n, \acute{v}_n .

Using \acute{u}_n, \acute{v}_n , we express u_n, v_n as

$$u_n(\mathbf{r}) = \acute{u}_n(\mathbf{r}) e^{\frac{i}{2} \chi(\mathbf{r})}, \quad v_n(\mathbf{r}) = \acute{v}_n(\mathbf{r}) e^{-\frac{i}{2} \chi(\mathbf{r})} \quad (64)$$

We obtain χ in the following manner. First, we note that χ has to satisfy constraints,

$$\oint_{C_\ell} \nabla \chi \cdot d\mathbf{r} = 2\pi w_{C_\ell}[\chi] \quad (65)$$

where $w_{C_\ell}[\chi]$ is the integer called the winding number. The value $w_{C_\ell}[\chi]$ needs to satisfy the condition similar to the one in Eq. (B16) to make the wave function single-valued.

We obtain χ for the ground state by minimizing the total energy under the above constraint. For this purpose, we use the following functional,

$$F[\nabla \chi] = E[\nabla \chi] + \sum_{\ell=1}^{N_{\text{loop}}} \lambda_\ell \left(\oint_{C_\ell} \nabla \chi \cdot d\mathbf{r} - 2\pi w_{C_\ell}[\chi] \right), \quad (66)$$

where

$$E[\nabla \chi] = \langle \text{Gnd} | H_{\text{eff}} | \text{Gnd} \rangle \quad (67)$$

is obtained using Eq. (64); λ_ℓ 's are Lagrange multipliers, and $\{C_1, \dots, C_{N_{\text{loop}}}\}$ are boundaries of plaques of the lattice (we assume we have the lattice version of the Hamiltonian); and N_{loop} is the total number of plaques of the lattice.

The minimum condition of $F[\nabla\chi]$ with respect to the variation of $\nabla\chi$ yields,

$$\frac{\delta E[\nabla\chi]}{\delta \nabla\chi} + \sum_{\ell=1}^{N_{\text{loop}}} \lambda_\ell \frac{\delta}{\delta \nabla\chi} \oint_{C_\ell} \nabla\chi \cdot d\mathbf{r} = 0 \quad (68)$$

Then, the current density is given by

$$\mathbf{J} = \frac{2e}{\hbar} \frac{\delta E}{\delta \nabla\chi} = -\frac{2e}{\hbar} \sum_{\ell=1}^{N_{\text{loop}}} \lambda_\ell \frac{\delta}{\delta \nabla\chi} \oint_{C_\ell} \nabla\chi \cdot d\mathbf{r} \quad (69)$$

This indicates that the current flowing in superconducting states is a collection of loop currents.

Expressing Eqs. (65) and (68) using the discrete lattice Hamiltonian, a system of equations for $\nabla\chi$ is obtained,

$$\sum_{k \leftarrow j} L_{k \leftarrow j}^\ell \tau_{k \leftarrow j} = 2\pi w_{C_\ell}[\chi] \quad (70)$$

$$\frac{\partial E(\{\tau_{k \leftarrow j}\})}{\partial \tau_{k \leftarrow j}} + \sum_{\ell=1}^{N_{\text{loop}}} \lambda_\ell \frac{\partial}{\partial \tau_{k \leftarrow j}} \sum_{k \leftarrow j} L_{k \leftarrow j}^\ell \tau_{k \leftarrow j} = 0 \quad (71)$$

where $k \leftarrow j$ indicates the bond that starts from site j and ends at site k , $\tau_{k \leftarrow j}$ is the difference of χ for the bond $k \leftarrow j$

$$\tau_{k \leftarrow j} = \chi_k - \chi_j \quad (72)$$

and $L_{k \leftarrow j}^\ell$ is defined as

$$L_{k \leftarrow j}^\ell = \begin{cases} -1 & \text{if } k \leftarrow j \text{ exists in } C_\ell \text{ in the clockwise direction} \\ 1 & \text{if } k \leftarrow j \text{ exists in } C_\ell \text{ in the counterclockwise direction} \\ 0 & \text{if } k \leftarrow j \text{ does not exist in } C_\ell \end{cases} \quad (73)$$

Note that a set of parameters $\{w_{C_\ell}[\chi]\}$ must be supplied as part of boundary conditions.

We take the branch of χ_j that satisfies the difference of value from the nearest neighbor site k is in the range,

$$-\pi \leq \chi_j - \chi_k < \pi \quad (74)$$

From $(\chi_j - \chi_k)$'s, we rebuild χ . The rebuilding process is as follows: first, we pick a value for the initial χ_1 (say $\chi_1 = 0$). After fixing the value of χ_1 , we calculate χ_2 by




FIG. 3. An example of a simply-connected lattice constructed from square lattice. 5×5 square lattice in **a** becomes a simply-connected lattice in **b** by removing 16 bonds.

$\chi_2 = \chi_1 + (\chi_2 - \chi_1)$, where the site 2 is connected to the site 1 by a nearest neighbor bond. The step where value χ_j is derived from the already evaluated value of χ_k is given by

$$\chi_j = \chi_k + (\chi_j - \chi_k) \quad (75)$$

where the sites j and k are connected by a bond in the path for the rebuilding of χ . This process is continued until values at all accessible sites are evaluated once and only once.

By this rebuilding process, a single path is constructed from the site 1 to other sites $k \neq 1$, which is achieved by making the region simply-connected by removing some bonds (see Fig. 3). We denote the path from the site 1 to other sites $k \neq 1$ by $C_{1 \rightarrow k}$. Then, the value χ_k is given by

$$\chi_k \approx \chi_1 + \int_{C_{1 \rightarrow k}} \nabla \chi \cdot d\mathbf{r} \quad (76)$$

When a magnetic field $\mathbf{B}^{\text{em}} = \nabla \times \mathbf{A}^{\text{em}}$ is applied, the energy functional in Eq. (67) is modified as

$$E[\nabla \chi] \rightarrow E \left[\nabla \chi - \frac{2e}{\hbar c} \mathbf{A}^{\text{em}} \right] \quad (77)$$

This leads to replace $\tau_{j \leftarrow i}$ in E by

$$\tau_{j \leftarrow i} \rightarrow \tau_{j \leftarrow i} - \frac{2e}{\hbar c} \int_i^j \mathbf{A}^{\text{em}} \cdot d\mathbf{r} \quad (78)$$

By this modification, a magnetic field can be taken into account.

Lastly in this section, we would like to consider the mass in the London moment. In the present formalism, the velocity field \mathbf{v} is given by

$$\mathbf{v} = \frac{e}{m_e c} \mathbf{A}^{\text{em}} - \frac{\hbar}{2m_e} \nabla \chi \quad (79)$$

since the velocity field from \tilde{u}_n, \tilde{v}_n is zero. This indicates that m in Eq. (1) is m_e in agreement with the value observed in the London moment.

III. MODEL CALCULATIONS

A. Model Hamiltonian

In this section, a model is solved using the particle number conserving version of the Bogoliubov-de Gennes equations. This model is originally built to study the cuprate superconductivity. Since the purpose of the present work is to explain the new theory, the model calculation below should be considered as an example exercise. The validity of the model as a model for the cuprate superconductivity will be dealt elsewhere.

The model is composed of two layers, a surface layer and a bulk layer (see Fig. 4). The superconductivity occurs in the bulk, however, many experimental results show contributions from the surface region, thus, the surface layer effects must be included to interpret experiments. The experimental results seem to indicate that the small polaron formation and coexisting magnetic moment effects are strong in the bulk, however, they are weak in the surface region.

Thus, we developed a model containing two layers to reproduce experimental results. It is also notable that although the electron-pairing does not occur in the bulk layer, supercurrent flows there in this model; thus, it demonstrates that the primary importance for superconductivity is the appearance of the non-trivial Berry connection.

B. Bulk layer

The model for the bulk layer is constructed based on the fact that the parent compound is a Mott insulator [60], which is well-described by the two dimensional Hubbard model with large on-site repulsion

$$U \gg t_1 \tag{80}$$

where U is the on-site repulsion parameter and t_1 is the nearest neighbor hopping parameter. The bulk CuO_2 plane of the cuprate is taken as a two-dimensional square lattice, where a




FIG. 4. Simply-connected lattice constructed by removing some bonds for the two-layer model. We consider the model with two 7×7 square lattices stacked in the z direction. There are four small polarons in the bulk layer indicated by “M” and “A”. Each “M” indicates a center of a spin-vortex with winding number $+1$, and “A” a center of a spin-vortex with winding number -1 .

copper atom resides at each site. Oxygens between copper atoms are not explicitly taken into account in the present model.

When holes are doped, they form small polarons in the bulk [25, 61]. At low temperatures, the mobility of the small polaron is very small; thus, the supercurrent is not due to the hopping of small polarons.

The inelastic neutron scattering experiments indicate spin-wave excitations with spin lying in CuO_2 plane [23]. Thus, the magnetic moment coexisting with superconductivity in the bulk. The magnetic excitation spectrum has a hourglass-shaped dispersion, suggesting the stripes of spins or spin-vortices [26, 62].

By taking into account the above observation, the following bulk Hamiltonian is constructed,

$$\begin{aligned}
H_{\text{EHFS}} = & -t_1 \sum_{\langle i,j \rangle_{1,\sigma}} [c_{i\sigma}^\dagger c_{j\sigma} + \text{H.c.}] + U \sum_j c_{j\uparrow}^\dagger c_{j\uparrow} c_{j\downarrow}^\dagger c_{j\downarrow} + J_h \sum_{\langle i,j \rangle_h} \mathbf{S}_i \cdot \mathbf{S}_j \\
& + \lambda \sum_h [c_{h+y\downarrow}^\dagger c_{h-x\uparrow} - c_{h+y\uparrow}^\dagger c_{h-x\downarrow} + c_{h+x\downarrow}^\dagger c_{h-y\uparrow} - c_{h+x\uparrow}^\dagger c_{h-y\downarrow} \\
& + i(c_{h-x\downarrow}^\dagger c_{h-y\uparrow} + c_{h-x\uparrow}^\dagger c_{h-y\downarrow}) + i(c_{h+y\downarrow}^\dagger c_{h+x\uparrow} + c_{h+y\uparrow}^\dagger c_{h+x\downarrow}) + \text{h.c.}] \\
& - \mu_{\text{bulk}} \sum_{j,\sigma} c_{j\sigma}^\dagger c_{j\sigma}
\end{aligned} \tag{81}$$

where $c_{j\sigma}^\dagger$ and $c_{j\sigma}$ are creation and annihilation operators for the electron at j th site with spin σ , respectively.

In this Hamiltonian, the doped holes are assumed to form small polarons and immobile, thus, the effectively-half-filled situation (EHFS) is realized, where the number of electrons and the number of accessible sites for the electron are equal. We take the hole-occupied sites as inaccessible sites for electrons, thus, not included in the sum.

The first term describes the electron hopping with transfer integral t_1 ; $\langle i, j \rangle_1$ indicates that the sum is taken over 1st nearest neighbor hoppings.

The second term describes the on-site Coulomb repulsion; U is significantly larger than t_1 (actually, we adopt $U = 8t_1$ in the present work).

The third term describes the antiferromagnetic exchange interaction between electron spins around each doped hole; $\langle i, j \rangle_h$ indicates that the sum is taken over pairs around holes, i.e., each hole (denoted by h) accompanies four sites around it (denoted by $h-x, h+x, h-y$ and $h+y$; they are the nearest neighbor sites of h in $-x, +x, -y$, and $+y$ directions, respectively), and there are six site-pairs that contribute to $\langle i, j \rangle_h$ for each doped hole. The parameter J_h is taken to be $0.5J_{\text{AF}}$ where $J_{\text{AF}} = \frac{4t_1^2}{U}$ is the antiferromagnetic exchange parameter for the parent compound [63].

The fourth term is the Rashba interaction term [64], which we only include around the holes [65]. The fifth term is the chemical potential term.

Components of the spin operator $\mathbf{S}_j = (S_j^x, S_j^y, S_j^z)$ are given by

$$\begin{aligned}
S_j^x &= \frac{1}{2}(c_{j\uparrow}^\dagger c_{j\downarrow} + c_{j\downarrow}^\dagger c_{j\uparrow}) \\
S_j^y &= \frac{i}{2}(-c_{j\uparrow}^\dagger c_{j\downarrow} + c_{j\downarrow}^\dagger c_{j\uparrow}) \\
S_j^z &= \frac{1}{2}(c_{j\uparrow}^\dagger c_{j\uparrow} - c_{j\downarrow}^\dagger c_{j\downarrow})
\end{aligned} \tag{82}$$

The chemical potential μ_{bulk} is chosen to control the number of small polarons in EHFS. Since H_{EHFS} is too difficult to handle as it is, we use a mean field version of it

$$\begin{aligned}
H_{\text{EHFS}}^{\text{HF}} = & -t_1 \sum_{\langle i,j \rangle_{1,\sigma}} [c_{i\sigma}^\dagger c_{j\sigma} + \text{H.c.}] \\
& + U \sum_j [(-\frac{2}{3}\langle S_j^z \rangle + \frac{1}{2})c_{j\uparrow}^\dagger c_{j\uparrow} + (\frac{2}{3}\langle S_j^z \rangle + \frac{1}{2})c_{j\downarrow}^\dagger c_{j\downarrow} \\
& - \frac{2}{3}(\langle S_j^x \rangle - i\langle S_j^y \rangle)c_{j\uparrow}^\dagger c_{j\downarrow} - \frac{2}{3}(\langle S_j^x \rangle + i\langle S_j^y \rangle)c_{j\downarrow}^\dagger c_{j\uparrow} - \frac{2}{3}\langle \mathbf{S}_j \rangle^2] \\
& + J_h \sum_{\langle i,j \rangle_h} [\frac{1}{2}(\langle S_i^z \rangle c_{j\uparrow}^\dagger c_{j\uparrow} + \langle S_j^z \rangle c_{i\uparrow}^\dagger c_{i\uparrow}) - \frac{1}{2}(\langle S_i^z \rangle c_{j\downarrow}^\dagger c_{j\downarrow} + \langle S_j^z \rangle c_{i\downarrow}^\dagger c_{i\downarrow}) \\
& + \frac{1}{2}\{(\langle S_i^x \rangle - i\langle S_i^y \rangle)c_{j\uparrow}^\dagger c_{j\downarrow} + (\langle S_j^x \rangle - i\langle S_j^y \rangle)c_{i\uparrow}^\dagger c_{i\downarrow}\} \\
& + \frac{1}{2}\{(\langle S_i^x \rangle + i\langle S_i^y \rangle)c_{j\downarrow}^\dagger c_{j\uparrow} + (\langle S_j^x \rangle + i\langle S_j^y \rangle)c_{i\downarrow}^\dagger c_{i\uparrow}\} - \langle \mathbf{S}_i \rangle \cdot \langle \mathbf{S}_j \rangle] \\
& + \lambda \sum_h [c_{h+y\downarrow}^\dagger c_{h-x\uparrow} - c_{h+y\uparrow}^\dagger c_{h-x\downarrow} + c_{h+x\downarrow}^\dagger c_{h-y\uparrow} - c_{h+x\uparrow}^\dagger c_{h-y\downarrow} \\
& + i(c_{h-x\downarrow}^\dagger c_{h-y\uparrow} + c_{h-x\uparrow}^\dagger c_{h-y\downarrow}) + i(c_{h+y\downarrow}^\dagger c_{h+x\uparrow} + c_{h+y\uparrow}^\dagger c_{h+x\downarrow}) + \text{h.c.}] \\
& - \mu_{\text{bulk}} \sum_{j,\sigma} c_{j\sigma}^\dagger c_{j\sigma} \tag{83}
\end{aligned}$$

where $\langle \hat{O} \rangle$ denotes the expectation value of the operator \hat{O} . This Hamiltonian yields states with spin-vortices around small polarons (see Fig.5). These spin-vortices induce loop currents called the “*spin-vortex-induced loop currents*”, due to the appearance of a non-trivial Berry connection.

C. Surface layer

The Hamiltonian for the surface layer is following

$$\begin{aligned}
H_{\text{surf}} = & -t_1 \sum_{\langle i,j \rangle_{1,\sigma}} [c_{i\sigma}^\dagger c_{j\sigma} + \text{H.c.}] - t_2 \sum_{\langle i,j \rangle_{2,\sigma}} [c_{i\sigma}^\dagger c_{j\sigma} + \text{H.c.}] + U \sum_j c_{j\uparrow}^\dagger c_{j\uparrow} c_{j\downarrow}^\dagger c_{j\downarrow} - \mu_{\text{surf}} \sum_{j,\sigma} c_{j\sigma}^\dagger c_{j\sigma} \tag{84}
\end{aligned}$$

This is the Hubbard model including the second nearest neighbor hopping. The second nearest neighbor hopping with the parameter t_2 is included to have a curved Fermi surface in the angle-resolved photoemission spectrum. We use $t_2 = -0.12t_1$ in this work.

The most important difference from the bulk Hamiltonian is the absence of the small polaron effect. The spin-orbit interaction term is also absent. The chemical potential μ_{surf} is so chosen that the bulk and surface hole densities are equal.




FIG. 5. Spin texture obtained. **a**: The spin-texture of the two layer. The spins in the surface layer are very small. **b**: The normalized spin-textures in the bulk (top), and that in the surface (bottom). Spin-vortices are created in the surface layer due to the influence of those in the bulk layer.

Since H_{surf} is too difficult to handle as it is, we construct a mean-field version via the $t - J$ model [66] for the Hubbard model,

$$\begin{aligned}
H_{\text{surf}}^{\text{HF}} = & -t_1 \sum_{\langle i,j \rangle_{1,\sigma}} [(1 - \langle n_{i,-\sigma} \rangle) c_{i\sigma}^\dagger c_{j\sigma} (1 - \langle n_{j,-\sigma} \rangle) + \text{H.c.}] \\
& - t_2 \sum_{\langle i,j \rangle_{2,\sigma}} [(1 - \langle n_{i,-\sigma} \rangle) c_{i\sigma}^\dagger c_{j\sigma} (1 - \langle n_{j,-\sigma} \rangle) + \text{H.c.}] \\
& + \sum_{\langle i,j \rangle_1} (\Delta_{ij} e^{-\frac{i}{2}\hat{x}_j} e^{-\frac{i}{2}\hat{x}_i} c_{i\uparrow}^\dagger c_{j\downarrow}^\dagger + \text{H.c.}) - \mu_{\text{surf}} \sum_{j,\sigma} c_{j\sigma}^\dagger c_{j\sigma}
\end{aligned} \tag{85}$$

where the pairing amplitude or pair potential Δ_{ij} is defined as

$$\Delta_{ij} = -\frac{4t_1^2}{U} \langle e^{\frac{i}{2}\hat{x}_j} e^{\frac{i}{2}\hat{x}_i} (c_{j\downarrow} c_{i\uparrow} - c_{j\uparrow} c_{i\downarrow}) \rangle = \Delta_{ji} \tag{86}$$

Due to the presence of the number changing operators, $e^{\frac{i}{2}\hat{x}_j}$ and $e^{\frac{i}{2}\hat{x}_i}$, Δ_{ij} is obtained in the particle number conserving formalism. The appearance of the number changing operator is due to the spin-vortex formation in the bulk; thus, the electron pair formation is a secondary effect. To have the superconducting state, stable loop currents that can generate a macroscopic persistent current should be realized.

The pair potential Δ_{ij} gives rise to d -wave pairing gap, and explains the Fermi arc [67]. We would like to emphasize that the appearance of the d -wave pairing gap is secondary effect enabled by the appearance of the non-trivial Berry connection in the bulk; this generates the collective mode for supercurrent, and particle number changing operators. The particle number changing operators make it possible to gain energy by the d -wave pairing gap formation.

D. Interlayer hopping

The bulk and surface layers are connected by the inter-layer hopping. The following is the inter-layer hopping Hamiltonian,

$$H_{\text{inter-layer}} = -t_3 \sum_{\langle i_s, j_b \rangle_{z, \sigma}} [(1 - n_{i_s, -\sigma}) c_{i_s \sigma}^\dagger c_{j_b \sigma} (1 - n_{j_b, -\sigma}) + \text{H.c.}] + \frac{4t_3^2}{U} \sum_{\langle i_s, j_b \rangle_z} \left(\mathbf{S}_{i_s} \cdot \mathbf{S}_{j_b} - \frac{1}{4} n_{i_s} n_{j_b} \right) \quad (87)$$

where an antiferromagnetic exchange interaction between spins in the bulk-layer and those in the surface-layer is included using a $t - J$ model approximation.

Since $H_{\text{inter-layer}}$ is still too difficult to handle, we use the following mean field version,

$$\begin{aligned} H_{\text{inter-layer}}^{\text{HF}} &= -t_3 \sum_{\langle i_s, j_b \rangle_{z, \sigma}} [(1 - \langle n_{i_s, -\sigma} \rangle) c_{i_s \sigma}^\dagger c_{j_b \sigma} (1 - \langle n_{j_b, -\sigma} \rangle) + \text{H.c.}] \\ &+ \frac{2t_3^2}{U} \sum_{\langle i_s, j_b \rangle_z} [(\langle S_{i_s}^x \rangle - i \langle S_{i_s}^y \rangle) c_{j_b \uparrow}^\dagger c_{j_b \downarrow} + (\langle S_{i_s}^x \rangle + i \langle S_{i_s}^y \rangle) c_{j_b \uparrow}^\dagger c_{j_b \downarrow} \\ &+ (\langle S_{i_s}^z \rangle - \frac{1}{2} \langle n_{i_s} \rangle) c_{j_b \uparrow}^\dagger c_{j_b \uparrow} - (\langle S_{i_s}^z \rangle + \frac{1}{2} \langle n_{i_s} \rangle) c_{j_b \downarrow}^\dagger c_{j_b \downarrow}] \\ &+ \frac{2t_3^2}{U} \sum_{\langle i_s, j_b \rangle_z} [(\langle S_{j_b}^x \rangle - i \langle S_{j_b}^y \rangle) c_{i_s \uparrow}^\dagger c_{i_s \downarrow} + (\langle S_{j_b}^x \rangle + i \langle S_{j_b}^y \rangle) c_{i_s \uparrow}^\dagger c_{i_s \downarrow} \\ &+ (\langle S_{j_b}^z \rangle - \frac{1}{2} \langle n_{j_b} \rangle) c_{i_s \uparrow}^\dagger c_{i_s \uparrow} - (\langle S_{j_b}^z \rangle + \frac{1}{2} \langle n_{j_b} \rangle) c_{i_s \downarrow}^\dagger c_{i_s \downarrow}] \\ &- \frac{4t_3^2}{U} \sum_{\langle i_s, j_b \rangle_z} \left(\langle \mathbf{S}_{i_s} \rangle \cdot \langle \mathbf{S}_{j_b} \rangle - \frac{1}{4} \langle n_{i_s} \rangle \langle n_{j_b} \rangle \right) \end{aligned} \quad (88)$$

We adopt $t_3 = 0.01t_1$ in the following calculations. Through this Hamiltonian, the non-trivial Berry connection generated in the bulk propagates in the surface.

E. The total Hamiltonian for the two-layer model

Over all, our model Hamiltonian is given by

$$\begin{aligned}
H_{\text{eff}} &= H_{\text{EHFS}}^{\text{HF}} + H_{\text{surf}}^{\text{HF}} + H_{\text{inter-layer}}^{\text{HF}} \\
&= \sum_{i,j,\sigma,\sigma'} h_{i\sigma,j\sigma'} c_{i\sigma}^\dagger c_{j\sigma'} + \sum_{i,j} [\Delta_{ij} e^{-\frac{i}{2}\hat{\chi}_j} e^{-\frac{i}{2}\hat{\chi}_i} c_{i\uparrow}^\dagger c_{j\downarrow}^\dagger + \Delta_{ij}^* e^{\frac{i}{2}\hat{\chi}_j} e^{\frac{i}{2}\hat{\chi}_i} c_{j\downarrow} c_{i\uparrow}] + E_{\text{const}}
\end{aligned} \tag{89}$$

F. Particle number conserving Bogoliubov-de Gennes equations

In order to obtain the particle number conserving Bogoliubov-de Gennes equations from H_{eff} in Eq. (89), we perform the following Bogoliubov transformation

$$\begin{aligned}
c_{i\uparrow} &= \sum_n^l [u_{i\uparrow}^n \gamma_n - (v_{i\uparrow}^n)^* \gamma_n^\dagger] e^{-\frac{i}{2}\hat{\chi}_i} \\
c_{i\downarrow} &= \sum_n^l [u_{i\downarrow}^n \gamma_n + (v_{i\downarrow}^n)^* \gamma_n^\dagger] e^{-\frac{i}{2}\hat{\chi}_i}
\end{aligned} \tag{90}$$

Using the Bogoliubov operators γ_n and γ_n^\dagger , H_{eff} is expressed as

$$H_{\text{eff}} = \sum_n^l E_n \gamma_n^\dagger \gamma_n + E'_{\text{const.}} \tag{91}$$

where $E'_{\text{const.}}$ is a constant and “ \sum_n^l ” denotes that the sum is take over $E_n > 0$, and the ground state $|\text{Gnd}\rangle$ satisfies

$$\gamma_n |\text{Gnd}\rangle = 0, \quad E_n > 0 \tag{92}$$

The Bogoliubov operators are fermion operators that satisfy

$$\{\gamma_n, \gamma_m^\dagger\} = \delta_{nm}, \quad \{\gamma_n^\dagger, \gamma_m^\dagger\} = 0, \quad \{\gamma_n, \gamma_m\} = 0 \tag{93}$$

thus, the commutation relations

$$[\gamma_n^\dagger, H_{\text{eff}}] = -E_n \gamma_n^\dagger, \quad [\gamma_n, H_{\text{eff}}] = E_n \gamma_n \tag{94}$$

are obtained.

The following commutation relations are obtained between H_{eff} and $c_{i\sigma}$, $c_{i\sigma}^\dagger$,

$$[c_{i\uparrow}, H_{\text{eff}}] = \sum_{j,\sigma'} h_{i\uparrow,j\sigma'} c_{j\sigma'} + \sum_j \Delta_{ij} e^{-\frac{i}{2}\hat{\chi}_j} e^{-\frac{i}{2}\hat{\chi}_i} c_{j\downarrow}^\dagger$$

$$\begin{aligned}
[c_{i\uparrow}^\dagger, H_{\text{eff}}] &= -\sum_{j,\sigma'} h_{j\sigma',i\uparrow} c_{j\sigma'}^\dagger - \sum_j \Delta_{ij}^* e^{\frac{i}{2}\hat{\chi}_j} e^{\frac{i}{2}\hat{\chi}_i} c_{j\downarrow} \\
[c_{i\downarrow}, H_{\text{eff}}] &= \sum_{j,\sigma'} h_{i\downarrow,j\sigma'} c_{j\sigma'} - \sum_j \Delta_{ji} e^{-\frac{i}{2}\hat{\chi}_j} e^{-\frac{i}{2}\hat{\chi}_i} c_{j\uparrow}^\dagger \\
[c_{i\downarrow}^\dagger, H_{\text{eff}}] &= -\sum_{j,\sigma'} h_{j\sigma',i\downarrow} c_{j\sigma'}^\dagger + \sum_j \Delta_{ji}^* e^{\frac{i}{2}\hat{\chi}_j} e^{\frac{i}{2}\hat{\chi}_i} c_{j\uparrow}
\end{aligned} \tag{95}$$

Using Eqs. (95), (90), and (94), the Bogoliubov-de Gennes equations are obtained,

$$\begin{aligned}
E_n u_{i\uparrow}^n &= \sum_{j\sigma'} e^{\frac{i}{2}\hat{\chi}_i} h_{i\uparrow,j\sigma'} e^{-\frac{i}{2}\hat{\chi}_j} u_{j\sigma'}^n + \sum_j \Delta_{ij} v_{j\downarrow}^n \\
E_n u_{i\downarrow}^n &= \sum_{j\sigma'} e^{\frac{i}{2}\hat{\chi}_i} h_{i\downarrow,j\sigma'} e^{-\frac{i}{2}\hat{\chi}_j} u_{j\sigma'}^n + \sum_j \Delta_{ji} v_{j\uparrow}^n \\
E_n v_{i\uparrow}^n &= -\sum_j e^{-\frac{i}{2}\hat{\chi}_i} h_{i\uparrow,j\uparrow}^* e^{\frac{i}{2}\hat{\chi}_j} v_{j\uparrow}^n + \sum_j e^{-\frac{i}{2}\hat{\chi}_i} h_{i\uparrow,j\downarrow}^* e^{\frac{i}{2}\hat{\chi}_j} v_{j\downarrow}^n + \sum_j \Delta_{ij}^* u_{j\downarrow}^n \\
E_n v_{i\downarrow}^n &= \sum_j e^{-\frac{i}{2}\hat{\chi}_i} h_{i\downarrow,j\uparrow}^* e^{\frac{i}{2}\hat{\chi}_j} v_{j\uparrow}^n - \sum_j e^{-\frac{i}{2}\hat{\chi}_i} h_{i\downarrow,j\downarrow}^* e^{\frac{i}{2}\hat{\chi}_j} v_{j\downarrow}^n + \sum_j \Delta_{ji}^* u_{j\uparrow}^n
\end{aligned} \tag{96}$$

In order to solve the above system of equations, we replace the operator $\hat{\chi}_j$ by its associated scalar value χ_j . Then, the Bogoliubov-de Gennes equations is put into the following matrix form,

$$\sum_j M_{ij} \phi_j^n = E_n \phi_i^n \tag{97}$$

where

$$M_{ij} = \begin{pmatrix} h_{i\uparrow,j\uparrow} e^{\frac{i}{2}(\chi_i - \chi_j)} & h_{i\uparrow,j\downarrow} e^{\frac{i}{2}(\chi_i - \chi_j)} & 0 & \Delta_{ij} \\ h_{i\downarrow,j\uparrow} e^{\frac{i}{2}(\chi_i - \chi_j)} & h_{i\downarrow,j\downarrow} e^{\frac{i}{2}(\chi_i - \chi_j)} & \Delta_{ji} & 0 \\ 0 & \Delta_{ij}^* & -h_{i\uparrow,j\uparrow}^* e^{-\frac{i}{2}(\chi_i - \chi_j)} & h_{i\uparrow,j\downarrow}^* e^{-\frac{i}{2}(\chi_i - \chi_j)} \\ \Delta_{ji}^* & 0 & h_{i\downarrow,j\uparrow}^* e^{-\frac{i}{2}(\chi_i - \chi_j)} & -h_{i\downarrow,j\downarrow}^* e^{-\frac{i}{2}(\chi_i - \chi_j)} \end{pmatrix} \tag{98}$$

and

$$\phi_i^n = \begin{pmatrix} u_{i\uparrow}^n \\ u_{i\downarrow}^n \\ v_{i\uparrow}^n \\ v_{i\downarrow}^n \end{pmatrix} \tag{99}$$

Note that solutions for E_n and $-E_n$ are connected; the solution for $-E_n$ is given by

$$\begin{pmatrix} -(v_{i\uparrow}^n)^* \\ (v_{i\downarrow}^n)^* \\ -(u_{i\uparrow}^n)^* \\ (u_{i\downarrow}^n)^* \end{pmatrix} \tag{100}$$

Therefore, we only need to obtain half of the solutions to Eq. (97).

Self-consistent solutions are obtained by solving the above equations using the self-consistent fields,

$$\begin{aligned}
\langle n_{i\uparrow} \rangle &= \langle c_{i\uparrow}^\dagger c_{i\uparrow} \rangle = \sum'_{n_1} \sum'_{n_2} \langle e^{\frac{i}{2}\hat{X}_i} [u_{i\uparrow}^{n_1} \gamma_{n_1} - (v_{i\uparrow}^{n_1})^* \gamma_{n_1}^\dagger]^\dagger [u_{i\uparrow}^{n_2} \gamma_{n_2} - (v_{i\uparrow}^{n_2})^* \gamma_{n_2}^\dagger] e^{-\frac{i}{2}\hat{X}_i} \rangle \\
&= \sum'_{n_1} \sum'_{n_2} \langle e^{\frac{i}{2}\hat{X}_i} [(u_{i\uparrow}^{n_1})^* \gamma_{n_1}^\dagger - (v_{i\uparrow}^{n_1}) \gamma_{n_1}] [u_{i\uparrow}^{n_2} \gamma_{n_2} - (v_{i\uparrow}^{n_2})^* \gamma_{n_2}^\dagger] e^{-\frac{i}{2}\hat{X}_i} \rangle \\
&= \sum'_{n_1} \sum'_{n_2} [(u_{i\uparrow}^{n_1})^* u_{i\uparrow}^{n_2} \langle \gamma_{n_1}^\dagger \gamma_{n_2} \rangle - (u_{i\uparrow}^{n_1})^* (v_{i\uparrow}^{n_2})^* \langle \gamma_{n_1}^\dagger \gamma_{n_2}^\dagger \rangle + (v_{i\uparrow}^{n_1}) u_{i\uparrow}^{n_2} \langle \gamma_{n_1} \gamma_{n_2} \rangle + (v_{i\uparrow}^{n_1}) (v_{i\uparrow}^{n_2})^* \langle \gamma_{n_1} \gamma_{n_2}^\dagger \rangle] \\
&= \sum'_{n_1} [|u_{i\uparrow}^{n_1}|^2 f(E_{n_1}) + |v_{i\uparrow}^{n_1}|^2 f(-E_{n_1})] = \sum_n |u_{i\uparrow}^n|^2 f(E_n) = \sum_n |v_{i\uparrow}^n|^2 f(-E_n) \\
\langle n_{i\downarrow} \rangle &= \sum'_{n_1} [|u_{i\downarrow}^{n_1}|^2 f(E_{n_1}) + |v_{i\downarrow}^{n_1}|^2 f(-E_{n_1})] = \sum_n |u_{i\downarrow}^n|^2 f(E_n) = \sum_n |v_{i\downarrow}^n|^2 f(-E_n) \tag{101}
\end{aligned}$$

$$\begin{aligned}
\langle S_i^x \rangle &= -\frac{1}{2} \sum'_n [v_{i\uparrow}^n (v_{i\downarrow}^n)^* + \text{c.c.}] f(-E_n) + \frac{1}{2} \sum'_n [u_{i\uparrow}^n (u_{i\downarrow}^n)^* + \text{c.c.}] f(E_n) \\
\langle S_i^y \rangle &= \frac{i}{2} \sum'_n [v_{i\uparrow}^n (v_{i\downarrow}^n)^* - \text{c.c.}] f(-E_n) - \frac{i}{2} \sum'_n [u_{i\uparrow}^n (u_{i\downarrow}^n)^* - \text{c.c.}] f(E_n) \\
\langle S_i^z \rangle &= \frac{1}{2} \sum'_n [v_{i\uparrow}^n (v_{i\uparrow}^n)^* - v_{i\downarrow}^n (v_{i\downarrow}^n)^*] f(-E_n) + \frac{1}{2} \sum'_n [u_{i\uparrow}^n (u_{i\uparrow}^n)^* - u_{i\downarrow}^n (u_{i\downarrow}^n)^*] f(E_n) \tag{102}
\end{aligned}$$

and

$$\Delta_{ij} = \frac{2t_1^2}{U} \sum'_n [u_{i\uparrow}^n (v_{j\downarrow}^n)^* + u_{j\downarrow}^n (v_{i\uparrow}^n)^* + u_{i\downarrow}^n (v_{j\uparrow}^n)^* + u_{j\uparrow}^n (v_{i\downarrow}^n)^*] \tanh \frac{E_n}{2k_B T} \tag{103}$$

where following relations are used

$$\langle \gamma_n^\dagger \gamma_n \rangle = f(E_n), \quad \langle \gamma_n \gamma_n^\dagger \rangle = 1 - f(E_n) = f(-E_n), \tag{104}$$

We solve the system of equations in Eq. (97) by the Car-Parrinello method [68], taking ϕ_i^n as “time-dependent” variable $\phi^n(t)$ at the i th site that follows the Newtonian dynamics

$$m_{\text{CP}} \ddot{\phi}_i^n(t) = - \sum_j M_{ij} \phi_j^n(t) + E_n \phi_i^n(t) - \eta_{\text{CP}} \dot{\phi}_i^n(t) \tag{105}$$

where m_{CP} and η_{CP} are mass and friction coefficient for the variable, respectively; here, the wave functions $\{\phi^n(t)\}$ are orthonormalized during the calculation.

G. Spin-vortices and multi-valued wave functions

The self-consistent wave functions obtained in the previous section, actually, become multi-valued with respect to electron coordinates due to the presence of spin-vortices. The spin-texture obtained for a model system composed of the two-layers shown in Fig. 4 is depicted in Fig. 5.

If $u_{i\sigma}^n$ and $v_{i\sigma}^n$ are obtained just by minimizing energy, the resulting state is a currentless state that satisfies the so-called ‘‘Bloch’s theorem’’ [69]. This corresponds to the solution with constant χ .

Let us examine this currentless state, more closely. Replacing the operator $\hat{\chi}_i$ in Eq. (90) by the scalar value of the underlying Berry connection, we obtain

$$\begin{pmatrix} c_{i\uparrow} \\ c_{i\downarrow} \end{pmatrix} = e^{-\frac{i}{2}\chi_i} \sum_n \begin{pmatrix} u_{i\uparrow}^n & -(v_{i\uparrow}^n)^* \\ u_{i\downarrow}^n & (v_{i\downarrow}^n)^* \end{pmatrix} \begin{pmatrix} \gamma_n \\ \gamma_n^\dagger \end{pmatrix} \quad (106)$$

and the currentless result corresponds to the solution with constant χ_j . Since the single-valued solution corresponds to multi-valued χ_j , the currentless $u_{i\sigma}^n$ and $v_{i\sigma}^n$ are actually multi-valued.

When the bulk Hamiltonian yields spin-vortices lying in the xy plane (in the CuO_2 plane), the on-site U term in $H_{\text{EHFS}}^{\text{HF}}$ in Eq. (83) becomes

$$U \sum_j \left[\frac{1}{2} (c_{j\uparrow}^\dagger c_{j\uparrow} + c_{j\downarrow}^\dagger c_{j\downarrow}) - \frac{2}{3} |\langle \mathbf{S}_j \rangle| (e^{-i\xi_j} c_{j\uparrow}^\dagger c_{j\downarrow} + e^{i\xi_j} c_{j\downarrow}^\dagger c_{j\uparrow}) - \frac{2}{3} \langle \mathbf{S}_j \rangle^2 \right] \quad (107)$$

We can absorb the spin-twisting factor $e^{\pm i\xi_j}$ using $\tilde{c}_{j\uparrow}^\dagger, \tilde{c}_{j\downarrow}$ as

$$c_{j\uparrow}^\dagger = e^{\frac{i}{2}\xi_j} \tilde{c}_{j\uparrow}^\dagger, \quad c_{j\downarrow} = e^{\frac{i}{2}\xi_j} \tilde{c}_{j\downarrow} \quad (108)$$

Then, Eq. (106) becomes

$$\begin{pmatrix} \tilde{c}_{i\uparrow} \\ \tilde{c}_{i\downarrow} \end{pmatrix} = \sum_n \begin{pmatrix} \tilde{u}_{i\uparrow}^n & -(\tilde{v}_{i\uparrow}^n)^* \\ \tilde{u}_{i\downarrow}^n & (\tilde{v}_{i\downarrow}^n)^* \end{pmatrix} \begin{pmatrix} \gamma_n \\ \gamma_n^\dagger \end{pmatrix} \quad (109)$$

This is essentially equivalent to the solution with constant χ in Eq. (106). This means that $\tilde{u}_{i\sigma}^n$ and $\tilde{v}_{i\sigma}^n$ are obtained from the energy minimization only calculation if $c_{j\uparrow}^\dagger, c_{j\downarrow}$ are replaced by $\tilde{c}_{j\uparrow}^\dagger, \tilde{c}_{j\downarrow}$. We do this in the following.

We obtain $\tilde{u}_{i\sigma}^n$ and $\tilde{v}_{i\sigma}^n$ from the following Hamiltonian, in which $c_{j\sigma}^\dagger$ and $c_{j\sigma}$ are replaced by $\tilde{c}_{j\sigma}^\dagger, \tilde{c}_{j\sigma}$,

$$\sum_j \tilde{M}_{ij} \tilde{\phi}_j^n = \tilde{E}_n \tilde{\phi}_i^n \quad (110)$$

where \tilde{M}_{ij} and $\tilde{\phi}_i^n$ are given by

$$\tilde{M}_{ij} = \begin{pmatrix} h_{i\uparrow,j\uparrow} e^{\frac{i}{2}(\xi_i - \xi_j)} & h_{i\uparrow,j\downarrow} e^{-\frac{i}{2}(\xi_i + \xi_j)} & 0 & \Delta_{ij} e^{\frac{i}{2}(\xi_i - \xi_j)} \\ h_{i\downarrow,j\uparrow} e^{-\frac{i}{2}(\xi_i + \xi_j)} & h_{i\downarrow,j\downarrow} e^{-\frac{i}{2}(\xi_i - \xi_j)} & \Delta_{ji} e^{-\frac{i}{2}(\xi_i - \xi_j)} & 0 \\ 0 & \Delta_{ij}^* e^{-\frac{i}{2}(\xi_i - \xi_j)} & -h_{i\uparrow,j\uparrow}^* e^{-\frac{i}{2}(\xi_i - \xi_j)} & h_{i\uparrow,j\downarrow}^* e^{-\frac{i}{2}(\xi_i + \xi_j)} \\ \Delta_{ji}^* e^{\frac{i}{2}(\xi_i - \xi_j)} & 0 & h_{i\downarrow,j\uparrow}^* e^{\frac{i}{2}(\xi_i + \xi_j)} & -h_{i\downarrow,j\downarrow}^* e^{\frac{i}{2}(\xi_i - \xi_j)} \end{pmatrix} \quad (111)$$

and

$$\tilde{\phi}_i^n = \begin{pmatrix} \tilde{u}_{i\uparrow}^n \\ \tilde{u}_{i\downarrow}^n \\ \tilde{v}_{i\uparrow}^n \\ \tilde{v}_{i\downarrow}^n \end{pmatrix} \quad (112)$$

respectively. Note that $\tilde{u}_{i\sigma}^n, \tilde{v}_{i\sigma}^n$ are single-valued since the multi-valued part $e^{\pm\frac{i}{2}\xi}$ are removed.

H. Evaluation of Berry connection

Using $\tilde{u}_{i\sigma}^n, \tilde{v}_{i\sigma}^n$, Eq. (106) is rewritten as

$$\begin{pmatrix} c_{i\uparrow} \\ c_{i\downarrow} \end{pmatrix} = \sum_n \begin{pmatrix} e^{-\frac{i}{2}(\chi_i + \xi_i)} \tilde{u}_{i\uparrow}^n & -e^{-\frac{i}{2}(\chi_i + \xi_i)} (\tilde{v}_{i\uparrow}^n)^* \\ e^{-\frac{i}{2}(\chi_i - \xi_i)} \tilde{u}_{i\downarrow}^n & e^{-\frac{i}{2}(\chi_i - \xi_i)} (\tilde{v}_{i\downarrow}^n)^* \end{pmatrix} \begin{pmatrix} \gamma_n \\ \gamma_n^\dagger \end{pmatrix} \quad (113)$$

Now, we need to obtain the multi-valued χ . The angular variable χ is obtained from conditions,

$$w_{C_\ell}[\xi] + w_{C_\ell}[\chi] = \text{even number for any loop } C_\ell \quad (114)$$

If the above conditions are satisfied, $e^{-\frac{i}{2}(\chi_i \pm \xi_i)}$ appeared in Eq. (113) become single-valued.

Due to the presence of spin-vortices ξ is multi-valued, and contains jump-of-value points of an integral multiple of 2π . To establish the condition in Eq. (114), ξ and χ must have the same jump-of-value points.

To have the same jump-of-value points for ξ and χ , we do the following: first, we rebuild ξ from differences of their values between bonds; ξ_j 's are obtained from the expectation values of the spin components S_j^x and S_j^y ; we choose a particular branch for ξ_j . The system we consider below has the antiferromagnetic background; this gives rise to the background




FIG. 6. Loops used to impose boundary conditions in Eq. (114). **a**: loops in the bulk and surface layers. **a**: loops in the bulk-surface interface region. Some of the loops in the system are removed since they are not independent; as a consequence, all the cells are accessible from outside without breaking any walls framed by loops.

value of ξ , $\xi_j^0 = \pi(j_x + j_y + j_z)$ in the units the lattice constant, where (j_x, j_y, j_z) is the xyz coordinates of the j th site.

In order to separate the antiferromagnetic contribution from ξ , we introduce angular variable η ,

$$\eta_j = \xi_j - \pi(j_x + j_y + j_z) \quad (115)$$

where η_j is η at the j th site. We take the branch of η_j that satisfies the following condition,

$$-\pi \leq \eta_j - \eta_k < \pi \quad (116)$$

where k is a nearest neighbor site of j .

From $(\eta_j - \eta_k)$'s, we construct $(\xi_j - \xi_k)$'s. After $(\xi_j - \xi_k)$'s are obtained, we rebuild ξ from them. The process is as follows: first, we pick a value for the initial ξ_1 . After fixing the value of ξ_1 , we calculate ξ_2 by $\xi_2 = \xi_1 + (\xi_2 - \xi_1)$, where the site 2 is connected to the site 1 by a nearest neighbor bond. The step where value ξ_j is derived from the already evaluated value of ξ_k is given by

$$\xi_j = \xi_k + (\xi_j - \xi_k) \quad (117)$$

where the sites j and k are connected by a bond in the path for the rebuilding of ξ . This process is continued until values at all accessible sites are evaluated once and only once.

By this rebuilding process, a single path is constructed from the site 1 to other sites $k \neq 1$, which is achieved by making the region singly-connected by removing some bonds (see Fig. 4). We denote the path from the site 1 to other sites $k \neq 1$ by $C_{1 \rightarrow k}$. Then, the value ξ_k is given as

$$\xi_k \approx \xi_1 + \int_{C_{1 \rightarrow k}} \nabla \xi \cdot d\mathbf{r} \quad (118)$$

Let us summarize the calculation so far.

1. Solve Eq. (97) self-consistently using the Car-Parrinello method in Eq. (105). Thereby, we obtain Eqs. (101), (102), and (103).
2. From Eq. (102), ξ_k 's are obtained. Next, ξ_k 's are rebuilt using Eq. (118). Thereby, we have ξ_k 's for which we know the positions of the jump-of-values.
3. From the rebuilt ξ_k 's, construct $\tilde{c}_{j\sigma}^\dagger, \tilde{c}_{j\sigma}$ in Eq. (108). Using Eqs. (101), (102), and (103) obtained in Step 1), solve Eq. (110). Thereby, $\tilde{u}_{i\sigma}^n, \tilde{v}_{i\sigma}^n$ are obtained.
4. $e^{\pm \frac{i}{2}\xi_i}, \tilde{u}_{i\sigma}^n, \tilde{v}_{i\sigma}^n, u_{i\sigma}^n$ and $v_{i\sigma}^n$ are obtained; $u_{i\uparrow}^n = e^{-\frac{i}{2}\xi_i} \tilde{u}_{i\uparrow}^n, u_{i\downarrow}^n = e^{\frac{i}{2}\xi_i} \tilde{u}_{i\downarrow}^n, v_{i\uparrow}^n = e^{\frac{i}{2}\xi_i} \tilde{v}_{i\uparrow}^n, v_{i\downarrow}^n = e^{-\frac{i}{2}\xi_i} \tilde{v}_{i\downarrow}^n$.

Next, we obtain χ_k 's that satisfy Eq. (114). The requirement in Eq. (114) is given by

$$w_{C_\ell}[\chi] = \frac{1}{2\pi} \sum_{i=1}^{N_\ell} \tau_{C_\ell(i+1) \leftarrow C_\ell(i)} \quad (119)$$

where $w_{C_\ell}[\chi]$ is the number supplied as a boundary condition. $C_\ell(i)$ is the i th site in the loop C_ℓ , where $C_{N_\ell+1} = C_1$ with N_ℓ is the number of sites in C_ℓ .

In order to impose conditions in Eq. (122), $\tau_{j \leftarrow i}$ is split into a multi-valued part $\tau_{j \leftarrow i}^0$ and single-valued part $f_{j \leftarrow i}$ as

$$\tau_{j \leftarrow i} = \tau_{j \leftarrow i}^0 + f_{j \leftarrow i} \quad (120)$$

where $\tau_{j \leftarrow i}^0$ is given to satisfy

$$w_{C_\ell}[\chi] = \frac{1}{2\pi} \sum_{i=1}^{N_\ell} \tau_{C_\ell(i+1) \leftarrow C_\ell(i)}^0 \quad (121)$$

and $f_{j \leftarrow i}$ is solved to satisfy

$$0 = \frac{1}{2\pi} \sum_{i=1}^{N_\ell} f_{C_\ell(i+1) \leftarrow C_\ell(i)} \quad (122)$$

The above is the equation corresponding to Eq. (70). They are imposed on independent loops in the system, as shown in Fig. 6.

For Eq. (71), we use the conservation of the local charge at sites instead. As will be explained later, it gives rise to a sufficient number of equations to obtain unknowns.

The conservation of the local charge at the k th site is

$$0 = \sum_{\ell \in b_k} J_{k \leftarrow \ell} \quad (123)$$

where b_k is a set of sites that are connected to the k th site. $J_{k \leftarrow \ell}$ is the current through the bond between sites k and ℓ in the direction $k \leftarrow \ell$,

$$\begin{aligned} J_{k \leftarrow \ell} &= \frac{2e}{\hbar} \frac{\partial E}{\partial \tau_{k \leftarrow \ell}} = \frac{2e}{\hbar} \frac{\partial}{\partial \tau_{k \leftarrow \ell}} \langle \text{Gnd} | \sum_{i,j,\sigma,\sigma'} h_{i\sigma,j\sigma'} c_{i\sigma}^\dagger c_{j\sigma'} | \text{Gnd} \rangle \\ &= \frac{2e}{\hbar} \frac{\partial}{\partial \tau_{k \leftarrow \ell}} \sum_{i,j,\sigma,\sigma'} h_{i\sigma,j\sigma'} \sum_{n_1}^{\prime} \sum_{n_2}^{\prime} \langle e^{\frac{i}{2} \hat{\chi}_i} [(u_{i\sigma}^{n_1})^* \gamma_{n_1}^\dagger - \sigma(v_{i\sigma}^{n_1}) \gamma_{n_1}] [u_{j\sigma'}^{n_2} \gamma_{n_2} - \sigma'(v_{j\sigma'}^{n_2})^* \gamma_{n_2}^\dagger] e^{-\frac{i}{2} \hat{\chi}_j} \rangle \\ &= \frac{2e}{\hbar} \sum_{i,j,\sigma,\sigma'} h_{i\sigma,j\sigma'} \sum_n^{\prime} \frac{\partial e^{\frac{i}{2} \tau_{i \leftarrow j}}}{\partial \tau_{k \leftarrow \ell}} [(u_{i\sigma}^n)^* (u_{j\sigma'}^n) f(E_n) + \sigma \sigma' (v_{i\sigma}^n) (v_{j\sigma'}^n)^* f(-E_n)] \\ &= \frac{2e}{\hbar} \sum_{n,i,j,\sigma}^{\prime} \frac{\partial e^{\frac{i}{2} \tau_{i \leftarrow j}}}{\partial \tau_{k \leftarrow \ell}} \left\{ h_{i\sigma,j\sigma} [(u_{i\sigma}^n)^* u_{j\sigma}^n f(E_n) + (v_{j\sigma}^n)^* v_{i\sigma}^n f(-E_n)] \right. \\ &\quad \left. + h_{i\sigma,j-\sigma} [(u_{i\sigma}^n)^* u_{j-\sigma}^n f(E_n) - (v_{j-\sigma}^n)^* v_{i\sigma}^n f(-E_n)] \right\} \end{aligned} \quad (124)$$

We employ an iterative improvement of the approximate solutions by using the linearized version of Eq. (123) given by

$$0 \approx \frac{2e}{\hbar} \sum_i \frac{\partial E(\{\tau_{j \leftarrow i}^0\})}{\partial \tau_{j \leftarrow i}} + \frac{2e}{\hbar} \sum_i \frac{\partial^2 E(\{\tau_{j \leftarrow i}^0\})}{\partial (\tau_{j \leftarrow i})^2} f_{j \leftarrow i} \quad (125)$$

A system of equations for $f_{j \leftarrow i}$'s composed of Eqs. (122) and (125) are solve for given $\tau_{j \leftarrow i}^0$'s by updating them iteratively

$$\tau_{j \leftarrow i}^{0 \text{ New}} = \tau_{j \leftarrow i}^{0 \text{ Old}} + f_{j \leftarrow i} \quad (126)$$

where $\tau_{j \leftarrow i}^{0 \text{ Old}}$ is $\tau_{j \leftarrow i}^0$ value that is used to obtain the current value of $f_{j \leftarrow i}$; $\tau_{j \leftarrow i}^{0 \text{ New}}$ will be used to obtain the next $f_{j \leftarrow i}$ value.

The numerical convergence is checked by the condition

$$\left| \frac{2e}{\hbar} \sum_i \frac{\partial E(\{\tau_{j \leftarrow i}^0\})}{\partial \tau_{j \leftarrow i}} \right| < \epsilon \quad (127)$$

where ϵ is a small number.

For the initial $\tau_{j \leftarrow i}^0$, we adopt the following,

$$\tau_{j \leftarrow i}^{0 \text{ init}} = \sum_h w_h \tan^{-1} \frac{j_y - h_y}{j_x - h_x} - \sum_h w_h \tan^{-1} \frac{i_y - h_y}{i_x - h_x} \quad (128)$$

where (j_x, j_y, j_z) and (i_x, i_y, i_z) are coordinates of the sites j and i , respectively, $h = (h_x, h_y, h_z)$ is the coordinate of the hole occupied site, and w_h is the winding number of χ around the hole at h .

The number of $\tau_{j \leftarrow i}$ to be evaluated is equal to the number of the bonds. The number of equations in Eq. (70) is equal to the number of the plaques. The number of equations from Eq. (123) for the conservation of charge is equal to the number of sites-1, due to the fact that the total charge is fixed in the calculation.

The equality of the number of unknowns and the number of equations gives

$$[\# \text{ bonds}] = [\# \text{ plaques}] + [\# \text{ sites} - 1] \quad (129)$$

It is interesting to note that this agrees with the Euler's theorem for the two-dimensional lattice

$$[\# \text{ edges}] = [\# \text{ faces}] + [\# \text{ vertices} - 1] \quad (130)$$

When the spin-orbit interaction exists the total energy depends on ξ_1 as shown in Fig. 7. In Fig. 8, obtained current using $\xi_1 = 0$ is depicted.

I. Diamagnetic current produced by an application of a magnetic field

One of the hallmarks of superconducting states is the diamagnetic response to an external magnetic field. Let us consider the situation where a magnetic field $\mathbf{B}^{\text{em}} = \nabla \times \mathbf{A}^{\text{em}}$ is applied. Then, the modified energy functional in Eq. (77) is used.

This leads to replace $\tau_{j \leftarrow i}$ in E by

$$u_{j \leftarrow i} = \tau_{j \leftarrow i} - \frac{2e}{\hbar c} \int_i^j \mathbf{A}^{\text{em}} \cdot d\mathbf{r} \quad (131)$$

where integration is performed along the bond $j \leftarrow i$.




FIG. 7. Dependence of the total energy on the value of ξ_1 . The zero of energy is taken to be the value for $\lambda = 0$ calculation. The units of the energy is t_1 .




FIG. 8. Loop current obtained by the calculation. **a**: Loop current in the system. Each “m” indicates a center of a loop current with winding number +1, and “a” a center of a loop-current with winding number -1. **b**: Loop current in the bulk layer viewed from above. **c**: Loop current in the surface layer viewed from above.




FIG. 9. Supercurrent under the application of an external magnetic field. $B = 0.01$ in the units of $t_1 = 1, \hbar = 1, e = 1$, and lattice constant in the plane = 1. **a**: The result from the calculation. **b**: The linear approximation given in Eq. (135). **c**: The difference between the calculated result and its linear approximation.

The calculation can be done similarly to the case for no magnetic field, starting from the initial value

$$u_{j \leftarrow i}^0 \text{ init} = \tau_{j \leftarrow i}^0 \text{ init} - \frac{2e}{\hbar c} \int_{j \leftarrow i} \mathbf{A}^{\text{em}} \cdot d\mathbf{r} \quad (132)$$

Note that during the evaluation process of $\nabla\chi$, the ambiguity in the gauge of \mathbf{A}^{em} is compensated, thus, the effective vector potential

$$\mathbf{A}^{\text{eff}} = \mathbf{A}^{\text{em}} - \frac{\hbar c}{2e} \nabla\chi \quad (133)$$

is invariant with respect to the choice of the gauge in \mathbf{A}^{em} .

Now we apply a uniform magnetic field perpendicular to the lattice. In the actual numerical calculations we have adopted

$$\mathbf{A}^{\text{em}} = \begin{pmatrix} -By \\ 0 \\ 0 \end{pmatrix} \quad (134)$$




FIG. 10. Supercurrent induced by the application of an external magnetic field. **a**: The induced current in the two layers. The scale for the current arrow is enlarged from Fig. 9. **b**: The induced current in the bulk layer. **c**: the induced current in the surface layer.

but we checked that the calculated current distribution is identical even other gauge is employed.

In Fig. 9**a**, current distributions with applying the magnetic field are depicted.

Numerical calculations indicate the following relation holds

$$J_{j\leftarrow i} \approx -\frac{4e^2}{\hbar^2} \frac{\partial^2 E[\{0\}]}{\partial (u_{j\leftarrow i})^2} \int_i^j \mathbf{A}^{\text{eff}} \cdot d\mathbf{r} \quad (135)$$

where $\{0\}$ means all $u_{j\leftarrow i}$'s are zero [70]. The result using the above approximation is shown in Fig. 9**b**, and the difference between the exact result and approximate one is shown in Fig. 9**c**. The approximate one is almost identical to the exact one, except the current just around the small polarons; it originates directly from the Rashba interaction.

The induced current is diamagnetic as seen in Fig. 10. This will become a screening current of the magnetic field in a large system size.




FIG. 11. Supercurrent obtained by feeding external current. **a**: Supercurrent with external current. Two sets of three arrows indicate sites where external current enters and exits. Each arrow indicates J^{EX} . The units of the current is $t_1 = 1, \hbar = 1$, and $e = 1$. **b**: The difference of currents with and without external current. **c**: The difference of currents with and without external current in the bulk layer viewed from above. Arrows are magnified from those in **b**.

J. Zero voltage current production by an external current feeding

Another hallmark of superconducting states is the zero voltage current flow through the sample.

Let us consider the external current feeding. When external currents are fed, the conservation of the local charge at site k in Eq. (123) is modified as

$$0 = \sum_{\ell \in b_k} J_{k \leftarrow \ell} + J_k^{\text{EX}} \quad (136)$$

where J_k^{EX} is the external current fed at k .

The results are shown in Fig. 12. The energy minimum with nonzero J^{EX} occurs for $\lambda \neq 0$ cases. This situation is similar to the one envisaged by Bloch for superconducting states [71]. When the system is at the energy minimum with nonzero J^{EX} , zero voltage current production is realized. In other words, persistent current flows through the sample.




FIG. 12. Total energy vs J^{EX} . The units of J^{EX} is $t_1 = 1, \hbar = 1$, and $e = 1$. Three different Rashba parameters λ are employed. The zero of the energy is that of $\lambda = 0$ and $J^{\text{EX}} = 0$.

The energy minimum occurs at zero J^{EX} for the $\lambda = 0$ case. This indicates the if the Rashba interaction is absent, the zero voltage current production does not occur in the present model.

IV. CONCLUDING REMARKS

We have presented a fundamentally revised theory for superconductivity, and the method to calculate properties of superconducting states based on it.

Since the early work of London, it has been repeatedly argued that theory of superconductivity needs to have a $U(1)$ gauge field different from the electromagnetic one to have the gauge invariant supercurrent. London called it “superpotential”, and the current standard theory calls it “Nambu-Goldston mode”. In the new theory, the Berry connection \mathbf{A}^{fic} provides the necessary gauge field.

We would like to emphasize that the vector potential provided by the Berry connection \mathbf{A}^{fic} plays a crucial role in realizing reversible superconducting-normal metal phase transition in a magnetic field [13, 18, 21, 57]. This phenomenon cannot be explained by the standard theory [35, 57].

Besides the reversible superconducting-normal metal phase transition in a magnetic field,

the new theory solves all four problems described in Introduction. It also provides a calculation framework for the two of the hallmarks of superconducting states, the diamagnetic response current to an external magnetic field, and the zero voltage current produced by an external current feeding, microscopically. This theory may be able to elucidate the cuprate superconductivity, where nanoscale inhomogeneity of electronic states is important.

Lastly, we would like to note that a similar formalism using the Berry connection is possible for superfluidity of bosons [56].

Appendix A: Berry connection from many-body wave functions and the modification of Maxwell's equations

Berry's derivation for the Berry phase uses the adiabatic assumption [72]. The adiabatic assumption here does not necessary means a slow change. We will explain this point below.

By following Berry, let us consider the state vector where given by

$$|n(\mathbf{R}(t))\rangle \tag{A1}$$

where \mathbf{R} is the adiabatic parameter. This indicates the parametric time-dependence occurs through time-dependent adiabatic parameter $\mathbf{R}(t)$.

Then, the time-dependent wave function becomes

$$\begin{aligned} |\Psi(t)\rangle &= e^{i \int_{\mathbf{R}(0)}^{\mathbf{R}(t)} \langle n(\mathbf{R}(t)) | \nabla_{\mathbf{R}} | n(\mathbf{R}(t)) \rangle \cdot d\mathbf{R}} |n(\mathbf{R}(t))\rangle e^{-\frac{i}{\hbar} \int_0^t dt' E_n(t')} \\ &= |n(\mathbf{R}(t))\rangle e^{i\gamma_n(t)} e^{-\frac{i}{\hbar} \int_0^t dt' E_n(t')} \end{aligned} \tag{A2}$$

The phase

$$\gamma_n(t) = i \int_{\mathbf{R}(0)}^{\mathbf{R}(t)} d\mathbf{R} \cdot \mathbf{A}_n(\mathbf{R}) \tag{A3}$$

is the Berry phase, and

$$\mathbf{A}_n(\mathbf{R}) = -i \langle n(\mathbf{R}) | \nabla_{\mathbf{R}} | n(\mathbf{R}) \rangle \tag{A4}$$

is the Berry connection.

The Berry connection gives rise to a vector potential on the dynamics of \mathbf{R} . Let us see this below, taking the Born-Oppenheimer approximation as an example.

In the Born-Oppenheimer approximation, \mathbf{R} is the nuclear coordinates. The momentum conjugate to \mathbf{R} is $-i\hbar\nabla_{\mathbf{R}}$. The electron coordinates is denoted by \mathbf{r} . The Born-Oppenheimer total wave function we consider is

$$f(\mathbf{R})\langle\mathbf{r}|n(\mathbf{R})\rangle \quad (\text{A5})$$

where $f(\mathbf{R})$ is the nuclear wave function and $\langle\mathbf{r}|n(\mathbf{R})\rangle$ is the electronic wave function with the nuclear coordinate as the adiabatic parameter.

Application of the momentum operator conjugate to \mathbf{R} on the total wave function yields

$$-i\hbar\nabla_{\mathbf{R}} [f(\mathbf{R})\langle\mathbf{r}|n(\mathbf{R})\rangle] = \langle\mathbf{r}|n(\mathbf{R})\rangle[-i\hbar\nabla_{\mathbf{R}}f(\mathbf{R})] - f(\mathbf{R})[i\hbar\nabla_{\mathbf{R}}\langle\mathbf{r}|n(\mathbf{R})\rangle] \quad (\text{A6})$$

The integration over \mathbf{r} after applying $\langle n(\mathbf{R})|\mathbf{r}\rangle$ from the left yields

$$[-i\hbar\nabla_{\mathbf{R}} + \hbar\mathbf{A}_n(\mathbf{R})]f(\mathbf{R}) \quad (\text{A7})$$

This indicates that the dynamics of \mathbf{R} is that in the gauge field with the vector potential $\hbar\mathbf{A}_n$. In other words, the change of the momentum operator

$$-i\hbar\nabla_{\mathbf{R}} \rightarrow -i\hbar\nabla_{\mathbf{R}} + \hbar\mathbf{A}_n \quad (\text{A8})$$

occurs due to the presence of other particles.

Now, we shall consider a wave function for N electron system,

$$\Phi(\mathbf{r}_1, \dots, \mathbf{r}_N, t) \quad (\text{A9})$$

where \mathbf{r}_j is the coordinate of the j th particle.

The Berry connection \mathbf{A}_n takes into account the effect of the electron dynamics on the nuclear dynamics through the wave function they share (i.e., $f(\mathbf{R})\langle\mathbf{r}|n(\mathbf{R})\rangle$).

Let us calculate the Berry connection arising from $\Phi(\mathbf{r}_1, \dots, \mathbf{r}_N, t)$ by regarding \mathbf{r}_1 as an adiabatic parameter. In other words, we calculate the gauge field that affects the dynamics on the particle 1 generated by other particles 2, \dots , N through the wave function they share (i.e., $\Phi(\mathbf{r}_1, \dots, \mathbf{r}_N, t)$).

We cannot consider \mathbf{r}_1 as a slow variable compared with other $\mathbf{r}_2, \dots, \mathbf{r}_N$; however, the whole system stays in the same state described by the wave function $\Phi(\mathbf{r}_1, \dots, \mathbf{r}_N, t)$.

We first construct a normalized wave function $|n_{\Phi}(\mathbf{r}_1)\rangle$ that contains \mathbf{r}_1 as an adiabatic parameter,

$$\langle\mathbf{r}_2, \dots, \mathbf{r}_N|n_{\Phi}(\mathbf{r}_1, t)\rangle = \frac{\Phi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N, t)}{|C_{\Phi}(\mathbf{r}_1, t)|^{\frac{1}{2}}} \quad (\text{A10})$$

Here, $|C_\Phi(\mathbf{r}_1, t)|$ is the normalization constant given by

$$|C_\Phi(\mathbf{r}_1, t)| = \int d^3r_2 \cdots d^3r_N \Phi(\mathbf{r}_1, \mathbf{r}_2, \cdots) \Phi^*(\mathbf{r}_1, \mathbf{r}_2, \cdots) \quad (\text{A11})$$

The wave function $\langle \mathbf{r}_2, \cdots, \mathbf{r}_N | n_\Phi(\mathbf{r}_1) \rangle$ is normalized to 1 with respect to integrals over $\mathbf{r}_2, \cdots, \mathbf{r}_N$, and single-valued with respect to \mathbf{r}_1 . Then, the *Berry connection for many-body wave functions* is calculated as

$$\mathbf{A}_\Phi^{\text{MB}}(\mathbf{r}_1, t) = -i \langle n_\Phi(\mathbf{r}_1, t) | \nabla_{\mathbf{r}_1} | n_\Phi(\mathbf{r}_1, t) \rangle \quad (\text{A12})$$

As in the Born-Oppenheimer case, $\mathbf{A}_\Phi^{\text{MB}}(\mathbf{r}_1, t)$ will modify the momentum operator as

$$-i\hbar \nabla_{\mathbf{r}_1} \rightarrow -i\hbar \nabla_{\mathbf{r}_1} + \hbar \mathbf{A}_\Phi^{\text{MB}}(\mathbf{r}_1) \quad (\text{A13})$$

The same modification also occurs for $\mathbf{r}_i, i = 2 \cdots, N$.

Let us consider the effect $\mathbf{A}_\Phi^{\text{MB}}(\mathbf{r}_1, t)$ using the path integral formalism of quantum mechanics [21]. According to the path integral formalism, a wave function is a sum of contributions from all paths each contributes an exponential whose phase is the classical action divided by \hbar for the path in question [73]. For the system of charged particles and electromagnetic field, the classical action S are composed of the following three terms,

$$S = S_1 + S_2 + S_3 \quad (\text{A14})$$

where

$$S_1 = \sum_i \frac{m}{2} \int dt \dot{\mathbf{r}}_i^2 \quad (\text{A15})$$

is the action for the particles,

$$S_2 = - \int d^3r dt \left[\rho \phi^{\text{em}}(\mathbf{r}, t) - \frac{1}{c} \mathbf{j} \cdot \mathbf{A}^{\text{em}}(\mathbf{r}, t) \right] = -q \sum_i \int dt \left[\phi^{\text{em}}(\mathbf{r}_i, t) - \frac{1}{c} \dot{\mathbf{r}}_i \cdot \mathbf{A}^{\text{em}}(\mathbf{r}_i, t) \right] \quad (\text{A16})$$

is the action for the interaction between the field and particles, and

$$S_3 = \frac{1}{8\pi} \int d^3r dt \left[(\mathbf{E}^{\text{em}})^2 - (\mathbf{B}^{\text{em}})^2 \right] = \frac{1}{8\pi} \int d^3r dt \left[\left(-\nabla \phi^{\text{em}} - \frac{1}{c} \frac{\partial \mathbf{A}^{\text{em}}}{\partial t} \right)^2 - (\nabla \times \mathbf{A}^{\text{em}})^2 \right] \quad (\text{A17})$$

is the action for the field. Here, ϕ^{em} and \mathbf{A}^{em} are the scalar and vector potentials for the electromagnetic field, respectively; ρ and \mathbf{j} are the electric charge and current densities, respectively.

In the following we consider the case where the electric field \mathbf{E}^{em} is absent, and only the magnetic field \mathbf{B}^{em} is present.

The Berry connection $\mathbf{A}_{\Phi}^{\text{MB}}$ modifies the momentum operator by providing a $U(1)$ field. As a consequence, the effective vector potential for the $U(1)$ field in the system becomes

$$\mathbf{A}^{\text{eff}} = \mathbf{A}^{\text{em}} + \mathbf{A}^{\text{fic}}, \quad \mathbf{A}^{\text{fic}} = -\frac{\hbar c}{q} \mathbf{A}_{\Phi}^{\text{MB}} \quad (\text{A18})$$

Here, we have used the ‘‘fictitious’’ vector potential \mathbf{A}^{fic} in place of $\mathbf{A}_{\Phi}^{\text{MB}}$.

As will be shown later, \mathbf{A}^{fic} is given by

$$\mathbf{A}^{\text{fic}} = -\frac{\hbar c}{2e} \nabla \chi \quad (\text{A19})$$

where χ is an angular variable with period 2π [19].

By including the Berry connection, S_2 becomes,

$$S'_2 = \frac{1}{c} \int d^3r dt \mathbf{j} \cdot \mathbf{A}^{\text{eff}}(\mathbf{r}, t) \quad (\text{A20})$$

This gives rise to ‘‘Lorentz force’’. The Lorentz force from \mathbf{A}^{fic} is zero in classical mechanics; however, \mathbf{A}^{fic} may affect the dynamics of charged particles through the Aharonov-Bohm effect [74] in quantum mechanics. Actually, this effect is the main concern of the present work. We call this term, the *Lorentz interaction* term.

The electromagnetic field energy is the energy stored in the space through the Lorentz interaction term. If we use S'_2 as the Lorentz interaction term, S_3 should be modified as

$$S'_3 = -\frac{1}{8\pi} \int d^3r dt (\mathbf{B}^{\text{eff}})^2 \quad (\text{A21})$$

where $\mathbf{B}^{\text{eff}} = \nabla \times \mathbf{A}^{\text{eff}}$.

$\mathbf{B}^{\text{fic}} = \nabla \times \mathbf{A}^{\text{fic}}$ may not be zero due to the fact that χ may be multi-valued.

The use of S'_2 and S'_3 , modify two of the Maxwell’s equations,

$$\nabla \cdot \mathbf{B}^{\text{eff}} = 0 \quad (\text{A22})$$

$$\nabla \times \mathbf{B}^{\text{eff}} = \frac{4\pi}{c} \mathbf{j} \quad (\text{A23})$$

The first one gives rise to a Dirac monopole as shown below: we rewrite it as

$$\nabla \cdot \mathbf{B}^{\text{em}} = -\nabla \cdot (\nabla \times \mathbf{A}^{\text{fic}}) \quad (\text{A24})$$

When the both sides of the above equation are integrated for a closed region with surface S_f , we have

$$\int_{S_f} d\mathbf{S} \cdot \mathbf{B}^{\text{em}} = - \int_{S_f} d\mathbf{S} \cdot (\nabla \times \mathbf{A}^{\text{fic}}) \quad (\text{A25})$$

We split S_f into two surfaces S_{f_1} and S_{f_2} with common boundary loop, $C = \partial(S_{f_1}) = -\partial(S_{f_2})$. Then, we have the following relation

$$\int_{S_{f_1}} d\mathbf{S} \cdot (\nabla \times \mathbf{A}^{\text{fic}}) + \int_{S_{f_2}} d\mathbf{S} \cdot (\nabla \times \mathbf{A}^{\text{fic}}) = \int_{\partial(S_{f_1})} d\mathbf{r} \cdot \mathbf{A}^{\text{fic}} + \int_{\partial(S_{f_1})} d\mathbf{r} \cdot \mathbf{A}^{\text{fic}} \quad (\text{A26})$$

We consider the case in which singularities exist in \mathbf{A}^{fic} : let us take a closed surface S with boundary $C = \partial S$, and \mathbf{A}^{fic} has a singularity in S . Then, the following relation is obtained

$$\int_C d\mathbf{r} \cdot \frac{\hbar c}{2e} \nabla \chi = \frac{\hbar c}{2e} n \quad (\text{A27})$$

where n is an integer.

If we have the case where $n = 0$ for $\partial(S_{f_1})$ term and $n = 1$ for $\partial(S_{f_2})$ term in Eq. (A26), respectively, Eq. (A25) yields

$$\int_{S_f} d\mathbf{S} \cdot \mathbf{B}^{\text{em}} = \frac{\hbar c}{2e} \quad (\text{A28})$$

This shows that a monopole with magnetic charge $\frac{\hbar c}{2e}$ exists in the region enclosed by S_f [75].

Let us show $\nabla \times \mathbf{B}^{\text{fic}} = 0$: it is well-known that $\nabla \chi$ in \mathbf{A}^{fic} can be decomposed as

$$\nabla \chi = \nabla \chi_0 + \nabla f, \quad \nabla^2 \chi_0 = 0 \quad (\text{A29})$$

where f is a single-valued ($\nabla \times \nabla f = 0$), and χ_0 may be multi-valued. Thus, we have

$$\nabla \times \mathbf{B}^{\text{fic}} = \nabla \times (\nabla \times \nabla \chi_0) = \nabla(\nabla^2 \chi_0) - \nabla^2 \nabla \chi_0 = 0 \quad (\text{A30})$$

As a consequence Eq. (A23) is equal to

$$\nabla \times \mathbf{B}^{\text{em}} = \frac{4\pi}{c} \mathbf{j} \quad (\text{A31})$$

Appendix B: Appearance of \mathbf{A}^{fic} from spin-twisting itinerant motion of electrons.

The electrons in the normal state of the BCS superconductor are assumed to be Bloch electrons. We consider the Bloch electron under the influence of the Rashba interaction, and show that the spin-twisting itinerant motion is realized [19].

We use the following Rashba interaction term

$$H_{so} = \boldsymbol{\lambda}(\mathbf{r}) \cdot \frac{\hbar\boldsymbol{\sigma}}{2} \times (\hat{\mathbf{p}} - q\mathbf{A}^{\text{em}}(\mathbf{r})), \quad (\text{B1})$$

where $\boldsymbol{\lambda}(\mathbf{r})$ is the spin-orbit coupling vector (its direction is the internal electric field direction), $\hat{\mathbf{p}} = -i\hbar\nabla$ is the momentum operator, and $q = -e$ is electron charge [76].

We take into account the coordinate dependence of the spin function by adopting the following spin function

$$\Sigma_1(\mathbf{r}) = e^{-\frac{i}{2}\chi(\mathbf{r})} \begin{pmatrix} e^{-i\frac{1}{2}\xi(\mathbf{r})} \sin \frac{\zeta(\mathbf{r})}{2} \\ e^{i\frac{1}{2}\xi(\mathbf{r})} \cos \frac{\zeta(\mathbf{r})}{2} \end{pmatrix} \quad (\text{B2})$$

for the Bloch electron, where $\zeta(\mathbf{r})$ and $\xi(\mathbf{r})$ are the polar and azimuthal angles of the spin direction, respectively. The expectation value of spin $\mathbf{s}(\mathbf{r}) = (s_x(\mathbf{r}), s_y(\mathbf{r}), s_z(\mathbf{r}))$ is given by

$$s_x(\mathbf{r}) = \frac{\hbar}{2} \cos \xi(\mathbf{r}) \sin \zeta(\mathbf{r}), \quad s_y(\mathbf{r}) = \frac{\hbar}{2} \sin \xi(\mathbf{r}) \sin \zeta(\mathbf{r}), \quad s_z = \frac{\hbar}{2} \cos \zeta(\mathbf{r}) \quad (\text{B3})$$

The Berry connection arising from the spin function $\Sigma_1(\mathbf{r})$ is

$$\mathbf{A}_1^{\text{fic}}(\mathbf{r}) = -i\frac{\hbar c}{e}\Sigma_1^\dagger \nabla \Sigma_1 = -\frac{\hbar c}{2e}\nabla\chi(\mathbf{r}) + \frac{\hbar c}{2e}\nabla\xi(\mathbf{r}) \cos \zeta(\mathbf{r}) \quad (\text{B4})$$

and the effective vector potential is

$$\mathbf{A}_1^{\text{eff}} = \mathbf{A}^{\text{em}} + \mathbf{A}_1^{\text{fic}} \quad (\text{B5})$$

We introduce the following wave vector \mathbf{k}_c for convenience,

$$\mathbf{k}_c = \mathbf{q}_c + \frac{e}{\hbar c}\mathbf{A}_1^{\text{eff}}(\mathbf{r}_c) \quad (\text{B6})$$

By following the standard method for the wave packet formalism[19, 77], the following equations of motion are obtained,

$$\dot{\mathbf{r}}_c = \frac{1}{\hbar} \frac{\partial \mathcal{E}}{\partial \mathbf{k}_c} + \boldsymbol{\lambda}(\mathbf{r}_c) \times \mathbf{s}(\mathbf{r}_c) \quad (\text{B7})$$

$$\dot{\mathbf{k}}_c = -\frac{e}{\hbar c}\dot{\mathbf{r}}_c \times \mathbf{B}^{\text{eff}}, \quad (\text{B8})$$

where \mathbf{B}^{eff} is the effective magnetic field,

$$\mathbf{B}^{\text{eff}} = \nabla \times \mathbf{A}_1^{\text{eff}} \quad (\text{B9})$$

Eqs. (B7) and (B8) indicate that the cyclotron motion in the band with energy

$$\mathcal{E}(\mathbf{k}) + \hbar \boldsymbol{\lambda}(\mathbf{r}) \times \mathbf{s}(\mathbf{r}) \cdot \mathbf{k} \quad (\text{B10})$$

occurs.

By following the Onsager's argument[78], let us quantize the cyclotron orbit. From Eq. (B6), the Bohr-Sommerfeld relation becomes

$$\oint_C \hbar \mathbf{q}_c \cdot d\mathbf{r}_c = \oint_C \left(\hbar \mathbf{k}_c - \frac{e}{c} \mathbf{A}_1^{\text{eff}} \right) \cdot d\mathbf{r}_c = 2\pi\hbar \left(n + \frac{1}{2} \right) \quad (\text{B11})$$

where n is an integer and C is the closed loop that corresponds to the section of Fermi surface enclosed by the cyclotron orbit.

Using Eq. (B8), we have

$$\begin{aligned} \oint_C \hbar \mathbf{k}_c \cdot d\mathbf{r}_c &= -\frac{e}{c} \oint_C d\mathbf{r}_c \cdot \mathbf{r}_c \times \mathbf{B}^{\text{eff}} \\ &= \frac{e}{c} \oint_C \mathbf{B}^{\text{eff}} \cdot \mathbf{r}_c \times d\mathbf{r}_c \\ &= \frac{2e}{c} \oint_C \mathbf{A}_1^{\text{eff}} \cdot d\mathbf{r}_c \end{aligned} \quad (\text{B12})$$

Thus, Eq. (B11) becomes

$$\frac{e}{c} \oint_C \mathbf{A}^{\text{em}} \cdot d\mathbf{r}_c + \frac{e}{c} \oint_C \mathbf{A}_1^{\text{fic}} \cdot d\mathbf{r}_c = 2\pi\hbar \left(n + \frac{1}{2} \right) \quad (\text{B13})$$

This is the well-known quantized condition for the cyclotron motion if $\mathbf{A}_1^{\text{fic}}$ is absent.

If $\mathbf{A}_1^{\text{fic}}$ is present, the above quantization condition is satisfied even if the magnetic field is absent. In this case, Eq. (B13) yields,

$$-\oint_C \frac{1}{2} \nabla \chi(\mathbf{r}) \cdot d\mathbf{r}_c + \oint_C \frac{1}{2} \nabla \xi(\mathbf{r}) \cos \zeta(\mathbf{r}) \cdot d\mathbf{r}_c = 2\pi \left(n + \frac{1}{2} \right) \quad (\text{B14})$$

which can be satisfied by the following two sets of conditions; one is $\zeta = \frac{\pi}{2}$, $w_C[\chi] = -1$, and $n = 0$; and the other is $\zeta = \frac{\pi}{2}$, $w_C[\chi] = 1$, and $n = -1$, where

$$w_C[\chi] = \frac{1}{2\pi} \oint_C \nabla \chi(\mathbf{r}) \cdot d\mathbf{r} \quad (\text{B15})$$

is the winding number of χ along loop C .

When $\zeta = \frac{\pi}{2}$, the single-valued condition for $\Sigma_1(\mathbf{r})$ in Eq. (B2) leads to the single-valuedness of $e^{-\frac{i}{2}\chi(\mathbf{r})}e^{\pm i\frac{1}{2}}$. This is satisfied if the following requirement

$$w_C[\chi] + w_C[\xi] = \text{even number} \quad (\text{B16})$$

is satisfied.

The condition $w_C[\chi] = \pm 1$ requires that $w_C[\xi]$ must be odd, thus, $w_C[\xi]$ is not zero. The nonzero value of $w_C[\xi]$ means that electrons perform spin-twisting itinerant motion. This indicates that the quantized cyclotron motion occurs without an external magnetic field by performing the spin-twisting itinerant motion.

When the quantized cyclotron motion without magnetic field occurs, $\mathbf{A}_1^{\text{fic}}$ is given by

$$\mathbf{A}_1^{\text{fic}}(\mathbf{r}) = -\frac{\hbar c}{2e}\nabla\chi(\mathbf{r}) \quad (\text{B17})$$

In addition to Σ_1 in Eq. (B2), we consider the spin function Σ_2 that is orthogonal to Σ_1 ,

$$\Sigma_2(\mathbf{r}) = e^{-\frac{i}{2}\chi(\mathbf{r})} \begin{pmatrix} ie^{-i\frac{1}{2}\xi(\mathbf{r})} \cos \frac{\zeta(\mathbf{r})}{2} \\ -ie^{i\frac{1}{2}\xi(\mathbf{r})} \sin \frac{\zeta(\mathbf{r})}{2} \end{pmatrix} \quad (\text{B18})$$

The fictitious vector potential from Σ_2 is

$$\mathbf{A}_2^{\text{fic}}(\mathbf{r}) = -i\frac{\hbar c}{e}\Sigma_2^\dagger\nabla\Sigma_2 = -\frac{\hbar c}{2e}\nabla\chi(\mathbf{r}) - \frac{\hbar c}{2e}\nabla\xi(\mathbf{r})\cos\zeta(\mathbf{r}) \quad (\text{B19})$$

and the effective vector potential is

$$\mathbf{A}_2^{\text{eff}} = \mathbf{A}^{\text{em}} + \mathbf{A}_2^{\text{fic}} \quad (\text{B20})$$

When the quantized cyclotron motion without magnetic field occurs, $\mathbf{A}_2^{\text{fic}}$ is given by

$$\mathbf{A}_2^{\text{fic}}(\mathbf{r}) = -\frac{\hbar c}{2e}\nabla\chi(\mathbf{r}) \quad (\text{B21})$$

just like $\mathbf{A}_1^{\text{fic}}$

Let us consider the case where the many-electron wave function Φ for N electrons is given as a Slater determinant of spin-orbitals $\phi_{1,1}(\mathbf{r})\Sigma_1(\mathbf{r}), \phi_{1,2}(\mathbf{r})\Sigma_2(\mathbf{r}), \dots, \phi_{\frac{N}{2},1}(\mathbf{r})\Sigma_1(\mathbf{r}),$ and $\phi_{\frac{N}{2},2}(\mathbf{r})\Sigma_2(\mathbf{r})$, where $\phi_{j,1}(\mathbf{r})$ and $\phi_{j,2}(\mathbf{r})$ are time-reversal partners and N is assumed to be even.

Then, $\mathbf{A}_\Phi^{\text{MB}}$ in Eq. (A12) is calculated as

$$\begin{aligned} \mathbf{A}_\Phi^{\text{MB}} &= \text{Im} \frac{\sum_{j=1}^{\frac{N}{2}} \left[\phi_{j,1}^*(\mathbf{r})\Sigma_1^\dagger(\mathbf{r})\nabla\phi_{j,1}(\mathbf{r})\Sigma_1(\mathbf{r}) + \phi_{j,2}^*(\mathbf{r})\Sigma_2^\dagger(\mathbf{r})\nabla\phi_{j,2}(\mathbf{r})\Sigma_2(\mathbf{r}) \right]}{\sum_{j=1}^{\frac{N}{2}} \left[\phi_{j,1}^*(\mathbf{r})\phi_{j,1}(\mathbf{r}) + \phi_{j,2}^*(\mathbf{r})\phi_{j,2}(\mathbf{r}) \right]} \\ &= \frac{e}{\hbar} \frac{\mathbf{A}_1^{\text{fic}} \sum_{j=1}^{\frac{N}{2}} \phi_{j,1}^*(\mathbf{r})\phi_{j,1}(\mathbf{r}) + \mathbf{A}_2^{\text{fic}} \sum_{j=1}^{\frac{N}{2}} \phi_{j,2}^*(\mathbf{r})\phi_{j,2}(\mathbf{r})}{\sum_{j=1}^{\frac{N}{2}} \left[\phi_{j,1}^*(\mathbf{r})\phi_{j,1}(\mathbf{r}) + \phi_{j,2}^*(\mathbf{r})\phi_{j,2}(\mathbf{r}) \right]} \end{aligned} \quad (\text{B22})$$

where “Im” indicates the imaginary part. Note that

$$\sum_{j=1}^{\frac{N}{2}} \left[\phi_{j,1}^*(\mathbf{r}) \nabla \phi_{j,1}(\mathbf{r}) + \phi_{j,2}^*(\mathbf{r}) \nabla \phi_{j,2}(\mathbf{r}) \right] \text{ is real} \quad (\text{B23})$$

since $\phi_{j,1}(\mathbf{r})$ and $\phi_{j,2}(\mathbf{r})$ are time-reversal partners.

When the quantized motion of spin-twisting itinerant motion occurs, $\mathbf{A}_1^{\text{fic}}$ and $\mathbf{A}_2^{\text{fic}}$ are given by $\mathbf{A}_1^{\text{fic}} = \mathbf{A}_2^{\text{fic}} = -\frac{\hbar c}{2e} \nabla \chi$. Thus, we have

$$\mathbf{A}_{\Phi}^{\text{MB}} = -\frac{1}{2} \nabla \chi \quad (\text{B24})$$

It is shown that when the pairing energy gap is formed, the smallest possible size of the loop current from the quantized motion is estimated to be ξ_{BCS} in Eq. (24) [21]. The loop currents can coexist with displacing their centers. This suggests that ξ_{BCS} -sized loop currents exist in the BCS superconductor with the current directions arranged in such a way that the macroscopic current is zero.

Appendix C: Reversible superconducting-normal metal phase transition in a magnetic field.

We will show that how the existence of \mathbf{A}^{fic} explains the reversible superconducting-normal metal phase transition in a magnetic field.

Let us consider the magnetic field part of the free energy of the Ginzburg-Landau theory is given by

$$F_{\text{mag}} = \int d^3r \frac{1}{8\pi} (\mathbf{B}^{\text{eff}})^2 \quad (\text{C1})$$

During the phase transition, the superconducting region shrinks and the magnetic field penetrating region increases. As a consequence $\partial_t \mathbf{B}^{\text{em}} \neq 0$ occurs. This will induce an electric field, and if normal current exists, the Joule heating should occur. Thus, for the reversible transition to occur, it should occur without generating normal current.

The change of F_{mag} due to $\partial_t \mathbf{B}^{\text{eff}} \neq 0$ is given by

$$\Delta F_{\text{mag}} = \int d^3r \int_t^{t+\Delta t} dt \frac{1}{4\pi} \mathbf{B}^{\text{eff}} \cdot \partial_t \mathbf{B}^{\text{eff}} \quad (\text{C2})$$

where Δt is the time interval for this change

If the above change of \mathbf{B}^{eff} is only due to the change of \mathbf{A}^{fic} , ΔF_{mag} is given by

$$\Delta F_{\text{mag}} = \frac{1}{c} \int d^3r \mathbf{j}_s \cdot \int_t^{t+\Delta t} dt \partial_t \mathbf{A}^{\text{fic}} \quad (\text{C3})$$

where

$$\mathbf{B}^{\text{eff}} = \nabla \times (\mathbf{A}^{\text{em}} + \mathbf{A}^{\text{fic}}), \quad \mathbf{j}_s = \frac{c}{4\pi} \nabla \times \mathbf{B}^{\text{eff}} \quad (\text{C4})$$

are used.

The supercurrent kinetic energy part of the free energy is given by

$$F_{\text{kin}} = \int d^3r \frac{m_e}{2} n_s \mathbf{v}^2 = \int d^3r \frac{e^2 n_s}{2m_e c^2} (\mathbf{A}^{\text{eff}})^2 \quad (\text{C5})$$

where the supercurrent density

$$\mathbf{j}_s = -en_s \mathbf{v} = -\frac{e^2 n_s}{m_e c} \mathbf{A}^{\text{eff}} \quad (\text{C6})$$

is used.

The change of the supercurrent kinetic energy ΔF_{kin} caused by $\partial_t \mathbf{A}^{\text{fic}}$ is

$$\Delta F_{\text{kin}} = -\frac{1}{c} \int d^3r \mathbf{j}_s \cdot \int_t^{t+\Delta t} dt \partial_t \mathbf{A}^{\text{fic}} \quad (\text{C7})$$

Therefore, according to Eqs. (C3) and (C7), the transition with $\Delta F_{\text{mag}} + \Delta F_{\text{kin}} = 0$ is realized via

$$\int_t^{t+\Delta t} dt \partial_t \mathbf{A}^{\text{fic}} \quad (\text{C8})$$

This occurs in a quantized manner by changing the winding numbers for χ , without Joule heating. This is the key step to realize the reversible superconducting-normal metal phase transition in a magnetic field. Other changes occur among \mathbf{j}_s , \mathbf{B}^{em} , n_s , to satisfy $\nabla \mathbf{B}^{\text{em}} = \frac{4\pi}{c} \mathbf{j}_s$ and $\partial_t n_s + \nabla \cdot \mathbf{j}_s = 0$. However, those changes can proceed without Joule heating [57].

Appendix D: Collective mode χ and associated number changing operators

We consider the quantization of the collective mode arising from χ . For this purpose, we use the Heisenberg operator formalism.

In order to have the phase arising from \mathbf{A}^{fic} in Eq. (A20), we express Φ as

$$\Phi = \Phi_0 e^{-\frac{i}{2} \sum_{j=1}^N \chi(\mathbf{r}_j)} \quad (\text{D1})$$

by explicitly extracting the Berry phase part $e^{-\frac{i}{2}\sum_{j=1}^N\chi(\mathbf{r}_j)}$.

This is actually the definition for Φ_0 through Φ and the Berry connection,

$$\Phi_0 = \Phi e^{\frac{i}{2}\sum_{j=1}^N\chi(\mathbf{r}_j)} \quad (\text{D2})$$

The kinetic energy part of the Hamiltonian is given by

$$K_0 = \frac{1}{2m_e} \sum_{j=1}^N \left(\frac{\hbar}{i} \nabla_j \right)^2 \quad (\text{D3})$$

where m_e is the electron mass. The current density calculated using the above kinetic energy

$$\mathbf{j}(\mathbf{r}) = -q \sum_{j=1}^N \frac{\hbar}{2im_e} \int d^3r_1 \cdots d^3r_N [\Phi_0^* \nabla_j \Phi_0 - \Phi_0 \nabla_j \Phi_0^*] \delta(\mathbf{r} - \mathbf{r}_j) \quad (\text{D4})$$

is zero; i.e., Φ_0 is a currentless state.

When the ground state does not have the non-trivial Berry connection, it is a currentless state. In this case, we may identify Φ as Φ_0 , and the collective coordinate χ disappears.

However, when the Berry connection is non-trivial, the collective coordinate χ is present. We treat χ as a collective dynamical variable, and quantize it with the help of the following Lagrangian,

$$\mathcal{L} = \langle \Phi | i\hbar \partial_t - H | \Phi \rangle = i\hbar \langle \Phi_0 | \partial_t | \Phi_0 \rangle + \frac{\hbar}{2} \int d\mathbf{r} n_p \dot{\chi} - \langle \Phi | H | \Phi \rangle \quad (\text{D5})$$

where n_p is the number density of the particles [79].

From the above Lagrangian, the conjugate momentum of χ is obtained as

$$p_\chi = \frac{\delta \mathcal{L}}{\delta \dot{\chi}} = \frac{\hbar}{2} n_p \quad (\text{D6})$$

thus, χ and n_p are canonical conjugate variables.

By following the canonical quantization condition

$$[\hat{p}_\chi(\mathbf{r}, t), \hat{\chi}(\mathbf{r}', t)] = -i\hbar \delta(\mathbf{r} - \mathbf{r}') \quad (\text{D7})$$

where \hat{p}_χ and $\hat{\chi}$ are operators corresponding to p_χ and χ respectively, we obtain the following relation

$$[\hat{n}(\mathbf{r}, t), \hat{\chi}(\mathbf{r}', t)] = -2i\delta(\mathbf{r} - \mathbf{r}') \quad (\text{D8})$$

where \hat{n} is the operator corresponding to n_p .

Strictly speaking, $\hat{\chi}$ is not a hermitian operator; however, it is known that when it is used as $e^{\pm\frac{i}{2}\hat{\chi}}$, the problem is avoided, practically [80].

We construct the following boson field operators from $\hat{\chi}$ and \hat{n} ,

$$\hat{\psi}^\dagger(\mathbf{r}) = (\hat{n}(\mathbf{r}))^{1/2} e^{\frac{i}{2}\hat{\chi}(\mathbf{r})}, \quad \hat{\psi}(\mathbf{r}) = e^{-\frac{i}{2}\hat{\chi}(\mathbf{r})} (\hat{n}(\mathbf{r}))^{1/2} \quad (\text{D9})$$

Using Eq. (D8), the following relations are obtained,

$$[\hat{\psi}(\mathbf{r}), \hat{\psi}^\dagger(\mathbf{r}')] = \delta(\mathbf{r} - \mathbf{r}'), \quad [\hat{\psi}(\mathbf{r}), \hat{\psi}(\mathbf{r}')] = 0, \quad [\hat{\psi}^\dagger(\mathbf{r}), \hat{\psi}^\dagger(\mathbf{r}')] = 0 \quad (\text{D10})$$

From Eq. (D10), the following relations are obtained

$$[e^{\pm\frac{i}{2}\hat{\chi}(\mathbf{r})}, \hat{\psi}^\dagger(\mathbf{r}')\hat{\psi}(\mathbf{r}')] = \mp e^{\pm\frac{i}{2}\hat{\chi}(\mathbf{r})}\delta(\mathbf{r} - \mathbf{r}') \quad (\text{D11})$$

which indicate that $e^{\pm\frac{i}{2}\hat{\chi}(\mathbf{r})}$ are number changing operators for the number density operator

$$\hat{n}(\mathbf{r}) = \hat{\psi}^\dagger(\mathbf{r})\hat{\psi}(\mathbf{r}) \quad (\text{D12})$$

-
- [1] F. London, Quantenmechanische deutung der theorie von weyl, Zeitschrift für Physik A Hadrons and nuclei **42**, 375 (1927).
 - [2] F. London, *Superfluids*, Vol. 1 (Wiley, New York, 1950).
 - [3] B. S. Deaver and W. M. Fairbank, Experimental evidence for quantized flux in superconducting cylinders, Phys. Rev. Lett. **7**, 43 (1961).
 - [4] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Theory of superconductivity, Phys. Rev. **108**, 1175 (1957).
 - [5] N. N. Bogoliubov, A new method in the theory of superconductivity I, Soviet Physics JETP **34**, 41 (1958).
 - [6] P. W. Anderson, Coherent excited states in the theory of superconductivity: Gauge invariance and the meissner effect, Phys. Rev. **110**, 827 (1958).
 - [7] P. W. Anderson, Random-phase approximation in the theory of superconductivity, Phys. Rev. **112**, 1900 (1958).
 - [8] Y. Nambu, Quasi-particles and gauge invariance in the theory of superconductivity, Phys. Rev. **117**, 648 (1960).

- [9] V. L. Ginzburg and L. D. Landau, On the theory of superconductivity, *Zh. Exsp. Teor. Fiz.* **20**, 1064 (1950).
- [10] S. Weinberg, Superconductivity for particular theorists, *Progress of Theoretical Physics Supplement* **86**, 43 (1986).
- [11] L. P. Gor'kov, Microscopic derivation of the Ginzburg-Landau equations in the theory of superconductivity, *Sov. Phys. JETP* **36**, 1364 (1959).
- [12] J. E. Hirsch, The london moment: what a rotating superconductor reveals about superconductivity, *Physica Scripta* **89**, 015806 (2013).
- [13] H. Koizumi, London moment, london's superpotential, nambu-goldstone mode, and berry connection from many-body wave functions, DOI: 10.1007/s10948-021-05827-9 *J. Supercond. Nov. Magn.* (2020), arXiv:2011.10701 [cond-mat.supr-con].
- [14] A. A. Abrikosov, On the magnetic properties of superconductors of the second group, *Sov. Phys. JETP* **5**, 1174 (1957).
- [15] B. D. Josephson, Possible new effects in superconductive tunneling, *Phys. Lett.* **1**, 251 (1962).
- [16] S. Shapiro, Josephson currents in superconducting tunneling: the effect of microwaves and other observations, *Phys. Rev. Lett.* **11**, 80 (1963).
- [17] H. Koizumi, Spin-vortex superconductivity, *J. Supercond. Nov. Magn.* **24**, 1997 (2011).
- [18] H. Koizumi and A. Ishikawa, Theory of supercurrent in superconductors, *International Journal of Modern Physics B* **34**, 2030001 (2020), <https://doi.org/10.1142/S0217979220300017>.
- [19] H. Koizumi, Possible occurrence of superconductivity by the π -flux Dirac string formation due to spin-twisting itinerant motion of electrons, *Symmetry* **12**, 776 (2020).
- [20] H. Koizumi and M. Tachiki, Supercurrent generation by spin-twisting itinerant motion of electrons: Re-derivation of the ac josephson effect including the current flow through the leads connected to josephson junction, *J. Supercond. Nov. Magn.* **28**, 61 (2015).
- [21] H. Koizumi, Superconductivity by Berry connection from many-body wave functions: revisit to Andreev–Saint-James reflection and Josephson effect, arXiv 2011.10701, to appear in *J. Supercond. Nov. Magn.* (2021), 2011.10701.
- [22] J. G. Bednorz and K. A. Müller, Possible high t_c superconductivity in the ba-la-cu-o system, *Z. Phys. B* **64**, 189 (1986).
- [23] M. A. Kastner, R. J. Birgeneau, G. Shirane, and Y. Endoh, Magnetic, transport, and optical

- properties of monolayer copper oxides, *Rev. Mod. Phys.* **70**, 897 (1998).
- [24] V. J. Emery and S. A. Kivelson, Importance of phase fluctuation in superconductors with small superfluid density, *Nature* **374**, 434 (1995).
- [25] A. Bianconi, N. L. Saini, A. Lanzara, M. Missori, T. Rossetti, H. Oyanagi, H. Yamaguchi, K. Oka, and T. Ito, Determination of the local lattice distortions in the CuO_2 plane of $\text{La}_{1.85}\text{Sr}_{0.15}\text{CuO}_4$, *Phys. Rev. Lett.* **76**, 3412 (1996).
- [26] R. Hidekata and H. Koizumi, Spin-vortices and spin-vortex-induced loop currents in the pseudogap phase of cuprates, *J. Supercond. Nov. Magn.* **24**, 2253 (2011).
- [27] J. Hirsch, Bcs theory of superconductivity: it is time to question its validity, *Physica Scripta* **80**, 035702 (2009).
- [28] W. H. Keesom and J. A. Kok, *Physica* **1**, 503 (1934).
- [29] W. H. Keesom and J. A. Kok, *Physica* **1**, 595 (1934).
- [30] W. H. Keesom and P. H. van Laer, *Physica* **4**, 487 (1937).
- [31] W. H. Keesom and P. H. van Laer, *Physica* **5**, 193 (1938).
- [32] P. H. van Laer and W. H. Keesom, On the reversibility of the transition processes between the superconductive and the normal state, *Physica* **5**, 993 (1938).
- [33] J. E. Hirsch, Momentum of superconducting electrons and the explanation of the Meissner effect, *Phys Rev. B* **95**, 014503 (2017).
- [34] J. E. Hirsch, Entropy generation and momentum transfer in the superconductor-normal and normal-superconductor phase transitions and the consistency of the conventional theory of superconductivity, *International Journal of Modern Physics B* **32**, 1850158 (2018).
- [35] J. E. Hirsch, Inconsistency of the conventional theory of superconductivity, *EPL* **130**, 17006 (2020).
- [36] R. Kubo, Statistical-mechanical theory of irreversible processes. I. General theory and simple applications to magnetic and conduction problems, *Journal of the Physical Society of Japan* **12**, 570 (1957).
- [37] A. Murani, N. Bourlet, H. le Sueur, F. Portier, C. Altimiras, D. Esteve, H. Grabert, J. Stockburger, J. Ankerhold, and P. Joyez, Absence of a dissipative quantum phase transition in josephson junctions, *Phys. Rev. X* **10**, 021003 (2020).
- [38] P. J. Hakonen and E. B. Sonin, Comment on “absence of a dissipative quantum phase transition in josephson junctions”, *Phys. Rev. X* **11**, 018001 (2021).

- [39] A. Murani, N. Bourlet, H. le Sueur, F. Portier, C. Altimiras, D. Esteve, H. Grabert, J. Stockburger, J. Ankerhold, and P. Joyez, Reply to “comment on ‘absence of a dissipative quantum phase transition in josephson junctions’”, *Phys. Rev. X* **11**, 018002 (2021).
- [40] R. Becker, G. Heller, and F. Sauter, Über die Stromverteilung in einer supraleitenden Kugel, *Zeitschrift für Physik* **85**, 772 (1933).
- [41] A. F. Hildebrandt, Magnetic field of a rotating superconductor, *Phys. Rev. Lett.* **12**, 190 (1964).
- [42] J. E. Zimmerman and J. E. Mercereau, Compton wavelength of superconducting electrons, *Phys. Rev. Lett.* **14**, 887 (1965).
- [43] N. F. Brickman, Rotating superconductors, *Phys. Rev.* **184**, 460 (1969).
- [44] J. Tate, B. Cabrera, S. B. Felch, and J. T. Anderson, Precise determination of the Cooper-pair mass, *Phys. Rev. Lett.* **62**, 845 (1989).
- [45] J. Tate, S. B. Felch, and B. Cabrera, Determination of the Cooper-pair mass in niobium, *Phys. Rev. B* **42**, 7885 (1990).
- [46] A. Verheijen, J. van Ruitenbeek, R. de Bruyn Ouboter, and L. de Jongh, The London moment for high temperature superconductors, *Physica B: Condensed Matter* **165-166**, 1181 (1990), IT-19.
- [47] A. A. Verheijen, J. M. van Ruitenbeek, R. de Bruyn Ouboter, and L. J. de Jongh, Measurement of the London moment in two high-temperature superconductors, *Nature* **345**, 418 (1990).
- [48] M. A. Sanzari, H. L. Cui, and F. Karwacki, London moment for heavy-fermion superconductors, *Applied Physics Letters* **68**, 3802 (1996).
- [49] R. Peierls, Spontaneously broken symmetries, *J. Phys. A* **24**, 5273 (1991).
- [50] A. J. Leggett, *Quantum Liquids: Bose Condensation And Cooper Pairing in Condensed-matter Systems* (Oxford Univ. Press, Oxford, 2006).
- [51] G. C. Wick, A. S. Wightman, and E. P. Wigner, Superselection rule for charge, *Phys. Rev. D* **1**, 3267 (1970).
- [52] R. Peierls, Broken symmetries, *Contemporary Phys.* **33**, 221 (1992).
- [53] B. D. Josephson, Supercurrents through barriers, *Adv. Phys.* **14**, 419 (1965).
- [54] K. Ueda, S. Matsuo, H. Kamata, Y. Sato, Y. Takeshige, K. Li, L. Samuelson, H. Xu, and S. Tarucha, Evidence of half-integer shapiro steps originated from nonsinusoidal current phase relation in a short ballistic inas nanowire josephson junction,

- Phys. Rev. Research **2**, 033435 (2020).
- [55] H. Zhang, M. W. A. de Moor, J. D. S. Bommer, D. Xu, G. Wang, N. van Loo, C.-X. Liu, S. Gazibegovic, J. A. Logan, D. Car, R. L. M. O. het Veld, P. J. van Veldhoven, S. Koelling, M. A. Verheijen, M. Pendharkar, D. J. Pennachio, B. Shojaei, J. S. Lee, C. J. Palmstrøm, E. P. A. M. Bakkers, S. D. Sarma, and L. P. Kouwenhoven, Large zero-bias peaks in InSb-Al hybrid semiconductor-superconductor nanowire devices, (2021), arXiv:2101.11456 [cond-mat.mes-hall].
- [56] H. Koizumi, Explanation of superfluidity using the Berry connection for many-body wave functions, J. Supercond. Nov. Magn. **33**, 1697 (2020).
- [57] H. Koizumi, Reversible superconducting-normal phase transition in a magnetic field and the existence of topologically-protected loop currents that appear and disappear without Joule heating, EPL **131**, 37001 (2020).
- [58] P. G. de Gennes, *Superconductivity of Metals and Alloys* (W. A. Benjamin, Inc., 1966).
- [59] J.-X. Zhu, *Bogoliubov-de Gennes Method and Its Applications* (Springer, 2016).
- [60] P. W. Anderson, *The Theory of Superconductivity in the High-Tc Cuprates* (Princeton Univ. Press, Princeton, 1997).
- [61] S. Miyaki, K. Makoshi, and H. Koizumi, Two-copper-atom units induce a pseudo jahn-teller polaron in hole-doped cuprate superconductors, J. Phys. Soc. Jpn. **77**, 034702 (2008).
- [62] J. M. Tranquada, H. Woo, T. G. Perring, H. Goka, G. D. Gu, G. Xu, M. Fujita, and K. Yamada, Quantum magnetic excitations from stripes in copper oxide superconductors, Nature **429**, 534 (2004).
- [63] A. Okazaki, H. Wakaura, H. Koizumi, M. A. Ghantous, and M. Tachiki, Superconducting transition temperature of the hole-doped cuprate as the stabilization temperature of super-current loops generated by spin-twisting itinerant motion of electrons, J. Supercond. Nov. Magn. **28**, 3221 (2015).
- [64] J. A. Riera, Spin polarization in the hubbard model with rashba spin-orbit coupling on a ladder, Phys. Rev. B **88**, 045102 (2013).
- [65] T. Morisaki, H. Wakaura, and H. Koizumi, Effect of rashba spin-orbit interaction on the stability of spin-vortex-induced loop current in hole-doped cuprate superconductors, J. Phys. Soc. Jpn. **86**, 104710 (2017).
- [66] F. C. Zhang and T. M. Rice, Effective hamiltonian for the superconducting cu oxides,

- Phys. Rev. B **37**, 3759 (1988).
- [67] A. Damascelli, Z. Hussain, and Z.-X. Shen, Angle-resolved photoemission studies of the cuprate superconductors, Rev. Mod. Phys. **75**, 473 (2003).
- [68] R. Car and M. Parrinello, Unified approach for molecular dynamics and density-functional theory, Phys. Rev. Lett. **55**, 2471 (1985).
- [69] D. Bohm, Note on a theorem of Bloch concerning possible causes of superconductivity, Phys. Rev **75**, 502 (1949).
- [70] D. Manabe and H. Koizumi, Supercurrent generation by spin-twisting itinerant motion of electrons, J. Supercond. Nov. Magn. **32**, 2303 (2019).
- [71] F. Bloch, Some remarks on the theory of superconductivity, Physics Today **19**, 27 (1966).
- [72] M. V. Berry, Quantal phase factors accompanying adiabatic changes, Proc. Roy. Soc. London Ser. A **391**, 45 (1984).
- [73] R. P. Feynman, *Quantum mechanics and path integrals* (McGraw-Hill companies, Inc., 1965).
- [74] Y. Aharonov and D. Bohm, Significance of electromagnetic potentials in the quantum theory, Phys. Rev. **115**, 167 (1959).
- [75] P. Dirac, Quantised singularities in the electromagnetic field, Proc. Roy. Soc. London **133**, 60 (1931).
- [76] E. I. Rashba, Sov. Phys. Solid State **2**, 1109 (1960).
- [77] G. Sundaram and Q. Niu, wave-packet dynamics in slowly perturbed crystals, Phys Rev. B **59**, 14915 (1999).
- [78] L. Onsager, Interpretation of the de Haas-van Alphen effect, Phil. Mag. Ser. 7 **43**, 1006 (1952).
- [79] A. K. Kerman and S. E. Koonin, Hamiltonian formulation of time-dependent variational principles for the many-body system, Ann. Phys. **100**, 332 (1976).
- [80] P. Carruthers and M. M. Nieto, Phase and angle variables in quantum mechanics, Rev. Mod. Phys. **40**, 411 (1968).