

Kramers' degeneracy for open systems in thermal equilibrium

Simon Lieu,^{1,2} Max McGinley,^{3,4} Oles Shtanko,^{1,2} Nigel R. Cooper,^{4,5} and Alexey V. Gorshkov^{1,2}

¹Joint Quantum Institute, NIST/University of Maryland, College Park, Maryland 20742, USA

²Joint Center for Quantum Information and Computer Science,
NIST/University of Maryland, College Park, Maryland 20742 USA

³Rudolf Peierls Centre for Theoretical Physics, Clarendon Laboratory, Oxford OX1 3PU, UK

⁴TCM Group, Cavendish Laboratory, University of Cambridge, JJ Thomson Avenue, Cambridge, CB3 0HE, UK

⁵Department of Physics and Astronomy, University of Florence, Via G. Sansone 1, 50019 Sesto Fiorentino, Italy
(Dated: May 10, 2021)

Kramers' degeneracy theorem underpins many interesting effects in quantum systems with time-reversal symmetry. We show that the generator of dynamics for Markovian open fermionic systems can exhibit an analogous degeneracy, protected by a combination of time-reversal symmetry *and* the microreversibility property of systems at thermal equilibrium—the degeneracy is lifted if either condition is not met. We provide simple examples of this phenomenon and show that the degeneracy is reflected in the standard Green's functions. Furthermore, we show that certain experimental signatures of topological edge modes in open many-body systems can be protected by microreversibility in the same way. Our results suggest that time-reversal symmetry of the system-bath Hamiltonian can affect open system dynamics only if the bath is in thermal equilibrium.

Kramers' theorem is one of the oldest and most celebrated theorems in quantum mechanics. It states that the Hamiltonian of a system with half-integer total spin will have a pairwise degenerate spectrum if time-reversal symmetry (TRS) is preserved [1, 2]. This simple claim carries deep implications, ranging from the quantum spin Hall effect [3, 4] to the robustness of superconductivity in disordered materials [5].

In this paper, we present an analogous degeneracy theorem that pertains to *dissipative* many-body fermionic systems. Dissipative systems propagate irreversibly in time, i.e. arbitrary initial states flow toward a (typically unique) steady state. Therefore it is not *a priori* obvious that TRS should have any bearing on dynamics in this context. Nevertheless, we show that in Markovian open systems—which are governed by a Lindblad master equation with generator \mathcal{L} —the spectrum of \mathcal{L} can feature Kramers-like degeneracies protected by a combination of TRS of the system-bath Hamiltonian *and* the microreversibility property of systems at thermal equilibrium [6–12]. An analogous non-equilibrium open system (e.g. one coupled to two reservoirs held at different temperatures) will not exhibit this degeneracy; see Fig. 1.

Many modern experimental platforms (including noisy quantum simulators) operate in a regime where coupling to an external environment cannot be ignored. Our work provides a precise description of how Kramers'-protected phenomena can survive coupling to a thermal bath. We show that, rather than being manifest in static properties of the steady state, these degeneracies can instead be inferred from time-dependent quantities describing the dynamical response about and/or approach to thermal equilibrium. As an example, we show that fermionic Green's functions and other related correlators [13] show signatures of the degeneracy, making our results directly observable in the solid state [14] and atomic systems [15].




FIG. 1. (a) A fermionic system (three small circles) coupled to an environment (big circle) in thermal equilibrium at temperature β^{-1} has a Kramers' degeneracy. (b) The same system (three small circles) coupled to two baths (two big circles) at different temperatures β_1^{-1} , β_2^{-1} will not be Kramers'-degenerate. This leads to a spin splitting of the Green's function (13), which can be experimentally detected using, e.g. spin-resolved tunneling spectroscopy (bottom panels). For both (a) and (b), the system-bath Hamiltonian obeys TRS: $[H_{SB}, T] = 0$.

One of the many areas where Kramers' theorem plays an important role is in topological phases of matter, where certain kinds of gapless edge modes can be attributed to TRS-protected degeneracies [3, 4, 16]. Although the irreversible effects of coupling such systems to an environment can spoil some of their TRS-protected properties [17], here we demonstrate that certain experimental signatures of TRS-protected edge modes can persist in the open regime.

These results complement recent work regarding symmetries and topological phenomena in open systems [18–30]. Many of those studies are based on non-Hermitian Hamiltonians [31–38], for which degeneracy theorems akin to Kramers' are known [39–41]. Our find-

ings are distinct from such ‘non-Hermitian topological phenomena’, which are primarily of relevance to photonic and classical mechanical systems [42]. Indeed, the Lindblad master equation formalism used here is much more widely applicable to dissipative quantum matter.

The study of microreversibility (or quantum detailed balance) in Markovian systems has a long and rich history [6–10], with relevance for open fluctuation theorems [43–45] and quantum thermodynamics [46–48]. Our work draws a connection between this mature field and the blossoming area of open topological systems.

Kramers’ theorem for Hamiltonians.—We review Kramers’ theorem in closed fermionic systems. Since fermions have half-integer spin, time reversal is implemented by an antiunitary operator T satisfying

$$T^2 = P, \quad P = (-1)^{\hat{N}}, \quad [H, T] = 0, \quad (1)$$

where \hat{N} is the total fermion number, and P is henceforth referred to as ‘parity’. Note that $[P, T] = 0$. Physical Hamiltonians must conserve parity, $[H, P] = 0$, and so Fock space can be decomposed into even and odd parity sectors $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$, wherein H and T are block diagonal

$$H = \begin{pmatrix} H_+ & 0 \\ 0 & H_- \end{pmatrix}, \quad T = \begin{pmatrix} U_+ & 0 \\ 0 & U_- \end{pmatrix} K. \quad (2)$$

Here, K takes the complex conjugate of any scalar to its right, and $U_{\pm}U_{\pm}^* = \pm\mathbb{I}$.

Kramers’ theorem states that for a TRS-invariant Hamiltonian (i.e. when $THT^{-1} = H$), the eigenvalues of H_- must be twofold degenerate. Specifically, any eigenstate $|\psi_-\rangle$ with odd parity has a time-reversed partner $T|\psi_-\rangle$ which is also an eigenstate of the same energy and is orthogonal to $|\psi_-\rangle$. In contrast, H_+ is generically non-degenerate.

Symmetries of open quantum systems.—When we come to study open quantum systems, it will be necessary to describe the state of the system using a density operator ρ whose dynamics is generated by a superoperator, rather than a pure wavefunction evolving under a Hamiltonian. For the purposes of this Letter, we will focus on Markovian open systems, whose dynamics is governed by a Lindblad master equation [49, 50]

$$\frac{d\rho}{dt} = \mathcal{L}(\rho) = -i[H, \rho] + \sum_i \left(2L_i\rho L_i^\dagger - \{L_i^\dagger L_i, \rho\} \right). \quad (3)$$

Here, the Hamiltonian part H describes the coherent evolution of the system, and the dissipators L_i are generically non-Hermitian operators describing non-unitary evolution due to coupling to an environment. The generating superoperator \mathcal{L} is often referred to as the ‘Lindbladian’.

In moving from closed to open systems, the formalisms we use to describe symmetries must be altered. We can define time-reversal and parity superoperators (\mathcal{T} and \mathcal{P} , respectively) which specify how

these symmetries transform the state $\rho(t)$. These act as $\mathcal{T}[\rho] = T\rho T^{-1}$ and $\mathcal{P}[\rho] = P\rho P^{-1}$. The space of operators can be split into even and odd superparity sectors $\mathcal{B}(\mathcal{H}) = \mathcal{B}(\mathcal{H})_+ \oplus \mathcal{B}(\mathcal{H})_-$, where $\mathcal{B}(\mathcal{H})_{\pm}$ contains operators satisfying $\mathcal{P}(A) = \pm A$. Specifically, $\mathcal{B}(\mathcal{H})_+$ contains operators of the form $|\text{even}\rangle\langle\text{even}|$ or $|\text{odd}\rangle\langle\text{odd}|$, and operators in $\mathcal{B}(\mathcal{H})_-$ are $|\text{even}\rangle\langle\text{odd}|$ or $|\text{odd}\rangle\langle\text{even}|$, which are traceless. (We use the convention $P|\text{even}\rangle = +|\text{even}\rangle$, $P|\text{odd}\rangle = -|\text{odd}\rangle$.) Since coherent superpositions of states with opposite fermion parity are not physical, the system density matrix must belong to $\mathcal{B}(\mathcal{H})_+$ (note that this does not prohibit classical mixtures of wavefunctions with opposite parity). Accordingly, physical generators must satisfy $[\mathcal{L}, \mathcal{P}] = 0$. By analogy to Eq. (2), the matrix representations of \mathcal{L} and \mathcal{T} then become block diagonal:

$$\mathcal{L} = \begin{pmatrix} \mathcal{L}_+ & 0 \\ 0 & \mathcal{L}_- \end{pmatrix}, \quad \mathcal{T} = \begin{pmatrix} U_+ & 0 \\ 0 & U_- \end{pmatrix} K. \quad (4)$$

From Eq. (1), we have $\mathcal{T}^2 = \mathcal{P}$, and hence $U_{\pm}U_{\pm}^* = \pm\mathbb{I}$. Evidently, the superoperator \mathcal{T} acts projectively on $\mathcal{B}(\mathcal{H})_-$, leaving operators invariant only up to a (-1) phase. This contrasts with systems made up of bosonic or spin degrees of freedom, where $T^2 = \pm 1$, and hence $\mathcal{T}^2 = \mathbb{I}$, even for half-integer spins.

Since $\rho(t)$ belongs to $\mathcal{B}(\mathcal{H})_+$, it is often stated that the odd-superparity part of the Lindbladian \mathcal{L}_- is unphysical. However, this is not true if we ask about the joint state of two fermionic systems. If system S and some probe R evolve independently under Lindbladians $\mathcal{L}_{S,R}$, then their joint state $\rho_{SR}(t)$ evolves as $\partial_t \rho_{SR} = (\mathcal{L}_S \otimes \text{id}_R + \text{id}_S \otimes \mathcal{L}_R)[\rho_{SR}]$. When the two are coupled, the overall superparity $\mathcal{P}_{\text{tot}} = \mathcal{P}_S \mathcal{P}_R$ must still be even (\mathcal{P}_S and \mathcal{P}_R are superparities for S and R , respectively), however the state $\rho_{SR}(t)$ may contain components which are odd under \mathcal{P}_S and \mathcal{P}_R separately, e.g. if a fermion is in a superposition between S and R . These components will evolve under $\mathcal{L}_{S,-}$ and $\mathcal{L}_{R,-}$. Later on, we will identify specific physical observables and experimental protocols that can be used to infer properties of \mathcal{L}_- (see also the Supplemental Material [51]).

Having discussed the mathematical structure of fermion parity symmetry and TRS in open systems, we will now establish a degeneracy theorem analogous to Kramers’, which applies to the spectrum of \mathcal{L}_- .

Kramers’ degeneracy for Lindbladians.—Because open quantum systems evolve irreversibly in time, TRS cannot be expected to play the same role as in closed systems. Indeed, if the Hamiltonian H is TRS-invariant in the sense of Eq. (1), then we have $\mathcal{T}^{-1}\mathcal{L}\mathcal{T} = -\mathcal{L}$. This relation is incompatible with a non-trivial dissipative part of Eq. (3), which is negative semi-definite [52]. One way in which an antiunitary symmetry can be imposed on open systems without such an inconsistency is to demand that the Hamiltonian be *odd* under TRS, i.e. $THT^{-1} = -H$, in which case $\mathcal{T}^{-1}\mathcal{L}\mathcal{T} = \mathcal{L}$ can be satisfied [53]. Although

such a symmetry is mathematically well-defined, it does not physically correspond to time-reversal in the closed system limit and suggests fine tuning of dynamics.

Instead, we observe that for systems in thermal equilibrium, time reversal symmetry of the system and environment degrees of freedom naturally gives rise to a “microreversibility” property, otherwise known as detailed balance. This condition relates the rate of each possible physical process to the rate of its time-reversed process. Mathematically, a Lindbladian that respects detailed balance satisfies a superoperator equation [6, 7, 9]

$$\mathcal{L}^\dagger = \mathcal{Q}^{-1} \mathcal{T}^{-1} \mathcal{L} \mathcal{T} \mathcal{Q}, \quad (5)$$

where \mathcal{Q} acts as $\mathcal{Q}[A] = qA$, and $q = \exp[-\beta H]/Z$ is the density matrix in the Gibbs ensemble at inverse temperature β with respect to the system Hamiltonian H (Z is the partition function). We have defined

$$\mathcal{L}^\dagger[\rho] = +i[H, \rho] + \sum_i \left(2L_i^\dagger \rho L_i - \{L_i^\dagger L_i, \rho\} \right), \quad (6)$$

which is the usual adjoint of \mathcal{L} with respect to the Hilbert-Schmidt inner product $\langle A, B \rangle := \text{Tr}[A^\dagger B]/\text{Tr} \mathbb{I}$.

A simple example of jump operators satisfying (5) is a pair $L_1 = \sqrt{\gamma_1} V$, $L_2 = \sqrt{\gamma_2} V^\dagger$, where V is TRS-invariant up to a phase, $\mathcal{T}[V] = e^{i\theta} V$ [54], and acts as a lowering operator $[H, V] = -\omega V$. The temperature is implicitly determined by $\gamma_1/\gamma_2 = e^{\beta\omega}$. In words, L_1 lowers the system energy by ω at a given rate, and L_2 does the opposite at a rate that differs by a Boltzman factor. This is the essence of the detailed balance condition. More generally, Eq. (5) is naturally satisfied when a time-reversal symmetric system is coupled to TRS-respecting environment at thermal equilibrium with temperature β^{-1} (we show this explicitly in the Supplemental Material [51]).

We now state the main claim of this work: Suppose a fermionic Lindbladian satisfies microreversibility via Eq. (5); then the odd-superparity superoperator \mathcal{L}_- [see Eq. (4)] is guaranteed to have a twofold degenerate spectrum. Our proof proceeds as follows. The odd parity part of the Lindbladian satisfies

$$\mathcal{L}_-^\dagger = \mathcal{Q}_-^{-1} \mathcal{T}_-^{-1} \mathcal{L}_- \mathcal{T}_- \mathcal{Q}_-, \quad (7)$$

where $\mathcal{T}_-^2 = -\mathbb{I}$, $\mathcal{T}_- = \mathcal{U}_- K$. We drop the “-” index. Define right and left eigenoperators

$$\mathcal{L}(r_i) = \Lambda_i r_i, \quad \mathcal{L}^\dagger(l_i) = \Lambda_i^* l_i, \quad (8)$$

with $\text{Tr}[l_i^\dagger r_j] = \delta_{ij}$. Substituting the expression for microreversibility into the left eigenoperator equation leads to

$$\mathcal{Q}^{-1} \mathcal{T}^{-1} \mathcal{L} \mathcal{T} \mathcal{Q}(l_i) = \Lambda_i^* l_i \Rightarrow \mathcal{L} \mathcal{T} \mathcal{Q}(l_i) = \Lambda_i \mathcal{T} \mathcal{Q}(l_i). \quad (9)$$

We find that r_i and $\mathcal{T} \mathcal{Q}(l_i)$ are both right eigenoperators of \mathcal{L} with eigenvalue Λ_i . However, one can show that

$\text{Tr}[l_i^\dagger \mathcal{T} \mathcal{Q}(l_i)] = 0$ (see the Supplemental Material [51]). Since $\text{Tr}[l_i^\dagger r_i] = 1$, we find that r_i and $\mathcal{T} \mathcal{Q}(l_i)$ are linearly independent eigenoperators, and hence the complex Lindblad spectrum in the odd superparity sector must be twofold degenerate.

In contrast to the Kramers’ theorem for closed systems, our result explicitly relies on the presence of thermal equilibrium. Intuition can be gained from microscopic considerations: Linear response about a thermal state can be formulated in terms of the (Kramers’ degenerate) eigenstates of the system-bath Hamiltonian. A system’s response in thermal equilibrium should thus be sensitive to the TRS of the microscopic Hamiltonian. Our work captures this behavior from the perspective of the system’s master equation.

Our analysis also suggests that a similar Kramers’ degeneracy is present in the spectrum of a thermal quantum channel superoperator, which can describe the evolution of a system coupled to a non-Markovian bath [51].

Example: random quadratic Hamiltonian.—We confirm the generalized Kramers’ theorem via an example. Consider a system of spin-1/2 fermions with N twofold degenerate single-particle orbitals. The most general particle-conserving quadratic Hamiltonian is

$$H = \sum_{ij;\sigma,\sigma'} H_{ij,\sigma\sigma'} f_{i,\sigma}^\dagger f_{j,\sigma'} = \sum_{k=1}^N \sum_{\tau=\pm} \epsilon_k c_{k,\tau}^\dagger c_{k,\tau} \quad (10)$$

with $i, j \in [1, \dots, N]$ and $\sigma, \sigma' \in \pm$. We impose a TRS on the Hamiltonian: $[H, T] = 0$, such that the single-particle spectrum is twofold degenerate, with $T f_{i,\sigma} T^{-1} = \sigma f_{i,-\sigma}$, $T c_{k,\tau} T^{-1} = \tau c_{k,-\tau}$. Let us define the following dissipators

$$L_{1,pq} = \sqrt{\gamma_{1,pq}} \sum_{\tau=\pm} \left(c_{q,\tau}^\dagger c_{p,\tau} + \tau c_{q,\tau}^\dagger c_{p,-\tau} \right), \quad (11)$$

$$L_{2,pq} = \sqrt{\gamma_{2,pq}} \sum_{\tau=\pm} \left(c_{p,\tau}^\dagger c_{q,\tau} + \tau c_{p,-\tau}^\dagger c_{q,\tau} \right), \quad (12)$$

for $\gamma_{1,pq} = g[n_\beta(\epsilon_p - \epsilon_q) + 1]$, $\gamma_{2,pq} = g n_\beta(\epsilon_p - \epsilon_q)$, $n_\beta(\omega) = (e^{\beta\omega} - 1)^{-1}$ [the Bose function], and $\epsilon_p > \epsilon_q$. Physically, such terms can appear due to coupling to a bosonic bath (since the dissipators are quadratic in fermions). L_1 represents a process that lowers the energy of the system; L_2 raises the energy. The L s satisfy: $[H, L_{1,pq}] = (\epsilon_q - \epsilon_p) L_{1,pq}$, $L_{2,pq} \sim L_{1,pq}^\dagger$, $\gamma_{1,pq}/\gamma_{2,pq} = e^{\beta(\epsilon_p - \epsilon_q)}$, and $[L_{1/2,pq}, T] = 0$, thus respecting microreversibility [see Fig. 2(a)]. We include these dissipators between each pair of energy levels in the system, and also consider uniform dephasing in the energy basis: $L_{d,i,\pm} = \sqrt{\Delta_d} c_{i,\pm}^\dagger c_{i,\pm}$ (preserves microreversibility). For a fixed number of fermions in the system, the thermal state is the unique steady state.

Fig. 2(b) plots the Lindblad spectrum associated with a random TRS-respecting Hamiltonian ($|H_{ij}| \in [0, \Delta]$; Δ sets the energy scale) coupled to a thermal bath for




FIG. 2. (a) Three doubly degenerate energy levels ($N = 3$) labeled by $\epsilon_{i,\pm}$. TRS-respecting dissipators cause jumps between energy levels at a rate consistent with detailed balance in thermal equilibrium at temperature β^{-1} . (b) Lindblad spectrum for the model described in the main text (for states with 0, 1, or 2 occupied fermions) at a temperature $\beta\Delta = 1$, with $N = 3$, $\Delta_d/\Delta = 0.1$, and $g/\Delta = 1$. The odd-superparity sector (blue and red dots) is twofold degenerate, while the even sector (black dots) is not. (c) Real part of the retarded Green's function for same parameters as in (b). The Kramers' degeneracy ensures that $G_{i,\sigma} = G_{i,-\sigma}$ in the expectation value of the thermal steady state with one occupied fermionic mode: black line and red dashed line correspond to $\text{Re}[G_{1,+}]$ and $\text{Re}[G_{1,-}]$, respectively. In general, Green's function pairs other than $G_{i,\pm}$ are non-degenerate: blue line corresponds to $\text{Re}[G_{2,+}]$. (d) A system that is coupled to *two* thermal baths $g_1/\Delta = 1, g_2/\Delta = 0.4$ at different temperatures: $\beta_1\Delta = 1, \beta_2\Delta = 10$ [otherwise same parameters as in (b)]. Microreversibility is broken, hence the Green's functions for $i, +$ and $i, -$ split.

states with up to two occupied fermionic modes. The spectrum of the odd superparity sector (blue and red dots) is indeed twofold degenerate, whilst the spectrum of the even sector (black dots) is not. Note that the Kramers' theorem does not imply a degeneracy of the steady state.

How can an open system violate microreversibility (and hence break its degeneracy)? One obvious way involves a system-environment coupling that directly violates TRS. However, a more subtle way to break degeneracy involves coupling the system to a non-equilibrium environment. Consider a system that is connected to *two* thermal baths, each at a different temperature. Even if all system-environment couplings respect TRS, the system will host a *non-equilibrium* (non-thermal) steady state, and hence microreversibility will be violated. We have numerically verified that the degeneracy of \mathcal{L}_- for the spin-1/2 system is broken when coupled to baths at different temperatures. [We note that such non-equilibrium splitting requires quartic terms (quadratic dissipators) in the master equation (see SM [51]).]

Physical observables.—As argued above, \mathcal{L}_- only governs dynamics in scenarios where fermions can move between the system (S) and some external 'probe' (R). Therefore, rather than looking at expectation values of the system density matrix $\text{Tr}[O_S \rho_S(t)]$ (which are determined by \mathcal{L}_+), we suggest that the Kramers'-like degeneracy of \mathcal{L}_- can be detected in the retarded Green's function of the steady state ρ_{SS} :

$$G_{i,\sigma}^R(t) = -i\Theta(t)\text{Tr}\{f_{i,\sigma}(t), f_{i,\sigma}^\dagger\}\rho_{\text{SS}}. \quad (13)$$

Here, $\Theta(t)$ is the Heaviside step function, and we work in the Heisenberg picture where operators evolve as $A(t) = e^{\mathcal{L}^\dagger t}[A]$ (although this expression should be slightly modified for open systems coupled to fermionic baths; see the Supplemental Material [51]). The Green's function can be measured in solid state systems using, e.g. photoemission or tunnelling spectroscopy, which indeed involve fermions moving in/out of the system; analogous methods have also been developed for ultracold atoms [15]. More concretely, $G_{i,\sigma}^R(t)$ is sensitive to the Kramers' degeneracy of \mathcal{L}_- because $f_{i,\sigma} \in \mathcal{B}(\mathcal{H})_-$ is superparity-odd, and so the time evolution of (13) is governed by \mathcal{L}_- . We show in the Supplemental Material [51] that the generalized Kramers' theorem ensures the relation: $G_{i,\sigma}^R(t) = G_{i,-\sigma}^R(t)$, which we confirm numerically in Fig. 2(c). However, when the system is coupled to two baths at different temperatures, microreversibility is broken and the Green's functions differ for opposite spins [Fig. 2(d)]. Evidently, the spin-degeneracy of the Green's function can be used as a diagnostic of thermal equilibrium in fermionic systems.

Degenerate zero-bias peak from microreversibility.—Kramers' degeneracy plays an important role in determining the stability of symmetry-protected topological edge modes of Hamiltonians [3, 55], e.g. the presence of spinful TRS ensures that a pair of Majorana zero modes in 1D cannot gap. For closed systems, degenerate Majorana modes are detectable via degenerate spin-resolved tunneling spectroscopy at the edge of the superconductor [13]. Here, we show that the spin-resolved tunneling spectra remain unsplit if the superconductor is coupled to a thermal bath with TRS-respecting terms, while a splitting can arise if microreversibility is violated.

Consider the following spin-1/2 Kitaev chain

$$H = -i \sum_{j=1, \sigma=\pm}^{j=N} [u_1 a_{j,\sigma} b_{j,\sigma} + u_2 a_{j,\sigma} b_{j,-\sigma}] - i \sum_{j=1, \sigma=\pm}^{j=N-1} [v_1 b_{j,\sigma} a_{j+1,\sigma} + v_2 b_{j,\sigma} a_{j+1,-\sigma}], \quad (14)$$

where $a_{j,\sigma}, b_{j,\sigma}$ are Majoranas corresponding to site j with spin σ . This model can be diagonalized as: $H = \sum_{j=1}^N \sum_{\tau=\pm} \epsilon_j d_{j,\tau}^\dagger d_{j,\tau}$, where $T d_{j,\tau} T^{-1} = \tau d_{j,-\tau}$. We include the same thermal dissipators as before [see Eq. (11)]




FIG. 3. Steady-state, spin-resolved spectral function at the boundary of the Majorana chain (14) coupled to two baths found via exact diagonalization. (a) Zero-bias peak due to the Majorana modes. The width of the peaks is set by the temperature. (b) Splitting between spin-up and spin-down spectral functions due to breaking of microreversibility via a non-equilibrium environment. Parameters: $N = 3, u_1/v_1 = 0.25, u_2/v_1 = 0.25, v_2/v_1 = 0.5, \beta v_1 = 3.3, g/v_1 = 0.05, g_{\text{neq}}/v_1 = 2$.

between each pair of energy levels p, q in the system at inverse temperature β . To break microreversibility, we consider a non-equilibrium bath that removes pairs of fermions on each site: $L_j = \sqrt{g_{\text{neq}}}\psi_{j,+}\psi_{j,-}$ where we define the complex fermion: $\psi_{j,\sigma} = a_{j,\sigma} + ib_{j,\sigma}$. These dissipators obey $[L_j, T] = 0$ but do not evolve the system toward a thermal state.

The splitting of spectral degeneracy can be experimentally detected using spin-resolved tunneling spectroscopy [14] (see Fig. 1). For a zero-temperature probe, the current-voltage characteristic of the tunnel junction is $\partial I_\sigma/\partial\mu \propto A_{1,\sigma}(-\mu)$, where $A_{i,\sigma}(\omega) = -\text{Im}[G_{i,\sigma}^R(\omega)]$ is the spectral function, I_σ is spin- σ current and μ is the chemical potential (see SM [51]). Fig. 3 plots the spin-up spectral function for the probe attached to the boundary, and the relative difference between the spin-up and spin-down spectral functions for the chain coupled to two baths. The zero-bias peak in the topological phase correspond to the boundary-localized Majoranas. The splitting that emerges between the spin up and down spectral functions is due to the non-equilibrium setup and is a direct evidence of breaking of Kramers' degeneracy. This splitting vanishes if the non-equilibrium bath is turned off and microreversibility is restored (not shown). Therefore, breaking of Kramers' degeneracy without a TRS-breaking Hamiltonian is observable in the experiment.

Conclusions and outlook.—Our work provides direct evidence that microreversibility is the natural generalization of time-reversal symmetry to open systems: For open systems, Kramers' degeneracy relies on both (1) TRS of the system-bath Hamiltonian and (2) thermal equilibrium of the bath. Future work should explore how different effects protected by TRS in closed systems generalize to open systems. For example, can the steady state of an open system spontaneously break microreversibility? This would lead to an enlarged steady state manifold that could be useful to store quantum infor-

mation [56]. Similarly, TRS plays an important role in symmetry-protected topological phases in closed systems [16]. Future studies should investigate whether microreversibility has any implications for the topological properties of the steady-state density matrix. Finally, we note that microreversibility may also have interesting implications for the random-matrix theory of open systems [57–60].

S.L. was supported by the NIST NRC Research Postdoctoral Associateship Award. M.M. was supported by an EPSRC studentship. O.S. and A.V.G. acknowledge funding by AFOSR MURI, NSF PFCQC program, AFOSR, U.S. Department of Energy Award No. DE-SC0019449, DoE ASCR Quantum Testbed Pathfinder program (award No. DE-SC0019040), DoE ASCR Accelerated Research in Quantum Computing program (award No. DE-SC0020312), and ARO MURI. NRC was supported by EPSRC Grants EP/P034616/1 and EP/P009565/1 and by an Investigator Award of the Simons Foundation.

-
- [1] H. A. Kramers, Koninkl. Ned. Akad. Wetenschap., Proc. **33**, 959 (1930).
 - [2] E. Wigner, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse **1932**, 546 (1932).
 - [3] C. L. Kane and E. J. Mele, *Phys. Rev. Lett.* **95**, 226801 (2005).
 - [4] B. A. Bernevig and S.-C. Zhang, *Phys. Rev. Lett.* **96**, 106802 (2006).
 - [5] P. Anderson, *J. Phys. Chem. Solids* **11**, 26 (1959).
 - [6] G. Agarwal, *Z. Phys. A* **258**, 409 (1973).
 - [7] W. A. Majewski, *J. Math. Phys.* **25**, 614 (1984).
 - [8] R. Alicki, *Rep. Math. Phys.* **10**, 249 (1976).
 - [9] R. Chetrite and K. Mallick, *J. Stat. Phys.* **148**, 480 (2012).
 - [10] F. Fagnola and V. Umantà, *Math. Notes* **84**, 108 (2008).
 - [11] D. Roberts, A. Lingenfelter, and A. Clerk, arXiv preprint arXiv:2011.02148 (2020).
 - [12] L. M. Sieberer, A. Chiochetta, A. Gambassi, U. C. Täuber, and S. Diehl, *Phys. Rev. B* **92**, 134307 (2015).
 - [13] H. Bruus and K. Flensberg, *Many-body quantum theory in condensed matter physics: an introduction* (Oxford university press, 2004).
 - [14] H. Oka, O. O. Brovko, M. Corbetta, V. S. Stepanyuk, D. Sander, and J. Kirschner, *Rev. Mod. Phys.* **86**, 1127 (2014).
 - [15] T.-L. Dao, A. Georges, J. Dalibard, C. Salomon, and I. Carusotto, *Phys. Rev. Lett.* **98**, 240402 (2007).
 - [16] F. Pollmann, A. M. Turner, E. Berg, and M. Oshikawa, *Phys. Rev. B* **81**, 064439 (2010).
 - [17] M. McGinley and N. R. Cooper, *Nature Physics* (2020).
 - [18] S. Diehl, E. Rico, M. A. Baranov, and P. Zoller, *Nature Physics* **7**, 971 (2011).
 - [19] S. Lieu, M. McGinley, and N. R. Cooper, *Phys. Rev. Lett.* **124**, 040401 (2020).
 - [20] M. van Caspel, S. E. T. Arze, and I. P. Castillo, *SciPost Phys* **6**, 26 (2019).

- [21] F. Tonielli, J. C. Budich, A. Altland, and S. Diehl, *Phys. Rev. Lett.* **124**, 240404 (2020).
- [22] C. Gneiting, A. Koottandavida, A. V. Rozhkov, and F. Nori, [arXiv:2007.05960](https://arxiv.org/abs/2007.05960) (2020).
- [23] F. Song, S. Yao, and Z. Wang, *Phys. Rev. Lett.* **123**, 170401 (2019).
- [24] T. Yoshida, K. Kudo, H. Katsura, and Y. Hatsugai, *Phys. Rev. Research* **2**, 033428 (2020).
- [25] C.-H. Liu, K. Zhang, Z. Yang, and S. Chen, *Phys. Rev. Research* **2**, 043167 (2020).
- [26] N. Okuma and M. Sato, [arXiv preprint arXiv:2011.08175](https://arxiv.org/abs/2011.08175) (2020).
- [27] S. Longhi, *Phys. Rev. B* **102**, 201103 (2020).
- [28] J.-S. Pan, L. Li, and J. Gong, [arXiv preprint arXiv:2010.14862](https://arxiv.org/abs/2010.14862) (2020).
- [29] S. Sayyad, J. Yu, A. G. Grushin, and L. M. Sieberer, [arXiv preprint arXiv:2011.10601](https://arxiv.org/abs/2011.10601) (2020).
- [30] A. McDonald, R. Hanai, and A. A. Clerk, [arXiv preprint arXiv:2103.01941](https://arxiv.org/abs/2103.01941) (2021).
- [31] K. Kawabata, K. Shiozaki, M. Ueda, and M. Sato, *Phys. Rev. X* **9**, 041015 (2019).
- [32] D. Leykam, K. Y. Bliokh, C. Huang, Y. D. Chong, and F. Nori, *Phys. Rev. Lett.* **118**, 040401 (2017).
- [33] K. Yokomizo and S. Murakami, *Phys. Rev. Lett.* **123**, 066404 (2019).
- [34] D. S. Borgnia, A. J. Kruchkov, and R.-J. Slager, *Phys. Rev. Lett.* **124**, 056802 (2020).
- [35] S. Yao and Z. Wang, *Phys. Rev. Lett.* **121**, 086803 (2018).
- [36] S. Malzard, C. Poli, and H. Schomerus, *Phys. Rev. Lett.* **115**, 200402 (2015).
- [37] L. Herviou, N. Regnault, and J. H. Bardarson, *SciPost Phys.* **7**, 69 (2019).
- [38] S. Lieu, *Phys. Rev. B* **98**, 115135 (2018).
- [39] M. Sato, K. Hasebe, K. Esaki, and M. Kohmoto, *Prog. Theor. Phys.* **127**, 937 (2012).
- [40] H. Zhou and J. Y. Lee, *Phys. Rev. B* **99**, 235112 (2019).
- [41] H. Kondo, Y. Akagi, and H. Katsura, [arXiv preprint arXiv:2006.10391](https://arxiv.org/abs/2006.10391) (2020).
- [42] E. J. Bergholtz, J. C. Budich, and F. K. Kunst, [arXiv:1912.10048](https://arxiv.org/abs/1912.10048) (2019).
- [43] J. Åberg, *Phys. Rev. X* **8**, 011019 (2018).
- [44] G. Manzano, J. M. Horowitz, and J. M. R. Parrondo, *Phys. Rev. X* **8**, 031037 (2018).
- [45] K. Temme, M. J. Kastoryano, M. B. Ruskai, M. M. Wolf, and F. Verstraete, *J. Math. Phys.* **51**, 122201 (2010).
- [46] R. Alicki, *J. Phys. A Math. Theor.* **12**, L103 (1979).
- [47] J. Goold, M. Huber, A. Riera, L. del Rio, and P. Skrzypczyk, *J. Phys. A Math. Theor.* **49**, 143001 (2016).
- [48] K. Brandner and U. Seifert, *Phys. Rev. E* **93**, 062134 (2016).
- [49] G. Lindblad, *Commun. Math. Phys.* **48**, 119 (1976).
- [50] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, *J. Math. Phys.* **17**, 821 (1976).
- [51] See the Supplemental Material for ***. Contains Refs. ***.
- [52] H. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2002).
- [53] A. Altland, M. Fleischhauer, and S. Diehl, [arXiv:2007.10448](https://arxiv.org/abs/2007.10448) (2020).
- [54] The phase θ carry no physical significance, since jump operators can be changed by a transform $L_i \rightarrow e^{i\chi} L_i$ without changing the Lindbladian (3).
- [55] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, *New J. Phys.* **12**, 065010 (2010).
- [56] S. Lieu, R. Belyansky, J. T. Young, R. Lundgren, V. V. Albert, and A. V. Gorshkov, *Phys. Rev. Lett.* **125**, 240405 (2020).
- [57] S. Denisov, T. Lapyeva, W. Tarnowski, D. Chruściński, and K. Życzkowski, *Phys. Rev. Lett.* **123**, 140403 (2019).
- [58] T. Can, V. Oganesyan, D. Orgad, and S. Gopalakrishnan, *Phys. Rev. Lett.* **123**, 234103 (2019).
- [59] L. Sá, P. Ribeiro, and T. c. v. Prosen, *Phys. Rev. X* **10**, 021019 (2020).
- [60] K. Wang, F. Piazza, and D. J. Luitz, *Phys. Rev. Lett.* **124**, 100604 (2020).
- [61] T. Prosen, *New J. Phys.* **10**, 043026 (2008).
- [62] B. Buča and T. Prosen, *New J. Phys.* **14**, 073007 (2012).
- [63] R. Kubo, *J. Phys. Soc. Jpn* **12**, 570 (1957).

Supplemental Material for “Kramers’ degeneracy for open systems in thermal equilibrium”

The Supplemental Material is organized as follows: In Sec. 1, we complete the steps in the proof of the Kramers’ theorem for Lindbladians. We also explain why the degeneracy only appears for fermionic systems (as opposed to spin systems) and describe an analogous Kramers’ degeneracy for thermal quantum channels. In Sec. 2, we show how the Kramers’ degeneracy manifests in the single-particle spectrum of “quadratic Lindbladians”. Sec. 3 shows that microreversibility of the Lindbladian arises naturally from TRS-invariant system-bath coupling in the case of a thermal bath. Sec. 4 describes linear response in open quantum systems, suggesting that the degeneracy can be probed via tunneling spectroscopy experiments.

1. KRAMERS’ THEOREM AND GREEN’S FUNCTION DEGENERACY

Orthogonal solutions. We complete the steps of the generalized Kramers’ theorem outlined in the main text. From the main text, we have seen that if the Lindbladian satisfies a microreversibility condition: $\mathcal{L}_-^\dagger = \mathcal{Q}_-^{-1}\mathcal{T}_-^{-1}\mathcal{L}_-\mathcal{T}_-\mathcal{Q}_-$ with $\mathcal{L}_-(r_i) = \Lambda_i r_i$, $\mathcal{L}_-^\dagger(l_i) = \Lambda_i^* l_i$, then r_i and $\mathcal{T}\mathcal{Q}(l_i)$ are both right eigenoperators of \mathcal{L}_- with eigenvalue Λ_i . We now proceed to show that these are indeed orthogonal solutions by showing that $\text{Tr}[l_i^\dagger \mathcal{T}\mathcal{Q}(l_i)] = 0$:

$$\text{Tr}[l_i^\dagger T(q l_i) T^{-1}] = \text{Tr}[(T(T(q l_i) T^{-1}) T^{-1})^\dagger T l_i T^{-1}] \quad (\text{S1})$$

$$= \text{Tr}[(T^2(q l_i) T^{-2})^\dagger T l_i T^{-1}] \quad (\text{S2})$$

$$= -\text{Tr}[(q l_i)^\dagger T l_i T^{-1}] \quad (\text{S3})$$

$$= -\text{Tr}[l_i^\dagger q T l_i T^{-1}] \quad (\text{S4})$$

$$= -\text{Tr}[l_i^\dagger T(q l_i) T^{-1}] \quad (\text{S5})$$

$$= 0. \quad (\text{S6})$$

In Eq. (S1), we have used the relation:

$$\text{Tr}[(T\psi T^{-1})^\dagger (T\phi T^{-1})] = \text{Tr}[(U\psi^* U^\dagger)^\dagger (U\phi^* U^\dagger)] \quad (\text{S7})$$

$$= \text{Tr}[U\psi^T U^\dagger U\phi^* U^\dagger] \quad (\text{S8})$$

$$= \text{Tr}[\psi^T \phi^*] \quad (\text{S9})$$

$$= \text{Tr}[\phi^\dagger \psi], \quad (\text{S10})$$

for any ψ, ϕ , where $T = UK$. In Eq. (S4), we have used $\mathcal{T}^2(q l_i) = q \mathcal{T}^2(l_i) = -q l_i$, i.e. l_i is an eigenoperator of the odd-superparity sector of the Lindbladian.

Definition of the Green’s function. Here we highlight an important subtlety regarding the definition of the Green’s function Eq. (13), and in particular the interpretation of the time-evolved fermion operator $\hat{f}_{i,\sigma}(t)$.

A general Green’s function of two observables $G_{AB}(t) = \text{Tr}(A(t)B\rho_{\text{SS}})$ describes the influence of a perturbation at time 0 on the outcome of a measurement at time t . Since fermion parity symmetry is fundamental, it is not possible to perturb the system by a fermion-parity-odd operator such as $f_{i,\sigma}^\dagger$; rather, in any physical protocol where the Green’s function is measured, the initial perturbation will involve some exchange of fermions between the system and some probe. Thus, we should understand that the operator B is a product of $\hat{f}_{i,\sigma}$ with some fermionic operator acting on the probe. (See Section 4 of the SM for an example of this construction in the context of tunneling spectroscopy.) We heuristically write $B \sim f_{i,\sigma}^\dagger f_{\text{pr}}$, where f_{probe} is an unspecified fermionic operator acting on the probe, ensuring that B itself is a superparity-even operator. Similarly, the observable A that we measure at time t will be a product of fermion-superparity-odd operators on the system and probe: $A \sim f_{i,\sigma} f_{\text{pr}}'$ for some other probe operator f_{pr}' .

With this in mind, the Heisenberg picture evolution of the observable A is given by $A(t) = e^{\mathcal{L}^\dagger t}[A]$, where \mathcal{L}^\dagger is defined in Eq. (6). The Lindbladian only involves operators acting on the system, and not the probe. Therefore, when considering the observable $A = f_{i,\sigma} f_{\text{pr}}'$, it is tempting to pull the probe operator f_{pr}' outside the evolution

superoperator, so that it can combine with the other probe operator in B . Indeed, this is precisely what is done when calculating fermionic Green's functions in closed systems where $\mathcal{L} = -i[H, \cdot]$. However, when the system is open, it is possible for fermions to move between the system and the environment (not to be confused with the probe), which leads to jump operators L_i that are superparity odd. In this case, we cannot ignore the presence of the probe operators, because $\mathcal{L}^\dagger[A]$ will include a term $L_i^\dagger f_{i,\sigma} f'_{\text{pr}} L_i$, which differs from the naïve expression $(L_i^\dagger f_{i,\sigma} L_i) f'_{\text{pr}}$ by a minus sign. In other words, even though f'_{pr} is a probe operator, its anticommutation with the jump operators means that the evolution of the product $f_{i,\sigma} f'_{\text{pr}}$ doesn't factorize as the product of time-evolved operators $f_{i,\sigma}(t)$ and f'_{pr} .

This can be remedied by defining a 'dummy' Majorana fermion operator η_d , having no dynamics of its own, and including it such that the correct anticommutation relations are obeyed. Specifically, one can verify that $\mathcal{L}^\dagger[f_{i,\sigma} f'_{\text{pr}}] = \eta_d \mathcal{L}^\dagger[\eta_d f_{i,\sigma}] f'_{\text{pr}}$, since $\eta_d^2 = 1$. This allows probe operators to be pulled out of the system evolution superoperator, such that Green's functions can be defined using operators that pertain to the system only. Specifically, the physically meaningful definition of the time-evolved operator $f_{i,\sigma}(t)$ appearing in Eq. (13) should be:

$$f_{i,\sigma}(t) := \eta_d e^{\mathcal{L}^\dagger t} [\eta_d f_{i,\sigma}]. \quad (\text{S11})$$

In practice, we will not explicitly write out the dummy Majorana fermion, but instead understand that it is implicitly included in any expression of the form $e^{\mathcal{L}^\dagger t} [f_{i,\sigma}]$. Alternatively, we can modify all the fermion-superparity-odd jump operators by $L_i \rightarrow \eta_d L_i$, in which case the usual expression $e^{\mathcal{L}^\dagger t} [f_{i,\sigma}]$ can be used without modification.

Degenerate Green's functions. Having dealt with the above issue, we now show that steady-state Green's functions corresponding to fermions with opposite spin are related in a simple way due to microreversibility. We first consider the quantity

$$\text{Tr}[e^{\mathcal{L}^\dagger t} (f_{i,\sigma}) f_{j,\tau}^\dagger q] = \text{Tr}[\mathcal{Q}(e^{\mathcal{L}^\dagger t} (f_{i,\sigma}) f_{j,\tau}^\dagger)] \quad (\text{S12})$$

$$= \text{Tr}[\mathcal{T} \mathcal{Q}(e^{\mathcal{L}^\dagger t} (f_{i,\sigma}) f_{j,\tau}^\dagger)]^* \quad (\text{S13})$$

$$= \text{Tr}[\mathcal{T} \mathcal{Q}(e^{\mathcal{L}^\dagger t} (f_{i,\sigma})) \mathcal{T}(f_{j,\tau}^\dagger)]^* \quad (\text{S14})$$

$$= \text{Tr}[e^{\mathcal{L} t} (\mathcal{T} \mathcal{Q}(f_{i,\sigma})) \mathcal{T}(f_{j,\tau}^\dagger)]^* \quad (\text{S15})$$

$$= \sigma \tau \text{Tr}[e^{\mathcal{L} t} (q f_{i,-\sigma}) f_{j,-\tau}^\dagger]^* \quad (\text{S16})$$

$$= \sigma \tau \text{Tr}[(q f_{i,-\sigma}) e^{\mathcal{L}^\dagger t} (f_{j,-\tau}^\dagger)]^*, \quad (\text{S17})$$

where we have used $\text{Tr}[A] = \text{Tr}[\mathcal{T}(A)]^*$ in (S13), $\mathcal{T}[AB] = \mathcal{T}[A]\mathcal{T}[B]$ in (S14), the definition of microreversibility in (S15), $\mathcal{T}[f_{i,\sigma}] = \sigma f_{i,-\sigma}$ in (S16), and $\text{Tr}[A e^{\mathcal{L} t} (B)] = \text{Tr}[e^{\mathcal{L}^\dagger t} (A) B]$ in (S17).

Let us now define the following generalizations of retarded Green's functions:

$$G_{i\sigma;j\tau} \equiv -i\Theta(t) \left(\text{Tr}[e^{\mathcal{L}^\dagger t} (f_{i,\sigma}) f_{j,\tau}^\dagger q] + \text{Tr}[f_{j,\tau}^\dagger e^{\mathcal{L}^\dagger t} (f_{i,\sigma}) q] \right) \quad (\text{S18})$$

$$= -i\Theta(t) \left(\sigma \tau \text{Tr}[f_{i,-\sigma} e^{\mathcal{L}^\dagger t} (f_{j,-\tau}^\dagger) q]^* + \sigma \tau \text{Tr}[e^{\mathcal{L}^\dagger t} (f_{j,-\tau}^\dagger) f_{i,-\sigma} q]^* \right) \quad (\text{S19})$$

$$= -i\Theta(t) \left(\sigma \tau \text{Tr}[e^{\mathcal{L}^\dagger t} (f_{j,-\tau}^\dagger) f_{i,-\sigma} q] + \sigma \tau \text{Tr}[e^{\mathcal{L}^\dagger t} (f_{j,-\tau}^\dagger) q f_{i,-\sigma}^\dagger] \right) \quad (\text{S20})$$

$$= -i\Theta(t) (\sigma \tau G_{j-\tau;i-\sigma}(t)), \quad (\text{S21})$$

where in (S20) we have used: $\text{Tr}[AB] = \text{Tr}[A^\dagger B^\dagger]^*$ and $[e^{\mathcal{L}^\dagger t} (A)]^\dagger = e^{\mathcal{L}^\dagger t} (A^\dagger)$. So indeed we find that steady-state Green's functions for opposite spin labels are related to each other in a simple way for systems with microreversibility. We have confirmed these expressions numerically for the example system described in the main text.

Non-fermionic systems. As mentioned in the main text, the Kramers' degeneracy of the Lindbladian only arises in fermionic systems, but not in bosonic or spin systems, even though the latter have a Kramers-degenerate Hamiltonian when the total spin is a half-integer. The differences between fermionic vs. non-fermionic open systems becomes apparent when we consider how linear response functions are constrained by microreversibility and TRS, in analogy to Eq. (S21). Regardless of particle statistics, detailed balance implies [Ref. [6], Eq. (2.2)]:

$$\text{Tr}[A(t) B q] = \text{Tr}[\tilde{B}(t) \tilde{A} q], \quad (\text{S22})$$

where $\tilde{A} := \mathcal{T}[A]^\dagger$ and similar for \tilde{B} . Suppose for the sake of simplicity that $\beta = 0$ and hence $q \propto \mathbb{I}$. Then, by decomposing A and B in terms of right and left eigenoperators of \mathcal{L} , respectively, the left-hand side of the above can

be written as a linear combination of terms $\text{Tr}[r_i(t)l_j] = e^{\Lambda_i t} \delta_{ij}$, and thus without loss of generality the condition becomes $\text{Tr}[r_i(t)l_i] = \text{Tr}[\tilde{l}_i(t)\tilde{r}_i]$. (The same arguments can in principle be generalized to $\beta > 0$ by defining a new set of operators l'_i which are left eigenoperators of \mathcal{L} with respect to the non-standard inner product $\langle A, B \rangle_q = \text{Tr}[A^\dagger B q]$.)

In a spin system, we find that $\tilde{l}_i = r_i$, which ensures that the above condition can be satisfied, regardless of whether or not \mathcal{L} possesses any degeneracies. The same relation cannot hold true for fermionic superparity-odd eigenoperators, because it would contradict with Eq. (S6) (when one sets $q \propto \mathbb{I}$). The only way for the Green's function identity to hold is if r_i and \tilde{l}_i are independent eigenoperators, which as we proved above implies that they are degenerate. Thus, fermionic systems differ from spin systems in that they cannot be made to satisfy (S22) without degeneracies in the spectrum of \mathcal{L} .

Kramers' degeneracy in quantum channels. We briefly note that a similar Kramers' degeneracy can be found in thermal quantum channels, which can describe the discrete time evolution of a system coupled to a *non-Markovian* bath. Define a quantum channel and its adjoint:

$$\mathcal{E}(x) = \sum_i E_i x E_i^\dagger, \quad \mathcal{E}^\dagger(x) = \sum_i E_i^\dagger x E_i. \quad (\text{S23})$$

The condition for trace preservation of the channel implies: $\mathcal{E}^\dagger(\mathbb{I}) = \sum_i E_i^\dagger E_i = \mathbb{I}$, which is the only condition that the Kraus operators (E_i) need to obey to be a proper channel. Suppose we further impose microreversibility: $\mathcal{E}^\dagger = \mathcal{Q}^{-1} \mathcal{T}^{-1} \mathcal{E} \mathcal{T} \mathcal{Q}$ where \mathcal{Q}, \mathcal{T} are defined as before. It is easy to show that the thermal state q is a steady state (eigenoperator of \mathcal{E} with eigenvalue 1). For physical fermionic channels, the channel superoperator can be split into even and odd superparity sectors: $\mathcal{E} = \text{Diag}[\mathcal{E}_+, \mathcal{E}_-]$. Our analysis implies that the odd superparity sector \mathcal{E}_- must be twofold degenerate.

2. KRAMERS' DEGENERACY IN QUADRATIC MODELS

We show that ‘‘quadratic Lindbladians’’ can host a twofold degeneracy in their single-particle spectrum in the absence of microreversibility, as long as the system-environment coupling respects time-reversal symmetry. This is in contrast to the (quartic) models studied in the main text, where spectral splitting can emerge due to a non-equilibrium environment even if all couplings respect TRS. The importance of microreversibility is therefore only apparent in quartic models.

Consider a quadratic Hamiltonian $H = \sum_{ij} H_{ij} \alpha_i \alpha_j$ in the presence of linear dissipators $L_\mu = \sum l_{\mu,i} \alpha_i$, where α_i are Majorana fermions. All terms in the master equation are quadratic in fermion operators, which implies that the Lindbladian can be split into superparity sectors: $\mathcal{L} = \mathcal{L}_+ + \mathcal{L}_-$. Define the superoperators: $e_j^\dagger(\rho) = [\alpha_j \rho + (\mathcal{P} \rho) \alpha_j]/2$, $e_j(\rho) = [\alpha_j \rho - (\mathcal{P} \rho) \alpha_j]/2$, where α_j are Majoranas, and \mathcal{P} is the parity superoperator. Then we can express \mathcal{L}_+ as

$$\mathcal{L}_+ = 4i \sum_{ij} (Z_{ji} e_i^\dagger e_j) + (Y_{ij} e_i^\dagger e_j^\dagger) = \sum_i \epsilon_i \beta_i^\dagger \beta_i', \quad (\text{S24})$$

where $Z = H + i\text{Re}[M]$, $Y = \text{Im}[M]$, $M = l^T l^*$, $\epsilon_i/(4i)$ are the eigenvalues of Z , and $\text{Re}[\epsilon_i] < 0$ [20, 61]. Analogously

$$\mathcal{L}_- = 4i \sum_{ij} (Z_{ji} e_i e_j^\dagger) + (Y_{ij} e_i e_j) = \sum_i \epsilon_i + \sum_i (-\epsilon_i) \eta_i^\dagger \eta_i'. \quad (\text{S25})$$

The first term $\epsilon_m \equiv \sum_i \epsilon_i$ is a negative offset, then excitations have a *positive* real energy. The many-body eigenvalues are thus built from single-particle eigenvalues $\{\epsilon_i\}$.

Consider a spin-1/2 Hamiltonian with TRS:

$$H = \frac{1}{2} (\alpha)^T H \alpha, \quad \alpha = (a_{1,+}, b_{1,+}, \dots, a_{1,-}, b_{1,-}, \dots)^T, \quad (\text{S26})$$

where a, b are Majoranas which transform via $T a_{i,\sigma} T^{-1} = \sigma a_{i,-\sigma}$, $T b_{i,\sigma} T^{-1} = -\sigma b_{i,-\sigma}$, and $H = U H^* U^\dagger$. We include arbitrary linear dissipators which transform into each other (up to a phase) upon action of the symmetry operator:

$$L_{i,+} = \sqrt{\gamma_i} f_{i,+}, \quad L_{i,-} = \sqrt{\gamma_i} f_{i,-}, \quad T f_{i,+} T^{-1} = f_{i,-}, \quad T f_{i,-} T^{-1} = -f_{i,+}. \quad (\text{S27})$$

The dissipators can be expanded in terms of Majoranas: $L_{i,+} = \vec{l}_{i,+} \cdot \alpha$, $L_{i,-} = \vec{l}_{i,-} \cdot \alpha$, which defines the matrix $M = l^T l^*$. We now show that $M = UM^*U^\dagger$:

$$UM^*U^\dagger = U(l^\dagger l)U^\dagger = \begin{pmatrix} \vec{l}_{i,-} \\ -\vec{l}_{i,+} \\ \vdots \end{pmatrix}^T \begin{pmatrix} \vec{l}_{i,-} \\ -\vec{l}_{i,+} \\ \vdots \end{pmatrix}^* = M, \quad (\text{S28})$$

where we have used $U\vec{l}_{i,+}^\dagger = \vec{l}_{i,-}^T$, $U\vec{l}_{i,-}^\dagger = -\vec{l}_{i,+}^T$ in the second equality. In the last equality, we have used the fact that the Lindbladian is invariant under a relabeling and a change of sign of the dissipators. This expression implies that the spectral matrix $Z = H + i\text{Re}[M]$ satisfies: $Z = UZ^T U^\dagger$, $U^2 = -\mathbb{I}$, which ensures that the single-particle spectrum is twofold degenerate. Note that we have not imposed microreversibility (detailed balance) for this result. Quadratic Lindbladians are thus special in the sense that the odd superparity sector can host a degeneracy in the absence of thermal equilibrium (unlike the examples in the main text).

3. MICROREVERSIBILITY FROM TRS-INVARIANT SYSTEM-BATH COUPLING

Here, we demonstrate that the Lindbladian of a Markovian open system will respect the microreversibility condition (5) if time-reversal symmetry is imposed on the system and bath as a whole. Before any Markovian approximation is made, the system and bath can be described by a Hamiltonian

$$H_{\text{tot}} = H_S \otimes \mathbb{I}_B + \mathbb{I}_S \otimes H_B + H_{SB}, \quad (\text{S29})$$

where H_S , H_B are the system and bath Hamiltonians, respectively, and H_{SB} couples the two. Without loss of generality, we can decompose the latter as [52]

$$H_{SB} = \sum_{\alpha} A_{\alpha} \otimes B_{\alpha}, \quad (\text{S30})$$

where A_{α} , B_{α} are Hermitian matrices.

We suppose that the bath is initialized in a thermal Gibbs state $\rho_B = Z_B^{-1} e^{-\beta H_B}$, where $\beta = 1/T$ is the inverse temperature, and $Z_B = \text{Tr} e^{-\beta H_B}$ is the partition function for the bath. Then define two-time correlators $\Gamma_{\alpha\beta}(t) = \text{Tr}[B_{\alpha}(t)B_{\beta}(0)\rho_B]$, where $B_{\alpha}(t) = e^{iH_B t} B_{\alpha} e^{-iH_B t}$. Hermiticity of B_{α} implies that

$$\Gamma_{\alpha\beta}(t)^* = \Gamma_{\beta\alpha}(-t). \quad (\text{S31})$$

The expectation values $\langle B_{\alpha} \rangle := \text{Tr} B_{\alpha} \rho_B$ can always be made to vanish by replacing $B_{\alpha} \rightarrow B_{\alpha} - \langle B_{\alpha} \rangle$, and adding a term A_{α} to H_S , which does not change H_{tot} . We also define the ‘lowering’ operators $A_{\alpha}(\omega)$, which are the components of A_{α} that decrease the energy of the system by ω . More concretely,

$$A_{\alpha}(\omega) = \sum_{\epsilon' - \epsilon = \omega} \Pi_{\epsilon} A_{\alpha} \Pi_{\epsilon'}, \quad (\text{S32})$$

where Π_{ϵ} is a projector onto the eigenspace of H_S with eigenvalue ϵ .

In order for the open system to be Markovian, the two-time correlation functions must decay over a timescale τ_m that is sufficiently short, and the system-bath coupling H_{SB} is sufficiently weak. If these criteria are met, then one can derive an expression for the Lindbladian in terms of the components of the microscopic Hamiltonian (S29) [52]

$$\mathcal{L}[\rho] = -i[H_S + H_{LS}, \rho] + \sum_{\omega} \sum_{\alpha\beta} \tilde{\Gamma}_{\alpha\beta}(\omega) \left(A_{\beta}(\omega) \rho A_{\alpha}(\omega)^\dagger - \frac{1}{2} \left\{ A_{\alpha}(\omega)^\dagger A_{\beta}(\omega), \rho \right\} \right), \quad (\text{S33})$$

where $\tilde{\Gamma}_{\alpha\beta}(\omega) = \int dt e^{i\epsilon t} \Gamma_{\alpha\beta}(t)$ is the Fourier transform of the two-time correlation functions, which is a Hermitian matrix due to (S31). Here, we have defined the Lamb shift Hamiltonian $H_{LS} = \sum_{\omega, \alpha, \beta} S_{\alpha\beta}(\omega) A_{\alpha}(\omega)^\dagger A_{\beta}(\omega)$, where $S_{\alpha\beta}(\omega) = \int dt \text{sgn}(t) e^{i\epsilon t} \Gamma_{\alpha\beta}(t)$, which is Hermitian and commutes with H_S .

Now, TRS of the combined system and bath implies that H_S and H_B are each TRS-invariant, and that $\mathcal{T}_{SB}[H_{\text{tot}}] = H_{\text{tot}}$, where the superoperator \mathcal{T}_{SB} acts as $\mathcal{T}_{SB}[O] = (U_S \otimes U_B) O^* (U_S^\dagger \otimes U_B^\dagger)$ for any operator O over the system-bath Hilbert space. The operator U_S is the unitary part of the TRS transformation acting on the system, as appears in

Eq. (2), and similarly U_B is a unitary operator acting on the bath, which we leave unspecified for full generality. In terms of the decomposition (S30), we have

$$U_S[A_\alpha]^* U_S^\dagger = \sum_\beta u_{\alpha\beta} A_\beta \quad U_B[B_\alpha]^* U_B^\dagger = \sum_\beta [u^{-1}]_{\beta\alpha} B_\beta \quad (\text{S34})$$

for some matrix u . Hermiticity of A_α, B_α implies that u is a real matrix, and $\mathcal{T}^2 = \mathcal{P}$ implies that $[uu^*]_{\alpha\beta} = p_\alpha \delta_{\alpha\beta}$, where $p_\alpha = +1$ (-1) if A_α is a fermion parity even (odd) operator. The above conditions generalise the notion of ‘weak’ symmetries described in Ref. [62] to include antiunitary symmetry operations, with the difference that we impose restrictions on the microscopic Hamiltonian, rather than the emergent master equation.

Now, if the bath is in thermal equilibrium at inverse temperature β , then this imposes a Kubo-Martin-Schwinger (KMS) condition (or ‘detailed balance’) on the spectral functions $\Gamma_{\alpha\beta}(\omega)$ [63]:

$$\tilde{\Gamma}_{\alpha\beta}(-\omega) = e^{-\beta\omega} \tilde{\Gamma}_{\beta\alpha}(\omega). \quad (\text{S35})$$

Furthermore, since H_B is TRS-invariant, we can use the transformation property (S34) to determine how the two-time correlation functions transform under TRS

$$\begin{aligned} \Gamma_{\alpha\beta}(t) &= Z_B^{-1} \text{Tr} \left[e^{iH_B t} B_\alpha e^{-iH_B t} B_\beta e^{-\beta H_B} \right] \\ &= Z_B^{-1} \text{Tr} \left[U_B \left(e^{iH_B(t+i\beta)} B_\alpha e^{-iH_B t} B_\beta \right)^* U_B^\dagger \right]^* \\ &= Z_B^{-1} \sum_{\gamma\delta} [u^{-1}]_{\gamma\alpha} [u^{-1}]_{\delta\beta} \text{Tr} \left[e^{-iH_B(t-i\beta)} B_\gamma e^{iH_B t} B_\delta \right]^* \\ &= Z_B^{-1} \sum_{\gamma\delta} [u^{-1}]_{\gamma\alpha} [u^{-1}]_{\delta\beta} \text{Tr} \left[B_\delta e^{-iH_B t} B_\gamma e^{iH_B(t+i\beta)} \right] \\ &= \sum_{\gamma\delta} [u^{-1}]_{\gamma\alpha} [u^{-1}]_{\delta\beta} \Gamma_{\delta\gamma}(t). \end{aligned} \quad (\text{S36})$$

The above can be Fourier transformed to obtain an analogous condition for $\tilde{\Gamma}_{\alpha\beta}(\omega)$.

With all these identities in hand, we are ready to begin our proof. We start by showing that the Gibbs state for the system $\rho_G = Z_S^{-1} e^{-\beta H_S}$ (where $Z_S = \text{Tr} e^{-\beta H_S}$ is the system partition function) is a steady state. We have

$$\begin{aligned} \mathcal{L}[\rho_G] &= \frac{1}{Z_S} \sum_\omega \sum_{\alpha\beta} \tilde{\Gamma}_{\alpha\beta}(\omega) \left(A_\beta(\omega) e^{-\beta H_S} A_\alpha(\omega)^\dagger - \frac{1}{2} \left\{ A_\alpha(\omega)^\dagger A_\beta(\omega), e^{-\beta H_S} \right\} \right) \\ &= \frac{1}{Z_S} \sum_\omega \sum_{\alpha\beta} \tilde{\Gamma}_{\alpha\beta}(\omega) A_\beta(\omega) e^{-\beta H_S} A_\alpha(\omega)^\dagger - \frac{1}{2} \tilde{\Gamma}_{\beta\alpha}(\omega) \left\{ A_\beta(-\omega) A_\alpha(-\omega)^\dagger, e^{-\beta H_S} \right\} \\ &= \frac{1}{Z_S} \sum_\omega \sum_{\alpha\beta} \left(\tilde{\Gamma}_{\alpha\beta}(\omega) e^{-\beta\omega} A_\beta(\omega) A_\alpha(\omega)^\dagger - \tilde{\Gamma}_{\beta\alpha}(\omega) A_\alpha(-\omega) A_\beta(-\omega)^\dagger \right) e^{-\beta H_S} = 0. \end{aligned} \quad (\text{S37})$$

In the first equality, we use $[H_{LS}, H_S] = 0$. The second equality involves a swap of labels α and β , and uses relation $A_\alpha(-\omega) = A_\alpha(\omega)^\dagger$, which can be verified using (S32). The third equality requires the relation $e^{-\beta H_S} A_\alpha(\omega)^\dagger = A_\alpha(\omega)^\dagger e^{-\beta(H_S + \omega)}$, and finally we replace $\omega \rightarrow -\omega$ in the second term and employ Eq. (S35).

Now we verify that Eq. (5) is satisfied. Recall that $\mathcal{Q}[A] = qA$ for any operator A , and here $q = \rho_G$ is the steady-state density matrix. We can separate out $\mathcal{L} = \mathcal{L}_c + \mathcal{L}_d$, where $\mathcal{L}_c[\rho] = -i[H_S + H_{LS}, \rho]$ contains the coherent part of the evolution, and the remainder \mathcal{L}_d contains the dissipative part. Then the coherent part of the right-hand side of (5) acting on an arbitrary density operator ρ gives (remembering that \mathcal{T} and \mathcal{T}^{-1} are antiunitary superoperators)

$$\begin{aligned} \mathcal{Q}^{-1} \mathcal{T}^{-1} \mathcal{L}_c \mathcal{T} \mathcal{Q}[\rho] &= \rho_G^{-1} \cdot \mathcal{T}^{-1} \left[-i[H_S + H_{LS}, \mathcal{T}[\rho_G \rho]] \right] \\ &= +i \rho_G^{-1} \left[\mathcal{T}^{-1}[H_S + H_{LS}], \rho_G \rho \right] = +i[H_S + H_{LS}, \rho_G^{-1} \rho_G \rho] = \mathcal{L}_c^\dagger[\rho], \end{aligned} \quad (\text{S38})$$

where we have used the fact that that $H_S + H_{LS}$ is TRS-invariant and commutes with ρ_G^{-1} . Now the dissipative part

is

$$\begin{aligned}
\mathcal{Q}^{-1} \mathcal{T}^{-1} \mathcal{L}_d \mathcal{T} \mathcal{Q}[\rho] &= \rho_G^{-1} \cdot \mathcal{T}^{-1} \left[\sum_{\omega} \sum_{\alpha\beta} \tilde{\Gamma}_{\alpha\beta}(\omega) \left[A_{\beta}(\omega) \rho_G \rho A_{\alpha}(\omega)^{\dagger} - \frac{1}{2} \left\{ A_{\alpha}(\omega)^{\dagger} A_{\beta}(\omega), \mathcal{T}[\rho_G \rho] \right\} \right] \right] \\
&= \rho_G^{-1} \sum_{\omega} \sum_{\alpha\beta\gamma\delta} [u^{-1}]_{\alpha\gamma} [u^{-1}]_{\beta\delta} \tilde{\Gamma}_{\alpha\beta}(\omega)^* \left(A_{\delta}(\omega) \rho_G \rho A_{\gamma}(\omega)^{\dagger} - \frac{1}{2} \left\{ A_{\gamma}(\omega)^{\dagger} A_{\delta}(\omega), \rho_G \rho \right\} \right) \\
&= \rho_G^{-1} \sum_{\omega} \sum_{\gamma\delta} \tilde{\Gamma}_{\gamma\delta}(\omega) \left(A_{\delta}(\omega) \rho_G \rho A_{\gamma}(\omega)^{\dagger} - \frac{1}{2} \left\{ A_{\gamma}(\omega)^{\dagger} A_{\delta}(\omega), \rho_G \rho \right\} \right) \\
&= \sum_{\omega} \sum_{\gamma\delta} \tilde{\Gamma}_{\gamma\delta}(\omega) \left(e^{-\beta\omega} A_{\delta}(\omega) \rho_G \rho A_{\gamma}(\omega)^{\dagger} - \frac{1}{2} \left\{ A_{\gamma}(\omega)^{\dagger} A_{\delta}(\omega), \rho_G \rho \right\} \right) \\
&= \sum_{\omega} \sum_{\gamma\delta} \tilde{\Gamma}_{\delta\gamma}(-\omega) A_{\delta}(\omega) \rho_G \rho A_{\gamma}(\omega)^{\dagger} - \frac{1}{2} \tilde{\Gamma}_{\gamma\delta}(\omega) \left\{ A_{\gamma}(\omega)^{\dagger} A_{\delta}(\omega), \rho_G \rho \right\}, \tag{S39}
\end{aligned}$$

having used the transformation properties of A_{α} under \mathcal{T}^{-1} [Eq. (S34) with u replaced by u^{-1}] in the second equality; Eq. (S36) in the third equality; and Eq. (S35) in the final equality. This can be compared to \mathcal{L}_d^{\dagger}

$$\begin{aligned}
\mathcal{L}_d^{\dagger}[\rho] &= \sum_{\omega} \sum_{\gamma\delta} \tilde{\Gamma}_{\gamma\delta}(\omega)^* \left(A_{\delta}(\omega)^{\dagger} \rho A_{\gamma}(\omega) - \frac{1}{2} \left\{ A_{\delta}(\omega)^{\dagger} A_{\gamma}(\omega), \rho \right\} \right) \\
&= \sum_{\omega} \sum_{\gamma\delta} \tilde{\Gamma}_{\delta\gamma}(\omega) \left(A_{\delta}(-\omega) \rho A_{\gamma}(-\omega)^{\dagger} - \frac{1}{2} \left\{ A_{\delta}(\omega)^{\dagger} A_{\gamma}(\omega), \rho \right\} \right), \tag{S40}
\end{aligned}$$

having used $A_{\alpha}(-\omega) = A_{\alpha}(\omega)^{\dagger}$. Direct comparison verifies that (S39) and (S40) are indeed equal, thus confirming that Eq. (5) is satisfied.

4. LINEAR RESPONSE IN OPEN QUANTUM SYSTEMS

This section shows the connection between the retarded Green's function in Eq. (13) in the main text and the current in tunneling spectroscopy experiments. We demonstrate that the standard formula for closed systems also applies to open quantum systems upon a generalization of operator time evolution.

Kubo formula. Consider a general Markovian dynamics in the form

$$\frac{d}{dt} \rho = \left(\mathcal{L}_0 + \lambda \mathcal{L}_1(t) \right) \rho, \tag{S41}$$

where \mathcal{L}_0 is the unperturbed Lindbladian, $\lambda \mathcal{L}_1(t)$ is time-dependent perturbation superoperator, and λ is a small perturbation parameter. Assume $\mathcal{L}_0(t) = \mathcal{L}_1 \Theta(t - t_0)$ for some initial time t_0 , where $\Theta(t - t')$ is the Heaviside step function. We also choose the initial state to be the steady state of the unperturbed system, $\rho(t_0) = \rho_{\text{SS}}$ defined as $\mathcal{L}_0 \rho_{\text{SS}} = 0$. In the lowest order of perturbative expansion over λ , the system's dynamics can be represented by a Dyson series,

$$\begin{aligned}
\rho(t) &= \rho_{\text{SS}} + \lambda \int_{t_0}^t dt' \exp(\mathcal{L}_0(t - t')) \mathcal{L}_1 \exp(\mathcal{L}_0 t') \rho_{\text{SS}} + O(\lambda^2) \\
&= \rho_{\text{SS}} + \lambda \int_{t_0}^t dt' \exp(\mathcal{L}_0(t - t')) \mathcal{L}_1 \rho_{\text{SS}} + O(\lambda^2). \tag{S42}
\end{aligned}$$

Now let us consider the time-dependent expectation value of a local observable O defined as $O(t) = \text{Tr}(O\rho(t))$. Its

dynamics can be expressed using Eq. (S42) as

$$\begin{aligned}
O(t) &= \langle O \rangle_{\text{SS}} + \lambda \int_{t_0}^t dt' \text{Tr} \left(O \exp(\mathcal{L}_0(t-t')) \mathcal{L}_1 \rho_{\text{SS}} \right) + O(\lambda^2) \\
&= \langle O \rangle_{\text{SS}} + \lambda \int_{t_0}^t dt' \text{Tr} \left(\rho_{\text{SS}} \mathcal{L}_1^\dagger \exp(\mathcal{L}_0^\dagger(t-t')) O \right) + O(\lambda^2) \\
&= \langle O \rangle_{\text{SS}} + \lambda \int_{t_0}^t dt' \text{Tr} \left(\rho_{\text{SS}} \mathcal{L}_1^\dagger O(t-t') \right) + O(\lambda^2),
\end{aligned} \tag{S43}$$

where $\langle O \rangle_{\text{SS}} = \text{Tr} O \rho_{\text{SS}}$, $O(t) = \exp(\mathcal{L}_0^\dagger t) O$, and $\mathcal{L}_0^\dagger, \mathcal{L}_1^\dagger$ are conjugate Liouvillian operators as defined in Eq. (6) in the main text. We focus on unitary perturbation, $\mathcal{L}_1 = -i[H_1, \cdot]$, where H_1 is perturbation Hamiltonian. Extending $t_0 \rightarrow -\infty$, we obtain

$$O(t) = \langle O \rangle_{\text{SS}} - i\lambda \int_{-\infty}^{\infty} dt' \Theta(t-t') \langle [O(t-t'), H_1] \rangle_{\text{SS}} + O(\lambda^2). \tag{S44}$$

For unitary dynamics, this expression coincides with the conventional Kubo formula.

Tunneling spectroscopy. To measure the spectral properties of the system, we assume that the system of interest is connected to a tunneling probe as shown in Fig. 1 in the main text. The probe contains a reservoir of electrons able to tunnel into the system, thus generating an electric current. The dependence of the current on the probe's chemical potential, i.e. differential conductance, can be used to find the spectral function of the system.

Let us consider Eq. (S41) applied to a joint fermionic system-probe configuration. Without the perturbation, we assume that the system and the probe are decoupled, i.e. the unperturbed Liouvillian has the form $\mathcal{L}_0 = \mathcal{L}_0^S + \mathcal{L}_0^R$, where \mathcal{L}_0^S and \mathcal{L}_0^R are Liouville operators acting on system or the probe, respectively. Also, we assume that the initial state is a product state, $\rho_{\text{SS}} = \rho_{\text{SS}}^S \otimes \rho_{\text{SS}}^R$. The coupling between the system and the probe is produced by the perturbation

$$H_1 = \sum_{\mu\nu} (T_{\mu\nu} f_\mu^\dagger b_\nu + \text{h.c.}) = B + B^\dagger, \tag{S45}$$

where f_μ and b_ν are Fock operators for the system and the probe respectively, $T_{\mu\nu}$ are tunnelling coefficients, and $B = \sum_{\mu\nu} T_{\mu\nu} f_\mu^\dagger b_\nu$. Here and below μ and ν are generalized indices that incorporate several quantum numbers, including the electron's position and spin.

The tunneling current is defined through the change of the probe's electron number $N_D = \sum_\nu b_\nu^\dagger b_\nu$, namely

$$I := i\lambda[H_1, N_D] = -i\lambda \sum_{\mu\nu} (T_{\mu\nu} f_\mu^\dagger b_\nu - \text{h.c.}) = -i\lambda(B - B^\dagger). \tag{S46}$$

Using the Kubo formula in Eq. (S44), we obtain

$$I(t) = 2\lambda^2 \text{Re} \int_{-\infty}^{\infty} dt' \Theta(t-t') \left(\langle [B^\dagger(t-t'), B] \rangle_{\text{SS}} - \langle [B(t-t'), B] \rangle_{\text{SS}} \right), \tag{S47}$$

where $B(t) = \sum_{\mu\nu} T_{\mu\nu} f_\mu^\dagger(t) b_\nu(t)$ including $f_\mu(t) := \eta_d \exp(\mathcal{L}_0^S t) [\eta_d f_\mu]$ and $b_\nu(t) := \eta_d \exp(\mathcal{L}_0^R t) [\eta_d b_\nu]$, where η_d is dummy Majorana fermion operator as defined in Eq. (S11) (see discussion in Section 1). Assuming that the probe is in the normal (i.e. not superconducting) state, the last term in Eq. (S47) vanishes.

$$\begin{aligned}
I(t) &= 2\lambda^2 \text{Re} \int_{-\infty}^{\infty} dt' \Theta(t-t') \sum_{\mu\nu} \sum_{\mu'\nu'} T_{\mu\nu}^* T_{\mu'\nu'} \left(\langle f_\mu^\dagger(t-t') f_{\mu'} \rangle_{\text{SS}} \langle b_\nu(t-t') b_{\nu'}^\dagger \rangle_{\text{SS}} - \langle f_\mu(t-t') f_{\mu'}^\dagger \rangle_{\text{SS}} \langle b_\nu^\dagger(t-t') b_{\nu'} \rangle_{\text{SS}} \right) \\
&= 2\lambda^2 \text{Re} \int_{-\infty}^0 d\tau \sum_{\mu\nu} \sum_{\mu'\nu'} T_{\mu\nu}^* T_{\mu'\nu'} \left(G_{\mu\mu'}^<(\tau) D_{\nu\nu'}^>(-\tau) - G_{\mu\mu'}^>(\tau) D_{\nu\nu'}^<(-\tau) \right) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} I(\omega),
\end{aligned} \tag{S48}$$

where $G_{\mu\mu'}^<(\tau) = -i \langle f_\mu^\dagger(\tau) f_{\mu'} \rangle_{\text{SS}}$ and $G_{\mu\mu'}^>(\tau) = -i \langle f_\mu^\dagger f_{\mu'}(\tau) \rangle_{\text{SS}}$ are the greater and the lesser Green's function, $D_{\nu\nu'}^<(\tau)$ and $D_{\nu\nu'}^>(\tau)$ are similar expressions for the probe in terms of b_ν , and $I(\omega)$ are the Fourier components of the

tunneling current defined as

$$I(\omega) = \lambda^2 \sum_{\mu\nu} \sum_{\mu'\nu'} T_{\mu\nu}^* T_{\mu'\nu'} \left(G_{\mu\mu'}^<(\omega) D_{\nu}^>(\omega) - G_{\mu\mu'}^>(\omega) D_{\nu}^<(\omega) \right). \quad (\text{S49})$$

Let us assume that the probe's reservoir of fermions is at thermal equilibrium at zero temperature and b_{ν} are eigenmodes of the reservoir. Then, the expression for probe's Green's functions takes the form

$$D_{\nu\nu'}^>(\omega) = -i\delta_{\nu\nu'} \tilde{A}_{\nu}(\mu + \omega) \left(1 - n_F(\mu + \omega) \right), \quad D_{\nu\nu'}^<(\omega) = i\delta_{\nu\nu'} \tilde{A}_{\nu}(\mu + \omega) n_F(\mu + \omega), \quad (\text{S50})$$

where $n_F(x) = \Theta(x)$ is zero-temperature Fermi distribution, μ is probe's chemical potential, and \tilde{A}_{ν} is the probe's spectral function.

Assuming $\tilde{A}_{\nu}(V + \omega) \simeq \tilde{A}_{\nu}$ depends weakly on the chemical potential V , and introducing coefficients $J_{\mu\mu'} = 2\lambda^2 \sum_{\nu} T_{\mu\nu}^* T_{\mu'\nu} \tilde{A}_{\nu}$, we obtain the expression for the differential conductance

$$\frac{\partial I(\omega)}{\partial \mu} = \sum_{\mu\mu'} J_{\mu\mu'} \text{Im} G_{\mu\mu'}^R(\omega) \delta(\omega + \mu), \quad (\text{S51})$$

where $\delta(x)$ is the Dirac delta function, $G_{\mu\mu'}^R(\omega) = G_{\mu\mu'}^<(\omega) + G_{\mu\mu'}^>(\omega)$ is the Fourier transform of the time-domain retarded Green's function defined in Eq. (13) in the main text. Choosing appropriate filtering $J_{i\sigma',j\sigma''} \propto \delta_{ij} \delta_{\sigma'\sigma''} \delta_{\sigma\sigma'}$, we can measure spin-polarized current characteristics,

$$\frac{\partial I_{\sigma}}{\partial \mu} \propto \text{Im} G_{i\sigma}^R(-\mu). \quad (\text{S52})$$

Thus, changing the chemical potential μ , we can probe the spectral function of the system.