

The complexity of a quantum system and the accuracy of its description

Yuri I. Ozhigov^{1,2},

*1. Moscow State University of M.V.Lomonosov,
Faculty VMK,*

2. Institute of physics and technology of K.A.Valiev

May 10, 2021

Abstract

The complexity of the quantum state of a multiparticle system and the maximum possible accuracy of its quantum description are connected by a relation similar to the coordinate-momentum uncertainty relation. The coefficient in this relation is equal to the maximum physically adequate dimension of the Hilbert space of states. This value is the binary exponent of the maximum number of qubits whose dynamics can be adequately described by quantum theory, and therefore it can be determined experimentally through Grover search algorithm. Such a restriction of the Copenhagen formalism is relevant for complex systems; it gives a natural description of unitary dynamics together with decoherence and measurement, but also implies the existence of a minimum non-zero amplitude size, as well as a restriction on the equality of bases in the state space. The quantization of the amplitude allows us to formally introduce a certain kind of determinism into quantum evolution, which is important for complex systems.

1 Introduction

Our understanding of quantum theory has evolved greatly since its inception. If until about 80-90 years of the 20th century, as a rule, simple, from the classical point of view, systems were studied: individual atoms, molecules or ensembles consisting of identical particles that could be reduced to separate independent simple objects, then in recent decades the focus of research has shifted towards more complex systems. In particular, the relevance of microbiology and virology has also aroused physicists' interest in studying objects related to living things, for example, the DNA molecule, which can no longer be attributed to simple systems.

Meanwhile, quantum theory, which is the basis of our understanding of the microcosm, and, therefore, an accurate understanding of complex systems, has a very rigid and well-defined mathematical apparatus based on the matrix technique. The predictions of quantum mechanics have always proved to coincide very precisely with experiments on simple systems that are traditional for physics, but for complex systems this theory meets a fundamental obstacle. The very procedure of obtaining theoretical predictions requires such unimaginable computational resources that we will never have them at our disposal.

If for simple systems the procedure of computation the quantum state had no relation to the physics of its evolution and was only a technical technique, then for complex systems the situation is different. Here, the computation process is the main part of the definition of the quantum state itself, and therefore should be considered as a physical process, and the device that implements this computation is an integral part of any experiment with complex systems at the quantum level.

This computing device is an abstract computer that simulates the evolution of the complex system under consideration. Thus, all restrictions on this computer, following from the theory of algorithms, have the status of physical laws; and these laws have absolute priority over physical laws in the usual sense in the case of complex systems and processes.

This is a new situation that did not exist in classical physics, where the procedure for obtaining theoretical predictions was not very complicated. In any case, the complexity there has almost always been within the reach of classical supercomputers, which are created mainly to cover processes from the point of view of classical physics - by the number of particles in the system under consideration. In quantum mechanics, the complexity increases exponentially with the number of particles, and the classical way of computing becomes unacceptable. This was strictly proved by the discovery of theoretically possible (from the point of view of the standard - Copenhagen quantum theory) processes that cannot be modeled on any classical supercomputers - the so-called fast quantum algorithms ([1], [2]).

The attempt to circumvent the complexity barrier with the help of a quantum computer proposed by R. Feynman ([4]) has given us a lot to understand the microcosm and some interesting applications, for example, in cryptography and metrology. However, this attempt did not solve the main problem: the scaling of a fully functional quantum computer is very questionable due to decoherence. Decoherence occurs as a result of spontaneous measurements of the states of the simulated system from the environment, which is traditionally considered in the framework of the concept of an open quantum system in contact with the environment (see [5]), so that the influence of the environment is reduced to uncontrolled measurements of the state of the original system.

Thus, decoherence is a fundamental factor that cannot be eliminated with the help of mathematical techniques, such as error correction codes (they begin to really work only for a quantum computer with more than a hundred qubits). If we set the task of modeling complex systems at the quantum level, decoherence should

be embedded in the quantum formalism itself, and not introduced into it as an extraneous influence. The deviation from the linear unitary law of evolution resulting from decoherence must be naturally justified mathematically.

There is a complexity barrier to matrix formalism. If we have a quantum state of a system of c particles with two possible states that cannot be simplified, and a is the total number of binary signs that determine the position of all these particles, then for the total number of bits $a + c$, the inequality $a + c \leq q$ must be satisfied, where the dimensionless constant q can be determined experimentally.

If we move from the strings of qubit values $\alpha_0, \alpha_1, \dots, \alpha_{n-1}$ to the numbers defined by their binary expansions $\sum_{j=0}^{n-1} a_j 2^j$, then $a = cn$ and we get the classical analogs of accuracy and complexity: $A = 2^a$, $C = 2^c$, for which a relation similar to the coordinate - momentum uncertainty relation is satisfied:

$$AC \leq Q, \tag{1}$$

where $Q = 2^q$.

We will argue in favor of such a ratio of complexity and accuracy in the general case, for complex particles; in particular, we will show that the quantization of the amplitude makes it possible to introduce a certain determinism into the quantum formalism, the nature of which is not reduced to the classical one.

2 The universal computer

A complex system is a system whose behavior cannot be reduced to independent particles. Predictions of the behavior of such a system can be obtained only by relying on quantum mechanics and computer ideology, since analytical techniques do not work here. Therefore, we must introduce the concept of a Universal Computer (MC), which adequately represents real processes - an abstract computing device, the laws of which will be for complex systems (and only for them!) have priority over physical laws, the scope of which is unconditionally limited to simple systems and processes.

A physical prototype of a universal mainframe computer will be built in the foreseeable future-there is no doubt about that. Such a device will be able to adequately represent the processes traditionally related to chemistry, as well as to those areas of physics in which quantum methods are successfully applied, for example, to electrodynamics. Nuclear physics does not yet belong to such processes, and its complexity radically exceeds the electrodynamics complexity (see [7]).

The prototype of a universal computer may be classical, but then it will not be able to correctly represent quantum dynamics due to the inability to reflect the long-range effect of quantum nonlocality; the second reason may be complex, if it turns out that the constant Q in nature is too large.

The difference between the classical and quantum versions of the MC is that if the classical one is able to directly calculate the amplitudes of all the components of the state, then the quantum MC only simulates the initial state, representing it as the state of the quantum part of the computer.

The accuracy of a state is the accuracy of its amplitudes, and the complexity is the number of basic states in the canonical representation (see below). In the quantum case, the unit of measurement of these quantities is one qubit, in the classical case-one bit. If a is the quantum precision of the state vector of a complex system, and c is its quantum complexity, then the classical analogues of these quantities are found by the formulas $A = 2^a$, $C = 2^c$, so in any case the inequalities will be satisfied

$$a + c \leq q, \quad AC \leq Q. \quad (2)$$

The fulfillment of these inequalities is a condition for the adequacy of this representation of the state vector for both quantum and classical universal computers.

Thus, we have, within the framework of this limitation of the computational capabilities of the MC, a balance between the complexity and accuracy of the representation of the state vector, which must be found specifically in each specific case.

For a very complex system, precision means the total number of its basic states in the canonical representation, each of which will be either occupied or free, like the fill numbers, so that the values of the amplitudes will only be binary. For a system of intermediate complexity, you can more accurately determine the amplitudes - see figure 1.

3 Complexity of the quantum state

In order to give a correct definition of the complexity of the quantum state of a system of n particles, we must first consider the canonical transformation - the main method of reducing such complexity.

Any coordinate on a unit segment is a real number represented as its binary expansion $2^{-l} \sum_{j=0}^{l-1} a_j 2^j$ with an accuracy of 2^{-l} , where $a_j = 0,1$ are the values of l qubits representing this coordinate¹ By ordering the qubit values lexicographically, we get a standard ordering of basis vectors, in which any operator will have a certain matrix.

Let the classical state of the particle i be the real vector x_i . Then the classical state of the system of n particles will be a vector of the form $\bar{x} = (x_1, x_2, \dots, x_n)$. Let

¹To represent a coordinate on any other segment, we need to apply a suitable linear transformation, for example, on the segment $[-2^{l/2}, 2^{l/2}]$ the approximate qubit representation looks like this: $2^{-l/2} \sum_{j=0}^{l-1} a_j 2^j - 2^{l/2-1}$.




Figure 1: Representation of the state vector. Bottom left - the main computing resource captures accuracy: $|\Psi\rangle = \lambda_0|0\rangle + \lambda_1|1\rangle$, bottom right - the main resource is captured by complexity: $|\Psi\rangle = \sum_{j=0}^{N-1} \epsilon|j\rangle$, $\epsilon \in \{0, \epsilon\}$, at the top-the resource is divided equally between accuracy and complexity.

\bar{x}' and \bar{x}'' be two vectors with non-empty sets of coordinates such that their Cartesian product coincides with \bar{x} . This means that we have split the set of particles into two non-empty subsets X' and X'' , so that the given vectors are the sets of coordinates of these subsets. Let $H(\bar{x})$ be the Hamiltonian of this system, having the form

$$H(\bar{x}) = H_1(\bar{x}') + H_2(\bar{x}''); \quad (3)$$

here, we assume by default that $H(\bar{x}')$ is $H(\bar{x}') \otimes I(\bar{x}'')$, that is, on particles not included in the first subset, this term of the Hamiltonian acts as an identical operator, and similarly with the second term. Then we call the Hamiltonian H reducible. Let X' be the maximum subset of the components of the vector \bar{x} in terms of the number of elements, such that the equality (3) holds, and the Hamiltonian $H_1(\bar{x}')$ is not reducible. Then X' is called the kernel of the given Hamiltonian.

We will assume by default that any quantum evolution begins with the basic state of the system under consideration. Since (3) implies the equality $\exp(-\frac{i}{\hbar}H) = \exp(-\frac{i}{\hbar}H_1) \otimes \exp(-\frac{i}{\hbar}H_2)$, we see that the core of the Hamiltonian is the maximum set of particles whose states in quantum evolution with the Hamiltonian H can be entangled; we denote the number of particles in this set by $\nu(H)$, and call it the *naive* complexity of this Hamiltonian.

Consider the transformation of the coordinates of the particles of the form

$$q_i = q_i(x_1, x_2, \dots, x_n), \quad i = 1, 2, \dots, n; \quad (4)$$

denote $\bar{q} = (q_1, q_2, \dots, q_n)$ and let $H_q = H(x_1(\bar{q}), x_2(\bar{q}), \dots, x_n(\bar{q}))$ is the original Hamiltonian written in terms of the new variables q_i , $\bar{x} = \bar{x}(\bar{q})$ is the inverse of (4). We will introduce virtual particles with coordinates q_1, q_2, \dots, q_n , which we will call quasiparticles.

The classical state of the source system is a set of specific values x_1, x_2, \dots, x_n . Then each classical state will correspond to the classical state of the same system, obtained using the formulas (4). The basis vector of a Hilbert space passes into another basis vector of the same space. The standard ordering of the basis vectors, according to the qubit representation of coordinates, will pass to another ordering, that is, the coordinate transformation (4) is a permutation of the basis vectors of the Hilbert space of quantum states.

In this case, the qubits representing the coordinate values will already have a different, new meaning. In the new coordinates, for quasiparticles, the Hamiltonian will have a different form H_q . We will call the coordinate transformation (4) canonical if $\nu(H_q)$ is minimal. In this case, the transition to quasiparticles will mean a reduction in the complexity of the original Hamiltonian. So, the canonical transformation is a permutation of the basis vectors that minimizes the complexity of the Hamiltonian.

The transition to the description of evolution in the form of quasiparticles has the form $H = \tau^{-1}H_q\tau$ where τ is the transformation of the transition to quasiparticles under the canonical transformation. Then the representation of the evolution

operator is $\exp(-\frac{i}{\hbar}Ht) = \tau^{-1}\exp(-\frac{i}{\hbar}H_qt)\tau$ requires less computational resources than the direct calculation of $\exp(-\frac{i}{\hbar}Ht)$, since the main resource is spent on the kernel, which for a quasi-partial representation will have a minimum size.

An example is a chain of interacting harmonic oscillators, for which the Fourier transform of their coordinates x_i is canonical, and the kernel of the quasiparticle representation-in terms of phonons - will generally consist of a single particle, that is, the phonons are all independent. Here we have the maximum reduction of the kernel.

A simpler example would be the Hamiltonian H of a closed chain of 4 interacting qubits, which is reduced by the canonical transformation $CNOT$ to a completely reduced Hamiltonian of the form $H_q = \sigma_x^{(1)} \otimes I_2 + I_1 \sigma_x^{(2)}$:

$$H = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix} = CNOT \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix} CNOT.$$

Note that the permutation of the basis vectors, which is a canonical transformation, must be an entangling and simultaneously disentangling operator in the state space, since it reduces the kernel of the Hamiltonian. So, for example, the $CNOT$ operator applied repeatedly to the state $|00\dots 0\rangle + |11\dots 1\rangle$ completely untangles this state.

The opposite example is given by the Tavis-Cummings Hamiltonian for n two-level atoms interacting with the resonant mode field in the optical cavity. Here there is a basic state of the field and atoms of the form $|n\rangle_{ph}|00\dots 0\rangle_{at}$, such that the column of the Hamiltonian matrix corresponding to this state consists of the numbers $g\sqrt{n}$ and one number $n\hbar\omega$, and such a column is the only one. No Hamiltonian of the form (3) can have this property even at $n = 2$, so there is no non-identical canonical transformation for the Tavis-Cummings Hamiltonian.

The quantum complexity of the Hamiltonian H , denoted by $c(H)$, is the minimal naive complexity of the operators $\tau^{-1}H\tau$ over all possible permutations of τ basis vectors. For the above examples, the complexity of the Hamiltonian is 1, that is, it can be completely reduced by the canonical transformation.

The complexity of the quantum state $|\Psi\rangle$ is determined in a similar way. Its *naive* complexity is $\nu(|\Psi\rangle)$ is defined as the number of particles in the maximum tensor divisor $|\Psi_1\rangle$ of the state $|\Psi\rangle = |\Psi_1\rangle \otimes |\Psi_2\rangle$.

The quantum complexity of the state $|\Psi\rangle$ is the minimal naive complexity of the state $\tau|\Psi\rangle$ over all permutations of τ particles. In other words, the quantum complexity of $c(|\Psi\rangle)$ of a state is the logarithm of the kernel length of that state in the canonical representation.

For example, the quantum complexity of the generalized GHZ state $|GHZ\rangle = \frac{1}{\sqrt{2}}(|00\dots 0\rangle + |11\dots 1\rangle)$ is equal to 1, since it can be untangled by successive CNOT operators. If we start from the basic state $|\Psi(0)\rangle$ in the canonical representation of the Hamiltonian H , then an evolution with H will only contain states with a

complexity of no more than $c(H)$. The classical complexity is defined as $C(\Psi) = 2^{c(|\Psi\rangle)}$.

The quantum complexity of the Hamiltonian has a clear algorithmic meaning. Suppose we model quantum evolution on a classical supercomputer with an unlimited parallelization resource. Then, for the reduced Hamiltonian, you can parallelize such a simulation by assigning separate nodes of the supercomputer to simulate separate groups of variables. Thus, the node that will be assigned the most difficult task will be the one that will be assigned the kernel of the Hamiltonian. We can say that the classical complexity of the Hamiltonian is the memory size of a single node of a classical supercomputer that can simulate quantum evolution with a given Hamiltonian. The quantum complexity of the Hamiltonian is the logarithm of this value.

4 Amplitude granularity

The qubit representation of the classical coordinates and impulses determines the grain size of the amplitudes. For any expansion of the quantum state $|\Psi\rangle$ over an arbitrary orthonormal set of basis states $|\psi_j\rangle$ of the form

$$|\Psi\rangle = \sum_{j \in J} \lambda_j |\psi_j\rangle, \quad \lambda_j \neq 0, \quad (5)$$

the amplitudes of λ_j must be bounded from below modulo some nonzero constant.

To preserve the principle of linearity in the region where the nonzero value of this constant does not play a role, we must assume that any amplitude has the form

$$\lambda_j = \epsilon n_j + i \epsilon m_j \quad (6)$$

where $n_j, m_j \in \mathbb{Z}$ are integers, $\epsilon > 0$ is a constant that is the quantum of the amplitude. This restriction of the matrix formalism entails the rejection of the absolute equivalence of any bases in the space of quantum states, which also manifests itself only at sufficiently small amplitudes λ_j , and, accordingly, large sets of coherent states J .

However, modeling does not provide for such equivalence of bases; it is only an algebraic technique that cannot be experimentally tested for complex systems. The smallest modulo possible nonzero amplitude is thus ϵ . This restriction is very well consistent with the probabilistic nature of the state vector, since to determine the value of $|\lambda_j|^2$ with an accuracy δ , it is necessary to conduct about $1/\delta$ measurements of equally prepared samples of the original system; for complex systems with small amplitudes of λ_j , this will be possible only if the minimum probability ϵ^2 is separated from zero to obtain the most unlikely outcome.

You can determine the value of ϵ by "smearing" the amplitude over as many basic classical states of the system as possible, for which the presence in them can simply be detected by measurement. If all the amplitudes are equal to: $\lambda_j = \epsilon$, we get for

the total number of coherent basis states the estimate $|J| = [1/\epsilon^2]$ - as the integer part of the inverse square of the amplitude quantum. In the (5) expansion, the amplitudes of λ_j have no physical dimension, the dimension has the basis states of $|\psi_j\rangle$. The number of qubits whose possible quantum states are physically realizable is equal to the constant q , for which we get the expression $q = \log_2([1/\epsilon^2])$.

We can now formulate a hypothesis about the relation of the form "accuracy - complexity" in the final form:

$$C(|\Psi\rangle)A(|\Psi\rangle) \leq Q, \quad (7)$$

where $Q = 2^q$, and q is the maximum number of completely entangled qubits that cannot be disentangled by any permutation of the basis states.

Consider, as an example, the state of a set of n qubits of the form

$$|\Psi_{GSA}(t)\rangle = \alpha \sum_{j \neq j_0, 0 \leq j < N} |j\rangle + |j\rangle, \quad (8)$$

where $\alpha = \cos(t)/\sqrt{N-1}$, $\equiv \sin(t)$ for some t , and $N = 2^n$. The quantum complexity of this state is n if $t \neq k\pi/2$ for no integer k . Indeed, this superposition has the property that all its basic components, except for exactly one, have the same non-zero amplitude, and one has a different amplitude from them.

This property is preserved under any permutation of the basis states, that is, under any quasiparticle representation. But if the state is reducible, then it should have the form $\lambda_1|i_1\rangle + \lambda_2|j_2\rangle + \dots \otimes (\lambda_3|j_3\rangle + \lambda_4|j_4\rangle + \dots)$ for some basis $|j_i\rangle$, and such a superposition cannot contain exactly 2 amplitude values for any basis states, since there must either be at least 3 different non-zero amplitude values of the components, or it must have only two different non-zero amplitude values that correspond to two groups of basis states containing an equal number of terms. Both of these possibilities are excluded for states of the form (8).

Note that the discrete representation of amplitudes in the form of (11) allows us to naturally include state measurements in its unitary evolution. The hard contact of the system in the state $|\Psi\rangle = \sum_{j \in J} \lambda_j |j\rangle$ with the measuring device means the inclusion of the states of this device in the system, that is, the transition to the state of the extended system of the form $|\Psi_{ex}\rangle = \sum_{j \in J, \nu_j \in \mathcal{N}_j} \mu_{j, \nu_j} |j, \nu_j\rangle$, where, under the condition of the applicability of quantum mechanics, all μ_{j, ν_j} must be minimal, which means that the quantum of the amplitude of eis equal. Since contact with the environment is a unitary evolution, we have $|\lambda_j|^2 = \sum_{\nu_j \in \mathcal{N}_j} |\mu_{j, \nu_j}|^2$ that is, the measurement will be an arbitrary choice of one of the states of the ν_j meter and we get the standard urn scheme from probability theory.

5 Experimental finding of a constant Q

You can find the approximate value of Q by the building the state of the form (8). These states have the memory of a quantum computer when implementing the Grover GSA algorithm with a single target state $|j_0 \text{rangle}$ (see [2]). Let $N = 2^n$. Let's say $t_0 = \arcsin(1/\sqrt{N})$. Then at $|\Psi(t_0)\rangle = \frac{1}{\sqrt{N}} \sum_{j=0}^{N-1} |j\rangle = (\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle))^{\otimes n}$, and the complexity of this state is 1; this is the initial state for the GSA algorithm. As soon as the first step of the algorithm is performed, t becomes equal to $t_0 + 2t_0$ and we get a state of complexity N , of the form (8). Already at the first step of the algorithm, the complexity jumps from one to the maximum value of N for n qubit particles. If $N = Q$, at the first step of the GSA algorithm, we will go beyond the limits of acceptable states with amplitudes of the form (11), which have the property (7).

Thus, we can estimate the constant Q from above, increasing the value of N to the limit when the GSA stops working correctly. Here, by correct operation, we mean the possibility to raise the amplitude of the target state by an amount of the order of $1/\sqrt{2}$ compared to the others, which can be fixed by quantum tomography, since the amplitudes of other states will have the order of $1/\sqrt{N}$. For a more rough estimate, fixing the jump in the amplitude by $1/\sqrt{N}$ compared to the main mass is also suitable, but this is only possible for small n , not exceeding 20.

The question of what will happen to the real state if the amplitudes calculated according to standard quantum theory become less than ϵ is formally open. However, it is natural to assume that small amplitudes should simply disappear, with a corresponding renormalization of the remaining state. This means that implementing GSA near the boundary of $N \approx Q$ we will get the target state very quickly, much faster than when implementing GSA in a normal model. However, this will only happen with an ideal implementation of GSA; in practice, the amplitudes of the main mass of states in $|\Psi_{GSA}(t)\rangle$ in (8) cannot be exactly the same, so that zeroing will not occur simultaneously for all states, which can greatly distort the picture.

6 Equilibrium states

We will begin by describing the classes of states for which the amplitude quantization is introduced in the most visual way.

For a complex number $z = a + ib$, $a, b \in R$, we introduce the notation $\{z\} = |a| + |b|$. For the quantum state $|\psi\rangle$, we define $\{\psi\} = \sum_{i=0}^{N-1} \{ \langle i|\psi\rangle \}$.

Let A be a linear operator, and $|j\rangle$ be some basis vector, $j \in \{0, 1, \dots, N-1\}$. Define $|a_j\rangle = A|j\rangle$. We call the state vector $|\Psi\rangle$ equilibrium with respect to the operator A , if all the numbers $\{a_j\}$ are the same for all the base components $|j \text{rangle}$

that are included in it with nonzero amplitudes.

As an example, consider the Hamiltonian of a one-dimensional particle moving in the potential V : $H = \frac{p^2}{2m} + V$. We will reduce the matrix of this Hamiltonian, assuming that there are no too long transitions of a given particle in space. Then the equilibrium states in the coordinate basis for this Hamiltonian will be exactly the states $|\Psi\rangle$, all the basis components of which have the same potential.

An important class of multiparticle equilibrium states are connected states. Here is an example of such a state. Consider k of two-level atoms in an optical cavity holding photons with the transition energy between the ground and excited levels of the atoms. We choose a basis consisting of vectors of the form $|n\rangle_{ph}|m_1, m_2, \dots, m_k\rangle_{at}$, where n is the number of photons in the cavity, $m_j \in \{0, 1\}$ is the state of the atom j , ground and excited. Let g_j , $j = 1, 2, \dots, k$ be the forces of the interaction of atoms with the field. Then the dynamics of the system of atoms and the field under the condition $g_j/\hbar\omega \ll 1$, where ω is the frequency of the cavity, will obey the Schrodinger equation with the Tavis-Cummings Hamiltonian in the RWA approximation:

$$H_{TC}^{RWA} = \hbar\omega(a^+a + \sum_{j=1}^k \sigma_j^+ \sigma_j) + a^+ \bar{\sigma} + a \bar{\sigma}^+, \quad \bar{\sigma} = \sum_{j=1}^k g_j \sigma_j, \quad (9)$$

where a, a^+ are the standard field operators of photon annihilation and creation, and σ_j, σ_j^+ are the atomic relaxation and excitation operators of the atom j . The connected states in such a system will be for $k = 2$ only for $g_1 = g_2$, and this will be either one of the basic states, or the states $|n\rangle_{ph}(\alpha|10\rangle_{at} + |01\rangle_{at})$, of which in all but the singlet state $\equiv -\alpha$ the atoms will interact with the field. All such states will be in equilibrium.

The general definition of connectivity looks like this.

Let H be a Hamiltonian in the state space of n qubits. If a qubit is associated with a real or virtual two-level particle, H can be, for example, the Tavis-Cummings Hamiltonian or some modification of it. Let S_n be a group of permutations of qubits that are naturally extended to operators on the entire space of quantum states \mathcal{H} , namely: on the basis states, the permutation $\eta \in S_n$ acts directly, and $\eta \sum_j |j\rangle = \sum_j \eta |j\rangle$.

Denote by G_H the subgroup S_n consisting of all permutations of qubits τ such that $[H, \tau] = 0$. Let $A \subseteq \{0, 1, \dots, 2^n - 1\}$ be a subset of the basis states of the n -qubit system. Its linear shell $L(A)$ is called a connected subspace with respect to H if for any two states $|i\rangle, |j\rangle \in A$ there exists a permutation of qubits $\tau \in G_H$ such that $\tau(i) = j$. The state of a $|\Psi\rangle$ n -qubit system is called connected with respect to H if it belongs to a connected subspace with respect to H , and $H|\Psi\rangle \neq is0$.

The connectedness of a state means that all its nonzero components are obtained from one another by permutations of those particles that behave in the same way with respect to a given Hamiltonian. The above example of the state $|n\rangle_{ph}(\alpha|10\rangle_{at} + |01\rangle_{at})$ will obviously be connected, since the permutation of atoms

interacting equally with the field does not change the Hamiltonian. The states of the form $|n\rangle_{ph}(\alpha|10\rangle_{at} + \beta|01\rangle_{at} + c|00\rangle_{at} + d|11\rangle_{at})$, for non-zero values of the amplitudes a, b, c, d will not be connected.

Lemma.

If $|\Psi\rangle = \sum_j \lambda_j |j\rangle$ is connected with respect to H , then any two columns of the matrix H with numbers j_1, j_2 , such that λ_{j_1} and λ_{j_2} are nonzero, differ from each other only by the permutation of elements. The same is true for the unitary evolution matrix $U_t = \exp(-\frac{i}{\hbar}Ht)$.

Indeed, for such basis states j_1 and j_2 , according to the definition of H - connectivity, there exists $\tau \in G_H$ such that $j_2 = \tau(j_1)$. The columns numbered j_1, j_2 consist of the amplitudes of the states $H|j_1\rangle$ and $H|j_2\rangle$, respectively. From the commutation condition, we have $\tau H|j_1\rangle = H\tau|j_1\rangle = H|j_2\rangle$, and this just means that the column j_2 is obtained from the column j_1 by the permutation of elements induced by τ . Moving on to the evolution matrix U_t , we see that the switching relation $\tau U_t |j_1\rangle = U_t \tau |j_1\rangle = U_t |j_2\rangle$ will hold for it as well, which is what is required. Lemma is proven.

It follows from Lemma that the states connected with respect to the Hamiltonian H are equilibrium with respect to H and with respect to the evolution operator $U_t = e^{-\frac{i}{\hbar}Ht}$ corresponding to this Hamiltonian.

7 Amplitude quanta and determinism

The exact description of the dynamics, even in the classical framework, has some degree of nondeterminism or stochasticity (see [8],[9]). The advantage of the quantum language lies in the precise limitation of this stochasticity to the state vector, which, according to the main thesis of Copenhagen mechanics, provides an exhaustive description of the dynamics of microparticles.

The possibility of introducing determinism into quantum theory, which interested researchers at the beginning of the history of quantum physics, has not lost the interest of researchers (see [10]) and is again becoming relevant for complex systems; for example, for systems of extreme complexity, the DNA molecule determines the trajectory of its owner with an accuracy unattainable in physical experiments. This higher type of determinism must have an analog for the simple systems described by standard quantum theory.

We describe a possible specific form of determinism at the level of amplitude quanta.

Our goal is to show that if the state $|\Psi\rangle$ is equilibrium with respect to the evolution operator U_t , then the amplitudes of all the basic states in $|\Psi\rangle$ can be divided into small portions - quanta of amplitude, so that for each quantum its trajectory will be uniquely determined when U_t acts on a previously fixed time interval t , in particular,

it will be uniquely determined, and with which other quantum of amplitude it will contract when summing the amplitudes to obtain the subsequent states.

This fact is also true for the arbitrary operator A , for which we will formulate the quantization of the amplitude.

Let $|\Psi\rangle$ be an arbitrary equilibrium state with respect to A , the decomposition of which in terms of the basis has the form

$$|\Psi\rangle = \sum_j \lambda_j |j\rangle. \quad (10)$$

We introduce the important concept of the amplitude quantum as a simple formalization of the transformation of a small portion of the amplitude between different base states when multiplying the state vector by the matrix A . Let $T = \{+1, -1, +i, -i\}$ be a set of 4 elements, which are called amplitude types: real positive, real negative, and similar imaginary ones. The product of types is defined in a natural way: as a product of numbers. The quantum of the amplitude of the size $\varepsilon > 0$ is called a list of the form

$$\kappa = (\varepsilon, id, |b_{in}\rangle, |b_{fin}\rangle, t_{in}, t_{fin}), \quad (11)$$

where $|b_{in}\rangle, |b_{fin}\rangle$ are two different basic states of the system of atoms and photons, id is a unique identification number that distinguishes this quantum from all others, $t_{in}, t_{fin} \in T$. Transition of the form $|b_{in}\rangle \rightarrow |b_{fin}\rangle$ is called a state transition, $t_{in} \rightarrow t_{fin}$ is called a type transition. Let's choose the identification numbers so that if they match, all the other attributes of the quantum also match, that is, the identification number uniquely determines the quantum of the amplitude. In this case, there must be an infinite number of quanta with any set of attributes, except for the identification number. Thus, we will identify the quantum of the amplitude with its identification number, without specifying this in the future. Let's introduce the notation:

$$t_{in}(\kappa) = t_{in}, \quad t_{fin}(\kappa) = t_{fin}, \quad s_{in}(\kappa) = b_{in}, \quad s_{fin}(\kappa) = b_{fin}.$$

The transitions of states and types of amplitude quanta actually indicate how a given state should change over time, and their choice depends on the choice of A ; the size of the amplitude quantum indicates the accuracy of a discrete approximation of the action of this operator using amplitude quanta.

The set of θ quanta of the amplitude of the size ε is called the quantization of the amplitude of this size, if the following condition is met:

Q. In the set θ , there are no such quanta of the amplitude κ_1 and κ_2 that their state transitions are the same, $t_{in}(\kappa_1) = t_{in}(\kappa_2)$ and $t_{fin}(\kappa_1) = -t_{fin}(\kappa_2)$, and there are no such quanta of amplitude κ_1 and κ_2 that $s_{in}(\kappa_1) = s_{in}(\kappa_2)$ and $t_{in}(\kappa_1) = -t_{in}(\kappa_2)$.

The condition **Q** means that during the transition described by the symbol " \rightarrow ", the total value of the amplitude quantum cannot contract with the total value

of a similar amplitude quantum, and also that the amplitude quanta do not contract with each other directly in the initial state record.

Quantization of the θ amplitude sets a pair of quantum states

$$|\theta_{in}\rangle = \sum_j \lambda_j |j\rangle, \quad |\theta_{fin}\rangle = \sum_i \mu_i |i\rangle, \quad (12)$$

according to the natural rule: for any basis states $|j\rangle, |i\rangle$, the equalities must be satisfied

$$\lambda_j = \langle j|\theta_{in}\rangle = \varepsilon \sum_{\kappa \in \theta: s_{in}(\kappa)=j} t_{in}(\kappa), \quad \mu_i = \langle i|\theta_{fin}\rangle = \tilde{\varepsilon} \sum_{\kappa \in \theta: s_{fin}(\kappa)=i} t_{fin}(\kappa), \quad (13)$$

where $\tilde{\varepsilon}$ is some normalization coefficient, so that the state $|\theta_{fin}\rangle$ has a unit norm, and $|\theta_{in}\rangle$ has an arbitrary nonzero one. The coefficient $\tilde{\varepsilon}$ does not have to coincide with ε , because when quantizing the amplitude, the usual norm of the state vector, in general, is not preserved; if we took $\tilde{\varepsilon} = \varepsilon$, then the value we entered $\{|\Psi\rangle\}$ at the transition $|\theta_{in}\rangle \rightarrow |\theta_{fin}\rangle$ could this is exactly due to the fact that some quanta of the amplitude "contract" with each other in the second sum from the formula (13).

Let's fix the dimension $dim(\mathcal{H})$ of the state space, and we will make estimates (from above) of the considered positive quantities: the time and the size of the amplitude quantum up to an order of magnitude, considering all constants to depend only on the independent constants: $dim(\mathcal{H})$ and from the minimum and maximum absolute values of the elements of the matrix A . At the same time, the term "strict order" will mean the evaluation of both the top and bottom positive numbers that depend only on independent constants.

To quantize the amplitude θ and the numbers i, j of the basis states via $n_{i,j}(\theta)$ denote the number of elements of the set $\mathcal{N}_{i,j}(\theta) = \{\kappa \in \theta : s_{in}(\kappa) = j, s_{fin}(\kappa) = i\}$.

Let $\theta(\varepsilon)$ be some function that maps some sequence of positive numbers - values of ε converging to zero, into quantizations of the amplitude of the size ε . This function will be called parametric quantization of the amplitude.

The parametric quantization of the amplitude $\theta(\varepsilon)$ is called consistent with the operator A if there exists a function $c(\varepsilon)$ such that for any basis states i, j

$$c(\varepsilon)n_{i,j}(\theta(\varepsilon)) \rightarrow \{\lambda_j\}\{\langle i|A|j\rangle\} \quad (\varepsilon \rightarrow 0) \quad (14)$$

for λ_j , defined in (13).

If A is the evolution operator U_t , then having a parametric quantization of the amplitudes $\theta(\varepsilon)$ consistent with A is a completely non-trivial property of the quantum states $|\theta_{in}(\varepsilon)\rangle$, which says that it is possible to introduce a hidden parameter corresponding to the dynamics set by the evolution matrix U_t , and making the quantum evolution U_t deterministic. Such a parameter is the quantum of the amplitude $\kappa \in \theta(\varepsilon)$, where the accuracy of the deterministic description is determined by the value ε .

The amplitude quantization theorem.

Let A be an arbitrary matrix. For every equilibrium state $|\Psi\rangle$ with respect to A , there exists a parametric quantization of the amplitudes $\theta(\varepsilon)$, consistent with the operator A , such that

$$|\theta_{in}(\varepsilon)\rangle \rightarrow |\Psi\rangle, \quad |\theta_{fin}(\varepsilon)\rangle \rightarrow A|\Psi\rangle, \quad (\varepsilon \rightarrow 0). \quad (15)$$

Proof.

Let us be given the equilibrium state with respect to A $|\Psi\rangle = \sum_j \lambda_j |j\rangle$ and the number $\varepsilon > 0$. For $|j\rangle$ with nonzero $\lambda_j \neq 0$ let

$$\lambda_j = \langle j|\Psi\rangle \approx \underbrace{\text{sign}_{re}(\varepsilon + \varepsilon + \dots + \varepsilon)}_{M_j} + \underbrace{\text{sign}_{im}i(\varepsilon + \varepsilon + \dots + \varepsilon)}_{N_j}, \quad (16)$$

where $\text{sign}_{re}\varepsilon M_j + \text{sign}_{im}i\varepsilon N_j \approx \lambda_j$ is the best approximation of the amplitude of λ_j with the accuracy of ε ; M_j, N_j are natural numbers, $\text{sign}_{re} (im) = \pm 1$. Thus, the first tendency relation from (15) will be satisfied, and the second relation must be satisfied if the parametric quantization is consistent with the Hamiltonian.

Let us approximate each element of the evolution matrix in the same way as we approximate the amplitudes of the initial state:

$$\langle i|A|j\rangle \approx \pm \underbrace{(\varepsilon + \varepsilon + \dots + \varepsilon)}_{R_{i,j}} \pm i \underbrace{(\varepsilon + \varepsilon + \dots + \varepsilon)}_{I_{i,j}}, \quad (17)$$

where $R_{i,j}, I_{i,j}$ are natural numbers; the real and imaginary parts are exactly ε each, and the signs before the real and imaginary parts are chosen based on the fact that this approximation should be as accurate as possible for the selected ε .

The amplitudes of the resulting state $A|\Psi\rangle$ are obtained by multiplying all possible expressions (16) by all possible expressions (17):

$$\lambda_j \langle i|A|j\rangle \approx (\text{sign}_{re} M_j \varepsilon + i \text{sign}_{im} N_j \varepsilon)(\pm R_{i,j} \varepsilon \pm i I_{i,j} \varepsilon). \quad (18)$$

We will expand the brackets in the right part of the expression (18), but we will not make abbreviations. Each occurrence of the expression ε^2 in the amplitudes of the resulting state after opening the brackets in the right part of (18) will be obtained by multiplying a certain occurrence of ε in the right part of (16) by a certain occurrence of ε in the right part of (17). The problem is that the same occurrence of ε in (16) corresponds to not one, but several occurrences of ε^2 in the result, and therefore we can not match the amplitude quanta directly to the occurrences of ε in (16).

How many occurrences of ε^2 in the amplitudes of the state $A|\Psi\rangle$ from the result of opening the brackets in (18) correspond to one occurrence of ε in the approximation of the amplitude $\lambda_j = \langle j|\Psi\rangle$ of the state $|\Psi\rangle$? This number - the multiplicity of the

given occurrence of ε - is equal to $\sum_i (R_{i,j} + I_{i,j})$. These numbers can be different for an arbitrary operator A and the state $|\Psi\rangle$. However, since $|\Psi\rangle$ is equilibrium with respect to A , $\sum_i (R_{i,j} + I_{i,j})$ will be the same for different j .

We introduce the notation $\nu = \sum_i (R_{i,j} + I_{i,j})$ - this is the number of occurrences of ε in any column from the expansion of the matrix (17); this number ν has the order $1/\varepsilon$ at $\varepsilon \rightarrow 0$.

Denote by $Z_{i,j}$ the set of occurrences of the letter ε in the right part of the expression (17), and let $Z_j = \bigcup_i Z_{i,j}$. Then the number of elements in the set Z_j will be equal to ν .

Consider the smaller value of the amplitude quantum: $\epsilon = \varepsilon/\nu$. We substitute in the expression (16) instead of each occurrence of ε its formal decomposition of the form $\varepsilon = \overbrace{\epsilon + \epsilon + \dots + \epsilon}^\nu$, obtaining the decomposition of the amplitudes of the initial state into smaller numbers:

$$\lambda_j = \langle j|\Psi\rangle \approx \text{sign}_{re} \left(\overbrace{\epsilon + \epsilon + \dots + \epsilon}^\nu + \overbrace{\epsilon + \epsilon + \dots + \epsilon}^\nu + \dots + \overbrace{\epsilon + \epsilon + \dots + \epsilon}^\nu \right) + \text{sign}_{im} i \left(\overbrace{\epsilon + \epsilon + \dots + \epsilon}^\nu + \overbrace{\epsilon + \epsilon + \dots + \epsilon}^\nu + \dots + \overbrace{\epsilon + \epsilon + \dots + \epsilon}^\nu \right). \quad (19)$$

Let $W_1^j, W_2^j, \dots, W_{M_j+N_j}^j$ be the sets of occurrences of the letter ϵ in the right part of the expression (19), marked with upper curly brackets. In each of these sets of ν elements, as in the previously defined sets of Z_j . Therefore, we can construct for each such set W_s^j a one-to-one mapping of the form $\xi : W_s^j \rightarrow Z_j$. For each occurrence of ε in (16), its descendants are naturally defined - the occurrences of ϵ in (19); the descendants for each occurrence will be ν .

We define the quantization of the amplitudes $\theta = \theta(\epsilon)$ so that the *id* of the amplitude quanta $\kappa \in \theta$ will simply be the occurrences of ϵ in the expansions of (19) for all j . Define, as required in (11), the initial state and initial type of this quantum as the state and type of this occurrence. It remains to determine the transitions of states and types. This definition is given in the following natural way.

Each pair of the form $(w_s^j, \xi(w_s^j))$, where $w_s^j \in W_s^j$, will correspond to the transition of states and the transition of types in a natural way. Namely, the state transition will have the form $j \rightarrow i$ for such i that $\xi(w_s^j) \in Z_{i,j}$; the transition of the types $t_{in} \rightarrow t_{fin}$ is defined so that t_{in} is the type of occurrence² w_s^j , and the type t_{fin} is the product of the occurrence type t_{in} by the occurrence type $\xi(w_s^j)$. The sets W_s^j

²The type of occurrence is determined naturally after opening the parentheses, for example, for the occurrence of $\dots - i\epsilon\dots$ the type is $-i$.




Figure 2: A. Multiplying the state vector by the matrix. The contribution of each occurrence of ε is multiplied by ε . B. θ - initial state shift. The size of the ε amplitude quantum is of the order of ε^2 .

do not intersect at different pairs j, s , so we consider all occurrences of the letter ε in the right part of (19) to be the domain of the function definition ξ (see Fig. 2).

Now let the transition of states and types for a given quantum $\kappa \in \theta$ correspond to the mapping ξ in the sense defined above. The condition **Q** will be met, since there are no reducing terms in the expression for the matrix element (17). So we defined the quantization of the amplitude.

Due to our definition of the function ξ , the distribution of amplitudes in the state $|\theta\Psi\rangle$ will be approximately proportional to the distribution of amplitudes in the state $A|\Psi\rangle$, and the accuracy will increase indefinitely with the decrease of ε to zero. In order to determine the value of the function $c(\varepsilon)$ necessary for consistency of θ with the operator A , we calculate the contribution of each occurrence of ε^2 to the right side of the equality (18) and compare it with the contribution of the corresponding letter ε to $|\theta\Psi\rangle$.

Let's fix any transition of the types $t_{in} \rightarrow t_{fin}$ and the transition of the states $s_{in} \rightarrow s_{fin}$. We will call the occurrence of ε^2 in the result of opening brackets in (18) corresponding to these transitions if $j = s_{in}$, $i = s_{fin}$, and this occurrence is obtained by multiplying the occurrence of ε of type t_{in} in the first factor of the right side of (18) by the occurrence of ε in the second factor of type t' , so that $t_{in}t' = t_{fin}$. For each such occurrence, ε^2 corresponds to exactly one quantum of the amplitude of the size ε from the quantization of the amplitude defined above via the function ξ , which has the same transitions of states and types: this quantum corresponds to the occurrence of ε , which is translated by a one-to-one mapping ξ to this occurrence ε^2 (see Fig. 3).




Figure 3: Determinism of trajectories when multiplying the state vector $|\Psi\rangle$ by the matrix A . Each quantum of the amplitude passes into a certain quantum of the amplitude of the resulting state, there are no branches.

So, the occurrences of ϵ^2 in (18) are in one-to-one correspondence with the occurrences of ϵ in (19). How many quanta of amplitude will the transition of states of the form $|j\rangle \rightarrow |i\rangle$ have? In each bracket of (19) such quanta will be approximately $\{\langle i|A|j\rangle\}/\epsilon$ pieces. And the total number of brackets is about $\{\lambda_j\}/\epsilon$ pieces. Therefore, the number $n_{i,j}$ will be approximately equal to $\{\lambda_j\}\{\langle i|A|j\rangle\}/\epsilon^2$, and the accuracy will grow indefinitely at $\epsilon \rightarrow 0$. So, $c(\epsilon) = \epsilon^2$.

Note that if $A = 0$, you can take $c(\epsilon) = 0$ and any quantization of the amplitude will be suitable.

The theorem is proved.

Note that if we abandon the condition **Q** and the condition of the equilibrium state $|\Psi\rangle$ with respect to A , we can also define matrix determinism, only we need to introduce the reducing terms $\epsilon - \epsilon$ into the matrix elements; then the formal entries of the amplitudes for all columns of A will contain the same number of terms, and the reasoning will be valid, but interference will now occur not only between the descendants of different base states, as for equilibrium states. $|\Psi\rangle$, and also between descendants of the same state.

Note also that a quantum computation in which only gates of the form *CNOT* and Hadamard are used has the property that for any gate the number of elements in all Z_j will be the same, so that for such calculations determinism is provided with interference only between images of different basis states. This is, in particular, the Grover GSA algorithm.

Moreover, for promising implementations of quantum gates on photons (see, for example, [6]), the states resulting from the implementation of gates are connected,

so that interference in the course of such quantum calculations also has the property noted above.

8 Conclusion

We have argued for limiting the standard matrix formalism of quantum theory in the field of complex systems, as an upper bound on the possible number of qubits that can have an irreducible entangled state. This approach does not contradict any experiments with multi-bit systems, but allows you to build models of such systems on existing supercomputers. This restriction also applies to the equivalence of bases in the state space, and implies the existence of a minimum nonzero amplitude in the superposition. The size of this quantum of amplitude can be approximately found in experiments on the implementation of Grover's algorithm. The quantization of the amplitude makes it possible to introduce some type of determinism into quantum theory, which is not reduced to a quasi-classical approximation.

9 Acknowledgements

The work was supported by the Russian Foundation for Basic Research, grant a-18-01-00695.

References

- [1] Shor P. Algorithms for Quantum Computation: Discrete Logarithms and Factoring // Foundations of Computer Science, 1994 Proceedings., 35th Annual Symposium on — IEEE, 1994. — P. 124–134. — ISBN 0-8186-6580-7 — doi:10.1109/SFCS.1994.365700
- [2] L.Grover, A fast quantum mechanical algorithm for database search, Proceedings, 28th Annual ACM Symposium on the Theory of Computing (STOC), May 1996, pages 212-219. Proceedings, Melville, NY, 2006, vol. 810.
- [3] R.Feynman, QED: The strange theory of light and matter, Princeton University Press, 1985.
- [4] Richard P. Feynman, Simulating Physics with Computers, International Journal of Theoretical Physics, Vol 21, Nos. 6/7, 1982, pp. 467-488.
- [5] H. Breuer and F. Petruccione, The Theory of Open Quantum Systems, Oxford (2002).
- [6] Azuma H., Quantum computation with the Jaynes-Cummings model, Prog. Theor. Phys. 126, 369-385 (2011).

- [7] Y.Ozhigov, About quantum computer software, *Quantum Information and Computation*, Vol. 20, No. 7&8 (2020) 570-580.
- [8] Flavio Del Santo, Nicolas Gisin , Reply to a "Comment on 'Physics without determinism: Alternative interpretations of classical physics' ", *Phys. Rev. A* 102, 036202 (2020)
- [9] Flavio Del Santo and Nicolas Gisin, Physics without determinism: Alternative interpretations of classical physics, *Phys. Rev. A* 100, 062107, 2019, DOI:<https://doi.org/10.1103/PhysRevA.100.062107>
- [10] Arthur Jabs, A conjecture concerning determinism, reduction, and measurement in quantum mechanics, *Quantum Studies: Mathematics and Foundtions*, Vol.3 (4) 279-292, 2016.